Maximum directed cuts in digraphs with degree restriction

Jenő Lehel *

ehel * Frédéric Maffray [†] Myriam Preissmann [‡]

November 15, 2018

Abstract

For integers $m, k \ge 1$, we investigate the maximum size of a directed cut in directed graphs in which there are m edges and each vertex has either indegree at most k or outdegree at most k.

1 Introduction

We deal with directed graphs, called here *digraphs*, without loops and parallel edges. An edge xy of a digraph is interpreted as an arc or an *arrow* going from the starting vertex or *tail* x to the end vertex or *head* y. The *indegree* and the *outdegree* of a vertex $v \in V(D)$ is respectively defined as $d_D^-(v) = |\{zv \in E(D)|z \in V(D)\}|$ and $d_D^+(v) = |\{vw \in E(D)|w \in V(D)\}|$.

Let X, Y be a partition of the vertex set V(D) of a digraph D. The edge set $\{xy \in E(D) | x \in X, y \in Y\}$ is called a *directed cut*. Clearly a directed cut of a digraph D does not contain a directed path on three vertices (a P_3). On the other hand every directed P_3 -free subgraph of D is the subgraph of some directed cut. Thus when estimating the size of maximum directed cuts we must find directed P_3 -free subgraphs as large as possible. The *size of a cut* is its cardinality, the *size of a digraph* is the cardinality of its edge set.

Discussions in [1] show that a digraph D of size m has a cut of size $\frac{1}{4}m + \Theta(m^{1/2})$. Furthermore, if the outdegree of each vertex of D is at most k, then D has a cut of size at least $(\frac{1}{4} + \frac{1}{8k+4})m$. In [6], lower bounds for the largest directed cuts were asked for a family of digraphs with constrained indegree or outdegree. Let $D(k, \ell)$ be the family of all digraphs in which every vertex has either indegree at most k or outdegree at most ℓ (that is $d^-(v) \leq k$ or $d^+(v) \leq \ell$,

^{*}Department of Mathematical Sciences, The University of Memphis, Tennessee, USA and Computer and Automation Research Institute of the Hungarian Academy of Sciences, Budapest, Hungary; jlehel@memphis.edu

 $^{^{\}dagger}{\rm CNRS},$ Laboratoire G-SCOP, 46 avenue Félix Viallet, 38031 Grenoble Cedex, France; frederic.maffray@g-scop.inpg.fr

[‡]CNRS, Laboratoire G-SCOP, 46 avenue Félix Viallet, 38031 Grenoble Cedex, France; myriam.preissmann@g-scop.inpg.fr

for all $v \in V(D)$). Note that a directed cut (or any directed P_3 -free graph) forms a graph that belongs to D(0,0).

In Section 2 we consider the case $k = \ell = 1$ and discuss the size of the maximum directed cut of digraphs in D(1,1). It was proved in [1] that every acyclic digraph of size m in D(1,1) has a directed cut of at least 2m/5 edges. From a result of Bondy and Locke [4] it is easy to see that the same lower bound holds for maximum directed cuts in triangle-free subcubic digraphs (a graph is *subcubic* if it has maximum degree at most three). Our main result in Theorem 1 is the extension of this bound for all digraphs in D(1,1) as follows: if D contains at most t pairwise disjoint directed triangles, then D has a directed cut of size at least (2m - t)/5. The proof yields a polynomial algorithm which actually finds a directed cut of that size (Corollary 2).

Theorem 1 implies that every digraph of size m in D(1, 1) has a directed cut with at least m/3 edges (a result first proved in [1]). Furthermore, every connected digraph of size m in D(1, 1) has a directed cut with at least 7m/20edges (see Theorem 5).

In Section 3 we consider digraphs in D(k, k) for any k. First we prove a decomposition property in Theorem 8: the edge set of every digraph in $D(p_1 + p_2, p_1 + p_2)$ can be partitioned into two subgraphs one in $D(p_1, p_1)$ and the other in $D(p_2, p_2)$. In Theorem 10 we prove the lower bound (2k-1)m/(2k+1) on the maximum size of a subgraph of D belonging to D(k-1, k-1). It is worth noting that the regular tournament on 2k + 1 vertices has no subgraph in D(k-1, k-1) with more than (2k-1/k)m/(2k+1) edges.

In Section 4 we show that if $D \in D(k, k)$ is acyclic and has m edges, then it contains a directed cut of size at least $(\frac{1}{4} + \frac{1}{8k+4})m$ (see Theorem 12). It is worth noting that in a digraph $D \in D(k, k)$ one cannot guarantee a directed cut of size larger than that proportion. This is shown by the regular tournament on 2k + 1 vertices, which has no directed cut of size more than $(\frac{1}{4} + \frac{1}{8k+4})\binom{2k+1}{2}$ (see in [1]). For k = 2 this ratio is 3m/10. In Theorem 13 we can show that actually every digraph $D \in D(2, 2)$ with m edges has a directed cut of size at least 3m/10. In the proof of Theorems 12 and 13 we use an elementary counting method similar to those applied in [1].

Section 5 concludes with open problems for further consideration. A challenging question whose answer we would like to see the most is whether Theorem 12 remains true for all digraphs in D(k, k), and for every $k \ge 3$.

2 Maximum directed cut of digraphs in D(1,1)

It was proved in [1] that every acyclic digraph of size m in D(1, 1) has a directed cut of at least 2m/5 edges. This is not true for all digraphs in D(1, 1). For example the directed triangle, which is a member of D(1, 1), has no directed cut with two edges. Hence there are digraphs of size m with maximum directed cut not larger than m/3. On the other hand, it was shown in [1] that the edge set of every digraph $D \in D(1, 1)$ has a decomposition into three directed cuts (see Theorem 7 below), hence D always contains a directed cut of size m/3. One might conclude that the ratio m/3 cannot be improved to 2m/5 in general, and the graph that consists of disjoint directed triangles is an obvious example showing that. Actually the following example shows that, for infinitely many values of m, there are even connected digraphs in D(1, 1) of size m, that contain no cut of size 3m/8.

Example 1. For i = 1, ..., k let H_i be a directed path with five vertices $(u_i, v_i, w_i, x_i, y_i)$ plus the chord $v_i x_i$, such that the H_i 's are pairwise disjoint. Add k + 1 directed triangles (y_i, u_{i+1}, z_i) , for i = 0, ..., k, where y_0, u_{k+1} and $z_0, z_1, ..., z_k$ are distinct new vertices. The obtained graph H has m = 8k + 3 edges and its maximum directed cut has size 3k + 1 = (3m - 1)/8.

In spite of the evidence that the maximum directed cut size to edge count ratio 2/5 cannot be achieved, we show in the next theorem that 1/3 improves to 2/5, in some sense, for all digraphs in D(1,1).

Theorem 1. Let D be a digraph in D(1,1) with m edges, and let t be the maximum number of pairwise disjoint directed triangles in D. Then D has a directed cut of size at least (2m - t)/5.

Proof. The claim is clearly true for $m \leq 3$ and t = 0. If m = 3 and t = 1, then D is the directed triangle and any edge of the triangle forms a directed cut of size 1 = (2m - t)/5. Now let D be a counterexample with $m \geq 4$ edges, and assume that the theorem is true for all digraphs in D(1, 1) with at most m - 1 edges. Clearly D is connected.

Let D^+ be the subgraph of D induced by $V^+ = \{v \in V(D) | d^+(v) \ge 2\}$; and let D^- be the subgraph of D induced by $V^- = \{v \in V(D) | d^-(v) \ge 2\}$. Notice that $v \in V^+$ implies that $d^-(v) \le 1$ and $v \in V^-$ implies $d^+(v) \le 1$.

Because $D \in D(1, 1)$, if two directed triangles of D have a common vertex, then they must share a common edge. Moreover, if a triangle intersects with at least two other triangles, then they all share the same common edge. The following property of triangles will be useful.

Claim 1.1. Every directed triangle of D is contained in D^+ or in D^- .

Assume that T = (x, y, z) is a directed triangle with $d^-(x) = 1$ and $d^+(y) = 1$. Remove the edges of T from D. The graph D' that remains has m' = m - 3 edges, and the maximum number t' of disjoint triangles in D' satisfies $t' \leq t - 1$. By induction, D' contains a directed cut K of size at least $(2m' - t')/5 \geq (2m - t)/5 - 1$. Obviously, $K \cup \{xy\}$ is still a directed P_3 -free subgraph of D containing (2m - t)/5 edges, a contradiction. Therefore, either $d^+(w) \geq 2$ for all $w \in \{x, y, z\}$ or $d^-(w) \geq 2$ for all $w \in \{x, y, z\}$. In the first case $T \subset D^+$ and in the second case $T \subset D^-$. Thus Claim 1.1 holds.

Let $A, B \subset E(D)$ be a pair of disjoint edge sets such that every directed P_3 in D that has one edge in A has its second edge in B. Note that this implies that A contains no directed P_3 . We call any such pair $A, B \subset E(D)$ a reducing pair. It is clear that if K is any directed cut in the digraph $D \setminus (A \cup B)$, then $K \cup A$ is a directed P_3 -free subgraph of D. The following claim will be used several times in the induction step.

Claim 1.2. D has no reducing pair $A, B \subset E(D)$ with $|B| \leq \frac{3}{2}|A|$.

Suppose that $A, B \subset E(D)$ is a reducing pair with $|B| \leq \frac{3}{2}|A|$. Let K be a largest directed cut in the digraph $D' = D \setminus (A \cup B)$. Then $K \cup A$ is a directed P_3 -free subgraph of D. Digraph D' has $m' = m - |A \cup B|$ edges, hence it follows by induction that $|K| \geq (2m' - t)/5$. We obtain

$$|K \cup A| \ge \frac{2m'-t}{5} + |A| \ge \frac{2m-t}{5} - \frac{2}{5}(|A|+|B|) + |A| \ge \frac{2m-t}{5},$$

which contradicts the assumption that D is a counterexample to the theorem. Thus Claim 1.2 holds.

Claim 1.3. Each of D^+ and D^- is a disjoint union of directed cycles. Furthermore, every vertex in D^+ or D^- is incident with exactly one edge of $D \setminus (D^+ \cup D^-)$.

Let C be any connected component of D^- . We show that C is a directed cycle. By the definition of D^- , C is either a rooted tree with all edges directed towards the root, or a *function graph* which is a rooted tree plus an edge from the root to some vertex of the tree.

If C is not a directed cycle, then it is either a singleton vertex v_0 or it has a leaf v_0 . In each case, because v_0 is in V^- , there exist distinct edges $e_1 = v_1v_0, e_2 = v_2v_0$ of D. Furthermore, $d^+(v_0) \leq 1$, thus at most one edge f_0 leaves v_0 . Since v_1, v_2 are not in C, they are not in V^- , hence at most one edge enters each, say f_1 and f_2 , respectively. Note that both edges exist in $A = \{e_1, e_2\}$, but any edge from the set $B = \{f_0, f_1, f_2\}$ might actually not exist. In either case, A, B form a reducing pair with $|B| \leq \frac{3}{2}|A|$, contradicting Claim 1.2. Thus every component of D^- is a directed cycle. Furthermore, if there are two edges e_1, e_2 of $D \setminus (D^+ \cup D^-)$ at some vertex $v_0 \in V^-$, then one obtains a contradiction using the same reducing pair.

An analogous argument shows that every connected component C of D^+ is a directed cycle with exactly one edge of $D \setminus (D^+ \cup D^-)$ at each vertex of C. Thus Claim 1.3 holds.

Note that, due to Claims 1.1 and 1.3, all directed triangles of D are among the cycles of D^+ and D^- .

Claim 1.4. All directed cycles in D^+ and D^- have odd length.

Suppose the contrary, and let $C = (x_1, x_2, \ldots, x_{2p})$ be a directed cycle, say in D^+ . For every $i = 1, \ldots, p$ let e_{2i-1}, e_{2i} be the two edges going out from x_{2i} , such that $e_{2i} = x_{2i}x_{2i+1}$, and call y_i the end vertex of e_{2i-1} . Then $y_i \in V \setminus V^+$, therefore there is at most one edge g_i going out from y_i . For every $i = 1, \ldots, p$, let f_{2i-1}, f_{2i} be the two edges going out from x_{2i-1} , such that $f_{2i-1} = x_{2i-1}x_{2i}$. Let $A = \{e_1, \ldots, e_{2p}\}$ and $B = \{f_1, \ldots, f_{2p}\} \cup \{g_1, \ldots, g_p\}$. Observe that A, B are disjoint and that B contains one edge of each directed P_3 of D that has an edge in A. Therefore A, B is a reducing pair, with $|B| \leq \frac{3}{2}|A|$, contradicting Claim 1.2. Thus Claim 1.4 holds.

Claim 1.5. D^+ and D^- have the same number of vertices, say this number is k, and $D \setminus (D^+ \cup D^-)$ is the union of k disjoint edges going from D^+ to D^- .

We shall prove that $V^0 = V(D) \setminus (V^- \cup V^+) = \emptyset$. Assume on the contrary that $V^0 \neq \emptyset$. By the connectivity of D, there is a vertex $y \in V^0$ adjacent to some vertex of $D^+ \cup D^-$. By symmetry, we may assume that yz is an edge for some $z \in V^-$. Let $C \subseteq D^-$ be the directed cycle containing z, let C have length $2\ell + 1$, with $\ell \ge 1$. We call $(\ell + 1)$ -set any subset $L \subset V(C)$ such that $V(C) \setminus L$ is a maximum independent set of C. Note that for any two vertices x, y of Cthere exists an $(\ell + 1)$ -set that contains both x, y.

If $d^-(y) \neq 0$, then let $e_0 = xy$, and let g_0 be an edge going into x if it exists. (Note that $y \notin V^-$ implies $x \notin V^-$.) Let $L \subset V(C)$ be an $(\ell + 1)$ -set of C not containing z and define:

$$B_1 = \{ f \in E(C) \mid f = ww' \text{ for some } w \in L \},\$$

$$A = \{ e_0 \} \cup (E(C) \setminus B_1) \cup \{ e \notin E(C) \mid e = vw \text{ for some } w \in L \},\$$

$$B_2 = \{ g \in (E(D) \setminus B_1) \mid g = uw \text{ such that } wv \in A \}.$$

Observe that A contains no directed P_3 and that every directed P_3 with one edge in A has its other edge in $B = B_1 \cup B_2$. So A, B is a reducing pair. Since $|A| = 2\ell + 2$, $|B_1| = \ell + 1$, and $|B_2| \le 2\ell + 2$, we have $|B| \le \frac{3}{2}|A|$, contradicting Claim 1.2.

If $d^-(y) = 0$, then let $L \subset V(C)$ be an $(\ell + 1)$ -set of C containing z, and define:

$$B_1 = \{f \in E(C) \mid f = ww' \text{ for some } w \in L\},\$$

$$A = (E(C) \setminus B_1) \cup \{e \notin E(C) \mid e = vw \text{ for some } w \in L\},\$$

$$B_2 = \{g \in (E(D) \setminus B_1) \mid g = uw \text{ such that } wv \in A\}.$$

Again, A and $B = B_1 \cup B_2$ form a reducing pair. We have $|A| = 2\ell + 1$, $|B_1| = \ell + 1$ and $|B_2| = 2\ell$ since no edge enters into y. Hence $|B| < \frac{3}{2}|A|$, contradicting Claim 1.2. Then Claim 1.5 follows from the second part of Claim 1.3.

Call M the (loopless) bipartite multigraph obtained by contracting of every directed cycle into one vertex.

Claim 1.6. *M* is a simple graph.

Suppose on the contrary that there are at least two edges from the cycle $C^+ \subseteq D^+$ to the cycle $C^- \subseteq D^-$. By Claim 1.5, C^+ is an odd cycle, thus there exist edges $ux, vy \in E(D)$ with $u, v \in V(C^+), x, y \in V(C^-)$ such that $(u, b_1, \ldots, b_{2q}, v)$ is a directed subpath of C^+ , and no vertex b_i has an edge to C^- (q = 0 means that uv is an edge of C^+).

Let C^- have length $2\ell + 1$ (it is an odd cycle by Claim 1.5). Obviously there exists an $(\ell + 1)$ -set $L \subset V(C^-)$ including x and excluding y. Define:

$$B_1 = \{ f \in E(C^-) \mid f = wz \text{ for some } w \in L \},\$$

$$A_0 = (E(C^-) \setminus B_1) \cup \{ e \notin E(C^-) \mid e = zw \text{ for some } w \in L \},\$$

 $B_2 = \{g \in (E(D) \setminus B_1) \mid g = wz \text{ such that } zw' \in A_0\}.$

Let $e'_0 = ub_1$, $f'_0 = vw$ where $w \in V(C^+)$, $g'_0 = vy$, and define:

$$\begin{array}{rcl} A'_0 &=& \{e'_0\} \cup \{e' \in E(D) \mid e' = b_{2i}z, \ 1 \le i \le q\}, \\ B'_1 &=& \{f'_0\} \cup \{f' \in E(D) \mid f' = b_{2i-1}z, \ 1 \le i \le q\}, \\ B'_2 &=& \{g' \in (E(D) \setminus B'_1) \mid g' = zw' \text{ such that } bz \in A'_0\} \setminus \{g'_0\}. \end{array}$$

Observe that the set $A = A_0 \cup A'_0$ contains no directed P_3 , and every directed P_3 with an edge in A has its other edge in $B = B_1 \cup B_2 \cup B'_1 \cup B'_2$ from D. Hence A, B form a reducing pair. We have $|A_0| = 2\ell + 1$, $|B_1| = \ell + 1$, $|B_2| = 2\ell + 1$, $|A'_0| = 2q + 1$, $|B'_1| = 2q + 1$, $|B'_2| = q$, so $|A| = 2(\ell + q + 1)$ and $|B| = 3(\ell + q + 1) = \frac{3}{2}|A|$, contradicting Claim 1.2. Thus Claim 1.6 holds.

Because every vertex of the contraction graph M has degree at least three, M has a cycle. To conclude the proof of the theorem we show that this leads to a contradiction.

Consider a shortest cycle $\gamma \subset M$, and let $\gamma = (C_1^+, C_1^-, C_2^+, C_2^-, \dots, C_p^+, C_p^-)$, where, for each $i \in \{1, \dots, p\}, C_i^+ \subseteq D^+$ and $C_i^- \subseteq D^-$ are cycles of D of odd length. The edges of γ correspond to a matching of D from the set $\cup_{i=1}^q \{u^i, v^i\}$, to the set $\cup_{i=1}^q \{x^i, y^i\}$, where $u^i, v^i \in C_i^+$ and $x^i, y^i \in C_i^-$. Furthermore, by Claim 1.6 and since γ has no chords in M, no more edges of D are induced between these cycles. We may assume, so we do, that $(u^i, b_1^i, \dots, b_{2q_i-1}^i, v^i)$, where $q_i \geq 1$, is a directed subpath of C_i^+ .

Let $2\ell_i + 1$ be the length of C_i^- . For every $i = 1, \ldots, p$ select an $(\ell_i + 1)$ -set $L_i \subset V(C_i^-)$ of C^- such that $x^i, y^i \in L_i$, and define the following sets:

 $B_{i}^{1} = \{f \in E(C_{i}^{-}) \mid f = wz \text{ for some } w \in L_{i}\},\$ $A_{i}^{1} = (E(C_{i}^{-}) \setminus B_{i}^{1}) \cup \{e \notin E(C_{i}^{-}) \mid e = zw \text{ for some } w \in L_{i}\},\$ $B_{i}^{2} = \{g \in (E(D) \setminus B_{i}^{1}) \mid g = uz \text{ such that } zw \in A_{i}^{1}\}.$

Let $A^1 = \bigcup_{i=1}^p A_i^1$ and $B^1 = \bigcup_{i=1}^p (B_i^1 \cup B_i^2)$. We have $|A^1| = \sum_{i=1}^p |A_i^1| = \sum_{i=1}^p (2\ell_i + 1)$, and because $|B_i^1| = \ell_i + 1, |B_i^2| = 2\ell_i + 1$, we obtain $|B^1| = \sum_{i=1}^p (3\ell_i + 2)$. For every $i = 1, \dots, p$, let $e_i = u^i b_1^i$, $f_i = b_{2q_i-1}^i v^i$, and define sets: $A_i^2 = \{e_i\} \cup \{e \in (E(D) \mid e = b_{2j}w, \ 1 \le j \le q_i - 1\},$ $B_i^3 = \{f \in E(D) \mid f = b_{2j-1}w, \ 1 \le j \le q_i\} \setminus \{f_i\},$ $B_i^4 = \{g \in (E(D) \setminus B_i^3) \mid g = wz \text{ such that } bw \in A_i^2\} \setminus \{f_i\}.$ Let $A^2 = \bigcup_{i=1}^p A_i^2$ and $B^3 = \bigcup_{i=1}^p (B_i^3 \cup B_i^4)$. We have $|A^2| = \sum_{i=1}^p |A_i^2| = \sum_{i=1}^p (2q_i - 1)$, and because $|B_i^3| = 2q_i - 1$ and $|B_i^4| = q_i - 1$, we obtain $|B^3| = \sum_{i=1}^p (3q_i - 2)$. Observe that the sets $A = A^1 \cup A^2$ and $B = B^1 \cup B^3$ form a reducing pair. Furthermore, $|A| = \sum_{i=1}^p (2\ell_i + 1 + 2q_i - 1) = 2\sum_{i=1}^p (\ell_i + q_i)$ and $|B| = \sum_{i=1}^p (3\ell_i + 2 + 3q_i - 2) = 3\sum_{i=1}^p (\ell_i + q_i) = \frac{3}{2}|A|$, contradicting Claim 1.2. This concludes the proof of the theorem.

The proof of the theorem can be formulated as an algorithm which, given any digraph $D \in D(1,1)$ with m edges and at most t disjoint directed triangles, constructs a directed cut K of size at least (2m-t)/5. We sketch such an algorithm here. Start from $K := \emptyset$. Then apply the following general step. Find the subgraphs D^+ and D^- . If there is a directed triangle that is not included in D^+ or D^- , with the notation of Claim 1.1, then set $K := K \cup \{xy\}$ and iterate with the subgraph $D \setminus \{xy, yz, zx\}$. (When iterating, the subgraphs $D^+, D^$ must be updated.) If there is no such directed triangle, then either D violates one of Claims 1.3-1.6, or D satisfies the conditions described after the proof of Claim 1.6; and in either case, the proof of the theorem shows how to find a reducing pair (A, B). Then set $K := K \cup A$ and iterate the general step with the subgraph $D \setminus (A \cup B)$. The algorithm terminates when D becomes edgeless. Then at termination K is a directed cut of size at least (2m-t)/5. It is easy to see that all the operations (updating D^+ and D^- , finding a directed triangle, checking whether D violates one of the claims, determining the structure described after the proof of Claim 1.6) can be done in polynomial time, and there are at most m iterations. Thus we obtain:

Corollary 2. There is a polynomial time algorithm which, given any digraph $D \in D(1,1)$ with m edges and at most t disjoint directed triangles, finds a directed cut in D of size at least (2m-t)/5.

Corollary 3. If $D \in D(1,1)$ has m edges, then it contains a directed cut of size at least m/3. Moreover D has no directed cut of size larger than m/3 if and only if D is the union of disjoint directed triangles.

Proof. The number of pairwise disjoint directed triangles satisfies $t \le m/3$, with equality if and only if D is a union of disjoint directed triangles. Now the claim follows by Theorem 1, because $(2m - t)/5 \ge (2m - m/3)/5 = m/3$.

Corollary 4. If $D \in D(1,1)$ has m edges and no directed triangle, then it contains a directed cut of size at least 2m/5.

Results by Bondy and Locke [4] on the bipartite density of (undirected) subcubic graphs are reminescent of our investigations concerning D(1,1). They proved in [4] that a triangle-free subcubic graph has a bipartite subgraph of size at least 4m/5. Observe that any triangle-free digraph of maximum degree at most three belongs to D(1,1), and it is obtained from a subcubic graph by orienting its edges. Hence their result implies that such a D has a directed cut of size at least 2m/5, the half of 4m/5. Corollary 4 shows that this bound

is valid for the much larger class of digraphs in D(1,1) containing no directed triangle.

Now we show that the lower bound m/3 in Corollary 3 can be surpassed for connected digraphs of D(1, 1).

Theorem 5. If $D \in D(1,1)$ is a connected digraph with m edges, and D is not a triangle, then it contains a directed cut of size at least 7m/20.

Proof. The proof works by induction on m. Let t be the maximum number of pairwise disjoint directed triangles of D. By the hypothesis, we have t = 0 if $m \leq 3$ and $t \leq 1$ if m = 4, 5, 6. Thus by Theorem 1, there is a cut of size at least 1, 1, 2, 2, 2, 3, respectively, for $m = 1, \ldots, 6$, which matches the corresponding value of $\lceil 7m/20 \rceil$. Now let $m \geq 7$, and assume that the claim is true for connected graphs with strictly less than m edges. Observe that any t disjoint directed triangles of D have a total of 3t edges, furthermore, by the connectivity of D, there are at least t-1 more edges between these triangles. Hence we have $m \geq 4t-1$.

If m > 4t - 1, or equivalently, if $t \le m/4$, then by Theorem 1, D has a cut of size at least $(2m - t)/5 \ge (2m - m/4)/5 = 7m/20$ edges as stated.

Assume now that m = 4t - 1. So D consists of t disjoint directed triangles connected by t - 1 edges in a tree-like manner. Since we cannot have t = 1and m = 3, we have $t \ge 2$. So there is a directed triangle T = (x, y, z) that is adjacent to exactly one edge, say xx', which is adjacent to another directed triangle T' = (x', y', z'). (The symmetric argument applies if the orientation of the edge between T and T' is x'x.) Removing from D the vertices and edges of T together with the two edges xx', x'y', we obtain a connected digraph D' with $m' = m - 5 \ge 2$ edges. By the induction hypothesis, D' has a cut H' of size at least 7m'/20 = (7m - 35)/20 > 7m/20 - 2 edges. Clearly $H' \cup \{xx', yz\}$ has no directed P_3 , which yields a cut of size at least 7m/20 in D.

Just like with Theorem 1, the proof of Theorem 5 can be formulated easily as a polynomial time algorithm (we omit the details). So we have:

Corollary 6. There is a polynomial time algorithm which, given any digraph $D \in D(1,1)$ with m edges, such that no component of D is a directed triangle, finds a directed cut in D of size at least 7m/20.

3 Decompositions of D(k, k)

The problem of covering the edges of a digraph with cuts was proposed in [1]. Upper bounds were given for digraphs in $D(k, \ell)$, and the only exact value was determined for $k = \ell = 1$.

Theorem 7 ([1]). The edge set of any digraph $D \in D(1,1)$ can be decomposed into at most three cuts.

Theorem 8. For integers $p_1, p_2 \ge 0$, the edge set of every digraph $D \in D(p_1 + p_2, p_1 + p_2)$ can be decomposed into two subgraphs $D_1 \in D(p_1, p_1)$ and $D_2 \in D(p_2, p_2)$.

Proof. Since D is in $D(p_1 + p_2, p_1 + p_2)$, its vertex set V(D) can be partitioned into two sets X, Y such that every vertex $x \in X$ satisfies $d^-(x) \leq p_1 + p_2$ and every vertex $y \in Y$ satisfies $d^+(y) \leq p_1 + p_2$. Consider the set of edges $B = \{yx \in E(D) | y \in Y, x \in X\}$. By the definition of X and Y, in the bipartite graph (X, Y; B) every vertex has degree at most $p_1 + p_2$. By a classical corollary of the König-Hall theorem (see e.g., [7, Prop. 5.3.1]), the edges of B can be colored with $p_1 + p_2$ colors so that any two adjacent edges have different colors. Let B_1 be the set of edges of B with the first p_1 colors and B_2 be the set of edges of B with the remaining colors.

For every vertex $x \in X$, the set $E^{-}(x)$ of edges with end x has size at most $p_1 + p_2$, so it can be partitioned into two sets $E_1(x)$ and $E_2(x)$ such that, for $j = 1, 2, |E_j(x)| \leq p_j$ and $E^{-}(x) \cap B_j \subseteq E_j(x)$. Likewise, for every vertex $y \in Y$, the set $E^{+}(y)$ of edges with origin y has size at most $p_1 + p_2$, so it can be partitioned into two sets $E_1(y)$ and $E_2(y)$ such that, for $j = 1, 2, |E_j(y)| \leq p_j$ and $E^{+}(y) \cap B_j \subseteq E_j(y)$.

Finally let the set $\{xy \in E(D) | x \in X, y \in Y\}$ be partitioned arbitrarily into two sets F_1, F_2 . Now, for j = 1, 2, let D_j be the subgraph of D whose edge set is $B_j \cup F_j \cup \bigcup_{x \in V} E_j(x)$. The definition of these sets implies that each edge of D lies in exactly one of D_1, D_2 and that $D_j \in D(p_j, p_j)$ for j = 1, 2. More precisely, for j = 1, 2, in D_j every vertex $x \in X$ satisfies $d^-(x) \leq p_j$ and every vertex $y \in Y$ satisfies $d^+(y) \leq p_j$.

Corollary 9. The edges of every digraph $D \in D(2,2)$ can be decomposed into two subgraphs $D_1, D_2 \in D(1,1)$.

From Corollary 9 and Theorem 7 it follows that every digraph $D \in D(2,2)$ can be covered with six directed cuts. If there was a decomposition of D into a cut and a digraph in D(1,1), then D would have a cut cover only with four cuts, by Theorem 7 again. Our next example shows that such a decomposition is not always possible.

Example 2. Take two disjoint copies of a regular tournament on five vertices, G_1, G_2 , and include all 25 edges directed from G_1 to G_2 . Thus we obtain a digraph $H \in D(2, 2)$. Assume that $K \subset E(H)$ is a cut such that $H' = H \setminus K$ is in D(1, 1). The regular tournament has no cut with more than three edges, hence G_1 has a vertex v_0 such that every edge going into v_0 is in $E(H) \setminus K$ and at least one edge going out of v_0 is in $E(H) \setminus K$. Thus $d_{H'}^-(v_0) = 2$, which implies that $v_0 z \in K$ for all $z \in V(G_2)$ in order to obtain $d_{H'}^+(v_0) \leq 1$. Then it follows that no edge of G_2 belongs to K, thus $d_{H'}^+(z) = 2$ and $d_{H'}^-(z) \geq 2$ for all $z \in V(G_2)$, a contradiction.

How large a subgraph belonging to D(1,1) can be found in a digraph $D \in D(2,2)$? Corollary 9 implies that D with m edges contains a subgraph in D(1,1)

with at least m/2 edges. A larger bound will follow from our more general result.

Theorem 10. Every digraph $D \in D(k, k)$ with m edges has a subgraph belonging to D(k-1, k-1) with at least (2k-1)m/(2k+1) edges.

Proof. Let $W = \{v \in V(D) | d_D^-(v) \leq k\}$ and $B = \{v \in V(D) | d_D^+(v) \leq k\}$. Because $D \in D(k, k)$, we have $V(D) = W \cup B$. We say that $v \in W$ is white, and $v \in B$ is black; note that a vertex may have both colors. An edge $xy \in E(D)$ is called a black tail arrow if $x \in B$, and it is called a white head arrow if $y \in W$. Note that an edge can be both a black tail and a white head arrow. Observe the symmetry of the colors with respect to reversing all arrows in D. Due to this symmetry, if a property is verified for white vertices, then the analogous property is true for black vertices with directions reversed.

Let $R \subset E(D)$ be a set of edges such that (a) the graph $D' = D \setminus R$ is in D(k-1, k-1), (b) R is minimum among all sets with property (a), and (c) R has the maximum number of black tail arrows and white head arrows (each arrow counted once) among all sets that satisfy (a) and (b). Clearly such a set R exists.

For each edge $e = xy \in R$, we define a *critical vertex* of e as follows:

x is a critical vertex for $e = xy \in R$ if $d_{D'}^+(x) = k - 1$ and $d_{D'}^-(x) \ge k$;

y is a critical vertex for $e = xy \in R$ if $d_{D'}^{-}(y) = k - 1$ and $d_{D'}^{+}(y) \ge k$.

The minimality of R means that at least one of x, y is critical for each edge e = xy of R. Note that both x, y may be critical for e. For each $e \in R$, let $\operatorname{Crit}(e) \subseteq \{x, y\}$ be the set of critical vertices of e. For any subset $X \subseteq R$, define $\operatorname{Crit}(X) = \bigcup_{e \in X} \operatorname{Crit}(e)$. From here on, the word critical vertex refers to elements of $\operatorname{Crit}(R)$. All critical vertices are in the set $\{v \in V(D) | d_D^-(v), d_D^+(v) \ge k\}$, however not every vertex in that set is critical for some edge of R. The main point of the proof is to establish that:

Claim 10.1. $|Crit(R)| \ge |R|$.

Assume we already know that $|\operatorname{Crit}(R)| \geq |R|$. Then the definition of critical vertices implies that, for every $v \in \operatorname{Crit}(R)$, there are at least 2k - 1 edges not in R and incident with v. Thus for the size of D we have the bound $m \geq |R| + (2k-1)|R|/2$, hence $|R| \leq 2m/(2k+1)$. So D' has at least $m - |R| \geq (2k-1)m/(2k+1)$ edges, and the theorem follows. Therefore the rest of the proof consists in proving Claim 10.1.

The definition of critical vertices implies easily the following two claims, whose proof is omitted.

Claim 10.2. If $x \in B$ and $d_R^+(x) \ge 2$, then $x \notin Crit(R)$. By symmetry, If $x \in W$ and $d_R^-(x) \ge 2$, then $x \notin Crit(R)$.

Claim 10.3. If $e = xy \in R$, $y \in B$, and $yz \in R$, then $Crit(e) = \{x\}$. By symmetry, if $e = xy \in R$, $x \in W$, and $ux \in R$, then $Crit(e) = \{y\}$.

Now we examine the subgraph formed by R. Let $A \subseteq R$ be any connected component of R (we use A and R to denote the digraphs defined by the edges in A and R, respectively). Note that $\operatorname{Crit}(R) = \operatorname{Crit}(A) \cup \operatorname{Crit}(R \setminus A)$.

Claim 10.4. If A has no cycle, then $|Crit(A)| \ge |V(A)| - 1 = |A|$.

In this case A is a tree with |A| + 1 vertices. We show that at most one non critical vertex may exist in A. Suppose on the contrary that u, v are two non-critical vertices in A, and let $P = (u, \ldots, v)$ be the (unique) shortest chain between them in A. Observe that P has length at least 2 (for otherwise its unique edge uv would satisfy $\operatorname{Crit}(uv) = \emptyset$), and that the inclusion into D' of the two edges of P incident to u and v does not increase their corresponding indegree or outdegree above k - 1.

For every white vertex w of $V(P) \setminus \{u, v\}$ select a white head arrow xw, and for every black vertex z of $V(P) \setminus \{u, v\}$ select a black tail arrow zx (for twocolored vertices take one such arrow arbitrarily). Let F be the set of selected arrows. So $|F| \leq |P| - 1$. Define $R^* = (R \setminus P) \cup F$. The graph $D^* = D \setminus R^*$ belongs to D(k-1, k-1), because the outdegree of every black vertex of $V(P) \setminus \{u, v\}$, and the indegree of every white vertex of $V(P) \setminus \{u, v\}$ is at most k-1, furthermore the corresponding degrees of u and v do not increase above k-1. The set R^* satisfies $|R^*| \leq |R| - 1$, contradicting the minimality of R. Thus Ahas at most one non-critical vertex, and Claim 10.4 holds.

Now we consider an arbitrary cycle C in R (if any).

Claim 10.5. *C* has no edge e = xy with $x \in W \setminus B$ and $y \in B \setminus W$.

Suppose that there is such an edge e = xy. Note that e is neither a white head arrow nor a black tail arrow in C. Hence C has at most |C| - 1 white head and black tail arrows. For every white vertex $w \in V(C)$ select a white head arrow zw, and for every black vertex $v \in V(C)$ select a black tail arrow vu (for two-colored vertices select one arrow arbitrarily). Let F be the set of |C| selected edges, and define $R^* = (R \setminus C) \cup F$. The set R^* satisfies $|R^*| \leq |R|$, and contains more white head and black tail arrows than R. Furthermore, the graph $D^* = D \setminus R^*$ belongs to D(k - 1, k - 1), because the outdegree of every black vertex of C, and the indegree of every white vertex of C is at most k - 1. This contradicts the choice of R. Thus Claim 10.5 holds.

Claim 10.6. Let u, x, y, v be four consecutive vertices of C.

- (1) If $xu, xy, vy \in C$ and x is black, then y is not white.
- (2) If $ux, xy, vy \in C$, then either x or y is not white.
- (3) If $ux, xy, yv \in C$ and y is black, then x is not white.

Suppose on the contrary that any of (1), (2), (3) fails. Then, in either case, the edge e = xy satisfies $Crit(e) = \emptyset$, which contradicts the minimality of R. Thus Claim 10.6 holds.

Claim 10.7. *C* is a directed cycle and it is monochromatic, i.e., its vertices are either all in $W \setminus B$ or all in $B \setminus W$.

Suppose first that C is not a directed cycle, and consider a longest directed subpath (x_1, \ldots, x_q) of C, where $q \ge 2$.

Suppose that q = 2, i.e., the directions of the edges alternate on C. Let $z_1w_1, z_1w_2, z_2w_2 \in C$. If $z_1 \in W \setminus B$, then by Claim 10.5 we have $w_1, w_2 \in W$. If $z_1 \in B$, then Claim 10.6 (1) implies that $w_2 \in B \setminus W$, and $z_2 \in B$ follows by Claim 10.5. Thus we obtain that either $w_1, z_1, w_2 \in W$ or (symmetrically) $z_1, w_2, z_2 \in B$. We show a contradiction in the first case, then, by symmetry, the second case is impossible as well. So assume that $w_1, z_1, w_2 \in W$ and set $e_i = z_1w_i, i = 1, 2$. Select an arbitrary white head arrow $f = xz_1 \in E(D)$. The set $R^* = (R \setminus \{e_1, e_2\}) \cup \{f\}$ satisfies $|R^*| \leq |R| - 1$, and the graph $D^* = D \setminus R^*$ belongs to D(k - 1, k - 1), because $d_{D^*}^-(w_i) = d_{D'}^-(w_i) \leq k - 1$ for i = 1, 2, and $d_{D^*}^-(z_1) \leq k - 1$. This contradicts the minimality of R. Therefore $q \geq 3$.

By Claim 10.6 (2), either x_q or x_{q-1} is not white on the directed path (x_1, x_2, \ldots, x_q) , for $q \ge 3$. If $x_q \in B \setminus W$, then $x_{q-1} \in B$ by Claim 10.5. Thus that in each case x_{q-1} is black.

Suppose that q = 3. Let $e_1 = x_1x_2$, $e_2 = x_1y_2 \in C$, with $e_1 \neq e_2$, and let y_3 be the second neighbor of y_2 on C different from x_1 .

Assume first that $y_2y_3 \in E(D)$. Then, by the argument above, x_2 and y_2 are both black. Observe that $x_1 \in W \setminus B$, since otherwise the edges e_1 and e_2 have no critical vertices. Now select an arbitrary white head arrow $f = zx_1 \in E(D)$. The set $R^* = (R \setminus \{e_1, e_2\}) \cup \{f\}$ satisfies $|R^*| = |R| - 1$, and the graph $D^* = D \setminus R^*$ belongs to D(k - 1, k - 1), contradicting the minimality of R.

Assume now that $y_3y_2 \in E(D)$ (where y_3y_2 might coincide with x_2x_3 , if C is a triangle). As before, we have $x_2 \in B$ and $x_1 \in W \setminus B$. Then, by Claim 10.5, $y_2 \in W$. Selecting a white head arrow f at x_1 and defining the set $R^* = (R \setminus \{e_1, e_2\}) \cup \{f\}$ we obtain a contradiction in the same way as before. Therefore $q \geq 4$.

We already know that x_{q-1} is black. Hence $x_{q-2} \in B \setminus W$ by Claim 10.6 (3). Applying Claim 10.6 (3) repeatedly, we obtain that x_{q-2}, \ldots, x_2 are in $B \setminus W$. Then we have $x_1 \in B$ by Claim 10.5. Hence the edge x_1x_2 has no critical vertex, because $d_{D'}^+(x_1) \leq k-2$ and $d_{D'}^+(x_2) \leq k-1$, contradicting the minimality of R. So we have established that C is a directed cycle.

Now assume without loss of generality that some vertex y of C is black. By Claim 10.6 (3), the predecessor $x \in V(C)$ of y is not white, i.e., it is in $B \setminus W$. Applying Claim 10.6 (3) repeatedly we obtain that every vertex of C is in $B \setminus W$. So C is monochromatic. Thus Claim 10.7 holds.

Claim 10.8. If a component A of R contains a cycle, then A is unicyclic and |Crit(A)| = |V(A)| = |A|.

Let A contain a cycle C. By Claim 10.7 and by symmetry, C is a black directed cycle.

For every edge $e = xy \in C$, by Claim 10.3, we have $x \in \operatorname{Crit}(e)$, and by Claim 10.2, we have $d_R^+(x) = 1$. Let A_0 be a subgraph of A that is maximal with the following property: A_0 contains C, for every $x \in V(A_0)$ there is a directed path in A_0 from x to some vertex of C, and every vertex $x \in V(A_0)$ is

black and satisfies $d_R^+(x) = 1$. Let us prove that $A_0 = A$. Note that A_0 exists, because C itself satisfies all the required properties.

Suppose that $A_0 \neq A$. Then, since A is connected, there is a vertex $x \in V(A) \setminus V(A_0)$ that is adjacent to some $y \in V(A_0)$. Because $d_R^+(y) = 1$ and $d_{A_0}^+(y) = 1$, we have $e = xy \in A$. Since y is black, Claim 10.3 implies Crit $(e) = \{x\}$. Observe that A_0 contains a directed P_3 from y containing black vertices. Hence by Claim 10.6 (3), we have $y \in B \setminus W$. If $x \in W \setminus B$, then let $f = ux \in R$ be any white head arrow. Define $R^* = (R \setminus \{e\}) \cup \{f\}$. The digraph $D^* = D \setminus R^*$ belongs to D(k-1, k-1), because $d_{D^*}^+(y) = d_{D'}^+(y) \leq k-1$, and $d_{D^*}^-(x) \leq k-1$. This contradicts the choice of R. So x is black. Because x is black and $x \in \operatorname{Crit}(e)$, Claim 10.2 implies $d_R^+(x) = 1$. Hence one could include x to A_0 , contradicting the maximality of A_0 . Therefore $A_0 = A$.

Since $A = A_0$, C is the only cycle in A and every vertex x of A is black and satisfies $d_R^+(x) = 1$. Claim 10.3 implies that every vertex of A is a critical vertex of its outgoing edge. So $|\operatorname{Crit}(A)| = |V(A)| = |A|$, and Claim 10.8 holds.

Claims 10.4 and 10.8 show that $|\operatorname{Crit}(A)| \geq |A|$ is true for every connected component A. If A_1, \ldots, A_t are the components of R, we have clearly $\operatorname{Crit}(R) = \operatorname{Crit}(A_1) \cup \cdots \cup \operatorname{Crit}(A_t)$. Thus we obtain $|\operatorname{Crit}(R)| \geq |R|$, which proves Claim 10.1. This concludes the proof of the theorem.

The regular tournament on 2k + 1 vertices has indegree equal to outdegree for every vertex, hence it is in D(k, k). To obtain a subgraph belonging to D(k-1, k-1) one has to remove at least k+1 from its $m = \binom{2k+1}{2}$ edges. This shows that the tournament has no subgraph in D(k-1, k-1) containing more than $\binom{2k+1}{2} - k + 1 = m - (1+1/k)m/(2k+1) = (2k-1/k)m/(2k+1)$ edges.

Corollary 11. Every digraph $D \in D(2,2)$ with m edges contains a subgraph belonging to D(1,1) with at least 3m/5 edges.

We note that for k = 1, Theorem 10 yields another proof that every digraph $D \in D(1,1)$ with m edges contains a directed cut of size at least m/3 (cf. Corollary 3).

4 Cuts in D(k,k)

In [1] it was observed that the k-regular orientation of the complete graph on 2k + 1 vertices has no directed cut of size more than $(\frac{1}{4} + \frac{1}{8k+4})\binom{2k+1}{2}$. Consequently, in a digraph $D \in D(k, k)$ with m edges one cannot guarantee a directed cut of size larger than $(\frac{1}{4} + \frac{1}{8k+4})m$. It was proved in [1] that every digraph with outdegree at most k does contain a directed cut of that size. Using the same methods we show that it is also true for the acyclic members of D(k, k).

The basic tool is a lemma in [10] that is proved there by elementary counting.

Lemma 1 ([10]). If a γ -colorable graph G has m edges, then it has a bipartite partial graph with at least $(\lfloor \gamma^2/4 \rfloor/{\gamma \choose 2})m$ edges.

Theorem 12. If $D \in D(k,k)$ is acyclic and has m edges, then D contains a directed cut of size at least $(\frac{1}{4} + \frac{1}{8k+4})m$.

Proof. Let D^+ be the subgraph of D induced by the set $X = \{v \in V(D) | d^+(v) \le k\}$ and let D^- be the subgraph of D induced by $V(D) \setminus X$. Because D is acyclic, every subgraph of D^+ has a source, thus its underlying graph G^+ is k-degenerate. Similarly, every subgraph of D^- has a sink, thus its underlying graph G^- is k-degenerate. Consequently, both graphs G^+ and G^- are (k + 1)-colorable, therefore the underlying graph G of D is (2k + 2)-colorable.

Applying Lemma 1 with $\gamma = 2k+2$, we obtain a bipartite partial graph of G with $\frac{k+1}{2k+1}m$ edges. In D at least half of the edges of that bipartite graph form a directed cut of size at least $\frac{k+1}{4k+2}m = (\frac{1}{4} + \frac{1}{8k+4})m$.

We do not know whether Theorem 12 remains true for all digraphs in D(k, k), and for every k. The coefficients are 1/3, 3/10 and 2/7 for k = 1, 2, and 3, respectively. By Theorem 7, a digraph $D \in D(1, 1)$ of size m has a cut with at least m/3 edges. The theorem below answers the question affirmatively for D(2, 2).

Theorem 13. Every digraph $D \in D(2,2)$ with m edges has a directed cut of size at least 3m/10.

Proof. We prove the theorem by induction on m. For m = 1 the theorem is true. Now suppose that $m \ge 2$ and that the theorem holds for every digraph with at most m-1 edges. Since D is in D(2,2), its vertex set can be partitioned into two sets X, Y such that every vertex $x \in X$ satisfies $d^{-}(x) \le 2$ and every vertex $y \in Y$ satisfies $d^{+}(y) \le 2$. Consider the set of edges $F = \{xy \in E | x \in X, y \in Y\}$.

First suppose that the underlying bipartite graph (X, Y; F) contains a cycle. Let C be any such cycle, say with length 2k, let X_C and Y_C be the set of vertices of C that lie in X and Y respectively, and let F_C be the set of edges of C. So $|F_C| = 2k$. Let E_C be the set of edges such that either their end is in X_C or their origin is in Y_C . By the definition of X, Y and the fact that D is in D(2,2), we have $|E_C| \leq 4k$. Consider the digraph $D' = D \setminus (E_C \cup F_C)$. Clearly, $D' \in D(2,2)$, and the number m' of edges of D' satisfies $m' \geq m - 6k$. By the induction hypothesis, D' has a directed cut of size at least 3m'/10. If the edges of F_C are added to such a directed cut, we obtain a directed cut of D, because D' does not contain any edge of E_C . This directed cut of D has size at least $3m'/10 + |F_C| \geq 3(m - 6k)/10 + 2k \geq 3m/10$. So the theorem holds for D.

Now suppose that the bipartite graph (X, Y; F) does not contain any cycle. Thus $|F| \leq n-1$, where *n* is the number of vertices of *D*. By the definition of *X* and *Y*, we have $m \leq 2|X| + 2|Y| + |F| \leq 2n + n - 1 = 3n - 1$, which implies that *D* has a vertex *v* of degree at most 5. Actually the same argument can be repeated with $D \setminus \{v\}$, and so on. Thus the underlying graph of *D* is 5-degenerate and therefore has chromatic number at most 6. Applying Lemma 1 with $\gamma = 6$ we obtain a bipartite subgraph with 3/5 edges, thus *D* has a directed cut of size at least 3m/10.

5 Problems

Let c_{max} be the ratio of the maximum directed cut size to the edge count m of a digraph. For connected digraphs of D(1,1), Theorem 5 improves the basic estimation $c_{max} \ge 1/3$ to $c_{max} \ge 7/20$ provided m > 3. On the other hand, Example 1 before Theorem 1 shows infinitely many connected digraphs of D(1,1) with $c_{max} < 3/8$. We conjecture that the bound $c_{max} \ge 7/20$ can be improved to 3/8 in the limit in the following sense.

Problem 1. For every $\varepsilon > 0$, there exists a constant m_{ε} such that $c_{max} > 3/8 - \varepsilon$ holds for every connected digraph of D(1,1) with $m > m_{\varepsilon}$ edges.

At some point of the investigation in D(1,1) we observed that the presence of source or sink vertices of the digraph increases the size of a maximum directed cut. Corollary 4 might have the following sharpening.

Problem 2. If a connected digraph $D \in D(1,1)$ with m edges contains no directed triangle and has s vertices with indegree or outdegree zero, then D has a directed cut of size at least (2m + s)/5.

Bondy and Locke [4] proved that a triangle-free subcubic graph has a cut (a bipartite subgraph) of size at least 4m/5. A characterization of all extremal graphs for that bound was given by Xu and Yu in [11]. The problem of characterizing the extremal graphs for the bound of Corollary 4 remains open:

Problem 3. Determine the list of all digraphs $D \in D(1,1)$ of size m that contain no directed triangle and have no directed cut with more than 2m/5 edges.

Bondy and Locke's result in [4] consists of a polynomial time algorithm that finds a cut with at least 4m/5 edges in a triangle-free subcubic graph. It is known that finding a maximum cut is NP-hard even in the restricted family of triangle-free cubic graphs (see Yannakakis [12]). Even the approximation of the max cut problem in cubic graphs within the ratio of 0.997 is NP-hard (see Berman and Karpinski [3]). On the other hand, Halperin, Livnat and Zwick [9] give a polynomial time approximation algorithm with ratio 0.9326.

Concerning digraphs in D(1,1), Corollary 6 gives a polynomial time algorithm that produces a cut of size at least 7m/20 in every digraph in D(1,1) of which no component is a directed triangle; and so this is an approximation algorithm with ratio 0.35. Can a better ratio be obtained? Actually, as far as we know, none of the known results implies that computing the exact value of a maximum directed cut is NP-hard in D(1,1). So we ask:

Problem 4. What is the complexity status of computing the size of a maximum directed cut in a digraph of D(1,1)? If it is NP-hard, what is the best value of ε for which there is a polynomial time approximation algorithm with ratio $1 - \varepsilon$ for this problem?

The same problem can be posed for digraphs of D(1,1) with no directed triangle, or with no triangle at all.

How large a subgraph belonging to D(1,1) can be found in a digraph $D \in D(2,2)$? Corollary 11 says that D with m edges contains a subgraph in D(1,1) with at least 3m/5 edges. This lower bound is probably not sharp.

Problem 5. Determine the largest constant λ such that in every digraph $D \in D(2,2)$ with m edges there exists a subgraph $D' \in D(1,1)$ of size at least λm .

If D is the regular tournament on five vertices, then $D \in D(2,2)$ and one needs to remove at least three edges to obtain a subgraph $D' \in D(1,1)$. This shows that in the problem above $\lambda \leq 7/10$.

From a result in [1] it follows that the edges of every graph $D \in D(2,2)$ can be decomposed into at most five directed cuts. Furthermore, four cuts are sufficient if D is acyclic. The regular tournament on five vertices shows that four cuts might be necessary. Indeed, it has 10 edges, and the size of a directed cut is at most 3. No example has been found to show that five directed cuts are necessary.

Problem 6. The edges of every digraph $D \in D(2,2)$ can be decomposed into at most four directed cuts.

Several problems remain open in D(2,2) pertaining to the ratio c_{max} .

Problem 7. If $D \in D(2,2)$ has m edges and contains no copy of the regular tournament on five vertices, then D has a directed cut of size at least m/3.

We do not know whether Theorem 12 pertaining to acyclic digraphs remains true for all digraphs in D(k, k), and for every k. The coefficients $\frac{1}{4} + \frac{1}{8k+4}$ are equal to 1/3, 3/10, and 2/7 for k = 1, 2, and 3, respectively. By Corollary 3, and by Theorem 13, a digraph D of size m has a cut with m/3 and 3m/10 edges, respectively for $D \in D(1, 1)$ and $D \in D(2, 2)$. The next case k = 3 is proposed here as a question. It is quite possible that the answer is negative. Even if it is not the case we conjecture that Theorem 12 does not extend for every k.

Problem 8. Is it true that every digraph of D(3,3) with m edges contains a directed cut of size at least 2m/7?

Digraphs with maximum outdegree k satisfy $c_{max} \geq \frac{1}{4} + \frac{1}{8k+4}$, and this is the best bound, as shown in [1]. It is worth noting that the same bound was obtained here in Theorem 12 for acyclic members of D(k, k). Furthermore, the regular tournament on 2k+1 vertices is an example of a digraph with no directed cut larger than $(\frac{1}{4} + \frac{1}{8k+4})m$. We believe that in the larger family D(k, k) there are more examples showing that this bound cannot be achieved, provided k is large enough.

Problem 9. There exists a k_0 such that for every $k \ge k_0$ there are digraphs in D(k,k) with $c_{max} < \frac{1}{4} + \frac{1}{8k+4}$.

In Theorem 10 we are dealing with the largest subgraph of $D \in D(k, k)$ that belongs to the "lower" class D(k-1, k-1). This leads naturally to the investigation of the minimum sets $R \subset E(D)$ to be removed from D in order to lower its class. The proof of Theorem 10 suggests that such minimum sets considered as digraphs have a particular structure reminiscent of forests. Repeating the procedure, one obtains a decomposition of the original digraph $D \in D(k, k)$ into at most k of these structures.

Practical applications motivate the study of decompositions of digraphs into directed stars (see [5]). The *directed star arboricity* (dst) introduced in [8] is defined as the minimum number of outstar forests (also called galaxy) the edge set of a digraph can be partitioned. For instance it is proved in [2] that a digraph D with indegree at most k has a decomposition into k outforests plus one galaxy. This result implies $dst(D) \leq 2k+1$, and it is conjectured in [2] that 2k is the tight bound, for $k \geq 2$.

As a general problem we propose here a similar decomposition theory of the digraphs of D(k, k) into appropriate forest-like structures.

Acknowledgment

The first author's participation in the present research was supported by the Professional Development Assignment obtained from The University of Memphis and by the funding of the Université de Grenoble 1, Joseph Fourier. He is grateful for the help and the hospitality of all colleagues at the Laboratoire Leibniz - IMAG during his stay in Grenoble in October and November 2006.

This research has been supported by the ADONET network of the European Community, which is a Marie Curie Training Network.

References

- N. Alon, A. Gyárfás, B. Bollobás, J. Lehel and A. Scott, Maximum directed cuts in acyclic digraphs, J. Graph Theory 55 (2007), 1–13.
- [2] O. Amini, F. Havet, F. Huc, and S. Thomassé, Directed star arboricity of digraphs, Journées Graphes et Algorithmes, Orléans, Novembre 2006.
- [3] P. Berman, M. Karpinski, On some tighter inapproximability results (extended abstract) Lecture Notes in Comput. Sci., 1644, Springer, Berlin (1999), 200–209.
- [4] J. A. Bondy and S.C. Locke, Largest bipartite subgraphs in triangle-free graphs with maximum degree three, J. Graph Theory 10 (1986), 477–504.
- [5] R. Brandt, and T.F. Gonzalez, Wavelength assignment in multifiber optical star networks under the multicasting communication mode, Journal of Interconnection Networks 6 (2005), 383–405.

- [6] M. Cropper, M. Jacobson, A. Gyárfás, and J. Lehel, The Hall-ratio of graphs and hypergraphs, Les cahiers du Laboratoire Leibniz, No 17, Dec. 2000.
- [7] R. Diestel. *Graph Theory*. Graduate Texts in Mathematics 173, Springer, 1997.
- [8] B. Guiduli, On incidence coloring and star arboricity of graphs, Discrete Math. 163 (1997), 275–278.
- [9] E. Halperin, D. Livnat, U. Zwick. Max Cut in cubic graphs. J. Algorithms 53 (2004), 169–185.
- [10] J. Lehel and Zs. Tuza, Triangle-free partial graphs and edge covering theorems, Discrete Math. 39 (1982), 59–65.
- [11] B. Xu and X. Yu, Triangle-free subcubic graphs with minimum bipartite density, manuscript.
- [12] M. Yannakakis, Node- and edge-deletion NP-complete problems. Conference Record of the Tenth Annual ACM Symposium on Theory of Computing (San Diego, Calif., 1978), ACM, New York (1978), pp. 253–264.