

ON STRING GRAPH LIMITS AND THE STRUCTURE OF A TYPICAL STRING GRAPH

SVANTE JANSON AND ANDREW J. UZZELL

ABSTRACT. We study limits of convergent sequences of string graphs, that is, graphs with an intersection representation consisting of curves in the plane. We use these results to study the limiting behavior of a sequence of random string graphs. We also prove similar results for several related graph classes.

1. INTRODUCTION

Given a graph property \mathcal{P} , it is interesting to study the structure of a typical graph that satisfies \mathcal{P} . A natural definition of a “typical” graph is a graph chosen uniformly at random from all graphs of a given order that satisfy \mathcal{P} . One can choose a sequence of random graphs in this way and then study its limiting behavior. The theory of graph limits concerns the asymptotic behavior of certain sequences of graphs, and therefore provides a natural framework for studying the structure of a typical graph with a given property. In this paper, we will study the structure of string graphs. We will study graph limits of string graphs and of related graph classes and draw some conclusions about random string graphs and random elements of these other classes.

A *planar curve* is the image of a continuous function $f : [0, 1] \rightarrow \mathbb{R}^2$. The points $f(0)$ and $f(1)$ are called the *endpoints* of the curve. A *string representation* of a graph G is a collection of planar curves $\{A_v : v \in V(G)\}$ such that $A_u \cap A_v \neq \emptyset$ if and only if $uv \in E(G)$. We say that a graph G is a *string graph* if it has a string representation and we let \mathcal{SG} denote the family of string graphs. String graphs have been studied by many authors, see e.g. [27, 9, 19, 25] and the further references given there.

It is intuitively clear that a graph has a string representation if and only if it has an intersection representation consisting of arcwise-connected sets. Alternatively, we may assume that the curves in the definition are homeomorphic images of $[0, 1]$, and there are several other variations of the definition that give the same class of graphs. Although such equivalences are well-known, we have not found a detailed proof of these equivalences in the literature, so we give a proof in Appendix A.

As mentioned above, we will also study several special classes of string graphs. First, an *outer-string representation* of a graph G is a string representation such that all of the curves A_v lie in a disk and such that each A_v has an endpoint on the boundary of the disk. We say that a graph G is an *outer-string graph* if it has an outer-string representation and let \mathcal{OSG} denote the family of outer-string graphs. (Outer-string graphs were first so called in [18], but were studied in the monograph [19], there denoted Ng^0 . Sinden [27] studied a special case in which the strings are required to meet the boundary of the disk in a prescribed order.) It is clear from the definition that every outer-string graph is a string graph. It was shown in [19] that the converse does not hold; one consequence of our results

Date: 12 March, 2014.

Partly supported by the Knut and Alice Wallenberg Foundation.

is that only a very small fraction of all string graphs are outer-string graphs (see Remark 4.5).

Next, we consider the class of graphs with an intersection representation consisting of strings lying between two parallel segments, with one endpoint on each segment. It has been discovered several times [12, 19, 21] that this class is equal to the class of incomparability graphs. (Recall that if $<$ is a partial order on $[n]$, then the *incomparability graph* of $<$ is the graph with vertex set $[n]$ in which $x \sim y$ if and only if neither $x < y$ nor $y < x$.) We let \mathcal{ICG} denote the class of incomparability graphs.

Finally, we say that a graph is a *two-clique graph* if it is the disjoint union of at most two cliques. We let \mathcal{TCG} denote the class of two-clique graphs.

It is easy to see that all of these classes are hereditary. They are related by the following theorem [19, Theorem 5.8]. For $k \geq 1$, the authors of [19] defined $k\mathcal{NG}$ to be the set of graphs G with the property that if H_1, \dots, H_k are cliques such that the vertex sets of G and of the H_i are all pairwise disjoint, then any graph F with $V(F) = V(G \cup H_1 \cup \dots \cup H_k)$ and $E(F) \supseteq E(G \cup H_1 \cup \dots \cup H_k)$ is a string graph.

Theorem 1.1 ([19]). *We have the following characterizations of the classes $k\mathcal{NG}$.*

- (i) $1\mathcal{NG}$ equals the class \mathcal{OSG} of outer-string graphs;
- (ii) $2\mathcal{NG}$ equals the class \mathcal{ICG} of incomparability graphs;
- (iii) $3\mathcal{NG}$ equals the class \mathcal{TCG} of two-clique graphs.

Moreover, $4\mathcal{NG} = \{K_0\}$, where K_0 denotes the null graph, and $k\mathcal{NG} = \emptyset$ for $k \geq 5$. \square

It is easy to see that $k\mathcal{NG} \subseteq (k-1)\mathcal{NG}$ for all $k \geq 2$, and it is shown in [19] that (for $k \leq 5$) all of these inclusions are strict.

1.1. Preliminaries. As noted above, we will use the theory of graph limits to study these graph classes. For the basic notions of graph limits, see the recent book by Lovász [22] or, e.g., [4, 8, 23]. We recall that certain sequences of graphs (G_n) , with $|V(G_n)| \rightarrow \infty$, are defined to be *convergent*. A convergent sequence of graphs has a limit, a *graph limit*; these objects can be defined in several different ways. A *graphon* is a (measurable) symmetric function $[0, 1]^2 \rightarrow [0, 1]$. Each graphon W defines a unique graph limit Γ (we say that W *represents* Γ), and every graph limit is represented by some graphon; however, the representing graphon is not unique. We say that two graphons W and W' are *equivalent*, and write $W \cong W'$, if they represent the same graph limit. (Hence, the graph limits correspond to equivalence classes of graphons, and may be defined in this way.)

The *entropy* of a graphon W is defined as

$$\text{Ent}(W) = \int_0^1 \int_0^1 h(W(x, y)) \, dx \, dy,$$

where $h(x) = -x \log_2(x) - (1-x) \log_2(1-x)$ is the usual binary entropy function, see [14] and [6]. The entropy $\text{Ent}(\Gamma)$ of a graph limit Γ is the entropy of any representing graphon; equivalent graphons have the same entropy, so this defines $\text{Ent}(\Gamma)$ uniquely.

If \mathcal{P} is a graph class (or graph property, we do not distinguish between these), then $\widehat{\mathcal{P}}$ denotes the set of all graph limits Γ such that there exists a sequence $G_n \in \mathcal{P}$ with $G_n \rightarrow \Gamma$. We will also, slightly abusing the notation, let $\widehat{\mathcal{P}}$ denote the set of all graphons that represent such a graph limit. Furthermore, let $\widehat{\mathcal{P}}^*$ denote the set of graph limits (or graphons) in $\widehat{\mathcal{P}}$ with maximal entropy. (This set is nonempty, except in the trivial case when the graph class \mathcal{P} is finite, see [14].) For the importance of the set $\widehat{\mathcal{P}}^*$ of maximum-entropy graph limits, see [14].

We define some special graphons and sets of graphons, see further [14]. For convenience, we define them on $[0, 1]^2$ instead of $[0, 1]^2$; this is clearly immaterial.

Definition 1.2. Fix $k \geq 1$. For each $i \in [k]$, let $I_i = [(i-1)/k, i/k)$.

- (i) Let R_k be the set of all graphons W such that $W(x, y) = 1/2$ on $\bigcup_{i \neq j} I_i \times I_j$ and $W(x, y) \in \{0, 1\}$ on each $I_i \times I_i$. We also let R_∞ consist of the constant graphon $W = 1/2$.
- (ii) For $s \in \{0, \dots, k\}$, let $W_{k,s}^*$ be the graphon that is $1/2$ on $\bigcup_{i \neq j} I_i \times I_j$, 1 on each $I_i \times I_i$, $i \leq s$, and 0 on each $I_i \times I_i$, $i > s$.
- (iii) For $a \in [0, 1]$, let W_a^k denote the graphon obtained from $W_{k,k}^*$ by changing it on $I_1 \times I_1$ such that $W_a^k = 1$ on $[0, a/k]^2 \cup [a/k, 1]^2$ and $W_a^k = 0$ on $[0, a/k] \times [a/k, 1] \cup [a/k, 1] \times [0, a/k]$.

Note that $W_{k,s}^* \in R_k$ for $s = 0, \dots, k$, and $W_a^k \in R_k$. Furthermore, $\text{Ent}(W) = 1 - 1/k$ for every $W \in R_k$; in particular, $\text{Ent}(W_{k,s}^*) = \text{Ent}(W_a^k) = 1 - 1/k$.

We note that $W_0^k = W_1^k = W_{k,k}^*$ and that $W_a^k \cong W_{1-a}^k$. However, if $b \notin \{a, 1-a\}$, then $W_a^k \not\cong W_b^k$, because the two graphons have different edge densities. Indeed, for any $a \in [0, 1]$, we have

$$(1.1) \quad \int_0^1 \int_0^1 W_a^k = \frac{a^2 + (1-a)^2}{k^2} + \frac{k-1}{k^2} + \frac{k(k-1)}{2k^2}.$$

Given $t \geq 1$ and $0 \leq s \leq t$, we define $\mathcal{C}(t, s)$ to be the set of graphs whose vertex sets can be partitioned into s (possibly empty) cliques and $t-s$ (possibly empty) independent sets. In particular, $\mathcal{C}(t, 0)$ is the class of t -colorable graphs. If \mathcal{P} is a hereditary property, the *coloring number* of \mathcal{P} , denoted $\text{col}(\mathcal{P})$, is the largest t for which $\mathcal{C}(t, s) \subseteq \mathcal{P}$ for some $s \leq t$, see e.g. the survey [2]. (We define $\text{col}(\mathcal{P}) = \infty$ if \mathcal{P} is the class of all unlabeled finite graphs; otherwise, $\text{col}(\mathcal{P})$ is finite.)

Let \mathcal{U} denote the set of unlabeled finite graphs and let \mathcal{U}_n denote the set of unlabeled graphs on n vertices. Given a graph property \mathcal{P} , let $\mathcal{P}_n = \mathcal{P} \cap \mathcal{U}_n$ and let \mathcal{P}_n^L denote the set of labeled graphs in \mathcal{P} with vertex set $[n]$. The function $n \mapsto |\mathcal{P}_n^L|$ is called the (unlabeled) *speed* of \mathcal{P} . (The labeled speed is defined similarly. For our purposes, it does not matter whether we consider labeled or unlabeled graphs, since the difference is at most a factor $n! = 2^{o(n^2)}$, which will be negligible.) The speed of graph properties has been studied in many papers, see e.g. [1, 3, 2, 25].

Hatami, Janson, and Szegedy [14, Theorem 1.9] proved the following result that relates the maximum-entropy graph limits of a hereditary property \mathcal{P} , the speed of \mathcal{P} , and the coloring number of \mathcal{P} .

Theorem 1.3 ([14]). *If \mathcal{P} is a hereditary class of graphs, there exists $r \in \mathbb{N} \cup \{\infty\}$ such that $\max_{\Gamma \in \widehat{\mathcal{P}}} \text{Ent}(\Gamma) = 1 - 1/r$ and every graph limit $\Gamma \in \widehat{\mathcal{P}}^*$ can be represented by a graphon $W \in R_r$; hence $\widehat{\mathcal{P}}^* = \widehat{\mathcal{P}} \cap R_r$. Moreover,*

$$\begin{aligned} r &= \sup\{t : W_{t,s}^* \in \widehat{\mathcal{P}} \text{ for some } s \leq t\} \\ &= \sup\{t : \mathcal{C}(t, s) \subseteq \mathcal{P} \text{ for some } s \leq t\} \\ &= \text{col}(\mathcal{P}). \end{aligned}$$

Moreover,

$$(1.2) \quad |\mathcal{P}_n^L| = 2^{\left(1 - \frac{1}{r} + o(1)\right) \binom{n}{2}}.$$

We note that (1.2) was originally proved independently by Alekseev [1] and by Bollobás and Thomason [3].

1.2. Main Results. One of our aims is to classify maximum-entropy graph limits of string graphs and of related families of graphs. In order to do so, we prove a somewhat more general result about the maximum-entropy graph limits of certain hereditary properties. Before we can state this result, we need to define a few special graphs.

Let $k \geq 3$. We define three graphs, each with vertex set consisting of the $k + \binom{k}{2}$ subsets of $[k]$ of size either one or two. We denote the vertex corresponding to a subset S by v_S . First, let G_k be the intersection graph of this family of subsets, that is, the graph in which two vertices are adjacent if and only if their corresponding subsets have non-empty intersection. Second, we define B_k in the same way, except that we do not allow edges between two vertices that correspond to subsets of size two. Hence, B_k is bipartite. Finally, for $k \geq 4$, we define H_k be the subgraph of G_k with $E(H_k) = E(G_k) \setminus \{v_{\{1,2\}}v_{\{1,3\}}, v_{\{1,2\}}v_{\{2,3\}}, v_{\{1,3\}}v_{\{2,3\}}\}$. (Equivalently, H_k is such that $v_{\{s\}} \sim v_{\{t,u\}}$ if and only if $s = t$ or $s = u$, while $v_{\{s_1,s_2\}} \sim v_{\{t_1,t_2\}}$ if and only if $\{s_1, s_2\} \cap \{t_1, t_2\} \neq \emptyset$ and $\max\{s_1, s_2, t_1, t_2\} \geq 4$.)

Given a family of graphs \mathcal{H} , we define $\text{Forb}^*(\mathcal{H})$ to be the class of graphs that do not contain a copy of any $H \in \mathcal{H}$ as an induced subgraph. It is easy to see that $\text{Forb}^*(\mathcal{H})$ is a hereditary class.

Theorem 1.4. *Let \mathcal{P} be a hereditary class of graphs and let $r = \text{col}(\mathcal{P})$. If $3 \leq r < \infty$ and $\mathcal{P} \subseteq \text{Forb}^*(\{G_{r+1}, B_{r+1}, H_{r+1}\})$, then*

$$\widehat{\mathcal{P}}^* \subseteq \{W_a^r : a \in [0, 1]\}$$

up to equivalence of graphons.

If $r = 2$ and $\mathcal{P} \subseteq \text{Forb}^(\{G_3, B_3\})$, then*

$$\widehat{\mathcal{P}}^* \subseteq \{W_a^2 : a \in [0, 1]\}$$

up to equivalence of graphons.

Note that the assumption $\mathcal{P} \subseteq \text{Forb}^*(\{G_{r+1}, B_{r+1}, H_{r+1}\})$ is equivalent to $G_{r+1}, B_{r+1}, H_{r+1} \notin \mathcal{P}$.

Using Theorem 1.4, we can characterize maximum-entropy limits of string graphs, outer-string graphs, and incomparability graphs as follows. See also Theorem 3.3 for another related result.

Theorem 1.5. *Up to equivalence of graphons,*

$$\widehat{\mathcal{S}\mathcal{G}}^* = \{W_a^4 : a \in [0, 1]\}.$$

We note that the inclusion \supseteq in Theorem 1.5 was shown in [14, Example 2.4], but the question of equality was left open there.

Theorem 1.6. *Up to equivalence of graphons,*

$$\widehat{\mathcal{O}\mathcal{S}\mathcal{G}}^* = \{W_a^3 : a \in [0, 1]\}.$$

Theorem 1.7. *Up to equivalence of graphons,*

$$\widehat{\mathcal{I}\mathcal{C}\mathcal{G}}^* = \{W_a^2 : a \in [0, 1]\}.$$

Theorem 1.8. *Up to equivalence of graphons,*

$$\widehat{\mathcal{T}\mathcal{C}\mathcal{G}}^* = \{W_a^1 : a \in [0, 1]\}.$$

The proof of Theorem 1.8 is trivial, so we omit it. (The other theorems are proved later.) In fact, it is easy to see that $\widehat{\mathcal{T}\mathcal{C}\mathcal{G}} = \widehat{\mathcal{T}\mathcal{C}\mathcal{G}}^* = \{W_a^1 : a \in [0, 1]\}$. For the graph classes \mathcal{P} in Theorems 1.5–1.7, $\widehat{\mathcal{P}} \supseteq \widehat{\mathcal{P}}^*$, i.e., these classes have graph limits that do not have maximum entropy. We leave it as an open problem to classify all graph limits for these classes.

1.3. Results from Graph Limit Theory. Now we assemble the tools from graph limit theory that we will need in order to prove the results in Section 1.2.

Let $W : [0, 1]^2 \rightarrow [0, 1]$ be a graphon and let G be a graph with vertex set $[n]$. We define

$$(1.3) \quad \Psi_{G,W}^*(x_1, \dots, x_n) = \prod_{ij \in E(G)} W(x_i, x_j) \prod_{ij \notin E(G)} (1 - W(x_i, x_j))$$

and recall that the induced subgraph density of G in W is defined as

$$(1.4) \quad p(G, W) = \int_{[0,1]^n} \Psi_{G,W}^*.$$

We further say that the graph G is W -constructible if there exist (not necessarily distinct) points $x_1, \dots, x_n \in [0, 1]$ such that $\Psi_{G,W}^*(x_1, \dots, x_n) > 0$, i.e., more explicitly,

$$(1.5) \quad W(x_i, x_j) = 0 \implies ij \notin E(G)$$

and

$$(1.6) \quad W(x_i, x_j) = 1 \implies ij \in E(G).$$

(If $0 < W(x_i, x_j) < 1$, then there is no restriction on ij .) If $\mathbf{x} = (x_1, \dots, x_n)$ is such that (1.5) and (1.6) hold, i.e. $\Psi_{G,W}^*(\mathbf{x}) > 0$, then we say that \mathbf{x} is a *witnessing vector* for G .

Given a graphon W , there is a standard definition of a W -random graph $G(n, W)$: let X_1, X_2, \dots be an i.i.d. sequence of uniform random variables in $[0, 1]$. For each n , $G(n, W)$ is a graph with vertex set $[n]$ in which the edge ij is present with probability $W(X_i, X_j)$, independently of all other edges. By [22, Proposition 11.32], $G(n, W) \rightarrow W$ almost surely.

Remark 1.9. It follows from (1.4) that for every graph G with vertex set $[n]$,

$$\mathbb{P}(G(n, W) = G) = p(G; W).$$

Hence,

$$(1.7) \quad \mathbb{P}(G(n, W) = G) = 0 \iff p(G, W) = 0 \iff \Psi_{G,W}^*(\mathbf{x}) = 0 \text{ for a.e. } \mathbf{x}.$$

This is thus a condition for a.e. \mathbf{x} , while we have defined G to be not W -constructible if $\Psi_{G,W}^*(\mathbf{x}) = 0$ for *every* \mathbf{x} . This looks like a minor technical difference, and indeed it is (although this turns out to be non-trivial to prove). Petrov's general removal lemma [26, Theorem 1] shows that if the properties in (1.7) hold, then W can be modified on a null set such that then $\Psi_{G,W}^* = 0$ everywhere, i.e., G is not W -constructible. (Note that the properties in (1.7) are preserved if W is modified on a null set, and more generally if W is replaced by an equivalent graphon, but the property that G is (not) W -constructible is not.)

Let \mathcal{P} be a hereditary class. It is easily seen, see [15, Theorem 3.3], that if W is a graphon, then $W \in \widehat{\mathcal{P}}$ if and only if $p(G; W) = 0$ for every graph $G \notin \mathcal{P}$. In particular, see Diaconis, Holmes, and Janson [7, Theorem 3.2], if \mathcal{H} is a set of graphs and $\mathcal{P} = \text{Forb}^*(\mathcal{H})$, then $\Gamma \in \widehat{\mathcal{P}}$ if and only if $p(H; \Gamma) = 0$ for all $H \in \mathcal{H}$. We will need the following extension of this, which is a consequence of Petrov's removal lemma [26, Theorem 1], see Section 7.

Lemma 1.10. *Let \mathcal{P} be a hereditary class of graphs. If $\Gamma \in \widehat{\mathcal{P}}$, then there exists a graphon W representing Γ such that if a graph G is W -constructible, then $G \in \mathcal{P}$. Moreover, if $\Gamma \in \widehat{\mathcal{P}}^*$ and $r = \text{col}(\mathcal{P})$, then there exists $W \in R_r$ such that if G is W -constructible, then $G \in \mathcal{P}$.*

We conclude this section with a special class of graphons that will be needed in the proof of Theorem 1.4. A *disjoint clique graphon* is a graphon of the form $W = \sum_{\alpha \in \mathcal{A}} \mathbf{1}_{A_\alpha \times A_\alpha}$ for a family $(A_\alpha)_{\alpha \in \mathcal{A}}$ of non-empty pairwise disjoint subsets of $[0, 1]$. Here the index set \mathcal{A} may be finite, countably infinite or uncountable. We say that W has *parts* A_α , $\alpha \in \mathcal{A}$.

Since a graphon is assumed to be (Lebesgue) measurable, the section $\{y : W(x, y) = 1\}$ is measurable for a.e. $x \in [0, 1]$. In particular, if W is a disjoint clique graphon and A_α one of its parts, then either A_α is a null set, and thus measurable, or there exists $x \in A_\alpha$ such that $\{y : W(x, y) = 1\}$ is measurable. Since the latter set equals A_α , we see that in both cases A_α is measurable. In other words, the parts of a disjoint clique graphon are all measurable.

Remark 1.11. It is easy to see that up to equivalence, we may eliminate all parts with measure 0, leaving only a countable number of parts A_n . Moreover, up to equivalence, a disjoint clique graphon is uniquely determined by the sequence $|A_n|$ of measures of these parts, arranged in (weakly) decreasing order. It is also easy to see that if \mathcal{DC} is the family of *disjoint clique graphs*, i.e. graphs that are disjoint unions of cliques, then the set of all disjoint clique graphons equals $\widehat{\mathcal{DC}}$. See further Janson [15, Section 7].

The rest of this paper is organized as follows. In Section 2, we prove Theorem 1.4. In Sections 3, 4, and 5, we prove Theorems 1.5, 1.6, and 1.7, respectively. In Section 6, we derive one more result each about the structure of typical string graphs and the structure of typical outer-string graphs, and make several conjectures. Finally, in Section 7, we prove Lemma 1.10.

2. PROOF OF THEOREM 1.4

The plan of the proof of Theorem 1.4 is straightforward. We want to show that if \mathcal{P} is as in the statement of the theorem, then any maximum-entropy element Γ of $\widehat{\mathcal{P}}$ can be represented by a graphon W that has a number of desirable properties, eventually leading to the conclusion that $W \cong W_a^r$ for some a . We use Lemma 1.10 to find a “nice” version of W ; then, in each case, we will show that if W does not have the desired property, then this implies that \mathcal{P} contains at least one of the graphs G_{r+1} , B_{r+1} , and H_{r+1} , which is a contradiction.

Proof of Theorem 1.4. Suppose that $\Gamma \in \widehat{\mathcal{P}}^*$ has entropy $1 - 1/r$ and let $W \in R_r$ be the graphon representing Γ whose existence is guaranteed by Lemma 1.10. Thus, every graph that is W -constructible belongs to \mathcal{P} . In particular, our assumption on \mathcal{P} implies that G_{r+1} , B_{r+1} , and H_{r+1} are *not* W -constructible. For (notational) convenience, we let all graphons be defined on $[0, 1]$ in this section.

Let W_i denote the restriction of W to $I_i \times I_i$. By rescaling the interval I_i to $[0, 1]$, we may regard W_i as a graphon. Recall that by the definition of R_r , W_i takes only the values 0 and 1.

Claim 1: At most one of the sets I_i can contain some point x such that $W(x, x) = 0$. If not, then without loss of generality there exist $x \in I_1$ and $y \in I_2$ such that $W(x, x) = W(y, y) = 0$. Let F be a bipartite graph. If (A, B) is a bipartition of $V(F)$ with $|A| = n_1$ and $|B| = n_2$, then we can construct a witnessing vector for F by choosing n_1 copies of x and n_2 copies of y . Thus, we may conclude that F is W -constructible. In particular, B_{r+1} is W -constructible, which is a contradiction.

Claim 2: $W = 1$ along the diagonal. If not, then without loss of generality there exists $z \in I_1$ such that $W(z, z) = 0$. We then partition the vertex set of G_{r+1} as follows: let $V_1 = \{1, 2, 3\}$, let $V_2 = \{\{1, 2\}, \{1, 3\}, \{2, 3\}\}$, and for $\ell = 3, \dots, r$, let $V_\ell = \{\{\ell+1, j\} : 1 \leq j \leq \ell+1\}$. (Here, we write $\{i, i\}$ for $\{i\}$.) Observe that each of

V_2, \dots, V_r induces a complete graph, while V_1 induces an empty graph. For $j = 2, \dots, r$, let $v_j \in I_j$ be arbitrary. By Claim 1, for each j , we have $W(v_j, v_j) = 1$. Then we can construct a witnessing vector for G_{r+1} by choosing three copies of z and $j + 1$ copies of the point v_j for $j = 2, \dots, r$. Thus, G_{r+1} is W -constructible, which is a contradiction.

Claim 3: If $x, y, z \in I_1$ and $W(x, y) = W(y, z) = 1$, then $W(x, z) = 1$. We partition the vertex set of G_{r+1} in a different way. Let $U_1 = \{1, 2, \{1, 2\}\}$ and for $\ell = 2, \dots, r$, let $U_\ell = \{\{\ell + 1, j\} : 1 \leq j \leq \ell + 1\}$. Observe that U_1 induces a copy of P_3 and that each of U_2, \dots, U_r induces a complete graph. Suppose that there exist $x, y, z \in I_1$ such that $W(x, y) = W(y, z) = 1$ but $W(x, z) = 0$. For $\ell = 2, \dots, r$, let $u_\ell \in I_\ell$ be arbitrary and note that, by Claim 2, $W(u_\ell, u_\ell) = 1$. Then the vector consisting of x, y, z , and $\ell + 1$ copies of each point u_ℓ is a witnessing vector for G_{r+1} , which implies that G_{r+1} is W -constructible. This contradiction proves the claim.

Claim 4: Each W_i is a disjoint clique graphon $\sum_{\alpha \in \mathcal{A}_i} \mathbf{1}_{A_{i,\alpha} \times A_{i,\alpha}}$, rescaled to I_i . It suffices to consider $i = 1$. The relation on I_1 defined by $x \equiv y$ if $W(x, y) = 1$ is symmetric since W is, reflexive by Claim 2, and transitive by Claim 3; hence it is an equivalence relation. Denote the equivalence classes by $(A_\alpha)_{\alpha \in \mathcal{A}}$; then $W_1 = \sum_{\alpha \in \mathcal{A}} \mathbf{1}_{A_\alpha \times A_\alpha}$.

Claim 5: Each W_i has at most two parts $A_{i,\alpha}$. If not, then without loss of generality W_1 has at least three parts A_1, A_2, A_3 . We partition the vertex set of G_{r+1} using the partition (V_1, \dots, V_r) defined in the proof of Claim 2 above. Let $a_i \in A_i$ for $i = 1, 2, 3$ and observe that $W(a_i, a_j) = \delta_{ij}$. For $j = 2, \dots, r$, let $v_j \in I_j$ be arbitrary. Observe that the vector consisting of a_1, a_2, a_3 , and $j + 1$ copies of each point v_j is a witnessing vector for G_{r+1} . Thus G_{r+1} is W -constructible, which is a contradiction.

Claim 6: $W = 1$ on at least $r - 1$ of the squares $I_i \times I_i$. If not, there are at least two restrictions W_i such that the partitions $(A_{i,\alpha})$ in Claim 4 have at least two parts, i.e., $|\mathcal{A}_i| \geq 2$. Suppose without loss of generality that $|\mathcal{A}_1|, |\mathcal{A}_2| \geq 2$, and let A_1, A_2 be parts of W_1 and B_1, B_2 be parts of W_2 . First, suppose that $r \geq 3$. We claim that H_{r+1} is W -constructible. To show this, we partition $V(H_{r+1})$ as follows. Let $X'_1 = \{1, \{1, 3\}\}$, let $X''_1 = \{\{2, 3\}\}$, let $X'_2 = \{2, \{1, 2\}\}$, and let $X''_2 = \{3\}$. For $\ell = 3, \dots, r$, let $X_\ell = \{\{\ell + 1, j\} : 1 \leq j \leq \ell + 1\}$. Observe that the sets $X'_1, X''_1, X'_2, X''_2, X_3, \dots, X_r$ all induce complete subgraphs, that there are no edges between X'_1 and X''_1 , and that there are no edges between X'_2 and X''_2 . Let $x'_1 \in A_1$, let $x''_1 \in A_2$, let $x'_2 \in B_1$, let $x''_2 \in B_2$, and, for $\ell = 3, \dots, r$, let $x_\ell \in I_\ell$. Let \mathbf{x} denote the vector consisting of two copies of x'_1 , one copy of x''_1 , two copies of x'_2 , one copy of x''_2 , and $\ell + 1$ copies of x_ℓ for $\ell = 3, \dots, r$. Observe that \mathbf{x} is a witnessing vector for H_{r+1} . Thus H_{r+1} is W -constructible, which is a contradiction. If $r = 2$, we may repeat the argument above to show that B_3 is W -constructible, which is again a contradiction. This proves the claim.

To summarize, we have shown that if $\Gamma \in \widehat{\mathcal{P}}^*$, then Γ may be represented by a graphon $W \in R_r$ such that there exists a measurable set $A_1 \subseteq I_1$ such that $W = 1$ on $(A_1 \times A_1) \cup ((I_1 \setminus A_1) \times (I_1 \setminus A_1))$, $W = 0$ on the rest of $I_1 \times I_1$, and $W = 1$ on $\bigcup_{j=2}^r I_j \times I_j$. That is, $W \cong W_a^r$ for some $a \in [0, 1]$, as claimed. \square

3. PROOF OF THEOREM 1.5

We begin by showing that the class of string graphs satisfies the hypotheses of Theorem 1.4.

Lemma 3.1. *The graphs G_5, B_5 , and H_5 are not string graphs.*

Proof. The result for B_5 has been discovered several times [27, 9, 19], while the result for G_5 was shown in [25, Lemma 3.2]. The argument that we give is a slight modification of the proof used in [25]. (It actually works for any graph F with $B_5 \subseteq F \subseteq G_5$.)

Suppose that any of G_5 , B_5 , and H_5 had a string representation $\{A_i, A_{ij}\}$. Then, we could select points $v_i \in A_i$ for each i and points $v_{ij} \in A_i \cap A_{ij}$, for each i and j with $i \neq j$. Then, for each i and j , we could let e_{ij} be a curve consisting of the portion of A_i between v_i and v_{ij} , the portion of A_{ij} between v_{ij} and v_{ji} , and the portion of A_j between v_{ji} and v_j . Because $e_{ij} \subseteq A_i \cup A_j \cup A_{ij}$, we have $e_{ij} \cap e_{k\ell} = \emptyset$ if i, j, k , and ℓ are all distinct. Hence, the points v_i and the curves e_{ij} define a drawing of K_5 in which no two independent edges cross, which contradicts the Hanani–Tutte theorem [13, 29]. \square

Let G be a graph and let $\mathcal{C} = \{V_1, \dots, V_k\}$ be a clique covering of G . We can define a graph $G_{\mathcal{C}}$ with $V(G_{\mathcal{C}}) = [k]$ and $E(G_{\mathcal{C}}) = \{ij : E(V_i, V_j) \neq \emptyset\}$. We say that a graph G admits a *planar clique covering* if there is a clique covering \mathcal{C} of G such that $G_{\mathcal{C}}$ is planar. Similarly, we say that G admits an *outerplanar clique covering* if there is a clique covering \mathcal{C} of G such that $G_{\mathcal{C}}$ is outerplanar.

Proof of Theorem 1.5. It is shown in [25] that $\text{col}(\mathcal{SG}) = 4$ (see also Lemma 4.2). This, Lemma 3.1, and Theorem 1.4 imply that $\widehat{\mathcal{SG}}^* \subseteq \{W_a^4 : a \in [0, 1]\}$ up to equivalence of graphons.

In the opposite direction, it is shown in [14, Example 2.4] that $W_a^4 \in \widehat{\mathcal{SG}}^*$ for every $a \in [0, 1]$. For later use, we repeat the argument.

It is shown in [19, Corollary 2.7] that $\mathcal{C}(4, 4) \subseteq \mathcal{SG}$ and more generally in [19, Theorem 2.3] that every graph that admits a planar clique covering is a string graph. (See also [14, Example 2.4].) Let $a \in [0, 1]$ and, for each n , let $G_n = G(n, W_a^4)$, where $G(n, W_a^4)$ is the W_a^4 -random graph defined in Section 1.3. Let $V_1' = \{i : X_i \in [0, a/4]\}$, let $V_1'' = \{i : X_i \in [a/4, 1/4]\}$, and, for $j = 2, 3, 4$, let $V_j = \{i : X_i \in I_j\}$. Observe that with probability one, V_1', V_1'', V_2, V_3 , and V_4 are all cliques and there are no edges between V_1' and V_1'' . This means that each G_n almost surely admits a planar clique covering \mathcal{C} with $G_{\mathcal{C}} = K_5 - e$, which means that each G_n is a string graph. Because $G_n \rightarrow W_a^4$ almost surely, it follows that $W_a^4 \in \widehat{\mathcal{SG}} \cap R_4$, which, by Theorem 1.3, implies that $W_a^4 \in \widehat{\mathcal{SG}}^*$, as claimed. This completes the proof. \square

Remark 3.2. Because $\text{col}(\mathcal{SG}) = 4$, it follows from Theorem 1.3 that

$$(3.1) \quad |\mathcal{SG}_n| = 2^{\binom{3}{4} + o(1)} \binom{n}{2},$$

as proved by Pach and Tóth [25].

A graph G is a *convex intersection graph* if it has an intersection representation consisting of convex sets in the plane. Let \mathcal{CVX} denote the class of convex intersection graphs. It is easy to see that \mathcal{CVX} is hereditary and that $\mathcal{CVX} \subseteq \mathcal{SG}$. It is shown in [19, Proposition 8.3.1] that \mathcal{CVX} is in fact a proper subclass of \mathcal{SG} . However, the next result shows that \mathcal{CVX} has the same asymptotic speed as the larger class \mathcal{SG} and furthermore shows that \mathcal{CVX} has the same maximum-entropy graph limits as \mathcal{SG} , i.e. $\widehat{\mathcal{CVX}}^* = \widehat{\mathcal{SG}}^*$. (The sets of all graph limits for these classes differ, i.e., $\widehat{\mathcal{CVX}} \subsetneq \widehat{\mathcal{SG}}$. For example, if $G \in \mathcal{SG} \setminus \mathcal{CVX}$, then the adjacency matrix of G defines a graphon, and thus a graph limit, that easily is seen to belong to $\widehat{\mathcal{SG}} \setminus \widehat{\mathcal{CVX}}$, cf. [15, Section 3] and [24, Proposition 4.10].)

Theorem 3.3. *We have*

$$(3.2) \quad \text{col}(\mathcal{CVX}) = 4$$

and

$$(3.3) \quad |\mathcal{CVX}_n| = 2^{\binom{\frac{3}{4} + o(1)}{2} \binom{n}{2}}.$$

Furthermore, up to equivalence of graphons,

$$(3.4) \quad \widehat{\mathcal{CVX}}^* = \{W_a^4 : a \in [0, 1]\}.$$

Proof. It is shown in [20] that $\mathcal{C}(4, 4) \subseteq \mathcal{CVX}$ and, more generally, that every graph that admits a planar clique covering is a convex intersection graph. Hence, we have $\text{col}(\mathcal{CVX}) \geq 4$, and since $\mathcal{CVX} \subseteq \mathcal{SG}$, we have $\text{col}(\mathcal{CVX}) \leq \text{col}(\mathcal{SG}) = 4$. This proves (3.2) and (3.3) follows by (1.2). Finally, (3.4) follows by the same argument as in the proof of Theorem 1.5. \square

4. PROOF OF THEOREM 1.6

Lemma 4.1. *The graphs G_4 , B_4 , and H_4 are not outer-string graphs.*

Proof. Suppose to the contrary that G_4 , B_4 , or H_4 has an outer-string representation. If necessary, we may assume (possibly by extending the outer-string representation to a slightly larger disk) that all curves meet the disk at distinct points. Then we may extend this representation to a string representation of G_5 , B_5 , or H_5 by adding curves $A_{\{5\}}$, $A_{\{1,5\}}$, \dots , $A_{\{4,5\}}$ outside of the disk with the appropriate intersection pattern. (See also Theorem 1.1.) However, this contradicts Lemma 3.1, and the claimed result follows. \square

We will also need the following result of Pach and Tóth [25, Lemma 3.1].

Lemma 4.2 ([25]). *If $r \geq 1$, then $G_{r+1} \in \mathcal{C}(r+1, s)$ for $s = 0, \dots, r+1$. In particular, if \mathcal{P} is a hereditary property and $\mathcal{P} \subseteq \text{Forb}^*(\{G_{r+1}\})$, then $\text{col}(\mathcal{P}) \leq r$.* \square

Lemma 4.3. *Every graph in the class $\mathcal{C}(3, 3)$ is an outer-string graph. Moreover, every graph that admits an outerplanar clique covering is an outer-string graph.*

Proof. As stated above, the corresponding results for string graphs were proved in [19, Theorem 2.3] and in [14, Example 2.4]. We modify the construction given in the latter.

Let G be a graph whose vertex set can be covered by three cliques V_1 , V_2 , and V_3 . Place distinct points v_1 , v_2 , and v_3 on the boundary of a disk D . For each $x \in V_i$ and $y \in V_j$ with $i \neq j$, add a curve $\tilde{A}_{xy} \subset D$ between v_i and v_j . We may place the curves in such a way that different curves do not meet apart from their endpoints. For each $x \in V_i$ and $y \in V_j$ with $i \neq j$, choose a point $a_{xy} \in \tilde{A}_{xy}$ that is different from both v_i and v_j . Let \tilde{A}_{xy}^* denote the portion of \tilde{A}_{xy} between v_i and a_{xy} and let \tilde{A}_{yx}^* denote the portion of \tilde{A}_{xy} between v_j and a_{xy} . Let, for $x \in V_i$,

$$A_x = \left(\bigcup_{\substack{xy \in E(G), \\ y \notin V_i}} \tilde{A}_{xy}^* \right).$$

It is easy to see that the collection of curves $\{A_v : v \in V(G)\}$ gives an outer-string representation of G . (We can regard each A_x as a curve starting and ending at v_i , traversing each \tilde{A}_{xy}^* in both directions, with these parts in arbitrary order.)

In general, if G admits an outerplanar clique covering with $\mathcal{C} = \{V_1, \dots, V_k\}$, we may place k distinct points on the boundary of the disk and repeat the construction described above to define an outer-string representation of G . \square

Lemma 4.4. *We have*

$$\text{col}(\mathcal{OSG}) = 3$$

and

$$(4.1) \quad |\mathcal{OSG}_n| = 2^{\binom{\frac{2}{3}+o(1)}{2} \binom{n}{2}}.$$

Proof. The fact that $\text{col}(\mathcal{OSG}) = 3$ is immediate from Lemmas 4.1, 4.2, and 4.3. This and (1.2) imply (4.1). \square

Remark 4.5. Because $G_4 \in \mathcal{C}(4,4) \subseteq \mathcal{SG}$, G_4 is an example of a string graph that is not an outer-string graph. There are many others: (3.1) and (4.1) imply that almost every string graph is not an outer-string graph.

Proof of Theorem 1.6. One inclusion is immediate from Lemma 4.1, Lemma 4.4, and Theorem 1.4. For the other, let $a \in [0,1]$. As in the proof of Theorem 1.5, one can define a sequence of graphs $G_n = G(n, W_a^3)$ such that for each n , with probability one, G_n admits a clique covering \mathcal{C} such that $G_{\mathcal{C}} = K_4 - e$, and such that $G_n \rightarrow W_a^3$ almost surely as $n \rightarrow \infty$. Because $K_4 - e$ is outerplanar, it follows from Lemma 4.3 that each G_n is an outer-string graph. Theorem 1.3 then implies that $W_a^3 \in \widehat{\mathcal{OSG}} \cap R_3 = \widehat{\mathcal{OSG}}^*$, as claimed. \square

5. PROOF OF THEOREM 1.7

Lemma 5.1. *The graphs G_3 and B_3 are not incomparability graphs.*

Proof. Observe that $B_3 = C_6$ and that G_3 is the complement of a triangle with a pendant edge attached to each vertex. It is a well-known result of Gallai [11] (see also [28]) that neither of these graphs is an incomparability graph.

Alternatively, suppose that G_3 or B_3 had an intersection representation consisting of strings between two parallel segments. By arguing as in the proof of Lemma 4.1, we could extend this representation to an outer-string representation of G_4 or B_4 , a contradiction. \square

Lemma 5.2. *We have*

$$\text{col}(\mathcal{ICG}) = 2$$

and

$$(5.1) \quad |\mathcal{ICG}_n| = 2^{\binom{\frac{1}{2}+o(1)}{2} \binom{n}{2}}.$$

Remark 5.3. The asymptotic speed of the class of incomparability graphs (equation (5.1)) was first determined by Kleitman and Rothschild [16, 17].

Proof of Lemma 5.2. It is easy to see that every bipartite graph is a comparability graph, which is equivalent to the statement that $\mathcal{C}(2,2) \subseteq \mathcal{ICG}$. It follows that $\text{col}(\mathcal{ICG}) \geq 2$. Lemmas 5.1 and 4.2 then imply that $\text{col}(\mathcal{ICG}) = 2$. Finally, (5.1) follows from (1.2). \square

In the proof of Theorem 1.7, instead of directly proving our results about incomparability graphs, we will find it convenient to prove the corresponding results for comparability graphs. We let \mathcal{CG} denote the class of comparability graphs.

Proof of Theorem 1.7. One inclusion is immediate from Lemma 5.1, Lemma 5.2, and Theorem 1.4. In order to show that

$$\widehat{\mathcal{ICG}}^* \supseteq \{W_a^2 : a \in [0,1]\},$$

it is enough to prove that $1 - W_a^2 \in \widehat{\mathcal{CG}}^*$ for every $a \in [0,1]$.

For $a \in \{0,1\}$, the result follows from the fact that every bipartite graph is a comparability graph. Hence, we may suppose that $a \in (0,1)$. For each n , we let $G_n = G(n, 1 - W_a^2)$. Let $A_n = \{i : X_i \in [0, a/2)\}$, $C_n = \{i : X_i \in [a/2, 1/2)\}$,

and $B_n = \{i : X_i \in [1/2, 1)\}$. Observe that with probability one, each of A_n , B_n , and C_n is an incomparable set, and $A_n \cup C_n$ is a complete bipartite graph. Hence, if we orient edges from A_n to B_n , from A_n to C_n and from B_n to C_n , then if ab and bc are directed edges, then ac is almost surely a directed edge. Thus, each G_n is almost surely a comparability graph. Because $G_n \rightarrow 1 - W_a^2$ almost surely and $\text{col}(\mathcal{CG}) = \text{col}(\mathcal{ICG}) = 2$, it follows from Theorem 1.3 that $1 - W_a^2 \in \widehat{\mathcal{CG}} \cap R_2 = \widehat{\mathcal{CG}}^*$, as claimed. \square

6. RANDOM STRING GRAPHS

Let \mathcal{P} be any graph class. The next result says that if the maximum entropy of an element of $\widehat{\mathcal{P}}$ determines the speed of \mathcal{P} , then the maximum-entropy elements of $\widehat{\mathcal{P}}$ also determine the asymptotic structure of a typical element of \mathcal{P} . This was proved in [14, Theorem 1.6] in the special case when $\widehat{\mathcal{P}}^*$ consists of a single element. Since the proof of the general case is nearly identical to the argument given in [14], we omit it.

Given $\delta > 0$, let $A_\delta(\mathcal{P}) = \{\Gamma \in \widehat{\mathcal{U}} : \delta_\square(\Gamma, \widehat{\mathcal{P}}^*) < \delta\}$, where δ_\square is the standard metric on the space of graph limits, see e.g. [22].

Theorem 6.1. *Let \mathcal{P} be a class of graphs and suppose that*

$$(6.1) \quad \lim_{n \rightarrow \infty} \frac{\log_2 |\mathcal{P}_n|}{\binom{n}{2}} = \max_{\Gamma \in \widehat{\mathcal{P}}} \text{Ent}(\Gamma).$$

For each n , let G_n be a uniformly random unlabeled element of \mathcal{P}_n . For any $\delta > 0$,

$$\lim_{n \rightarrow \infty} \mathbb{P}(G_n \in A_\delta(\mathcal{P})) = 1.$$

Moreover, the same conclusion holds if we let each G_n be a uniformly random labeled element of \mathcal{P}_n . \square

Remark 6.2. By Theorem 1.3, equation (6.1) holds whenever \mathcal{P} is a hereditary property.

It is immediate from Theorem 6.1 and Theorems 1.5 and 1.6, respectively, that if G_n is a uniformly random (unlabeled or labeled) string graph, then G_n converges in probability as $n \rightarrow \infty$ to the set $\{W_a^4 : a \in [0, 1]\}$, and similarly that a sequence of uniformly random outer-string graphs converges in probability to the set $\{W_a^3 : a \in [0, 1]\}$. In principle, it is possible that the sequence G_n converges in distribution to a limit that is a non-degenerate random graph limit in $\{W_a^4\}$, or that the distributions oscillate and do not converge at all, but it seems very probable that there exists some $a \in [0, 1/2]$ such that $G_n \xrightarrow{\mathbb{P}} W_a^4$. However, assuming that this really holds, what is the limiting a ? We believe that $a = 1/2$, and thus that the following stronger results should also hold.

Conjecture 6.3. *If, for each n , G_n is a uniformly random unlabeled element of \mathcal{SG}_n , then $G_n \xrightarrow{\mathbb{P}} W_{1/2}^4$. Moreover, the same conclusion holds if each G_n is a uniformly random labeled element of \mathcal{SG}_n^L .*

Conjecture 6.4. *If, for each n , G_n is a uniformly random unlabeled element of \mathcal{OSG}_n , then $G_n \xrightarrow{\mathbb{P}} W_{1/2}^3$. Moreover, the same conclusion holds if each G_n is a uniformly random labeled element of \mathcal{OSG}_n^L .*

We believe that these results should hold for two reasons. First, in the proof of Theorem 1.5, the value $a = 1/2$ gives the largest number of partitions of the vertex set. (And similarly for outer-string graphs, see the proof of Theorem 1.6.) Second, the corresponding results for incomparability graphs (at least in the labeled case) and for two-clique graphs are known to hold. The result for labeled incomparability

graphs follows from the corresponding result for partial orders, which was proved in [17] (see also [5]). We believe that the same result should hold for unlabeled incomparability graphs; it is a folklore result that almost every partial order has trivial automorphism group, and we believe that the same is true of incomparability graphs, but we do not know a proof of this statement. The result for two-clique graphs is trivial; for a sketch of a nearly identical argument, see [15, Example 7.9].

Theorem 6.5. *If, for each n , G_n is a uniformly random labeled element of \mathcal{ICG}_n^L , then $G_n \xrightarrow{\mathbb{P}} W_{1/2}^2$. \square*

Theorem 6.6. *If, for each n , G_n is a uniformly random unlabeled element of \mathcal{TCG}_n , then $G_n \xrightarrow{\mathbb{P}} W_{1/2}^1$. If each G_n is a uniformly random labeled element of \mathcal{TCG}_n^L , then G_n converges in probability to the random graphon W_T^1 , where $T \sim U(0, 1/2)$. \square*

It is easy to see that the limiting distribution of a uniformly random string graph, if it exists, determines the limiting distribution of the edge density of a uniformly random string graph (and similarly for outer-string graphs). It also determines the limiting distribution of the proportion of vertices of degree approximately cn , where $c \in [0, 1]$ is a constant; it is convenient to state this using the distribution of the degree of a uniformly random vertex. (See e.g. [7, Section 4].) Thus, Conjectures 6.3 and 6.4 imply the following conjectures, including the somewhat surprising statement that the degree distribution of a typical string graph (and of a typical outer-string graph) is bimodal.

Given a graph G on n vertices, let X_n denote the degree of a uniformly random vertex of G .

Conjecture 6.7. *If, for each n , G_n is a uniformly random unlabeled element of \mathcal{SG}_n , then*

$$(6.2) \quad \frac{e(G_n)}{\binom{n}{2}} \xrightarrow{\mathbb{P}} \frac{19}{32}.$$

Furthermore, X_n/n converges in distribution to a random variable Y such that $y \in \{1/2, 5/8\}$ almost surely and such that

$$\mathbb{P}(Y = 1/2) = \frac{1}{4} \quad \text{and} \quad \mathbb{P}(Y = 5/8) = \frac{3}{4}.$$

Moreover, the same conclusions hold if each G_n is a uniformly random labeled element of \mathcal{SG}_n^L .

Conjecture 6.8. *If, for each n , G_n is a uniformly random unlabeled element of \mathcal{OSG}_n , then*

$$(6.3) \quad \frac{e(G_n)}{\binom{n}{2}} \xrightarrow{\mathbb{P}} \frac{11}{18}.$$

Furthermore, X_n/n converges in distribution to a random variable Y such that $y \in \{1/2, 2/3\}$ almost surely and such that

$$\mathbb{P}(Y = 1/2) = \frac{1}{3} \quad \text{and} \quad \mathbb{P}(Y = 2/3) = \frac{2}{3}.$$

Moreover, the same conclusions hold if each G_n is a uniformly random labeled element of \mathcal{OSG}_n^L .

Let us also remark that because $W_{1/2}^4$ has minimum edge density in the set $\{W_a^4 : a \in [0, 1]\}$ (cf. (1.1)), the assertion (6.2) is actually equivalent to Conjecture 6.3; moreover, (6.2) is also equivalent to saying that the edge density of a random string graph converges in expectation to $19/32$. Similarly, (6.3) is equivalent to the analogous statements about outer-string graphs.

7. PROOF OF LEMMA 1.10

We find it convenient to make two new definitions, closely related to concepts in Section 1.3 but from a slightly different point of view. Recall (1.3)–(1.4).

Definition 7.1. Let W be a graphon and G a graph with vertex set $[n]$.

(i) W is G -free if $p(G; W) = 0$. By (1.4), this is equivalent to

$$(7.1) \quad \Psi_{G,W}^*(x_1, \dots, x_n) = 0 \quad \text{for a.e. } x_1, \dots, x_n \in [0, 1].$$

(ii) W is completely G -free if

$$(7.2) \quad \Psi_{G,W}^*(x_1, \dots, x_n) = 0 \quad \text{for all } x_1, \dots, x_n \in [0, 1].$$

In other words, G is not W -constructible.

We emphasize that in (ii), the sequence x_1, \dots, x_n is completely arbitrary, and may contain repetitions. In particular, we require $\Psi_{G,W}^*(x, \dots, x) = 0$ for every $x \in [0, 1]$, which by (1.3) implies $W(x, x) \in \{0, 1\}$ for every $x \in [0, 1]$.

As said in the introduction, by [15, Theorem 3.3], if \mathcal{P} is hereditary, then a graphon $W \in \widehat{\mathcal{P}}$ if and only if $p(G; W) = 0$ for every graph $G \notin \mathcal{P}$; in other words, if and only if W is G -free for every $G \notin \mathcal{P}$. The following theorem says that this can be strengthened to completely G -free.

Theorem 7.2. *Let \mathcal{P} be a hereditary graph property and Γ a graph limit. Then the following are equivalent.*

(i) $\Gamma \in \widehat{\mathcal{P}}$.

(ii) Every graphon representing Γ is G -free for every $G \notin \mathcal{P}$.

(iii) There exists a graphon representing Γ that is G -free for every $G \notin \mathcal{P}$.

(iv) There exists a graphon representing Γ that is completely G -free for every $G \notin \mathcal{P}$.

Moreover, suppose that $[0, 1] = \bigcup_{k=1}^r A_k$ is a partition into finitely many measurable sets with positive measures, and that we are given a subset $E \subseteq [r] \times [r]$ and real numbers $a_{kl} \in [0, 1]$ for $(k, l) \in E$. Suppose further that the properties above hold and that there is a graphon W representing Γ such that

$$(7.3) \quad W(x, y) = a_{kl} \quad \text{for all } (k, l) \in E \text{ and all } (x, y) \in A_k \times A_l.$$

Then there exists a graphon W representing Γ that is completely G -free for every $G \notin \mathcal{P}$ and is such that (7.3) holds.

Obviously, it suffices to assume that (7.3) holds for a.e. $(x, y) \in A_k \times A_l$, since we can begin by redefining W on a set of measure 0 so that (7.3) holds for all $(x, y) \in A_k \times A_l$.

Proof. The equivalences (i) \iff (ii) \iff (iii) follow, as said above, from [15, Theorem 3.3] and Definition 7.1. The implication (iv) \implies (iii) is trivial. It thus suffices to prove (iii) \implies (iv), and the final statement.

For this, suppose that Γ is represented by a graphon W that is G -free for every $G \notin \mathcal{P}$. Let \mathcal{I} be the set of all pairs $ij = (i, j)$ of distinct positive integers, and for every labeled graph G , define $M_G \subset [0, 1]^{\mathcal{I}}$ by

$$(7.4) \quad M_G = \left\{ (w_{ij}) \in [0, 1]^{\mathcal{I}} : w_{ij} = w_{ji} \text{ for all } i, j \in \mathbb{N} (i \neq j) \right. \\ \left. \text{and } \prod_{ij \in E(G)} w_{ij} \prod_{ij \notin E(G)} (1 - w_{ij}) = 0. \right\}$$

Furthermore, define

$$(7.5) \quad M = \bigcap_{G \notin \mathcal{P}} M_G.$$

If $G \notin \mathcal{P}$, then W is assumed to be G -free, which by (1.3) and (7.4) means that the infinite vector $(W(x_i, x_j))_{ij} \in M_G$ for a.e. sequence $x_1, x_2, \dots \in [0, 1]$ (with the product measure). Since the number of labeled graphs is countable, it follows that $(W(x_i, x_j))_{ij} \in M$ for a.e. $x_1, x_2, \dots \in [0, 1]$.

By the general removal lemma of Petrov [26, Theorem 1(2)] (taking $K = X = [0, 1]$ and $k = 2$ there), there exists a graphon W' such that $W'(x, y) = W(x, y)$ a.e. (so W' also represents Γ) and moreover $(W'(x_i, x_j))_{ij} \in M$ for all $x_1, x_2, \dots \in [0, 1]$. By (7.4)–(7.5) and (1.3), it follows that W' is completely G -free for every $G \notin \mathcal{P}$, which proves (iv).

For the final statement, we may, by a suitable measure-preserving bijection $[0, 1] \rightarrow [0, 1)$, assume that the graphons are defined on $[0, 1)$ and that the sets A_k are half-open intervals $[a_k, b_k)$. We look into the proof by Petrov [26]; there is defined a set $Y \subseteq [0, 1)^2$ consisting of all pairs $(x_1, x_2) \in [0, 1)^2$ such that if U is any open set containing $f(x_1, x_2)$, $I_{x, m} = \{y : \lfloor my \rfloor = \lfloor mx \rfloor\}$ (an interval of length $1/m$ containing x), and μ is Lebesgue measure in the plane, then

$$(7.6) \quad \lim_{m \rightarrow \infty} m^2 \mu \{(y_1, y_2) \in I_{x_1, m} \times I_{x_2, m} : f(y_1, y_2) \in U\} = 1.$$

The construction of W' is such that $W' = W$ on Y . It follows immediately from (7.3) that if $(k, l) \in E$, then $A_k^\circ \times A_l^\circ \subseteq Y$, where $A_k^\circ = (a_k, b_k)$, and thus $W' = W = a_{kl}$ on $A_k^\circ \times A_l^\circ$. Finally, pick $a'_k \in (a_k, b_k)$, and define $\phi : [0, 1) \rightarrow [0, 1)$ by $\phi(a_k) = a'_k$ and $\phi(x) = x$ for $x \notin \{a_1, \dots, a_r\}$. Then $W''(x, y) := W'(\phi(x), \phi(y))$ equals W a.e., is completely G -free for every $G \notin \mathcal{P}$, and satisfies (7.3). \square

Remark 7.3. We derived the last statement in Theorem 7.2 by using the proof by Petrov [26]. It is also possible to use only the statement together with a more complicated construction, which we sketch here: Consider the finite set $B = \{\{i, j\} : i, j \in [r]\}$ of unordered pairs or singletons and the function $g : [0, 1]^2 \rightarrow B$ defined by $g = \{i, j\}$ on $A_i \times A_j$. (We consider unordered pairs $\{i, j\}$ since we need g to be symmetric.) We apply Petrov's theorem to the function $(W(x, y), g(x, y))$ mapping $[0, 1]^2$ into the compact set $K = [0, 1] \times B$, and define $M \subset K^\mathbb{Z} = [0, 1]^\mathbb{Z} \times B^\mathbb{Z}$ by the conditions in (7.4) for all $G \notin \mathcal{P}$ together with conditions corresponding to

- (i) $g(x, y) \cap g(x, z) \neq \emptyset$;
- (ii) $g(x, y) = \{i, j\}$ and $g(y, z) = \{i, k\}$ with $j \neq k \implies g(x, z) = \{j, k\}$;
- (iii) $g(x, y) = \{k, l\}$ and $(k, l) \in E \implies W(x, y) = a_{kl}$.

Petrov's theorem yields W' and g' satisfying corresponding conditions everywhere, and it can be seen that (at least if $r \geq 3$, as we may assume by splitting some A_k) there exists a partition $[0, 1] = \bigcup_k A'_k$ such that $g'(x, y) = \{k, l\}$ and thus $W'(x, y) = a_{kl}$ on $A'_k \times A'_l$, and furthermore $A_k \setminus A'_k$ has measure 0 for each k . Finally we redefine $W'(x, y)$ when x or $y \in A_k \setminus A'_k$ for some k . We omit the details. This argument can also be used when the condition $W(x, y) = a_{kl}$ in (7.3) is replaced by $W(x, y) \in A_{kl}^*$ for some compact sets A_{kl}^* .

Proof of Lemma 1.10. By Theorem 7.2, there exists a graphon W representing Γ that is completely G -free for every $G \notin \mathcal{P}$. If G is W -constructible, then W is not completely G -free, so $G \in \mathcal{P}$.

Moreover, if $\Gamma \in \widehat{\mathcal{P}}^*$ and $r = \text{col}(\mathcal{P})$, then Theorem 1.3 shows that there exists a representing graphon $W \in R_r$. If $r = \infty$, i.e., $\mathcal{P} = \mathcal{U}$, then it is easy to see that the constant graphon $W = 1/2$ possesses the required properties. Otherwise, we can apply the last statement in Theorem 7.2 with $A_k = I_k = [(k-1)/r, k/r)$, $E = \{(k, \ell) : k \neq \ell\}$ and $a_{k\ell} = 1/2$ for $(k, \ell) \in E$. This shows that there exists a representing graphon W' that is completely G -free for every $G \notin \mathcal{P}$ with $W' = 1/2$ on $\bigcup_{i \neq j} I_i \times I_j$. Furthermore, $W' = W \in \{0, 1\}$ a.e. on each $I_i \times I_i$. Define W'' as the modification of W' obtained by letting $W''(x, y) = 0$ if $x, y \in I_i$ for some i , and

$0 < W'(x, y) < 1$, and otherwise $W'' = W'$. Then $W''' = W'$ a.e., so W''' represents Γ , $W''' \in R_r$ and W''' is still completely G -free for every $G \notin \mathcal{P}$. This completes the proof. \square

ACKNOWLEDGMENTS

We would like to thank János Pach and Graham Brightwell for helpful discussions.

REFERENCES

- [1] V.E. Alekseev. Range of values of entropy of hereditary classes of graphs (in Russian). *Diskret. Mat.*, 4:148–157, 1992.
- [2] B. Bollobás. Hereditary and monotone properties of combinatorial structures. In *Surveys in Combinatorics 2007*, volume 346 of *London Math. Soc. Lecture Note Ser.*, pages 1–39. Cambridge Univ. Press, Cambridge, 2007.
- [3] B. Bollobás and A. Thomason. Hereditary and monotone properties of graphs. In R.L. Graham and J. Nešetřil, editors, *The Mathematics of Paul Erdős II*, volume 14 of *Algorithms Combin.*, pages 70–78. Springer Berlin Heidelberg, 1997.
- [4] C. Borgs, J.T. Chayes, L. Lovász, V.T. Sós, and K. Vesztegombi. Convergent sequences of dense graphs I: Subgraph frequencies, metric properties and testing. *Adv. Math.*, 219(6):1801–1851, 2008.
- [5] G. Brightwell, H.J. Prömel, and A. Steger. The average number of linear extensions of a partial order. *J. Combin. Theory Ser. A*, 73(2):193–206, 1996.
- [6] S. Chatterjee and S.R.S. Varadhan. The large deviation principle for the Erdős–Rényi random graph. *European J. Combin.*, 32(7):1000–1017, 2011.
- [7] P. Diaconis, S. Holmes, and S. Janson. Threshold graph limits and random threshold graphs. *Internet Math.*, 5(3):267–320, 2008.
- [8] P. Diaconis and S. Janson. Graph limits and exchangeable random graphs. *Rend. Mat. Appl. (7)*, 28(1):33–61, 2008.
- [9] G. Ehrlich, S. Even, and R.E. Tarjan. Intersection graphs of curves in the plane. *J. Combin. Theory Ser. B*, 21(1):8–20, 1976.
- [10] R. Engelking. *General topology*, volume 6 of *Sigma Ser. Pure Math.* Heldermann Verlag, Berlin, second edition, 1989. Translated from the Polish by the author.
- [11] T. Gallai. Transitiv orientierbare Graphen. *Acta Math. Acad. Sci. Hung.*, 18(1-2):25–66, 1967.
- [12] M.C. Golumbic, D. Rotem, and J. Urrutia. Comparability graphs and intersection graphs. *Discrete Math.*, 43(1):37–46, 1983.
- [13] Ch. Chojnacki (H. Hanani). Über wesentlich unplättbare Kurven im dreidimensionalen Raume. *Fund. Math.*, 23:135–142, 1934.
- [14] H. Hatami, S. Janson, and B. Szegedy. Graph properties, graph limits, and entropy. Preprint, <http://arxiv.org/abs/1312.5626>, 2013.
- [15] S. Janson. Graph limits and hereditary properties. Preprint, <http://arxiv.org/abs/1102.3571v2>, 2013.
- [16] D. Kleitman and B. Rothschild. The number of finite topologies. *Proc. Amer. Math. Soc.*, 25(2):276–282, 1970.
- [17] D.J. Kleitman and B.L. Rothschild. Asymptotic enumeration of partial orders on a finite set. *Trans. Amer. Math. Soc.*, 205:205–220, 1975.
- [18] J. Kratochvíl. String graphs. I. The number of critical nonstring graphs is infinite. *J. Combin. Theory Ser. B*, 52(1):53–66, 1991.
- [19] J. Kratochvíl, M. Goljan, and P. Kučera. *String Graphs*. Academia Praha, Prague, 1986.
- [20] J. Kratochvíl and A. Kuběna. On intersection representations of co-planar graphs. *Discrete Math.*, 178(1–3):251–255, 1998.
- [21] L. Lovász. Perfect graphs. In *Selected Topics in Graph Theory, vol. 2*, pages 55–87. Academic Press, London, 1983.
- [22] L. Lovász. *Large networks and graph limits*, volume 60 of *Amer. Math. Soc. Colloq. Publ.* Amer. Math. Soc., Providence, RI, 2012.
- [23] L. Lovász and B. Szegedy. Limits of dense graph sequences. *J. Combin. Theory Ser. B*, 96(6):933–957, 2006.
- [24] L. Lovász and B. Szegedy. Regularity partitions and the topology of graphons. In *An irregular mind*, volume 21 of *Bolyai Soc. Math. Stud.*, pages 415–446. János Bolyai Math. Soc., Budapest, 2010.
- [25] J. Pach and G. Tóth. How many ways can one draw a graph? *Combinatorica*, 26(5):559–576, 2006.

- [26] F. Petrov. General removal lemma. Preprint, <http://arxiv.org/abs/1309.3795>, 2013.
- [27] F.W. Sinden. Topology of thin film RC circuits. *Bell System Tech. J.*, 45(9):1639–1662, 1966.
- [28] W.T. Trotter. *Combinatorics and Partially Ordered Sets: Dimension Theory*. Johns Hopkins studies in the mathematical sciences. Johns Hopkins Univ. Press, Baltimore, 2001.
- [29] W.T. Tutte. Toward a theory of crossing numbers. *J. Combin. Theory*, 8(1):45–53, 1970.

APPENDIX A. STRING REPRESENTATIONS AND ARCWISE-CONNECTED SETS

We have defined string graphs as graphs having an intersection representation using curves in the plane. It is well-known that the string graphs can also be defined in terms of intersection representations of other types of planar sets. We collect some of these equivalences, and for completeness give a proof.

Lemma A.1. *The following families of subsets of the plane \mathbb{R}^2 define the same class of intersection graphs, i.e., the string graphs:*

- (i) *curves (continuous images of $[0, 1]$),*
- (ii) *simple curves (homeomorphic images of $[0, 1]$),*
- (iii) *simple closed curves (homeomorphic images of S^1),*
- (iv) *piecewise-linear simple closed curves,*
- (v) *pathwise-connected sets,*
- (vi) *arcwise-connected sets,*
- (vii) *open connected sets,*
- (viii) *open simply connected sets.*

Furthermore, in (i)–(iv) we may assume that the intersection representation is such that every pair of strings intersects finitely many times, such that no three curves cross at a single point, and such that if two strings cross, then they cross properly.

Recall that a topological space is *pathwise-connected* if any two points in it can be connected by a curve, and *arcwise-connected* if any two points can be connected by a simple curve. These notions are actually equivalent for Hausdorff spaces (and thus for subsets of the plane), see [10, Problem 6.3.12]. Any open connected set in the plane is pathwise-connected.

Proof. The implications (iv) \implies (iii) \implies (i) \implies (v), (ii) \implies (i), (ii) \implies (vi) \implies (v), and (viii) \implies (vii) \implies (v) are trivial. To complete the proof it thus suffices to show that (v) \implies (ii),(iv) and (iv) \implies (viii).

(v) \implies (i),(ii),(iii),(iv). Suppose that G has an intersection representation consisting of pathwise-connected sets $\{A_v : v \in V(G)\}$. For convenience, we may assume that G has no isolated vertices, since these can be added at the end (or by trivial modifications in the argument below). For each i , choose a point $p_i \in A_i$. For each $j \neq i$ such that $A_i \cap A_j \neq \emptyset$, choose a point $p_{ij} \in A_i \cap A_j$. By hypothesis, for each i and for each $j \neq i$ such that $A_i \cap A_j \neq \emptyset$, we may choose a curve $C_i^j \subseteq A_i$ with endpoints p_i and p_{ij} . (Note that this notation is *not* symmetric: in general, $C_i^j \neq C_j^i$.) Define $A'_i = \bigcup_j C_i^j$. Then $A'_i \subseteq A_i$, each A'_i is pathwise-connected and compact, and $\{A'_i\}$ yields an intersection representation of G . Hence, by replacing A_i by A'_i , we may assume that each set A_i is compact, which implies that there exists $\varepsilon > 0$ such that if $A_i \cap A_j = \emptyset$, then $d(A_i, A_j) \geq 3\varepsilon$. Furthermore, we may at each point p_{ij} add a small line segment (of length less than ε and with p_{ij} as one endpoint) to both A_i and A_j without creating any new intersecting pairs of sets; this guarantees that any pair of sets that intersects will intersect in infinitely many points.

We now start again with these modified sets A_i and choose new points $p_i \in A_i$ and $p_{ij} \in A_i \cap A_j$, for each $j \neq i$ such that $A_i \cap A_j \neq \emptyset$. We choose these points such that $p_{ij} = p_{ji}$ when these points are defined. However, we require that $p_{ij} \neq p_{kl}$

otherwise; that $p_i \neq p_{jk}$ for all i, j , and k ; and that $p_i \neq p_j$ for $i \neq j$. (Note that this is possible by the modifications above.) Once again, choose a curve $C_i^j \subseteq A_i$ with endpoints p_i and p_{ij} . Since $C_i^j \subseteq A_i$, $d(C_i, C_j) \geq \varepsilon$ whenever $A_i \cap A_j = \emptyset$, but $C_i \cap C_j \supseteq \{p_{ij}\}$ if $A_i \cap A_j \neq \emptyset$.

For each i and for each $j \neq i$ such that $A_i \cap A_j \neq \emptyset$, let \tilde{C}_i^j be a piecewise-linear approximation of C_i^j (with the same endpoints p_i and p_{ij} , and at least two linear segments) to within distance $\varepsilon/3$. (That is, for all $x \in \tilde{C}_i^j$, there exists $y \in C_i^j$ such that $d(x, y) < \varepsilon/3$.) Furthermore, if necessary, we may randomly perturb the endpoints of the segments in each \tilde{C}_i^j (except p_i and p_{ij}) such that every such endpoint is contained in only one curve \tilde{C}_i^j , p_i is not contained in any \tilde{C}_k^ℓ with $k \neq i$, p_{ij} is contained only in \tilde{C}_i^j and \tilde{C}_j^i , no point outside of $\{p_i\}$ lies on three curves \tilde{C}_k^ℓ , and all segments have pairwise different directions. Note that our choice of ε means that if $A_i \cap A_j = \emptyset$, then for any k and ℓ , we have $\tilde{C}_i^k \cap \tilde{C}_j^\ell = \emptyset$.

For each i and for each j such that \tilde{C}_i^j is defined, let $N = N_i^j$ denote the number of segments in the curve \tilde{C}_i^j . Let $s_i^j(m)$ denote the m th segment in order of increasing distance from p_i . Furthermore, let $p_i = x_i^j(0), \dots, x_i^j(N) = p_{ij}$ denote the sequence of endpoints of segments in the curve \tilde{C}_i^j , so that $x_i^j(m-1)$ and $x_i^j(m)$ are the endpoints of $s_i^j(m)$.

For each i , for each j such that \tilde{C}_i^j is defined, and for $m \in \{0, \dots, N\}$, let $u_i^j(m)$ be a segment of length η centered at $x_i^j(m)$, where $\eta < \varepsilon/6$ is a small number. If $m \in \{1, \dots, N-1\}$, then we define $u_i^j(m)$ to lie on the angle-bisector of $s_i^j(m)$ and $s_i^j(m+1)$. We define $u_i^j(0)$ to be perpendicular to $s_i^j(1)$ and $u_i^j(N)$ to be perpendicular to $s_i^j(N)$. For each m , let $x_i^j(m)^+$ be the right-hand endpoint of $u_i^j(m)$ as \tilde{C}_i^j is oriented from p_i to p_{ij} , and let $x_i^j(m)^-$ denote the left-hand endpoint. Let $T_i^j(m)$ denote the trapezoid with vertices $x_i^j(m-1)^+$, $x_i^j(m-1)^-$, $x_i^j(m)^-$, and $x_i^j(m)^+$.

For each i and for each j such that \tilde{C}_i^j is defined, let $B_i^j = \cup_{m=1}^N T_i^j(m)$. Note that the choices of ε and η still ensure that if $A_i \cap A_j = \emptyset$, then for any k and ℓ , we have $B_i^k \cap B_j^\ell = \emptyset$. If η is small enough, then the boundary $D_i^j = \partial B_i^j$ is a piecewise-linear simple closed curve, and no point lies on three of these curves, except that p_i lies on D_i^j for each j such that this curve is defined; making a small random perturbation of the segments through p_i , we may assume that no point at all lies on three of these curves, while D_i^j still intersects D_i^k for any j and k such that these curves are defined. Note also that $p_{ij} \in D_i^j$. Furthermore, if necessary after another random perturbation of the segments through p_{ij} , all segments have pairwise different directions and since each string consists of finitely many segments, all pairs of curves D_i^j have finitely many intersection points. This and the fact that every endpoint is contained in only one such curve (if η is small enough) means that if two curves cross, then they cross properly.

For each i , let $D_i = \bigcup_j D_i^j$, where once again the union is taken over all j for which B_i^j is defined. Once again, $D_i \cap D_j \neq \emptyset$ if and only if $A_i \cap A_j \neq \emptyset$. Each D_i is a union of polygonal curves, and D_i is connected. Furthermore, viewing D_i as a graph, each vertex has degree 2 or 4, so the graph is Eulerian. Moreover, viewing an Eulerian circuit as a directed curve, it is easy to see that the graph has an Eulerian circuit that does not properly cross itself. Hence, making a small modification at each point where D_i intersects itself, we may replace D_i by a piecewise-linear simple closed curve γ_i , without creating or destroying any other intersections.

This yields an intersection representation $\{\gamma_i\}$ of G consisting of piecewise-linear simple closed curves (as in (iv)), which furthermore satisfies the properties in the

final part of the statement. By choosing a point q_i on each γ_i , not on any other of the curves, and deleting a small open interval about q_i , we can replace the closed curves γ_i by simple curves γ'_i as in (ii). This completes the proof that (v) \implies (i),(ii),(iii),(iv).

(iv) \implies (viii): Let $\{A_i\}$ be an intersection representation of G consisting of piecewise-linear simple curves. Extend the first and last segments a tiny bit η past the original endpoints. Then, similarly to the proof above, define a region $B_i \supset A_i$ as the union of small trapezoids of width at most η surrounding the segments in A_i . Then, if η is small enough, the interiors B_i° form an intersection representation of G consisting of open simply connected sets. \square

A similar result holds for outer-string graphs. We give one version, leaving further variations to the reader.

Lemma A.2. *Let D be a closed disk in the plane \mathbb{R}^2 . The following families of subsets of D define the same class of intersection graphs, i.e., the outer-string graphs:*

- (i) *curves with one endpoint in ∂D ,*
- (ii) *simple curves with one endpoint in ∂D ,*
- (iii) *piecewise-linear simple curves with one endpoint in ∂D ,*
- (iv) *pathwise-connected sets A with $A \cap \partial D \neq \emptyset$,*

Furthermore, in (i)–(iii) we may assume that both endpoints of the strings are on the boundary ∂D , but that otherwise the strings lie in the interior D° , and that the intersection representation is such that every pair of strings intersects finitely many times, such that no three curves cross at a single point, and such that if two strings cross, then they cross properly.

Proof. The implications (iii) \implies (ii) \implies (i) \implies (iv) are trivial.

(iv) \implies (iii): Suppose that G has an intersection representation $\{A_i\}$ with pathwise-connected sets as in (iv), and argue as in the proof of Lemma A.1, now choosing $p_i \in \partial D$. (If necessary we may add small arcs of ∂D to the A_i in order to get all p_i distinct.) The construction above yields piecewise-linear closed curves γ_i lying in the interior of a slightly larger disk D' . Let q_i be the point furthest from the center of D where the radius through p_i intersects γ_i (such a point exists, and is close to p_i); we may then remove a small interval about q_i from γ_i and replace it by two line segments to the boundary $\partial D'$. This yields an intersection representation as in (iii), which further satisfies the conditions in the final statement. \square

DEPARTMENT OF MATHEMATICS, UPPSALA UNIVERSITY, P.O. BOX 480, SE-751 06 UPPSALA, SWEDEN

E-mail address: svante.janson@math.uu.se

E-mail address: andrew.uzzell@math.uu.se