arXiv:1512.08130v1 [math.CO] 26 Dec 2015

Extracting list colorings from large independent sets

H. A. Kierstead and Landon Rabern^{*}

September 10, 2018

Abstract

We take an application of the Kernel Lemma by Kostochka and Yancey [10] to its logical conclusion. The consequence is a sort of magical way to draw conclusions about list coloring (and online list coloring) just from the existence of an independent set incident to many edges. We use this to prove an Ore-degree version of Brooks' Theorem for online list-coloring. The Ore-degree of an edge xy in a graph G is $\theta(xy) = d_G(x) + d_G(y)$. The Ore-degree of G is $\theta(G) = \max_{xy \in E(G)} \theta(xy)$. We show that every graph with $\theta \ge 18$ and $\omega \le \frac{\theta}{2}$ is online $\lfloor \frac{\theta}{2} \rfloor$ -choosable. In addition, we prove an upper bound for online list-coloring triangle-free graphs: $\chi_{OL} \le \Delta + 1 - \lfloor \frac{1}{4} \lg(\Delta) \rfloor$. Finally, we characterize Gallai trees as the connected graphs G with no independent set incident to at least |G| edges.

1 Introduction

In [10], Kostochka and Yancey applied the Kernel Lemma to a coloring problem in a novel manner. Our Main Lemma generalizes and strengthens their idea. The basic idea is that given an independent set that is incident to many edges, we can find a reducible configuration. In this way, we can reduce coloring problems to the mere existence of a large independent set. Before stating the Main Lemma we need to introduce some notation, and define the concepts of f-choosable and online f-choosable graphs.

Let G = (V, E) be a graph. We write |G| for |V| and ||G|| for |E|. For $A, B \subseteq V$, put $||A, B||_G := \sum_{v \in A} |N_G(v) \cap B|$. Also, put $||A||_G := ||G[A]||$. Note that $||A, B||_G =$ $||A \setminus B, B \setminus A||_G + 2 ||A \cap B||_G$. In particular, $||A, B||_G = ||B, A||_G$ and $||V, V||_G = 2 ||V||_G$. When the graph G is clear from context we drop the G from the notation and write $||A, B||_G$ and ||A||. For $v \in V$, let $N[v] = N(v) \cup \{v\}$.

Let G = (V, E) be a graph. A list assignment on G is a function L from V to the subsets of \mathbb{N} . A graph G is L-colorable if there is $\pi: V \to \mathbb{N}$ such that $\pi(v) \in L(v)$ for each $v \in V$ and $\pi(x) \neq \pi(y)$ for each $xy \in E$. For $f: V \to \mathbb{N}$, a list assignment L is an f-assignment if |L(v)| = f(v) for each $v \in V$. We say that G is f-choosable if G is L-colorable for every f-assignment L.

^{*}School of Mathematical and Statistical Sciences, Arizona State University

Online choosability was independently introduced by Zhu [21] and Schauz [18] (Schauz called it *paintability*). Let G = (V, E) be a graph and $f: V \to \mathbb{N}$. We say that G is online f-choosable if |G| = 0 or both $f(v) \ge 1$ for all $v \in V$ and for every $S \subseteq V$ there is an independent set $I \subseteq S$ such that G - I is online f'-choosable where f'(v) := f(v) for $v \in V \setminus S$ and f'(v) := f(v) - 1 for $v \in S \setminus I$.

Observe that if G is online f-choosable then it is f-choosable: For any f-list assignment L on G, let $v \in V(G)$, $\alpha \in L(v)$, and $S = \{x \in V(G) : \alpha \in L(x)\}$. As G is online f-choosable, there is an independent set $I \subseteq S$ with G - I online f'-choosable. Color each vertex of I with α , and let L' be the f'-list assignment on G - I obtained by removing α from every list. Arguing inductively (with a trivial basis) yields an L'-coloring of G - I, and an L-coloring of G.

When f(v) := k - 1 for all $v \in V(G)$, we say that G is online k-list-critical if G is not online f-choosable, but every proper subgraph H of G is online $f|_{V(H)}$ -choosable.

Main Lemma. Let G = (V, E) be a nonempty graph and $f: V \to \mathbb{N}$ with $f(v) \leq d_G(v) + 1$ for all $v \in V$. If there is an independent $A \subseteq V$ such that

$$||A, V|| \ge \sum_{v \in V} d_G(v) + 1 - f(v),$$

then G has a nonempty induced subgraph H that is (online) f_H -choosable where $f_H(v) := f(v) + d_H(v) - d_G(v)$ for $v \in V(H)$.

As a simple first application, we show that Brooks' Theorem can be derived from the simple bound on the independence number it implies. In particular, the proof shows that Brooks' Theorem for list coloring [20] and online list coloring [7] follow from Brooks' Theorem for ordinary coloring.

Brooks' Theorem for Independence Number. If G is a graph with $\Delta := \Delta(G) \ge 3$ and $K_{\Delta+1} \not\subseteq G$, then $\alpha(G) \ge \frac{|G|}{\Delta}$.

To prove Brooks' Theorem, consider a minimal counterexample G. By minimality G is regular: if $d(x) < \Delta$ then G - x has a Δ -coloring, and some color is used on no neighbor of x, a contradiction. Hence for any maximum independent set A in G, Brooks' Theorem for Independence Number gives $||A, V(G)|| \ge |A|\Delta \ge |G|$. Applying the Main Lemma with $f(v) := d_G(v)$ gives a nonempty induced subgraph H of G that is (online) d_H -choosable. So, after Δ -coloring G - H by minimality of G, we can finish the coloring on H, a contradiction.

A bound like Brooks' Theorem in terms of the Ore-degree was given by Kierstead and Kostochka [8] and subsequently the required lower bound on Δ was improved in [15, 12, 17]. For example, we have the following.

Definition 1. The Ore-degree of an edge xy in a graph G is $\theta(xy) := d(x) + d(y)$. The Ore-degree of a graph G is $\theta(G) := \max_{xy \in E(G)} \theta(xy)$.

Theorem 1.1. Every graph with $\theta \ge 10$ and $\omega \le \frac{\theta}{2}$ is $\lfloor \frac{\theta}{2} \rfloor$ -colorable.

Another method for achieving the tightest of these results on Ore-degree was given by Kostochka and Yancey [10]. Their proof combined their new lower bound on the number of edges in a color critical graph together with their list coloring lemma derived via the kernel lemma. The Main Lemma improves this latter lemma and, in a similar way, we use it in combination with our lower bound on the number of edges in online list-critical graphs [9] to prove an Ore-degree version of Brooks' Theorem for online list coloring.

Now we introduce the key graph theoretic parameter for this paper.

Definition 2. The maximum independent cover number of a graph G is the maximum mic(G) of $\sum_{v \in I} d_G(v)$ (= ||I, V(G)||) over all independent sets I of G. A set I that witnesses this maximum is said to be *optimal*.

We work in terms of a class of graphs that is more general than the class of online k-listcritical graphs. Basically, we want graphs G that have no induced subgraph H such that every online (k-1)-list-coloring of G-H can be extended to H. We can replace k-1 with $\delta(G)$ and still get a generalization of online k-list-critical since an online k-list-critical graph has minimum degree at least k-1. Doing so, we get the following graph class that does not depend on k, but still does everything we need.

Definition 3. A graph G is OC-reducible to H if H is a nonempty induced subgraph of G which is online f_H -choosable where $f_H(v) := \delta(G) + d_H(v) - d_G(v)$ for all $v \in V(H)$. If G is not OC-reducible to any nonempty induced subgraph, then it is OC-irreducible.

The Main Lemma can be used to give another lower bound on the number of edges in a critical graph G. Viewed differently, it gives an upper bound on mic(G).

Theorem 2.4. Every OC-irreducible graph G satisfies $mic(G) \le 2 ||G|| - (\delta(G) - 1) |G| - 1$.

This quickly gives the aforementioned Ore degree version of Brooks' Theorem for list coloring.

Theorem 3.13. Every graph with $\theta \ge 18$ and $\omega \le \frac{\theta}{2}$ is $\left|\frac{\theta}{2}\right|$ -choosable.

Note that using Kostochka and Stiebitz's lower bound on the number of edges in a list critical graph [13] gives a weakened version of Theorem 3.13 with $\theta \ge 54$ instead of $\theta \ge 18$. Similarly, we get the online version.

Theorem 3.12. Every graph with $\theta \geq 18$ and $\omega \leq \frac{\theta}{2}$ is online $\left|\frac{\theta}{2}\right|$ -choosable.

We expect that Theorems 3.12 and 3.13 actually hold for $\theta \ge 10$. In the regular coloring case, it was shown in [12] that the only exception when $\theta \ge 8$ is the graph O_5 ; where O_n is the graph formed from the disjoint union of $K_n - xy$ and K_{n-1} by joining $\lfloor \frac{n-1}{2} \rfloor$ vertices of the K_{n-1} to x and the other $\lceil \frac{n-1}{2} \rceil$ vertices of the K_{n-1} to y. Again, the expectation is that the same result will hold for Theorems 3.12 and 3.13.

A simple probabilistic argument gives a reasonable bound on mic(G) for triangle-free graphs and we get the following.

Corollary 5.3. Triangle-free graphs are online $(\Delta + 1 - \lfloor \frac{1}{4} \lg(\Delta) \rfloor)$ -choosable.

In the final section, we characterize Gallai trees as the connected graphs G that have mic(G) = |G| - 1, and some further characterizations.

2 Proving the Main Lemma

A kernel in a digraph D is an independent set $I \subseteq V(D)$ such that each vertex in $V(D) \setminus I$ has an edge into I. A digraph in which every induced subdigraph has a kernel is kernel-perfect.

Kernel Lemma. Let G = (V, E) be a graph and $f: V \to \mathbb{N}$. If G has a kernel-perfect orientation such that $f(v) \ge d^+(v) + 1$ for each $v \in V$, then G is online f-choosable.

Lemma 2.1 (Kostochka and Yancey [10]). Let A be an independent set in a graph G and let $B := V(G) \setminus A$. Any digraph D created from G by replacing each edge in G[B] by a pair of opposite arcs and orienting the edges between A and B arbitrarily is kernel-perfect.

Proof. Let G be a minimum counterexample, and let D be a digraph created from G that is not kernel-perfect. To get a contradiction it suffices to construct a kernel in D, since each subdigraph has a kernel by minimality. Either A is a kernel or there is some $v \in B$ which has no outneighbors in A. In the latter case, each neighbor of v in G has an inedge to v, so a kernel in D - v - N(v) together with v is a kernel in D.

The following lemma is folklore and can be derived from Hall's theorem by vertex splitting. It also follows by taking an arbitrary orientation and repeatedly reversing paths if doing so gets a gain.

Lemma 2.2. Let G = (V, E) be a graph and $g: V \to \mathbb{N}$. Then G has an orientation such that $d^{-}(v) \ge g(v)$ for all $v \in V$ if and only if for every $X \subseteq V$, we have

$$||X|| + ||X, V \smallsetminus X|| \ge \sum_{v \in X} g(v).$$

For independent $A \subseteq V(G)$, we write G_A for the bipartite subgraph G - E(G - A) of G. Thus G is the edge-disjoint union of G_A and $G[V(G) \smallsetminus A]$.

Lemma 2.3. Let G = (V, E) be a graph and $f: V \to \mathbb{N}$ with $f(v) \leq d_G(v) + 1$ for all $v \in V$. If there is an independent $A \subseteq V$ such that each $X \subseteq V(G_A)$ satisfies

$$||X||_{G_A} + ||X, V(G_A) \setminus X||_{G_A} \ge \sum_{v \in X} (d_G(v) + 1 - f(v)).$$

then G is online f-choosable.

Proof. Applying Lemma 2.2 on G_A with $g(v) := d_G(v) + 1 - f(v)$ for all $v \in V(G_A)$ gives an orientation of G_A where $d^-(v) \ge d_G(v) + 1 - f(v)$ for each $v \in V(G_A)$. Make an orientation D of G by using this orientation of G_A for the edges between A and $V(G) \smallsetminus A$ and replacing each edge in G - A by a pair of opposite arcs. For $v \in V(D)$, where $d_{G-A}(v) = 0$ if $v \in A$,

$$d^+(v) \le d_{G-A}(v) + d_{G_A}(v) - (d_G(v) + 1 - f(v)) = f(v) - 1,$$

so $f(v) \ge d^+(v) + 1$. By Lemma 2.1, D is kernel-perfect, so the Kernel Lemma implies G is online f-choosable.

Proof of Main Lemma. Let $A \subseteq V$ be an independent set with

$$||A, V|| \ge \sum_{v \in V} (d_G(v) + 1 - f(v)).$$

Choose a nonempty induced subgraph H of G with $||H_A|| \ge \sum_{v \in V(H)} (d_H(v) + 1 - f_H(v))$ minimizing |H| (we can make this choice since G is a such a subgraph). Suppose H is not online f_H -choosable. Then, by Lemma 2.3, we have $X \subseteq V(H_A)$ with

$$||X||_{H_A} + ||X, V(H_A) \setminus X||_{H_A} < \sum_{v \in X} (d_H(v) + 1 - f_H(v)).$$

Now $X \neq V(H)$ by our assumption on $||H_A||$, hence Z := H - X is a nonempty induced subgraph of G with

$$\begin{split} \|Z_A\| &= \|H_A\| - \|X\|_{H_A} - \|X, V(H_A) \smallsetminus X\|_{H_A} \\ &> \sum_{v \in V(H)} \left(d_H(v) + 1 - f_H(v) \right) - \sum_{v \in X} \left(d_H(v) + 1 - f_H(v) \right) \\ &= \sum_{v \in V(Z)} \left(d_Z(v) + 1 - f_Z(v) \right), \end{split}$$

contradicting the minimality of |H|.

As a special case we get:

Theorem 2.4. Every OC-irreducible graph G = (V, E) satisfies

$$\min(G) \le 2 \|G\| - (\delta(G) - 1) |G| - 1.$$

Proof. As G is OC-irreducible, it has no proper, induced, online f_H -choosable subgraph H, where $f_H(v) = \delta(G) + d_H(v) - d_G(v)$. Let $A \subseteq V$ be an optimal set. By the Main Lemma

$$\operatorname{mic}(G) = \|A, V\| < \sum_{v \in V} (d_G(v) + 1 - \delta(G)) = 2 \|G\| + |G|(1 - \delta(G)).$$

3 Ore version of Brooks' Theorem for (online) list coloring

For a graph G, let $\mathcal{H}(G)$ be the subgraph of G induced on the vertices of degree greater than $\delta(G)$ and $\mathcal{L}(G)$ the subgraph of G induced on the vertices of degree $\delta(G)$.

Lemma 3.1. All OC-irreducible graphs G with $\mathcal{H}(G)$ edgeless and $\Delta(G) = \delta(G) + 1$ satisfy

1.
$$mic(G) < |\mathcal{H}(G)| + |G|$$
, and

2.
$$(\delta - 1) |\mathcal{H}(G)| < |\mathcal{L}(G)|$$
, and $2 ||G|| < (\delta(G) + \frac{1}{\delta(G)}) |G|$.

Proof. Put $\delta := \delta(G)$ and $\Delta := \Delta(G)$. As $\Delta = \delta + 1$, $2 ||G|| = \delta |G| + |\mathcal{H}(G)|$. As $\mathcal{H}(G)$ is edgeless, and using Theorem 2.4,

$$(\delta + 1) |\mathcal{H}(G)| \le \operatorname{mic}(G) < 2 ||G|| + |G| (1 - \delta) = |\mathcal{H}(G)| + |G|.$$

Thus (1) holds, and $\delta |\mathcal{H}(G)| < |G|, (\delta - 1) |\mathcal{H}(G)| < |\mathcal{L}(G)|, \text{ and } 2 ||G|| < (\delta + 1/\delta) |G|.$

To break up our computations we reformulate Lemma 3.1 as an upper bound on σ where

$$\sigma(G) := \left(\delta(G) - 1 + \frac{2}{\delta(G)}\right) |\mathcal{L}(G)| - 2 \|\mathcal{L}(G)\|.$$

Lemma 3.2. If G is an OC-irreducible graph such that $\mathcal{H}(G)$ is edgeless and $\Delta(G) = \delta(G) + 1$, then $\sigma(G) < \left(4 - \frac{2}{\delta(G)}\right) |\mathcal{H}(G)|$.

Proof. Put $\delta := \delta(G)$. As $(1+\delta) |\mathcal{H}(G)| = ||\mathcal{H}(G), \mathcal{L}(G)|| = \delta |\mathcal{L}(G)| - 2 ||\mathcal{L}(G)||$,

$$\sigma(G) = \left(\delta - 1 + \frac{2}{\delta}\right) |\mathcal{L}(G)| + (1 + \delta) |\mathcal{H}(G)| - \delta |\mathcal{L}(G)|$$

$$= \left(\frac{2}{\delta} - 1\right) |\mathcal{L}(G)| + (1 + \delta) |\mathcal{H}(G)|$$

$$< \left(\frac{2}{\delta} - 1\right) (\delta - 1) |\mathcal{H}(G)| + (1 + \delta) |\mathcal{H}(G)| \qquad \text{(Lemma 3.1.2)}$$

$$= \left(4 - \frac{2}{\delta}\right) |\mathcal{H}(G)|.$$

We need the following bound from [9]. Put $\alpha_k := \frac{1}{2} - \frac{1}{(k-1)(k-2)}$ and let c(G) be the number of components in G.

Corollary 3.3. If G is an OC-irreducible graph with $\delta(G) \ge 6$ and $\omega(G) \le \delta(G)$ such that $\mathcal{H}(G)$ is edgeless, then $\sigma(G) \ge (\delta(G) - 2)\alpha_{\delta(G)+1} |\mathcal{H}(G)| + 2(1 - \alpha_{\delta(G)+1})c(\mathcal{L}(G)).$

By combining Lemma 3.2 with Corollary 3.3 we can prove the Ore version of Brooks' Theorem for online list coloring for $\Delta \ge 11$. With a bit more work we will improve this to $\Delta \ge 10$. First, we can squeeze a bit more out of Theorem 2.4 by considering independent sets of low vertices that have no high neighbors. Such sets can be added to $V(\mathcal{H}(G))$ to get a cut with more edges. To apply this idea we need the following counting lemma. For a graph G and $t \in \mathbb{N}$, let $V_t = \{v \in V(G) : d(v) = t\}$, $G_t = G[V_t]$ and $\beta_t = \alpha(G_t)$.

To apply this idea, we need to better understand the structure of $\mathcal{L}(G)$ when G is OCirreducible. A *Gallai tree* is a graph such that each block is a clique or odd cycle. A *Gallai forest* is a disjoint union of Gallai trees. A classical result of Gallai [6] says that if G is a k-critical graph, then $\mathcal{L}(G)$ is a Gallai forest. Borodin [3] and independently Erdős, Rubin and Taylor [5] generalized Gallai's result to show that a connected graph H is f-choosable where $f(v) = d_H(v)$ for all $v \in V(H)$ if and only if H is not a Gallai tree. In [7], Hladkỳ, Král and Schauz extended this result to online f-choosability. **Lemma 3.4** (Hladkỳ, Král and Schauz). A connected graph H is online f-choosable where $f(v) = d_H(v)$ for all $v \in V(H)$ if and only if H is not a Gallai tree.

In Section 6, we give another proof of Lemma 3.4 using our Main Lemma. We need Gallai's result for OC-irreducible graphs.

Lemma 3.5. If G is OC-irreducible, then $\mathcal{L}(G)$ is a Gallai forest.

Proof. Let G be an OC-irreducible graph and H a component of $\mathcal{L}(G)$. Since G is OCirreducible, H is not online f_H -choosable where $f_H(v) := \delta(G) + d_H(v) - d_G(v)$ for all $v \in V(H)$. But $d_G(v) = \delta(G)$ for $v \in V(H)$, so $f_H(v) = d_H(v)$ for all $v \in V(H)$. By Lemma 3.4, H is a Gallai tree.

Lemma 3.6. Fix $k \ge 6$. Let G be a Gallai forest with maximum degree at most k - 1 not containing K_k . We have the following inequality:

$$(k-1)\beta_{k-1}(G) + \sum_{v \in V(G)} k - 1 - d(v) \ge \frac{2(k-3)}{k-2} |G| - \frac{(k-1)(k-4)}{k-2} c(G).$$

Proof. It will suffice to prove that for any Gallai tree T with maximum degree at most k-1 we have:

$$(k-1)\beta_{k-1}(T) + \sum_{v \in V(T)} (k-1-d(v)) \ge \frac{2(k-3)}{k-2} |T| - \frac{(k-1)(k-4)}{k-2}$$

Suppose not and choose a counterexample T minimizing |T|. First, if T has only one block it is easy to see that the inequality is satisfied. Let B be an endblock of T and say xis the cutvertex in B. Suppose $\chi(B) \leq k-3$. Put $T^* := T - (B-x)$. By minimality of |T|, T^* satisfies the inequality. Adding B - x back increases the left side by $(k - \chi(B)) |B - x| ||x, B - x|| \geq 2(|B| - 1)$, but only increases the right side by $\frac{2(k-3)}{k-2}(|B| - 1)$, so T is not a counterexample, a contradiction. Thus B is either K_{k-2} or K_{k-1} .

Consider T' := T - B. First suppose $d_T(x) = k - 1$. Note that none of x's neighbors in T' have degree k - 1 in T' and thus are in no maximum independent set of degree k - 1 vertices in T'. Therefore, we can add x to any such independent set, giving $\beta_{k-1}(T) > \beta_{k-1}(T')$. Hence, after applying minimality to T', we see that adding back B increases the left side by k - 1 + |B - x| - ||x, T'|| = 2k - 4 if B is K_{k-1} and by k - 1 + 2|B - x| - ||x, T'|| = 3k - 9 if B is K_{k-2} . Since the right side increases by only $\frac{2(k-3)}{k-2}|B|$ in both cases, T satisfies the inequality, a contradiction.

Otherwise, $d_T(x) \leq k-2$. So B is K_{k-2} and $d_T(x) = k-2$. Now adding B back, increases the left side by 2(k-3) + 1 - 1 and increases the right side by only 2(k-3), so again T satisfies the inequality, a contradiction.

Lemma 3.7. If G is an OC-irreducible graph such that $\mathcal{H}(G)$ is edgeless, $\Delta(G) = \delta(G) + 1 \ge 7$ and $K_{\Delta(G)} \not\subseteq G$, then $|\mathcal{H}(G)| < \frac{\delta(G)(\delta(G)-3)}{(\delta(G)-1)(\delta(G)-5)}c(\mathcal{L}(G)).$

Proof. Put $\delta := \delta(G)$. By Lemma 3.1.1 we have

$$\operatorname{mic}(G) < |\mathcal{L}(G)| + 2|\mathcal{H}(G)| < |\mathcal{L}(G)| + \frac{2}{\delta - 1}|\mathcal{L}(G)| = \frac{\delta + 1}{\delta - 1}|\mathcal{L}(G)|.$$

By Lemma 3.5, $\mathcal{L}(G)$ is a Gallai forest. Pick $I \subseteq V_{k-1}$ with $|I| = \beta_{k-1}(G)$, and set $J = I \cup \mathcal{H}(G)$. As J is independent, applying Lemma 3.6 to the Gallai forest $\mathcal{L}(G)$ gives

$$\operatorname{mic}(G) \geq \|J, V(G)\| \geq \delta\beta_{k-1}(\mathcal{L}(G)) + \sum_{v \in V(\mathcal{L}(G))} \delta - d_{\mathcal{L}(G)}(v)$$
$$\geq 2\frac{\delta - 2}{\delta - 1} |\mathcal{L}(G)| - \frac{\delta(\delta - 3)}{\delta - 1} c(\mathcal{L}(G)).$$

By Lemma 3.1.2, $|\mathcal{L}(G)| > (\delta - 1) |\mathcal{H}(G)|$. Combining this with the inequalities above proves the lemma.

Lemma 3.8. Every OC-irreducible graph G with $\delta(G) + 1 = \Delta(G) \ge 10$ such that $\mathcal{H}(G)$ is edgeless contains $K_{\Delta(G)}$.

Proof. Suppose not and let G be a counterexample. Put $\delta := \delta(G)$. Then

$$\left(4 - \frac{2}{\delta}\right) |\mathcal{H}(G)| > \sigma(G) \qquad (\text{Lemma 3.2})$$

$$\geq (\delta - 2)\alpha_{\delta+1} |\mathcal{H}(G)| + 2(1 - \alpha_{\delta+1})c(\mathcal{L}(G)) \qquad (\text{Corollary 3.3})$$

$$\geq |\mathcal{H}(G)| \left((\delta - 2)\alpha_{\delta+1} + 2(1 - \alpha_{\delta+1})\frac{(\delta - 1)(\delta - 5)}{\delta(\delta - 3)} \right) \qquad \text{(Lemma 3.7)}$$
$$4 - \frac{2}{\delta} > \alpha_{\delta+1}(\delta - 2) + 2(1 - \alpha_{\delta+1})\frac{(\delta - 1)(\delta - 5)}{\delta(\delta - 3)}.$$

Thus $\delta \leq 8$, a contradiction.

The following lemma from [18] allows us to infer online list colorability of the whole from online list colorability of parts.

Lemma 3.9. Let G be a graph and $f: V(G) \to \mathbb{N}$. If H is an induced subgraph of G such that G - H is online $f|_{V(G-H)}$ -choosable and H is online f_H -choosable where $f_H(v) := f(v) + d_H(v) - d_G(v)$, then G is online f-choosable.

Theorem 3.10. If G is a graph with $\Delta(G) \geq 10$ not containing $K_{\Delta(G)}$ such that $\mathcal{H}(G)$ is edgeless, then G is online $(\Delta(G) - 1)$ -choosable.

Proof. Suppose not and choose a counterexample G minimizing |G|. Then G is online f-critical where $f(v) := \Delta(G) - 1$ for all $v \in V(G)$. Hence $\delta(G) \ge \Delta(G) - 1$ and we may apply Lemma 3.8 to get a nonempty induced subgraph H of G that is online f_H -choosable where $f_H(v) := \Delta(G) - 1 + d_H(v) - d_G(v)$ for all $v \in V(H)$. But then applying Lemma 3.9 shows that G is $(\Delta(G) - 1)$ -choosable, a contradiction.

Combining Theorem 3.10 with the following version of Brooks' Theorem for online list coloring (first proved in [7]) we get Theorem 3.12.

Lemma 3.11. Every graph with $\Delta \geq 3$ not containing $K_{\Delta+1}$ is online Δ -choosable.

Theorem 3.12. Every graph with $\theta \ge 18$ and $\omega \le \frac{\theta}{2}$ is online $\lfloor \frac{\theta}{2} \rfloor$ -choosable.

Proof. Suppose not and choose a counterexample G minimizing |G|. Put $k := \lfloor \frac{\theta(G)}{2} \rfloor$. Then G is online f-critical where f(v) := k for all $v \in V(G)$. Hence $\delta(G) \ge k$ and thus $\Delta(G) \le k + 1$. If $\Delta(G) = k$, then the theorem follows from Lemma 3.11. Hence we must have $\Delta(G) = k + 1$. Therefore $\mathcal{H}(G)$ is edgeless, $\Delta(G) \ge 10$ and $\omega(G) \le \Delta(G) - 1$. Applying Theorem 3.10 shows that G is online $(\Delta(G) - 1)$ -choosable, a contradiction.

The same result for list coloring is an immediate consequence.

Theorem 3.13. Every graph with $\theta \ge 18$ and $\omega \le \frac{\theta}{2}$ is $\lfloor \frac{\theta}{2} \rfloor$ -choosable.

4 Ore Brooks for maximum degree four

Kostochka and Yancey's bound [10] shows that if G is 4-critical, then $||G|| \ge \left\lceil \frac{5|G|-2}{3} \right\rceil$. If we try to analyze 4-critical graphs with edgeless high vertex subgraphs by putting this lower bound on the number of edges together with the results on orientations and list coloring obtained in [10], the bounds miss each other. Using the improved bound from Lemma 3.1 we get an exact bound on the number of edges in such a graph.

Lemma 4.1. For a critical graph G with $\Delta(G) \leq \chi(G) = 4$ such that $\mathcal{H}(G)$ is edgeless we have $||G|| = \left\lceil \frac{5|G|-2}{3} \right\rceil$ and |G| is not a multiple of 3.

Proof. Since G is 4-critical, applying Lemma 3.1.1 gives $2 ||G|| < (3 + \frac{1}{3}) |G| = \frac{10}{3} |G|$. By Kostochka and Yancey's bound we have $\left\lceil \frac{5|G|-2}{3} \right\rceil \le ||G|| < \frac{5}{3} |G|$. Hence $||G|| = \left\lceil \frac{5|G|-2}{3} \right\rceil$ and |G| is not a multiple of 3.

It is easy to see that contracting a diamond in a critical graph G with $\Delta(G) \leq \chi(G) = 4$ such that $\mathcal{H}(G)$ is edgeless gives another such graph. So, the following characterization of these graphs is natural. Recently, Postle [14] proved this using an extension of the potential method of Kostochka and Yancey.

Theorem 4.2 (Postle [14]). Every critical graph G with $\Delta(G) \leq \chi(G) = 4$ such that $\mathcal{H}(G)$ is edgeless, except K_4 , has an induced diamond. In particular, any such G can be reduced to K_4 by a sequence of diamond contractions.

5 Online choosability of triangle-free graphs

We write lg(x) for the base 2 logarithm of x. We can get a reasonably good lower bound on mic(G) for triangle-free graphs using a simple probabilistic technique of Shearer and its modification by Alon (see [1]).

Lemma 5.1. Every triangle-free graph G = (V, E) satisfies $mic(G) \ge \frac{1}{4} \sum_{v \in V} \lg(d(v))$.

Proof. Let W be a random independent set in G chosen uniformly from all independent sets in G. It suffices to show that $E(||W, V||) \ge \frac{1}{4} \sum_{v \in V} \lg(d(v))$. For each $v \in V$ put

$$X_v := \begin{cases} d(v) & \text{if } v \in W \\ \|v, W\| & \text{if } v \notin W \end{cases}$$

Then $||W, V|| = \frac{1}{2} \sum_{v \in V} X_v$. By linearity of expectation it suffices to prove

$$E(X_v) \ge \frac{1}{2} \lg(d(v)). \tag{1}$$

To prove (1), let $H := G[V \setminus N[v]]$, fix an independent set S in H, and set $X := N(v) \setminus N(S)$. Put x := |X|. It suffices to prove that all such S satisfy

$$E(X_v \mid W \cap V(H) = S) \ge \frac{\lg(d(v))}{2}.$$
(2)

Suppose (2) fails for S. As G is triangle-free, the independent sets W with $W \cap V(H) = S$ are exactly $S \cup \{v\}$ and $S \cup X_0$ where $X_0 \subseteq X$. Thus

$$\frac{\lg(d(v))}{2} > E\left(X_v \mid W \cap V(H) = S\right) = \frac{x2^{x-1} + d(v)}{2^x + 1}$$

So $2^x \lg(d(v)) + \lg(d(v)) > x2^x + 2d(v)$. Putting $t := \lg(d(v)) - x$ and rearranging yields

$$2^{x}t = 2^{x} \left(\lg(d(v)) - x \right) > 2d(v) - \lg(d(v)) > d(v).$$

Now we have the contradiction

$$\frac{t}{2^t} = \frac{2^x t}{d(v)} > 1.$$

Theorem 5.2. If G is a triangle-free graph and $f: V(G) \to \mathbb{N}$ by $f(v) := d_G(v) + 1 - \lfloor \frac{1}{4} \lg(d_G(v)) \rfloor$, then G has a nonempty induced subgraph H that is online f_H -choosable where $f_H(v) := f(v) + d_H(v) - d_G(v)$ for $v \in V(H)$.

Proof. Immediate upon applying Main Lemma to G since

$$\sum_{v \in V(G)} d_G(v) + 1 - f(v) = \sum_{v \in V(G)} \left\lfloor \frac{1}{4} \lg(d_G(v)) \right\rfloor \le \operatorname{mic}(G).$$

Corollary 5.3. If G is a triangle-free graph with $\Delta(G) \leq t$ for some $t \in \mathbb{N}$, then G is online $(t+1-\lfloor \frac{1}{4} \lg(t) \rfloor)$ -choosable.

Proof. Suppose not and choose a counterexample G and $t \in \mathbb{N}$ so as to minimize |G|. Put $f(v) := d_G(v) + 1 - \lfloor \frac{1}{4} \lg(d_G(v)) \rfloor$. By Theorem 5.2, G has a nonempty induced subgraph H that is online f_H -choosable where $f_H(v) := f(v) + d_H(v) - d_G(v)$ for $v \in V(H)$. Since $t+1-\lfloor \frac{1}{4} \lg(t) \rfloor \ge d_G(v)+1-\lfloor \frac{1}{4} \lg(d_G(v)) \rfloor$ for all $v \in V(G)$, we have that H is g(v)-choosable where $g(v) := t+1-\lfloor \frac{1}{4} \lg(t) \rfloor + d_H(v) - d_G(v)$. Now applying minimality of |G| and Lemma 3.9 gives a contradiction.

The best, known bounds for the chromatic number of triangle-free graphs are Kostochka's upper bound of $\frac{2}{3}\Delta + 2$ in [11] (see [16] for a proof in English) for small Δ and Johansson's upper bound of $\frac{9\Delta}{\ln(\Delta)}$ for large Δ . Johansson's proof also works for list coloring, but not for online list coloring. To the best of our knowledge Corollary 5.3 is the best, known-upper bound for online list colorings of triangle-free graphs. Additionally, Corollary 5.3 improves on Johansson's bound for list coloring for $\Delta \leq 8000$. The bound can surely be improved by a more complicated computation of $\operatorname{mic}(G)$, but not beyond around $\Delta + 1 - \lfloor 2\ln(\Delta) \rfloor$ via this method as can be seen by examples of triangle-free graphs with independence number near $\frac{2\ln(\Delta)}{\Delta}n$.

6 Gallai Forests

Recall that a Gallai forest is a graph such that each block is a clique or odd cycle. In this section we add to the many characterizations of Gallai forests.

6.1 Graphs with minimum mic

Lemma 6.1. Let G = (V, E) be a connected graph with a connected induced subgraph H. Then $\operatorname{mic}(G) \ge \operatorname{mic}(H) + |G - H|$. In particular, $\operatorname{mic}(G) \ge |G| - 1$.

Proof. Argue by induction on |G-H|. Plainly, |G-H| > 0. Let $v \notin V(H)$ be a noncutvertex in G/H. By induction $\operatorname{mic}(G-v) \ge \operatorname{mic}(H) + |G-H-v|$. Let A be an optimal set in G-v. Put $A' := A \cup \{v\}$ if ||v, A|| = 0; else put A' := A. Then A' is independent in G and $||A', V|| \ge \operatorname{mic}(H) + |G-H|$. Letting |H| = 1 yields $\operatorname{mic}(G) \ge |G| - 1$. \Box

Lemma 6.2. Every Gallai tree G = (V, E) satisfies mic(G) = |G| - 1.

Proof. Argue by induction on the number of blocks. If G has only one block then G is a clique or an odd cycle, and the lemma holds. Else let B be an endblock with cutvertex x. Then G' := G - (B - x) is a Gallai tree with fewer blocks. Let $A = I \cup J$ be an optimal set in G, where $I \subseteq V(B)$ and $J \subseteq V(G')$. Lemma 6.1 and induction yield the equality

$$|G| - 1 \le \operatorname{mic}(G) = ||A, V||_G = ||I, V(B)||_B + ||J, V(G')||_{G'} \le |B| + |G'| - 2 = |G| - 1. \quad \Box$$

The next lemma has many different proofs [5, 4, 7]. Although it is known as Rubin's Block Lemma [5], the lemma was implicit in the much earlier work of Gallai [6] and Dirac.

Lemma 6.3 (Rubin's Block Lemma). If G is a 2-connected graph that is not complete and not an odd cycle, then G contains an even cycle with at most one chord.

Theorem 6.4. A connected graph G = (V, E) is a Gallai tree if and only if mic(G) = |G| - 1; otherwise $mic(G) \ge |G|$.

Proof. By Lemmas 6.1 and 6.2, it suffices to show that $\operatorname{mic}(G) \geq |G|$ when G is not a Gallai tree. In this case Rubin's Block Lemma implies G has an induced even cycle C with only one possible chord xy. As C has an equitable 2-coloring, it has an independent set A with |A| = |C|/2 that contains at most one of x and y. Then A is independent in H := G[C], so $\operatorname{mic}(H) \geq |H|$. By Lemma 6.1, $\operatorname{mic}(G) \geq \operatorname{mic}(H) + |G - H| \geq |G|$.

6.2 *f*-AT graphs

For a graph G, we define $d_0: V(G) \to \mathbb{N}$ by $d_0(v) := d_G(v)$. The d_0 -choosable graphs were first characterized by Borodin [3] and independently by Erdős, Rubin and Taylor [5]. The connected graphs which are not d_0 -choosable are precisely the Gallai trees (connected graphs in which every block is complete or an odd cycle). Hladkỳ, Král and Schauz [7] generalized this classification to online d_0 -choosable graphs. In fact, they proved a classification in terms of Alon-Tarsi orientations as follows. A subgraph H of a directed multigraph D is called *Eulerian* if $d_H^-(v) = d_H^+(v)$ for every $v \in V(H)$. We call H even if ||H|| is even and odd otherwise. Let EE(D) be the number of even, spanning, Eulerian subgraphs of D and EO(D) the number of odd, spanning, Eulerian subgraphs of D. Note that the edgeless subgraph of D is even and hence we always have EE(D) > 0.

Let G be a graph and $f: V(G) \to \mathbb{N}$. We say that G is f-Alon-Tarsi (for brevity, f-AT) if G has an orientation D where $f(v) \ge d_D^+(v) + 1$ for all $v \in V(D)$ and $EE(D) \neq EO(D)$. One simple way to achieve $EE(D) \neq EO(D)$ is to have D be acyclic since then we have EE(D) = 1 and EO(D) = 0. In this case, ordering the vertices so that all edges point the same direction and coloring greedily shows that G is f-choosable. If we require f to be constant, we get the familiar coloring number col(G); that is, col(G) is the smallest k for which G has an acyclic orientation D with $k \ge d_D^+(v) + 1$ for all $v \in V(D)$. Alon and Tarsi [2] generalized from the acyclic case to arbitrary f-AT orientations.

Lemma 6.5. If a graph G is f-AT for $f: V(G) \to \mathbb{N}$, then G is f-choosable.

Schauz [19] extended this result to online f-choosability.

Lemma 6.6. If a graph G is f-AT for $f: V(G) \to \mathbb{N}$, then G is online f-choosable.

Hladkỳ, Král and Schauz [7] proved the following.

Theorem 6.7. A connected graph is d_0 -AT if and only if it is not a Gallai tree.

6.3 d_0 -KP graphs

Acyclic orientations are also a special case of kernel-perfect orientations. A graph H is f-KP if H has a kernel-perfect oriented supergraph H' with vertex set V(H), where $f(v) > d_{H'}^+(v)$ for all $v \in V(H')$. This supergraph for f-KP gives us more power. For example $K_4 - e$ has no kernel-perfect orientation D with $d_G(v) \ge d_D^+(v) + 1$ for every vertex v, but if H is the result of doubling the edge that is in two triangles, then there is a kernel-perfect orientation D of H with $d_G(v) \ge d_D^+(v) + 1$. We could allow a supergraph for f-AT as well, but this doesn't give us any more power. We will now prove the classification of connected d_0 -KP graphs.

Lemma 6.8. A connected graph G is d_0 -KP if and only if some nonempty induced subgraph of G is d_0 -KP.

Proof. The 'only if' direction is trivial. For the other direction, suppose G has a nonempty induced subgraph that is d_0 -KP and choose H to be a maximal such subgraph. If H = G, we are done, so suppose not. Let $S = N_G(V(H)) \setminus V(H)$. Then $S \neq \emptyset$ since G is connected. We show that $G[V(H) \cup S]$ is d_0 -KP, violating maximality of H. Start with a kernel-perfect oriented supergraph H' of H where $d_H(v) > d_{H'}^+(v)$ for all $v \in V(H')$. Now create an oriented supergraph Q' of $Q := G[V(H) \cup S]$ by directing all edges from V(H) to S into Sand replacing each edge in Q[S] with arcs going both ways. Clearly, this oriented graph is kernel-perfect. Each vertex in Q'[S] has at least one in-edge coming from V(H), so we have $d_Q(v) > d_{Q'}^+(v)$ for all $v \in V(Q')$ as desired. \Box

Corollary 6.9. A connected graph is d_0 -KP if and only if it is not a Gallai tree.

Proof. If G is a Gallai tree, then it is not d_0 -choosable and hence not d_0 -KP by the Kernel Lemma. For the other direction, let G be a connected graph that is not a Gallai tree. By Theorem 6.4, we have $\operatorname{mic}(G) \geq |G|$. Applying Main Lemma with $f(v) := d_G(v)$ for all $v \in V(G)$ gives a nonempty induced subgraph H of G that is f_H -KP where $f_H(v) :=$ $f(v) + d_H(v) - d_G(v) = d_H(v)$ for $v \in V(H)$. Now Lemma 6.8 shows that G is d_0 -KP. \Box

References

- [1] N. Alon and J. Spencer, The probabilistic method, vol. 57, Wiley-Interscience, 2004. 5
- [2] N. Alon and M. Tarsi, Colorings and orientations of graphs, Combinatorica 12 (1992), no. 2, 125–134. 6.2
- [3] O.V. Borodin, Criterion of chromaticity of a degree prescription, Abstracts of IV All-Union Conf. on Th. Cybernetics, 1977, pp. 127–128. 3, 6.2
- [4] R.C. Entringer, A Short Proof of Rubin's Block Theorem, Annals of Discrete Mathematics 27 - Cycles in Graphs (B.R. Alspach and C.D. Godsil, eds.), North-Holland Mathematics Studies, vol. 115, North-Holland, 1985, pp. 367–368. 6.1

- [5] P. Erdős, A.L. Rubin, and H. Taylor, *Choosability in graphs*, Proc. West Coast Conf. on Combinatorics, Graph Theory and Computing, Congressus Numerantium, vol. 26, 1979, pp. 125–157. 3, 6.1, 6.2
- [6] T. Gallai, Kritische graphen I., Math. Inst. Hungar. Acad. Sci 8 (1963), 165–192 (in German). 3, 6.1
- [7] Jan Hladký, Daniel Král, and Uwe Schauz, Brooks' theorem via the Alon-Tarsi theorem, Discrete Math. **310** (2010), no. 23, 3426–3428. MR 2721105 (2012a:05115) 1, 3, 3, 6.1, 6.2, 6.2
- [8] H.A. Kierstead and A.V. Kostochka, Ore-type versions of Brooks' theorem, Journal of Combinatorial Theory, Series B 99 (2009), no. 2, 298–305. 1
- [9] H.A. Kierstead and L. Rabern, Improved lower bounds on the number of edges in list critical and online list critical graphs, arXiv preprint arXiv:1406.7355 (2014). 1, 3
- [10] Alexandr Kostochka and Matthew Yancey, Ore's conjecture on color-critical graphs is almost true, J. Combin. Theory Ser. B 109 (2014), 73–101. MR 3269903 (document), 1, 1, 2.1, 4
- [11] A.V. Kostochka, A modification of a Catlin's algorithm, Methods and Programs of Solutions Optimization Problems on Graphs and Networks 2 (1982), 75–79. 5
- [12] A.V. Kostochka, L. Rabern, and M. Stiebitz, Graphs with chromatic number close to maximum degree, Discrete Mathematics 312 (2012), no. 6, 1273–1281. 1, 1
- [13] A.V. Kostochka and M. Stiebitz, A new lower bound on the number of edges in colourcritical graphs and hypergraphs, Journal of Combinatorial Theory, Series B 87 (2003), no. 2, 374–402. 1
- [14] L. Postle, Characterizing 4-Critical Graphs of Ore-Degree at most Seven, arXiv:1409.5116 (2014). 4, 4.2
- [15] L. Rabern, Δ-critical graphs with small high vertex cliques, Journal of Combinatorial Theory, Series B 102 (2012), no. 1, 126–130. 1
- [16] _____, Destroying noncomplete regular components in graph partitions, Journal of Graph Theory 72 (2013), no. 2, 123–127. 5
- [17] _____, Partitioning and coloring graphs with degree constraints, Discrete Mathematics
 9 (2013), no. 313, 1028–1034.
- [18] U. Schauz, Mr. Paint and Mrs. Correct, The Electronic Journal of Combinatorics 16 (2009), no. 1, R77. 1, 3

- [19] _____, Flexible color lists in Alon and Tarsi's theorem, and time scheduling with unreliable participants, The Electronic Journal of Combinatorics 17 (2010), no. 1, R13.
 6.2
- [20] V.G. Vizing, Vextex coloring with given colors, Metody Diskretn. Anal. 29 (1976), 3–10 (in Russian).
- [21] X. Zhu, On-line list colouring of graphs, The Electronic Journal of Combinatorics 16 (2009), no. 1, R127. 1