

# The Dominant Eigenvalue of an Essentially Nonnegative Tensor

Liping Zhang<sup>a\*</sup>    Liqun Qi<sup>b†</sup>    Ziyang Luo<sup>c‡</sup>

<sup>a</sup> Department of Mathematical Sciences, Tsinghua University, Beijing, China

<sup>b</sup> Department of Applied Mathematics, The Hong Kong Polytechnic University, Hong Kong

<sup>c</sup> Department of Mathematics, Beijing Jiaotong University, Beijing, China

**Abstract.** It is well known that the dominant eigenvalue of a real essentially nonnegative matrix is a convex function of its diagonal entries. This convexity is of practical importance in population biology, graph theory, demography, analytic hierarchy process and so on. In this paper, the concept of essentially nonnegativity is extended from matrices to higher order tensors, and the convexity and log convexity of dominant eigenvalues for such a class of tensors are established. Particularly, for any nonnegative tensor, the spectral radius turns out to be the dominant eigenvalue and hence possesses these convexities. Finally, an algorithm is given to calculate the dominant eigenvalue, and numerical results are reported to show the effectiveness of the proposed algorithm.

**Key words.** Essentially nonnegative tensor, dominant eigenvalue, convex function, spectral radius, algorithm.

**AMS subject classifications.** 74B99, 15A18, 15A69

## 1. Introduction

Tensors are increasingly ubiquitous in various areas of applied, computational, and industrial mathematics and have wide applications in data analysis and mining, information science, signal/image processing, and computational biology, etc; see the workshop report [26] and references therein. A tensor can be regarded as a higher-order generalization of a

---

\*Corresponding author. E-mail address: lzhang@math.stinghua.edu.cn

†Email address: maqilq@polyu.edu.hk

‡E-mail address: starkeynature@hotmail.com

matrix, which takes the form

$$\mathcal{A} = (A_{i_1 \dots i_m}), \quad A_{i_1 \dots i_m} \in R, \quad 1 \leq i_1, \dots, i_m \leq n.$$

Such a multi-array  $\mathcal{A}$  is said to be an  $m$ -order  $n$ -dimensional square real tensor with  $n^m$  entries  $A_{i_1 \dots i_m}$ . In this regard, a vector is a first-order tensor and a matrix is a second-order tensor. Tensors of order more than two are called higher-order tensors.

Analogous with that of matrices, the theory of eigenvalues and eigenvectors is one of the fundamental and essential components in tensor analysis. 72 references on eigenvalues of tensors can be found in the bibliography [22]. Wide range of practical applications can be found the references there. Compared with that of matrices, eigenvalue problems for higher-order tensors are nonlinear due to their multilinear structure. Various types of eigenvalues are defined for higher-order tensors in the setting of multilinear algebra. For example, the eigenvalue, the  $H$ -eigenvalue, the  $E$ -eigenvalue, the  $Z$ -eigenvalue, the  $N$ -eigenvalue defined by Qi for even order symmetric tensors [21], the  $l^p$  eigenvalues for general order symmetric tensors, and the mode- $i$  eigenvalues for general square tensors defined by Lim [16], the  $M$ -eigenvalue for a partially symmetric fourth-order tensor, defined by Qi, Dai and Han [23], the  $D$ -eigenvalue for a fourth-order symmetric tensor and a second-order symmetric tensor, defined by Qi, Wang and Wu [24], eigenvalues of general square tensors extended by Qi in [22] Chang, Pearson and Zhang in [5] and equivalent eigenvalue pair classes by Cartwright and Sturmfels [3]. Here, we are concerned with the one in [5, 22] as reviewed below.

**Definition 1.1** *Let  $C$  be the complex field. A pair  $(\lambda, x) \in C \times (C^n \setminus \{0\})$  is called an eigenvalue-eigenvector pair of  $\mathcal{A}$ , if they satisfy:*

$$\mathcal{A}x^{m-1} = \lambda x^{[m-1]}, \tag{1}$$

where  $n$ -dimensional column vectors  $\mathcal{A}x^{m-1}$  and  $x^{[m-1]}$  are defined as

$$\mathcal{A}x^{m-1} := \left( \sum_{i_2, \dots, i_m=1}^n A_{i_1 i_2 \dots i_m} x_{i_2} \cdots x_{i_m} \right)_{1 \leq i_1 \leq n} \quad \text{and} \quad x^{[m-1]} := (x_i^{m-1})_{1 \leq i \leq n},$$

respectively.

Nonnegative tensors, arising from multilinear pagerank [16], spectral hypergraph theory [1, 2, 14], and higher-order Markov chains [18], etc., form a singularly important class of tensors and have attracted more and more attention since they share some intrinsic properties with those of the nonnegative matrices. One of those properties is the Perron-Frobenius theorem on eigenvalues. In [4], Chang, Pearson, and Zhang generalized the Perron-Frobenius theorem for nonnegative matrices to irreducible nonnegative tensors. In [11], Friedland,

Gaubert and Han generalized the Perron-Frobenius theorem to weakly irreducible nonnegative tensors. Further generalization of the Perron-Frobenius theorem to nonnegative tensors can be found in [20, 27, 28]. Numerical methods for finding the spectral radius of nonnegative tensors are subsequently proposed. Ng, Qi, and Zhou [18] provided an iterative method to find the largest eigenvalue of an irreducible nonnegative tensor by extending the Collatz method [9] for calculating the spectral radius of an irreducible nonnegative matrix. The Ng-Qi-Zhou method is efficient but it is not always convergent for irreducible nonnegative tensors. Chang, Pearson and Zhang [6] extended the notion of primitive matrices into the realm of tensors, and established the convergence of the Ng-Qi-Zhou method for primitive tensors. Zhang and Qi [29] established global linear convergence of the Ng-Qi-Zhou method for essentially positive tensors. Liu, Zhou and Ibrahim [17] proposed an always convergent algorithm for computing the largest eigenvalue of an irreducible nonnegative tensors. Zhang, Qi, and Xu [30] established its explicit linear convergence rate for weakly positive tensors.

The essentially nonnegative tensor we defined in this paper is ultimately related to the nonnegative tensor and includes the latter one as a special case. It is a higher order generalization of the so-called essentially nonnegative matrix, whose off-diagonal entries are all nonnegative. Such a class of matrices possesses nice properties on eigenvalues. It follows from the famous Perron-Frobenius theorem for nonnegative matrices that for any essentially nonnegative matrix  $A$ , there exists a real eigenvalue with a nonnegative eigenvector, which is the largest one among real parts of all other eigenvalues of  $A$ . This special eigenvalue, termed as  $r(A)$ , is often called the *dominant eigenvalue* of  $A$ . Moreover,  $r(A)$  is known as a convex function of the diagonal entries of  $A$ . This convexity is a fundamental property for essentially nonnegative matrices [8, 10, 13, 19] and has numerous applications, not only in many branches of mathematics, such as graph theory [25], differential equations [19], but also in practical fields, e.g., population biology [19], demography [7], and analytic hierarchy process as well [12]. A natural question arises: does this convexity maintain for higher-order essentially nonnegative tensors? In this paper, we will give an affirmative answer to this question.

Similar to the essentially nonnegative matrix, an essentially nonnegative tensor has a real eigenvalue with the property that it is greater than or equal to the real part of every eigenvalue of  $\mathcal{A}$ . We also call it the *dominant eigenvalue* of  $\mathcal{A}$ , and denoted by  $\lambda(\mathcal{A})$ . Particularly, if  $\mathcal{A}$  is nonnegative, we have  $\rho(\mathcal{A}) = \lambda(\mathcal{A})$ , where  $\rho(\mathcal{A})$  is the spectral radius of  $\mathcal{A}$ . By employing the technique proposed in [19], we manage to obtain that the dominant eigenvalue is a convex function of the diagonal elements for any essentially nonnegative tensor. In addition, it is also a convex function of all elements of a tensor in some special convex set of tensors. Furthermore, the log convexity is also exploited for essentially nonnegative tensors with whose entries are either identically zero or log convex of some real univariate functions. Finally, we propose an algorithm to calculate the dominant eigenvalue, conver-

gence of the proposed algorithm is established and numerical results are reported to show the effectiveness of the proposed algorithm.

This paper is organized as follows. In Section 2, we recall some preliminary results, introduce the concept of essentially nonnegative tensors, and characterize some basic properties of such tensors. In Section 3, we show that the spectral radius of nonnegative tensors is a convex function of the diagonal elements, and so is the dominant eigenvalue of essentially nonnegative tensors. Section 4 is devoted to the log convexity of the dominant eigenvalue. In Section 5, we give an algorithm to calculate the dominant eigenvalue, and some numerical results are reported. Some concluding remarks are made in Section 6.

## 2. Preliminaries and essentially nonnegative tensors

We start this section with some fundamental notions and properties on tensors. An  $m$ -order  $n$ -dimensional tensor  $\mathcal{A}$  is called nonnegative (or, respectively, positive) if  $A_{i_1 \dots i_m} \geq 0$  (or, respectively,  $A_{i_1 \dots i_m} > 0$ ). The  $m$ -order  $n$ -dimensional unit tensor, denoted by  $\mathcal{I}$ , is the tensor whose entries are  $\delta_{i_1 \dots i_m}$  with  $\delta_{i_1 \dots i_m} = 1$  if and only if  $i_1 = \dots = i_m$  and otherwise zero. The symbol  $\mathcal{A} \geq \mathcal{B}$  means that  $\mathcal{A} - \mathcal{B}$  is a nonnegative tensor. A tensor  $\mathcal{A}$  is called reducible, if there exists a nonempty proper index subset  $I \subset \{1, 2, \dots, n\}$  such that

$$A_{i_1 \dots i_m} = 0, \quad \forall i_1 \in I, \quad \forall i_2, \dots, i_m \notin I.$$

Otherwise, we say  $\mathcal{A}$  is irreducible. We call  $\rho(\mathcal{A})$  the spectral radius of tensor  $\mathcal{A}$  if

$$\rho(\mathcal{A}) = \max\{|\lambda| : \lambda \text{ is an eigenvalue of } \mathcal{A}\},$$

where  $|\lambda|$  denotes the modulus of  $\lambda$ . An immediate consequence on the spectral radius follows directly from Corollary 3 in [21].

**Lemma 2.1** *Let  $\mathcal{A}$  be an  $m$ -order  $n$ -dimensional tensor. Suppose that  $\mathcal{B} = a(\mathcal{A} + b\mathcal{I})$ , where  $a$  and  $b$  are two real numbers. Then  $\mu$  is an eigenvalue of  $\mathcal{B}$  if and only if  $\mu = a(\lambda + b)$  and  $\lambda$  is an eigenvalue of  $\mathcal{A}$ . In this case, they have the same eigenvectors. Moreover,  $\rho(\mathcal{B}) \leq |a|(\rho(\mathcal{A}) + |b|)$ .*

Let  $P := \{x \in \mathbb{R}^n : x_i \geq 0, 1 \leq i \leq n\}$ , and  $\text{int}(P) = \{x \in \mathbb{R}^n : x_i > 0, 1 \leq i \leq n\}$ . The Perron-Frobenius theorem for nonnegative tensors is as below, following by [4, Theorem 1.4].

**Theorem 2.1** *If  $\mathcal{A}$  is an irreducible nonnegative tensor of order  $m$  and dimension  $n$ , then there exist  $\lambda_0 > 0$  and  $x_0 \in \text{int}(P)$  such that*

$$\mathcal{A}x_0^{m-1} = \lambda_0 x_0^{[m-1]}.$$

Moreover, if  $\lambda$  is an eigenvalue with a nonnegative eigenvector, then  $\lambda = \lambda_0$ . If  $\lambda$  is an eigenvalue of  $\mathcal{A}$ , then  $|\lambda| \leq \lambda_0$ .

The well-known Collatz minimax theorem [9] for irreducible nonnegative matrices has been extended to irreducible nonnegative tensors in [4, Theorem 4.2].

**Theorem 2.2** *Assume that  $\mathcal{A}$  is an irreducible nonnegative tensor of order  $m$  dimension  $n$ . Then*

$$\min_{x \in \text{int}(P)} \max_{x_i > 0} \frac{(\mathcal{A}x^{m-1})_i}{x_i^{m-1}} = \lambda_0 = \max_{x \in \text{int}(P)} \min_{x_i > 0} \frac{(\mathcal{A}x^{m-1})_i}{x_i^{m-1}},$$

where  $\lambda_0$  is the unique positive eigenvalue corresponding to a positive eigenvector.

For nonnegative tensors, Yang and Yang [28] asserted that the spectral radius is an eigenvalue, which is a generalization of the weak Perron-Frobenius theorem for nonnegative matrices. We state it [28, Theorem 2.3 and Lemma 5.8] in the following theorem.

**Theorem 2.3** *Assume that  $\mathcal{A}$  is a nonnegative tensor of order  $m$  dimension  $n$ , then  $\rho(\mathcal{A})$  is an eigenvalue of  $\mathcal{A}$  with a nonzero nonnegative eigenvector. Moreover, for any  $x \in \text{int}(P)$  we have*

$$\min_{1 \leq i \leq n} \frac{(\mathcal{A}x^{m-1})_i}{x_i^{m-1}} \leq \rho(\mathcal{A}) \leq \max_{1 \leq i \leq n} \frac{(\mathcal{A}x^{m-1})_i}{x_i^{m-1}}.$$

The following inequality and continuity of the spectral radius were given in [28, Lemma 3.5] and the proof of [28, Theorem 2.3], respectively.

**Lemma 2.2** *Let  $\mathcal{A}$  be a nonnegative tensor of order  $m$  and dimension  $n$ , and  $\varepsilon > 0$  be a sufficiently small number. Suppose  $\mathcal{A} \leq \mathcal{B}$ , then  $\rho(\mathcal{A}) \leq \rho(\mathcal{B})$ . Furthermore, if  $\mathcal{A}_\varepsilon = \mathcal{A} + \mathcal{E}$  where  $\mathcal{E}$  denotes the tensor with every entry being  $\varepsilon$ , then*

$$\lim_{\varepsilon \rightarrow 0} \rho(\mathcal{A}_\varepsilon) = \rho(\mathcal{A}).$$

Based on the above results, we can easily get the following lemma.

**Lemma 2.3** *Suppose that  $\mathcal{A}$  is an irreducible nonnegative tensor of order  $m$  dimension  $n$  and that there exists a nonzero vector  $x \in P$  and a real number  $\beta$  such that*

$$\mathcal{A}x^{m-1} \leq \beta x^{[m-1]}. \tag{2}$$

Then  $\beta > 0$ ,  $x \in \text{int}(P)$ , and  $\rho(\mathcal{A}) \leq \beta$ . Furthermore,  $\rho(\mathcal{A}) < \beta$  unless equality holds in (2).

**Proof.** Assume on the contrary that for  $x \in \text{int}(P)$  there exists a nonempty proper index subset  $I \subset \{1, 2, \dots, n\}$  such that  $x_i = 0$  for  $i \in I$  and  $x_i > 0$  for  $i \notin I$ . It follows from (2) that

$$A_{i_1 \dots i_m} = 0, \quad \forall i_1 \in I, \quad \forall i_2, \dots, i_m \notin I.$$

A contradiction to the irreducibility of  $\mathcal{A}$  comes, which henceforth implies that  $x \in \text{int}(P)$ . Together with Lemma 2.2 in [18],  $\mathcal{A}x^{m-1} \in \text{int}(P)$  is established. It further deduces that  $\beta > 0$ , and then the last statement holds from Lemma 5.9 in [28]. This completes the proof. ■

A simple but useful result follows immediately from Lemmas 2.2 and 2.3.

**Lemma 2.4** *Let  $\mathcal{A}$  and  $\mathcal{B}$  be irreducible nonnegative tensors of order  $m$  dimension  $n$ . If  $\mathcal{A} \leq \mathcal{B}$  and  $\mathcal{A} \neq \mathcal{B}$ , then  $\rho(\mathcal{A}) < \rho(\mathcal{B})$ .*

**Proof.** By Lemma 2.2,  $\rho(\mathcal{A}) \leq \rho(\mathcal{B})$ . Since  $\mathcal{B}$  is irreducible, Theorem 2.1 implies that there exists  $x \in \text{int}(P)$  such that

$$\mathcal{A}x^{m-1} \leq \mathcal{B}x^{m-1} = \rho(\mathcal{B})x^{[m-1]}. \quad (3)$$

Since  $x \in \text{int}(P)$  and  $\mathcal{A} \neq \mathcal{B}$ , equality cannot hold in (3). The desired strict inequality  $\rho(\mathcal{A}) < \rho(\mathcal{B})$  holds from Lemma 2.3. ■

The remaining of this section is devoted to the essentially nonnegative tensor, with the introduction of its definition and some basic properties.

**Definition 2.1** *Let  $\mathcal{A}$  be an  $m$ -order and  $n$ -dimensional tensor.  $\mathcal{A}$  is said to be essentially nonnegative if all its off-diagonal entries are nonnegative.*

**Theorem 2.4** *Let  $\mathcal{A}$  be an  $m$ -order and  $n$ -dimensional essentially nonnegative tensor. Then there exists  $\alpha > 0$  such that  $\alpha\mathcal{I} + \mathcal{A}$  is nonnegative. Moreover,  $\mathcal{A}$  has a real eigenvalue  $\lambda(\mathcal{A})$  with corresponding eigenvector in  $P$  and  $\lambda(\mathcal{A}) \geq \text{Re}\lambda$  for every eigenvalue  $\lambda$  of  $\mathcal{A}$ . Furthermore,*

$$\lambda(\mathcal{A}) = \rho(\alpha\mathcal{I} + \mathcal{A}) - \alpha.$$

**Proof.** Take

$$\alpha = \max_{1 \leq i \leq n} |A_{i \dots i}| + 1.$$

Clearly,  $\alpha > 0$  and  $\alpha\mathcal{I} + \mathcal{A}$  is nonnegative. By Lemma 2.1 and Theorem 2.3, we have

$$\rho(\alpha\mathcal{I} + \mathcal{A}) = \alpha + \lambda_1, \quad (4)$$

where  $\lambda_1$  is an eigenvalue of  $\mathcal{A}$  with corresponding eigenvector in  $P$ . Thus, (4) implies  $\lambda_1 \in \mathbb{R}$ . Let  $\lambda(\mathcal{A}) = \lambda_1$ , It follows from Lemma 2.1 that,

$$\begin{aligned}\lambda(\mathcal{A}) + \alpha &= \max\{|\alpha + \lambda| : \lambda \text{ is an eigenvalue of } \mathcal{A}\} \\ &\geq |\alpha + \lambda| \geq \alpha + \operatorname{Re}\lambda.\end{aligned}$$

The desired result arrives. ■

We call such an eigenvalue in the above theorem the *dominant eigenvalue* of  $\mathcal{A}$ . Throughout this paper,  $\rho(\mathcal{A})$  and  $\lambda(\mathcal{A})$  will denote the spectral radius and dominant eigenvalue respectively of a tensor  $\mathcal{A}$ . In the next section, we will show that both  $\rho(\mathcal{A})$  and  $\lambda(\mathcal{A})$  are convex functions of the diagonal elements of  $\mathcal{A}$ .

### 3. Convexity of the spectral radius and the dominant eigenvalue

Based on Theorems 2.1 and 2.3, we proceed with the convexity of the dominant eigenvalue of essentially nonnegative tensors in this section. It can be verified that the diagonal entries have nothing to do with the irreducibility of a tensor. Specifically, let  $\mathcal{A}$  be an essentially nonnegative tensor of order  $m$  and dimension  $n$ , define a nonnegative tensor  $\mathcal{B}$  by  $B_{i_1 \dots i_m} = 0$  if  $i_1 = \dots = i_m$  and the others are  $A_{i_1 \dots i_m}$ . Then  $\mathcal{A}$  is irreducible if and only if  $\mathcal{B}$  is. Equivalently,  $\mathcal{A}$  is irreducible if and only if  $\mathcal{A} + \alpha\mathcal{I}$  is, whenever it is nonnegative. Thus, by Lemma 2.2 and Theorem 2.4, it is sufficient to consider the class of irreducible nonnegative tensors.

**Theorem 3.1** *If  $\mathcal{A}$  is a given irreducible nonnegative tensor of order  $m$  and dimension  $n$ , and  $\mathcal{D}$  is allowed to vary in the class of nonnegative diagonal tensors, then the spectral radius  $\rho(\mathcal{A} + \mathcal{D})$  is a convex function of the diagonal entries of  $\mathcal{D}$ . That is, for nonnegative diagonal tensors  $\mathcal{C}$  and  $\mathcal{D}$  we have*

$$\rho(\mathcal{A} + t\mathcal{C} + (1-t)\mathcal{D}) \leq t\rho(\mathcal{A} + \mathcal{C}) + (1-t)\rho(\mathcal{A} + \mathcal{D}), \quad \forall t \in [0, 1]. \quad (5)$$

Moreover, equality holds in (5) for some  $t \in (0, 1)$  if and only if  $\mathcal{D} - \mathcal{C}$  is a scalar multiple of the unite tensor  $\mathcal{I}$ .

**Proof.** Since both  $\mathcal{A} + \mathcal{C}$  and  $\mathcal{A} + \mathcal{D}$  are irreducible nonnegative tensors, by Theorem 2.1 and Theorem 2.3 we have  $\rho(\mathcal{A} + \mathcal{C}) > 0$ ,  $\rho(\mathcal{A} + \mathcal{D}) > 0$ , and there exist  $x, y \in \operatorname{int}(P)$  such that

$$(\mathcal{A} + \mathcal{C})x^{m-1} = \rho(\mathcal{A} + \mathcal{C})x^{[m-1]}, \quad (\mathcal{A} + \mathcal{D})y^{m-1} = \rho(\mathcal{A} + \mathcal{D})y^{[m-1]}.$$

That is, for  $i = 1, 2, \dots, n$  we have

$$\begin{aligned}\rho(\mathcal{A} + \mathcal{C}) &= C_{i\dots i} + \sum_{i_2 \dots i_m = 1}^n A_{i i_2 \dots i_m} \frac{x_{i_2} \cdots x_{i_m}}{x_i}, \\ \rho(\mathcal{A} + \mathcal{D}) &= D_{i\dots i} + \sum_{i_2 \dots i_m = 1}^n A_{i i_2 \dots i_m} \frac{y_{i_2} \cdots y_{i_m}}{y_i},\end{aligned}$$

and hence  $\rho(\mathcal{A} + \mathcal{C}) - C_{i\dots i} > 0$  and  $\rho(\mathcal{A} + \mathcal{D}) - D_{i\dots i} > 0$ . The inequality between geometric and arithmetic means yields

$$\begin{aligned}\left( \sum_{i_2 \dots i_m = 1}^n A_{i i_2 \dots i_m} \frac{x_{i_2} \cdots x_{i_m}}{x_i} \right)^t \left( \sum_{i_2 \dots i_m = 1}^n A_{i i_2 \dots i_m} \frac{y_{i_2} \cdots y_{i_m}}{y_i} \right)^{1-t} &\leq t(\rho(\mathcal{A} + \mathcal{C}) - C_{i\dots i}) \\ &\quad + (1-t)(\rho(\mathcal{A} + \mathcal{D}) - D_{i\dots i}).\end{aligned}\quad (6)$$

Therefore, Hölder's inequality and Theorem 2.2 give from (6)

$$\begin{aligned}\rho(\mathcal{A} + t\mathcal{C} + (1-t)\mathcal{D}) &\leq \max_{1 \leq i \leq n} \left\{ tC_{i\dots i} + (1-t)D_{i\dots i} + \sum_{i_2 \dots i_m = 1}^n A_{i i_2 \dots i_m} \frac{z_{i_2} \cdots z_{i_m}}{z_i} \right\} \\ &\leq t\rho(\mathcal{A} + \mathcal{C}) + (1-t)\rho(\mathcal{A} + \mathcal{D}),\end{aligned}$$

where  $z_i = x_i^t y_i^{1-t}$  for  $i = 1, \dots, n$ . This shows (5) holds.

The inequality between geometric and arithmetic means implies that equality in (5) holds for  $t \in (0, 1)$  if and only if  $\rho(\mathcal{A} + \mathcal{C}) - C_{i\dots i} = \rho(\mathcal{A} + \mathcal{D}) - D_{i\dots i}$  for  $i = 1, \dots, n$ , i.e.,  $\mathcal{D} - \mathcal{C} = \gamma \mathcal{I}$  where  $\gamma = \rho(\mathcal{A} + \mathcal{D}) - \rho(\mathcal{A} + \mathcal{C})$ . This completes the proof.  $\blacksquare$

The convexity involved in Theorem 3.1 can be extended to the case of essentially non-negative tensors as follows.

**Corollary 3.1** *If  $\mathcal{A}$  is a given irreducible essentially nonnegative tensor of order  $m$  dimension  $n$  and  $\mathcal{D}$  is allowed to vary in the class of diagonal tensors, then the dominant eigenvalue  $\lambda(\mathcal{A} + \mathcal{D})$  is a convex function of the diagonal entries of  $\mathcal{D}$ . That is, for diagonal tensors  $\mathcal{C}$  and  $\mathcal{D}$  we have*

$$\lambda(\mathcal{A} + t\mathcal{C} + (1-t)\mathcal{D}) \leq t\lambda(\mathcal{A} + \mathcal{C}) + (1-t)\lambda(\mathcal{A} + \mathcal{D}), \quad \forall t \in [0, 1]. \quad (7)$$

Moreover, equality holds in (7) for some  $t \in (0, 1)$  if and only if  $\mathcal{D} - \mathcal{C}$  is a scalar multiple of the unite tensor  $\mathcal{I}$ .

**Proof.** Take

$$\alpha = 1 + \max_{1 \leq i \leq n} \{|A_{i\dots i}| + |C_{i\dots i}| + |D_{i\dots i}|\}.$$

Then  $\alpha\mathcal{I} + \mathcal{A} + \mathcal{C}$  and  $\alpha\mathcal{I} + \mathcal{A} + \mathcal{D}$  are all irreducible nonnegative tensors. By Theorem 2.4 and Theorem 3.1, we have for  $0 \leq t \leq 1$

$$\begin{aligned}\lambda(\mathcal{A} + t\mathcal{C} + (1-t)\mathcal{D}) + \alpha &= \rho(\alpha\mathcal{I} + \mathcal{A} + t\mathcal{C} + (1-t)\mathcal{D}) \\ &\leq t\rho(\alpha\mathcal{I} + \mathcal{A} + \mathcal{C}) + (1-t)\rho(\alpha\mathcal{I} + \mathcal{A} + \mathcal{D}) \\ &= t\lambda(\mathcal{A} + \mathcal{C}) + (1-t)\lambda(\mathcal{A} + \mathcal{D}) + \alpha,\end{aligned}$$

which yields (7). This completes the proof.  $\blacksquare$

Invoking the continuity presented in Lemma 2.2, it is easy to see that Theorem 3.1 and Corollary 3.1 hold even when  $\mathcal{A}$  is reducible. Moreover, Theorem 3.1 and Corollary 3.1 give necessary and sufficient conditions for the strict convexity. It is worth pointing out that the convexity of the dominant eigenvalue only works on the diagonal elements other than on all elements of the essentially nonnegative tensor, unless for some special cases. By collecting all symmetric essentially nonnegative tensors of order  $m$  and dimension  $n$ , we can get a closed convex cone, says  $\mathcal{S}(m, n)$ . The dominant eigenvalue of any tensor in  $\mathcal{S}(m, n)$  remains convex of all elements of the corresponding tensor in the domain  $\mathcal{S}(m, n)$ , as the following proposition shows.

**Proposition 3.1** *For any  $\mathcal{A}, \mathcal{B} \in \mathcal{S}(m, n)$ , and any  $t \in [0, 1]$ , we have*

$$\lambda(t\mathcal{A} + (1-t)\mathcal{B}) \leq t\lambda(\mathcal{A}) + (1-t)\lambda(\mathcal{B}).$$

**Proof.** For any  $\mathcal{A}, \mathcal{B} \in \mathcal{S}(m, n)$ , there exists an integer  $k > 0$  such that  $\mathcal{A} + k\mathcal{I}$  and  $\mathcal{B} + k\mathcal{I}$  are nonnegative and symmetric and hence for any of their convex combinations. The Perron-Frobenius theorem then ensures that  $\rho(\mathcal{A} + k\mathcal{I})$ ,  $\rho(\mathcal{B} + k\mathcal{I})$  and  $\rho(t\mathcal{A} + (1-t)\mathcal{B} + k\mathcal{I})$  ( $t \in [0, 1]$ ) all act as eigenvalues of the corresponding nonnegative symmetric tensor. By the variational approach, it follows that

$$\begin{aligned}&\rho(t\mathcal{A} + (1-t)\mathcal{B} + k\mathcal{I}) \\ &= \max \left\{ (t\mathcal{A} + (1-t)\mathcal{B} + k\mathcal{I})x^m : \sum_{i=1}^n x_i^m = 1 \right\} \\ &\leq t \max \left\{ (\mathcal{A} + k\mathcal{I})x^m : \sum_{i=1}^n x_i^m = 1 \right\} + (1-t) \max \left\{ (\mathcal{B} + k\mathcal{I})x^m : \sum_{i=1}^n x_i^m = 1 \right\} \\ &= t\rho(\mathcal{A} + k\mathcal{I}) + (1-t)\rho(\mathcal{B} + k\mathcal{I}).\end{aligned}$$

Combining with the fact that  $\rho(\mathcal{A} + k\mathcal{I}) = \lambda(\mathcal{A}) + k$ , the desired convexity follows.  $\blacksquare$

## 4. Log convexity of the spectral radius and the dominant eigenvalue

If a function  $f(x)$  is positive on its domain and  $\log f(x)$  is convex, then  $f(x)$  is called *log convex*. It is known [15] that the sum or product of log convex functions is also log convex. In this section we extend Kingman's theorem [15] for matrices to tensors. Our motivation for the following proof comes from [19].

**Theorem 4.1** *For  $t \in [0, 1]$  assume that  $\mathcal{F}(t) = (F_{i_1 \dots i_m}(t))$  is an  $m$ -order  $n$ -dimensional irreducible nonnegative tensor, and suppose that for  $1 \leq i_1, \dots, i_m \leq n$ ,  $F_{i_1 \dots i_m}(t)$  is either identically zero or positive and a log convex function of  $t$ . Then  $\rho(\mathcal{F}(t))$  is a log convex function of  $t$  for  $t \in [0, 1]$ . That is, if  $\mathcal{F}(0) = \mathcal{A}$ ,  $\mathcal{F}(1) = \mathcal{B}$ , and a nonnegative tensor  $\mathcal{G}(t) = (A_{i_1 \dots i_m}^{1-t} B_{i_1 \dots i_m}^t)$ , then*

$$\rho(\mathcal{F}(t)) \leq \rho(\mathcal{G}(t)) \leq \rho(\mathcal{A})^{1-t} \rho(\mathcal{B})^t. \quad (8)$$

Moreover, the first equality occurs in (8) for some  $t$  with  $t \in (0, 1)$  if and only if

$$\mathcal{F}(t) = \mathcal{G}(t),$$

and the second equality occurs in (8) for some  $t$  with  $t \in (0, 1)$  if and only if there exists a constant  $\sigma > 0$  and a positive diagonal matrix  $D = \text{diag}(d_1, \dots, d_n)$  such that

$$\mathcal{B} = \sigma \mathcal{A} \cdot D^{-(m-1)} \cdot \overbrace{D \cdots D}^{m-1} \text{ with } B_{i_1 i_2 \dots i_m} = \sigma A_{i_1 i_2 \dots i_m} d_{i_1}^{-(m-1)} d_{i_2} \cdots d_{i_m}.$$

**Proof.** Clearly,  $\mathcal{G}(0) = \mathcal{F}(0) = \mathcal{A}$  and  $\mathcal{G}(1) = \mathcal{F}(1) = \mathcal{B}$ . The log convexity assumption on  $F_{i_1 \dots i_m}(t)$  implies that, for  $t \in [0, 1]$ ,

$$\mathcal{F}(t) \leq \mathcal{G}(t),$$

which, together with Lemma 2.2, implies

$$\rho(\mathcal{F}(t)) \leq \rho(\mathcal{G}(t)). \quad (9)$$

Since  $\mathcal{F}(t)$  is irreducible, if equality holds in 9 for some  $t_0$  with  $0 < t_0 < 1$ , Lemma 2.4 implies that  $\mathcal{F}(t_0) = \mathcal{G}(t_0)$ .

Since  $\mathcal{F}(0)$  and  $\mathcal{F}(1)$  are irreducible nonnegative, Theorem 2.1 shows that there exist  $x, y \in \text{int}(P)$  such that

$$\mathcal{A}x^{m-1} = \rho(\mathcal{A})x^{[m-1]}, \quad \mathcal{B}y^{m-1} = \rho(\mathcal{B})y^{[m-1]}.$$

For a fixed  $t \in (0, 1)$ , define  $z = x^{1-t}y^t$ , i.e.,  $z_i = x_i^{1-t}y_i^t$  for  $1 \leq i \leq n$ . Then, the  $i$ th component of  $\mathcal{G}(t)z^{m-1}$  satisfies

$$(\mathcal{G}(t)z^{m-1})_i = \sum_{i_2 \dots i_m=1}^n A_{i i_2 \dots i_m}^{1-t} B_{i i_2 \dots i_m}^t z_{i_2} \cdots z_{i_m}.$$

Hence, Hölder's inequality gives

$$\begin{aligned} (\mathcal{G}(t)z^{m-1})_i &\leq \left( \sum_{i_2 \dots i_m=1}^n A_{i i_2 \dots i_m} x_{i_2} \cdots x_{i_m} \right)^{1-t} \left( \sum_{i_2 \dots i_m=1}^n B_{i i_2 \dots i_m} y_{i_2} \cdots y_{i_m} \right)^t \\ &= \rho(\mathcal{A})^{1-t} \rho(\mathcal{B})^t z_i^{m-1}. \end{aligned} \quad (10)$$

It follows from Lemma 2.3 and (10) that

$$\rho(\mathcal{G}(t)) \leq \rho(\mathcal{A})^{1-t} \rho(\mathcal{B})^t.$$

Furthermore, equality holds in 10 for some  $t \in (0, 1)$  if and only if, for  $1 \leq i \leq n$ ,

$$B_{i i_2 \dots i_m} y_{i_2} \cdots y_{i_m} = \sigma_i A_{i i_2 \dots i_m} x_{i_2} \cdots x_{i_m}. \quad (11)$$

Summing (11) over  $i_2 \dots i_m$  yields

$$\rho(\mathcal{B}) y_i^{m-1} = \sigma_i \rho(\mathcal{A}) x_i^{m-1}. \quad (12)$$

Take

$$\sigma = \frac{\rho(\mathcal{B})}{\rho(\mathcal{A})}, \quad d_i = \frac{x_i}{y_i},$$

Then, combining (11) and (12) we obtain

$$B_{i i_2 \dots i_m} = \sigma A_{i i_2 \dots i_m} d_i^{-(m-1)} d_{i_2} \cdots d_{i_m},$$

i.e.,

$$\mathcal{B} = \sigma \mathcal{A} \cdot D^{-(m-1)} \cdot \overbrace{D \cdots D}^{m-1}.$$

This completes the proof. ■

By Theorems 2.3 and 2.4, the above theorem also holds for the dominant eigenvalue of  $\mathcal{F}(t)$ , when  $\mathcal{F}(t)$  is essentially nonnegative with  $t \in [0, 1]$ .

## 5. An algorithm for calculating the dominant eigenvalue

Let  $\mathcal{A}$  be an essentially nonnegative tensor of order  $m$  and dimension  $n$ . In this section we propose an algorithm to calculate the dominant eigenvalue of an essentially nonnegative tensor. This algorithm is a modification of the Ng-Qi-Zhou algorithm given in [18]. By Lemma 2.2 and Theorem 2.4, we modify the Ng-Qi-Zhou algorithm such that for any essentially nonnegative tensor, the sequence generated by the modified algorithm always converges to its dominant eigenvalue.

Define two functions from  $\text{int}(P)$  to  $P$ :

$$F(x) := \min_{x_i \neq 0} \frac{(\mathcal{W}x^{m-1})_i}{x_i^{m-1}}, \quad G(x) := \max_{x_i \neq 0} \frac{(\mathcal{W}x^{m-1})_i}{x_i^{m-1}}, \quad (13)$$

where  $\mathcal{W}$  is an irreducible nonnegative tensor. The details of the modified algorithm are given as follows.

**Algorithm 5.1:**

**Step 0.** Given a sufficiently small number  $\varepsilon > 0$ . Let

$$\mathcal{W} = \mathcal{A} + \alpha \mathcal{I} + \mathcal{E}, \quad (14)$$

where

$$\alpha = \max_{1 \leq i \leq n} |A_{i\dots i}| + 1,$$

and  $\mathcal{E}$  is the tensor with every entry being  $\varepsilon$ . Choose any  $x^{(0)} \in \text{int}(P)$ . Set  $y^{(0)} = \mathcal{W}(x^{(0)})^{m-1}$  and  $k := 0$ .

**Step 1.** Compute

$$x^{(k+1)} = \frac{(y^{(k)})^{\lfloor \frac{1}{m-1} \rfloor}}{\left\| (y^{(k)})^{\lfloor \frac{1}{m-1} \rfloor} \right\|}, \quad y^{(k+1)} = \mathcal{B}(x^{(k+1)})^{m-1}.$$

According to (13), compute  $F(x^{(k+1)})$  and  $G(x^{(k+1)})$ .

**Step 2.** If  $G(x^{(k+1)}) - F(x^{(k+1)}) < \varepsilon$ , stop. Output  $\varepsilon$ -approximation of the dominant eigenvalue of  $\mathcal{A}$ :

$$\lambda^{(k+1)} = \frac{1}{2} (G(x^{(k+1)}) + F(x^{(k+1)})) - \alpha, \quad (15)$$

and the corresponding eigenvector  $x^{(k+1)}$ . Otherwise, set  $k := k + 1$  and go to Step 1.

Clearly, the tensor  $\mathcal{W}$  defined by (14) is positive and hence it is primitive. By Theorems 2.1 and 2.2, Algorithm 5.1 is well-defined. As an immediate consequence of Lemma 2.2, Theorem 2.4, and Theorem 3.3 in [20], we have the following convergence theorem.

**Theorem 5.1** *Let  $\mathcal{A}$  be an essentially nonnegative tensor of order  $m$  and dimensional  $n$ , and let  $\mathcal{W}$  be defined by (14) where  $\varepsilon$  is a sufficiently small number. Then the sequences  $\{F(x^{(k)})\}$  and  $\{G(x^{(k)})\}$ , generated by Algorithm 5.1, converge to  $\lambda_\varepsilon$ , where  $\lambda_\varepsilon$  is the unique*

positive eigenvalue of  $\mathcal{W}$ . Moreover, the sequence  $\{x^{(k)}\}$  converges to  $x_\varepsilon^*$  and  $x_\varepsilon^*$  is a positive eigenvector of  $\mathcal{W}$  corresponding to the largest eigenvalue  $\lambda_\varepsilon$ . Furthermore,

$$\lim_{\varepsilon \rightarrow 0} \lambda_\varepsilon = \lambda^*, \quad \lim_{\varepsilon \rightarrow 0} x_\varepsilon^* = x^*,$$

where  $\lambda^*$  is the spectral radius of  $\mathcal{A} + \alpha\mathcal{I}$  and  $x^*$  is the corresponding eigenvector. In particular, the dominant eigenvalue of  $\mathcal{A}$  is  $\lambda(\mathcal{A}) = \lambda^* - \alpha$  and  $x^*$  is also the eigenvector corresponding to  $\lambda(\mathcal{A})$ .

**Proof.** It follows from (14) that  $\mathcal{W}$  is positive, and hence it is irreducible. Therefore, for any nonzero  $x \in P$ , we have  $\mathcal{W}x^{m-1} \in \text{int}(P)$ , which shows that the tensor  $\mathcal{W}$  is primitive. Hence, by Theorem 3.3 in [20],

$$\lim_{k \rightarrow \infty} F(x^{(k)}) = \lim_{k \rightarrow \infty} \{G(x^{(k)}) = \lambda_\varepsilon, \quad \lim_{k \rightarrow \infty} x^{(k)} = x_\varepsilon^*.$$

Therefore,  $\lambda_\varepsilon - \alpha$  is an  $\varepsilon$ -approximation of the dominant eigenvalue of  $\mathcal{A}$  from Theorem 2.4. Furthermore, It follows from Lemma 2.2 that

$$\lim_{\varepsilon \rightarrow 0} \lambda_\varepsilon = \lambda^*, \quad \lim_{\varepsilon \rightarrow 0} x_\varepsilon^* = x^*.$$

It is easy to see that  $\lambda^* - \alpha$  is the dominant eigenvalue of  $\mathcal{A}$  with corresponding eigenvector  $x^*$ . ■

The above theorem shows that the convergence of Algorithm 5.1 is established for any essentially nonnegative tensor without the irreducible and primitive assumption. In order to show the effectiveness of Algorithm 5.1, we used MATLAB 7.4 to test it on the following three examples.

**Example 5.1** Consider the 3-order 3-dimensional essentially nonnegative tensor

$$\mathcal{A} = [A(1, :, :), A(2, :, :), A(3, :, :)],$$

where

$$\begin{aligned} A(:, :, 1) &= \begin{pmatrix} -1.51 & 8.35 & 1.03 \\ 4.04 & 3.72 & 1.45 \\ 6.71 & 6.43 & 1.35 \end{pmatrix} \\ A(:, :, 2) &= \begin{pmatrix} 9.02 & 0.78 & 6.89 \\ 9.71 & -5.32 & 1.85 \\ 2.09 & 4.17 & 2.98 \end{pmatrix} \\ A(:, :, 3) &= \begin{pmatrix} 9.55 & 1.57 & 6.91 \\ 5.63 & 5.55 & 1.43 \\ 5.76 & 8.29 & -0.15 \end{pmatrix} \end{aligned}$$

**Example 5.2** Let a 3-order 3-dimensional tensor  $\mathcal{A}$  be defined by  $A_{133} = A_{233} = A_{311} = A_{322} = 1$ ,  $A_{111} = A_{222} = -1$  and zero otherwise.

**Example 5.3** Let a 3-order 4-dimensional tensor  $\mathcal{A}$  be defined by  $A_{111} = A_{222} = A_{333} = A_{444} = -1$ ,  $A_{112} = A_{114} = A_{121} = A_{131} = A_{212} = A_{332} = A_{443} = 1$ , and zero otherwise.

Clearly, the essentially nonnegative tensors defined as Examples 5.1 and 5.2 respectively are irreducible. While, the essentially nonnegative tensor defined as Example 5.3 is reducible. We take  $\varepsilon = 10^{-9}$  and terminate our iteration when one of the conditions  $G(x^{(k)}) - F(x^{(k)}) \leq 10^{-9}$  and  $k \geq 100$  is satisfied. Algorithm 5.1 produces the dominant eigenvalue  $\lambda(\mathcal{A}) = 36.2757$  with eigenvector  $x^* = (1.0000; 0.8351; 0.9415)$  for Example 5.1, the dominant eigenvalue  $\lambda(\mathcal{A}) = 1$  with eigenvector  $x^* = (0.5000; 0.5000; 1.000)$  for Example 5.2, and the dominant eigenvalue  $\lambda(\mathcal{A}) = 0.8225$  with eigenvector  $x^* = (1.0000; 0.7408; 0.9714; 0.5330)$  for Example 5.3. The details of numerical results are reported in Tables 1 and 2. We list the output details at each iteration for Example 5.1 in Table 1. We also report the number of iterations (*No.Iter*), the elapsed CPU time (*CPU(sec)*), the lower bound  $\underline{\lambda}^{(k)} = F(x^{(k)}) - \alpha$  and the upper bound  $\overline{\lambda}^{(k)} = G(x^{(k)}) - \alpha$  for  $k \geq 1$ , the error  $\Delta^{(k)} = \|\mathcal{A}(x^{(k)})^{m-1} - \lambda^{(k)}(x^{(k)})^{[m-1]}\|_\infty$ , and the approximation  $\lambda^{(k)}$  defined by (15) of the dominant eigenvalue in Tables 1 and 2.

From Tables 1 and 2, we see that the sequence generated by Algorithm 5.1 converges to the dominant eigenvalue of the essentially nonnegative tensor without irreducibility. Algorithm 5.1 is promising for calculating the dominant eigenvalues of the test three examples.

Table 1: Detailed output of Algorithm 5.1 for Example 5.1

$k$	$\underline{\lambda}^{(k)}$	$\overline{\lambda}^{(k)}$	$\lambda^{(k)}$	$\overline{\lambda}^{(k)} - \underline{\lambda}^{(k)}$	$\Delta^{(k)}$
1	35.9969	36.5635	36.2802	0.5666	0.2833
2	36.2554	36.3030	36.2792	0.0476	0.0211
3	36.2747	36.2776	36.2762	0.0030	0.0015
4	36.2757	36.2758	36.2757	9.1725e-005	4.5870e-005
5	36.2757	36.2757	36.2757	6.7568e-006	2.9868e-006
6	36.2757	36.2757	36.2757	4.6425e-007	2.2441e-007
7	36.2757	36.2757	36.2757	1.9041e-008	1.4348e-008
8	36.2757	36.2757	36.2757	8.8998e-010	8.1036e-009

Table 2: Output of Algorithm 5.1 for Examples 5.1, 5.2, and 5.3

Example	No.Iter	CPU(sec)	$\underline{\lambda}^{(k)}$	$\overline{\lambda}^{(k)}$	$\lambda^{(k)}$	$\overline{\lambda}^{(k)} - \underline{\lambda}^{(k)}$	$\Delta^{(k)}$
5.1	8	0.013	36.2757	36.2757	36.2757	8.8998e-010	8.1036e-009
5.2	31	0.035	1.0000	1.0000	1.0000	9.6831e-010	4.1210e-009
5.3	37	0.078	0.8225	0.8225	0.8225	7.3324e-010	1.0635e-008

## 6. Conclusions

In this paper, we have introduced the concepts of essentially nonnegative tensors, which is closely related to nonnegative tensors. The main contribution is the convexity and log convexity of the dominant eigenvalue of an essentially nonnegative tensor, and hence the same for the spectral radius of a nonnegative tensor. By modifying the Ng-Qi-Zhou algorithm [18], we have proposed an algorithm (Algorithm 5.1) for calculating the dominant eigenvalue. Its convergence can be established for any essentially nonnegative tensor without the assumptions of irreducibility and primitiveness. Numerical results indicate that Algorithm 5.1 is promising.

**Acknowledgements.** Liping Zhang's work was supported by the National Natural Science Foundation of China(Grant No. 10871113). Liqun Qi's work was supported by the Hong Kong Research Grant Council. Ziyang Luo's work was supported by the National Basic Research Program of China (2010CB732501).

## References

- [1] S.R. Bulò and M. Pelillo, *A generalization of the Motzkin-Straus theorem to hypergraphs*, Optim. Lett. 3 (2009) 187-295.
- [2] S.R. Bulò and M. Pelillo, *New bounds on the clique number of graphs based on spectral hypergraph theory*, in: T. Stützle ed., Learning and Intelligent Optimization, Springer Verlag, Berlin, (2009) pp. 45-48.
- [3] D. Cartwright and B. Sturmfels, The number of eigenvalues of a tensor, to appear in: Linear Algebra and Applications.
- [4] K. C. Chang, K. Pearson and T. Zhang, *Perron Frobenius Theorem for nonnegative tensors*, Commu. Math. Sci. 6 (2008) 507-520.

- [5] K. C. Chang, K. Pearson and T. Zhang, *On eigenvalue problems of real symmetric tensors*, J. Math. Anal. Appl. 350 (2009) 416-422.
- [6] K.C. Chang, K. Pearson and T. Zhang, *Primitivity, the convergence of the NZQ method, and the largest eigenvalue for nonnegative tensors*, SIAM Journal on Matrix Analysis and Applications 32 (2011) 806-819.
- [7] J. E. Cohen, *Derivatives of the spectral radius as a function of nonnegative matrix elements*, Math. Proc. Cambridge Philos. Soc. 83 (1978) 183-190.
- [8] J. E. Cohen, *Convexity of the dominant eigenvalue of an essentially nonnegative matrix*, Proc. Amer. Math. Soc. 81 (1981) 657-658.
- [9] L. Collatz, *Einschliessungssatz für die charakteristischen Zahlen von Matrizen*, Math. Zeit. 48 (1942) 221-226.
- [10] S. Friedland, *Convex spectral functions*, Linear and Multilinear Algebra 9 (1981) 299-316.
- [11] S. Friedland, S. Gaubert and L. Han, *Perron-Frobenius theorem for nonnegative multilinear forms and extensions*, to appear in: Linear Algebra and Applications.
- [12] P. T. Harker, *Derivatives of the Perron root of a positive reciprocal matrix: with application to the analytic hierarchy process*, Applied Mathematics and Application 22 (1987) 217-232.
- [13] R. Horn and C. H. Johnson, *Matrix Analysis*, Cambridge University Press, Cambridge, 1996.
- [14] S. Hu and L. Qi, *Algebraic connectivity of an even uniform hypergraph*, to appear in: Journal of Combinatorial Optimization.
- [15] J. F. C. Kingman, *A convexity property of positive matrices*, Quart. J. Math. 12 (1961) 283-284.
- [16] L.-H. Lim, *Singular values and eigenvalues of tensors: a variational approach*, in Proceedings of the IEEE International Workshop on Computational Advances in Multi-Sensor Addaptive Processing (CAMSAP'05), Vol. 1, IEEE Computer Society Press, Piscataway, NJ, 2005, pp. 129-132.
- [17] Y. Liu, G. Zhou and N.F. Ibrahim, *An always convergent algorithm for the largest eigenvalue of an irreducible nonnegative tensor*, Journal of Computational and Applied Mathematics 235 (2010) 286-292.

- [18] M. Ng, L. Qi and G. Zhou, *Finding the largest eigenvalue of a nonnegative tensor*, SIAM J. Matrix Anal. Appl. 31 (2009) 1090-1099.
- [19] R. D. Nussbaum, *Convexity and Log convexity for the spectral radius*, Linear Algebra and its Applications 73 (1986) 59-122.
- [20] K. Pearson, *Essentially positive tensors*, International Journal of Algebra 4 (2010) 421 - 427.
- [21] L. Qi, *Eigenvalues of a real supersymmetric tensor*, J. Symbolic Comput. 40 (2005) 1302-1324.
- [22] L. Qi, *Bibliography of eigenvalues of tensors*,  
<http://www.polyu.edu.hk/ama/staff/new/qilq/TensorComp.htm>
- [23] L. Qi, H.H. Dai and D. Han, *Conditions for strong ellipticity and M-eigenvalues*, Frontiers of Mathematics in China 4 (2009) 349-364.
- [24] L. Qi, Y. Wang and E. X. Wu, *D-eigenvalues of diffusion kurtosis tensor*, J. Comput. Appl. Math. 221 (2008) 150-157.
- [25] R. Roth, *On the eigenvectors belonging to the minimum eigenvalue of an essentially nonnegative symmetric matrix with bipartite graph*, Linear Algebra and its Applications 118 (1989) 1-10.
- [26] C. Van Loan, *Future directions in tensor-based computation and modeling*, Workshop Report in Arlington, Virginia at National Science Foundation, February 20-21, 2009. <http://www.cs.cornell.edu/cv/TenWork/Home.htm>.
- [27] Q. Yang and Y. Yang, *Further results for Perron-Frobenius Theorem for nonnegative tensors II*, to appear in: SIAM Journal on Matrix Analysis.
- [28] Y. Yang and Q. Yang, *Further results for Perron-Frobenius Theorem for nonnegative tensors*, SIAM Journal on Matrix Analysis 31 (2010) 2517-2530.
- [29] L. Zhang and L. Qi, *Linear convergence of an algorithm for computing the largest eigenvalue of a nonnegative tensor*, to appear in: Numerical Linear Algebra with Applications.
- [30] L. Zhang, L. Qi and Y. Xu, *Linear convergence of the LZI algorithm for weakly positive tensors*, to appear in: Journal of Computational Mathematics.