Identifying codes in line graphs^{\ddagger}

Florent Foucaud^a, Sylvain Gravier^b, Reza Naserasr^{a,b}, Aline Parreau^b, Petru Valicov^a

^aLaBRI - Université de Bordeaux - CNRS, 351 cours de la Libération, 33405 Talence cedex, France. ^bInstitut Fourier 100, rue des Maths, BP 74, 38402 St Martin d'Hères cedex, France.

Abstract

An identifying code of a graph is a subset of its vertices such that every vertex of the graph is uniquely identified by the set of its neighbours within the code. We study the edge-identifying code problem, i.e. the identifying code problem in line graphs. If $\gamma^{ID}(G)$ denotes the size of a minimum identifying code of an identifiable graph G, we show that the usual bound $\gamma^{ID}(G) \geq \lceil \log_2(n+1) \rceil$, where n denotes the order of G, can be improved to $\Theta(\sqrt{n})$ in the class of line graphs. Moreover, this bound is tight. We also prove that the upper bound $\gamma^{ID}(\mathcal{L}(G)) \leq 2|V(G)| - 5$, where $\mathcal{L}(G)$ is the line graph of G, holds (with two exceptions). This implies that a conjecture of R. Klasing, A. Kosowski, A. Raspaud and the first author holds for a subclass of line graphs. Finally, we show that the edge-identifying code problem is NP-complete, even for the class of planar bipartite graphs of maximum degree 3 and arbitrarily large girth.

Keywords: Identifying codes, Dominating sets, Line graphs, NP-completeness.

1. Introduction

An *identifying code* of a graph G is a subset C of vertices of G such that for each vertex x, the set of vertices in C at distance at most 1 from x, is nonempty and uniquely identifies x. More formally:

Definition 1. Given a graph G, a subset C of V(G) is an identifying code of G if C is both:

- a dominating set of G, i.e. for each vertex $v \in V(G)$, $N[v] \cap \mathcal{C} \neq \emptyset$, and
- a separating set of G, i.e. for each pair $u, v \in V(G)$ $(u \neq v)$, $N[u] \cap C \neq N[v] \cap C$.

Here N[v] is the closed neighbourhood of v in G. This concept was introduced in 1998 in [13] and is a well-studied one (see e.g. [1, 4, 5, 8, 9, 12, 16]).

A vertex x is a twin of another vertex y if N[x] = N[y]. A graph G is called twin-free if no vertex has a twin. The first observation regarding the concept of identifying codes is that a graph is *identifiable* if and only if it is twin-free. As usual for many other graph theory concepts, a natural problem in the study of identifying codes is to find one of a minimum size. Given a graph G, the smallest size of an identifying code of G is called *identifying code number* of G and denoted by $\gamma^{\text{ID}}(G)$. The main lines of research here are to find the exact value of $\gamma^{\text{ID}}(G)$ for interesting graph classes, to approximate it and to give lower or upper bounds in terms of simpler graph parameters. Examples of classic results are as follows:

Theorem 2. [12] If G is a twin-free graph with at least two edges, then $\gamma^{ID}(G) \leq |V(G)| - 1$.

The collection of all twin-free graphs reaching this bound is classified in [8].

A better upper bound in terms of both number of vertices and maximum degree $\Delta(G)$ of a graph G is also conjectured:

 $^{^{\}circ}$ This research is supported by the ANR Project IDEA • ANR-08-EMER-007, 2009-2011.

Preprint submitted to J. of Graph Theory

Conjecture 3. [9] There exists a constant c such that for every twin-free graph G,

$$\gamma^{\scriptscriptstyle ID}(G) \le |V(G)| - \frac{|V(G)|}{\Delta(G)} + c.$$

Some support for this conjecture is provided in [8, 9, 10].

The parameter $\gamma^{\text{ID}}(G)$ is also bounded below by a function of |V(G)| where equality holds for infinitely many graphs.

Theorem 4. [13] For any twin-free graph G, $\gamma^{ID}(G) \ge \lceil \log_2(|V(G)| + 1) \rceil$.

The collection of all graphs attaining this lower bound is classified in [16].

From a computational point of view, it is shown that given a graph G, finding the exact value of $\gamma^{\text{ID}}(G)$ is in the class of NP-hard problems. It in fact remains NP-hard for many subclasses of graphs [1, 4]. Furthermore, approximating $\gamma^{\text{ID}}(G)$ is not easy either as shown in [14, 11, 17]: it is NP-hard to approximate $\gamma^{\text{ID}}(G)$ within a $o(\log(|V(G)|))$ -factor.

The problem of finding identifying codes in graphs can be viewed as a special case of the more general combinatorial problem of finding *transversals* in hypergraphs (a transversal is a set of vertices intersecting each hyperedge). More precisely, to each graph G one can associate the hypergraph $\mathcal{H}(G)$ whose vertices are vertices of G and whose hyperedges are all the sets of the form N[v] and $N[u] \ominus N[v]$ (symmetric difference of N[u] and N[v]). Finding an identifying code for G is then equivalent to finding a transversal for $\mathcal{H}(G)$. Though the identifying code problem is captured by this more general problem, the structural properties of the graph from which the hypergraph is built allow one to obtain stronger results which are not true for general hypergraphs. In this work, we show that even stronger results can be obtained if we consider hypergraphs coming from line graphs. These stronger results follow from the new perspective of identifying edges by edges.

Given a graph G and an edge e of G, we define N[e] to be the set of edges adjacent to e together with e itself. An edge-identifying code of a graph G is a subset C_E of edges such that for each edge e the set $N[e] \cap C_E$ is nonempty and uniquely determines e. More formally:

Definition 5. Given a graph G, a subset C_E of E(G) is an edge-identifying code of G if C_E is both:

- an edge-dominating set of G, i.e. for each edge $e \in E(G)$, $N[e] \cap C_E \neq \emptyset$, and
- an edge-separating set of G, *i.e.* for each pair $e, f \in E(G)$ $(e \neq f), N[e] \cap C_E \neq N[f] \cap C_E$.

We will say that an edge e separates edges f and g if either e belongs to N[f] but not to N[g], or viceversa. When considering edge-identifying codes we will assume the edge set of the graph is nonempty. The line graph $\mathcal{L}(G)$ of a graph G is the graph with vertex set E(G), where two vertices of $\mathcal{L}(G)$ are adjacent if the corresponding edges are adjacent in G. It is easily observed that the notion of edge-identifying code of G is equivalent to the notion of (vertex-)identifying code of the *line graph* of G. Thus a graph G admits an edge-identifying code if and only if $\mathcal{L}(G)$ is twin-free. A pair of twins in $\mathcal{L}(G)$ can correspond in G to a pair of: 1. parallel edges; 2. adjacent edges whose non-common ends are of degree 1; 3. adjacent edges whose non common ends are of degree 2 but they are connected to each other. Hence we will consider simple graphs only. A pair of edges of type 2 or type 3 is called *pendant* (see Figure 1) and thus a graph is *edge-identifiable* if and only if it is *pendant-free*. The smallest size of an edge-identifying code of an edge-identifiable graph G is denoted by $\gamma^{\text{EID}}(G)$ and is called *edge-identifying code number* of G.

As we will use it often throughout the paper, given a graph G and a set S_E of its edges, we define the graph induced by S_E to be the graph with the set of all endpoints of the edges of S_E as its vertex set and S_E as its edge set.

To warm up, we notice that five edges of a perfect matching of the Petersen graph P, form an edgeidentifying code of this graph (see Figure 2). The lower bound of Theorem 4 proves that $\gamma^{\text{EID}}(P) \ge 4$. Later, by improving this bound for line graphs, we will see that in fact $\gamma^{\text{EID}}(P) = 5$ (see Theorem 12 and Theorem 16).




Figure 1: Two possibilities for a pair of pendant edges (thick edges) in G




Figure 2: An edge-identifying code of the Petersen graph

The outline of the paper is as follows: in Section 2, we introduce some useful lemmas and give the edge-identifying code number of some basic families of graphs. In Section 3, we improve the general lower bound for the class of line graphs, then in Section 4 we improve the upper bound. Finally, in Section 5 we show that determining $\gamma^{\text{EID}}(G)$ is also in the class of NP-hard problems even when restricted to planar subcubic bipartite graphs of arbitrarily large girth, but the problem is 4-approximable in polynomial time.

2. Preliminaries

In this section we first give some easy tools which help for finding minimum-size edge-identifying codes of graphs. We then apply these tools to determine the exact values of γ^{EID} for some basic families of graphs. We recall that C_n is the cycle on n vertices, P_n is the path on n vertices, K_n is the complete graph on nvertices and $K_{n,m}$ is the complete bipartite graph with parts of size n and m. We recall that the *girth* of a graph is the length of one of its shortest cycles. An *edge cover* of a graph G is a subset S_E of its edges such that the union of the endpoints of S_E equals V(G). A *matching* is a set of pairwise non-adjacent edges, and a *perfect matching* is a matching which is also an edge cover.

Lemma 6. Let G be a simple graph with girth at least 5. Let C_E be an edge cover of G such that the graph $(V(G), C_E)$ is pendant-free. Then C_E is an edge-identifying code of G. In particular, if G has a perfect matching M, M is an edge-identifying code of G.

Proof. The code C_E is an edge-dominating set of G because it covers all the vertices of G. To complete the proof, we need to prove that C_E is also an edge-separating set. Let e_1 , e_2 be two edges of G. If $e_1, e_2 \in C_E$, then $C_E \cap N[e_1] \neq C_E \cap N[e_2]$ because $(V(G), C_E)$ is pendant-free. Otherwise, we can assume that $e_2 \notin C_E$. If $e_1 \in C_E$ and $C_E \cap N[e_1] = C_E \cap N[e_2]$, then e_2 must be adjacent to e_1 . Let u be their common vertex and $e_2 = uv$. Since C_E is an edge cover, there is an edge $e_3 \in C_E$ which is incident to v. However, e_3 cannot be adjacent to e_1 because G is triangle-free. Therefore e_3 separates e_1 and e_2 . Finally, we assume neither of e_1 and e_2 is in C_E . Then there are two edges of C_E , say e_3 and e_4 , adjacent to the two ends of e_1 . But since G has neither C_3 nor C_4 as a subgraph, e_3 and e_4 cannot both be adjacent to e_2 and, therefore, e_1 and e_2 are separated.

We note that in the previous proof the absence of C_4 is only used when the endpoints of e_1, e_2, e_3, e_4 could induce a C_4 which would not be adjacent to any other edge of C_E . Thus, we have the following stronger statement:

Lemma 7. Let G be a triangle-free graph. Let C_E be a subset of edges of G that covers vertices of G, such that C_E is pendant-free. If for no pair xy, uv of isolated edges in C_E , the set $\{x, y, u, v\}$ induces a C_4 in G, then C_E is an edge-identifying code of G.

We will also need the following lemma about pendant-free trees.

Lemma 8. If T is a pendant-free tree on more than two vertices, then T has two vertices of degree 1, each adjacent to a vertex of degree 2.

Proof. Take a longest path in T, then it is easy to verify that both ends of this path satisfy the condition of the lemma.

We are now ready to determine the value of γ^{EID} of some families of graphs.

Proposition 9. We have $\gamma^{EID}(K_n) = \begin{cases} 5, & \text{if } n = 4 \text{ or } 5 \\ n-1, & \text{if } n \ge 6 \end{cases}$. Furthermore, let \mathcal{C}_E be an edge-identifying code of K_n of size n-1 ($n \ge 6$) and let G_1, G_2, \ldots, G_k be the connected components of $(V(K_n), \mathcal{C}_E)$. Then exactly one component, say G_i , is isomorphic to K_1 and every other component G_j ($j \ne i$) is isomorphic to a cycle of length at least 5.

Proof. We note that $\mathcal{L}(K_4)$ is isomorphic to $K_6 \setminus M$, where M is a perfect matching of K_6 . One can check that this graph has identifying code number 5. By a case analysis, we can show that K_5 does not admit an edge-identifying code of size 4. Indeed, since an edge-identifying code must be pendant-free, there are only two graphs possible for an edge-identifying code of this size: a path P_5 or a cycle C_4 . In both cases, there are edges which are not separated. Edges of a C_5 form an edge-identifying code of size 5 of K_5 , hence $\gamma^{\text{EID}}(K_5) = 5$. Furthermore, it is not difficult to check that the set of edges of a cycle of length n - 1 ($n \ge 6$) identifies all edges of K_n . Thus we have $\gamma^{\text{EID}}(K_n) \le n - 1$. The fact that $\gamma^{\text{EID}}(K_n) \ge n - 1$ follows from the second part of the theorem which is proved as follows.

Let C_E be an edge-identifying code of K_n of size n-1 or less $(n \ge 6)$. Let $G' = (V(K_n), C_E)$. Let G_1, G_2, \ldots, G_k be the connected components of G'. Since G' has n vertices but at most n-1 edges, at least one component of G' is a tree. On the other hand we claim that at most one of these components can be a tree and that such tree would be isomorphic to K_1 . Let G_i be a tree. First we show that $|V(G_i)| \le 2$. If not, by Lemma 8 there is a vertex x of degree 1 in G_i with a neighbour u of degree 2. Let v be the other neighbour of u. Then the edges xv and uv are not identified. If $V(G_i) = \{x, y\}$ then for any other vertex u, the edges ux and uy are not separated. Finally, if there are G_i and G_j with $V(G_i) = \{x\}$ and $V(G_j) = \{y\}$, then the edge xy is not dominated by C_E . Thus exactly one component of G', say G_1 , is a tree and $G_1 \cong K_1$. This implies that $\gamma^{\text{EID}}(K_n) \ge n-1$. Therefore, $\gamma^{\text{EID}}(K_n) = n-1$ and, furthermore, each G_i , $(i \ge 2)$, is a graph with a unique cycle.

It remains to prove that each G_i , $i \ge 2$ is isomorphic to a cycle of length at least 5. By contradiction suppose one of these graphs, say G_2 , is not isomorphic to a cycle. Since G_2 has a unique cycle, it must contain a vertex v of degree 1. Let t be the neighbour of v in G_2 and let u be the vertex of G_1 . Then the edges tv and tu are not separated by \mathcal{C}_E . Finally we note that such cycle cannot be of length 3 or 4, because C_3 is not pendant-free and in C_4 , the two chords (which are edges of K_n) would not be separated.

Proposition 10. $\gamma^{\text{EID}}(K_{n,n}) = \left\lceil \frac{3n-1}{2} \right\rceil$ for $n \ge 3$.

Proof. Let X and Y be the two parts of $K_{n,n}$. If n is even, then let $\{A_i\}_{i=1}^{\frac{n}{2}}$ be a partition of vertices such that each A_i has exactly two vertices in X and two in Y. Let G_i be a subgraph of $K_{n,n}$ isomorphic to P_4 and with A_i as its vertices. If n is odd, let A_1 be of size 2 and having exactly one element from X and one element from Y and let also G_1 be the subgraph (isomorphic to K_2) induced by A_1 . Then we define A_i 's

and G_i 's $(i \ge 2)$ as in the previous case (for $K_{n-1,n-1}$). By Lemma 7, the set of edges in the G_i 's induces an edge-identifying code of $K_{n,n}$ of size $\lceil \frac{3n-1}{2} \rceil$. To complete the proof we show that there cannot be any smaller edge-identifying code.

Let \mathcal{C}_E be an edge-identifying code of G. Let G_1, G_2, \ldots, G_k be the connected components of $(X \cup Y, \mathcal{C}_E)$. The proof will be completed if we show that except possibly one, every G_i must have at least four vertices. To prove this claim we first note that there is no connected pendant-free graph on three vertices. We now suppose G_1 and G_2 are both of order 2. Then the two edges connecting G_1 and G_2 are not separated. If G_1 is of order 1 and G_2 is of order 2, then the edge connecting G_1 to G_2 is not identified from the edge of G_2 . If G_1 and G_2 are both of order 1, then both of their vertices must be in the same part of the graph as otherwise the edge connecting them is not dominated by \mathcal{C}_E . But now for any vertex x which is not in the same part as G_1 and G_2 , the edges connecting x to G_1 and G_2 are not separated.

The following examples show that if true, the upper bound of Conjecture 3 is tight even in the class of line graphs. These examples were first introduced in [9] but without using the notion of edge-identifying codes.

Proposition 11. Let G be a k-regular multigraph $(k \ge 3)$. Let G_1 be obtained from G by subdividing each edge exactly once. Then $\gamma^{\text{EID}}(G_1) = (k-1)|V(G)| = |E(G_1)| - \frac{|E(G_1)|}{2k-2} = |V(\mathcal{L}(G_1))| - \frac{|V(\mathcal{L}(G_1))|}{\Delta(\mathcal{L}(G_1))}$.

Proof. Let x be a vertex of G_1 of degree at least 3 (an original vertex from G). For each edge e_i^x incident to x, let $e_i'^x$ be the edge adjacent to e_i^x but not incident to x and let $A_x = \{e_i'^x\}_{i=1}^k$. Then $\{A_x \mid x \in V(G)\}$ is a partition of $E(G_1)$. For any edge-identifying code \mathcal{C}_E of G_1 , if two elements of A_x , say $e_1'^x$ and $e_2'^x$, are both not in \mathcal{C}_E , then e_1^x and e_2^x are not separated. Thus $|\mathcal{C}_E \cap A_x| \ge k-1$. This proves that $|\mathcal{C}_E| \ge (k-1)|V(G)|$.

We now build an edge-identifying code of this size by choosing one edge of each set A_x , in such a way that for each vertex x originally from G, exactly one edge incident to x is chosen. Then the set of non-chosen edges will be an edge-identifying code. To select this set of edges, one can consider the incident bipartite multigraph H of G: the vertex set of H is $V \cup V'$ where V and V' are copies of V(G) and there is an edge xx'in H if $x \in V$, $x' \in V'$ and $xx' \in E(G)$. The multigraph H is k-regular and bipartite, thus it has a perfect matching M. For each vertex $x \in V$, let $\rho(x)$ be the vertex in V' such that $x\rho(x) \in M$. Let now e'_M be the edge of G_1 that belongs to the set A_x and is incident to $\rho(x)$ (in G_1). Finally, let $\mathcal{C}_E = E(G_1) \setminus \{e'_M\}_{x \in V(G)}$. Exactly one element of each A_x is not in \mathcal{C}_E , and for each vertex x, exactly one edge incident to x is not in \mathcal{C}_E . This implies that \mathcal{C}_E is an edge-identifying code.

For a simple example of the previous construction, let G be the multigraph on two vertices with k parallel edges. Then $G_1 \cong K_{2,n}$ and therefore $\gamma^{\text{EID}}(K_{2,n}) = 2n - 2$.

Hypercubes, being the natural ground of code-like structures, have been a center of focus for determining the smallest size of their identifying codes. The hypercube of dimension d, denoted \mathcal{H}_d , is a graph whose vertices are elements of \mathbb{Z}_2^d with two vertices being adjacent if their difference is in the standard basis: $\{(1,0,0,\ldots,0), (0,1,0,\ldots,0), \ldots, (0,0,\ldots,0,1)\}$. The hypercube of dimension d can also be viewed as the cartesian product of the hypercube of dimension d-1 and K_2 . In this way of building \mathcal{H}_d we add a new coordinate to the left of the vectors representing the vertices of \mathcal{H}_{d-1} .

The problem of determining the identifying code number of hypercubes has proved to be a challenging one from both theoretical and computational points of view. Today the precise identifying code number is known for only seven hypercubes [3]. In contrast, we show here that finding the edge-identifying code number of a hypercube is not so difficult. We first introduce the following general theorem.

Theorem 12. Let G be a connected pendant-free graph. We have:

$$\gamma^{\text{EID}}(G) \ge \frac{|V(G)|}{2}.$$

Proof. Let C_E be an edge-identifying code of G. Let G' be the subgraph induced by C_E and let G_1, \ldots, G_s be the connected components of G'. Let n_i be the order of G_i and k_i be its size (thus $\sum_{i=1}^s k_i = |\mathcal{C}_E|$). Let $X = V(G) \setminus V(G')$ and n'_i be the number of vertices in X that are joined to a vertex of G_i in G. We show

that $n'_i + n_i \leq 2k_i$. If $k_i = 1$, then clearly $n'_i = 0$ and $n'_i + n_i = 2 = 2k_i$. If G_i is a tree, then $n_i = k_i + 1$ and, by Lemma 8, G_i must have two vertices of degree 2 each having a vertex of degree 1 as a neighbour. Then no vertex of X can be adjacent to one of these two vertices in G. Moreover, each other vertex of G_i can be adjacent to at most one vertex in X. So $n'_i \leq k_i - 1$, and finally $n_i + n'_i \leq 2k_i$. If G_i is not at tree, we have $n_i \leq k_i$ and $n'_i \leq n_i$ and, therefore, $n'_i + n_i \leq 2k_i$. Finally, since G is connected, each vertex in X is connected to at least one G_i . Hence by counting the number vertices of G we have:

$$|V(G)| \le \sum_{i=1}^{s} (n_i + n'_i) \le 2 \sum_{i=1}^{s} k_i \le 2|\mathcal{C}_E|.$$

Theorem 12 together with Lemma 7 leads to the following result:

Corollary 13. Let G be a triangle-free pendant-free graph. Suppose G has a perfect matching M with the property that for any pair xy, uv of edges in M, the set $\{x, y, u, v\}$ does not induce a C_4 . Then M is an optimal edge-identifying code and $\gamma^{\text{EID}}(G) = \frac{|V(G)|}{2}$.

We note that in particular, if the girth of a graph G is at least 5 and G admits a perfect matching M, then M is a minimum-size identifying code of G. For example, the edge-identifying code of the Petersen graph given in Figure 2 is optimal.

As another application of Corollary 13, we give the edge-identifying code of all hypercubes of dimension $d \ge 4$.

Proposition 14. For $d \geq 4$, we have $\gamma^{EID}(\mathcal{H}_d) = 2^{d-1}$.

Proof. By Theorem 12, we have $\gamma^{\text{EID}}(\mathcal{H}_d) \geq 2^{d-1}$. We will construct by induction a perfect matching M_d of \mathcal{H}_d such that no pair of edges induces a C_4 , for $d \geq 4$. By Lemma 7, M_d will be an edge-identifying code of \mathcal{H}_d , proving the result. Two such matchings of \mathcal{H}_4 , which are also disjoint, are presented in Figure 3. The matching M_5 can now be built using each of these two matchings of \mathcal{H}_4 — one matching per copy of \mathcal{H}_4 in \mathcal{H}_5 . It is easily verified that M_5 has the required property. Furthermore, M_5 has the extra property that for each edge uv of M_5 , u and v do not differ on the first coordinate (recall that we build \mathcal{H}_5 from \mathcal{H}_4 by adding a new coordinate on the left, hence the first coordinate is a the new one). We now build the matching M_d of \mathcal{H}_d ($d \geq 6$) from M_{d-1} in such a way that no two edges of M_d belong to a 4-cycle in \mathcal{H}_d and that for each edge uv of M_d , u and v do not differ on the first coordinate. To do this, let \mathcal{H}'_1 be the copy of \mathcal{H}_{d-1} in \mathcal{H}_d induced by the set of vertices whose first coordinate is 0. Similarly, let \mathcal{H}'_2 be the copy of \mathcal{H}_{d-1} in \mathcal{H}_d induced by the other vertices. Let \mathcal{M}'_1 be a copy of M_{d-1} in \mathcal{H}'_1 and let \mathcal{M}'_2 be a matching in \mathcal{H}'_2 obtained from \mathcal{M}'_1 by the following transformation: for $e = uv \in \mathcal{M}'_1$, define $\psi(e) = \sigma(u)\sigma(v)$ where $\sigma(x) = x + (1, 0, 0, \ldots, 0)$. It is now easy to check that the new matching $M_d = \mathcal{M}'_1 \cup \mathcal{M}'_2$ has both properties we need.

We note that the formula of Proposition 14 does not hold for d = 2 and d = 3. For d = 2 the hypercube \mathcal{H}_2 is isomorphic to C_4 and thus $\gamma^{\text{EID}}(\mathcal{H}_2) = 3$. For d = 3, we note that an identifying code of size 4, if it exists, must be a matching with no pair of edges belonging to a 4-cycle. But this is not possible. An identifying code of size 5 is shown in Figure 4, therefore $\gamma^{\text{EID}}(\mathcal{H}_3) = 5$.

3. Lower Bounds

Recall from Theorem 4 that $\gamma^{\text{ID}}(G)$ is bounded below by a function of the order of G. As mentioned before, this bound is tight. Let \mathcal{C} be a set of c isolated vertices. We can build a graph G of order $2^c - 1$ such that \mathcal{C} is an identifying code of G. To this end, for every subset X of \mathcal{C} with $|X| \geq 2$, we associate a new vertex which is joined to all vertices in X and only to those vertices. Then, it is easily seen that \mathcal{C} is an identifying code of this graph. However, the graph built in this way is far from being a line graph as it contains $K_{1,t}$, even for large values of t. In fact this lower bound turns out to be far from being tight




Figure 3: Two disjoint edge-identifying codes of \mathcal{H}_4




Figure 4: An optimal edge-identifying code of \mathcal{H}_3

for the family of line graphs. In this section we give a tight lower bound on the size of an edge-identifying code of a graph in terms of the number of its edges. Equivalently we have a lower bound for the size of an identifying code in a line graph in terms of its order. This lower bound is of the order $\Theta(\sqrt{n})$ and thus is a much improved lower bound with respect to the general bound of Theorem 4.

Let G be a pendant-free graph and let C_E be an edge-identifying code of G. To avoid trivialities such as having isolated vertices we may assume G is connected. We note that this does not mean that the subgraph induced by C_E is also connected, in fact we observe almost the contrary, i.e. in most cases, an edge-identifying code of a minimum size will induce a disconnected subgraph of G. We first prove a lower bound for the case when an edge-identifying code induces a connected subgraph.

Theorem 15. If an edge-identifying code C_E of a nontrivial graph G induces a connected subgraph of G which is not isomorphic to K_2 , then G has at most $\binom{|\mathcal{C}_E|+2}{2} - 4$ edges. Furthermore, equality can only hold if \mathcal{C}_E induces a path.

Proof. Let G' be the subgraph induced by C_E . Since we assumed G' is connected, and since G' is pendantfree, it cannot have three vertices. Since we assumed G' $\cong K_2$, we conclude that G' has at least four vertices. For each vertex x of G', let C_E^x be the set of all edges incident to x in G'. Let e = uv be an edge of G, then one or both of u and v must be in V(G'). Therefore, depending on which of these vertices belong to C_E , e is uniquely determined by either C_E^u (if $u \in V(G')$ and $v \notin V(G')$), or C_E^v (if $u \notin V(G')$ and $v \in V(G')$), or $C_E^u \cup C_E^v$ (if both $u, v \in V(G')$). The total number of sets of this form can be at most $|V(G')| + \binom{|V(G')|}{2} = \binom{|V(G')|+1}{2}$, thus if $|V(G')| \leq |C_E|$ we are done. Otherwise, since G' is connected, $|V(G')| = |C_E| + 1$ and G' is a pendant-free tree on at least 4 vertices. If v is a vertex of degree 1 adjacent to u, then we have $C_E^v = \{uv\}$ but $uv \in C_E^u$ and, therefore, $C_E^u = C_E^u \cup C_E^v$. On the other hand, by Lemma 8, there are two vertices of degree 2 that have neighbours of degree 1. Let u be such a vertex, let v be its neighbour of degree 1 and x be its other neighbour. Then $C_E^v = \{uv\}$ and $C_E^u = \{uv, ux\}$ and, therefore, $C_E^u \cup C_E^x = C_E^v \cup C_E^x$. Thus the total number of distinct sets of the form $C_E^v or C_E^v \cup C_E^v$ is at most $\binom{|C_E|+2}{2} - 4$. But if equality holds there can only be two vertices of degree 1 in G' and hence C_E is a path. We note that if this bound is tight, then G' is a path. Furthermore, for each path P_{k+1} one can build many graphs which have P_{k+1} as an edge-identifying code and have $\binom{k+2}{2} - 4$ edges. The set of all these graphs will be denoted by \mathcal{J}_k . An example of such a graph is obtained from K_{k+2} by removing a certain set of four edges as shown in Figure 5. Note that every other member of \mathcal{J}_k is obtained from the previous example by splitting the vertex that does not belong to P_{k+1} (but without adding any new edge).




Figure 5: An extremal graph of \mathcal{J}_k with its connected edge-identifying code

Next we consider the case when the subgraph induced by C_E is not necessarily connected.

Theorem 16. Let G be a pendant-free graph and let C_E be an edge-identifying code of G with $|C_E| = k$. Then we have:

$$|E(G)| \le \begin{cases} \left(\frac{4}{3}k\right), & \text{if } k \equiv 0 \mod 3\\ \left(\frac{4}{3}(k-1)+1\right)+1, & \text{if } k \equiv 1 \mod 3\\ \left(\frac{4}{3}(k-2)+2\right)+2, & \text{if } k \equiv 2 \mod 3. \end{cases}$$

Proof. Let G be a graph with maximum number of edges among all graphs with $\gamma^{\text{EID}}(G) = k$. It can be easily checked that for k = 1, 2 or 3, the maximum number of edges of G is 1, 3 or 6 respectively. For $k \ge 4$, we prove a slightly stronger statement: given an edge-identifying code \mathcal{C}_E of G of size k, all but at most two of the connected components of the subgraph induced by \mathcal{C}_E must be isomorphic to P_4 . When there is only one component not isomorphic to P_4 , it must be isomorphic to a P_2 , a P_5 or a P_6 . If there are two such components, then they can be two copies of P_2 , a P_2 with a P_5 , or just two copies of P_5 . This depends on the value of k mod 3.

To prove our claim let G be a graph as defined above, let C_E be an edge-identifying code of size k of G and let G' be the subgraph induced by C_E . For each vertex $u \in V(G) \setminus V(G')$, we can assume that u has degree 1: if u has degree d > 1, with neighbours v_1, \ldots, v_d necessarily in V(G'), then replace u by d vertices of degree 1: u_1, \ldots, u_d , connecting u_i to v_i . Then the number of edges does not change, and the code C_E remains an edge-identifying code of size k, thus it suffices to prove our claim for this new graph. Let G'_1, G'_2, \ldots, G'_r be the connected components of G' with $|V(G'_i)| = n'_i$. For each $i \in \{1, \ldots, r\}$, let G_i be the graph induced by the vertices of G'_i and the vertices connected to G'_i only. To each vertex x of G' we assign the set \mathcal{C}^x_E of edges in G' incident to x.

We first note that no G'_i can be of order 3, because there is no connected pendant-free graph on three vertices. If u and v are vertices from two disjoint components of G' with each component being of order at least 4, then the pair u, v is uniquely determined by $\mathcal{C}^u_E \cup \mathcal{C}^v_E$, thus by maximality of G, uv is an edge of G. If a component of G' is isomorphic to K_2 , assuming u and u' are vertices of this component, then for any other vertex v of G' exactly one of uv or u'v is an edge of G.

We now claim that each G'_i with $n'_i \ge 4$ is a path. By contradiction, if a G'_i is not a path, we replace G_i by a member $J_{n'_i-1}$ of $\mathcal{J}_{n'_i-1}$ with $P_{n'_i}$ being its edge-identifying code. Then we join each vertex of $P_{n'_i}$ to each vertex of each G'_j (with $j \ne i$ and $n'_j \ge 4$) and to exactly one vertex of each G_j with $n'_j = 2$. We note that the new graph still admits an edge-identifying code of size k. However, it has more edges than G.

Indeed, while the number of edges connecting G'_i and the G'_j 's $(j \neq i)$ is not decreased, the number of edges in G_i is increased when we replace G_i by $J_{n'_i-1}$. This can be seen by applying Theorem 15 on G_i .

We now show that none of the G'_i 's can have more than six vertices. By contradiction, suppose G'_1 is a component with $n'_1 \ge 7$ vertices (thus $n'_1 - 1$ edges). We build a new graph G^*_1 from G as follows. We take disjoint copies of $J_3 \in \mathcal{J}_3$ and $J_{n'_1-4} \in \mathcal{J}_{n'_1-4}$ with P_4 and $P_{n'_1-3}$ being, respectively, their edge-identifying codes. We now let $V(G^*_1) = V(J_3) \cup V(J_{n'_1-4}) \cup (V(G) \setminus V(G_1))$. The edges of J_3 , $J_{n'_1-4}$ and $G - G_1$ are also edges of G^*_1 . We then add edges between these three parts as follows. We join every vertex of P_4 to each vertex of $P_{n'_1-3}$. For $i = 2, 3, \ldots, r$ if $n'_i \ge 4$, join every vertex of G'_i to each vertex of $P_4 \cup P_{n'_1-3}$. If $n'_i = 2$, we choose exactly one vertex of G'_i and join it to each vertex of $P_4 \cup P_{n'_1-3}$. The construction of G^*_1 ensures that it still admits an edge-identifying code of size k, but it has more edges than G. In fact, the number of edges is increased in two ways. First, because $P_4 \cup P_{n'_1-3}$ has one more vertex than G'_1 , the number of edges induced by $J_3 \cup J_{n'_1-4}$ is $6 + \binom{n'_1-2}{2} - 4 + 4 \times (n'_1-3) = \frac{n'_1^2}{2} + \frac{3n'_1}{2} - 7$ which is strictly

more than $|E(G'_1)| = \frac{n'_1^2}{2} + \frac{n'_1}{2} - 4$ for $n'_1 \ge 3$. Since $n'_1 \ge 7$, this contradicts the maximality of G.

With a similar method, the following transformations strictly increase the number of edges while the new graph still admits an edge-identifying code of size k:

- 1. Two components of G' each on six vertices transform into two graphs of \mathcal{J}_3 and a graph of \mathcal{J}_4 .
- 2. One component of G' on six vertices and another component on five vertices transform into three graphs of \mathcal{J}_3 .
- 3. One component of G' on six vertices and one on two vertices transform into two graphs of \mathcal{J}_3 .
- 4. Three components of G' each on five vertices transform into four graphs of \mathcal{J}_3 .
- 5. Two components of G' on five vertices and one on two vertices transform into three graphs of \mathcal{J}_3 .
- 6. A component of G' on five vertices and two on two vertices transform into two graphs of \mathcal{J}_4 .
- 7. Three components of G' each isomorphic to P_2 transform into a graph of \mathcal{J}_3 .

For the proof of case 7, we observe that the number of edges identified by the three P_2 's would be the same as the number of edges identified by the P_4 . However, since $k \ge 4$, there must be some other component in G'. Moreover, the number of vertices of the three P_2 's, which are joined to the vertices of the other components of G', is three, whereas the number of these vertices of the P_4 , is four. Hence the maximality of G is contradicted.

We note that cases 1, 2 and 3 imply that if a component of G' is isomorphic to P_6 , every other component is isomorphic to P_4 . Then cases 4, 5 and 6 imply that if a component is isomorphic to P_5 , then at most one other component is not isomorphic to P_4 and such component is necessarily either a P_2 or a P_5 . Finally, case 7 shows that there can be at most two components both isomorphic to P_2 .

We conclude that each of the components of G' is isomorphic to P_4 except for possibly two of them. These exceptions are dependent on the value of $k \mod 3$ as we described. The formulas of the theorem can be derived using these structural properties of G. For instance, in the case $k \equiv 0 \mod 3$, each component of G' is isomorphic to P_4 . There are $\frac{k}{3}$ such components. For each component G'_i , there are six edges in the graph G_i . That gives 2k edges. The other edges of G are edges between two components of G'. By maximality of G, between two components of G', there are exactly 16 edges. There are $\binom{k}{2}$ pairs of components of G'. Hence, the number of edges in G is:

$$2k + 16\binom{\frac{k}{3}}{2} = \binom{\frac{4}{3}k}{2}.$$

The other cases can be proved with the same method.

We note that this bound is tight and the examples were in fact built inside the proof. More precisely, for $k \equiv 0 \mod 3$ we take $\frac{k}{3}$ disjoint copies of elements of \mathcal{J}_3 each having a P_4 as an edge-identifying code. We then add an edge between each pair of vertices coming from two distinct such P_4 's. We note that the union of these P_4 's is a minimum edge-identifying code of the graph. If $k \not\equiv 0 \mod 3$, then we build a similar construction. This time we use elements from \mathcal{J}_3 with at most two exceptions that are elements of \mathcal{J}_4 or \mathcal{J}_5 .

The above theorem can be restated in the language of line graphs as follows.

Corollary 17. Let G be a twin-free line graph on $n \ge 4$ vertices. Then we have $\gamma^{ID}(G) \ge \frac{3\sqrt{2}}{4}\sqrt{n}$.

Proof. Suppose G is the line graph of a pendant-free graph H ($\mathcal{L}(H) = G$). Let $k = \gamma^{\text{ID}}(G) = \gamma^{\text{EID}}(H)$, and let n be the number of vertices of G (n = |E(H)|). Then, after solving the quadratic inequalities of Theorem 16 for k, we have:

$$\begin{split} k \geq &\frac{3}{8} + \frac{3\sqrt{8n+1}}{8}, \text{ for } k \equiv 0 \mod 3, \\ k \geq &\frac{5}{8} + \frac{3\sqrt{8n-7}}{8}, \text{ for } k \equiv 1 \mod 3, \\ k \geq &\frac{3}{8} + \frac{3\sqrt{8n-15}}{8}, \text{ for } k \equiv 2 \mod 3. \end{split}$$

It is then easy to check that the right-hand side of each of the three inequalities is at least as $\frac{3\sqrt{2}}{4}\sqrt{n}$ for $n \ge 3$.

Remark. Note that the lower bound of $\gamma^{\text{ID}}(G) \geq \Theta(\sqrt{|V(G)|})$, which holds for the class of line graphs, is also implied by Theorem 12. However, the bound of 17 is more precise. In [2], Beineke characterized line graphs by a list of nine forbidden induced subgraphs. Considering Beineke's characterization, the lower bound of Corollary 17 can be restated as follows: $\gamma^{\text{ID}}(G) \geq \Theta(\sqrt{|V(G)|})$ holds if G has no induced subgraph from Beineke's list. It is then natural to ask what is a minimal list of forbidden induced subgraphs for which a similar claim would hold. Note that the claw graph, $K_{1,3}$, belongs to Beineke's list of forbidden subgraphs. However, we remark that the bound $\gamma^{\text{ID}}(G) \geq \Theta(\sqrt{|V(G)|})$ does not hold for the class of claw-free graphs. Examples can be built as follows: let A be a set of size k and let B be the set of nonempty subsets of A. Let G be the graph built on $A \cup B$, where A and B each induce a complete graph and a vertex a of A is joined to a vertex b of B if $a \in b$. This graph is claw-free and it is easy to find an identifying code of size at most $2k = \Theta(\log |V(G)|)$ in G.

4. Upper bounds

The most natural question in the study of identifying codes in graphs is to find an identifying code as small as possible. A general bound, only in terms of the number of vertices of a graph, is provided by Theorem 2. Furthermore, the class of all graphs with $\gamma^{\text{ID}}(G) = |V(G)| - 1$ is classified in [8]. It is easy to check that none but six of these graphs are line graphs. Thus we have the following corollary (where $G \bowtie H$ denotes the complete join of graphs G and H):

Corollary 18. If G is a twin-free line graph with $G \notin \{P_3, P_4, C_4, P_4 \bowtie K_1, C_4 \bowtie K_1, \mathcal{L}(K_4)\}$, then we have $\gamma^{ID}(G) \leq |V(G)| - 2$.

Since $\gamma^{\text{EID}}(K_{2,n}) = 2n - 2$, $\gamma^{\text{ID}}(\mathcal{L}(K_{2,n})) = |V(\mathcal{L}(K_{2,n}))| - 2$ and the bound of Corollary 18 is tight for an infinite family of graphs. Conjecture 3 proposes a better bound in terms of both the number of vertices and the maximum degree of a graph. As pointed out in Proposition 11, most of the known extremal graphs for Conjecture 3 are line graphs. In this section, after proving some general bound for the edge-identifying code number of a pendant-free graph we will show that Conjecture 3 holds for the class of line graphs of high enough density.

We recall that a graph on *n* vertices is 2-degenerated if its vertices can be ordered v_1, v_2, \ldots, v_n such that each vertex v_i is joined to at most two vertices in $\{v_1, v_2, \ldots, v_{i-1}\}$. Our main idea for proving upper bounds is to show that given a pendant-free graph G, any (inclusionwise) minimal edge-identifying code C_E induces a 2-degenerated subgraph of G and hence $|\mathcal{C}_E| \leq 2|V(G)| - 3$. Our proofs are constructive and one could build such small edge-identifying codes.

Theorem 19. Let G be a pendant-free graph and let C_E be a minimal edge-identifying code of G. Then G', the subgraph induced by C_E , is 2-degenerated.

Proof. Let uv be an edge of G' with $d_{G'}(u), d_{G'}(v) \geq 3$. By minimality of \mathcal{C}_E the subset $\mathcal{C}' = \mathcal{C}_E - uv$ of E(G) is not an edge-identifying code of G. By the choice of uv, \mathcal{C}' is still an edge-dominating set, thus there must be two edges, e_1 and e_2 , that are not separated by \mathcal{C}' . Hence one of them, say e_1 , is incident either to u or to v (possibly to both) and the other one (e_2) is incident to neither one.

We consider two cases: either $e_1 = uv$ or e_1 is incident to only one of u and v. In the first case, e_2 is adjacent to every edge of \mathcal{C}' which uv is adjacent to. Since for each vertex of uv there are at least two edges in \mathcal{C}' incident to this vertex, the subgraph induced by u, v and the vertices of e_2 must be isomorphic to K_4 and there should be no other edge of \mathcal{C}' incident to any vertex of this K_4 (see Figure 6(a)).

In the other case, suppose e_1 is adjacent to uv at u. Let x and y be two neighbours of u in G' other than v. Then it follows that $e_2 = xy$ and, therefore, $d_{G'}(u) = 3$. Let z be the other end of e_1 . We consider two subcases: either $z \notin \{x, y\}$, or, without loss of generality, z = x. Suppose $z \notin \{x, y\}$. Recall that uv is the only edge separating e_1 and e_2 , but e_1 must be separated from ux. Thus $zy \in \mathcal{C}_E$. Similarly, e_1 must be separated from uy, so $zx \in \mathcal{C}_E$. Furthermore, $d_{G'}(x) = d_{G'}(y) = d_{G'}(z) = 2$ and $\{x, y, z, u\}$ induces a C_4 in G' (see Figure 6(b)). Now suppose $e_1 = ux$, since uv is the only edge separating e_1 and e_2 , then uy and possibly xy are the only edges in G' incident to y, so $d_{G'}(y) \leq 2$ and $d_{G'}(u) = 3$ (see Figures 6(c) and 6(d)).




Figure 6: Case distinctions in the proof of Theorem 19. Black vertices have fixed degree in G'. Thick edges belong to C_E .

To summarize, we proved that given an edge uv, in a minimal edge-identifying code C_E , we have one of the following cases.

- One of u or v is of degree at most 2 in G'.
- Edge uv is an edge of a connected component of G' isomorphic to K_4^- (that is K_4 with an edge removed), see Figure 6(a).
- $d_{G'}(u) = 3$ (considering the symmetry between u and v) in which case either u is incident to a C_4 whose other vertices are of degree 2 in G' (Figure 6(b)), or to a vertex of degree 1 in G' (Figure 6(c)) or to a triangle with one vertex y of degree 2 in G' and y is not adjacent to v (Figure 6(d)).

In either case, there exists a vertex x of degree at most 2 in G' such that when x is removed, at least one of the vertices u, v has degree at most 2 in the remaining subgraph of G'. In this way we can define an order of elimination of the vertices of G' showing that G' is 2-degenerated.

By further analysis of our proof we prove the following:

Corollary 20. If G is a pendant-free graph on n vertices not isomorphic to K_4 or K_4^- , then $\gamma^{\text{EID}}(G) \leq 2n-5$.

Proof. We first prove that if G is a pendant-free graph on n vertices not isomorphic to K_4 , then $\gamma^{\text{EID}}(G) \leq 2n-4$. Let \mathcal{C}_E be a minimal edge-identifying code and let G' be the subgraph induced by \mathcal{C}_E . Then, by Theorem 19, G' is 2-degenerated. Let $v_n, v_{n-1}, \ldots, v_1$ be a sequence of vertices of G' obtained by a process of eliminating vertices of degree at most 2. Since v_1 and v_2 can induce at most a K_2 , we notice that there could only be at most 2n-3 edges in G'. Furthermore, if there are exactly 2n-3 edges in G', then $v_1v_2 \in \mathcal{C}_E$ and each vertex $v_i, 3 \leq i \leq n$, has exactly two neighbours in $\{v_1, \ldots, v_{i-1}\}$. Hence, the subgraph induced by $\{v_1, v_2, v_3, v_4\}$ is isomorphic to K_4^- . Considering symmetries, there are three possibilities for the subgraph induced by $\{v_1, \ldots, v_5\}$ (recall that v_5 is of degree 2 in this subgraph): see Figure 7. In each of these three cases, the edge uv has both ends of degree at least 3. Thus, we can apply the argument used in the proof of Theorem 19 on G' and uv, showing that we have one of the four configurations of Figure 6. But none of them matches with the configurations of Figure 7, a contradiction.




Figure 7: The three maximal 2-degenerated graphs on five vertices

Now we show that if $\gamma^{\text{EID}}(G) = 2n - 4$, then $G \cong K_4^-$. This can be easily checked if G has at most four vertices, so we may assume $n \ge 5$. Let G'' be the subgraph of G' induced by $\{v_1, v_2, v_3, v_4, v_5\}$. If G'' has seven edges, then it is isomorphic to one of the graphs of Figure 7, and we are done just like in the last case. Therefore, we can assume that G'' has exactly six edges and, since it is 2-degenerated, by an easy case analysis, it must be isomorphic to one of the graphs of Figure 8.




Figure 8: The five possibilities of 2-degenerated graphs on five vertices with six edges

If G'' is a graph in part (i), (ii) or (iii) of Figure 8, then again one could repeat the arguments of the proof of Theorem 19 with G' and the edge uv of the corresponding figure, to obtain a contradiction.

Suppose G'' is isomorphic to the graph of Figure 8(iv). Since G'' is not pendant-free, there must be at least one more vertex in G'. Let v_6 be as in the sequence obtained by the 2-degeneracy of G'. Since G' has exactly 2n - 4 edges, v_6 must have exactly two neighbours in G''. By the symmetry of the four vertices of degree 2 in G'', we may assume $uv_6 \in C_E$. Then u and v are both of degree at least 3 in G'. Therefore, we could again repeat the argument of Theorem 19 with G' and uv, where only one of the configurations of this theorem, namely 6(d), matches G''. Furthermore, if this happens then $v'v_6$ should also be an edge of G'. Now u and v' are both of degree at least 3 and we apply the argument of Theorem 19 with G' and uv' to obtain a contradiction.

Finally, let G'' be isomorphic to the graph of Figure 8(v). We claim that every other vertex v_i $(i \ge 6)$ is adjacent, in G', only to u and v. By contradiction suppose v_6 is adjacent to t. Then using the technique of Theorem 19 applied on G' and tu (respectively tv), we conclude that v_6 is adjacent to u (respectively v).

Since $|E(G')| = |\mathcal{C}_E| = 2n - 4$, G' is a spanning subgraph of G. But then it is easy to verify that $\mathcal{C}_E \setminus \{xu, xv\}$ is an edge-identifying code of G — a contradiction.

We note that $\gamma^{\text{EID}}(K_{2,n}) = 2n - 2 = 2|V(K_{2,n})| - 6$ thus this bound cannot be improved much. Corollary 20 implies that Conjecture 3 holds for a large subclass of line graphs:

Corollary 21. If G is a pendant-free graph on n vertices and with average degree $\bar{d}(G) \geq 5$, then we have $\gamma^{ID}(\mathcal{L}(G)) \leq n - \frac{n}{\Delta(\mathcal{L}(G))}$.

Proof. Let u be a vertex of degree $d(u) \ge \bar{d}(G) \ge 5$. Since G is pendant-free there is at least one neighbour v of u that is of degree at least 2. Thus there is an edge uv in G with $d(u) + d(v) \ge \bar{d}(G) + 2$ and, therefore, $\Delta(\mathcal{L}(G)) \ge \bar{d}(G)$. Hence, considering Corollary 20, it is enough to show that $2|V(G)| - 5 \le |E(G)| - \frac{|E(G)|}{\bar{d}(G)}$.

To this end, since $\bar{d}(G) \ge 5$, we have $4|V(G)| \le (\bar{d}(G) - 1)|V(G)|$, therefore,

$$4|V(G)| - 10 \le (d(G) - 1)|V(G)|.$$

Mutiplying both sides by $\frac{\overline{d}(G)}{2}$ we have:

$$(2|V(G)| - 5)\overline{d}(G) \le (\overline{d}(G) - 1)\frac{\overline{d}(G)}{2}|V(G)| = (\overline{d}(G) - 1)|E(G)|.$$

5. Complexity

This section is devoted to the study of the decision problem associated to the concept of edge-identifying codes. Let us first define the decision problems we use. The IDCODE problem is defined as follows:

IDCODE

INSTANCE: A graph G and an integer k. QUESTION: Does G have an identifying code of size at most k?

IDCODE was proved to be NP-complete even when restricted to the class of bipartite graphs of maximum degree 3 (see [4]) or to the class of planar graphs of maximum degree 4 and arbitrarily large girth (see [1]). The EDGE-IDCODE problem is defined as follows:

EDGE-IDCODE

INSTANCE: A graph G and an integer k. QUESTION: Does G have an edge-identifying code of size at most k?

We will prove that EDGE-IDCODE is NP-hard in some restricted class of graphs by reduction from PLANAR ($\leq 3, 3$)-SAT, which is a variant of the SAT problem and is defined as follows [7]:

PLANAR ($\leq 3, 3$)-SAT

INSTANCE: A collection \mathcal{Q} of clauses over a set X of boolean variables, where each clause contains at least two and at most three distinct literals (a variable x or its negation \overline{x}). Moreover, each variable appears in exactly three clauses: twice in its non-negated form, and once in its negated form. Finally, the bipartite incidence graph of \mathcal{Q} , denoted $B(\mathcal{Q})$, is planar ($B(\mathcal{Q})$ has vertex set $\mathcal{Q} \cup X$ and $\mathcal{Q} \in \mathcal{Q}$ is adjacent to $x \in X$ if x or \overline{x} appears in clause \mathcal{Q}).

QUESTION: Can Q be satisfied, i.e. is there a truth assignment of the variables of X such that each clause contains at least one true literal?

PLANAR ($\leq 3, 3$)-SAT is known to be NP-complete [7]. We are now ready to prove the main result of this section.

Theorem 22. EDGE-IDCODE is NP-complete even when restricted to bipartite planar graphs of maximum degree 3 and arbitrarily large girth.

Proof. The problem is clearly in NP: given a subset C of edges of G, one can check in polynomial time whether it is an edge-identifying code of G by computing the sets $C \cap N[e]$ for each edge e and comparing them pairwise.

We now reduce PLANAR ($\leq 3, 3$)-SAT to EDGE-IDCODE. We first give the proof for the case of girth 8 and show that it can be easily extended to an arbitrarily large girth.

We first need to define a generic sub-gadget (denoted P-gadget) that will be needed for the reduction. In order to have more compact figures, we will use the representation of this gadget as drawn in Figure 9. We will say that a P-gadget is *attached* at some vertex x if x is incident to edge a of the gadget as depicted in the figure. When speaking of a P-gadget as a subgraph of a graph G, we always mean that it forms an induced subgraph of G, that is, there are no other edges within the gadget than $\{a, b, c, d, e\}$ in Figure 9. Moreover, vertex x is the only vertex of the P-gadget which may be joined by an edge to other vertices outside the gadget.




Figure 9: The generic P-gadget

We make the following claims.

Claim 1. In any graph containing a P-gadget, at least three edges of this gadget must belong to any edgeidentifying code.

Claim 1 is true because d is the only edge separating b and c. Similarly c is the only edge separating d and e. Finally, in order to separate d and c, one has to take at least one of a, b or e.

Claim 2. If G is a pendant-free graph obtained from a graph H with a P-gadget attached at a vertex x of H, then any edge-identifying code of G must contain an edge of H incident to x.

Claim 2 follows from the fact that edge a must be separated from edge b.

We are now ready to describe the reduction.

Given an instance $\mathcal{Q} = \{Q_1, \ldots, Q_m\}$ of PLANAR ($\leq 3, 3$)-SAT over the set of boolean variables $X = \{x_1, \ldots, x_n\}$ together with an embedding of its bipartite incidence graph $B(\mathcal{Q})$ in the plane, we build the graph $G_{\mathcal{Q}}$ as follows.

For each variable x_j and clause Q_i we build the subgraphs G_{x_j} and G_{Q_i} respectively, as shown in Figure 10. We recall that a given variable x_j appears in positive form in exactly two clauses, say Q_p , Q_q , and in negative form in exactly one clause, say Q_r . We then unify^{*} vertex x_j^1 of G_{x_j} with vertex l_{p_k} of G_{Q_p} which corresponds to x_j . We do a similar unification for vertices x_j^2 and $\overline{x_j}^1$ with corresponding vertices from G_{Q_q} and G_{Q_r} . The intuition is that vertices of the form l_{i_j} in the clause gadgets will represent literals of the clauses, and vertices of the form x_i^j , $\overline{x_i}^j$ of the vertex gadgets represent positive and negative occurences of a variable, respectively.

This can be done while ensuring the planarity of $G_{\mathcal{Q}}$, using the given planar embedding of $B(\mathcal{Q})$. Moreover, $G_{\mathcal{Q}}$ is bipartite because $B(\mathcal{Q})$ is bipartite, there are no odd cycles in the variable and clause gadgets and there is no path of odd length between l_{i_j} 's. Finally, it is easy to see that $G_{\mathcal{Q}}$ has maximum degree 3 and girth 8. Since a clause gadget has fourty-five vertices and a variable gadget, fourty-two vertices, $G_{\mathcal{Q}}$ has 45m + 42n vertices and, therefore, the construction has polynomial size in terms of the size of \mathcal{Q} .




Figure 10: Reduction gadgets for clause Q_i and variable x_j

We will need two additional claims in order to complete the proof.

Claim 3. In a variable gadget G_{x_j} , in order to separate the four pairs of edges $\{d_i, e_i\}$ for $1 \le i \le 4$, at least two edges of $A = \{d_i, e_i \mid 1 \le i \le 4\} \cup \{t_j^1, \overline{t_j}^1, t_j^2, \overline{t_j}^2\}$ belong to any edge-identifying code C. Moreover, if $|C \cap A| = 2$, then either $C \cap A = \{t_j^1, t_j^2\}$ or $C \cap A = \{\overline{t_j}^1, \overline{t_j}^2\}$.

^{*}We use the term "unify" instead of the usual term "identify" in order to avoid confusion with identifying codes.

The first part of Claim 3 follows from the fact that the two edges of each of the pairs $\{d_1, e_1\}$ and $\{d_3, e_3\}$ must be separated. The second follows from an easy case analysis.

The following claim follows directly from Claim 2.

Claim 4. Let $v_1v_2v_3v_4$ be a path of four vertices of G_Q where each of the vertices v_2 and v_3 has its own P-gadget attached and both v_2 and v_3 have degree 3. Then, at least one of the three edges of the path belong to any identifying code of the graph. If exactly one belongs to a code, it must be v_2v_3 .

We now claim that Q is satisfiable if and only if G_Q has an edge-identifying code of size at most k = 25m + 22n.

For the sufficient side, given a truth assignment of the variables satisfying \mathcal{Q} , we build an edge-identifying code \mathcal{C} as follows. For each P-gadget, edges a, c, d are in \mathcal{C} . For each clause gadget G_{Q_i} , edge c_0 belongs to \mathcal{C} . For each literal l_{i_k} of Q_i , $1 \leq k \leq 3$, if l_{i_k} is true, edge a_{i_k} belongs to \mathcal{C} ; otherwise, edge b_{i_k} belongs to \mathcal{C} . If Q_i has only two literals and vertex l_{i_k} is the vertex not corresponding to a literal of Q_i , then edge b_{i_k} belongs to \mathcal{C} . Now, one can see that all edges of G_{Q_i} are dominated. Furthermore, all pairs of edges of G_{Q_i} are separated. This can be easily seen for all pairs besides $\{c_1, c_2\}$. For this pair, since we are considering a satisfying assignment of \mathcal{Q} , in every clause Q_i of \mathcal{Q} , there exists a true literal. Hence, for each clause Q_i , at least one edge a_{i_j} with $1 \leq j \leq 3$, must be in the code and, therefore, the pair $\{c_1, c_2\}$ is separated.

Next, in each variable gadget G_{x_j} , if x_j is true, edges t_j^1 and t_j^2 belong to \mathcal{C} . Otherwise, edges $\overline{t_j}^1$ and $\overline{t_j}^2$ belong to \mathcal{C} . Edges f_1, f_2, f_3, f_4 and f_5 also belong to \mathcal{C} . Because of this choice, all edges of $G_{x_j} \setminus \{t_j^1, t_j^2, \overline{t_j}^1\}$ are dominated. Since each of the three edges $t_j^1, t_j^2, \overline{t_j}^1$ is incident to a vertex of a *P*-gadget of some clause gadget, they are also dominated. Moreover, all pairs of edges containing at least one edge of $G_{x_j} \setminus \{t_j^1, t_j^2, \overline{t_j}^1\}$ are clearly separated. Now, since for each *P*-gadget of the clause gadgets, edge *a* is in \mathcal{C} , $t_j^1, t_j^2, \overline{t_j}^1$ are separated from all edges in $G_{\mathcal{Q}}$.

We conclude that C is an edge-identifying code of size k.

For the necessary side, let \mathcal{C}' be an edge-identifying code of $G_{\mathcal{Q}}$ with $|\mathcal{C}'| \leq k$. It follows from Claim 1 that at least three edges of each of the seven P-gadgets of a clause gadget G_{Q_i} must belong to \mathcal{C}' . Moreover, by Claim 2, edge c_0 is forced to be in any code. Finally, by Claim 2, for each vertex l_{i_k} $(1 \leq k \leq 3)$ of G_{Q_i} , at least one of the edges a_{i_k} and b_{i_k} is in \mathcal{C}' .

Note that this is a total of at least twenty-five edges per clause gadget.

Similarly, it follows from Claim 1 that in each variable gadget G_{x_j} , at least fifteen edges of \mathcal{C}' are contained in the *P*-gadgets of G_{x_j} . Following Claim 2, all edges f_i $(1 \le i \le 5)$ belong to \mathcal{C}' . Note that this is a total of at least twenty edges in each variable gadget. We have considered 25m + 20n edges of \mathcal{C}' so far. Hence 2n edges remain to be considered. It follows from Claim 3 that for each variable gadget, at least two additional edges belong to \mathcal{C}' (in order to separate the pairs $\{d_i, e_i\}$, for $1 \le i \le 4$). Therefore, since $|\mathcal{C}'| \le k$, in each variable gadget, exactly two of these edges belong to \mathcal{C}' . Hence, following the second part of Claim 3, either $\{t_j^1, t_j^2\}$ or $\{\overline{t_j}^1, \overline{t_j}^2\}$ is a subset of \mathcal{C}' .

Remark that we have now considered all k = 25m + 22n edges of \mathcal{C}' . Therefore, in each clause gadget G_{Q_i} , exactly one of the edges a_{i_k} and b_{i_k} of G_{Q_i} belongs to \mathcal{C}' . We can now build the following truth assignment: for each variable gadget, if $\{t_j^1, t_j^2\}$ is a subset of \mathcal{C}' ,

We can now build the following truth assignment: for each variable gadget, if $\{t_j^1, t_j^2\}$ is a subset of C', x_j is set to TRUE. Otherwise, $\{\overline{t_j}^1, \overline{t_j}^2\}$ is a subset of C' and x_j is set to FALSE. Let us prove that this assignment satisfies Q.

In each clause gadget G_{Q_i} , note that edges c_1 and c_2 must be separated by \mathcal{C}' ; this means that one edge a_{i_k} from $\{a_{i_1}, a_{i_2}, a_{i_3}\}$ belongs to \mathcal{C}' . Hence, as noted in the previous paragraph, $b_{i_k} \notin \mathcal{C}'$ and by Claim 4, in the path formed by edges $\{a_{i_k}, b_{i_k}, t_j^1\}$, t_j^1 belongs to the code (without loss of generality, we suppose that $l_{i_k} = x_j$ and t_j^1 is the edge of G_{x_j} incident to vertex l_{i_k} of G_{Q_i}). Therefore, in the constructed truth assignment, literal l_{i_k} has value TRUE, and the clause is satisfied. Repeating this argument for each clause shows that the formula is satisfied.

Now, it remains to show that similar arguments can be used to prove the final statement of the theorem for larger girth. Consider some integers $\lambda \geq 1$ and $\mu \geq 2$. We build the graph $G_Q(\lambda, \mu)$ using modified variable gadgets $G_{x_j}(\mu)$ and modified clause gadgets $G_{Q_i}(\lambda)$, which are depicted in Figure 11. The construction is the same as in the previous proof and $G_Q(\lambda, \mu)$ has $(36\lambda + 9)m + (30\mu - 18)n$ vertices. We claim that the girth of $G_Q(\lambda, \mu)$ is now at least min $\{4\mu, 8(\lambda + 1)\}$. Indeed, $G_{x_j}(\mu)$ has a cycle of size exactly 4μ and since the girth of B(Q) is at least 4, it follows that the minimum length of a cycle between some clause gadgets (at least two) and some variable gadgets (at least two) is at least $4(2\lambda + 1) + 2 + 2 = 8(\lambda + 1)$.

Now, using a similar proof as the proof for girth 8, it can be shown that Q is satisfiable if and only if $G_Q(\lambda,\mu)$ has an identifying code of size at most $k = (21\lambda + 4)m + (17\mu - 12)n$.

Recall that a graph is *perfect* if and only if for each of its induced subgraphs H, the chromatic number of H equals the clique number of H. It is known that a line graph $\mathcal{L}(G)$ is perfect if and only if G has no odd cycles of length more than 3, see [18]. Moreover, one can check that the line graphs of the graphs constructed in the previous proof are planar, have maximum degree 4 and clique number 3. Therefore, the following corollary follows:

Corollary 23. *IDCODE is NP-complete even when restricted to perfect 3-colorable planar line graphs of maximum degree 4.*

Note that by Theorem 12 and Corollary 20, we have $\frac{|V(G)|}{2} \leq \gamma^{\text{EID}}(G) \leq |\mathcal{C}_E| \leq 2|V(G)| - 3$ for any pendant-free graph G and any inclusionwise minimal edge-identifying code \mathcal{C}_E of G. Since one can construct such a code in polynomial time, this gives a polynomial-time 4-approximation algorithm for the optimization problem associated to EDGE-IDCODE:

Theorem 24. The optimization problems associated to EDGE-IDCODE in general graphs and to IDCODE when restricted to line graphs are 4-approximable in polynomial-time.

We remark that it is NP-hard to approximate the optimization version of IDCODE within a factor of $o(\log(n))$ in general graphs on n vertices (see [14, 17]).

In the following, by slightly restricting the class of graphs considered in Theorem 22, we show that EDGE-IDCODE becomes linear-time solvable in this restricted class.


Let us first introduce some necessary concepts.

A graph property \mathcal{P} is expressable in *counting monadic second-order logic*, CMSOL for short (see [6] for further reference), if \mathcal{P} can be defined using:

- vertices, edges, sets of vertices and sets of edges of a graph
- the binary adjacency relation adj where adj(u, v) holds if and only if u, v are two adjacent vertices
- the binary incidence relation *inc*, where inc(v, e) holds if and only if edge e is incident to vertex v
- the equality operator = for vertices and edges
- the membership relation \in , to check whether an element belongs to a set
- the unary cardinality operator *card* for sets of vertices
- the logical operators OR, AND, NOT (denoted by \lor, \land, \neg)
- the logical quantifiers \exists and \forall over vertices, edges, sets of vertices or sets of edges

It has been shown that CMSOL is particularly useful when combined with the concept of the graph parameter *tree-width* (we refer the reader to [6] for a definition). Some important classes of graphs have bounded tree-width. For example, trees have tree-width at most 1, series-parallel graphs have tree-width at most 2 and outerplanar graphs have tree-width at most 3.

The following result shows that many graph properties can be checked in linear time for graphs of bounded tree-width.



(b) Variable gadget $G_{x_j}(\mu)$

Figure 11: Reduction gadgets for clause Q_i and variable x_j for arbitrarily large girth

Theorem 25 ([6]). Let \mathcal{P} be a graph property expressable in CMSOL and let c be a constant. Then, for any graph G of tree-width at most c, it can be checked in linear time whether G has property \mathcal{P} .

We now show that CMSOL can be used in the context of edge-identifying codes:

Proposition 26. Given a graph G and an integer k, let $\mathcal{EID}(G, k)$ be the property that $\gamma^{\text{EID}}(G) \leq k$. Property $\mathcal{EID}(G, k)$ can be expressed in CMSOL. *Proof.* Let V = V(G) and E = E(G). We define the CMSOL relation dom(e, f) which holds if and only if e, f are edges of E and e, f dominate each other, i.e. e and f are incident to the same vertex. We have $dom(e, f) := \exists x \in V, (inc(x, e) \land inc(x, f)).$

Now we define $\mathcal{EID}(G, k)$ as follows:

$$\begin{split} \mathcal{EID}(G,k) &:= \quad \exists C, C \subseteq E, card(C) \leq k, \left(\forall e \in E, \exists f \in C, dom(e,f) \right) \land \\ & \left(\forall e \in E, \forall f \in E, e \neq f, \exists g \in C, \left((dom(e,g) \land \neg dom(f,g)) \lor (dom(f,g) \land \neg dom(e,g)) \right) \right). \end{split}$$

This together with Theorem 25 implies the following corollary.

Corollary 27. *EDGE-IDCODE* can be solved in linear time for all classes of graphs having their tree-width bounded by a constant.

This result implies, in particular, that one can find the edge-identifying code number of a tree in linear time. Note that a similar approach has been used in [15] to show that this holds for IDCODE as well.

The proof of Theorem 25 is constructive and gives a linear-time algorithm, but it is very technical and hides a large constant depending on the size of the CMSOL expression. Therefore, it would be interesting to give a simpler and more practical linear-time algorithm for EDGE-IDCODE in trees. Observe that this has been done in [1] for the case of vertex-identifying codes.

References

- D. Auger. Minimal identifying codes in trees and planar graphs with large girth, European Journal of Combinatorics 31(5):1372–1384, 2010.
- [2] L. W. Beineke. Characterizations of derived graphs, Journal of Combinatorial Theory 9(2)2:129–135, 1970.
- [3] I. Charon, G. Cohen, O. Hudry and A. Lobstein. New identifying codes in the binary Hamming space, European Journal of Combinatorics 31(2):491–501, 2010.
- [4] I. Charon, O. Hudry and A. Lobstein. Minimizing the size of an identifying or locating-dominating code in a graph is NP-hard. *Theoretical Computer Science* 290(3):2109–2120, 2003.
- [5] I. Charon, O. Hudry and A. Lobstein. Extremal cardinalities for identifying and locating-dominating codes in graphs. Discrete Mathematics 307(3-5):356–366, 2007.
- [6] B. Courcelle. The monadic second-order logic of graphs. I. Recognizable sets of finite graphs. Information and Computation 85(1):12–75, 1990.
- [7] E. Dahlhaus, D. S. Johnson, C. H. Papadimitriou, P. D. Seymour and M. Yannakakis. The complexity of multiterminal cuts. SIAM Journal of Computing 23(4):864–894, 1994.
- [8] F. Foucaud, E. Guerrini, M. Kovše, R. Naserasr, A. Parreau and P. Valicov. Extremal graphs for the identifying code problem. *European Journal of Combinatorics* 32(4):628–638, 2011.
- [9] F. Foucaud, R. Klasing, A. Kosowski and A. Raspaud. On the size of identifying codes in triangle-free graphs. Discrete Applied Mathematics 160(10-11):1532–1546, 2012.
- [10] F. Foucaud and G. Perarnau. Bounds on identifying codes in terms of degree parameters. The Electronic Journal of Combinatorics 19:P32, 2012.
- [11] S. Gravier, R. Klasing and J. Moncel. Hardness results and approximation algorithms for identifying codes and locatingdominating codes in graphs. Algorithmic Operations Research 3(1):43–50, 2008.
- [12] S. Gravier and J. Moncel. On graphs having a $V \setminus \{x\}$ set as an identifying code. Discrete Mathematics 307(3-5):432–434, 2007.
- [13] M. G. Karpovsky, K. Chakrabarty and L. B. Levitin. On a new class of codes for identifying vertices in graphs. *IEEE Transactions on Information Theory* 44:599–611, 1998.
- [14] M. Laifenfeld, A. Trachtenberg and T. Y. Berger-Wolf. Identifying codes and the set cover problem. Proceedings of the 44th Annual Allerton Conference on Communication, Control and Computing, Monticello, USA, September 2006.
- [15] J. Moncel. Codes Identifiants dans les Graphes. PhD Thesis, Université Joseph-Fourier Grenoble I, France, June 2005. Available online at http://tel.archives-ouvertes.fr/tel-00010293.
- [16] J. Moncel. On graphs on n vertices having an identifying code of cardinality $\lceil \log_2(n+1) \rceil$. Discrete Applied Mathematics 154(14):2032–2039, 2006.

- [17] J. Suomela. Approximability of identifying codes and locating-dominating codes. Information Processing Letters 103(1):28–33, 2007.
- [18] L. E. Trotter. Line perfect graphs. Mathematical Programming 12(1):255–259, 1977.