Injective tests of low complexity in the plane

Dominique LECOMTE and Rafael ZAMORA¹

July 3, 2021

 Université Paris 6, Institut de Mathématiques de Jussieu, Projet Analyse Fonctionnelle Couloir 16-26, 4ème étage, Case 247, 4, place Jussieu, 75 252 Paris Cedex 05, France dominique.lecomte@upmc.fr

Université de Picardie, I.U.T. de l'Oise, site de Creil, 13, allée de la faïencerie, 60 107 Creil, France

 ¹ Université Paris 6, Institut de Mathématiques de Jussieu, Projet Analyse Fonctionnelle Couloir 15-16, 5ème étage, Case 247, 4, place Jussieu, 75 252 Paris Cedex 05, France rafael.zamora@imj-prg.fr

Abstract. We study injective versions of the characterization of sets potentially in a Wadge class of Borel sets, for the first Borel and Lavrentieff classes. We also study the case of oriented graphs in terms of continuous homomorphisms, injective or not.

²⁰¹⁰ Mathematics Subject Classification. Primary: 03E15, Secondary: 26A21, 54H05

Keywords and phrases. acyclic, Borel, class, dichotomy, difference, homomorphism, injective, locally countable, oriented graph, reduction, Wadge

1 Introduction

The reader should see [K] for the standard descriptive set theoretic notation used in this paper. This work is a contribution to the study of analytic subsets of the plane. We are looking for results of the following form: either a situation is simple, or it is more complicated than a situation in a collection of known complicated situations. The notion of complexity we consider is the following, and defined in [Lo3].

Definition 1.1 (Louveau) Let X, Y be Polish spaces, B be a Borel subset of $X \times Y$, and Γ be a class of Borel sets closed under continuous pre-images. We say that B is **potentially in** Γ (denoted $B \in pot(\Gamma)$) if there are finer Polish topologies σ and τ on X and Y, respectively, such that B, viewed as a subset of the product $(X, \sigma) \times (Y, \tau)$, is in Γ .

The quasi-order \leq_B of Borel reducibility was intensively considered in the study of analytic equivalence relations during the last decades. The notion of potential complexity is a natural invariant for \leq_B : if $E \leq_B F$ and $F \in \text{pot}(\Gamma)$, then $E \in \text{pot}(\Gamma)$ too. However, as shown in [L1]-[L6] and [L8], \leq_B is not the right notion of comparison to study potential complexity, in the general context, because of cycle problems. A good notion of comparison is as follows. Let X, Y, X', Y' be topological spaces and $A, B \subseteq X \times Y, A', B' \subseteq X' \times Y'$. We write

$$\begin{split} (X,Y,A,B) &\leq (X',Y',A',B') \Leftrightarrow \\ & \exists f \colon X \to X' \ \exists g \colon Y \to Y' \text{ continuous with } A \subseteq (f \times g)^{-1}(A') \text{ and } B \subseteq (f \times g)^{-1}(B'). \end{split}$$

Our motivating result is the following (see [L8]).

Definition 1.2 We say that a class Γ of subsets of zero-dimensional Polish spaces is a Wadge class of Borel sets if there is a Borel subset \mathbf{A} of ω^{ω} such that for any zero-dimensional Polish space X, and for any $A \subseteq X$, A is in Γ if and only if there is $f: X \to \omega^{\omega}$ continuous such that $A = f^{-1}(\mathbf{A})$. In this case, we say that \mathbf{A} is Γ -complete.

If Γ is a class of sets, then $\check{\Gamma} := \{\neg A \mid A \in \Gamma\}$ is the **dual class** of Γ , and Γ is **self-dual** if $\Gamma = \check{\Gamma}$. We set $\Delta(\Gamma) := \Gamma \cap \check{\Gamma}$.

Theorem 1.3 (Lecomte) Let Γ be a Wadge class of Borel sets, or the class Δ_{ξ}^{0} for some countable ordinal $\xi \geq 1$. Then there are concrete disjoint Borel relations \mathbb{S}_{0} , \mathbb{S}_{1} on 2^{ω} such that, for any Polish spaces X, Y, and for any disjoint analytic subsets A, B of $X \times Y$, exactly one of the following holds:

(a) the set A is separable from B by a $pot(\Gamma)$ set,

(b) $(2^{\omega}, 2^{\omega}, \mathbb{S}_0, \mathbb{S}_1) \leq (X, Y, A, B).$

It is natural to ask whether we can have f and g injective if (b) holds. Debs proved that this is the case if Γ is a non self-dual Borel class of rank at least three (i.e., a class Σ_{ξ}^{0} or Π_{ξ}^{0} with $\xi \geq 3$). As mentioned in [L8], there is also an injectivity result for the non self-dual Wadge classes of Borel sets of level at least three. Some results in [L4] and [L8] show that we cannot have f and g injective if (b) holds and Γ is a non self-dual Borel class of rank one or two, or the class of clopen sets, because of cycle problems again.

The work of Kechris, Solecki and Todorčević indicates a way to try to solve this problem. Let us recall one of their results in this direction. All the relations considered in this paper will be binary.

Definition 1.4 *Let X be a set, and A be a relation on X*.

(a) $\Delta(X) := \{(x, y) \in X^2 \mid x = y\}$ is the diagonal of X.

(b) We say that A is irreflexive if A does not meet $\Delta(X)$.

(c) $A^{-1} := \{(x, y) \in X^2 \mid (y, x) \in A\}$, and $s(A) := A \cup A^{-1}$ is the symmetrization of A.

(d) We say that A is symmetric if $A = A^{-1}$.

(e) We say that A is a graph if A is irreflexive and symmetric.

(f) We say that A is **acyclic** if there is no injective sequence $(x_i)_{i \leq n}$ of points of X with $n \geq 2$, $(x_i, x_{i+1}) \in A$ for each i < n, and $(x_n, x_0) \in A$.

(g) We say that A is **locally countable** if A has countable horizontal and vertical sections (this also makes sense in a rectangular product $X \times Y$).

Notation. Let $(s_n)_{n \in \omega}$ be a sequence of finite binary sequences with the following properties:

(a) $(s_n)_{n \in \omega}$ is **dense** in $2^{<\omega}$. This means that for each $s \in 2^{<\omega}$, there is $n \in \omega$ such that s_n extends s (denoted $s \subseteq s_n$).

(b)
$$|s_n| = n$$
.

We put $\mathbb{G}_0 := \{(s_n 0\gamma, s_n 1\gamma) \mid n \in \omega \land \gamma \in 2^\omega\}$. The following result is proved in [K-S-T].

Theorem 1.5 (*Kechris, Solecki, Todorčević*) Let X be a Polish space, and A be an analytic graph on X. We assume that A is acyclic or locally countable. Then exactly one of the following holds:

- (a) there is $c: X \to \omega$ Borel such that $A \subseteq (c \times c)^{-1} (\neg \Delta(\omega))$,
- (b) there is $f: 2^{\omega} \to X$ injective continuous such that $s(\mathbb{G}_0) \subseteq (f \times f)^{-1}(A)$.

This seems to indicate that there is a hope to get f and g injective in Theorem 1.3.(b) for the first classes of the hierarchy if we assume acyclicity or local countability. This is the main purpose of this paper, and leads to the following notation. Let X, Y, X', Y' be topological spaces and $A, B \subseteq X \times Y$, $A', B' \subseteq X' \times Y'$. We write

 $\begin{aligned} (X,Y,A,B) &\sqsubseteq (X',Y',A',B') \Leftrightarrow \\ &\exists f \colon X \to X' \ \exists g \colon Y \to Y' \text{ injective continuous with } A \subseteq (f \times g)^{-1}(A') \text{ and } B \subseteq (f \times g)^{-1}(B'). \end{aligned}$

We want to study the Borel and Wadge classes of the locally countable Borel relations: the Borel classes of rank one or two, the Lavrentieff classes built with the open sets (the classes of differences of open sets), their dual classes and their ambiguous classes. We will also study the Lavrentieff classes built with the F_{σ} sets and their dual classes.

Definition 1.6 Let $\eta < \omega_1$. If $(O_\theta)_{\theta < \eta}$ is an increasing sequence of subsets of a set X, then

$$D((O_{\theta})_{\theta < \eta}) := \{ x \in X \mid \exists \theta < \eta \ parity(\theta) \neq parity(\eta) \ and \ x \in O_{\theta} \setminus \big(\bigcup_{\theta' < \theta} O_{\theta'} \big) \}.$$

Now $D_{\eta}(\Sigma_{\xi}^{0})(X) := \{ D((O_{\theta})_{\theta < \eta}) \mid \forall \theta < \eta \ O_{\theta} \in \Sigma_{\xi}^{0}(X) \}$, for each $1 \le \xi < \omega_{1}$. The classes $D_{\eta}(\Sigma_{\xi}^{0}), \check{D}_{\eta}(\Sigma_{\xi}^{0})$ and $\Delta(D_{\eta}(\Sigma_{\xi}^{0}))$ form the difference hierarchy.

Some recent work of the first author shows that having f and g injective in Theorem 1.3.(b) can be used to get results of reduction on the whole product, under some acyclicity and also topological assumptions. Some of the results in the present paper will be used by the first author in a future article on this topic. This work is also motivated by the work of Louveau on oriented graphs in [Lo4].

Definition 1.7 Let X be a set, and A be a relation on X.

- (a) We say that A is antisymmetric if $A \cap A^{-1} \subseteq \Delta(X)$.
- (b) We say that A is an oriented graph if A is irreflexive and antisymmetric.

It follows from results of Wadge and Martin that inclusion well-orders

 $\{\Gamma \cup \check{\Gamma} \mid \Gamma \text{ Wadge class of Borel sets}\},\$

giving rise to an ordinal assignment $w(\Gamma)$. If G is an analytic oriented graph, then we can define w(G) as the least $w(\Gamma)$ such that G is separable from G^{-1} by a pot(Γ) set C. It is well defined by the separation theorem. Moreover, it is useless in the definition to distinguish between dual classes, for if C separates G from G^{-1} , then so does $\neg C^{-1}$, which is potentially in $\check{\Gamma}$. The main property of this assignment is that $w(G) \leq w(H)$ if there is a Borel homomorphism from G into H. Louveau also considers a rough approximation of w(G), which is the least countable ordinal ξ for which G is separable from G^{-1} by a pot(Δ_{ξ}^{0}) set. He proves the following.

Theorem 1.8 (Louveau) Let $\xi \in \{1, 2\}$. Then there is a concrete analytic oriented graph \mathbb{G}_{ξ} on 2^{ω} such that, for any Polish space X, and for any analytic oriented graph G on X, exactly one of the following holds:

- (a) the set G is separable from G^{-1} by a pot $(\mathbf{\Delta}^0_{\mathbf{\xi}})$ set,
- (b) there is $f: 2^{\omega} \to X$ continuous such that $\mathbb{G}_{\xi} \subseteq (f \times f)^{-1}(G)$.

Our main results are the following.

• We generalize Theorem 1.8 to all the $\Delta_{\mathcal{E}}^{0}$'s, and all the Wadge classes of Borel sets.

Theorem 1.9 Let Γ be a Wadge class of Borel sets, or the class Δ_{ξ}^{0} for some countable ordinal $\xi \ge 1$. Then there is a concrete Borel oriented graph \mathbb{G}_{Γ} on 2^{ω} such that, for any Polish space X, and for any analytic oriented graph G on X, exactly one of the following holds:

- (a) the set G is separable from G^{-1} by a pot (Γ) set,
- (b) there is $f: 2^{\omega} \to X$ continuous such that $\mathbb{G}_{\Gamma} \subseteq (f \times f)^{-1}(G)$.

We also investigate the injective version of this, for the first classes of the hierarchies again.

• In the sequel, it will be very convenient to say that a relation A on a set X is s-acyclic if s(A) is acyclic.

Theorem 1.10 Let $\Gamma \in \{D_{\eta}(\Sigma_1^0), \check{D}_{\eta}(\Sigma_1^0), D_n(\Sigma_2^0), \check{D}_n(\Sigma_2^0) \mid 1 \leq \eta < \omega_1, 1 \leq n < \omega\} \cup \{\Delta_2^0\}$. Then there are concrete disjoint Borel relations \mathbb{S}_0 , \mathbb{S}_1 on 2^{ω} such that, for any Polish space X, and for any disjoint analytic relations A, B on X with s-acyclic union, exactly one of the following holds:

- (a) the set A is separable from B by a $pot(\Gamma)$ set,
- (b) $(2^{\omega}, 2^{\omega}, \mathbb{S}_0, \mathbb{S}_1) \sqsubseteq (X, Y, A, B).$

In fact, we prove a number of extensions of this result. It also holds

- for $\eta = 0$ if we replace 2^{ω} with 1,

- with f = g if $\Gamma \notin \{D_{\eta}(\Sigma_1^0), \check{D}_{\eta}(\Sigma_1^0) \mid \eta < \omega_1\}$; if $\Gamma \in \{D_{\eta}(\Sigma_1^0), \check{D}_{\eta}(\Sigma_1^0) \mid \eta < \omega_1\}$, then there is an antichain basis with two elements for the square reduction (it is rather unusual to have an antichain basis but no minimum object in this kind of dichotomy),

- if we assume that $A \cup B$ is locally countable instead of s-acyclic when $\Gamma \subseteq \Pi_2^0$ (this also holds in rectangular products $X \times Y$),

- if we only assume that A is s-acyclic or locally countable when $\Gamma = \Pi_2^0$.

The situation is more complicated for the ambiguous classes.

Theorem 1.11 Let $\Gamma \in \{\Delta(D_{\eta}(\Sigma_1^0)) \mid 1 \leq \eta < \omega_1\}$. Then there is a concrete finite antichain \mathcal{A} , made of tuples $(2^{\omega}, 2^{\omega}, \mathbb{S}_0, \mathbb{S}_1)$ where $\mathbb{S}_0, \mathbb{S}_1$ are disjoint Borel relations $\mathbb{S}_0, \mathbb{S}_1$ on 2^{ω} , such that, for any Polish space X, and for any disjoint analytic relations A, B on X whose union is contained in a potentially closed s-acyclic relation R, exactly one of the following holds:

(a) the set A is separable from B by a $pot(\Gamma)$ set,

(b) there is $(2^{\omega}, 2^{\omega}, \mathbb{A}, \mathbb{B}) \in \mathcal{A}$ with $(2^{\omega}, 2^{\omega}, \mathbb{A}, \mathbb{B}) \sqsubseteq (X, Y, A, B)$.

Here again, we can say more. This also holds

- if we assume that R is locally countable instead of s-acyclic (this also holds in rectangular products $X \times Y$),

- in all those cases, A has size three if η is a successor ordinal, and size one if η is a limit ordinal (it is quite remarkable that the situation depends on the fact that η is limit or not, it confirms the difference observed in the description of Wadge classes of Borel sets in terms of operations on sets present in [Lo1]),

- with f = g, but in order to ensure this A must have size six if η is a successor ordinal, and size two if η is a limit ordinal.

• We characterize when part (b) in the injective reduction property holds.

Theorem 1.12 Let $\Gamma \in \{D_{\eta}(\Sigma_1^0), \check{D}_{\eta}(\Sigma_1^0), D_n(\Sigma_2^0), \check{D}_n(\Sigma_2^0) \mid 1 \leq \eta < \omega_1, 1 \leq n < \omega\} \cup \{\Delta_2^0\}$. Then there are concrete disjoint Borel relations \mathbb{S}_0 , \mathbb{S}_1 on 2^{ω} such that, for any Polish space X, and for any disjoint analytic relations A, B on X, the following are equivalent:

(1) there is an s-acyclic relation $R \in \Sigma_1^1$ such that $A \cap R$ is not separable from $B \cap R$ by a pot (Γ) set, (2) $(2^{\omega}, 2^{\omega}, \mathbb{S}_0, \mathbb{S}_1) \sqsubseteq (X, Y, A, B)$.

The same kind of extensions as before hold (except that we cannot assume local countability instead of s-acyclicity for the classes of rank two).

Theorem 1.13 Let $\Gamma \in \{\Delta(D_{\eta}(\Sigma_1^0)) \mid 1 \leq \eta < \omega_1\}$. Then there is a concrete finite antichain \mathcal{A} , made of tuples $(2^{\omega}, 2^{\omega}, \mathbb{S}_0, \mathbb{S}_1)$ where $\mathbb{S}_0, \mathbb{S}_1$ are disjoint Borel relations $\mathbb{S}_0, \mathbb{S}_1$ on 2^{ω} , such that, for any Polish space X, and for any disjoint analytic relations A, B on X, the following are equivalent: (1) there is a potentially closed s-acyclic relation $R \in \Sigma_1^1$ such that $A \cap R$ is not separable from $B \cap R$ by a $pot(\Gamma)$ set,

(2) there is $(2^{\omega}, 2^{\omega}, \mathbb{A}, \mathbb{B}) \in \mathcal{A}$ with $(2^{\omega}, 2^{\omega}, \mathbb{A}, \mathbb{B}) \subseteq (X, Y, A, B)$.

Here again, the same kind of extensions as before hold.

• The injective versions of Theorem 1.9 mentioned earlier are as follows.

Theorem 1.14 Let $\Gamma \in \{D_{\eta}(\Sigma_1^0), \check{D}_{\eta}(\Sigma_1^0), D_n(\Sigma_2^0), \check{D}_n(\Sigma_2^0) \mid 1 \leq \eta < \omega_1, 1 \leq n < \omega\} \cup \{\Delta_2^0\}$. Then there is a concrete Borel oriented graph \mathbb{G}_{Γ} on 2^{ω} such that, for any Polish space X, and for any analytic s-acyclic oriented graph G on X, exactly one of the following holds:

(a) the set G is separable from G^{-1} by a pot (Γ) set,

(b) there is $f: 2^{\omega} \to X$ injective continuous such that $\mathbb{G}_{\Gamma} \subseteq (f \times f)^{-1}(G)$.

This result also holds if we assume that G is locally countable instead of s-acyclic when $\Gamma \subseteq \Pi_2^0$.

Theorem 1.15 Let $\Gamma \in \{\Delta(D_{\eta}(\Sigma_1^0)) \mid 1 \leq \eta < \omega_1\}$. Then there is a concrete finite antichain \mathcal{A} , made of Borel oriented graphs on 2^{ω} , such that, for any Polish space X, and for any analytic oriented graph G on X contained in a potentially closed s-acyclic relation, exactly one of the following holds:

- (a) the set G is separable from G^{-1} by a pot (Γ) set,
- (b) we can find $\mathbb{G}_{\Gamma} \in \mathcal{A}$ and $f: 2^{\omega} \to X$ injective continuous such that $\mathbb{G}_{\Gamma} \subseteq (f \times f)^{-1}(G)$.

The same kind of extensions as before hold, except that A has size three if η is a successor ordinal, and size two if η is a limit ordinal.

• At the end of the paper, we study the limits of our results and give negative results.

2 Generalities

The acyclic and the locally countable cases

In [K-S-T], Section 6, the authors introduce the notion of an almost acyclic analytic graph, in order to prove an injective version of the \mathbb{G}_0 -dichotomy for acyclic or locally countable analytic graphs. We now give a similar definition, in order to prove injective versions of Theorem 1.3 for the first classes of the hierarchies. This definition is sufficient to cover all our cases, even if it is not always optimal.

Definition 2.1 Let X be a Polish space, and A be a relation on X. We say that A is **quasi-acyclic** if there is a sequence $(C_n)_{n\in\omega}$ of $pot(\mathbf{\Pi}_1^0)$ relations on X with disjoint union A such that, for any s(A)-path $(z_i)_{i\leq 2}$ with $z_0 \neq z_2$, and for any $n_1, ..., n_k \in \omega$, $C'_{n_i} \in \{C_{n_i}, C_{n_i}^{-1}\}$ $(1 \leq i \leq k), x_1, y_1, ..., x_k, y_k$ in $X \setminus \{z_i \mid i \leq 2\}$, if $(z_0, x_1), (z_2, y_1) \in C'_{n_1}, (x_1, x_2), (y_1, y_2) \in C'_{n_2}, ..., (x_{k-1}, x_k), (y_{k-1}, y_k) \in C'_{n_k}$ all hold, then $x_k \neq y_k$.

Lemma 2.2 Let X be a Polish space, and A be a Borel relation on X. We assume that A is either s-acyclic and pot(Σ_2^0), or locally countable. Then A is quasi-acyclic.

Proof. Assume first that A is s-acyclic and $pot(\Sigma_2^0)$. Then we can write $A = \bigcup_{n \in \omega} C_n$, where $(C_n)_{n \in \omega}$ is a disjoint sequence of potentially closed relations on X. The acyclicity of s(A) shows that A is quasi-acyclic.

Assume now that A is locally countable. By 18.10 in [K], A can be written as $\bigcup_{q\in\omega} G_q$, where G_q is the Borel graph of a partial function f_q , and we may assume that the G_q 's are pairwise disjoint. By 18.12 in [K], the projections of the G_q 's are Borel. By Lemma 2.4.(a) in [L2], there is, for each q, a countable partition $(D_p^q)_{p\in\omega}$ of the domain of f_q into Borel sets on which f_q is injective. So the C_n 's are the $\operatorname{Gr}(f_{q|D_n^q})$'s.

Topologies

Let Z be a recursively presented Polish space (see [M] for the basic notions of effective theory).

(1) The topology Δ_Z on Z is generated by $\Delta_1^1(Z)$. This topology is Polish (see (iii) \Rightarrow (i) in the proof of Theorem 3.4 in [Lo3]). The topology τ_1 on Z^2 is Δ_Z^2 . If $2 \le \xi < \omega_1^{CK}$, then the topology τ_{ξ} on Z^2 is generated by $\Sigma_1^1 \cap \Pi_{<\xi}^0(\tau_1)$.

(2) The **Gandy-Harrington topology** on Z is generated by $\Sigma_1^1(Z)$ and denoted Σ_Z . Recall the following facts about Σ_Z (see [L7]).

- (a) Σ_Z is finer than the initial topology of Z.
- (b) We set $\Omega_Z := \{z \in Z \mid \omega_1^z = \omega_1^{\mathbb{C}K}\}$. Then Ω_Z is $\Sigma_1^1(Z)$ and dense in (Z, Σ_Z) .
- (c) $W \cap \Omega_Z$ is a clopen subset of (Ω_Z, Σ_Z) for each $W \in \Sigma_1^1(Z)$.
- (d) (Ω_Z, Σ_Z) is a zero-dimensional Polish space.

3 The classes $D_\eta(\mathbf{\Sigma}_1^0)$ and $\check{D}_\eta(\mathbf{\Sigma}_1^0)$

Examples

In Theorem 1.3, either \mathbb{S}_0 or \mathbb{S}_1 is not locally countable if Γ is not self-dual. If $\Gamma \subseteq \Delta_2^0$, we can find disjoint analytic locally countable relations A, B on 2^{ω} such that A is not separable from B by a pot(Γ) set, as we will see. This shows that, in order to get partial reductions with injectivity, we have to use examples different from those in [L8], so that we prove the following.

Notation. We introduce examples in the style of \mathbb{G}_0 in order to prove a dichotomy for the classes $D_\eta(\Sigma_1^0)$, where $\eta \ge 1$ is a countable ordinal.

• If $t \in 2^{<\omega}$, then $N_t := \{ \alpha \in 2^{\omega} \mid t \subseteq \alpha \}$ is the usual basic clopen set.

• As in Section 2 in [L2] we inductively define $\varphi_{\eta}: \omega^{<\omega} \to \{-1\} \cup (\eta+1)$ by $\varphi_{\eta}(\emptyset) = \eta$ and

$$\varphi_{\eta}(sn) = \begin{cases} -1 \text{ if } \varphi_{\eta}(s) \leq 0, \\\\ \theta \text{ if } \varphi_{\eta}(s) = \theta + 1, \\\\ \text{ an odd ordinal such that the sequence } \left(\varphi_{\eta}(sn)\right)_{n \in \omega} \text{ is cofinal in } \varphi_{\eta}(s) \\\\ \text{ and strictly increasing if } \varphi_{\eta}(s) > 0 \text{ is limit.} \end{cases}$$

If no confusion is possible, then we will write φ instead of φ_{η} . We set $T_{\eta} := \{s \in \omega^{<\omega} \mid \varphi_{\eta}(s) \neq -1\}$, which is a wellfounded tree.

• Let $(p_q)_{q \in \omega}$ be the sequence of prime numbers, and $\langle . \rangle_\eta : T_\eta \to \omega$ be the following bijection. We define $I: T_\eta \to \omega$ by $I(\emptyset) := 0$ and $I(s) := p_0^{s(0)+1} \dots p_{|s|-1}^{s(|s|-1)+1}$ if $s \neq \emptyset$. As I is injective, there is an increasing bijection $J: I[T_\eta] \to \omega$. We set $\langle . \rangle_\eta := J \circ I$. Note that $\langle sq \rangle_\eta - \langle s \rangle_\eta \ge q+1$ if $sq \in T_\eta$. Indeed, $I(s0), \dots, I(s(q-1))$ are strictly between I(s) and I(sq).

• Let $\psi : \omega \to 2^{<\omega}$ be the map defined by $\emptyset, \emptyset, 0, 0, 1, 1, 0^2, 0^2, 01, 01, 10, 10, 1^2, 1^2, ...,$ so that $|\psi(q)| \le q$ and $\psi[\{2n \mid n \in \omega\}], \psi[\{2n+1 \mid n \in \omega\}] = 2^{<\omega}$.

• For each $s \in T_{\eta}$, we define $(t_s^0, t_s^1) \in (2 \times 2)^{<\omega}$ by $t_{\emptyset}^{\varepsilon} = \emptyset$, and $t_{sq}^{\varepsilon} = t_s^{\varepsilon} \psi(q) 0^{< sq >_{\eta} - < s >_{\eta} - |\psi(q)| - 1_{\varepsilon}}$. Note that this is well defined, $|t_s^{\varepsilon}| = < s >_{\eta}$ and $\operatorname{Card}(\{l < < s >_{\eta} | t_s^0(l) \neq t_s^1(l)\}) = |s|$ for each $s \in T_{\eta}$.

• We set $\mathcal{T}^{\eta} := \{ (t_s^0 w, t_s^1 w) \mid s \in T_{\eta} \land w \in 2^{<\omega} \}$. The following properties are satisfied.

- \mathcal{T}^{η} is a tree on 2×2, and $\lceil \mathcal{T}^{\eta} \rceil \subseteq \mathbb{E}_0 := \{(\alpha, \beta) \in 2^{\omega} \times 2^{\omega} \mid \exists m \in \omega \ \forall n > m \ \alpha(n) = \beta(n)\}$ is locally countable.

- If
$$(s,t) \in \mathcal{T}^{\eta}$$
 and $s(l) \neq t(l)$, then $s(l) < t(l)$.

- For each $l \in \omega$, there is exactly one sequence $(u, v) \in \mathcal{T}^{\eta} \cap (2^{l+1} \times 2^{l+1})$ such that $u(l) \neq v(l)$ since $t_{sq}^0 (< sq >_{\eta} -1) \neq t_{sq}^1 (< sq >_{\eta} -1)$ (in fact, (u, v) is of the form (t_s^0, t_s^1) for some s). In particular, $s(\mathcal{T}^{\eta} \cap (2^{l+1} \times 2^{l+1})) \setminus \Delta(2^{l+1})$ is a connected acyclic graph on 2^{l+1} , inductively.

• We set, for $\varepsilon \in 2$,

$$\mathbb{N}_{\varepsilon}^{\eta} := \Big\{ (t_s^0 \gamma, t_s^1 \gamma) \mid s \in T_{\eta} \land \operatorname{parity}(|s|) = \varepsilon \land \gamma \in 2^{\omega} \Big\}.$$

If $s \in T_{\eta}$, then $f_s : N_{t_s^0} \to N_{t_s^1}$ is the partial homeomophism with clopen domain and range defined by $f_s(t_s^0\gamma) := t_s^1\gamma$, so that $\mathbb{N}_{\varepsilon}^{\eta} = \bigcup_{s \in T_{\eta}, \text{parity}(|s|) = \varepsilon} \operatorname{Gr}(f_s)$. We set $C_s := \bigcup_{q \in \omega} \operatorname{Gr}(f_{sq})$ when it makes sense (i.e., when $\varphi_{\eta}(s) \ge 1$). For $\eta = 0$, we set $\mathbb{N}_0^{\eta} := 1^2$ and $\mathbb{N}_1^{\eta} := \emptyset$ (in 1²).

Lemma 3.1 Let η be a countable ordinal, and C be a nonempty clopen subset of 2^{ω} .

(a) If $\varphi_{\eta}(s) \ge 1$ and G is a dense G_{δ} subset of 2^{ω} , then $\overline{C_s} \cap (C \cap G)^2 \subseteq \overline{C_s \cap (C \cap G)^2}$. (b) $\mathbb{N}_0^{\eta} \cap C^2$ is not separable from $\mathbb{N}_1^{\eta} \cap C^2$ by a pot $(D_{\eta}(\Sigma_1^0))$ set. **Proof.** (a) It is enough to prove that if $q \in \omega$, then $\operatorname{Gr}(f_{sq}) \cap C^2 \subseteq \overline{\operatorname{Gr}(f_{sq}) \cap (C \cap G)^2}$. This comes from the proof of Lemma 3.5 in [L1], but we recall it for self-containedness. Let U, V be open subsets of C such that $\operatorname{Gr}(f_{sq}) \cap (U \times V) \neq \emptyset$. Then $N_{t_{sq}^1} \cap V \cap G$ is a dense G_{δ} subset of $N_{t_{sq}^1} \cap V$, so that $f_{sq}^{-1}(V \cap G)$ is a dense G_{δ} subset of $f_{sq}^{-1}(V)$. Thus $G \cap f_{sq}^{-1}(V)$ and $G \cap f_{sq}^{-1}(V \cap G)$ are dense G_{δ} subsets of $f_{sq}^{-1}(V)$. This gives α in this last set and $U \cap f_{sq}^{-1}(V)$. Therefore $(\alpha, f_{sq}(\alpha))$ is in $\operatorname{Gr}(f_{sq}) \cap (C \cap G)^2 \cap (U \times V)$.

(b) We may assume that $\eta \ge 1$. We argue by contradiction, which gives $P \in \text{pot}(D_{\eta}(\Sigma_{1}^{0}))$, and a dense G_{δ} subset of 2^{ω} such that $P \cap G^{2} \in D_{\eta}(\Sigma_{1}^{0})(G^{2})$. So let $(O_{\theta})_{\theta < \eta}$ be a sequence of open relations on 2^{ω} such that $P \cap G^{2} = (\bigcup_{\theta < \eta, \text{parity}(\theta) \neq \text{parity}(\eta)} O_{\theta} \setminus (\bigcup_{\theta' < \theta} O_{\theta'})) \cap G^{2}$.

• Let us show that if $\theta \leq \eta$, $s \in T_{\eta}$ and $\varphi(s) = \theta$, then $\operatorname{Gr}(f_s) \cap (C \cap G)^2 \subseteq \neg O_{\theta}$ if $\theta < \eta$, and $\operatorname{Gr}(f_s) \cap (C \cap G)^2$ is disjoint from $\bigcup_{\theta' < \theta} O_{\theta'}$ if $\theta = \eta$. The objects $s = \emptyset$ and $\theta = \eta$ will give the contradiction.

• We argue by induction on θ . Note that if $s \in T_{\eta}$, |s| is even if and only if $\varphi(s)$ has the same parity as η . If $\theta = 0$, then |s| has the same parity as η , thus $\operatorname{Gr}(f_s) \cap (C \cap G)^2 \subseteq \mathbb{N}^{\eta}_{\operatorname{parity}(\eta)} \cap G^2 \subseteq \neg O_0$.

• Assume that the result has been proved for $\theta' < \theta$. If θ is the successor of θ' , then the induction assumption implies that $\operatorname{Gr}(f_{sq}) \cap (C \cap G)^2 \subseteq \neg O_{\theta'}$ for each q. So $C_s \cap (C \cap G)^2 \subseteq \neg O_{\theta'}$ and $\overline{C_s \cap (C \cap G)^2} \subseteq \neg O_{\theta'}$. By (a), we get $\overline{C_s} \cap (C \cap G)^2 \subseteq \overline{C_s \cap (C \cap G)^2}$, which gives the desired inclusion if $\theta = \eta$ since $\operatorname{Gr}(f_s) \subseteq \overline{C_s}$.

If $\theta < \eta$ and |s| is even, then $\varphi(s)$ has the same parity as η and the parity of θ' is opposite to that of η . Note that $\operatorname{Gr}(f_s) \cap (C \cap G)^2 \subseteq \mathbb{N}_0^{\eta} \cap G^2 \subseteq \bigcup_{\theta'' < \eta, \operatorname{parity}(\theta'') \neq \operatorname{parity}(\eta)} O_{\theta''} \setminus (\bigcup_{\theta''' < \theta''} O_{\theta'''}) \subseteq \neg O_{\theta}$.

If |s| is odd, then the parity of $\varphi(s)$ is opposite to that of η and θ' has the same parity as η . But if $s \in T_{\eta}$ has odd length, then

$$\operatorname{Gr}(f_s) \cap (C \cap G)^2 \subseteq \mathbb{N}_1^{\eta} \cap G^2 \subseteq G^2 \setminus (\bigcup_{\theta'' < \eta} O_{\theta''}) \cup \bigcup_{\theta'' < \eta, \operatorname{parity}(\theta'') = \operatorname{parity}(\eta)} O_{\theta''} \setminus (\bigcup_{\theta''' < \theta''} O_{\theta'''}).$$

This gives the result.

• If θ is limit, then $(\varphi(sn))_{n\in\omega}$ is cofinal in $\varphi(s)$, and $\operatorname{Gr}(f_{sn}) \cap (C \cap G)^2 \subseteq \neg O_{\varphi(sn)}$ by the induction assumption. If $\theta_0 < \varphi(s)$, then there is $n(\theta_0)$ such that $\varphi(sn) > \theta_0$ if $n \ge n(\theta_0)$. Thus $\operatorname{Gr}(f_{sn}) \cap (C \cap G)^2 \subseteq \neg O_{\theta_0}$ as soon as $n \ge n(\theta_0)$. But

$$\operatorname{Gr}(f_s) \cap (C \cap G)^2 \subseteq (C \cap G)^2 \cap \overline{C_s} \setminus C_s = \overline{C_s \cap (C \cap G)^2} \setminus C_s \subseteq \overline{\bigcup_{n \ge n(\theta_0)} \operatorname{Gr}(f_{sn}) \cap (C \cap G)^2} \subseteq \neg O_{\theta_0}.$$

Thus $\operatorname{Gr}(f_s) \cap (C \cap G)^2 \subseteq \neg(\bigcup_{\theta' < \theta} O_{\theta'}).$

If $\theta < \eta$, as |s| has the same parity as η , we get $\operatorname{Gr}(f_s) \cap (C \cap G)^2 \subseteq \mathbb{N}_{\operatorname{parity}(\eta)}^{\eta} \cap G^2$, so that $\operatorname{Gr}(f_s) \cap (C \cap G)^2 \subseteq \neg O_{\theta}$.

A topological characterization

Notation. Let $1 \leq \xi < \omega_1^{\mathbb{C}K}$. Theorem 4.1 in [L6] shows that if A_0, A_1 are disjoint Σ_1^1 relations on ω^{ω} , then A_0 is separable from A_1 by a pot (Σ_{ξ}^0) set exactly when $A_0 \cap \overline{A_1}^{\tau_{\xi}} = \emptyset$. We now define the versions of $A_0 \cap \overline{A_1}^{\tau_{\xi}}$ for the classes $D_{\eta}(\Sigma_{\xi}^0)$. So let $\varepsilon \in 2$ and $\eta < \omega_1^{\mathbb{C}K}$. We define $\bigcap_{\theta < 0} F_{\theta, \varepsilon}^{\varepsilon} := (\omega^{\omega})^2$, and, inductively,

$$F_{\eta,\xi}^{\varepsilon} := \overline{A_{|\operatorname{parity}(\eta) - \varepsilon|} \cap \bigcap_{\theta < \eta} F_{\theta,\xi}^{\varepsilon}}^{\tau_{\xi}}$$

We will sometimes denote by $F_{\eta,\xi}^{\varepsilon}(A_0, A_1)$ the sets $F_{\eta,\xi}^{\varepsilon}$ previously defined. By induction, we can check that $F_{\eta,\xi}^{\varepsilon}(A_1, A_0) = F_{\eta,\xi}^{1-\varepsilon}(A_0, A_1)$.

Fix a bijection $l \mapsto ((l)_0, (l)_1)$ from ω onto ω^2 , with inverse map $(m, p) \mapsto < m, p >$. We define, for $u \in \omega^{\leq \omega}$ and $n \in \omega$, $(u)_n \in \omega^{\leq \omega}$ by $(u)_n(p) := u(< n, p >)$ if < n, p > < |u|.

Theorem 3.2 Let $1 \le \xi < \omega_1^{CK}$, $\eta = \lambda + 2k + \varepsilon < \omega_1^{CK}$ with λ limit, $k \in \omega$ and $\varepsilon \in 2$, and A_0 , A_1 be disjoint Σ_1^1 relations on ω^{ω} . Then the following are equivalent:

(1) the set A_0 is not separable from A_1 by a pot $(D_{\eta}(\Sigma_{\varepsilon}^0))$ set,

(2) the Σ_1^1 set $F_{n,\xi}^{\varepsilon}$ is not empty.

Proof. This result is essentially proved in [L8]. However, the formula for $F_{\eta,\xi}^{\varepsilon}$ is more concrete here, since the more general and abstract case of Wadge classes is considered in [L8]. So we give some details.

• In [Lo-SR], the following class of sets is introduced. Let $1 \le \xi < \omega_1$ and Γ , Γ' be two classes of sets. Then $A \in S_{\xi}(\Gamma, \Gamma') \iff A = \bigcup_{p \ge 1} (A_p \cap C_p) \cup (B \setminus \bigcup_{p \ge 1} C_p)$, where $A_p \in \Gamma$, $B \in \Gamma'$, and $(C_p)_{p \ge 1}$ is a sequence of pairwise disjoint Σ_{ξ}^0 sets. The authors prove the following:

$$\Sigma_{\xi}^{0} = S_{\xi}(\{\emptyset\}, \{\emptyset\}),$$

$$D_{\theta+1}(\Sigma_{\xi}^{0}) = S_{\xi}(\check{D}_{\theta}(\Sigma_{\xi}^{0}), \Sigma_{\xi}^{0}) \text{ if } \theta < \omega_{1},$$

$$D_{\lambda}(\Sigma_{\xi}^{0}) = S_{\xi}(\bigcup_{p \geq 1} D_{\theta_{p}}(\Sigma_{\xi}^{0}), \{\emptyset\}) \text{ if } \lambda = \sup_{p \geq 1} \theta_{p} \text{ is limit.}$$

They also code the non self-dual Wadge classes of Borel sets by elements of ω_1^{ω} as follows (we sometimes identify ω_1^{ω} with $(\omega_1^{\omega})^{\omega}$). The relations "*u* is a second type description" and "*u* describes Γ " (written $u \in \mathcal{D}$ and $\Gamma_u = \Gamma$ - ambiguously) are the least relations satisfying the following properties.

(a) If $u = 0^{\infty}$, then $u \in \mathcal{D}$ and $\Gamma_u = \{\emptyset\}$.

(b) If $u = \xi^{1}v$, with $v \in \mathcal{D}$ and $v(0) = \xi$, then $u \in \mathcal{D}$ and $\Gamma_u = \check{\Gamma}_v$.

(c) If $u = \xi^{-2} < u_p$ > satisfies $\xi \ge 1$, $u_p \in \mathcal{D}$, and $u_p(0) \ge \xi$ or $u_p(0) = 0$, then $u \in \mathcal{D}$ and $\Gamma_u = S_{\xi}(\bigcup_{p>1} \Gamma_{u_p}, \Gamma_{u_0})$.

They prove that Γ is a non self-dual Wadge class of Borel sets exactly when there is $u \in \mathcal{D}$ such that $\Gamma(\omega^{\omega}) = \Gamma_u(\omega^{\omega})$.

• In [L8], the elements of \mathcal{D} are coded by elements of ω^{ω} . An inductive operator \mathfrak{H} over ω^{ω} is defined and there is a partial function $c:\omega^{\omega} \to \omega_1^{\omega}$ with $c[\mathfrak{H}^{\infty}] = \mathcal{D}$ (see Lemma 6.2 in [L8]). Another operator \mathfrak{J} on $(\omega^{\omega})^3$ is defined in [L8] to code the non self-dual Wadge classes of Borel sets and their elements (see Lemma 6.5 in [L8]). We will need a last inductive operator \mathfrak{K} , on $(\omega^{\omega})^6$, to code the sets that will play the role of the Σ_1^1 sets $F_{\eta,\xi}^{\varepsilon}$'s, via a universal set \mathcal{U} for the class $\Pi_1^1(\omega^{\omega} \times \omega^{\omega})$. More precisely, if $(\alpha, a_0, a_1, b_0, b_1, r) \in \mathfrak{K}^{\infty}$, then b_0, b_1 and r are completely determined by (α, a_0, a_1) and in practice α will be in \mathfrak{H}^{∞} , so that we will write $r = r(\alpha, a_0, a_1) = r(u, a_0, a_1)$ if $u = c(\alpha)$. Our Σ_1^1 sets A_0, A_1 are coded by a_0, a_1 , in the sense that $A_{\varepsilon} = \neg \mathcal{U}_{a_{\varepsilon}}$. By Lemma 6.6 in [L8], there is a recursive map $\mathcal{A}: (\omega^{\omega})^2 \to \omega^{\omega}$ such that $\neg \mathcal{U}_{\mathcal{A}(\alpha,r)} = (\neg \mathcal{U}_{(r)_0}) \cap \bigcap_{p\geq 1} \overline{\neg \mathcal{U}_{(r)_p}}^{\tau_{|\alpha|}}$ if $\alpha \in \Delta_1^1$ codes a wellordering, where $r \mapsto ((r)_p)_{p\in\omega}$ is a bijection from ω^{ω} onto $(\omega^{\omega})^{\omega}$. In the sequel, all the closures will be for τ_{ξ} .

• We argue by induction on η . As $D_0(\Sigma_{\xi}^0) = \{\emptyset\}$, A_0 is separable from A_1 by a $D_0(\Sigma_{\xi}^0)$ set when $A_0 = \emptyset$, which is equivalent to $F_{0,\xi}^0 = \overline{A_0} = \emptyset$. As $D_1(\Sigma_{\xi}^0) = \Sigma_{\xi}^0$, A_0 is separable from A_1 by a $D_1(\Sigma_{\xi}^0)$ set when $A_0 \cap \overline{A_1} = \emptyset$ by Theorem 4.1 in [L6], which is equivalent to $F_{1,\xi}^1 = \overline{A_0 \cap \overline{A_1}} = \emptyset$.

Let us do these two basic cases in the spirit of the material from [L8] previously described, which will be done also for the other more complex cases.

- Note that $D_0(\Sigma_{\xi}^0) = \{\emptyset\} = \Gamma_{0^{\infty}}$. Let $\alpha \in \Delta_1^1$ such that $(\alpha)_n$ codes a wellordering of order type 0 for each $n \in \omega$. A look at the definition of \mathfrak{H} shows that $\alpha \in \mathfrak{H}^{\infty}$. Another look at Definition 6.3 in [L8] shows that α is normalized (this will never be a problem in the sequel as well). Lemma 6.5 in [L8] gives $\beta, \gamma \in \omega^{\omega}$ with $(\alpha, \beta, \gamma) \in \mathfrak{J}^{\infty}$. Lemma 6.7 in [L8] gives $b_0, b_1, r \in \omega^{\omega}$ with $(\alpha, \alpha_1, a_0, b_0, b_1, r) \in \mathfrak{K}^{\infty}$. By Theorem 6.10 in [L8], A_1 is separable from A_0 by a pot $(\check{D}_0(\Sigma_{\xi}^0))$ set if and only if $\neg \mathcal{U}_r = \emptyset$. A look at the definition of \mathfrak{K} shows that $r = a_0$, so that $\neg \mathcal{U}_r = A_0$.

- Now $D_1(\Sigma_{\xi}^0) = \Sigma_{\xi}^0 = S_{\xi}(\check{\{\emptyset\}}, \{\emptyset\}) = S_{\xi}(\Gamma_{010^{\infty}}, \Gamma_{0^{\infty}}) = S_{\xi}(\bigcup_{p \ge 1} \Gamma_{010^{\infty}}, \Gamma_{0^{\infty}}) = \Gamma_{v_1}$, where $v_1 := \xi 2 < 0^{\infty}, 010^{\infty}, 010^{\infty}, \dots >$. As above, A_1 is separable from A_0 by a pot $(\check{D}_1(\Sigma_{\xi}^0))$ set if and only if $\neg \mathcal{U}_r = \emptyset$. A look at the definition of \mathfrak{K} shows that $r = b_0 = \mathcal{A}(\alpha_1, < a_0, a_1, a_1, \dots >)$, where $|\alpha_1| = \xi$. Thus $\neg \mathcal{U}_r = A_0 \cap \overline{A_1}$.

In the general case, there is $v_{\eta} \in \mathcal{D}$ such that $D_{\eta}(\Sigma_{\xi}^{0}) = \Gamma_{v_{\eta}}$ and A_{1} is separable from A_{0} by a $\text{pot}(\check{D}_{\eta}(\Sigma_{\xi}^{0}))$ set if and only if $\neg \mathcal{U}_{r(v_{\eta},a_{1},a_{0})} = \emptyset$. Moreover,

- (a) if $v_{\eta} = 0^{\infty}$, then $r(v_{\eta}, a_1, a_0) = a_0$,
- (b) if $v_{\eta} = \xi^{-1} v$, then $r(v_{\eta}, a_1, a_0) = a_1$,

(c) if $v_{\eta} = \xi^{-2} < u_p >$ and $r_p = r(u_p, a_1, a_0)$, then $r(v_{\eta}, a_1, a_0) = r(u_0, b_1, b_0)$, where by definition $b_i := \mathcal{A}(\alpha_1, < a_i, r_1, r_2, ... >)$.

It is enough to prove that $F_{\eta,\xi}^{\varepsilon} = \overline{\neg \mathcal{U}_{r(v_{\eta},a_1,a_0)}}$, and we may assume that $\eta \geq 2$ by the previous discussion.

• If η is a limit ordinal, then fix a sequence $(\eta_p)_{p\in\omega}$ of even ordinals cofinal in η . Note that

$$D_{\eta}(\boldsymbol{\Sigma}_{\xi}^{0}) = S_{\xi}(\bigcup_{p \ge 1} D_{\eta_{p}}(\boldsymbol{\Sigma}_{\xi}^{0}), \{\emptyset\}) = S_{\xi}(\bigcup_{p \ge 1} \boldsymbol{\Gamma}_{u_{p}}, \boldsymbol{\Gamma}_{u_{0}}) = \boldsymbol{\Gamma}_{v_{\eta}},$$

where $v_{\eta} = \xi^{\gamma} 2^{\gamma} < u_p >$.

Therefore, if $r_p := r(u_p, a_1, a_0)$, then $F_{\theta_p,\xi}^{\varepsilon} = \overline{-\mathcal{U}_{r_p}}$ if $p \ge 1$, by the induction hypothesis. On the other hand, $r(u_0, b_1, b_0) = b_0$. But $b_0 = \mathcal{A}(\alpha_1, < a_0, r_1, r_2, \dots >)$, so that

$$\neg \mathcal{U}_{b_0} = (\neg \mathcal{U}_{a_0}) \cap \bigcap_{p \ge 1} \overline{\neg \mathcal{U}_{r_p}}$$

as required.

• If $\eta = \theta + 1$, then

$$D_{\eta}(\boldsymbol{\Sigma}_{\xi}^{0}) = S_{\xi}(\check{D}_{\theta}(\boldsymbol{\Sigma}_{\xi}^{0}), \boldsymbol{\Sigma}_{\xi}^{0}) = S_{\xi}(\bigcup_{p \ge 1} \boldsymbol{\Gamma}_{u_{p}}, \boldsymbol{\Gamma}_{u_{0}}) = \boldsymbol{\Gamma}_{v_{\eta}},$$

where $v_{\eta} = \xi \widehat{} 2 \widehat{} < u_p >$. Therefore, if $r_p := r(u_p, a_1, a_0)$, then $F_{\theta,\xi}^{\varepsilon} = \overline{\neg \mathcal{U}_{r_p}}$ if $p \ge 1$, by the induction hypothesis (there is a double inversion of the superscript, one because the parity of θ is different from that of η , and the other one because there is a complement, so that the roles of A_0, A_1 are exchanged). By the case $\eta = 1$ applied to b_0 and $b_1, \neg \mathcal{U}_{r(u_0,b_1,b_0)} = \neg \mathcal{U}_{b_0} \cap \overline{\neg \mathcal{U}_{b_1}}$. Note that

$$\neg \mathcal{U}_{b_i} = (\neg \mathcal{U}_{a_i}) \cap \bigcap_{p \ge 1} \overline{\neg \mathcal{U}_{r_p}} = (\neg \mathcal{U}_{a_i}) \cap F_{\theta,\xi}^{\varepsilon}$$

since $b_i = \mathcal{A}(\alpha_1, < a_i, r_1, r_2, ... >)$. If $r := r(v_\eta, a_1, a_0)$, then

$$\neg U_r = (\neg \mathcal{U}_{a_0}) \cap F_{\theta,\xi}^{\varepsilon} \cap \overline{\neg \mathcal{U}_{a_1} \cap F_{\theta,\xi}^{\varepsilon}} = A_0 \cap F_{\theta,\xi}^{\varepsilon},$$

because $F_{\theta,\xi}^{\varepsilon} = \overline{A_1 \cap \bigcap_{\rho < \theta} F_{\rho,\xi}^{\varepsilon}} \subseteq \overline{A_1 \cap \overline{A_1 \cap \bigcap_{\rho < \theta} F_{\rho,\xi}^{\varepsilon}}} \subseteq \overline{A_1 \cap F_{\theta,\xi}^{\varepsilon}}$ (since the parity of θ is different from ε). Finally, $\overline{\neg U_r} = \overline{A_0 \cap F_{\theta,\xi}^{\varepsilon}} = F_{\eta,\xi}^{\varepsilon}$, as required.

The main result

We set, for $\eta < \omega_1$ and $\varepsilon \in 2$, $\mathbb{B}_{\varepsilon}^{\eta} := \{(0\alpha, 1\beta) \mid (\alpha, \beta) \in \mathbb{N}_{\varepsilon}^{\eta}\}.$

Theorem 3.3 Let $\eta \ge 1$ be a countable ordinal, X be a Polish space, and A_0, A_1 be disjoint analytic relations on X such that $A_0 \cup A_1$ is quasi-acyclic. The following are equivalent: (1) the set A_0 is not separable from A_1 by a pot $(D_n(\Sigma_1^0))$ set,

(2) there is $(\mathbb{A}_0, \mathbb{A}_1) \in \{(\mathbb{N}_0^{\eta}, \mathbb{N}_1^{\eta}), (\mathbb{B}_0^{\eta}, \mathbb{B}_1^{\eta})\}$ such that $(2^{\omega}, 2^{\omega}, \mathbb{A}_0, \mathbb{A}_1) \sqsubseteq (X, X, A_0, A_1)$, via a square map,

(3) $(2^{\omega}, 2^{\omega}, \mathbb{N}_0^{\eta}, \mathbb{N}_1^{\eta}) \sqsubseteq (X, X, A_0, A_1).$

Proof. (1) \Rightarrow (2) Let $\varepsilon := \text{parity}(\eta)$, and $(C_p)_{p \in \omega}$ be a witness for the quasi-acyclicity of $A_0 \cup A_1$. We may assume that $X = \omega^{\omega}$. Indeed, we may assume that X is zero-dimensional, and thus a closed subset of ω^{ω} . As A_0 is not separable from A_1 by a $\text{pot}(D_\eta(\Sigma_1^0))$ set in X^2 , it is also the case in $(\omega^{\omega})^2$, which gives $f : 2^{\omega} \to \omega^{\omega}$. As $\Delta(2^{\omega}) \subseteq \mathbb{N}_0^{\eta}$ and $\{(0\alpha, 1\alpha) \mid \alpha \in 2^{\omega}\} \subseteq \mathbb{B}_0^{\eta}$, the range of $\Delta(2^{\omega})$ by $f \times f$ is a subset of X^2 , so that f takes values in X. We may also assume that A_0, A_1 are Σ_1^1 , and that the relation " $(x, y) \in C_p$ " is Δ_1^1 in (x, y, p). By Theorem 3.2,

$$F_{\eta}^{\varepsilon} = \overline{A_0 \cap \bigcap_{\theta < \eta} F_{\theta}^{\varepsilon}}^{\tau_1}$$

is a nonempty Σ_1^1 relation on X (where $F_{\eta}^{\varepsilon} := F_{\eta,1}^{\varepsilon}$, for simplicity).

We set, for $\theta \leq \eta$, $F_{\theta} := A_{|\text{parity}(\theta) - \varepsilon|} \cap \bigcap_{\theta' < \theta} F_{\theta'}^{\varepsilon}$, so that $F_{\theta}^{\varepsilon} = \overline{F_{\theta}}^{\tau_1}$. We put, for $\theta \leq \eta$,

$$D_{\theta} := \{ (t_s^0 w, t_s^1 w) \in \mathcal{T}^{\eta} \mid \varphi(s) = \theta \},\$$

so that $(D_{\theta})_{\theta \leq \eta}$ is a partition of \mathcal{T}^{η} . As $D_{\eta} = \Delta(2^{<\omega})$, $G_{l+1} := s((\bigcup_{\theta < \eta} D_{\theta}) \cap (2^{l+1} \times 2^{l+1}))$ is a connected acyclic graph on 2^{l+1} for each $l \in \omega$.

Case 1 $F_\eta \not\subseteq \Delta(X)$.

Let $(x, y) \in F_{\eta} \setminus \Delta(X)$, and O_0, O_1 be disjoint Δ_1^0 sets with $(x, y) \in O_0 \times O_1$. We can replace F_{η}, A_0 and A_1 with their intersection with $O_0 \times O_1$ if necessary and assume that they are contained in $O_0 \times O_1$.

• We construct the following objects:

- sequences $(x_s)_{s\in 2^{<\omega}}$, $(y_s)_{s\in 2^{<\omega}}$ of points of X,
- sequences $(X_s)_{s\in 2^{<\omega}}$, $(Y_s)_{s\in 2^{<\omega}}$ of Σ_1^1 subsets of X,
- a sequence $(U_{s,t})_{(s,t)\in\mathcal{T}^{\eta}}$ of Σ_1^1 subsets of X^2 , and $\Phi:\mathcal{T}^{\eta}\to\omega$.

We want these objects to satisfy the following conditions:

$$\begin{array}{l} (1) \ x_s \in X_s \ \land \ y_s \in Y_s \ \land \ (x_s, y_t) \in U_{s,t} \\ (2) \ X_{s\varepsilon} \subseteq X_s \subseteq \Omega_X \cap O_0 \ \land \ Y_{s\varepsilon} \subseteq Y_s \subseteq \Omega_X \cap O_1 \ \land \ U_{s,t} \subseteq C_{\Phi(s,t)} \cap \Omega_{X^2} \cap (X_s \times Y_t) \\ (3) \ \dim_{\operatorname{GH}}(X_s), \ \dim_{\operatorname{GH}}(Y_s), \ \dim_{\operatorname{GH}}(U_{s,t}) \leq 2^{-|s|} \\ (4) \ X_{s0} \cap X_{s1} = Y_{s0} \cap Y_{s1} = \emptyset \\ (5) \ U_{s\varepsilon,t\varepsilon} \subseteq U_{s,t} \\ (6) \ U_{s,t} \subseteq F_{\theta} \ \text{if} \ (s,t) \in D_{\theta} \end{array}$$

• Assume that this has been done. Let $\alpha \in 2^{\omega}$. The sequence $(X_{\alpha|n})_{n \in \omega}$ is a decreasing sequence of nonempty clopen subsets of Ω_X with vanishing diameters, which defines $f_0(\alpha) \in \bigcap_{n \in \omega} X_{\alpha|n}$. As the Gandy-Harrington topology is finer than the original topology, $f_0: 2^{\omega} \to O_0$ is continuous. By (4), f_0 is injective. Similarly, we define $f_1: 2^{\omega} \to O_1$ injective continuous. Finally, we define $f: 2^{\omega} \to X$ by $f(\varepsilon \alpha):=f_{\varepsilon}(\alpha)$, so that f is also injective continuous since O_0, O_1 are disjoint.

If $(0\alpha, 1\beta) \in \mathbb{B}_0^\eta$, then there is $\theta \leq \eta$ of the same parity as η such that $(\alpha, \beta) | n \in D_\theta$ if $n \geq n_0$. In this case, by (1)-(3) and (5)-(6), $(U_{(\alpha,\beta)|n})_{n\geq n_0}$ is a decreasing sequence of nonempty clopen subsets of $A_0 \cap \Omega_{X^2}$ with vanishing diameters, so that its intersection is a singleton $\{F(\alpha, \beta)\} \subseteq A_0$. As $(x_{\alpha|n}, y_{\beta|n})$ converges (for Σ_{X^2} , and thus for Σ_X^2) to $F(\alpha, \beta)$, $(f(0\alpha), f(1\beta)) = F(\alpha, \beta) \in A_0$. If $(0\alpha, 1\beta) \in \mathbb{B}_1^\eta$, then the parity of θ is opposite to that of η and, similarly, $(f(0\alpha), f(1\beta)) \in A_1$.

• So let us prove that the construction is possible. Note that $(t_{\emptyset}^0, t_{\emptyset}^1) = (\emptyset, \emptyset), \mathcal{T}^{\eta} \cap (2^0 \times 2^0) = \{(\emptyset, \emptyset)\}$ and $(\emptyset, \emptyset) \in D_{\eta}$. Let $(x_{\emptyset}, y_{\emptyset}) \in F_{\eta} \cap \Omega_{X^2}$, and $\Phi(\emptyset, \emptyset) \in \omega$ such that $(x_{\emptyset}, y_{\emptyset}) \in C_{\Phi(\emptyset,\emptyset)}$. As $\Omega_{X^2} \subseteq \Omega_X^2$, $x_{\emptyset}, y_{\emptyset} \in \Omega_X$. We choose Σ_1^1 subsets $X_{\emptyset}, Y_{\emptyset}$ of X with GH-diameter at most 1 such that

$$(x_{\emptyset}, y_{\emptyset}) \in X_{\emptyset} \times Y_{\emptyset} \subseteq (\Omega_X \cap O_0) \times (\Omega_X \cap O_1),$$

as well as a Σ_1^1 subset $U_{\emptyset,\emptyset}$ of X^2 with GH-diameter at most 1 such that

$$(x_{\emptyset}, y_{\emptyset}) \in U_{\emptyset,\emptyset} \subseteq F_{\eta} \cap C_{\Phi(\emptyset,\emptyset)} \cap \Omega_{X^2} \cap (X_{\emptyset} \times Y_{\emptyset}),$$

which completes the construction for the length l=0.

Assume that we have constructed our objects for the sequences of length l. Let $u \in \omega^{<\omega}$ and $q \in \omega$ with $l+1 = \langle uq \rangle_{\eta}$, which gives $w \in \omega^{<\omega}$ with $(t_{uq}^0, t_{uq}^1) = (t_u^0 w 0, t_u^1 w 1)$. We set

$$\begin{split} U &:= \{ x \in X \mid \exists (x'_s)_{s \in 2^l} \in \Pi_{s \in 2^l} X_s \ \exists (y'_s)_{s \in 2^l} \in \Pi_{s \in 2^l} Y_s \ x = x'_{t_u^0 w} \land \\ \forall (s,t) \in \mathcal{T}^\eta \cap (2^l \times 2^l) \ (x'_s, y'_t) \in U_{s,t} \}, \\ V &:= \{ y \in X \mid \exists (x'_s)_{s \in 2^l} \in \Pi_{s \in 2^l} X_s \ \exists (y'_s)_{s \in 2^l} \in \Pi_{s \in 2^l} Y_s \ y = y'_{t_u^1 w} \land \\ \forall (s,t) \in \mathcal{T}^\eta \cap (2^l \times 2^l) \ (x'_s, y'_t) \in U_{s,t} \}. \end{split}$$


Note that U, V are Σ_1^1 and $(x_{t_u^0 w}, y_{t_u^1 w}) \in F_{\varphi(u)} \cap (U \times V) \subseteq \bigcap_{\theta < \varphi(u)} \overline{F_{\theta}}^{\tau_1} \cap (U \times V)$. This gives $(x_{t_u^0 w0}, y_{t_u^1 w1}) \in F_{\varphi(uq)} \cap (U \times V) \cap \Omega_{X^2}$. Let $(x_{s0})_{s \in 2^l \setminus \{t_u^0 w\}}$ be witnesses for the fact that $x_{t_u^0 w0} \in U$, and $(x_{s1})_{s \in 2^l \setminus \{t_u^1 w\}}$ be witnesses for the fact that $x_{t_u^1 w1} \in V$.

We need to show that $x_{s0} \neq x_{s1}$ (and similarly for y_{s0} and y_{s1}). First observe that if $s \neq t \in 2^l$, then $x_{s\varepsilon} \in X_s$ and $x_{t\varepsilon'} \in X_t$, so that $x_{s\varepsilon} \neq x_{t\varepsilon'}$ by condition 4. Similarly, $y_{s\varepsilon} \neq y_{t\varepsilon'}$. As $\varphi(u)$ and $\varphi(uq)$ do not have the same parity, there is $\epsilon \in 2$ such that $(x_{t_u^0 w0}, y_{t_u^1 w1}) \in A_{\epsilon}$ and

$$(x_{t_u^0w1}, y_{t_u^1w1}) \in U_{t_u^0w, t_u^1w} \subseteq A_{1-\epsilon}.$$

As A_0 and A_1 are disjoint, $x_{t_u^0w0} \neq x_{t_u^0w1}$. Similarly, $y_{t_u^0w0} \neq y_{t_u^0w1}$.

So we may assume that $l \ge 1$ and $s \ne t_u^0 w$. The fact that G_l is a connected graph provides a G_l -path from s to $t_u^0 w$. This path gives us two $s(A_0 \cup A_1)$ -paths by the definition of U and V, one from y_{s0} to $x_{t_u^0 w0}$, and another one from y_{s1} to $x_{t_u^0 w1}$. Moreover, the same $\Phi(s', t')$'s are involved in these two pathes since they are induced by the same G_l -path. Observe that $(x_{t_u^0 w0}, y_{t_u^1 w1}), (x_{t_u^0 w1}, y_{t_u^1 w1})$ are in $s(A_0 \cup A_1)$. Also, since $x_{s\varepsilon} \in O_0$ and $y_{t\varepsilon'} \in O_1$, no "x" is equal to no "y". Thus, by quasi-acyclicity, $y_{s0} \ne y_{s1}$. Similarly, one can prove that $x_{s0} \ne x_{s1}$. The following picture illustrates the situation when l=1:



Let $\Phi(t_u^0w0, t_u^1w1) \in \omega$ such that $(x_{t_u^0w0}, y_{t_u^1w1}) \in C_{\Phi(t_u^0w0, t_u^1w1)}$, and $\Phi(s\varepsilon, t\varepsilon) := \Phi(s, t)$ if (s, t) is in $\mathcal{T}^\eta \cap (2^l \times 2^l)$ and $\varepsilon \in 2$. It remains to take disjoint Σ_1^1 sets $X_{s0}, X_{s1} \subseteq X_s$ (respectively $Y_{s0}, Y_{s1} \subseteq Y_s$) with the required properties, as well as $V_{s\varepsilon, t\varepsilon'}$, accordingly.

Case 2 $F_\eta \subseteq \Delta(X)$.

Let us indicate the differences with Case 1. We set $S := \{x \in X \mid (x, x) \in F_{\eta}\}$, which is a nonempty Σ_1^1 set by our assumption.

• We construct the following objects:

- a sequence $(x_s)_{s \in 2^{<\omega}}$ of points of S,
- a sequence $(X_s)_{s \in 2^{<\omega}}$ of Σ_1^1 subsets of X,
- a sequence $(U_{s,t})_{(s,t)\in\mathcal{T}^{\eta}}$ of Σ_1^1 subsets of X^2 , and $\Phi:\mathcal{T}^{\eta}\to\omega$.

We want these objects to satisfy the following conditions:

 $\begin{array}{l} (1) \ x_s \in X_s \ \land \ (x_s, x_t) \in U_{s,t} \\ (2) \ X_{s\varepsilon} \subseteq X_s \subseteq \Omega_X \cap S \ \land \ U_{s,t} \subseteq C_{\Phi(s,t)} \cap \Omega_{X^2} \cap (X_s \times X_t) \\ (3) \ \dim_{\operatorname{GH}}(X_s), \ \dim_{\operatorname{GH}}(U_{s,t}) \leq 2^{-|s|} \\ (4) \ X_{s0} \cap X_{s1} = \emptyset \\ (5) \ U_{s\varepsilon, t\varepsilon} \subseteq U_{s,t} \\ (6) \ U_{s,t} \subseteq F_{\theta} \ \text{if} \ (s,t) \in D_{\theta} \end{array}$

• Assume that this has been done. As in Case 1, we get $f: 2^{\omega} \to X$ injective continuous such that $\mathbb{N}_{\epsilon}^{\eta} \subseteq (f \times f)^{-1}(A_{\epsilon})$ for each $\epsilon \in 2$.

• So let us prove that the construction is possible. Let $(x_{\emptyset}, y_{\emptyset}) \in F_{\eta} \cap \Omega_{X^2}$. As $F_{\eta} \subseteq \Delta(X)$, $y_{\emptyset} = x_{\emptyset} \in S$. Let $\Phi(\emptyset, \emptyset) \in \omega$ with $(x_{\emptyset}, x_{\emptyset}) \in C_{\Phi(\emptyset,\emptyset)}$. As $\Omega_{X^2} \subseteq \Omega_X^2$, $x_{\emptyset} \in \Omega_X$. We choose a Σ_1^1 subset X_{\emptyset} of X with GH-diameter at most 1 such that $x_{\emptyset} \in X_{\emptyset} \subseteq \Omega_X \cap S$, as well as a Σ_1^1 subset $U_{\emptyset,\emptyset}$ of X^2 with GH-diameter at most 1 such that $(x_{\emptyset}, x_{\emptyset}) \in U_{\emptyset,\emptyset} \subseteq F_{\eta} \cap C_{\Phi(\emptyset,\emptyset)} \cap \Omega_{X^2} \cap (X_{\emptyset} \times X_{\emptyset})$, which completes the construction for the length l = 0.

For the inductive step, we set

$$\begin{split} U &:= \{ x \in X \mid \exists (x'_s)_{s \in 2^l} \in \Pi_{s \in 2^l} X_s \ x = x'_{t^0_u w} \land \forall (s,t) \in \mathcal{T}^\eta \cap (2^l \times 2^l) \ (x'_s, x'_t) \in U_{s,t} \}, \\ V &:= \{ x \in X \mid \exists (x'_s)_{s \in 2^l} \in \Pi_{s \in 2^l} X_s \ x = x'_{t^1_u w} \land \forall (s,t) \in \mathcal{T}^\eta \cap (2^l \times 2^l) \ (x'_s, x'_t) \in U_{s,t} \}. \end{split}$$

Again, we need to check that $x_{t_q^0} \neq x_{t_q^1}$ if $q \in \omega$. Note first that $A_1 \cap S^2$ is irreflexive, since otherwise it contains $(x, x) \in A_1 \cap F_\eta \subseteq A_1 \cap A_0$. By construction, $(x_{t_q^0}, x_{t_q^1}) \in F_{\varphi(q)} \subseteq A_1$, and we are done.

(2) \Rightarrow (3) Note that $(2^{\omega}, 2^{\omega}, \mathbb{N}_0^{\eta}, \mathbb{N}_1^{\eta}) \sqsubseteq (2^{\omega}, 2^{\omega}, \mathbb{B}_0^{\eta}, \mathbb{B}_1^{\eta})$, with witnesses $\alpha \to 0\alpha$ and $\beta \to 1\beta$.

(3) \Rightarrow (1) This comes from Lemma 3.1.

Proposition 3.4 Let η be a countable ordinal. The pairs $(\mathbb{N}_0^{\eta}, \mathbb{N}_1^{\eta})$ and $(\mathbb{B}_0^{\eta}, \mathbb{B}_1^{\eta})$ are incomparable for the square reduction.

Proof. There is no map $f: 2^{\omega} \to 2^{\omega}$ such that $\mathbb{N}_{\varepsilon}^{\eta} \subseteq (f \times f)^{-1}(\mathbb{B}_{\varepsilon}^{\eta})$ since $\Delta(2^{\omega})$ is a subset of \mathbb{N}_{0}^{η} .

There is no injection $f: 2^{\omega} \to 2^{\omega}$ for which there is $\alpha \in 2^{\omega}$ such that $f(0\alpha) = f(1\alpha)$. Using this fact, assume, towards a contradiction, that there is $f: 2^{\omega} \to 2^{\omega}$ injective continuous such that $\mathbb{B}^{\eta}_{\varepsilon} \subseteq (f \times f)^{-1}(\mathbb{N}^{\eta}_{\varepsilon})$. Let $(0t^{0}_{s}\gamma, 1t^{1}_{s}\gamma) \in \mathbb{B}^{\eta}_{\varepsilon}$, so that $(f(0t^{0}_{s}\gamma), f(1t^{1}_{s}\gamma)) = (t^{0}_{v}\gamma', t^{1}_{v}\gamma') \in \mathbb{N}^{\eta}_{\varepsilon}$.

We claim that $\varphi(s) \leq \varphi(v)$. We proceed by induction on $\varphi(s)$. Notice that is is obvious for $\varphi(s) = 0$. Suppose that it holds for all $\theta < \varphi(s)$. Note that we can find $p_k \in \omega$ and $\gamma_k \in 2^{\omega}$ such that $(t_{sp_k}^0 \gamma_k, t_{sp_k}^1 \gamma_k) \in \mathbb{N}_{1-\varepsilon}^{\eta}$ and $(t_{sp_k}^0 \gamma_k, t_{sp_k}^1 \gamma_k) \to (t_s^0 \gamma, t_s^1 \gamma)$. By continuity,

$$(t_{v_k}^0 \gamma', t_{v_k}^1 \gamma') := (f(0t_{sp_k}^0 \gamma_k), f(1t_{sp_k}^1 \gamma_k)) \to (t_v^0 \gamma', t_v^1 \gamma').$$

In particular, for k large, $(t_v^0, t_v^1) \subseteq (t_{v_k}^0, t_{v_k}^1)$. This implies that the sequence v_k is a strict extension of v. Therefore $\varphi(v_k) < \varphi(v)$. By the induction hypothesis, $\varphi(sp_k) \le \varphi(v_k) < \varphi(v)$. If $\varphi(s) = \theta + 1$, then $\theta = \varphi(sp_k) < \varphi(v)$, so we are done. If $\varphi(s)$ is a limit ordinal, then $(\varphi(sp_k))_{k \in \omega}$ is cofinal in it, so we are done too.

Finally, let $\alpha \in 2^{\omega}$, so that $(0\alpha, 1\alpha) = (0t_{\emptyset}^{0}\alpha, 1t_{\emptyset}^{1}\alpha) \in \mathbb{B}_{0}^{\eta}$. Then $(f(0\alpha), f(1\alpha)) = (t_{v}^{0}\gamma', t_{v}^{1}\gamma')$ with $\varphi(v) = \eta$, so that $v = \emptyset$, which contradicts the injectivity of f.

Consequences

Lemma 3.5 Let Γ be a class of sets contained in Δ_2^0 which is either a Wadge class or Δ_2^0 , X be a Polish space, and A, B be disjoint analytic relations on X. Then exactly one of the following holds:

- (a) the set A is separable from B by a $pot(\Gamma)$ set,
- (b) there are K_{σ} sets $A' \subseteq A$ and $B' \subseteq B$ such that A' is not separable from B' by a pot (Γ) set.

Proof. Assume that (a) does not hold. Theorems 1.9 and 1.10 in [L8] give Σ_2^0 relations $\mathbb{S}_0, \mathbb{S}_1$ on 2^{ω} and $g, h: 2^{\omega} \to X$ continuous with $\mathbb{S}_0 \subseteq (g \times h)^{-1}(A)$ and $\mathbb{S}_1 \subseteq (g \times h)^{-1}(B)$. We set $A' := (g \times h) [\mathbb{S}_0]$ and $B' := (g \times h) [\mathbb{S}_1]$.

Corollary 3.6 Let $\eta < \omega_1$, X be a Polish space, and A, B be disjoint analytic relations on X such that $A \cup B$ is s-acyclic or locally countable. Then exactly one of the following holds:

- (a) the set A is separable from B by a pot $(D_n(\Sigma_1^0))$ set,
- (b) $(2^{\omega}, 2^{\omega}, \mathbb{N}_0^{\eta}, \mathbb{N}_1^{\eta}) \sqsubseteq (X, X, A, B)$ if $\eta \ge 1$ and $(1, 1, \mathbb{N}_0^{\eta}, \mathbb{N}_1^{\eta}) \sqsubseteq (X, X, A, B)$ if $\eta = 0$.

Proof. By Lemma 3.1, \mathbb{N}_0^{η} is not separable from \mathbb{N}_1^{η} by a pot $(D_{\eta}(\Sigma_1^0))$ set. This shows that (a) and (b) cannot hold simultaneously. So assume that (a) does not hold. We may assume that $\eta \ge 1$. By Lemma 3.5, we may assume that A, B are Σ_2^0 . By Lemma 2.2, we may also assume that $A \cup B$ is quasi-acyclic. It remains to apply Theorem 3.3.

Corollary 3.7 Let η be a countable ordinal, X, Y be Polish spaces, and A, B be disjoint analytic subsets of $X \times Y$ such that $A \cup B$ is locally countable. Then exactly one of the following holds:

- (a) the set A is separable from B by a pot $(D_{\eta}(\Sigma_1^0))$ set,
- (b) $(2^{\omega}, 2^{\omega}, \mathbb{N}_0^{\eta}, \mathbb{N}_1^{\eta}) \sqsubseteq (X, Y, A, B)$ if $\eta \ge 1$ and $(1, 1, \mathbb{N}_0^{\eta}, \mathbb{N}_1^{\eta}) \sqsubseteq (X, Y, A, B)$ if $\eta = 0$.

Proof. We may assume that $\eta \ge 1$. As in the proof of Corollary 3.6, (a) and (b) cannot hold simultaneously. So assume that (a) does not hold. We put $Z := X \oplus Y$, $A' := \{((x, 0), (y, 1)) \in Z^2 \mid (x, y) \in A\}$ and $B' := \{((x, 0), (y, 1)) \in Z^2 \mid (x, y) \in B\}$. Then Z is Polish, A', B' are disjoint analytic relations on Z, $A' \cup B'$ is locally countable, and A' is not separable from B' by a pot $(D_{\eta}(\Sigma_1^0))$ set.

Corollary 3.6 gives $f', g': 2^{\omega} \to Z$ injective continuous such that $\mathbb{N}_0^{\eta} \subseteq (f' \times g')^{-1}(A')$, and also $\mathbb{N}_1^{\eta} \subseteq (f' \times g')^{-1}(B')$. We set $f(\alpha) := \Pi_0[f'(\alpha)]$, and $g(\beta) := \Pi_0[g'(\beta)]$. As $\Delta(2^{\omega}) \subseteq \mathbb{N}_0^{\eta}$, f' takes values in $X \times \{0\}$ and g' takes values in $Y \times \{1\}$. This implies that $f: 2^{\omega} \to X$, $g: 2^{\omega} \to Y$ are injective continuous. We are done since $\mathbb{N}_0^{\eta} \subseteq (f \times g)^{-1}(A)$ and $\mathbb{N}_1^{\eta} \subseteq (f \times g)^{-1}(B)$. \Box

Notation. If A is a relation on 2^{ω} , then we set $G_A := \{(0\alpha, 1\beta) \mid (\alpha, \beta) \in A\}$.

Lemma 3.8 Let A be an antisymmetric s-acyclic relation on 2^{ω} . Then G_A is s-acyclic.

Proof. We argue by contradiction, which gives $n \ge 2$ and an injective $s(G_A)$ -path $(\varepsilon_i z_i)_{i\le n}$ such that $(\varepsilon_0 z_0, \varepsilon_n z_n) \in s(G_A)$. This implies that $\varepsilon_i \ne \varepsilon_{i+1}$ if i < n and n is odd. Thus $(z_i)_{i\le n}$ is a s(A)-path such that $(z_{2j})_{2j\le n}$ and $(z_{2j+1})_{2j+1\le n}$ are injective and $(z_0, z_n) \in s(A)$. As s(A) is acyclic, the sequence $(z_i)_{i\le n}$ is not injective. We erase z_{2j+1} from this sequence if $z_{2j+1} \in \{z_{2j}, z_{2j+2}\}$ and $2j+1 \le n$, which gives a sequence $(z'_i)_{i\le n'}$ which is still a s(A)-path with $(z'_0, z'_{n'}) \in s(A)$, and moreover satisfies $z'_i \ne z'_{i+1}$ if i < n'.

If n' < 2, then n = 3, $z_0 = z_1$ and $z_2 = z_3$. As A is antisymmetric and $\varepsilon_3 = \varepsilon_1 \neq \varepsilon_2 = \varepsilon_0$, we get $z_0 = z_2$, which is absurd. If $n' \ge 2$, then $(z'_i)_{i \le n'}$ is not injective again. We choose a subsequence of it with at least three elements, made of consecutive elements, such that the first and the last elements are equal, and of minimal length with these properties. The acyclicity of s(A) implies that this subsequence has exactly three elements, say $(z'_i, z'_{i+1}, z'_{i+2} = z'_i)$.

If $z'_i = z_{2j+1}$, then $z'_{i+1} = z_{2j+2}$, $z'_{i+2} = z_{2j+4}$ and $z_{2j+3} = z_{2j+2}$. As A is antisymmetric and $\varepsilon_{2j+3} = \varepsilon_{2j+1} \neq \varepsilon_{2j+2} = \varepsilon_{2j+4}$, we get $z_{2j+2} = z_{2j+4}$, which is absurd. If $z'_i = z_{2j}$, then $z'_{i+1} = z_{2j+2}$, and $z'_{i+2} = z_{2j+3}$. As A is antisymmetric and $\varepsilon_{2j+3} = \varepsilon_{2j+1} \neq \varepsilon_{2j+2} = \varepsilon_{2j}$, we get $z_{2j} = z_{2j+2}$, which is absurd.

Corollary 3.9 Let $\eta \ge 1$ be a countable ordinal, X be a Polish space, and A, B be disjoint analytic relations on X. The following are equivalent:

(1) there is an s-acyclic relation $R \in \Sigma_1^1$ such that $A \cap R$ is not separable from $B \cap R$ by a pot $(D_\eta(\Sigma_1^0))$ set,

(2) there is a locally countable relation $R \in \Sigma_1^1$ such that $A \cap R$ is not separable from $B \cap R$ by a $pot(D_\eta(\Sigma_1^0))$ set,

(3) $(2^{\omega}, 2^{\omega}, \mathbb{N}^{\eta}_0, \mathbb{N}^{\eta}_1) \sqsubseteq (X, X, A, B),$

(4) there is $(\mathbb{A}_0, \mathbb{A}_1) \in \{(\mathbb{N}_0^{\eta}, \mathbb{N}_1^{\eta}), (\mathbb{B}_0^{\eta}, \mathbb{B}_1^{\eta})\}$ such that $(2^{\omega}, 2^{\omega}, \mathbb{A}_0, \mathbb{A}_1) \sqsubseteq (X, X, A, B)$, via a square map.

A similar result holds for $\eta = 0$ with 1 instead of 2^{ω} .

Proof. (1) \Rightarrow (3),(4) and (2) \Rightarrow (3),(4) This is a consequence of Corollary 3.6 and its proof.

(4) \Rightarrow (1) By the remarks before Lemma 3.1, $\mathbb{N}_0^{\eta} \cup \mathbb{N}_1^{\eta}$ has s-acyclic levels. This implies that $\mathbb{N}_0^{\eta} \cup \mathbb{N}_1^{\eta}$ is s-acyclic. As $\mathbb{N}_0^{\eta} \cup \mathbb{N}_1^{\eta}$ is antisymmetric, $\mathbb{B}_0^{\eta} \cup \mathbb{B}_1^{\eta}$ is s-acyclic too, by Lemma 3.8. Thus we can take $R := (f \times f)[\mathbb{A}_0 \cup \mathbb{A}_1]$ since the s-acyclicity is preserved by images by the square of an injection, and by Lemma 3.1.

(4) \Rightarrow (2) We can take $R := (f \times f)[\mathbb{A}_0 \cup \mathbb{A}_1]$ since $\mathbb{A}_0 \cup \mathbb{A}_1$ is locally countable, by Lemma 3.1.

(3) \Rightarrow (2) We can take $R := (f \times f)[\mathbb{N}_0^{\eta} \cup \mathbb{N}_1^{\eta}]$ since $\mathbb{N}_0^{\eta} \cup \mathbb{N}_1^{\eta}$ is locally countable, by Lemma 3.1. \Box

Remark. There is a version of Corollary 3.9 for $\check{D}_{\eta}(\Sigma_1^0)$ instead of $D_{\eta}(\Sigma_1^0)$, obtained by exchanging the roles of A and B. This symmetry is also present in Theorem 3.3.

We now give some complements when $\eta = 1$. At the beginning of this section, we mentioned the fact that our examples are in the style of \mathbb{G}_0 . If $\eta = 1$, then \mathbb{G}_0 itself is involved.

Corollary 3.10 Let X be a Polish space, and A, B be disjoint analytic relations on X such that

- either $A \cup B$ is s-acyclic or locally countable,

- or A is contained in a potentially closed s-acyclic or locally countable relation.

Then exactly one of the following holds:

(a) the set A is separable from B by a $pot(\mathbf{\Pi}_1^0)$ set,

(b) $(2^{\omega}, 2^{\omega}, \mathbb{G}_0, \Delta(2^{\omega})) \sqsubseteq (X, X, A, B).$

Corollary 3.11 Let X, Y be Polish spaces, and A, B be disjoint analytic subsets of $X \times Y$ such that $A \cup B$ is locally countable or A is contained in a potentially closed locally countable set. Then exactly one of the following holds:

- (a) the set A is separable from B by a $pot(\mathbf{\Pi}_1^0)$ set,
- (b) $(2^{\omega}, 2^{\omega}, \mathbb{G}_0, \Delta(2^{\omega})) \sqsubseteq (X, Y, A, B).$

4 The class $\Delta (D_{\eta}(\Sigma_1^0))$

Examples

Notation. We set, for each countable ordinal $\eta \ge 1$ and each $\varepsilon \in 2$,

$$\mathbb{S}_{\varepsilon}^{\eta} := \Big\{ (t_s^0 \gamma, t_s^1 \gamma) \mid s \in T_{\eta} \setminus \{ \emptyset \} \land \operatorname{parity}(|s|) = 1 - \big| \operatorname{parity}(s(0)) - \varepsilon \big| \land \gamma \in 2^{\omega} \Big\}.$$

Lemma 4.1 Let $\eta \geq 1$ be a countable ordinal, and C be a nonempty clopen subset of 2^{ω} . Then $\mathbb{S}_0^{\eta} \cap C^2$ is not separable from $\mathbb{S}_1^{\eta} \cap C^2$ by a pot $\left(\Delta\left(D_{\eta}(\boldsymbol{\Sigma}_1^0)\right)\right)$ set.

Proof. We use the notation in the proof of Lemma 3.1. We argue by contradiction, which gives P in $pot(\Delta(D_{\eta}(\Sigma_{1}^{0})))$, and a dense G_{δ} subset of 2^{ω} such that $P \cap G^{2}, G^{2} \setminus P \in D_{\eta}(\Sigma_{1}^{0})(G^{2})$. So let, for each $\varepsilon \in 2$, $(O_{\theta}^{\varepsilon})_{\theta < \eta}$ be a sequence of open relations on 2^{ω} such that

$$P \cap G^2 = \left(\bigcup_{\theta < \eta, \text{parity}(\theta) \neq \text{parity}(\eta)} O^0_{\theta} \setminus \left(\bigcup_{\theta' < \theta} O^0_{\theta'}\right)\right) \cap G^2$$

and $G^2 \setminus P = \left(\bigcup_{\theta < \eta, \text{parity}(\theta) \neq \text{parity}(\eta)} O^1_{\theta} \setminus \left(\bigcup_{\theta' < \theta} O^1_{\theta'} \right) \right) \cap G^2.$

• Note that $\mathbb{S}_{\varepsilon}^{\eta} = \bigcup_{s \in T_{\eta} \setminus \{\emptyset\}, \operatorname{parity}(|s|)=1-|\operatorname{parity}(s(0))-\varepsilon|} \operatorname{Gr}(f_s)$. Let us show that if $\theta \leq \eta$, $s \in T_{\eta}$ and $\varphi(s) = \theta$, then $\operatorname{Gr}(f_s) \cap (C \cap G)^2 \subseteq \neg O_{\theta}^{1-\operatorname{parity}(s(0))}$ if $\theta < \eta$, and $\operatorname{Gr}(f_s) \cap (C \cap G)^2$ is disjoint from $\bigcup_{\theta' < \theta} (O_{\theta'}^0 \cup O_{\theta'}^1)$ if $\theta = \eta$. The objects $s = \emptyset$ and $\theta = \eta$ will give the contradiction.

• We argue by induction on θ . Note that $\operatorname{Gr}(f_s) \cap (C \cap G)^2 \subseteq \mathbb{S}_{1-|\operatorname{parity}(|s|)-\operatorname{parity}(s(0))|}^{\eta} \cap G^2$ if $\theta = 0$ since $s \neq \emptyset$. As $\mathbb{S}_{\varepsilon}^{\eta} \cap G^2 \subseteq \neg O_0^{|\operatorname{parity}(\eta)-\varepsilon|}$ for each $\varepsilon \in 2$ and |s| has the same parity as η if $\theta = 0$, we are done.

• Assume that the result has been proved for $\theta' < \theta$. If θ is the successor of θ' , then the induction assumption implies that $\operatorname{Gr}(f_{sq}) \cap (C \cap G)^2 \subseteq \neg O_{\theta'}^{1-\operatorname{parity}((sq)(0))}$ for each q. We set, for each $\varepsilon \in 2$, $C_s^{\varepsilon} := \bigcup_{k \in \omega} \operatorname{Gr}(f_{s(2k+\varepsilon)})$, so that $\operatorname{Gr}(f_s) \subseteq \overline{C_s^{\varepsilon}}$, by the choice of ψ . If $s = \emptyset$, then

 $C^{\varepsilon}_{\emptyset} \cap (C \cap G)^2 \!\subseteq\! \neg O^{1-\varepsilon}_{\theta'},$

 $\operatorname{Gr}(f_s) \cap (C \cap G)^2 \subseteq \overline{C_{\emptyset}^{\varepsilon}} \cap (C \cap G)^2 \subseteq \overline{C_{\emptyset}^{\varepsilon}} \cap (C \cap G)^2 \subseteq \neg O_{\theta'}^{1-\varepsilon}$, which gives the desired inclusion for $\theta = \eta$.

If $s \neq \emptyset$, then $\operatorname{Gr}(f_{sq}) \cap (C \cap G)^2 \subseteq \neg O_{\theta'}^{1-\operatorname{parity}(s(0))}$ for each q, so that

$$\operatorname{Gr}(f_s) \cap (C \cap G)^2 \subseteq \overline{C_s} \cap (C \cap G)^2 \subseteq \overline{C_s \cap (C \cap G)^2} \subseteq \neg O_{\theta'}^{1-\operatorname{parity}(s(0))}.$$

Thus

$$\operatorname{Gr}(f_s) \cap (C \cap G)^2 \subseteq (G^2 \setminus O_{\theta'}^{1-\operatorname{parity}(s(0))}) \cap \neg (O_{\theta}^{1-\operatorname{parity}(s(0))} \setminus O_{\theta'}^{1-\operatorname{parity}(s(0))}) \subseteq \neg O_{\theta}^{1-\operatorname{parity}(s(0))}$$

since $\operatorname{parity}(\theta) = |\operatorname{parity}(|s|) - \operatorname{parity}(\eta)|.$

• If θ is limit, then $(\varphi(sn))_{n\in\omega}$ is cofinal in $\varphi(s)$, and $\operatorname{Gr}(f_{sn}) \cap (C \cap G)^2 \subseteq \neg O_{\varphi(sn)}^{1-\operatorname{parity}((sn)(0))}$, by the induction assumption. If $\theta_0 < \varphi(s)$, then there is $n(\theta_0)$ such that $\varphi(sn) > \theta_0$ if $n \ge n(\theta_0)$. Thus $\operatorname{Gr}(f_{sn}) \cap (C \cap G)^2 \subseteq \neg O_{\theta_0}^{1-\operatorname{parity}((sn)(0))}$ if $n \ge n(\theta_0)$. If $s = \emptyset$, then, for each $\varepsilon \in 2$,

$$\operatorname{Gr}(f_s) \cap (C \cap G)^2 \subseteq \underbrace{(C \cap G)^2 \cap \overline{C_s^{\varepsilon}} \setminus C_s^{\varepsilon} = \overline{C_s^{\varepsilon} \cap (C \cap G)^2} \setminus C_s^{\varepsilon}}_{\subseteq \bigcup_{n \ge n(\theta_0), \operatorname{parity}(n) = \varepsilon} \operatorname{Gr}(f_{sn}) \cap (C \cap G)^2} \subseteq \neg O_{\theta_0}^{1 - \varepsilon}.$$

Thus $\operatorname{Gr}(f_s) \cap (C \cap G)^2 \subseteq \neg \left(\bigcup_{\theta' < \eta} (O^0_{\theta'} \cup O^1_{\theta'})\right)$. If $s \neq \emptyset$, then $\operatorname{Gr}(f_{sn}) \cap (C \cap G)^2 \subseteq \neg O^{1-\operatorname{parity}(s(0))}_{\theta_0}$ for each n, so that $\operatorname{Gr}(f_s) \cap (C \cap G)^2 \subseteq \overline{C_s} \cap (C \cap G)^2 \subseteq \overline{C_s \cap (C \cap G)^2} \subseteq \neg O^{1-\operatorname{parity}(s(0))}_{\theta_0}$. As $\operatorname{parity}(|s|) = \operatorname{parity}(\eta)$, $\operatorname{Gr}(f_s) \cap (C \cap G)^2 \subseteq \neg O^{1-\operatorname{parity}(s(0))}_{\theta}$ as above. \Box

A topological characterization

Notation. We define, for $1 \le \xi < \omega_1^{\mathsf{CK}}$ and $\eta < \omega_1^{\mathsf{CK}}$, $\bigcap_{\theta < 0} G_{\theta,\xi} := (\omega^{\omega})^2$, and, inductively, $G_{\eta,\xi} := \left\{ \begin{array}{l} \bigcap_{\theta < \eta} G_{\theta,\xi} \text{ if } \eta \text{ is limit (possibly 0),} \\ \overline{A_0 \cap G_{\theta,\xi}}^{\tau_{\xi}} \cap \overline{A_1 \cap G_{\theta,\xi}}^{\tau_{\xi}} \text{ if } \eta = \theta + 1. \end{array} \right.$ **Theorem 4.2** Let $1 \le \xi < \omega_1^{CK}$, $1 \le \eta = \lambda + 2k + \varepsilon < \omega_1^{CK}$ with λ limit, $k \in \omega$ and $\varepsilon \in 2$, and A_0 , A_1 be disjoint Σ_1^1 relations on ω^{ω} . Then the following are equivalent:

(1) the set A_0 is not separable from A_1 by a pot $\left(\Delta\left(D_\eta(\Sigma_{\xi}^0)\right)\right)$ set,

(2) the Σ_1^1 set $G_{\eta,\xi}$ is not empty.

Proof. The proof is in the spirit of that of Theorem 3.2. The proof of Theorem 1.10.(2) in [L8] gives α suitable such that $c(\alpha)$ codes the class $D_{\eta}(\Sigma_{\xi}^{0})$. By Theorem 6.26 in [L8] and Theorem 3.2, (1) is equivalent to $R'(\alpha, a_0, a_1) \neq \emptyset$, where

$$R'(\alpha, a_0, a_1) := \begin{cases} F_{\theta, \xi}^0 \cap F_{\theta, \xi}^1 \text{ if } \eta = \theta + 1, \\ \bigcap_{p \ge 1} F_{\theta_p, \xi}^0 \text{ if } \eta = \sup_{p \ge 1} \theta_p \text{ is limit } \wedge \theta_p \text{ is odd.} \end{cases}$$

So it is enough to prove that

$$G_{\eta,\xi} = \begin{cases} F_{\theta,\xi}^0 \cap F_{\theta,\xi}^1 \text{ if } \eta = \theta + 1, \\ \bigcap_{p \ge 1} F_{\theta_p,\xi}^0 \text{ if } \eta = \sup_{p \ge 1} \theta_p \text{ is limit } \wedge \theta_p \text{ is odd.} \end{cases}$$

We argue by induction on η . Note first that $G_{1,\xi} = \overline{A_0} \cap \overline{A_1} = F_{0,\xi}^0 \cap F_{0,\xi}^1$. Then, inductively,

$$G_{\theta+2,\xi} = \overline{A_0 \cap G_{\theta+1,\xi}} \cap \overline{A_1} \cap \overline{G_{\theta+1,\xi}} = \overline{A_0 \cap F_{\theta,\xi}^0} \cap F_{\theta,\xi}^1 \cap \overline{A_1} \cap F_{\theta,\xi}^0 \cap F_{\theta,\xi}^1$$
$$= \overline{A_0 \cap F_{\theta,\xi}^{1-\operatorname{parity}(\theta)}} \cap \overline{A_1 \cap F_{\theta,\xi}^{\operatorname{parity}(\theta)}} = F_{\theta+1,\xi}^0 \cap F_{\theta+1,\xi}^1.$$

If λ is limit, then

$$\begin{aligned} G_{\lambda+1,\xi} = &\overline{A_0 \cap G_{\lambda,\xi}} \cap \overline{A_1 \cap G_{\lambda,\xi}} = \overline{A_0 \cap \bigcap_{\theta < \lambda} G_{\theta,\xi}} \cap \overline{A_1 \cap \bigcap_{\theta < \lambda} G_{\theta,\xi}} \\ = &\overline{A_0 \cap \bigcap_{\theta < \lambda} G_{\theta+1,\xi}} \cap \overline{A_1 \cap \bigcap_{\theta < \lambda} G_{\theta+1,\xi}} \\ = &\overline{A_0 \cap \bigcap_{\theta < \lambda} F_{\theta,\xi}^0} \cap \overline{F_{\theta,\xi}^1} \cap \overline{A_1 \cap \bigcap_{\theta < \lambda} F_{\theta,\xi}^0} \cap \overline{F_{\theta,\xi}^1} \\ = &\overline{A_0 \cap \bigcap_{\theta < \lambda} F_{\theta,\xi}^0} \cap \overline{A_1 \cap \bigcap_{\theta < \lambda} F_{\theta,\xi}^1} = F_{\lambda,\xi}^0 \cap F_{\lambda,\xi}^1 \end{aligned}$$

and $G_{\lambda,\xi} = \bigcap_{\theta < \lambda} G_{\theta,\xi} = \bigcap_{\theta < \lambda} G_{\theta+1,\xi} = \bigcap_{\theta < \lambda} F_{\theta,\xi}^0 \cap F_{\theta,\xi}^1 = \bigcap_{\theta < \lambda} F_{\theta,\xi}^0 = \bigcap_{p \ge 1} F_{\theta_p,\xi}^0$.

The main result

We prove a version of Theorem 3.3 for the class $\Delta(D_{\eta}(\Sigma_1^0))$. We set, for $1 \leq \eta < \omega_1$ and $\varepsilon \in 2$, $\mathbb{C}_{\varepsilon}^{\eta} := \{(0\alpha, 1\beta) \mid (\alpha, \beta) \in \mathbb{S}_{\varepsilon}^{\eta}\}.$

Theorem 4.3 Let $\eta \ge 1$ be a countable ordinal, X be a Polish space, and A_0, A_1 be disjoint analytic relations on X such that $A_0 \cup A_1$ is contained in a potentially closed quasi-acyclic relation. The following are equivalent:

(1) the set A_0 is not separable from A_1 by a pot $\left(\Delta\left(D_{\eta}(\Sigma_1^0)\right)\right)$ set,

(2) there is $(\mathbb{A}_0, \mathbb{A}_1) \in \{(\mathbb{N}_1^{\eta}, \mathbb{N}_0^{\eta}), (\mathbb{B}_1^{\eta}, \mathbb{B}_0^{\eta}), (\mathbb{N}_0^{\eta}, \mathbb{N}_1^{\eta}), (\mathbb{B}_0^{\eta}, \mathbb{B}_1^{\eta}), (\mathbb{S}_0^{\eta}, \mathbb{S}_1^{\eta}), (\mathbb{C}_0^{\eta}, \mathbb{C}_1^{\eta})\}$ for which the inequality $(2^{\omega}, 2^{\omega}, \mathbb{A}_0, \mathbb{A}_1) \sqsubseteq (X, X, A_0, A_1)$ holds, via a square map, (3) there is $(\mathbb{A}_0, \mathbb{A}_1) \in \{(\mathbb{N}_1^{\eta}, \mathbb{N}_0^{\eta}), (\mathbb{N}_0^{\eta}, \mathbb{N}_1^{\eta}), (\mathbb{S}_0^{\eta}, \mathbb{S}_1^{\eta})\}$ such that $(2^{\omega}, 2^{\omega}, \mathbb{A}_0, \mathbb{A}_1) \sqsubseteq (X, X, A_0, A_1)$. **Proof.** (1) \Rightarrow (2) The proof is partly similar to that of Theorem 3.3. Let R be a potentially closed quasi-acyclic relation containing $A_0 \cup A_1$, and $(C_n)_{n \in \omega}$ be a witness for the fact that R is quasi-acyclic. We may assume that X is zero-dimensional (and thus a closed subset of ω^{ω}) and R is closed. In fact, we may assume that $X = \omega^{\omega}$. Indeed, as A_0 is not separable from A_1 by a pot $\left(\Delta\left(D_\eta(\Sigma_1^0)\right)\right)$ set in X^2 , it is also the case in $(\omega^{\omega})^2$, which gives $f: 2^{\omega} \to \omega^{\omega}$. Note that

$$\mathbb{A}_0 \cup \mathbb{A}_1 \subseteq (f \times f)^{-1} (A_0 \cup A_1) \subseteq (f \times f)^{-1} (X^2),$$

which implies that $\overline{\mathbb{A}_0 \cup \mathbb{A}_1} \subseteq (f \times f)^{-1}(X^2)$. As $\Delta(2^{\omega}) \subseteq \mathbb{N}_0^{\eta} \cap \overline{\mathbb{S}_0^{\eta} \cup \mathbb{S}_1^{\eta}}$ and

$$\{(0\alpha,1\alpha) \mid \alpha \in 2^{\omega}\} \subseteq \mathbb{B}_0^{\eta} \cap \overline{\mathbb{C}_0^{\eta} \cup \mathbb{C}_1^{\eta}},\$$

the range of $\Delta(2^{\omega})$ by $f \times f$ is a subset of X^2 , so that f takes values in X. We may also assume that A_0, A_1 are Σ_1^1 , and that the relation " $(x, y) \in C_p$ " is Δ_1^1 in (x, y, p). By Theorem 4.2, G_{η} is a nonempty Σ_1^1 relation on X (we denote $G_{\eta} := G_{\eta,1}$ and $F_{\eta}^{\varepsilon} := F_{\eta,1}^{\varepsilon}$, for simplicity). We also consider F_{θ} with $F_{\theta}^{\varepsilon} := \overline{F_{\theta}}^{\tau_1}$. In the sequel, all the closures will refer to the topology τ_1 , so that, for example,

$$G_{\eta} \cup A_0 \cup A_1 \subseteq \overline{A_0 \cup A_1} \subseteq R = \bigcup_{n \in \omega} C_n.$$

• Let us show that $A_{\epsilon} \cap G_{\eta} \subseteq F_{\eta}^{|\operatorname{parity}(\eta)-\epsilon|}$ if $\epsilon \in 2$. We argue by induction on η . If $\eta = 1$, then $A_{\epsilon} \cap G_1 \subseteq A_{\epsilon} \cap \overline{A_{1-\epsilon}} \subseteq F_1^{1-\epsilon}$. If η is limit, then $A_{\epsilon} \cap G_{\eta} \subseteq A_{\epsilon} \cap \bigcap_{\theta < \eta} F_{\theta}^{\epsilon} \subseteq F_{\eta}^{\epsilon}$. Finally, if $\eta = \theta + 1$, then without loss of generality suppose that θ is even, so that η is odd and

$$A_{\epsilon} \cap G_{\eta} \subseteq A_{\epsilon} \cap \overline{A_{1-\epsilon} \cap G_{\theta}} \subseteq A_{\epsilon} \cap F_{\theta}^{1-\epsilon}$$

Note that this last set is contained in $F_{\eta}^{1-\epsilon},$ as required.

So, if $A_{\epsilon} \cap G_{\eta} \neq \emptyset$ for some $\epsilon \in 2$ and e is the correct digit, then $F_{\eta}^{e} \neq \emptyset$. Theorem 3.3 gives $(\mathbb{A}_{0}, \mathbb{A}_{1}) \in \{(\mathbb{N}_{1}^{\eta}, \mathbb{N}_{0}^{\eta}), (\mathbb{B}_{1}^{\eta}, \mathbb{B}_{0}^{\eta}), (\mathbb{B}_{0}^{\eta}, \mathbb{B}_{1}^{\eta})\}$ for which $(2^{\omega}, 2^{\omega}, \mathbb{A}_{0}, \mathbb{A}_{1}) \sqsubseteq (X, X, A_{0}, A_{1})$, via a square map.

• Thus, in the sequel, we suppose that $G_\eta \cap (A_0 \cup A_1) = \emptyset$. We put

$$D_{\eta} := \left\{ \left(t_s^0 w, t_s^1 w \right) \in \mathcal{T}^{\eta} \mid s = \emptyset \right\} = \Delta(2^{<\omega})$$

and, for $\theta < \eta$ and $\epsilon \in 2$,

$$D_{\theta}^{\epsilon} := \Big\{ (t_s^0 w, t_s^1 w) \in \mathcal{T}^{\eta} \mid s \in T_{\eta} \setminus \{ \emptyset \} \land \varphi(s) = \theta \land \operatorname{parity}(|s|) = 1 - \big| \operatorname{parity}(s(0)) - \epsilon \big| \Big\},$$

so that $\{D_{\eta}\} \cup \{D_{\theta}^{\epsilon} \mid \theta < \eta \land \epsilon \in 2\}$ defines a partition of \mathcal{T}^{η} .

Case 1 $G_\eta \not\subseteq \Delta(X)$.

Let $(x, y) \in G_{\eta} \setminus \Delta(X)$, and O_0, O_1 be disjoint Δ_1^0 sets with $(x, y) \in O_0 \times O_1$. We can replace G_{η}, A_0 and A_1 with their intersection with $O_0 \times O_1$ if necessary and assume that they are contained in $O_0 \times O_1$. Let us indicate the differences with the proof of Theorem 3.3.

• Condition (6) is changed as follows:

(6)
$$U_{s,t} \subseteq \begin{cases} G_{\eta} \text{ if } (s,t) \in D_{\eta} \\ A_{\epsilon} \cap G_{\theta} \text{ if } (s,t) \in D_{\theta} \end{cases}$$

• If $(0\alpha, 1\beta) \in \mathbb{C}_{\epsilon}^{\eta}$, then there is $\theta < \eta$ such that $(\alpha, \beta) | n \in D_{\theta}^{\epsilon}$ if $n \ge n_0$. In this case, $(U_{(\alpha,\beta)|n})_{n \ge n_0}$ is a decreasing sequence of nonempty clopen subsets of $A_{\epsilon} \cap \Omega_{X^2}$ with vanishing diameters, so that its intersection is a singleton $\{F(\alpha, \beta)\} \subseteq A_{\epsilon}$, and $(f(0\alpha), f(1\beta)) = F(\alpha, \beta) \in A_{\epsilon}$.

• So let us prove that the construction is possible. Let $(x_{\emptyset}, y_{\emptyset}) \in G_{\eta} \cap \Omega_{X^2}$. We choose a Σ_1^1 subset $U_{\emptyset,\emptyset}$ of X^2 such that $(x_{\emptyset}, y_{\emptyset}) \in U_{\emptyset,\emptyset} \subseteq G_{\eta} \cap C_{\Phi(\emptyset,\emptyset)} \cap \Omega_{X^2} \cap (X_{\emptyset} \times Y_{\emptyset})$, which completes the construction for the length l = 0. Assume that we have constructed our objects for the sequences of length l. Note that $(x_{t_u^0 w}, y_{t_u^1 w}) \in G_{\varphi(u)} \cap (U \times V) \subseteq G_{\varphi(uq)+1} \cap (U \times V) \subseteq \overline{A_{\epsilon} \cap G_{\varphi(uq)}} \cap (U \times V)$, where ϵ satisfies $(t_{uq}^0, t_{uq}^1) \in D_{\varphi(uq)}^{\epsilon}$. This gives $(x_{t_u^0 w0}, y_{t_u^1 w1}) \in A_{\epsilon} \cap G_{\varphi(uq)} \cap (U \times V) \cap \Omega_{X^2}$. If $u = \emptyset$, then $(t_u^0 w1, t_u^1 w1) \in D_{\eta}$, so that $(x_{t_u^0 w1}, y_{t_u^1 w1}) \in U_{t_u^0 w, t_u^1 w} \subseteq G_{\eta}$ and $(x_{t_u^0 w0}, y_{t_u^1 w1}) \in A_{\epsilon}$. As $G_{\eta} \cap (A_0 \cup A_1) = \emptyset$, $x_{t_u^0 w0} \neq x_{t_u^0 w1}$. Similarly, $y_{t_u^0 w0} \neq y_{t_u^0 w1}$. If $u \neq \emptyset$, then we argue as in the proof of Theorem 3.3 to see that $x_{s0} \neq x_{s1}$ (and similarly for y_{s0} and y_{s1}).

Case 2 $G_\eta \subseteq \Delta(X)$.

Let us indicate the differences with the proof of Theorem 3.3 and Case 1. We set

$$S := \{ x \in X \mid (x, x) \in G_{\eta} \},\$$

which is a nonempty Σ_1^1 set by our assumption. We get $f: 2^{\omega} \to X$ injective continuous such that $\mathbb{S}_{\epsilon}^{\eta} \subseteq (f \times f)^{-1}(A_{\epsilon})$ for each $\epsilon \in 2$. In this case, $A_0 \cap S^2$ and $A_1 \cap S^2$ are irreflexive.

(2) \Rightarrow (3) Note that $(2^{\omega}, 2^{\omega}, \mathbb{N}_0^{\eta}, \mathbb{N}_1^{\eta}) \subseteq (2^{\omega}, 2^{\omega}, \mathbb{B}_0^{\eta}, \mathbb{B}_1^{\eta})$ and $(2^{\omega}, 2^{\omega}, \mathbb{S}_0^{\eta}, \mathbb{S}_1^{\eta}) \subseteq (2^{\omega}, 2^{\omega}, \mathbb{C}_0^{\eta}, \mathbb{C}_1^{\eta})$, with witnesses $\alpha \to 0\alpha$ and $\beta \to 1\beta$.

 $(3) \Rightarrow (1)$ This comes from Lemmas 3.1 and 4.1.

Proposition 4.4 *Let* $\eta \ge 1$ *be a countable ordinal.*

(a) If η is a successor ordinal, then the pairs $(\mathbb{N}_1^{\eta}, \mathbb{N}_0^{\eta}), (\mathbb{B}_1^{\eta}, \mathbb{B}_0^{\eta}), (\mathbb{N}_0^{\eta}, \mathbb{N}_1^{\eta}), (\mathbb{B}_0^{\eta}, \mathbb{B}_1^{\eta}), (\mathbb{S}_0^{\eta}, \mathbb{S}_1^{\eta})$ and $(\mathbb{C}_0^{\eta}, \mathbb{C}_1^{\eta})$ are incomparable for the square reduction.

(b) If η is a limit ordinal, then $(2^{\omega}, 2^{\omega}, \mathbb{S}_0^{\eta}, \mathbb{S}_1^{\eta}) \sqsubseteq (2^{\omega}, 2^{\omega}, \mathbb{N}_1^{\eta}, \mathbb{N}_0^{\eta}), (2^{\omega}, 2^{\omega}, \mathbb{N}_0^{\eta}, \mathbb{N}_1^{\eta})$ and

 $(2^{\omega}, 2^{\omega}, \mathbb{C}^{\eta}_0, \mathbb{C}^{\eta}_1) \sqsubseteq (2^{\omega}, 2^{\omega}, \mathbb{B}^{\eta}_1, \mathbb{B}^{\eta}_0), (2^{\omega}, 2^{\omega}, \mathbb{B}^{\eta}_0, \mathbb{B}^{\eta}_1),$

via a square map, and the pairs $(\mathbb{S}_0^{\eta}, \mathbb{S}_1^{\eta})$ and $(\mathbb{C}_0^{\eta}, \mathbb{C}_1^{\eta})$ are incomparable for the square reduction.

Proof. (a) We set, for $\theta \leq \eta$, $C_{\theta} := \bigcup_{\varphi(s) > \theta} \operatorname{Gr}(f_s)$.

Claim. Let $\theta \leq \eta$. Then C_{θ} is a closed relation on 2^{ω} .

Indeed, this is inspired by the proof of Theorem 2.3 in [L2].

We first show that $C^l := \bigcup_{s \in \omega^{\leq l}, \varphi(s) \geq \theta} \operatorname{Gr}(f_s)$ is closed, by induction on $l \in \omega$. This is clear for l = 0. Assume that the statement is true for l. Note that $C^{l+1} = C^l \cup \bigcup_{s \in \omega^{l+1}, \varphi(s) \geq \theta} \operatorname{Gr}(f_s)$. Let $p_m \in C^{l+1}$ such that $(p_m)_{m \in \omega}$ converges to p. By induction assumption, we may assume that, for each m, there is $(s_m, n_m) \in \omega^l \times \omega$ such that $\varphi(s_m n_m) \geq \theta$ and $p_m \in \operatorname{Gr}(f_{s_m n_m})$. As the $\operatorname{Gr}(f_{s_n})$'s are closed, we may assume that there is $i \leq l$ such that the sequence $((s_m n_m)|i)_{m \in \omega}$ is constant and the sequence $((s_m n_m)(i))_{m \in \omega}$ tends to infinity. This implies that $p \in \operatorname{Gr}(f_{(s_0 n_0)|i}) \subseteq C^{l+1}$, which is therefore closed.

Now let $p_m \in C_{\theta}$ such that $(p_m)_{m \in \omega}$ converges to p. The previous fact implies that we may assume that, for each m, there is s'_m such that $\varphi(s'_m) \ge \theta$ and $p_m \in \operatorname{Gr}(f_{s'_m})$, and that the sequence $(|s'_m|)_{m \in \omega}$ tends to infinity. Note that there is l such that the set of $s'_m(l)$'s is infinite. Indeed, assume, towards a contradiction, that this is not the case. Then $\{s \in T_\eta \mid \exists m \in \omega \mid s \subseteq s'_m\}$ is an infinite finitely branching subtree of T_η . By König's lemma, it has an infinite branch, which contradicts the wellfoundedness of T_η . So we may assume that there is l such that the sequence $(s'_m|l)_{m \in \omega}$ is constant and the sequence $(s'_m(l))_{m \in \omega}$ tends to infinity. This implies that $p \in \operatorname{Gr}(f_{s'_0|l}) \subseteq C_{\theta}$.

• By Lemma 3.1, \mathbb{N}_0^{η} is not separable from \mathbb{N}_1^{η} by a pot $(D_{\eta}(\Sigma_1^0))$ set, and, by Lemma 4.1, \mathbb{S}_0^{η} is not separable from \mathbb{S}_1^{η} by a pot $(\Delta(D_{\eta}(\Sigma_1^0)))$ set.

• Let us show that \mathbb{N}_0^{η} is separable from \mathbb{N}_1^{η} by a $\check{D}_{\eta}(\Sigma_1^0)$ set. In fact, it is enough to see that $\mathbb{N}_0^{\eta} \in \check{D}_{\eta}(\Sigma_1^0)$ if η is odd and $\mathbb{N}_1^{\eta} \in D_{\eta}(\Sigma_1^0)$ if η is even. If η is odd, then

$$\mathbb{N}_{0}^{\eta} = \bigcup_{s \in T_{\eta}, \varphi(s) \text{ odd }} \operatorname{Gr}(f_{s}) = C_{\eta} \cup \bigcup_{\theta < \eta, \theta \text{ odd }} C_{\theta} \setminus C_{\theta+1}.$$

We set, for $\theta < \eta$, $O_{\theta} := \neg C_{\theta+1}$, which defines an increasing sequence of open relations on 2^{ω} with $\mathbb{N}_{0}^{\eta} = \neg O_{\eta-1} \cup \bigcup_{\theta < \eta, \theta \text{ odd}} O_{\theta} \setminus O_{\theta-1}$. Thus $\mathbb{N}_{0}^{\eta} \in \check{D}_{\eta}(\Sigma_{1}^{0})$. Similarly, if η is even, then $\mathbb{N}_{1}^{\eta} = \bigcup_{s \in T_{\eta}, f_{\eta}(s) \text{ odd}} \operatorname{Gr}(f_{s}) = \bigcup_{\theta < \eta, \theta \text{ odd}} C_{\theta} \setminus C_{\theta+1}$. We set, for $\theta < \eta$, $O_{\theta} := \neg C_{\theta+1}$, which defines an increasing sequence of open relations on 2^{ω} with $\mathbb{N}_{1}^{\eta} = \bigcup_{\theta < \eta, \theta \text{ odd}} O_{\theta} \setminus O_{\theta-1}$. Thus $\mathbb{N}_{1}^{\eta} \in D_{\eta}(\Sigma_{1}^{0})$. This shows that $(2^{\omega}, 2^{\omega}, \mathbb{N}_{1}^{\eta}, \mathbb{N}_{0}^{\eta})$ is not \sqsubseteq -below $(2^{\omega}, 2^{\omega}, \mathbb{N}_{0}^{\eta}, \mathbb{N}_{1}^{\eta})$, and consequently that $(2^{\omega}, 2^{\omega}, \mathbb{N}_{0}^{\eta}, \mathbb{N}_{1}^{\eta})$ is not \sqsubseteq -below $(2^{\omega}, 2^{\omega}, \mathbb{N}_{1}^{\eta}, \mathbb{N}_{0}^{\eta})$.

• Let us show that $\mathbb{S}_{\varepsilon}^{\eta}$ is separable from $\mathbb{S}_{1-\varepsilon}^{\eta}$ by a $\check{D}_{\eta}(\Sigma_{1}^{0})$ set if $\varepsilon \in 2$. We set, for $\theta \leq \eta$,

$$C^{\varepsilon}_{\theta} := \bigcup_{\varphi(s) \ge \theta, \text{ parity}(s(0)) = \varepsilon} \operatorname{Gr}(f_s).$$

As in the claim, $(C^{\varepsilon}_{\theta})_{\theta < \eta}$ is a decreasing sequence of closed sets.

Note that

$$\begin{split} \mathbb{S}_{\varepsilon}^{\eta} &= \bigcup_{s \in T_{\eta} \setminus \{\emptyset\}, \text{ parity}(|s|) = 1 - |\text{parity}(s(0)) - \varepsilon|} \operatorname{Gr}(f_{s}) \\ &= \bigcup_{s \in T_{\eta} \setminus \{\emptyset\}, \text{ parity}(\varphi(s)) - \text{parity}(\eta)| = 1 - |\text{parity}(s(0)) - \varepsilon|} \operatorname{Gr}(f_{s}) \\ &= \bigcup_{s \in T_{\eta} \setminus \{\emptyset\}, \text{ parity}(s(0)) = |1 - ||\text{parity}(\varphi(s)) - \text{parity}(\eta)| - \varepsilon||} \operatorname{Gr}(f_{s}) \\ &= \bigcup_{\theta < \eta, \varphi(s) = \theta} \bigcup_{parity}(s(0)) = |1 - ||\text{parity}(\theta) - \text{parity}(\eta)| - \varepsilon||} \operatorname{Gr}(f_{s}) \\ &= \bigcup_{\theta < \eta} \left(\bigcup_{\varphi(s) \ge \theta, \text{ parity}(s(0)) = |1 - ||\text{parity}(\theta) - \text{parity}(\eta)| - \varepsilon||} \operatorname{Gr}(f_{s}) \right) \\ &\qquad \left(\bigcup_{\varphi(s) \ge \theta + 1, \text{ parity}(s(0)) = |1 - ||\text{parity}(\theta) - \text{parity}(\eta)| - \varepsilon||} \operatorname{Gr}(f_{s}) \right) \\ &= \bigcup_{\theta < \eta} C_{\theta}^{1 - ||\text{parity}(\theta) - \text{parity}(\eta)| - \varepsilon|} \setminus C_{\theta + 1}^{1 - ||\text{parity}(\theta) - \text{parity}(\eta)| - \varepsilon|}. \end{split}$$

Assume first that $\eta = \theta_0 + 1$ is a successor ordinal. We define an increasing sequence $(O_\theta)_{\theta < \eta}$ of open sets as follows:

$$O_{\theta} := \begin{cases} \neg (C_{\theta+1}^{1-\varepsilon} \cup C_{\theta}^{\varepsilon}) \text{ if } \theta < \theta_{0}, \\ \neg C_{\theta}^{\varepsilon} \text{ if } \theta = \theta_{0}, \end{cases}$$

so that $D := \neg D((O_{\theta})_{\theta < \eta}) \in \check{D}_{\eta}(\Sigma_1^0).$

We now check that D separates $\mathbb{S}^{\eta}_{\varepsilon}$ from $\mathbb{S}^{\eta}_{1-\varepsilon}$. If $\theta < \eta$ has a parity opposite to that of η , then either $\theta = \theta_0$ and $C^{\varepsilon}_{\theta} \setminus C^{\varepsilon}_{\theta+1} \subseteq C^{\varepsilon}_{\theta_0} \subseteq \neg(\bigcup_{\theta' < \eta} O_{\theta'}) \subseteq D$. Or $\theta < \theta_0$, $\theta+1 < \theta_0 < \eta$ has the same parity as η , and $C^{\varepsilon}_{\theta} \setminus C^{\varepsilon}_{\theta+1} \subseteq O_{\theta+1} \setminus (\bigcup_{\theta' \leq \theta} O_{\theta'}) \subseteq D$. If now $\theta < \eta$ has the same parity as η , then $C^{1-\varepsilon}_{\theta} \setminus C^{1-\varepsilon}_{\theta+1} \subseteq O_{\theta} \setminus (\bigcup_{\theta' < \theta} O_{\theta'}) \subseteq D$. Thus $\mathbb{S}^{\eta}_{\varepsilon} \subseteq D$. Similarly, $\mathbb{S}^{\eta}_{1-\varepsilon} \subseteq \neg D$. If η is a limit ordinal, then we set $O_{\theta} := \neg(C^{1-\varepsilon}_{\theta+1} \cup C^{\varepsilon}_{\theta})$ and argue similarly. This shows that $(2^{\omega}, 2^{\omega}, \mathbb{N}^{\eta}_{\varepsilon}, \mathbb{N}^{\eta}_{1-\varepsilon})$ is not \sqsubseteq -below $(2^{\omega}, 2^{\omega}, \mathbb{S}^{\eta}_{0}, \mathbb{S}^{\eta}_{1})$ for each $\varepsilon \in 2$.

• Let us prove that $(2^{\omega}, 2^{\omega}, \mathbb{S}_0^{\eta}, \mathbb{S}_1^{\eta})$ is not \sqsubseteq -below $(2^{\omega}, 2^{\omega}, \mathbb{N}_{\varepsilon}^{\eta}, \mathbb{N}_{1-\varepsilon}^{\eta})$ if $\varepsilon \in 2$ and η is a successor ordinal. Let us do it for $\varepsilon = 0$, the other case being similar. We argue by contradiction, which gives f, g injective continuous with $\mathbb{S}_{\varepsilon}^{\eta} \subseteq (f \times g)^{-1}(\mathbb{N}_{\varepsilon}^{\eta})$ for each $\varepsilon \in 2$. We set, for $\theta < \eta$ and $\varepsilon \in 2$,

$$U_{\theta}^{\varepsilon} := \bigcup_{\substack{\theta \leq \theta' < \eta, \varphi(s) = \theta', \text{ parity}(s(0)) = |1 - || \text{parity}(\theta') - \text{parity}(\eta)| - \varepsilon ||}} \text{Gr}(f_s).$$

Note that the sequence $(U_{\theta}^{\varepsilon})_{\theta < \eta}$ is decreasing, $\mathbb{S}_{\varepsilon}^{\eta} = U_{0}^{\varepsilon}$,

$$\overline{U_{\theta}^{0} \cup U_{\theta}^{1}} = C_{\theta}^{0} \cup C_{\theta}^{1} = U_{\theta}^{0} \cup U_{\theta}^{1} \cup \Delta(2^{\omega}) = C_{\theta},$$

and $C^0_{\theta+1} \cup C^1_{\theta+1} = \overline{U^0_{\theta}} \cap \overline{U^1_{\theta}}$ if $\theta < \eta$ since

$$\overline{U_{\theta}^{\varepsilon}} = C_{\theta+1}^{0} \cup C_{\theta+1}^{1} \cup \bigcup_{\varphi(s)=\theta, \text{ parity}(s(0))=|1-||\text{parity}(\theta)-\text{parity}(\eta)|-\varepsilon||} \text{Gr}(f_{s})$$

as in the claim. Let us prove that $U_{\theta}^{0} \cup U_{\theta}^{1} \subseteq (f \times g)^{-1}(C_{\theta})$ if $\theta < \eta$. We argue by induction on θ , and the result is clear for $\theta = 0$. If $\theta = \theta' + 1$ is a successor ordinal, then

$$U^0_{\theta} \cup U^1_{\theta} \subseteq C^0_{\theta} \cup C^1_{\theta} = \overline{U^0_{\theta'}} \cap \overline{U^1_{\theta'}} \subseteq (f \times g)^{-1} (\overline{\mathbb{N}^\eta_0 \cap C_{\theta'}} \cap \overline{\mathbb{N}^\eta_1 \cap C_{\theta'}}) \subseteq (f \times g)^{-1} (C_{\theta}).$$

If θ is a limit ordinal, then $U^0_{\theta} \cup U^1_{\theta} \subseteq \bigcap_{\theta' < \theta} (U^0_{\theta'} \cup U^1_{\theta'}) \subseteq (f \times g)^{-1}(\bigcap_{\theta' < \theta} C_{\theta'}) = (f \times g)^{-1}(C_{\theta})$. This implies that $C^0_{\eta} \cup C^1_{\eta} \subseteq (f \times g)^{-1}(C_{\eta})$. In particular, $\Delta(2^{\omega})$ is sent into itself by $f \times g$ and f = g. As $\eta = \theta + 1$ is a successor ordinal, $U^0_{\theta} \subseteq (f \times f)^{-1}(\mathbb{N}^{\eta}_0 \cap C_{\theta}) \subseteq (f \times f)^{-1}(\Delta(2^{\omega}))$, which contradicts the injectivity of f. • So we proved that $\mathcal{A} := \{ (\mathbb{N}_1^{\eta}, \mathbb{N}_0^{\eta}), (\mathbb{N}_0^{\eta}, \mathbb{N}_1^{\eta}), (\mathbb{S}_0^{\eta}, \mathbb{S}_1^{\eta}) \}$ is a \sqsubseteq -antichain if η is a successor ordinal. For the same reasons, $\mathcal{B} := \{ (\mathbb{B}_1^{\eta}, \mathbb{B}_0^{\eta}), (\mathbb{B}_0^{\eta}, \mathbb{B}_1^{\eta}), (\mathbb{C}_0^{\eta}, \mathbb{C}_1^{\eta}) \}$ is a \sqsubseteq -antichain if η is a successor ordinal. Moreover, no pair in \mathcal{A} is below a pair in \mathcal{B} for the square reduction since $\Delta(2^{\omega}) \subseteq \mathbb{N}_0^{\eta} \cap \overline{\mathbb{S}_0^{\eta} \cup \mathbb{S}_1^{\eta}}$ and the element of the pairs in \mathcal{B} are contained in the clopen set $N_0 \times N_1$.

It remains to prove that we cannot find $(\mathbb{A}, \mathbb{B}), (\mathbb{A}', \mathbb{B}') \in \mathcal{A}$ and a continuous injection $f: 2^{\omega} \to 2^{\omega}$ such that $G_{\mathbb{A}} \subseteq (f \times f)^{-1}(\mathbb{A}')$ and $G_{\mathbb{B}} \subseteq (f \times f)^{-1}(\mathbb{B}')$. We argue by contradiction. If $(\mathbb{A}, \mathbb{B}) \neq (\mathbb{A}', \mathbb{B}')$ and $\varepsilon \in 2$, then we define continuous injections $f_{\varepsilon}: 2^{\omega} \to 2^{\omega}$ by $f_{\varepsilon}(\alpha) := f(\varepsilon \alpha)$. Note that $f_0 \times f_1$ reduces (\mathbb{A}, \mathbb{B}) to $(\mathbb{A}', \mathbb{B}')$, which contradicts the fact that \mathcal{A} is a \sqsubseteq -antichain. Thus $(\mathbb{A}, \mathbb{B}) = (\mathbb{A}', \mathbb{B}')$, and $(\mathbb{A}, \mathbb{B}) = (\mathbb{S}_0^{\eta}, \mathbb{S}_1^{\eta})$ by Proposition 3.4. As in the proof of Proposition 3.4, $\varphi(s) \leq \varphi(v)$. If $\alpha \in 2^{\omega}$, then $(0\alpha, 1\alpha)$ is the limit of $(0t_{p_k}^0 \gamma_k, 1t_{p_k}^1 \gamma_k)$. Note that $(f(0t_{p_k}^0 \gamma_k), f(1t_{p_k}^1 \gamma_k)) = (t_{v_k}^0 \gamma'_k, t_{v_k}^1 \gamma'_k)$ and $\varphi(p_k) \leq \varphi(v_k)$. As $(\varphi(p_k))_{k \in \omega}$ is cofinal in $\varphi(\emptyset) = \eta$, so is $(\varphi(v_k))_{k \in \omega}$. This implies that $(f(0\alpha), f(1\alpha)) \in \Delta(2^{\omega})$, which contradicts the injectivity of f.

(b) Let us prove that $(2^{\omega}, 2^{\omega}, \mathbb{S}_0^{\eta}, \mathbb{S}_1^{\eta}) \sqsubseteq (2^{\omega}, 2^{\omega}, \mathbb{N}_{\varepsilon}^{\eta}, \mathbb{N}_{1-\varepsilon}^{\eta})$ with a square map if $\varepsilon \in 2$. Let us do it for $\varepsilon = 0$, the other case being similar. We construct a map $\phi : 2^{<\omega} \to 2^{<\omega}$ satisfying the following:

$$\begin{array}{l} (1) \forall l \in \omega \ \exists k_l \in \omega \ \phi[2^l] \subseteq 2^{k_l} \\ (2) \ \phi(s) \subsetneqq \phi(s\varepsilon) \\ (3) \ \phi(s0) \neq \phi(s1) \\ (4) \ \forall s \in T_\eta \setminus \{\emptyset\} \ \left(\mathsf{parity}(|s|) = 1 - \left| \mathsf{parity}(s(0)) - \varepsilon \right| \right) \Rightarrow \exists v_s \in T_\eta \ \mathsf{parity}(|v_s|) = \varepsilon \land \\ (a) \ \forall w \in 2^{<\omega} \ \exists w' \in 2^{<\omega} \ \left(\phi(t_s^0 w), \phi(t_s^1 w) \right) = (t_{v_s}^0 w', t_{v_s}^1 w') \\ (b) \ \varphi(s) \leq \varphi(v_s) \end{array}$$

Assume that this is done. Then the map $f: \alpha \mapsto \lim_{n \to \infty} \phi(\alpha|n)$ is as desired. So let us check that the construction of ϕ is possible. We construct $\phi(s)$ by induction on the length of s.

- We set
$$k_0 := 0$$
 and $\phi(\emptyset) := \emptyset$.

- Note that $\langle 0 \rangle_{\eta} = 1$ and $(t_0^0, t_0^1) = (0, 1)$. As $\eta \ge 1$ is limit, $\varphi(1) > \varphi(0)$ are odd ordinals, so that $\varphi(10) \ge \varphi(0)$ is an even ordinal. We set $k_1 := \langle 10 \rangle_{\eta}$, $\phi(\varepsilon) := t_{10}^{\varepsilon}$ and $v_0 := 10$. This completes the construction of $\phi[2^1]$, and our conditions are satisfied since $k_1 > 0$.

- We next want to construct $\phi(s)$ for $s \in 2^{l+1}$, with $l \ge 1$, assuming that we have constructed $\phi(s)$ if $|s| \le l$. Note that there is exactly one sequence u such that $(t_u^0, t_u^1) \in 2^{l+1}$. We first define simultaneously $\phi(t_u^0)$ and $\phi(t_u^1)$, and then extend the definition to the other sequences in 2^{l+1} .

If $|u| \geq 2$, then there are $u_0 \in \omega^{<\omega}$ and $w \in 2^{<\omega}$ such that $t_u^{\varepsilon} = t_{u_0}^{\varepsilon} w\varepsilon$. By condition (4), $(\phi(t_{u_0}^0 w), \phi(t_{u_0}^0 w)) = (t_v^0 w', t_v^1 w')$ for some $v \in \omega^{<\omega}$ and $w' \in 2^{<\omega}$. Let $q \in \omega$ such that $w' \subseteq \psi(q)$ and $\varphi(u) \leq \varphi(vq)$. We can find such a q because if $\varphi(v) = \nu + 1$, then $\varphi(vq) = \nu$, but $\varphi(u) < \varphi(u_0) \leq \nu + 1$ so that $\varphi(u) \leq \nu$. If $\varphi(v)$ is limit, then $(\varphi(vq))_{q \in \omega}$ is cofinal in $\varphi(v)$ and $\varphi(u) < \varphi(u_0) \leq \varphi(v)$. We set $\phi(t_{u_0}^{\varepsilon} w\varepsilon) := t_{vq}^{\varepsilon}$. By definition, there is $N \in \omega$ such that $t_{vq}^{\varepsilon} = t_v^{\varepsilon} w' 0^N \varepsilon$. We set $\phi(s\varepsilon) := \phi(s) 0^N \varepsilon$, for any $s \in 2^l$. Conditions (1)-(3) clearly hold. So let us check condition (4). First note that $(\phi(t_u^0), \phi(t_u^1)) = (t_{vq}^0, t_{vq}^1)$ by definition, so that (4) holds for u since $|u| - |u_0| = |vq| - |v| = 1$.

Suppose now that there are $u_1 \in \omega^{<\omega}$, $z \in 2^{<\omega}$ and $e \in 2$ such that $(s,t) = (t_{u_1}^0 ze, t_{u_1}^1 ze)$. By the induction hypothesis, $(\phi(t_{u_1}^0 ze), \phi(t_{u_1}^1 ze)) = (\phi(t_{u_1}^0 z)0^N e, \phi(t_{u_1}^1 z)0^N e) = (t_{v_{u_1}}^0 z'0^N e, t_{v_{u_1}}^1 z'0^N e)$. Thus conditions (4) is checked.

Otherwise, |u| = 1 and $u = \langle p \rangle$ for some $p \in \omega \setminus \{0\}$. Let $w := t_u^0 | l$. Note there are infinitely many q's such that $\phi(w) \subseteq \psi(q)$. As η is a limit ordinal, $(\varphi(q))_{q \in \omega}$ is strictly increasing. Thus q can be chosen so that $\varphi(p) \leq \varphi(q)$. If p is odd, then we set $\phi(t_u^{\varepsilon}) := t_{<q>}^{\varepsilon}$. If p is even, then we set $\phi(t_u^{\varepsilon}) := t_{q0}^{\varepsilon}$. Let w^0 and w^1 be the sequences such that $\phi(t_u^{\varepsilon}) = \phi(w)w^{\varepsilon}\varepsilon$. Note that they are different if p is even. As in the previous case, we define $\phi(s\varepsilon) := \phi(s)w^{\varepsilon}\varepsilon$, for any $s \in 2^l$. Notice how the choice of w^{ε} only depends on the last coordinate of $s\varepsilon$. The conditions are verified as before for $(\phi(t_u^0), \phi(t_u^1))$. For the other cases,

 $\left(\phi(t_{u_1}^0 z e), \phi(t_{u_1}^1 z e)\right) = \left(\phi(t_{u_1}^0 z) w^e e, \phi(t_{u_1}^1 z) w^e e\right) = (t_{v_{u_1}}^0 w' w^e e, t_{v_{u_1}}^1 w' w^e e),$

by the induction hypothesis. So the conditions are checked.

It remains to note that $(2^{\omega}, 2^{\omega}, \mathbb{C}^{\eta}_0, \mathbb{C}^{\eta}_1) \sqsubseteq (2^{\omega}, 2^{\omega}, \mathbb{B}^{\eta}_{\varepsilon}, \mathbb{B}^{\eta}_{1-\varepsilon})$ with a square map if $\varepsilon \in 2$, with witness $\varepsilon \alpha \mapsto \varepsilon f(\alpha)$.

Consequences

Corollary 4.5 Let $\eta \ge 1$ be a countable ordinal, X be a Polish space, and A, B be disjoint analytic relations on X such that $A \cup B$ is contained in a potentially closed s-acyclic or locally countable relation. Then exactly one of the following holds:

(a) the set A is separable from B by a pot $\left(\Delta\left(D_{\eta}(\Sigma_{1}^{0})\right)\right)$ set,

(b) there is $(\mathbb{A}_0, \mathbb{A}_1) \in \{(\mathbb{N}_1^{\eta}, \mathbb{N}_0^{\eta}), (\mathbb{N}_0^{\eta}, \mathbb{N}_1^{\eta}), (\mathbb{S}_0^{\eta}, \mathbb{S}_1^{\eta})\}$ with $(2^{\omega}, 2^{\omega}, \mathbb{A}_0, \mathbb{A}_1) \sqsubseteq (X, X, A, B)$.

Proof. By Lemmas 3.1 and 4.1, (a) and (b) cannot hold simultaneously. So assume that (a) does not hold. By Lemma 2.2, we may assume that $A \cup B$ is contained in a potentially closed quasi-acyclic relation. It remains to apply Theorem 4.3.

Corollary 4.6 Let $\eta \ge 1$ be a countable ordinal, X, Y be Polish spaces, and A, B be disjoint analytic subsets of $X \times Y$ such that $A \cup B$ is contained in a potentially closed locally countable set. Then exactly one of the following holds:

- (a) the set A is separable from B by a pot $\left(\Delta\left(D_{\eta}(\Sigma_{1}^{0})\right)\right)$ set,
- (b) there is $(\mathbb{A}_0, \mathbb{A}_1) \in \{(\mathbb{N}_1^\eta, \mathbb{N}_0^\eta), (\mathbb{N}_0^\eta, \mathbb{N}_1^\eta), (\mathbb{S}_0^\eta, \mathbb{S}_1^\eta)\}$ with $(2^\omega, 2^\omega, \mathbb{A}_0, \mathbb{A}_1) \sqsubseteq (X, X, A, B)$.

Proof. As in the proof of Corollary 4.5, (a) and (b) cannot hold simultaneously. Then we argue as in the proof of Corollary 3.7. $A' \cup B'$ is contained in a potentially closed locally countable relation, and A' is not separable from B' by a pot $\left(\Delta\left(D_{\eta}(\Sigma_{1}^{0})\right)\right)$ set. Corollary 4.5 gives $f', g': 2^{\omega} \to Z$.

Corollary 4.7 Let $\eta \ge 1$ be a countable ordinal, X be a Polish space, and A, B be disjoint analytic relations on X. The following are equivalent:

(1) there is a potentially closed s-acyclic relation $R \in \Sigma_1^1$ such that $A \cap R$ is not separable from $B \cap R$ by a pot $\left(\Delta\left(D_\eta(\Sigma_1^0)\right)\right)$ set,

(2) there is a potentially closed locally countable relation $R \in \Sigma_1^1$ such that $A \cap R$ is not separable from $B \cap R$ by a pot $\left(\Delta\left(D_\eta(\Sigma_1^0)\right)\right)$ set,

(3) there is $(\mathbb{A}_0, \mathbb{A}_1) \in \{(\mathbb{N}_1^{\eta}, \mathbb{N}_0^{\eta}), (\mathbb{N}_0^{\eta}, \mathbb{N}_1^{\eta}), (\mathbb{S}_0^{\eta}, \mathbb{S}_1^{\eta})\}$ with $(2^{\omega}, 2^{\omega}, \mathbb{A}_0, \mathbb{A}_1) \sqsubseteq (X, X, A, B)$,

(4) there is $(\mathbb{A}_0, \mathbb{A}_1) \in \{(\mathbb{N}_1^{\eta}, \mathbb{N}_0^{\eta}), (\mathbb{B}_1^{\eta}, \mathbb{B}_0^{\eta}), (\mathbb{N}_0^{\eta}, \mathbb{N}_1^{\eta}), (\mathbb{B}_0^{\eta}, \mathbb{B}_1^{\eta}), (\mathbb{S}_0^{\eta}, \mathbb{S}_1^{\eta}), (\mathbb{C}_0^{\eta}, \mathbb{C}_1^{\eta})\}$ such that the inequality $(2^{\omega}, 2^{\omega}, \mathbb{A}_0, \mathbb{A}_1) \sqsubseteq (X, X, A, B)$ holds, via a square map.

Proof. (1) \Rightarrow (3),(4) and (2) \Rightarrow (3),(4) This is a consequence of Corollary 4.5 and its proof.

(4) \Rightarrow (1) By the remarks before Lemma 3.1, $\mathbb{N}_0^{\eta} \cup \mathbb{N}_1^{\eta}$ has s-acyclic levels. This implies that $\mathbb{N}_0^{\eta} \cup \mathbb{N}_1^{\eta}$ and $\mathbb{S}_0^{\eta} \cup \mathbb{S}_1^{\eta}$ are s-acyclic. As $\mathbb{N}_0^{\eta} \cup \mathbb{N}_1^{\eta}$ is antisymmetric, $\mathbb{B}_0^{\eta} \cup \mathbb{B}_1^{\eta}$ and $\mathbb{C}_0^{\eta} \cup \mathbb{C}_1^{\eta}$ are s-acyclic too, by Lemma 3.8. Thus we can take $R := (f \times f)[\mathbb{A}_0 \cup \mathbb{A}_1]$ since the s-acyclicity is preserved by images by the square of an injection, and by Lemmas 3.1 and 4.1.

(3),(4) \Rightarrow (2) We can take $R := (f \times f)[\mathbb{A}_0 \cup \mathbb{A}_1]$ since $\mathbb{A}_0 \cup \mathbb{A}_1$ is locally countable, by Lemmas 3.1 and 4.1.

5 Background

We now give some material to prepare the study of the Borel classes of rank two.

Potential Wadge classes

In Theorem 1.3, $\mathbb{S}_0 \cup \mathbb{S}_1$ is a subset of the body of a tree T on 2^2 which does not depend on Γ . We first describe a simple version of T, which is sufficient to study the Borel classes (see [L6]). We identify $(2^l)^2$ and $(2^2)^l$, for each $l \in \omega + 1$.

Definition 5.1 (1) Let $\mathcal{F} \subseteq \bigcup_{l \in \omega} (2^l)^2 \equiv (2^2)^{<\omega}$. We say that \mathcal{F} is a frame if

 $\begin{array}{l} (a) \; \forall l \in \omega \; \exists ! (s_l, t_l) \in \mathcal{F} \cap (2^l)^2, \\ (b) \; \forall p, q \in \omega \; \forall w \in 2^{<\omega} \; \exists N \in \omega \; (s_q 0 w 0^N, t_q 1 w 0^N) \in \mathcal{F} \; and \; (|s_q 0 w 0^N| - 1)_0 = p, \\ (c) \; \forall l > 0 \; \exists q < l \; \exists w \in 2^{<\omega} \; (s_l, t_l) = (s_q 0 w, t_q 1 w). \end{array}$

(2) If $\mathcal{F} = \{(s_l, t_l) \mid l \in \omega\}$ is a frame, then we will call T the tree on 2^2 generated by \mathcal{F} :

$$T := \left\{ (s,t) \in (2^2)^{<\omega} \mid s = \emptyset \lor \left(\exists q \in \omega \ \exists w \in 2^{<\omega} \ (s,t) = (s_q 0 w, t_q 1 w) \right) \right\}.$$

The existence condition in (a) and the density condition (b) ensure that $\lceil T \rceil$ is big enough to contain sets of arbitrary high potential complexity. The uniqueness condition in (a) and condition (c) ensure that $\lceil T \rceil$ is small enough to make the reduction in Theorem 1.3 possible. The last part of condition (b) gives a control on the verticals which is very useful to construct complicated examples.

In the sequel, T will be the tree generated by a fixed frame \mathcal{F} (Lemma 3.3 in [L6] ensures the existence of concrete frames). Note that $\lceil T \rceil \subseteq N_0 \times N_1$, which will be useful in the sequel (recall that N_s is the basic clopen set of sequences beginning with $s \in 2^{<\omega}$).

Acyclicity

We will use some material from [L6] and [L8], where some possibly different notions of acyclicity of the levels of T are involved. We will check that they coincide in our case.

Definition 5.2 Let X be a set, and A be a relation on X.

(a) An A-path is a finite sequence $(x_i)_{i \leq n}$ of points of X such that $(x_i, x_{i+1}) \in A$ if i < n.

(b) We say that A is connected if for any $x, y \in X$ there is an A-path $(x_i)_{i \leq n}$ with $x_0 = x$ and $x_n = y$.

(c) An A-cycle is an A-path $(x_i)_{i \leq L}$ with $L \geq 3$, $(x_i)_{i < L}$ is injective and $x_L = x_0$ (so that A is acyclic if and only if there is no A-cycle).

Lemma 5.3 Let $l \in \omega$, and $T_l := T \cap (2^l)^2$ be the lth level of T.

(a) $s(T_l)$ is connected and acyclic. In particular, [T] is s-acyclic.

(b) A tree S on 2^2 has acyclic levels in the sense of [L6] if and only if S has suitable levels in the sense of [L8], and this is the case of T.

Proof. (a) We argue by induction on l. The statement is clear for l = 0. For the inductive step we use the fact that $T_{l+1} = \{(s\varepsilon, t\varepsilon) \mid (s, t) \in T_l \land \varepsilon \in 2\} \cup \{(s_l0, t_l1)\}$. As the map $s\varepsilon \mapsto s$ defines an isomorphism from $\{(s\varepsilon, t\varepsilon) \mid (s, t) \in T_l\}$ onto T_l , we are done. A cycle for $s(\lceil T \rceil)$ gives a cycle for $s(T_l)$, for l big enough to ensure the injectivity of the initial segments.

(b) Assume that S has acyclic levels in the sense of [L6]. This means that, for each l, the graph G_{S_l} with set of vertices $2^l \oplus 2^l$ (with typical element $\overline{x_{\varepsilon}} := (x_{\varepsilon}, \varepsilon) \in 2^l \times 2$) and set of edges

$$\left\{\left\{\overline{x_0}, \overline{x_1}\right\} \mid \vec{x} := (x_0, x_1) \in S_l\right\}$$

is acyclic. We have to see that S has suitable levels in the sense of [L8]. This means that, for each l, the following hold:

- S_l is finite,
- $-\exists \varepsilon \in 2 \ x_{\varepsilon}^{0} \neq x_{\varepsilon}^{1} \text{ if } \vec{x^{0}} \neq \vec{x^{1}} \in S_{l},$

- consider the graph G^{S_l} with set of vertices S_l and set of edges

$$\{\{\vec{x^0}, \vec{x^1}\} \mid \vec{x^0} \neq \vec{x^1} \land \exists \varepsilon \in 2 \ x^0_\varepsilon = x^1_\varepsilon\};\$$

then for any G^{S_l} -cycle $(\vec{x^n})_{n \leq L}$, there are $\varepsilon \in 2$ and k < m < n < L such that $x_{\varepsilon}^k = x_{\varepsilon}^m = x_{\varepsilon}^n$.

The first two properties are obvious. So assume that $(\vec{x^n})_{n \leq L}$ is a G^{S_l} -cycle for which we cannot find $\varepsilon \in 2$ and k < m < n < L such that $x_{\varepsilon}^k = x_{\varepsilon}^m = x_{\varepsilon}^n$.

Case 1 $x_0^0 = x_0^1$.

Subcase 1.1 L is odd.

Note that $L \ge 5$. Indeed, $L \ge 3$ since $(\vec{x^n})_{n \le L}$ is a G^{S_l} -cycle. So we just have to see that $L \ne 3$. As $x_0^0 = x_0^1$ and $\vec{x^0} \ne \vec{x^1}$, $x_1^0 \ne x_1^1$. By the choice of $(\vec{x^n})_{n \le L}$, $x_0^1 \ne x_0^2$. Thus $x_1^1 = x_1^2$. By the choice of $(\vec{x^n})_{n \le L}$, $x_1^2 \ne x_1^3$. Thus $x_0^2 = x_0^3$ and $x_0^3 \ne x_0^0$. Therefore $\vec{x^3} \ne \vec{x^0}$ and $L \ne 3$.

Then $\overline{x_0^0}, \overline{x_1^1}, \overline{x_0^2}, ..., \overline{x_1^{L-2}}, \overline{x_0^{L-1}}$ is a G_{S_l} -cycle, by the choice of $(\vec{x^n})_{n \leq L}$.

Subcase 1.2 L is even, in which case $L \ge 4$.

Then $\overline{x_0^0}, \overline{x_1^1}, \overline{x_0^2}, ..., \overline{x_1^{L-1}}, \overline{x_0^L}$ is a G_{S_l} -cycle, by the choice of $(\vec{x^n})_{n \leq L}$.

Case 2 $x_0^0 \neq x_0^1$.

The same arguments work, we just have to exchange the indexes.

• Conversely, assume that $(\overline{x_{\varepsilon_n}^n})_{n \leq L}$ is a G_{S_l} -cycle. Then L is even, and actually $L \geq 4$.

Case 1 $\varepsilon_0 = 0$.

Then $(x_{\varepsilon_0}^0, x_{\varepsilon_1}^1), (x_{\varepsilon_2}^2, x_{\varepsilon_1}^1), ..., (x_{\varepsilon_{L-2}}^{L-2}, x_{\varepsilon_{L-1}}^{L-1}), (x_{\varepsilon_L}^L, x_{\varepsilon_{L-1}}^{L-1}), (x_{\varepsilon_0}^0, x_{\varepsilon_1}^1)$ is a G^{S_l} -cycle of length L+1. If $\varepsilon \in 2$, then each ε th coordinate appears exactly twice before the last element of the cycle.

Case 2 $\varepsilon_0 = 1$.

The same argument works, we just have to exchange the coordinates.

• By Proposition 3.2 in [L6], T has acyclic levels in the sense of [L6].

6 The classes Π^0_2 and Σ^0_2

Example

We will use an example for $\Gamma = \Pi_2^0$ different from that in [L6], so that we prove the following.

Lemma 6.1 $[T] \cap \mathbb{E}_0$ is not separable from $[T] \setminus \mathbb{E}_0$ by a pot (Π_2^0) set.

Proof. We argue by contradiction, which gives $P \in \text{pot}(\Pi_2^0)$, and also a dense G_δ subset G of 2^ω such that $P \cap G^2 \in \Pi_2^0(G^2)$. Let $(O_n)_{n \in \omega}$ be a sequence of dense open subsets of 2^ω with intersection G. Note that $[T] \cap \mathbb{E}_0 \cap G^2 = [T] \cap P \cap G^2 \in \Delta_2^0([T] \cap G^2)$. By Baire's theorem, it is enough to prove that $[T] \cap \mathbb{E}_0 \cap G^2$ is dense and co-dense in the nonempty space $[T] \cap G^2$. So let $q \in \omega$ and $w \in 2^{<\omega}$. Pick $u_0 \in 2^\omega$ such that $N_{s_q 0 w u_0} \subseteq O_0$, $v_0 \in 2^\omega$ such that $N_{t_q 1 w u_0 v_0} \subseteq O_0$, $u_1 \in 2^\omega$ such that $N_{s_q 0 w u_0 v_0 u_1} \subseteq O_1$, $v_1 \in 2^\omega$ such that $N_{t_q 1 w u_0 v_0 u_1} \subseteq O_1$, and so on.

Then $(s_q 0wu_0v_0u_1v_1..., t_q 1wu_0v_0u_1v_1...) \in [T] \cap \mathbb{E}_0 \cap G^2$. Similarly, pick $N_0 \in \omega$ such that $(s_q 0w0^{N_0}, t_q 1w0^{N_0}) \in \mathcal{F}, u_0 \in 2^{\omega}$ such that $N_{s_q 0w0^{N_0} 0u_0} \subseteq O_0, v_0 \in 2^{\omega}$ such that $N_{t_q 1w0^{N_0} 1u_0v_0} \subseteq O_0, N_1 \in \omega$ such that $(s_q 0w0^{N_0} 0u_0v_00^{N_1}, t_q 1w0^{N_0} 1u_0v_00^{N_1}) \in \mathcal{F}, u_1 \in 2^{\omega}$ such that

 $N_{s_q 0 w 0^{N_0} 0 u_0 v_0 0^{N_1} 0 u_1} \subseteq O_1,$

 $v_1 \in 2^{\omega}$ such that $N_{t_a 1 w 0^{N_0} 1 u_0 v_0 0^{N_1} 1 u_1 v_1} \subseteq O_1$, and so on. Then

$$(s_q 0w 0^{N_0} 0u_0 v_0 0^{N_1} 0u_1 v_1 \dots, t_q 1w 0^{N_0} 1u_0 v_0 0^{N_1} 1u_1 v_1 \dots) \in [T] \cap G^2 \setminus \mathbb{E}_0.$$

This finishes the proof.

The main result

We reduce the study of disjoint analytic sets to that of disjoint Borel sets of low complexity, for the first classes we are considering.

Lemma 6.2 Let X be a Polish space, and A, B be disjoint analytic relations on X. Then exactly one of the following holds:

- (a) the set A is separable from B by a pot(Π_2^0) set,
- (b) there is a K_{σ} relation $A' \subseteq A$ which is not $pot(\mathbf{\Pi}_2^0)$ such that $\overline{A'} \setminus A' \subseteq B$.

Proof. Theorem 1.10 in [L8] and Lemmas 6.1, 5.3 give $g, h : 2^{\omega} \to X$ continuous such that the inclusions $[T] \cap \mathbb{E}_0 \subseteq (g \times h)^{-1}(A)$ and $[T] \setminus \mathbb{E}_0 \subseteq (g \times h)^{-1}(B)$ hold. We set $A' := (g \times h) [[T] \cap \mathbb{E}_0]$, $B' := (g \times h) [[T] \setminus \mathbb{E}_0]$ and $C' := (g \times h) [[T]]$. Note that A' is a K_{σ} subset of $A, B' \subseteq B$, so that the compact set C' is the disjoint union of A' and B'. As $[T] \cap \mathbb{E}_0$ is dense in [T], C' is also the closure of A'. As $[T] \cap \mathbb{E}_0 = [T] \cap (g \times h)^{-1}(A')$, A' is not pot (Π_2^0) , by Lemma 6.1.

Theorem 6.3 Let X be a Polish space, and A, B be disjoint analytic relations on X such that A is quasi-acyclic. Then one of the following holds:

(a) the set A is separable from B by a pot(Π_2^0) set,

(b) there is $f: 2^{\omega} \to X$ injective continuous such that the inclusions $\lceil T \rceil \cap \mathbb{E}_0 \subseteq (f \times f)^{-1}(A)$ and $\lceil T \rceil \setminus \mathbb{E}_0 \subseteq (f \times f)^{-1}(B)$ hold.

Proof. Assume that (a) does not hold. By Lemma 6.2, we may assume that B is the complement of A. Let $(C_n)_{n \in \omega}$ be a witness for the fact that A is quasi-acyclic. Note that there are disjoint Borel subsets O_0, O_1 of X such that $A \cap (O_0 \times O_1)$ is not pot (Π_2^0) . We may assume that X is zero-dimensional, the C_n 's are closed, and O_0, O_1 are clopen, refining the topology if necessary. We can also replace A and the C_n 's with their intersection with $O_0 \times O_1$ and assume that they are contained in $O_0 \times O_1$.

• We may assume that X is recursively presented, $O_0, O_1 \in \Delta_1^1$ and the relation " $(x, y) \in C_n$ " is Δ_1^1 in (x, y, n). As Δ_X is Polish finer than the topology on $X, A \notin \Pi_2^0(X^2, \tau_1)$. We now perform the following derative on A. We set, for $F \in \Pi_1^0(X^2, \tau_1), F' := \overline{F \cap A}^{\tau_1} \cap \overline{F \setminus A}^{\tau_1}$ (see 22.30 in [K]).

Then we inductively define, for any ordinal ξ , F_{ξ} by

$$\begin{cases} F_0 := X^2 \\ F_{\xi+1} := F'_{\xi} \\ F_{\lambda} := \bigcap_{\xi < \lambda} F_{\xi} \text{ if } \lambda \text{ is limit} \end{cases}$$

(see 22.27 in [K]). As (F_{ξ}) is a decreasing sequence of closed subsets of the Polish space (X^2, τ_1) , there is $\theta < \omega_1$ such that $F_{\theta} = F_{\theta+1}$. In particular, $F_{\theta} = F_{\theta+1} = F'_{\theta} = \overline{F_{\theta} \cap A}^{\tau_1} \cap \overline{F_{\theta} \setminus A}^{\tau_1}$, so that $F_{\theta} \cap A$ and $F_{\theta} \setminus A$ are τ_1 -dense in F_{θ} .

• Let us prove that F_{θ} is not empty. We argue by contradiction:

$$X^2 = \neg F_{\theta} = \bigcup_{\xi \le \theta} \neg F_{\xi} = \bigcup_{\xi \le \theta} (\neg F_{\xi} \cap \bigcap_{\eta < \xi} F_{\eta}) = \bigcup_{\xi < \theta} F_{\xi} \setminus F_{\xi+1},$$

so that $A = \bigcup_{\xi < \theta} A \cap F_{\xi} \setminus F_{\xi+1}$. But $A \cap F_{\xi} \setminus F_{\xi+1} = A \cap F_{\xi} \setminus (\overline{F_{\xi} \cap A^{\tau_1}} \cap \overline{F_{\xi} \setminus A^{\tau_1}}) = F_{\xi} \setminus \overline{F_{\xi} \setminus A^{\tau_1}}$. This means that $(F_{\xi} \setminus F_{\xi+1})_{\xi < \theta}$ is a countable partition of (X^2, τ_1) into Δ_2^0 sets, and that A is Δ_2^0 on each piece of the partition. This implies that A is $\Delta_2^0(X^2, \tau_1)$, which is absurd.

• Let us prove that F_{θ} is Σ_1^1 . We use 7C in [Mo]. We define a set relation by

$$\varphi(x, y, P) \Leftrightarrow (x, y) \notin (\neg P)'.$$

Note that φ is monotone, and thus operative. It is also Π_1^1 on Π_1^1 . By 3E.2, 3F.6 and 4B.2 in [Mo], we can apply 7C.8 in [Mo], so that $\varphi^{\infty}(x, y)$ is Π_1^1 . An induction shows that $\varphi^{\xi}(x, y)$ is equivalent to " $(x, y) \notin F_{\xi+1}$ ". Thus $(x, y) \notin F_{\theta}$ is equivalent to $(x, y) \notin \bigcap_{\xi} F_{\xi} = \bigcap_{\xi} F_{\xi+1}, (x, y) \in \bigcup_{\xi} \neg F_{\xi+1}$ and $\varphi^{\infty}(x, y)$.

• We are ready to prove the following key property:

$$\forall q \in \omega \ \forall U, V \in \Sigma_1^1(X) \ F_\theta \cap (U \times V) \neq \emptyset \Rightarrow \exists n \ge q \ F_\theta \cap C_n \cap (U \times V) \neq \emptyset.$$

Indeed, this property says that $I := F_{\theta} \cap (\bigcup_{n \geq q} C_n)$ is Σ_X^2 -dense in F_{θ} for each $q \in \omega$. We fix $q \in \omega$, and prove first that I is τ_1 -dense in F_{θ} . So let $U, V \in \Delta_1^1$ such that $F_{\theta} \cap (U \times V)$ is nonempty. As $F_{\theta} \setminus A$ is τ_1 -dense in F_{θ} , we get $(x, y) \in (F_{\theta} \setminus A) \cap (U \times V)$. As $F_{\theta} \cap A$ is τ_1 -dense in F_{θ} , we get $(x_k, y_k) \in F_{\theta} \cap A$ converving to (x, y) for τ_1 . Pick $n_k \in \omega$ such that $(x_k, y_k) \in C_{n_k}$. As C_{n_k} is closed, and thus τ_1 -closed, we may assume that the sequence $(n_k)_{k \in \omega}$ is strictly increasing. Now $(x_k, y_k) \in I \cap (U \times V)$ if k is big enough. In order to get the statement for Σ_X^2 , we have to note that I is Σ_1^1 since F_{θ} is Σ_1^1 and the relation " $(x, y) \in C_n$ " is Δ_1^1 in (x, y, n). This implies that $\overline{I}^{\tau_1} = \overline{I}^{\Sigma_X^2}$, by a double application of the separation theorem. Therefore $F_{\theta} \subseteq \overline{I}^{\tau_1} = \overline{I}^{\Sigma_X^2}$ and I is Σ_X^2 -dense in F_{θ} .

• We set, for $\vec{u} = (u_0, u_1) \in T \setminus \{\vec{\emptyset}\},\$

$$n(\vec{u}) := \operatorname{Card}(\{i < |\vec{u}| \mid u_0(i) \neq u_1(i)\}),\\ \vec{t}(\vec{u}) := (s_q 0, t_q 1) \text{ if } \vec{u} = (s_q 0 w, t_q 1 w).$$

- We are ready for the construction of f. We construct the following objects:
 - sequences $(x_s)_{s \in 2^{<\omega} \setminus \{\emptyset\}, s(0)=0}, (y_s)_{s \in 2^{<\omega} \setminus \{\emptyset\}, s(0)=1}$ of points of X,
 - sequences $(X_s)_{s \in 2^{<\omega} \setminus \{\emptyset\}, s(0)=0}$, $(Y_s)_{s \in 2^{<\omega} \setminus \{\emptyset\}, s(0)=1}$ of Σ_1^1 subsets of X, - a map $\Phi: \{\vec{t}(\vec{u}) \mid \vec{u} \in T \setminus \{\vec{\emptyset}\}\} \rightarrow \omega$.

We want these objects to satisfy the following conditions:

 $\begin{array}{l} (1) \ x_s \in X_s \ \land \ y_s \in Y_s \\ (2) \ X_{s\varepsilon} \subseteq X_s \subseteq \Omega_X \cap O_0 \ \land \ Y_{s\varepsilon} \subseteq Y_s \subseteq \Omega_X \cap O_1 \\ (3) \ \dim_{\operatorname{GH}}(X_s), \ \dim_{\operatorname{GH}}(Y_s) \le 2^{-|s|} \\ (4) \ (x_{u_0}, y_{u_1}) \in F_{\theta} \cap C_{\Phi(\vec{t}(\vec{u}))} \\ (5) \ (X_{u_0} \times Y_{u_1}) \cap (\bigcup_{n < n(\vec{u})} C_n) = \emptyset \\ (6) \ X_{s0} \cap X_{s1} = Y_{s0} \cap Y_{s1} = \emptyset \end{array}$

• Assume that this has been done. As in the proof of Theorem 3.3, we get $f: N_{\varepsilon} \to O_{\varepsilon}$ injective continuous, so that $f: 2^{\omega} \to X$ is injective continuous. If $(\alpha, \beta) \in [T] \cap \mathbb{E}_0$, then $\Phi(\vec{t}((\alpha, \beta)|n)) = N$ if n is big enough. In this case, by (4), $(x_{\alpha|n}, y_{\beta|n}) \in C_N$ which is closed, so that $(f(\alpha), g(\beta)) \in C_N \subseteq A$. If $(\alpha, \beta) \in [T] \setminus \mathbb{E}_0$, then the sequence $(n((\alpha, \beta)|n))_{n>0}$ tends to infinity. Thus $(f(\alpha), g(\beta))$ is not in $\bigcup_{n \in \omega} C_n = A$ by (5).

• So let us prove that the construction is possible. The key property gives $\Phi(0,1) \ge 1$ and (x_0, y_1) in $F_{\theta} \cap C_{\Phi(0,1)} \cap \Omega_{X^2}$. As $\Omega_{X^2} \subseteq \Omega_X^2$, $x_0, y_1 \in \Omega_X$. We choose Σ_1^1 subsets X_0, Y_1 of X with GHdiameter at most 2^{-1} such that $(x_0, y_1) \in X_0 \times Y_1 \subseteq ((\Omega_X \cap O_0) \times (\Omega_X \cap O_1)) \setminus C_0$, which completes the construction for the length l = 1.

Let $l \ge 1$. We now want to build x_s, X_s, y_s, Y_s for $s \in 2^{l+1}$, as well as $\Phi(s_l 0, t_l 1)$. Note that $(x_{s_l}, y_{t_l}) \in F_{\theta} \cap (U \times V)$, where

$$\begin{split} U &:= \{ x_{s_l}' \in X_{s_l} \mid \exists (x_s')_{s \in 2^l \setminus \{s_l\}, s(0) = 0} \in \Pi_{s \in 2^l \setminus \{s_l\}, s(0) = 0} X_s \ \exists (y_s')_{s \in 2^l, s(0) = 1} \in \Pi_{s \in 2^l, s(0) = 1} Y_s \\ \forall \vec{u} \in T \cap (2^l \times 2^l) \ (x_{u_0}', y_{u_1}') \in F_\theta \cap C_{\Phi(\vec{t}(\vec{u}))} \}, \\ V &:= \{ y_{t_l}' \in Y_{t_l} \mid \exists (x_s')_{s \in 2^l, s(0) = 0} \in \Pi_{s \in 2^l, s(0) = 0} X_s \ \exists (y_s')_{s \in 2^l \setminus \{t_l\}, s(0) = 1} \in \Pi_{s \in 2^l \setminus \{t_l\}, s(0) = 1} Y_s \\ \forall \vec{u} \in T \cap (2^l \times 2^l) \ (x_{u_0}', y_{u_1}') \in F_\theta \cap C_{\Phi(\vec{t}(\vec{u}))} \}. \end{split}$$

The key property gives $\Phi(s_l0, t_l1) > \max(n(s_l0, t_l1), \max_{q < l} \Phi(s_q0, t_q1))$ and


$$(x_{s_l0}, y_{t_l1}) \in F_\theta \cap C_{\Phi(s_l0, t_l1)} \cap (U \times V).$$

The fact that $x_{s_l0} \in U$ gives witnesses $(x_{s0})_{s \in 2^l \setminus \{s_l\}, s(0)=0}$ and $(y_{s0})_{s \in 2^l, s(0)=1}$. Similarly, the fact that $y_{t_l1} \in V$ gives $(x_{s1})_{s \in 2^l, s(0)=0}$ and $(y_{s1})_{s \in 2^l \setminus \{t_l\}, s(0)=1}$. Note that $x_{s_l0} \neq x_{s_l1}$ because

$$(x_{s_l0}, y_{t_l1}) \in C_{\Phi(s_l0, t_l1)},$$

 $(x_{s_l1}, y_{t_l1}) \in C_{\Phi(\vec{t}(s_l1, t_l1))}$, and $\Phi(s_l0, t_l1) > \Phi(\vec{t}(s_l1, t_l1))$. Similarly, $y_{t_l0} \neq y_{t_l1}$. If $s \in 2^l$, then the connectedness of $s(T_l)$ gives an injective s(T)-path p_s from s to s_l . This gives a s(A)-path from x_{s0} to x_{s1} if s(0) = 0, and a s(A)-path from y_{s0} to y_{s1} if s(0) = 1. Using the quasi-acyclicity of A, we see, by induction on the length of p_s , that $x_{s0} \neq x_{s1}$ and $y_{s0} \neq y_{s1}$.

The following picture illustrates the situation when l = 2.



Then we take small enough Σ_1^1 neighborhoods of the $x_{s\varepsilon}$'s and $y_{s\varepsilon}$'s to complete the construction. \Box

Consequences

Corollary 6.4 Let X be a Polish space, and A, B be disjoint analytic relations on X such that A is either s-acyclic, or locally countable. Then exactly one of the following holds:

(a) the set A is separable from B by a pot(Π_2^0) set,

(b) there is $f: 2^{\omega} \to X$ injective continuous such that the inclusions $\lceil T \rceil \cap \mathbb{E}_0 \subseteq (f \times f)^{-1}(A)$ and $\lceil T \rceil \setminus \mathbb{E}_0 \subseteq (f \times f)^{-1}(B)$ hold.

Proof. By Lemma 6.1, $[T] \cap \mathbb{E}_0$ is not separable from $[T] \setminus \mathbb{E}_0$ by a $\text{pot}(\Pi_2^0)$ set. This shows that (a) and (b) cannot hold simultaneously. So assume that (a) does not hold. By Lemma 6.2, we may assume that *A* is Σ_2^0 and *B* is the complement of *A*. By Lemma 2.2, we may also assume that *A* is quasi-acyclic. It remains to apply Theorem 6.3.

Corollary 6.5 Let X, Y be Polish spaces, and A, B be disjoint analytic subsets of $X \times Y$ such that A is locally countable. Then exactly one of the following holds:

- (a) the set A is separable from B by a $pot(\mathbf{\Pi}_2^0)$ set,
- (b) $(2^{\omega}, 2^{\omega}, \lceil T \rceil \cap \mathbb{E}_0, \lceil T \rceil \setminus \mathbb{E}_0) \sqsubseteq (X, Y, A, B).$

Proof. As in the proof of Corollary 6.4, (a) and (b) cannot hold simultaneously. So assume that (a) does not hold. We argue as in the proof of Corollary 3.7. Corollary 6.4 gives $f': 2^{\omega} \to Z$.

Corollary 6.6 Let X be a Polish space, and A, B be disjoint analytic relations on X. The following are equivalent:

(1) there is an s-acyclic relation $R \in \Sigma_1^1$ such that $A \cap R$ is not separable from $B \cap R$ by a pot (Π_2^0) set,

(2) there is $f: 2^{\omega} \to X$ injective continuous with $[T] \cap \mathbb{E}_0 \subseteq (f \times f)^{-1}(A)$ and $[T] \setminus \mathbb{E}_0 \subseteq (f \times f)^{-1}(B)$.

Proof. (1) \Rightarrow (2) We apply Corollary 6.4.

(2) \Rightarrow (1) We can take $R := (f \times f) [[T]].$

Remark. There is a version of Corollary 6.6 for Σ_2^0 instead of Π_2^0 , obtained by exchanging the roles of *A* and *B*. This symmetry is not present in Theorem 6.3.

Corollary 6.7 Let X be a Polish space, and A, B be disjoint analytic relations on X such that A is contained in a $pot(F_{\sigma})$ s-acyclic relation, or $A \cup B$ is s-acyclic. Then exactly one of the following holds:

(a) the set A is separable from B by a pot(Σ_2^0) set,

(b) there is $f: 2^{\omega} \to X$ injective continuous such that the inclusions $\lceil T \rceil \setminus \mathbb{E}_0 \subseteq (f \times f)^{-1}(A)$ and $\lceil T \rceil \cap \mathbb{E}_0 \subseteq (f \times f)^{-1}(B)$ hold.

Proof. Let R be a pot (F_{σ}) s-acyclic relation containing A. Then there is no pot (Σ_2^0) set P separating $A \cap R = A$ from $B \cap R$, since otherwise $P \cap R \in \text{pot}(\Sigma_2^0)$ and separates A from B. Corollary 6.6 gives $f: 2^{\omega} \to X$ injective continuous with $[T] \cap \mathbb{E}_0 \subseteq (f \times f)^{-1}(B)$ and $[T] \setminus \mathbb{E}_0 \subseteq (f \times f)^{-1}(A)$.

If $A \cup B$ is s-acyclic, then we apply Corollary 6.4.

Remarks. (1) Corollary 6.7 also holds when $A \cup B$ is locally countable, but we did not mention it in the statement since (a) always holds in this case. Indeed, by reflection, $A \cup B$ is contained in a locally countable Borel set C. As A, B are disjoint analytic sets, there is a Borel set D separating A from B. Thus $C \cap D$ is a locally countable Borel set separating A from B. But a locally countable Borel set has Σ_2^0 vertical sections, and is therefore pot(Σ_2^0) (see [Lo2]).

(2) There is a version of Corollary 6.7 for $\Gamma = \Sigma_1^0$, where we replace the class F_{σ} with the class of open sets. We do not state it since (a) always holds in this case. Indeed, a potentially open s-acyclic relation is a countable union of Borel rectangles for which at least one side is a singleton, so that this union is potentially clopen, just like any of its Borel subsets.

7 The class Δ_2^0

Example

We set, for each $\varepsilon \in 2$,

 $\mathbb{E}_{0}^{\varepsilon} := \{ (\alpha, \beta) \in 2^{\omega} \times 2^{\omega} \mid \exists m > 0 \ \alpha(m) \neq \beta(m) \land \forall n > m \ \alpha(n) = \beta(n) \land (m-1)_{0} \equiv \varepsilon \pmod{2} \}.$

Lemma 7.1 $[T] \cap \mathbb{E}^0_0$ is not separable from $[T] \cap \mathbb{E}^1_0$ by a pot $(\mathbf{\Delta}^0_2)$ set.

Proof. The proof is similar to that of Lemma 6.1. We argue by contradiction, which gives D in $\operatorname{pot}(\Delta_2^0)$, and also a dense G_{δ} subset G of 2^{ω} such that $D \cap G^2 \in \Delta_2^0(G^2)$. Let $(O_n)_{n \in \omega}$ be a sequence of dense open subsets of 2^{ω} with intersection G. Note that $[T] \cap \mathbb{E}_0^0 \cap G^2 \subseteq [T] \cap D \cap G^2$, $[T] \cap \mathbb{E}_0^1 \cap G^2 \subseteq [T] \cap G^2 \setminus D$ and $[T] \cap D \cap G^2 \in \Delta_2^0([T] \cap G^2)$. By Baire's theorem, it is enough to prove that $[T] \cap \mathbb{E}_0^0 \cap G^2$ and $[T] \cap \mathbb{E}_0^1 \cap G^2$ are dense in $[T] \cap G^2$. Let us do it for $[T] \cap \mathbb{E}_0^0 \cap G^2$, the other case being similar. So let $q \in \omega$ and $w \in 2^{<\omega}$. Pick $N \in \omega$ such that $(s_q 0w0^{N_0}, t_q 1w0^N)$ is in \mathcal{F} and $(|s_q 0w0^N| - 1)_0 = 0$. Then we argue as in the proof of Lemma 6.1: pick $u_0 \in 2^{\omega}$ with $N_{s_q 0w0^N 0u_0} \subseteq O_0, v_0 \in 2^{\omega}$ with $N_{t_q 1w0^N 1u_0 v_0} \subseteq O_0, u_1 \in 2^{\omega}$ with $N_{s_q 0w0^N 0u_0 v_0 u_1 v_1} \subseteq O_1, v_1 \in 2^{\omega}$ with $N_{t_q 1w0^N 1u_0 v_0 u_1 v_1} \subseteq O_1$, and so on. Then $(s_q 0w0^N 0u_0 v_0 u_1 v_1..., t_q 1w0^N 1u_0 v_0 u_1 v_1...)$ is in $[T] \cap \mathbb{E}_0^0 \cap G^2$.

The main result

We will prove a version of Theorem 6.3 for the class Δ_2^0 .

Theorem 7.2 Let X be a Polish space, and A, B be disjoint analytic relations on X such that $A \cup B$ is quasi-acyclic. Then one of the following holds:

(a) the set A is separable from B by a pot(Δ_2^0) set,

(b) there is $f: 2^{\omega} \to X$ injective continuous such that the inclusions $[T] \cap \mathbb{E}_0^0 \subseteq (f \times f)^{-1}(A)$ and $[T] \cap \mathbb{E}_0^1 \subseteq (f \times f)^{-1}(B)$ hold.

Proof. The proof is similar to that of of Theorem 6.3. Assume that (a) does not hold. By Lemma 3.5, we may assume that A, B are Σ_2^0 . Let $(C_n)_{n \in \omega}$ be a witness for the fact that $A \cup B$ is quasi-acyclic. As A, B are Σ_2^0 , we may assume that each C_n is either contained in A, or contained in B. Note that there are disjoint Borel subsets O_0, O_1 of X such that $A \cap (O_0 \times O_1)$ is not separable from $B \cap (O_0 \times O_1)$ by a pot (Δ_2^0) set. We may assume that X is zero-dimensional, the C_n 's are closed, and O_0, O_1 are clopen, refining the topology if necessary. We can also replace A, B and the C_n 's with their intersection with $O_0 \times O_1$ and assume that they are contained in $O_0 \times O_1$. This gives a sequence $(C_n^0)_{n \in \omega}$ (resp., $(C_n^1)_{n \in \omega}$) of pairwise disjoint closed relations on X with union A (resp., B).

• We may assume that X is recursively presented, O_0, O_1 are Δ_1^1 and the relation " $(x, y) \in C_n^{\varepsilon}$ " is Δ_1^1 in (x, y, ε, n) . As Δ_X is Polish finer than the topology on X, A is not separable from B by a $\Delta_2^0(X^2, \tau_1)$ set. We set, for $F \in \Pi_1^0(X^2, \tau_1), F' := \overline{F \cap A^{\tau_1} \cap \overline{F \cap B^{\tau_1}}}$ (see 22.30 in [K]). Then

$$F_{\theta} = F_{\theta+1} = F_{\theta}' = \overline{F_{\theta} \cap A}^{\tau_1} \cap \overline{F_{\theta} \cap B}^{\tau_1},$$

so that $F_{\theta} \cap A$ and $F_{\theta} \cap B$ are τ_1 -dense in F_{θ} .

• Let us prove that F_{θ} is not empty. We argue by contradiction, so that $A = \bigcup_{\xi < \theta} A \cap F_{\xi} \setminus F_{\xi+1}$. But $A \cap F_{\xi} \setminus F_{\xi+1} = A \cap F_{\xi} \setminus (\overline{F_{\xi} \cap A}^{\tau_1} \cap \overline{F_{\xi} \cap B}^{\tau_1}) \subseteq F_{\xi} \setminus \overline{F_{\xi} \cap B}^{\tau_1} \subseteq \neg B$. This means that $(F_{\xi} \setminus F_{\xi+1})_{\xi < \theta}$ is a countable partition of (X^2, τ_1) into Δ_2^0 sets, and that A is separable from B by a Δ_2^0 set on each piece of the partition. This implies that A is separable from B by a $\Delta_2^0(X^2, \tau_1)$ set, which is absurd.

• As in the proof of Theorem 6.3, F_{θ} is Σ_1^1 , and the following key property holds:

$$\forall \varepsilon \in 2 \ \forall q \in \omega \ \forall U, V \in \varSigma_1^1(X) \ F_\theta \cap (U \times V) \neq \emptyset \Rightarrow \exists n \ge q \ F_\theta \cap C_n^\varepsilon \cap (U \times V) \neq \emptyset.$$

- We construct again sequences $(x_s), (y_s), (X_s), (Y_s)$ and Φ satisfying the following conditions:
 - $\begin{array}{l} (1) \ x_s \in X_s \ \land \ y_s \in Y_s \\ (2) \ X_{s\varepsilon} \subseteq X_s \subseteq \Omega_X \cap O_0 \ \land \ Y_{s\varepsilon} \subseteq Y_s \subseteq \Omega_X \cap O_1 \\ (3) \ \dim_{\operatorname{GH}}(X_s), \ \dim_{\operatorname{GH}}(Y_s) \leq 2^{-|s|} \\ (4) \ (x_{u_0}, y_{u_1}) \in F_{\theta} \cap C^{\varepsilon}_{\Phi(\vec{t}(\vec{u}))} \ \text{if} \ (|\vec{t}(\vec{u})| 2)_0 \equiv \varepsilon \ (\text{mod } 2), \ \text{with the convention} \ (-1)_0 = 0 \\ (5) \ X_{s0} \cap X_{s1} = Y_{s0} \cap Y_{s1} = \emptyset \end{array}$

• Assume that this has been done. If $(\alpha, \beta) \in [T] \cap \mathbb{E}_0^0$, then $\Phi(\vec{t}((\alpha, \beta)|n)) = N$ if n is big enough. In this case, by (4), $(x_{\alpha|n}, y_{\beta|n}) \in C_N^0$ which is closed, so that $(f(\alpha), g(\beta)) \in C_N^0 \subseteq A$. Similarly, if $(\alpha, \beta) \in [T] \cap \mathbb{E}_0^1$, then $(f(\alpha), g(\beta)) \in C_N^1 \subseteq B$. • So let us prove that the construction is possible. The key property gives $\Phi(0,1) \in \omega$ and (x_0, y_1) in $F_{\theta} \cap C^0_{\Phi(0,1)} \cap \Omega_{X^2}$. We choose Σ^1_1 subsets X_0, Y_1 of X with GH-diameter at most 2^{-1} such that $(x_0, y_1) \in X_0 \times Y_1 \subseteq (\Omega_X \cap O_0) \times (\Omega_X \cap O_1)$, which completes the construction for the length l = 1.

Let $l \ge 1$. We now want to build x_s, X_s, y_s, Y_s for $s \in 2^{l+1}$, as well as $\Phi(s_l 0, t_l 1)$. Fix $\eta \in 2$ such that $(l-1)_0 \equiv \eta \pmod{2}$. Note that $(x_{s_l}, y_{t_l}) \in F_{\theta} \cap (U \times V)$, where

$$U := \{ x'_{s_l} \in X_{s_l} \mid \exists (x'_s)_{s \in 2^l \setminus \{s_l\}, s(0) = 0} \in \Pi_{s \in 2^l \setminus \{s_l\}, s(0) = 0} X_s \quad \exists (y'_s)_{s \in 2^l, s(0) = 1} \in \Pi_{s \in 2^l, s(0) = 1} Y_s \\ \forall \vec{u} \in T \cap (2^l \times 2^l) \quad (x'_{u_0}, y'_{u_1}) \in F_{\theta} \cap C^{\varepsilon}_{\Phi(\vec{t}(\vec{u}))} \text{ if } (|\vec{t}(\vec{u})| - 2)_0 \equiv \varepsilon \pmod{2} \},$$

$$\begin{split} V := \{ y_{t_l}' \in Y_{t_l} \mid \exists (x_s')_{s \in 2^l, s(0) = 0} \in \Pi_{s \in 2^l, s(0) = 0} X_s \ \exists (y_s')_{s \in 2^l \setminus \{t_l\}, s(0) = 1} \in \Pi_{s \in 2^l \setminus \{t_l\}, s(0) = 1} Y_s \\ \forall \vec{u} \in T \cap (2^l \times 2^l) \ (x_{u_0}', y_{u_1}') \in F_\theta \cap C_{\Phi(\vec{t}(\vec{u}))}^{\varepsilon} \text{ if } (|\vec{t}(\vec{u})| - 2)_0 \equiv \varepsilon \pmod{2} \}. \end{split}$$

The key property gives $\Phi(s_l 0, t_l 1) > \max_{q < l} \Phi(s_q 0, t_q 1)$ and

$$(x_{s_l0}, y_{t_l1}) \in F_\theta \cap C^\eta_{\Phi(s_l0, t_l1)} \cap (U \times V).$$

Note that $x_{s_l0} \neq x_{s_l1}$ because $(x_{s_l0}, y_{t_l1}) \in C^{\eta}_{\Phi(s_l0, t_l1)}, (x_{s_l1}, y_{t_l1}) \in C^{\varepsilon}_{\Phi(\vec{t}(s_l1, t_l1))}$ if

$$(|\vec{t}(s_l1,t_l1)|-2)_0 \equiv \varepsilon \pmod{2},$$

and $\Phi(s_l 0, t_l 1) > \Phi(\vec{t}(s_l 1, t_l 1))$. Similarly, $y_{t_l 0} \neq y_{t_l 1}$. If $s \in 2^l$, then there is an injective s(T)-path p_s from s to s_l . This gives a $s(A \cup B)$ -path from x_{s0} to x_{s1} if s(0) = 0, and a $s(A \cup B)$ -path from y_{s0} to y_{s1} if s(0) = 1. Using the quasi-acyclicity of $s(A \cup B)$, we see, by induction on the length of p_s , that $x_{s0} \neq x_{s1}$ and $y_{s0} \neq y_{s1}$.

Consequences

Corollary 7.3 Let X be a Polish space, and A, B be disjoint analytic relations on X such that

- either $A \cup B$ is either s-acyclic or locally countable

- or A is contained in a pot(Δ_2^0) s-acyclic or locally countable relation.

Then exactly one of the following holds:

(a) the set A is separable from B by a pot(Δ_2^0) set,

(b) there is $f: 2^{\omega} \to X$ injective continuous such that the inclusions $[T] \cap \mathbb{E}_0^0 \subseteq (f \times f)^{-1}(A)$ and $[T] \cap \mathbb{E}_0^1 \subseteq (f \times f)^{-1}(B)$ hold.

Proof. By Lemma 7.1, $[T] \cap \mathbb{E}_0^0$ is not separable from $[T] \cap \mathbb{E}_0^1$ by a pot (Δ_2^0) set. This shows that (a) and (b) cannot hold simultaneously. So assume that (a) does not hold.

- If $A \cup B$ is s-acyclic or locally countable, then by Lemma 3.5, we may assume that A, B are Σ_2^0 . By Lemma 2.2, we may also assume that $A \cup B$ is quasi-acyclic. It remains to apply Theorem 7.2.

- Assume that R is $pot(\Delta_2^0)$ and contains A. Then there is no $pot(\Delta_2^0)$ set P separating $A \cap R = A$ from $B \cap R$, since otherwise $P \cap R \in pot(\Delta_2^0)$ separates A from B. It remains to apply the first point. This finishes the proof.

Corollary 7.4 Let X, Y be Polish spaces, and A, B be disjoint analytic subsets of $X \times Y$ such that $A \cup B$ is locally countable or A is contained in a pot (Δ_2^0) locally countable set. Then exactly one of the following holds:

- (a) the set A is separable from B by a pot(Δ_2^0) set,
- (b) $(2^{\omega}, 2^{\omega}, \lceil T \rceil \cap \mathbb{E}^0_0, \lceil T \rceil \cap \mathbb{E}^1_0) \sqsubseteq (X, Y, A, B).$

Proof. As in the proof of Corollary 7.3, (a) and (b) cannot hold simultaneously. Then we argue as in the proof of Corollary 3.7. The set $A' \cup B'$ is locally countable or A' is contained in a pot $(\mathbf{\Delta}_2^0)$ locally countable set, and A' is not separable from B' by a pot $(\mathbf{\Delta}_2^0)$ set. Corollary 7.3 gives $f': 2^{\omega} \to Z$. \Box

Corollary 7.5 Let X be a Polish space, and A, B be disjoint analytic relations on X. The following are equivalent:

(1) there is an s-acyclic or locally countable relation $R \in \Sigma_1^1$ such that $A \cap R$ is not separable from $B \cap R$ by a pot (Δ_2^0) set,

(2) there is $f: 2^{\omega} \to X$ injective continuous with $[T] \cap \mathbb{E}_0^0 \subseteq (f \times f)^{-1}(A)$ and $[T] \cap \mathbb{E}_0^1 \subseteq (f \times f)^{-1}(B)$.

Proof. (1) \Rightarrow (2) We apply Corollary 7.3.

(2) \Rightarrow (1) We can take $R := (f \times f) [[T] \cap \mathbb{E}_0].$

8 The classes $D_n(\Sigma_2^0)$ and $\check{D}_n(\Sigma_2^0)$

Examples

Notation. Let $\eta \ge 1$ be a countable ordinal, and $S_{\eta}: \omega \to \eta$ be onto. We set

$$C_0 := \{ \alpha \in 2^\omega \mid \exists m \in \omega \; \forall p \ge m \; \alpha(p) = 0 \}$$

and, for $1 \leq \theta < \eta$, $C_{\theta} := \{ \alpha \in 2^{\omega} \mid \exists m \in \omega \; \forall p \in \omega \; \alpha (< m, p >) = 0 \land S_{\eta}((m)_{0}) \leq \theta \}$, so that $(C_{\theta})_{\theta < \eta}$ is an increasing sequence of Σ_{2}^{0} subsets of 2^{ω} . We then set $D_{\eta} := D((C_{\theta})_{\theta < \eta})$.

Lemma 8.1 The set D_{η} is $D_{\eta}(\Sigma_2^0)$ -complete.

Proof. By 21.14 in [K], it is enough to see that D_{η} is not $D_{\eta}(\Sigma_2^0)$ since it is $D_{\eta}(\Sigma_2^0)$. We will prove more. Let us say that a pair (θ, F) is **suitable** if $\theta \leq \eta$, F is a chain of finite binary sequences, $I_F := \bigcap_{s \in F} \{\alpha \in N_s \mid (\alpha)_{|s|} = 0^\infty\}$ is not empty and $S_{\eta}((|s|)_0) \geq \theta$ for each $s \in F$. Let us prove that $I_F \cap D((C_{\theta'})_{\theta' < \theta})$ is not $D_{\theta}(\Sigma_2^0)$ if (θ, F) is suitable. This will give the result since (η, \emptyset) is suitable.

We argue by induction on θ . If $\theta = 1$, then the Σ_2^0 set $I_F \cap C_0$ is dense and co-dense in the closed set I_F , so that it is not Π_2^0 , by Baire's theorem. Assume the result proved for $\theta' < \theta$. We argue by contradiction, which gives an increasing sequence $(H_{\theta'})_{\theta' < \theta}$ of Σ_2^0 sets with

$$I_F \cap D\bigl((C_{\theta'})_{\theta' < \theta}\bigr) = \neg D\bigl((H_{\theta'})_{\theta' < \theta}\bigr).$$

As $\neg (\bigcup_{\theta' < \theta} C_{\theta'})$ is comeager in I_F , $I_F \cap \bigcup_{\theta' < \theta} H_{\theta'}$ too, which gives $\theta' < \theta$ with parity opposite to that of θ and $s' \supseteq \max_{s \in F} s$ such that $S_\eta((|s'|)_0) = \theta'$ and $\emptyset \neq I_F \cap N_{s'} \subseteq H_{\theta'}$. We set $F' := F \cup \{s'\}$, so that (θ', F') is suitable. By induction assumption, $I_{F'} \cap D((C_{\theta''})_{\theta'' < \theta'})$ is not $\check{D}_{\theta'}(\Sigma_2^0)$. But $I_{F'} \cap D((C_{\theta''})_{\theta'' < \theta'}) = I_{F'} \setminus D((H_{\theta''})_{\theta'' < \theta'}) \in \check{D}_{\theta'}(\Sigma_2^0)$ since $I_{F'} \subseteq C_{\theta'}$, which is absurd. \Box

Notation. We now fix an effective frame in the sense of Definition 2.1 in [L8], which are frames in the sense of Definition 5.1. Lemma 2.3 in [L8] proves the existence of such an effective frame. Note that $(s_1, t_1) = (0, 1)$, so that $s_1(0) \neq t_1(0)$. But $s_{l+1}(l) = t_{l+1}(l)$ if $l \ge 1$. Indeed, it is enough to see that $((l)_1)_1_0 + ((l)_1)_1_1 < l$ in this case, by the proof of Lemma 2.3 in [L8]. As $(q)_0 + (q)_1 \le q$, and $(q)_0 + (q)_1 < q$ if $q \ge 2$, we may assume that $((l)_1)_1 \in 2$. If $((l)_1)_1 = 0$, then we are done since $l \ge 1$. If $((l)_1)_1 = 1$, then $l \ge 2$ and we are done too.

• The shift map $S: 2^L \to 2^{L-1}$ is defined by $S(\alpha)(m) := \alpha(m+1)$ when $1 \le L \le \omega$, with the convention $\omega - 1 := \omega$.

• The symmetric difference $\alpha \Delta \beta$ of $\alpha, \beta \in 2^L$ is the element of 2^L defined by $(\alpha \Delta \beta)(m) = 1$ exactly when $\alpha(m) \neq \beta(m)$, if $L \leq \omega$.

• We set $\mathbb{N}_{\eta} := \{ (\alpha, \beta) \in [T] \mid \mathcal{S}(\alpha \Delta \beta) \notin D_{\eta} \}.$

Lemma 8.2 The $\check{D}_{\eta}(\Sigma_2^0)$ set \mathbb{N}_{η} is not separable from $[T] \setminus \mathbb{N}_{\eta}$ by a pot $(D_{\eta}(\Sigma_2^0))$ set.

Proof. As $\lceil T \rceil$ is closed, D_{η} is $D_{\eta}(\Sigma_2^0)$ and S, Δ are continuous, \mathbb{N}_{η} is $\check{D}_{\eta}(\Sigma_2^0)$. By Lemma 2.6 in [L8], it is enough to check that D_{η} is ccs (see Definition 2.5 in [L8]). We just have to check that the C_{θ} 's are ccs. So let $\alpha, \alpha_0 \in 2^{\omega}$ and $F : 2^{\omega} \to 2^{\omega}$ satisfying the conclusion of Lemma 2.4.(b) in [L8]. Note that $\alpha \in C_0$ exactly when $\{m \in \omega \mid \alpha(m) = 1\}$ is finite, so that C_0 is ccs. If $\theta \ge 1$, then $\alpha \notin C_{\theta}$ exactly when, for each $m, S_{\eta}((m)_0) \le \theta$ or there is p with $\alpha(< m, p >) = 1$. As $(B_{\alpha}(< m, p >))_0 = (< m, p >)_0 = m, C_{\theta}$ is ccs too.

The main result

Notation. From now on, $\eta < \omega$. We set, for $2 \le \theta \le \eta$ and $(s, t) \in (2 \times 2)^{<\omega} \setminus \{(\emptyset, \emptyset)\},$

$$m^{\theta}_{s,t} := \min \big\{ m \in \omega \mid \big(\mathcal{S}(s\Delta t) \big)_m \subseteq 0^{\infty} \land S_{\eta} \big((m)_0 \big) \! < \! \theta \big\}.$$

We also set $s^- := < s(0), ..., s(|s|-2) > \text{if } s \in 2^{<\omega}$.

• We define the following relation on $(2 \times 2)^{<\omega}$:

$$\begin{split} (s,t) \ R \ (s',t') \Leftrightarrow (s,t) &\subseteq (s',t') \ \land \ \left(|s| \leq 1 \ \lor \ \left(|s| \geq 2 \ \land \ \exists 2 \leq \theta \leq \eta \ m_{s,t}^{\theta} \neq m_{s^-,t^-}^{\theta} \land \\ \forall (s,t) &\subseteq (s'',t'') \subseteq (s',t') \ \forall \theta < \theta' \leq \eta \ m_{s,t}^{\theta'} = m_{s^-,t^-}^{\theta'} = m_{s'',t''}^{\theta'} \right) \lor \\ \left(|s| \geq 2 \ \land \ s(|s|-1) \neq t(|s|-1) \land \\ \forall (s,t) &\subseteq (s'',t'') \subseteq (s',t') \ \forall 2 \leq \theta \leq \eta \ m_{s,t}^{\theta} = m_{s^-,t^-}^{\theta} = m_{s'',t''}^{\theta'} \right) \lor \\ \left(|s| \geq 2 \ \land \ \forall (s,t) \subseteq (s'',t'') \subseteq (s',t') \ (\forall 2 \leq \theta \leq \eta \ m_{s,t}^{\theta} = m_{s^-,t^-}^{\theta} = m_{s'',t''}^{\theta'} \right) \land \\ s''(|s''|-1) = t''(|s''|-1) \right) \end{split}$$

Note that R is a **tree relation**, which means that it is a partial order (it contains the diagonal, is antisymmetric and transitive) with minimum element (\emptyset, \emptyset) , the set of predecessors of any sequence is finite and lineary ordered by R. Moreover, R is **distinguished** in \subseteq , which means that (s, t) R(s', t') if $(s, t) \subseteq (s', t') \subseteq (s'', t'')$ and (s, t) R(s'', t'') (see [D-SR]).

• We set

$$\begin{split} D_{\eta} &:= \{ (s,t) \in T \mid |s| \ge 2 \Rightarrow m_{s,t}^{\eta} \neq m_{s^{-},t^{-}}^{\eta} \} \text{ if } \eta \ge 2, \\ D_{\theta} &:= \{ (s,t) \in T \mid |s| \ge 2 \land m_{s,t}^{\theta} \neq m_{s^{-},t^{-}}^{\theta} \land \forall \theta < \theta' \le \eta \ m_{s,t}^{\theta'} = m_{s^{-},t^{-}}^{\theta'} \} \text{ if } 2 \le \theta < \eta \\ D_{1} &:= \{ (s,t) \in T \mid |s| \ge 2 \land \forall 2 \le \theta \le \eta \ m_{s,t}^{\theta} = m_{s^{-},t^{-}}^{\theta} \land s(|s|-1) \ne t(|s|-1) \}, \\ D_{0} &:= \{ (s,t) \in T \mid |s| \ge 2 \land s(|s|-1) = t(|s|-1) \}, \end{split}$$

so that the $(D_{\theta})_{\theta < \eta}$ is a partition of T.

Theorem 8.3 Let $1 \le \eta < \omega$. Let X be a Polish space, and A_0, A_1 be disjoint analytic relations on X such that $A_0 \cup A_1$ is s-acyclic. Then exactly one of the following holds:

(a) the set A₀ is separable from A₁ by a pot(D_η(Σ₂⁰)) set,
(b) (2^ω, 2^ω, ℕ_η, [T] \ℕ_η) ⊑ (X, X, A₀, A₁), via a square map.

Proof. By Lemma 8.2, (a) and (b) cannot hold simultaneously. So assume that (a) does not hold. Note first that we may assume that $A_0 \cup A_1$ is compact and A_1 is $D_\eta(\Sigma_2^0)$. Indeed, Theorems 1.9 and 1.10 in [L8] give $\mathbb{S} \in D_\eta(\Sigma_2^0)(\lceil T \rceil)$ and $f', g' : 2^\omega \to X$ continuous such that the inclusions $\mathbb{S} \subseteq (f' \times g')^{-1}(A_1)$ and $\lceil T \rceil \setminus \mathbb{S} \subseteq (f' \times g')^{-1}(A_0)$ hold. Let $(\Sigma_\theta)_{\theta < \eta}$ be an increasing sequence of $\Sigma_2^0(\lceil T \rceil)$ sets with $\mathbb{S} = D((\Sigma_\theta)_{\theta < \eta}), K := (f' \times g')[\lceil T \rceil]$, and $R_\theta := (f' \times g')[\Sigma_\theta]$. Note that K is compact, R_θ is K_σ , $D((R_\theta)_{\theta < \eta}) \subseteq A_1, K \setminus D((R_\theta)_{\theta < \eta}) \subseteq A_0, D((R_\theta)_{\theta < \eta}) = K \cap A_1, K \setminus D((R_\theta)_{\theta < \eta}) = K \cap A_0$, so that $D((R_\theta)_{\theta < \eta})$ is not separable from $K \setminus D((R_\theta)_{\theta < \eta})$ by a pot $(\check{D}_\eta(\Sigma_2^0))$ set. So we can replace A_1, A_0 with $D((R_\theta)_{\theta < \eta}), K \setminus D((R_\theta)_{\theta < \eta})$, respectively. • We may also assume that X is zero-dimensional and there are disjoint clopen subsets O_0, O_1 of X such that $A_0 \cap (O_0 \times O_1)$ is not separable from $A_1 \cap (O_0 \times O_1)$ by a pot $(D_\eta(\Sigma_2^0))$ set. So, without loss of generality, we will assume that $A_0 \cup A_1 \subseteq O_0 \times O_1$. We may also assume that X is recursively presented, $A_0, A_1, O_0, O_1, R_\theta$ are Δ_1^1 , and R_θ is the union of $\Delta_1^1 \cap \Pi_1^0 \subseteq \Sigma_1^1 \cap \Pi_1^0(\tau_1) \subseteq \Sigma_1^0(\tau_2)$ sets.

We set, for $\theta < \eta$, $N_{\theta} := R_{\theta} \setminus (\bigcup_{\theta' < \theta} R_{\theta'}) \cap \bigcap_{\theta' < \theta} \overline{N_{\theta'}}^{\tau_2}$. Note that the N_{θ} 's are pairwise disjoint, which will be useful in the construction to get the injectivity of our reduction maps. We use the notation of Theorem 3.2. For simplicity, we set $F_{\theta}^{\varepsilon} := F_{\theta,2}^{\varepsilon}$.

Claim. (a) Assume that $k+1 < \eta$. Then $F_k^{\varepsilon} = \overline{N_k}^{\tau_2} \cup E_k$, where $E_k \subseteq \neg R_{k+1}$ is τ_2 -closed. (b) $A_0 \cap \bigcap_{\theta < \eta} F_{\theta}^{\varepsilon} = N_{\eta} := K \setminus (\bigcup_{\theta < \eta} R_{\theta}) \cap \bigcap_{\theta < \eta} \overline{N_{\theta}}^{\tau_2}$.

(a) Indeed, we argue by induction on k to prove (a). In the proof of this claim, all the closures will refer to τ_2 . Note first that $R_0 \subseteq A_{\varepsilon} \subseteq R_0 \cup \neg R_1$, so that $F_0^{\varepsilon} = \overline{A_{\varepsilon}} = \overline{R_0} \cup E_0 = \overline{N_0} \cup E_0$. Then, inductively,

$$F_{k+1}^{\varepsilon} = \overline{A_{1-|\text{parity}(k)-\varepsilon|} \cap F_{k}^{\varepsilon}} = \overline{A_{1-|\text{parity}(k)-\varepsilon|} \cap (\overline{N_{k}} \cup E_{k})}$$
$$= \overline{\left((R_{k+1} \setminus R_{k}) \cup (R_{k+3} \setminus R_{k+2})...\right) \cap (\overline{N_{k}} \cup E_{k})} = \overline{N_{k+1}} \cup E_{k+1}.$$
(b) Note then that $F_{\eta-1}^{\varepsilon} = \overline{A_{1} \cap \bigcap_{k+1 < \eta} F_{k}^{\varepsilon}} = \overline{A_{1} \cap \bigcap_{k+1 < \eta} (\overline{N_{k}} \cup E_{k})} = \overline{N_{\eta-1}}$, so that

$$A_0 \cap \bigcap_{\theta < \eta} F_{\theta}^{\varepsilon} = K \setminus (\bigcup_{\theta < \eta} R_{\theta}) \cap \bigcap_{\theta < \eta} \overline{N_{\theta}}.$$

This proves the claim.

- We construct the following objects:
- sequences $(x_s)_{s \in 2^{<\omega}, 0 \subseteq s}$, $(y_s)_{s \in 2^{<\omega}, 1 \subseteq s}$ of points of X,
- sequences $(X_s)_{s \in 2^{<\omega}, 0 \subseteq s}, (Y_s)_{s \in 2^{<\omega}, 1 \subseteq s}$ of Σ_1^1 subsets of X,
- a sequence $(U_{s,t})_{(s,t)\in T\setminus\{(\emptyset,\emptyset)\}}$ of Σ_1^1 subsets of X^2 .

We want these objects to satisfy the following conditions:

$$(1) x_{s} \in X_{s} \land y_{s} \in Y_{s} \land (x_{s}, y_{t}) \in U_{s,t}$$

$$(2) X_{s\varepsilon} \subseteq X_{s} \subseteq \Omega_{X} \cap O_{0} \land Y_{s\varepsilon} \subseteq Y_{s} \subseteq \Omega_{X} \cap O_{1} \land U_{s,t} \subseteq \Omega_{X^{2}} \cap (X_{s} \times Y_{t})$$

$$(3) \operatorname{diam}_{\operatorname{GH}}(X_{s}), \operatorname{diam}_{\operatorname{GH}}(Y_{s}), \operatorname{diam}_{\operatorname{GH}}(U_{s,t}) \leq 2^{-|s|}$$

$$(4) X_{s0} \cap X_{s1} = Y_{s0} \cap Y_{s1} = \emptyset$$

$$(5) ((s,t) R (s',t') \land \exists \theta \leq 2 (s,t), (s',t') \in D_{\theta}) \Rightarrow U_{s',t'} \subseteq U_{s,t}$$

$$(6) U_{s,t} \subseteq N_{\theta} \text{ if } (s,t) \in D_{\theta}$$

$$(7) (s,t) R (s',t') \Rightarrow U_{s',t'} \subseteq \overline{U_{s,t}}^{\tau_{1}}$$

 \diamond

• Assume that this has been done. As in the proof of Theorem 3.3, we get $f: 2^{\omega} \to X$ injective continuous. If $(\alpha, \beta) \in \mathbb{N}_{\eta}$, then we can find $\theta < \eta$ of parity opposite to that of η and $(n_k)_{k \in \omega}$ strictly increasing such that $(\alpha, \beta)|n_k \in D_{\theta}$ and $(\alpha, \beta)|n_k R (\alpha, \beta)|n_{k+1}$ for each $k \in \omega$. In this case, by (1)-(3) and (5)-(6), $(U_{(\alpha,\beta)|n_k})_{k\in\omega}$ is a decreasing sequence of nonempty clopen subsets of $A_0 \cap \Omega_{X^2}$ with vanishing diameters, so that its intersection is a singleton $\{F(\alpha, \beta)\} \subseteq A_0$. As $(x_{\alpha|n}, y_{\beta|n})$ converges (for Σ_{X^2} and thus for Σ_X^2) to $F(\alpha, \beta), (f(\alpha), f(\beta)) = F(\alpha, \beta) \in A_0$. If $(\alpha, \beta) \in [T] \setminus \mathbb{N}_{\eta}$, then we argue similarly to see that $(f(\alpha), f(\beta)) \in A_1$.

• So let us prove that the construction is possible. Let $(x_0, y_1) \in N_\eta \cap \Omega_{X^2}$, X_0, Y_1 be Σ_1^1 subsets of X with diameter at most 2^{-1} such that $x_0 \in X_0 \subseteq \Omega_X \cap O_0$ and $y_1 \in Y_1 \subseteq \Omega_X \cap O_1$, and $U_{0,1}$ be a Σ_1^1 subset of X^2 with diameter at most 2^{-1} such that $(x_0, y_1) \in U_{0,1} \subseteq N_\eta \cap \Omega_{X^2} \cap (X_0 \times Y_1)$. This completes the construction for l = 1 since $(0, 1) \in D_\eta$.

- Note that $(0^2, 1^2) \in D_\eta$ since $m_{0,1}^\eta = 0$ and $m_{0^2, 1^2}^\eta = 1$ if $\eta \ge 2$. We set $S_0 := \overline{U_{0,1}}^{\tau_1} \cap (X_0 \times Y_1)$ and $S_1 := S_0 \cap N_0 \cap \Omega_{X^2}$. As $U_{0,1} \subseteq \overline{N_0}^{\tau_2}$, $S_0 \subseteq \overline{S_1}^{\tau_1}$. In particular, $\Pi_{\varepsilon}[S_1]$ is Σ_X -dense in $\Pi_{\varepsilon}[S_0]$ for each $\varepsilon \in 2$, by continuity of the projections. As $(x_0, y_1) \in U_{0,1} \cap (\Pi_0[S_0] \times \Pi_1[S_0])$, this implies that $U_{0,1} \cap (\Pi_0[S_1] \times \Pi_1[S_1])$ is not empty and contains some (x_{0^2}, y_{1^2}) (the projections maps are open). This gives $y_{10} \in X$ with $(x_{0^2}, y_{10}) \in S_1$, and $x_{01} \in X$ with $(x_{01}, y_{1^2}) \in S_1$. As $U_{0,1} \subseteq N_\eta$ and $S_1 \subseteq N_0$, $x_{0^2} \neq x_{01}$ and $y_{10} \neq y_{1^2}$. It remains to choose Σ_1^1 subsets $X_{0^2}, X_{01}, Y_{10}, Y_{1^2}$ of X with diameter at most 2^{-2} such that $(x_{0\varepsilon}, y_{1\varepsilon}) \in X_{0\varepsilon} \times Y_{1\varepsilon} \subseteq X_0 \times Y_1$ and $X_{0^2} \cap X_{01} = Y_{10} \cap Y_{1^2} = \emptyset$, as well as Σ_1^1 subsets $U_{0^2,1^2}, U_{0^2,10}, U_{01,1^2}$ of X^2 with diameter at most 2^{-2} such that $(x_{0\varepsilon}, y_{1\varepsilon}) \in U_{0\varepsilon,1\varepsilon} \subseteq \overline{U_{0,1}}^{\tau_1} \cap N_0 \cap \Omega_{X^2} \cap (X_{0\varepsilon} \times Y_{1\varepsilon})$. This completes the construction for l = 2.

- Assume that our objects are constructed for the level $l \ge 2$, which is the case for l = 2. Note that $(s_l 0, t_l 1) \notin D_0$, and we already noticed that $s_l(l-1) = t_l(l-1)$ since $l \ge 2$, so that $(s_l, t_l) \in D_0$. We set $(\tilde{s}, \tilde{t}) := (s_{l-1}0, t_{l-1}1)$ (which is not in D_0), and

$$S_{0} := \left\{ \left((\overline{x}_{s})_{s \in 2^{l}, 0 \subseteq s}, (\overline{y}_{t})_{t \in 2^{l}, 1 \subseteq t} \right) \in X^{2^{l}} \mid \forall (s, t) \in T \cap (2^{l} \times 2^{l}) \setminus \{ (\tilde{s}, \tilde{t}) \} \quad (\overline{x}_{s}, \overline{y}_{t}) \in U_{s, t} \land (\overline{x}_{\tilde{s}}, \overline{y}_{\tilde{t}}) \in \overline{N_{0}}^{\tau_{2}} \cap \overline{U_{\tilde{s}, \tilde{t}}}^{\tau_{1}} \cap (X_{\tilde{s}} \times Y_{\tilde{t}}) \right\},$$

$$S_1 := \left\{ \left((\overline{x}_s)_{s \in 2^l, 0 \subseteq s}, (\overline{y}_t)_{t \in 2^l, 1 \subseteq t} \right) \in S_0 \mid (\overline{x}_{\tilde{s}}, \overline{y}_{\tilde{t}}) \in N_0 \cap \Omega_{X^2} \right\}.$$

We equip X^{2^l} with the product of the Gandy-Harrington topologies. Let us show that S_1 is dense in S_0 . Let $(\mathcal{U}_s)_{s \in 2^l, 0 \subseteq s}$ and $(\mathcal{V}_t)_{t \in 2^l, 1 \subseteq t}$ be sequences of Σ_1^1 sets with

$$\left((\Pi_{s \in 2^l, 0 \subset s} \mathcal{U}_s) \times (\Pi_{t \in 2^l, 1 \subset t} \mathcal{V}_t) \right) \cap S_0 \neq \emptyset$$

with witness $((x'_s), (y'_t))$, $\mathcal{A}_{\varepsilon} := \{s \in 2^l \mid s(l-1) = \varepsilon\}$, and

$$U := \{ \overline{x}_{\tilde{s}} \in \mathcal{U}_{\tilde{s}} \mid \exists (\overline{x}_s)_{s \in \mathcal{A}_0 \setminus \{\tilde{s}\}} \in \Pi_{s \in \mathcal{A}_0 \setminus \{\tilde{s}\}} \mathcal{U}_s \ \exists (\overline{y}_t)_{t \in \mathcal{A}_0} \in \Pi_{t \in \mathcal{A}_0} \mathcal{V}_t \\ \forall (s,t) \in T \cap (\mathcal{A}_0 \times \mathcal{A}_0) \ (\overline{x}_s, \overline{y}_t) \in U_{s,t} \},$$

$$V := \{ \overline{y}_{\tilde{t}} \in \mathcal{V}_{\tilde{t}} \mid \exists (\overline{x}_s)_{s \in \mathcal{A}_1} \in \Pi_{s \in \mathcal{A}_1} \mathcal{U}_s \ \exists (\overline{y}_t)_{t \in \mathcal{A}_1 \setminus \{\tilde{t}\}} \in \Pi_{t \in \mathcal{A}_1 \setminus \{\tilde{t}\}} \mathcal{V}_t \\ \forall (s, t) \in T \cap (\mathcal{A}_1 \times \mathcal{A}_1) \ (\overline{x}_s, \overline{y}_t) \in U_{s, t} \}.$$

Then $(x_{\tilde{s}}', y_{\tilde{t}}') \in \overline{N_0}^{\tau_2} \cap \overline{U_{\tilde{s},\tilde{t}}}^{\tau_1} \cap (U \times V)$. This gives $(\overline{x}_{\tilde{s}}, \overline{y}_{\tilde{t}})$ in $N_0 \cap \overline{U_{\tilde{s},\tilde{t}}}^{\tau_1} \cap (U \times V) \cap \Omega_{X^2}$. We choose witnesses $(\overline{x}_s)_{s \in \mathcal{A}_0 \setminus \{\tilde{s}\}}, (\overline{y}_t)_{t \in \mathcal{A}_0}$ (resp., $(\overline{x}_s)_{s \in \mathcal{A}_1}, (\overline{y}_t)_{t \in \mathcal{A}_1 \setminus \{\tilde{t}\}}$) for the fact that $\overline{x}_{\tilde{s}} \in U$ (resp., $\overline{y}_{\tilde{t}} \in V$). Then $((\overline{x}_s), (\overline{y}_t)) \in ((\Pi_{s \in 2^l, 0 \subseteq s} \mathcal{U}_t) \times (\Pi_{t \in 2^l, 1 \subseteq t} \mathcal{V}_t)) \cap S_1$, as desired.

The sets $U_{\varepsilon} := \prod_{s_l} [S_{\varepsilon}]$ and $V_{\varepsilon} := \prod_{t_l} [S_{\varepsilon}]$ are Σ_1^1 sets. As S_1 is dense in S_0 , U_1 (resp., V_1) is dense in U_0 (resp., V_0). Note that $(x_{s_l}, y_{t_l}) \in U_{s_l, t_l} \cap (U_0 \times V_0)$. As U_1 (resp., V_1) is dense in U_0 (resp., V_0), U_{s_l, t_l} meets $U_1 \times V_1$.

Let $(s_l0, t_l1)^R$ be the *R*-predecessor of (s_l0, t_l1) . Assume first that $(s_l0, t_l1) \in D_\eta$. Then $(s_l0, t_l1)^R \in D_\eta$ too. Note that $U_{s_l,t_l} \subseteq \overline{U_{(s_l0,t_l1)^R}}^{\tau_1}$ since $(s_l0, t_l1)^R R (s_l, t_l)$. Thus $\overline{U_{(s_l0,t_l1)^R}}^{\tau_1}$ meets $U_1 \times V_1$. This gives $(x_{s_l0}, y_{t_l1}) \in U_{(s_l0,t_l1)^R} \cap (U_1 \times V_1)$. We choose witnesses $(x_{s0})_{s \in 2^l \setminus \{s_l\}, 0 \subseteq s}$, $(y_{t0})_{t \in 2^l, 1 \subseteq t}$ (resp., $(x_{s1})_{s \in 2^l, 0 \subseteq s}$, $(y_{t1})_{t \in 2^l \setminus \{t_l\}, 1 \subseteq t}$) for the fact that $x_{s_l0} \in U_1$ (resp., $y_{t_l1} \in V_1$). As $(x_{s_l0}, y_{t_l1}) \in U_{(s_l0,t_l1)^R} \subseteq N_\eta$ and $(x_{s_l\varepsilon}, y_{t_l\varepsilon}) \in N_0, x_{s_l0} \neq x_{s_l1}$ and $y_{t_l0} \neq y_{t_l1}$. As in the proof of Theorem 3.3, the s-acyclicity of $A_0 \cup A_1$ and the fact that O_0, O_1 are disjoint ensure the fact that $x_{s0} \neq x_{s1}$ and $y_{t0} \neq y_{t_1}$ for s, t arbitrary with the right first coordinate. Then we choose Σ_1^1 subsets $X_{s\varepsilon}, Y_{t\varepsilon}$ of X with diameter at most 2^{-l-1} such that $(x_{s\varepsilon}, y_{t\varepsilon}) \in X_{s\varepsilon} \times Y_{t\varepsilon} \subseteq X_s \times Y_t$ and $X_{s0} \cap X_{s1} = Y_{s0} \cap Y_{s1} = \emptyset$, as well as Σ_1^1 subsets $U_{s\varepsilon,t\varepsilon'}$ of X^2 , with diameter at most 2^{-l-1} , containing $(x_{s\varepsilon}, y_{t\varepsilon'})$ and contained in $X_{s\varepsilon} \times Y_{t\varepsilon}$, such that

 $\begin{aligned} &- U_{s_l0,t_l1} \subseteq U_{(s_l0,t_l1)^R}, \\ &- U_{\tilde{s}\varepsilon,\tilde{t}\varepsilon} \subseteq \overline{U_{\tilde{s},\tilde{t}}}^{\tau_1} \cap N_0 \cap \Omega_{X^2}, \\ &- U_{s\varepsilon,t\varepsilon} \subseteq U_{s,t} \text{ if } (s,t) \neq (\tilde{s},\tilde{t}). \end{aligned}$

The argument is the same if $(s_l0, t_l1), (s_l0, t_l1)^R \in D_\theta$. So it remains to study the case where $(s_l0, t_l1) \in D_{\theta'}$ and $(s_l0, t_l1)^R \in D_\theta$, and $\theta' < \theta$. In this case, note that $U_{(s_l0, t_l1)^R} \cap (U_1 \times V_1)$ is not empty and contained in $N_\theta \subseteq \overline{N_{\theta'}}^{\tau_2}$. This gives $(x_{s_l0}, y_{t_l1}) \in N_{\theta'} \cap \overline{U_{(s_l0, t_l1)^R}}^{\tau_1} \cap \Omega_{X^2} \cap (U_1 \times V_1)$, and we conclude as before.

Consequences

Corollary 8.4 Let $1 \le \eta < \omega$, X be a Polish space, and A, B be disjoint analytic relations on X such that A is contained in a pot(Δ_2^0) s-acyclic relation. Then exactly one of the following holds:

- (a) the set A is separable from B by a pot $(D_{\eta}(\boldsymbol{\Sigma}_{2}^{0}))$ set,
- (b) $(2^{\omega}, 2^{\omega}, \mathbb{N}_n, [T] \setminus \mathbb{N}_n) \sqsubseteq (X, X, A, B)$, via a square map.

Proof. Let R be a $pot(\Delta_2^0)$ s-acyclic relation containing A. By Lemma 8.2, (a) and (b) cannot hold simultaneously. So assume that (a) does not hold. Then A is not separable from $B \cap R$ by a $pot(D_\eta(\Sigma_2^0))$ set. This allows us to apply Theorem 8.3.

Corollary 8.5 Let $1 \le \eta < \omega$, X be a Polish space, and A, B be disjoint analytic relations on X. The following are equivalent:

(1) there is $R \in \Sigma_1^1$ s-acyclic such that $A \cap R$ is not separable from $B \cap R$ by a pot $(D_\eta(\Sigma_2^0))$ set, (2) there is $f: 2^\omega \to X$ injective continuous such that $\mathbb{N}_\eta \subseteq (f \times f)^{-1}(A)$ and $[T] \setminus \mathbb{N}_\eta \subseteq (f \times f)^{-1}(B)$. **Proof.** (1) \Rightarrow (2) We apply Theorem 8.3.

(2) \Rightarrow (1) We can take $R := (f \times f) [[T]].$

9 Oriented graphs

Proof of Theorem 1.9. Theorem 1.3 provides Borel relations \mathbb{S}_0 , \mathbb{S}_1 on 2^{ω} . We saw that $\mathbb{S}_0 \cup \mathbb{S}_1$ is a subset of the body of a tree T, which does not depend on Γ , and is contained in $N_0 \times N_1$. We set $\mathbb{G}_{\Gamma} := \mathbb{S}_0 \cup (\mathbb{S}_1)^{-1}$, so that \mathbb{G}_{Γ} is Borel. As $\mathbb{S}_0 \cup \mathbb{S}_1 \subseteq N_0 \times N_1$ and \mathbb{S}_0 , \mathbb{S}_1 are disjoint, \mathbb{G}_{Γ} is an oriented graph. If (a) and (b) hold, then \mathbb{G}_{Γ} is separable from \mathbb{G}_{Γ}^{-1} by a pot(Γ) set S. Note that S also separates $\mathbb{S}_0 = \mathbb{G}_{\Gamma} \cap (N_0 \times N_1)$ from $\mathbb{S}_1 = \mathbb{G}_{\Gamma}^{-1} \cap (N_0 \times N_1)$, which is absurd. Thus (a) and (b) cannot hold simultaneously.

Assume now that (a) does not hold. Then there are $g, h : 2^{\omega} \to X$ continuous such that the inclusions $\mathbb{S}_0 \subseteq (g \times h)^{-1}(G)$ and $\mathbb{S}_1 \subseteq (g \times h)^{-1}(G^{-1})$ hold. It remains to set $f(0\alpha) := g(0\alpha)$ and $f(1\beta) := h(1\beta)$.

Proof of Theorem 1.14. We argue as in the proof of Theorem 1.9. The things to note are the following:

- if G is s-acyclic or locally countable, then s(G) too,

- as noted in [Lo4], if G is separable from G^{-1} by a $\text{pot}(\Gamma)$ set S, then $S^{-1} \in \text{pot}(\Gamma)$ separates G^{-1} from G, and $\neg S^{-1} \in \text{pot}(\check{\Gamma})$ separates G from G^{-1} , so that we can restrict our attention to the classes $D_{\eta}(\Sigma_{\xi}^{0})$ and Δ_{2}^{0} .

• If Γ has rank two, then Theorem 8.3 and Corollary 7.3 provide Borel relations \mathbb{S}_0 , \mathbb{S}_1 on 2^{ω} .

• If $\Gamma = D_{\eta}(\Sigma_1^0)$, then Corollaries 3.6 and 3.9 provide $f: 2^{\omega} \to X$ injective continuous such that one of the following holds:

(a) $\mathbb{N}_{0}^{\eta} \subseteq (f \times f)^{-1}(G)$ and $\mathbb{N}_{1}^{\eta} \subseteq (f \times f)^{-1}(G^{-1})$,

(b)
$$\mathbb{B}_0^\eta \subseteq (f \times f)^{-1}(G)$$
 and $\mathbb{B}_1^\eta \subseteq (f \times f)^{-1}(G^{-1})$.

The case (a) cannot happen since G^{-1} is irreflexive.

Proof of Theorem 1.15. Note first that $\mathbb{S}_0^{\eta} \cup (\mathbb{S}_1^{\eta})^{-1}, \mathbb{C}_0^{\eta} \cup (\mathbb{C}_1^{\eta})^{-1}, \mathbb{B}_0^{\eta} \cup (\mathbb{B}_1^{\eta})^{-1}$ and $\mathbb{B}_1^{\eta} \cup (\mathbb{B}_0^{\eta})^{-1}$ are Borel oriented graphs with locally countable closure. As in the proof of Theorem 1.9, \mathbb{G} is not separable from \mathbb{G}^{-1} by a pot $\left(\Delta\left(D_{\eta}(\boldsymbol{\Sigma}_1^0)\right)\right)$ set if $\mathbb{G} \in \{\mathbb{C}_0^{\eta} \cup (\mathbb{C}_1^{\eta})^{-1}, \mathbb{B}_0^{\eta} \cup (\mathbb{B}_1^{\eta})^{-1}, \mathbb{B}_1^{\eta} \cup (\mathbb{B}_0^{\eta})^{-1}\}$. By Lemma 3.1, $\mathbb{S}_0^{\eta} \cup (\mathbb{S}_1^{\eta})^{-1}$ is not separable from $(\mathbb{S}_0^{\eta})^{-1} \cup \mathbb{S}_1^{\eta}$ by a pot $\left(\Delta\left(D_{\eta}(\boldsymbol{\Sigma}_1^0)\right)\right)$ set.

• Assume now that (a) does not hold. Corollaries 4.5 and 4.7 provide

$$(\mathbb{A},\mathbb{B}) \in \{(\mathbb{N}_1^{\eta},\mathbb{N}_0^{\eta}), (\mathbb{B}_1^{\eta},\mathbb{B}_0^{\eta}), (\mathbb{N}_0^{\eta},\mathbb{N}_1^{\eta}), (\mathbb{B}_0^{\eta},\mathbb{B}_1^{\eta}), (\mathbb{S}_0^{\eta},\mathbb{S}_1^{\eta}), (\mathbb{C}_0^{\eta},\mathbb{C}_1^{\eta})\}$$

and $f: 2^{\omega} \to X$ injective continuous such that $\mathbb{A} \subseteq (f \times f)^{-1}(G)$ and $\mathbb{B} \subseteq (f \times f)^{-1}(G^{-1})$.

The pair (\mathbb{A}, \mathbb{B}) cannot be in $\{(\mathbb{N}_1^{\eta}, \mathbb{N}_0^{\eta}), (\mathbb{N}_0^{\eta}, \mathbb{N}_1^{\eta})\}$ since G and G^{-1} are irreflexive. It is enough to show the existence of $f: 2^{\omega} \to 2^{\omega}$ injective continuous such that $\mathbb{B}_0^{\eta} \cup (\mathbb{B}_1^{\eta})^{-1} \subseteq (f \times f)^{-1} (\mathbb{B}_1^{\eta} \cup (\mathbb{B}_0^{\eta})^{-1})$ to see that (b) holds.

- We use the notation of the proof of Proposition 4.4. Let us show that

$$F_{\theta}^{\operatorname{parity}(\eta)} := F_{\theta,1}^{\operatorname{parity}(\eta)} \subseteq C_{\theta}$$

if $\theta < \eta$ (where $A_{\varepsilon} = \mathbb{N}_{\varepsilon}^{\eta}$ and the closures refer to τ_1). We argue by induction on θ . Note first that

$$F_0^{\text{parity}(\eta)} = \overline{\mathbb{N}_{\text{parity}(\eta)}^{\eta}} = \overline{\bigcup_{\text{parity}(\varphi(s))=0} \operatorname{Gr}(f_s)} \subseteq \overline{C_0} = C_0,$$

by the proof of Proposition 4.4. Then, inductively,

$$F_{\theta}^{\text{parity}(\eta)} = \underbrace{\mathbb{N}_{|\text{parity}(\theta)-\text{parity}(\eta)|}^{\eta} \cap \bigcap_{\theta' < \theta} F_{\theta'}^{\text{parity}(\eta)}}_{\bigcup_{\text{parity}(\varphi(s))=\text{parity}(\theta)} \text{Gr}(f_s) \cap \bigcap_{\theta' < \theta} \bigcup_{\varphi(s) \ge \theta'} \text{Gr}(f_s) = \overline{C_{\theta}} = C_{\theta},$$

by the proof of Proposition 4.4.

- From this we deduce that $\mathbb{N}_0^\eta \cap \bigcap_{\theta < \eta} F_{\theta}^{\operatorname{parity}(\eta)}$ is contained in

$$\left(\bigcup_{\operatorname{parity}(\varphi(s))=\operatorname{parity}(\eta)}\operatorname{Gr}(f_s)\right)\cap\bigcap_{\theta<\eta}C_{\theta}\subseteq\operatorname{Gr}(f_{\emptyset})=\Delta(2^{\omega}).$$

As $\mathbb{N}_0^{\eta} \cup \mathbb{N}_1^{\eta}$ is locally countable and $\mathbb{N}_0^{\eta} \cap \bigcap_{\theta < \eta} F_{\theta}^{\operatorname{parity}(\eta)} \subseteq \Delta(2^{\omega})$, the proof of Theorem 3.3 gives $h: 2^{\omega} \to 2^{\omega}$ injective continuous such that $\mathbb{N}_0^{\eta} \subseteq (h \times h)^{-1}((\mathbb{N}_0^{\eta})^{-1})$ and $\mathbb{N}_1^{\eta} \subseteq (h \times h)^{-1}((\mathbb{N}_1^{\eta})^{-1})$ (we are in the case 2 of this proof). The map $f: \varepsilon \alpha \mapsto (1-\varepsilon)h(\alpha)$ is as desired.

• As $\Delta(2^{\omega})$ is contained in the closure of $\mathbb{S}_0^{\eta} \cup (\mathbb{S}_1^{\eta})^{-1}$, this last relation is not below the two others.

- Assume, towards a contradiction, that $\mathbb{B}_0^{\eta} \cup (\mathbb{B}_1^{\eta})^{-1}$ is below $\mathbb{S}_0^{\eta} \cup (\mathbb{S}_1^{\eta})^{-1}$. This gives $s \in 2^{<\omega}$ and $\varepsilon \in 2$ such that $(N_{0s}, N_{1s}, \mathbb{B}_0^{\eta} \cap (N_{0s} \times N_{1s}), \mathbb{B}_1^{\eta} \cap (N_{0s} \times N_{1s})) \sqsubseteq (2^{\omega}, 2^{\omega}, (\mathbb{S}_{\varepsilon}^{\eta})^{1-2\varepsilon}, (\mathbb{S}_{1-\varepsilon}^{\eta})^{1-2\varepsilon})$. By Lemma 3.1, $\mathbb{N}_0^{\eta} \cap \mathbb{N}_s^2$ is not separable from $\mathbb{N}_1^{\eta} \cap \mathbb{N}_s^2$ by a pot $(D_{\eta}(\Sigma_1^0))$ set. As $\mathbb{N}_0^{\eta} \cup \mathbb{N}_1^{\eta}$ is locally countable and $\mathbb{N}_0^{\eta} \cap \bigcap_{\theta < \eta} F_{\theta}^{\operatorname{parity}(\eta)} \subseteq \Delta(2^{\omega})$, the proof of Theorem 3.3 gives $h : 2^{\omega} \to N_s$ injective continuous such that $\mathbb{N}_{\epsilon}^{\eta} \subseteq (h \times h)^{-1}(\mathbb{N}_{\epsilon}^{\eta} \cap \mathbb{N}_s^2)$ for each $\epsilon \in 2$ (we are in the case 2 of this proof). This implies that $(2^{\omega}, 2^{\omega}, \mathbb{B}_0^{\eta}, \mathbb{B}_1^{\eta}) \subseteq (N_{0s}, N_{1s}, \mathbb{B}_0^{\eta} \cap (N_{0s} \times N_{1s}), \mathbb{B}_1^{\eta} \cap (N_{0s} \times N_{1s}))$ and

$$(2^{\omega},2^{\omega},\mathbb{B}^{\eta}_{0},\mathbb{B}^{\eta}_{1})\sqsubseteq \left(2^{\omega},2^{\omega},(\mathbb{S}^{\eta}_{\varepsilon})^{1-2\varepsilon},(\mathbb{S}^{\eta}_{1-\varepsilon})^{1-2\varepsilon}\right).$$

By Corollary 3.9, $(2^{\omega}, 2^{\omega}, \mathbb{N}^{\eta}_0, \mathbb{N}^{\eta}_1) \sqsubseteq (2^{\omega}, 2^{\omega}, \mathbb{B}^{\eta}_0, \mathbb{B}^{\eta}_1)$, so that

$$(2^{\omega}, 2^{\omega}, \mathbb{N}^{\eta}_{0}, \mathbb{N}^{\eta}_{1}) \sqsubseteq (2^{\omega}, 2^{\omega}, (\mathbb{S}^{\eta}_{\varepsilon})^{1-2\varepsilon}, (\mathbb{S}^{\eta}_{1-\varepsilon})^{1-2\varepsilon}).$$

But this contradicts the proof of Proposition 4.4.

- We will show that $(2^{\omega}, 2^{\omega}, \mathbb{C}_0^{\eta}, \mathbb{C}_1^{\eta}) \sqsubseteq (2^{\omega}, 2^{\omega}, \mathbb{S}_0^{\eta}, \mathbb{S}_1^{\eta})$. Using the proof of the previous point, this will show that $\mathbb{B}_0^{\eta} \cup (\mathbb{B}_1^{\eta})^{-1}$ is not below $\mathbb{C}_0^{\eta} \cup (\mathbb{C}_1^{\eta})^{-1}$.

We use the notation of the proof of Proposition 4.4. Let us show that $G_{\theta} := G_{\theta,1} \subseteq C_{\theta}$ if $1 \le \theta \le \eta$ (where $A_{\varepsilon} = \mathbb{S}_{\varepsilon}^{\eta}$ and the closures refer to τ_1). We argue by induction on θ . Note first that

$$G_1 = \overline{\mathbb{S}_0^{\eta}} \cap \overline{\mathbb{S}_1^{\eta}} = \overline{U_0^0} \cap \overline{U_0^1} = C_1^0 \cup C_1^1 = C_1^0$$

by the proof of Proposition 4.4. Then, inductively,

$$G_{\theta+1} = \overline{\mathbb{S}_0^{\eta} \cap G_{\theta}} \cap \overline{\mathbb{S}_1^{\eta} \cap G_{\theta}} \subseteq \overline{U_0^0 \cap C_{\theta}} \cap \overline{U_0^1 \cap C_{\theta}} \subseteq C_{\theta+1}$$

and $G_{\lambda} = \bigcap_{\theta < \lambda} G_{\theta} \subseteq \bigcap_{\theta < \lambda} C_{\theta} = C_{\lambda}$ if λ is limit.

From this we deduce that $G_{\eta} \subseteq C_{\eta} = \operatorname{Gr}(f_{\emptyset}) = \Delta(2^{\omega})$. As $\mathbb{S}_{0}^{\eta} \cup \mathbb{S}_{1}^{\eta}$ is locally countable and $G_{\eta} \subseteq \Delta(2^{\omega})$, the proof of Theorem 4.3 gives $h: 2^{\omega} \to N_{s}$ injective continuous such that the inclusion $\mathbb{S}_{\epsilon}^{\eta} \subseteq (h \times h)^{-1}(\mathbb{S}_{\epsilon}^{\eta} \cap N_{0}^{2})$ holds for each $\epsilon \in 2$ (we are in the case 2 of this proof). The maps defined by $f(0\alpha) := h(\alpha)$, $f(1\alpha) := 1\alpha$, $g(1\beta) := h(\beta)$ and $g(0\beta) := 1\beta$, are as desired.

- Assume, towards a contradiction, that $\mathbb{C}_0^{\eta} \cup (\mathbb{C}_1^{\eta})^{-1}$ is below $\mathbb{S}_0^{\eta} \cup (\mathbb{S}_1^{\eta})^{-1}$, with witness f. This gives $s \in 2^{<\omega} \setminus \{\emptyset\}$ and $\varepsilon \in 2$ such that $\mathbb{C}_{\epsilon}^{\eta} \cap (N_{0s} \times N_{1s}) \subseteq (f \times f)^{-1} ((\mathbb{S}_{|\epsilon-\varepsilon|}^{\eta})^{1-2\varepsilon})$ for each $\epsilon \in 2$. As in the previous point, there is $h: 2^{\omega} \to N_s$ injective continuous such that

$$\mathbb{S}^{\eta}_{\epsilon} \subseteq (h \times h)^{-1} (\mathbb{S}^{\eta}_{\epsilon} \cap N_s^2)$$

for each $\epsilon \in 2$. This implies that if we set $k(\epsilon \alpha) := \epsilon h(\alpha)$ and $l := f \circ k$, then

$$\mathbb{C}^{\eta}_{\epsilon} \subseteq (k \times k)^{-1} \big(\mathbb{C}^{\eta}_{\epsilon} \cap (N_{0s} \times N_{1s}) \big)$$

and $\mathbb{C}^{\eta}_{\epsilon} \subseteq (l \times l)^{-1} ((\mathbb{S}^{\eta}_{|\epsilon-\varepsilon|})^{1-2\varepsilon})$. As in the proof of Proposition 4.4, we see that the image of

$$\{(0\alpha, 1\alpha) \mid \alpha \in 2^{\omega}\}$$

by $l \times l$ is contained in the diagonal of 2^{ω} , which is not possible by injectivity of l.

- Assume that η is a successor ordinal. The previous points show that if $\mathbb{C}_0^\eta \cup (\mathbb{C}_1^\eta)^{-1}$ is below $\mathbb{B}_0^\eta \cup (\mathbb{B}_1^\eta)^{-1}$, then $(2^\omega, 2^\omega, \mathbb{C}_0^\eta, \mathbb{C}_1^\eta) \sqsubseteq (2^\omega, 2^\omega, (\mathbb{B}_{\varepsilon}^\eta)^{1-2\varepsilon}, (\mathbb{B}_{1-\varepsilon}^\eta)^{1-2\varepsilon})$ for some $\varepsilon \in 2$. We saw that there is $h: 2^\omega \to N_0$ injective continuous such that $\mathbb{N}_{\varepsilon}^\eta \subseteq (h \times h)^{-1}(\mathbb{N}_{\varepsilon}^\eta \cap N_0^2)$ for each $\varepsilon \in 2$. The maps defined by $f(0\alpha) := h(\alpha)$, $f(1\alpha) := 1\alpha$, $g(1\beta) := h(\beta)$ and $g(0\beta) := 1\beta$ are witnesses for the fact that $(2^\omega, 2^\omega, \mathbb{B}_0^\eta, \mathbb{B}_1^\eta) \sqsubseteq (2^\omega, 2^\omega, \mathbb{N}_0^\eta, \mathbb{N}_1^\eta)$, so that $(2^\omega, 2^\omega, \mathbb{C}_0^\eta, \mathbb{C}_1^\eta) \sqsubseteq (2^\omega, 2^\omega, (\mathbb{N}_{\varepsilon}^\eta)^{1-2\varepsilon}, (\mathbb{N}_{1-\varepsilon}^\eta)^{1-2\varepsilon})$. The maps $\alpha \mapsto 0\alpha$ and $\beta \mapsto 1\beta$ are witnesses for the fact that $(2^\omega, 2^\omega, \mathbb{S}_0^\eta, \mathbb{S}_1^\eta) \sqsubseteq (2^\omega, 2^\omega, \mathbb{C}_0^\eta, \mathbb{C}_1^\eta)$. Thus $(2^\omega, 2^\omega, \mathbb{S}_0^\eta, \mathbb{S}_1^\eta) \sqsubseteq (2^\omega, 2^\omega, (\mathbb{N}_{\varepsilon}^\eta)^{1-2\varepsilon}, (\mathbb{N}_{1-\varepsilon}^\eta)^{1-2\varepsilon})$, which contradicts the proof of Proposition 4.4.

- Assume that η is a limit ordinal. Let us show that $\mathbb{C}_0^{\eta} \cup (\mathbb{C}_1^{\eta})^{-1}$ is below $\mathbb{B}_0^{\eta} \cup (\mathbb{B}_1^{\eta})^{-1}$. The proof of Proposition 4.4 provides $h: 2^{\omega} \to 2^{\omega}$ injective continuous such that $\mathbb{S}_{\varepsilon}^{\eta} \subseteq (h \times h)^{-1}(\mathbb{N}_{\varepsilon}^{\eta})$ for each $\varepsilon \in 2$. It remains to set $f(\varepsilon \alpha) := \varepsilon h(\alpha)$.

10 Negative results

- By Theorem 15 in [L4], we cannot completely remove the assumption that A is s-acyclic or locally countable in Corollary 6.4. We can wonder whether there is an antichain basis if this assumption is removed (for this class Π_2^0 or any other one appearing in this section). This also shows that we cannot simply assume the disjointness of the analytic sets A, B in Theorem 6.3 and Corollaries 6.5, 6.7.

- We can use the proof of the previous fact to get a negative result for the class Δ_2^0 .

Theorem 10.1 There is no tuple (X, Y, A, B), where X, Y are Polish and A, B are disjoint analytic subsets of $X \times Y$, such that for any tuple $(\mathcal{X}, \mathcal{Y}, \mathcal{A}, \mathcal{B})$ of this type, exactly one of the following holds:

- (a) \mathcal{A} is separable from \mathcal{B} by a pot $(\mathbf{\Delta}_2^0)$ set,
- (b) $(\mathbb{X}, \mathbb{Y}, \mathbb{A}, \mathbb{B}) \sqsubseteq (\mathcal{X}, \mathcal{Y}, \mathcal{A}, \mathcal{B}).$

Proof. We argue by contradiction. By Lemma 7.1, we get $(\mathbb{X}, \mathbb{Y}, \mathbb{A}, \mathbb{B}) \sqsubseteq (2^{\omega}, 2^{\omega}, \lceil T \rceil \cap \mathbb{E}_0^0, \lceil T \rceil \cap \mathbb{E}_0^1)$. This shows that \mathbb{A}, \mathbb{B} are locally countable. As (a) and (b) cannot hold simultaneously, \mathbb{A} is not separable from \mathbb{B} by a pot $(\mathbf{\Delta}_2^0)$ set. By Corollary 7.4 we get

$$(2^{\omega}, 2^{\omega}, [T] \cap \mathbb{E}^0_0, [T] \cap \mathbb{E}^1_0) \sqsubseteq (\mathbb{X}, \mathbb{Y}, \mathbb{A}, \mathbb{B}),$$

so that we may assume that $(\mathbb{X}, \mathbb{Y}, \mathbb{A}, \mathbb{B}) = (2^{\omega}, 2^{\omega}, [T] \cap \mathbb{E}^0_0, [T] \cap \mathbb{E}^1_0)$.

• In the proof of Theorem 15 in [L4], the author considers a set $A = \bigcup_{s \in (\omega \setminus \{0\}) \le \omega} \operatorname{Gr}(l_{s|G})$, where the l_s 's are partial continuous open maps from 2^{ω} into itself with dense open domain, and G is the intersection of their domain. Moreover, the l_s 's have the properties that $l_s(x) \neq l_t(x)$ if $t \neq s$, and $l_s(x)$ is the limit of $(l_{sk}(x))_{k \in \omega}$, for each $x \in G$. We set, for $\varepsilon \in 2$, $A_{\varepsilon} := \bigcup_{s \in (\omega \setminus \{0\}) \le \omega} \operatorname{Gr}(l_s|_G)$, so that A_0 and A_1 are disjoint Borel sets.

Let us check that A_0 is not separable from A_1 by a $\text{pot}(\Delta_2^0)$ set. We argue by contradiction, which gives $D \in \text{pot}(\Delta_2^0)$ and a dense G_{δ} subset H of 2^{ω} such that $D \cap H^2 \in \Delta_2^0(H^2)$. We may assume that $H \subseteq G$. Note that $H \cap \bigcap_{s \in (\omega \setminus \{0\}) \le \omega} l_s^{-1}(H)$ is a dense G_{δ} subset of 2^{ω} , and thus contains a point x. The vertical section A_x is contained in H. In particular, the disjoint sections $(A_0)_x$ and $(A_1)_x$ are separable by a Δ_2^0 subset \mathcal{D} of the Polish space H. It remains to note that $\mathcal{D} \cap \overline{A_x}^H$ is a dense and co-dense Δ_2^0 subset of $\overline{A_x}^H$, which contradicts Baire's theorem.

This gives $u: N_0 \to 2^{\omega}$ and $v: N_1 \to 2^{\omega}$ with $[T] \cap \mathbb{E}_0^{\varepsilon} \subseteq (u \times v)^{-1}(A_{\varepsilon})$.

• We set $B_1 := [T] \cap (\mathbb{E}_0^0 \cup \mathbb{E}_0^1)$. Note that $B_1 \notin \text{pot}(G_\delta)$, since otherwise $[T] \cap \mathbb{E}_0^0$ and $[T] \cap \mathbb{E}_0^1$ are two disjoint $\text{pot}(G_\delta)$ sets, and thus $\text{pot}(\mathbf{\Delta}_2^0)$ -separable. Then we can follow the proof of Theorem 15 in [L4]. This proof gives $U: F \to G$ and $V: F \to 2^{\omega}$ injective continuous satisfying the inclusion $\bigcup_{n \in \omega} \operatorname{Gr}(f_n) \subseteq (U \times V)^{-1}(A)$.

The only thing to check is that there is (c,d) in $\bigcup_{n\in\omega} \omega^n \times \omega^{n+1}$ and a nonempty open subset R of $D_{f_{c,d}}$ such that $(U(x), V(f_{c,d}(x))) \notin \operatorname{Gr}(l_{\emptyset})$ for each $x \in R$. We argue by contradiction, which gives a dense G_{δ} subset K of F such that $\bigcup_{n\in\omega} \operatorname{Gr}(f_{n|K}) \subseteq (U_{|K} \times V)^{-1}(\operatorname{Gr}(l_{\emptyset|G}))$. As $(U_{|K} \times V)^{-1}(\operatorname{Gr}(l_{\emptyset|G}))$ is the graph of a partial Borel map, $\bigcup_{n\in\omega} \operatorname{Gr}(f_{n|K})$ too. Therefore $\bigcup_{n\in\omega} \operatorname{Gr}(f_{n|K}) \in \operatorname{pot}(\mathbf{\Pi}_1^0) \setminus \operatorname{pot}(G_{\delta})$, which is absurd. This shows that we cannot completely remove the assumption that $A \cup B$ is s-acyclic or locally countable in Corollary 7.3. This also shows that we cannot simply assume the disjointness of the analytic sets A, B in Theorem 7.2 and Corollary 7.4.

- By Theorem 2.16 in [L3], we cannot completely remove the assumption that $A \cup B$ is s-acyclic or locally countable in Corollary 3.10. This also shows that we cannot simply assume disjointness in Theorem 3.3 and Corollary 3.11.

We saw that there is a version of Corollary 6.7 for $\Gamma = \Sigma_1^0$, where we replace the class F_{σ} with the class of open sets. We cannot replace the class F_{σ} with the class of closed sets.

Proposition 10.2 There is no triple (X, A, B), where X is Polish and A, B are disjoint analytic relations on X such that A is contained in a potentially closed s-acyclic or locally countable relation such that, for each triple (X, A, B) of the same type, exactly one of the following holds:

- (a) the set \mathcal{A} is separable from \mathcal{B} by a pot (Σ_1^0) set,
- $(b) (\mathbb{X}, \mathbb{X}, \mathbb{A}, \mathbb{B}) \sqsubseteq (\mathcal{X}, \mathcal{X}, \mathcal{A}, \mathcal{B}).$

Proof. We argue by contradiction, which gives a triple. Note that \mathbb{A} is not separable from \mathbb{B} by a pot (Σ_1^0) set. Theorem 9 in [L5] gives $F, G: 2^{\omega} \to \mathbb{X}$ continuous such that $\Delta(2^{\omega}) \subseteq (F \times G)^{-1}(\mathbb{A})$ and $\mathbb{G}_0 \subseteq (F \times G)^{-1}(\mathbb{B})$. We set $\mathbb{A}' := (F \times G)[\Delta(2^{\omega})]$, $\mathbb{B}' := (F \times G)[\mathbb{G}_0]$ and $\mathbb{C}' := (F \times G)[\overline{\mathbb{G}_0}]$. Note that \mathbb{A}', \mathbb{C}' are compact and \mathbb{C}' is the locally countable disjoint union of \mathbb{A}' and \mathbb{B}' . In particular, \mathbb{B}' is $D_2(\Sigma_1^0)$, $\mathbb{A}' \subseteq \mathbb{A}, \mathbb{B}' \subseteq \mathbb{B}$, and \mathbb{A}' is not separable from \mathbb{B}' by a pot (Σ_1^0) set. So we may assume that \mathbb{A}, \mathbb{B} are Borel with locally countable union which is the closure of \mathbb{B} . Corollary 3.10 gives $f', g': 2^{\omega} \to \mathbb{X}$ injective continuous such that $\mathbb{G}_0 = \overline{\mathbb{G}_0} \cap (f' \times g')^{-1}(\mathbb{B})$. In particular,

$$\Delta(2^{\omega}) \subseteq (f' \times g')^{-1}(\overline{\mathbb{B}} \setminus \mathbb{B}) = (f' \times g')^{-1}(\mathbb{A}).$$

This means that we may assume that $\mathbb{X} = 2^{\omega}$, $\mathbb{A} = \Delta(2^{\omega})$ and $\mathbb{B} = \mathbb{G}_0$.

The proof of Theorem 10 in [L5] provides a Borel graph \mathcal{B} on $X := 2^{\omega}$ with no Borel countable coloring such that any locally countable Borel digraph contained in \mathcal{B} has a Borel countable coloring. Consider the closed symmetric acyclic locally countable relation $\mathcal{A} := \Delta(2^{\omega})$. As there is no Borel countable coloring of \mathcal{B} , \mathcal{A} is not separable from \mathcal{B} by a pot (Σ_1^0) set. If f, g exist, then f = g since \mathbb{A} is contained in $(f \times g)^{-1}(\mathcal{A})$. This implies that f is a homomorphism from \mathbb{G}_0 into \mathcal{B} . The digraph $(f \times f)[\mathbb{G}_0]$ is locally countable and Borel since f is injective. Thus it has a Borel countable coloring, and \mathbb{G}_0 too, which is absurd.

For oriented graphs, we cannot completely remove the assumption that G is s-acyclic or locally countable in Theorem 1.14. Let us check it for $\Gamma = \Delta_2^0$.

Proposition 10.3 *There is no tuple* (X, \mathbb{G}) *, where* X *is Polish and* \mathbb{G} *is an analytic oriented graph on* X*, such that for any tuple* $(\mathcal{X}, \mathcal{G})$ *of this type, exactly one of the following holds:*

- (a) the set \mathcal{G} is separable from \mathcal{G}^{-1} by a pot $(\mathbf{\Delta}_2^0)$ set,
- (b) there is $f: 2^{\omega} \to X$ injective continuous such that $\mathbb{G} \subseteq (f \times f)^{-1}(\mathcal{G})$.

Proof. We use the notation of the proof of Theorem 10.1, and argue by contradiction. Recall the analytic s-acyclic oriented graph $\mathcal{G}_{\Delta_2^0} = (\lceil T \rceil \cap \mathbb{E}_0^0) \cup (\lceil T \rceil \cap \mathbb{E}_0^1)^{-1}$ considered in the proof of Theorem 1.14. Note that there is $f_0: \mathbb{X} \to 2^{\omega}$ injective continuous such that $\mathbb{G} \subseteq (f_0 \times f_0)^{-1}(\mathcal{G}_{\Delta_2^0})$. In particular, \mathbb{G} is s-acyclic and Theorem 1.14 applies. This shows that we may assume that $(\mathbb{X}, \mathbb{G}) = (2^{\omega}, \mathcal{G}_{\Delta_2^0})$.

If R is a relation on 2^{ω} , then we set $G_R := \{(0\alpha, 1\beta) \mid (\alpha, \beta) \in R\}$. As A_0 is not separable from A_1 by a pot (Δ_2^0) set, G_{A_0} is not separable from G_{A_1} by a pot (Δ_2^0) set. As $G_{A_0} \cup G_{A_1} \subseteq N_0 \times N_1$ and G_{A_0}, G_{A_1} are disjoint, $\mathbb{H} := G_{A_0} \cup (G_{A_1})^{-1}$ is a Borel oriented graph, and \mathbb{H} is not separable from \mathbb{H}^{-1} by a pot (Δ_2^0) set, as in the proof of Theorem 1.9. If $f : 2^{\omega} \to 2^{\omega}$ is injective continuous and $(\lceil T \rceil \cap \mathbb{E}_0^0) \cup (\lceil T \rceil \cap \mathbb{E}_0^1)^{-1} \subseteq \mathbb{H}$, then on a nonempty clopen set $S := N_{s_q} \times N_{t_q}$, the first coordinate is either preserved, or changed.

As in the proof of Lemma 7.1, we see that $[T] \cap \mathbb{E}_0^0 \cap S$ is not separable from $[T] \cap \mathbb{E}_0^1 \cap S$ by a pot $(\mathbf{\Delta}_2^0)$ set. By Corollary 7.3, there is $f: 2^{\omega} \to 2^{\omega}$ injective continuous such that

$$[T] \cap \mathbb{E}_0^{\varepsilon} \subseteq (f \times f)^{-1} ([T] \cap \mathbb{E}_0^{\varepsilon} \cap S)$$

for each $\varepsilon \in 2$. This proves the existence of $g: 2^{\omega} \to 2^{\omega}$ injective continuous such that

$$\lceil T \rceil \cap (\mathbb{E}^0_0 \cup \mathbb{E}^1_0) \subseteq (g \times g)^{-1}(G_A).$$

This gives $u: N_0 \to 2^{\omega}$ and $v: N_1 \to 2^{\omega}$ injective continuous such that $[T] \cap (\mathbb{E}^0_0 \cup \mathbb{E}^1_0) \subseteq (u \times v)^{-1}(A)$ since the maps $\varepsilon \alpha \mapsto \alpha$ are injective. But we saw that this is not possible in the proof of Theorem 10.1.

Question. Are there versions of our results for the classes $D_{\eta}(\Sigma_2^0)$, $\check{D}_{\eta}(\Sigma_2^0)$ (when $\omega \leq \eta < \omega_1$) and $\Delta(D_{\eta}(\Sigma_2^0))$ (when $2 \leq \eta < \omega_1$)?

11 References

[D-SR] G. Debs and J. Saint Raymond, Borel liftings of Borel sets: some decidable and undecidable statements, *Mem. Amer. Math. Soc.* 187, 876 (2007)

[K] A. S. Kechris, Classical descriptive set theory, Springer-Verlag, 1995

[K-S-T] A. S. Kechris, S. Solecki and S. Todorčević, Borel chromatic numbers, *Adv. Math.* 141 (1999), 1-44

[L1] D. Lecomte, Classes de Wadge potentielles et théorèmes d'uniformisation partielle, *Fund. Math.* 143 (1993), 231-258

[L2] D. Lecomte, Uniformisations partielles et critères à la Hurewicz dans le plan, *Trans. Amer. Math. Soc.* 347, 11 (1995), 4433-4460

[L3] D. Lecomte, Tests à la Hurewicz dans le plan, Fund. Math. 156 (1998), 131-165

[L4] D. Lecomte, Complexité des boréliens à coupes dénombrables, *Fund. Math.* 165 (2000), 139-174

[L5] D. Lecomte, On minimal non potentially closed subsets of the plane, *Topology Appl.* 154, 1 (2007), 241-262

[L6] D. Lecomte, How can we recognize potentially Π_{ξ}^{0} subsets of the plane?, *J. Math. Log.* 9, 1 (2009), 39-62

[L7] D. Lecomte, A dichotomy characterizing analytic graphs of uncountable Borel chromatic number in any dimension, *Trans. Amer. Math. Soc.* 361 (2009), 4181-4193

[L8] D. Lecomte, Potential Wadge classes, Mem. Amer. Math. Soc., 221, 1038 (2013)

[Lo1] A. Louveau, Some results in the Wadge hierarchy of Borel sets, *Cabal Sem. 79-81, Lect. Notes in Math.* 1019 (1983), 28-55

[Lo2] A. Louveau, A separation theorem for Σ_1^1 sets, *Trans. A. M. S.* 260 (1980), 363-378

[Lo3] A. Louveau, Ensembles analytiques et boréliens dans les espaces produit, *Astérisque (S. M. F.)* 78 (1980)

[Lo4] A. Louveau, Some dichotomy results for analytic graphs, manuscript

[Lo-SR] A. Louveau and J. Saint Raymond, The strength of Borel Wadge determinacy, *Cabal Seminar 81-85, Lecture Notes in Math.* 1333 (1988), 1-30

[M] Y. N. Moschovakis, *Descriptive set theory*, North-Holland, 1980