TOPOLOGICAL EMBEDDINGS INTO RANDOM 2-COMPLEXES

MICHAEL FARBER AND TAHL NOWIK

ABSTRACT. We consider 2-dimensional random simplicial complexes Y in the multi-parameter model. We establish the multi-parameter threshold for the property that every 2-dimensional simplicial complex S admits a topological embedding into Y asymptotically almost surely. Namely, if in the procedure of the multi-parameter model, each *i*-dimensional simplex is taken independently with probability $p_i = p_i(n)$, from a set of n vertices, then the threshold is $p_0 p_1^3 p_2^2 = \frac{1}{n}$. This threshold happens to coincide with the previously established thresholds for uniform hyperbolicity and triviality of the fundamental group.

Our claim in one direction is in fact slightly stronger, namely, we show that if $p_0 p_1^3 p_2^2$ is sufficiently larger than $\frac{1}{n}$ then every S has a fixed subdivision S' which admits a simplicial embedding into Y asymptotically almost surely. The main geometric result we prove to this end is that given $\epsilon > 0$, there is a subdivision S' of S such that every subcomplex $T \subseteq S'$ has $\frac{f_0(T)}{f_1(T)} > \frac{1}{3} - \epsilon$ and $\frac{f_0(T)}{f_2(T)} > \frac{1}{2} - \epsilon$, where $f_i(T)$ denotes the number of simplices in T of dimension i.

In the other direction we show that if $p_0 p_1^3 p_2^2$ is sufficiently smaller than $\frac{1}{n}$, then asymptotically almost surely, the torus does not admit a topological embedding into Y. Here we use a result of Z. Gao which bounds the number of different triangulations of a surface.

1. INTRODUCTION

The theory of random graphs is widely used for modeling large systems in various scientific applications. It has been recently realized that the higher dimensional generalization, namely, large random simplicial complexes, offers even greater opportunities for mathematical modeling as it allows taking into account multiple interactions and not only pairwise relationships. The interest in *large* complexes is manifested by studying the *asymptotic* behavior of the random complexes, as the number of vertices grows.

The first model of random simplicial complexes was studied by Linial and Meshulam [LM] and Meshulam and Wallach [MW]. This model for simplicial complexes of dimension r assumes that for a given vertex set, all simplices of dimension less than r are present, and the top dimensional

Date: December 9, 2019.

Michael Farber was partially supported by a grant from the Leverhulme Foundation.

simplices are added independently with a given probability p. For the study of asymptotic properties, the probability p is assumed to depend on the number n of vertices. Many authors have contributed to the development of the Linial-Meshulam model by computing thresholds for various topological properties such as vanishing of the Betti numbers, simple connectivity, asphericity, and cohomological dimension of the fundamental group.

In [CF1]–[CF3] Costa and Farber developed a more general model of random r-dimensional simplicial complexes. In this model a random simplicial complex depends on a string of probability parameters (p_0, p_1, \ldots, p_r) , one for each dimension. In this multi-parameter setting, the thresholds for topological properties are not numbers depending on n, but rather boundaries of domains depending on n. In [CF1]–[CF3] domains corresponding to different asymptotic properties of the fundamental group are described. Another major result of [CF1]–[CF3] is the notion of a *critical dimension* which fully portrays the properties of the Betti numbers in the multi-parameter model. The Linial-Meshulam model can be considered as a special case of the multi-parameter model by taking $p_0 = p_1 = \cdots = p_{r-1} = 1$.

In [FMN], the authors studied an even more general model of random simplicial complexes, which is of two types, the *lower* and *upper* models, as described in Section 2. The lower and upper models are Spanier-Whitehead dual to each other, and the multi-parameter model mentioned above is a special case of the lower model. The notion of critical dimension for the upper model is also developed in [FMN].

In the present work we consider random 2-dimensional simplicial complexes Y in the multiparameter lower model. We find the threshold for the following property: Every 2-dimensional simplicial complex S asymptotically almost surely admits a topological embedding into Y. We prove that the threshold for this property is $p_0 p_1^3 p_2^2 = \frac{1}{n}$, that is, if $p_0 p_1^3 p_2^2$ is sufficiently larger than $\frac{1}{n}$, namely, if there is an $\epsilon > 0$ such that $n p_0 p_1^{3+\epsilon} p_2^{2+\epsilon} \to \infty$, then this property holds, whereas if $p_0 p_1^3 p_2^2$ is sufficiently smaller than $\frac{1}{n}$, namely, if $n p_0 p_1^3 p_2^2 \to 0$, then the property does not hold.

The first half of this theorem generalizes a result of [CCFK] regarding the Linial-Meshulam model, where it is proved for $p_2 \ge n^{-\frac{1}{2}+\epsilon}$. This was strengthened by Gundert and Wagner in [GW], showing it for $p_2 \ge cn^{-\frac{1}{2}}$. Versions of this result for higher dimensional complexes are also developed in [GW]. We also mention that our above threshold also happens to be the threshold for both uniform hyperbolicity and triviality of the fundamental group, as shown in [CF2].

2. Background

In this section we describe our model for random 2-dimensional simplicial complexes. Let Δ_n^2 denote the full 2-dimensional simplicial complex with vertex set $[n] = \{1, \ldots, n\}$, i.e. Δ_n^2 includes all subsets of [n] of size 1,2,3. They are regarded as simplices of dimension 0,1,2, respectively. A probability triple \mathfrak{p} is a triple of sequences $(p_0(n), p_1(n), p_2(n))$ with $0 \leq p_i(n) \leq 1$. We will however always suppress the argument n from the expression $p_i(n)$ and simply write p_i . A given argument n will always be implied by the context. A natural example for a probability triple is obtained as follows. Given a fixed $\alpha = (\alpha_0, \alpha_1, \alpha_2)$ with $\alpha_i \geq 0$, let \mathfrak{p}_{α} be the probability triple given by $\mathfrak{p}_{\alpha} = (n^{-\alpha_0}, n^{-\alpha_1}, n^{-\alpha_2})$.

We use a probability triple \mathfrak{p} to define a sequence $(\Omega_n, \mathbb{P}_{\mathfrak{p}})$ of probability spaces, where Ω_n is the set of all simplicial subcomplexes of Δ_n^2 , and the probability measure $\mathbb{P}_{\mathfrak{p}}$ on Ω_n is defined as follows. As an intermediate step we randomly select a subset $X \subseteq \Delta_n^2$ which is not necessarily a simplicial complex. Each *i*-dimensional simplex $\sigma \in \Delta_n^2$ is included in X independently with probability p_i , i.e. the probability for obtaining X is $\prod_{\sigma \in X} p_{\dim \sigma} \prod_{\sigma \notin X} q_{\dim \sigma}$, where $q_i = 1 - p_i$. Now, the simplicial complex produced by this random process is the maximal simplicial complex Y contained in X. It is shown in [CF1] and [FMN] that the probability for obtaining any given simplicial complex $Y \subseteq \Delta_n^2$ by this random process is

$$\mathbb{P}_{\mathfrak{p}}(Y) = \prod_{\sigma \in Y} p_{\dim \sigma} \prod_{\sigma \in E(Y)} q_{\dim \sigma}$$

where $E(Y) = \{ \sigma \in \Delta_n^2 : \sigma \notin Y \text{ but } \partial \sigma \subseteq Y \}$. Indeed, Y is the maximal simplicial complex contained in X iff every $\sigma \in Y$ is in X and every $\sigma \in E(Y)$ is not in X. To see this, note that if $\sigma \in E(Y)$, then $Y \cup \{\sigma\}$ is also a simplicial complex. Also note that if $\sigma \notin Y$, then a minimal $\tau \subseteq \sigma$ such that $\tau \notin Y$, is in E(Y), (including when τ is a vertex). When we write $Y \in (\Omega_n, \mathbb{P}_p)$ we will mean that Y is randomly drawn from Ω_n using the probability measure \mathbb{P}_p .

In [FMN] this model is termed the *lower model* for random simplicial complexes, in contrast to the *upper model*, which looks at the *minimal* simplicial complex *containing* X. The setting in [FMN] is more general, as the dimension of the simplices σ is not restricted and each simplex σ is assigned its own probability p_{σ} .

For a 2-complex S let $f_i(S)$, i = 0, 1, 2, denote the number of simplices in S of dimension i (for us simplicial complexes are *finite* by definition). In this work we will apply Theorem 1(B) of [CF2] which states the following. If \mathfrak{p} satisfies that for every subcomplex T of S

$$\lim_{n \to \infty} n^{f_0(T)} p_0^{f_0(T)} p_1^{f_1(T)} p_2^{f_2(T)} = \infty,$$

then S is simplicially embeddable into $Y \in (\Omega_n, \mathbb{P}_p)$ a.a.s. (asymptotically almost surely, i.e. with probability converging to 1 as $n \to \infty$). In [CF2] the corresponding claim for complexes of general dimension r is proved.

3. Statement of result

In this work we look at random 2-complexes $Y \in (\Omega_n, \mathbb{P}_p)$ for a given probability triple $\mathfrak{p} = (p_0, p_1, p_2)$, and ask whether every 2-complex S is a.a.s. topologically embeddable into Y. We show that probability triples for which $p_0 p_1^3 p_2^2 = \frac{1}{n}$ are the threshold for this property. That is, triples \mathfrak{p} for which $p_0 p_1^3 p_2^2$ is sufficiently larger than $\frac{1}{n}$ have the property that every 2-complex S is a.a.s. topologically embeddable into $Y \in (\Omega_n, \mathbb{P}_p)$, and triples \mathfrak{p} for which $p_0 p_1^3 p_2^2$ is sufficiently smaller than $\frac{1}{n}$ have the property that every 2-complex S is a.a.s. topologically embeddable into $Y \in (\Omega_n, \mathbb{P}_p)$, and triples \mathfrak{p} for which $p_0 p_1^3 p_2^2$ is sufficiently smaller than $\frac{1}{n}$ do not have this property. The precise statement of this claim appears in Theorem 3.1 below. It follows that for probability triples of the form \mathfrak{p}_{α} the plane $\alpha_0 + 3\alpha_1 + 2\alpha_2 = 1$ is the threshold for this property. This is stated in Corollary 3.2.

Theorem 3.1. Let $\mathfrak{p} = (p_0, p_1, p_2)$ be a probability triple.

(1) If

$$\lim_{n \to \infty} n p_0 p_1^{3+\epsilon} p_2^{2+\epsilon} = \infty$$

for some fixed $\epsilon > 0$, then every 2-dimensional simplicial complex S has a simplicial subdivision S' which is a.a.s. simplicially embeddable into $Y \in (\Omega_n, \mathbb{P}_p)$. In particular, S is a.a.s. toplogically embeddable into $Y \in (\Omega_n, \mathbb{P}_p)$.

(2) If

$$\lim_{n \to \infty} n p_0 p_1^3 p_2^2 = 0$$

then the torus is a.a.s. not topologically embeddable into $Y \in (\Omega_n, \mathbb{P}_p)$.

Corollary 3.2. Let $\alpha = (\alpha_0, \alpha_1, \alpha_2)$ with $\alpha_i \ge 0$ and let $\mathfrak{p}_{\alpha} = (n^{-\alpha_0}, n^{-\alpha_1}, n^{-\alpha_2})$.

- If α₀ + 3α₁ + 2α₂ < 1 then every 2-dimensional simplicial complex S has a simplicial subdivision S' which is a.a.s. simplicially embeddable into Y ∈ (Ω_n, ℙ_{p_α}). In particular, S is a.a.s. toplogically embeddable into Y ∈ (Ω_n, ℙ_{p_α}).
- (2) If $\alpha_0 + 3\alpha_1 + 2\alpha_2 > 1$ then the torus is a.a.s. not topologically embeddable into $Y \in (\Omega_n, \mathbb{P}_{\mathfrak{p}_\alpha})$.



FIGURE 1. $\Delta \longrightarrow$ two depictions of $\mathfrak{s}^1 \Delta \longrightarrow \mathfrak{s}^2 \Delta \longrightarrow \mathfrak{s}^3 \Delta$

The main effort of this work is the proof of Proposition 3.3 below, for which we use a particular subdivision scheme for 2-complexes which is suitable for our purposes. The first subdivision is barycentric, but all subsequent subdivisions divide each triangle into four triangles as shown in Figure 1. Geometrically, we view the kth subdivision $\mathfrak{s}^k \Delta$ of a 2-simplex Δ as a regular hexagon divided into $6 \cdot 4^{k-1}$ equilateral triangles.

For a 2-complex S let $f_i(S)$, i = 0, 1, 2, denote the number of simplices in S of dimension i, and define $\mu_i(S) = \frac{f_0(S)}{f_i(S)}$, i = 1, 2, whenever $f_i(S) \neq 0$.

Proposition 3.3. Let S be a 2-complex, and let $\epsilon > 0$. Then for sufficiently large k, every subcomplex T of $\mathfrak{s}^k S$ has

(1) $\mu_1(T) > \frac{1}{3} - \epsilon.$ (2) $\mu_1(T) > \frac{1}{3} - \epsilon.$

(2)
$$\mu_2(T) > \frac{1}{2} - \epsilon$$

The proofs of parts (1) and (2) of Proposition 3.3 appear in Sections 6 and 5 respectively. Once Proposition 3.3 is established, the proof of Theorem 3.1(1) is immediate. Indeed, by Proposition 3.3, given $\epsilon > 0$ there exists a subdivision S' of S such that every subcomplex T of S' satisfies $\mu_1(T) > \frac{1}{3+\epsilon}$ and $\mu_2(T) > \frac{1}{2+\epsilon}$. For every subcomplex T of S' we have

$$n^{f_0(T)} p_0^{f_0(T)} p_1^{f_1(T)} p_2^{f_2(T)} = \left(n p_0 p_1^{\frac{1}{\mu_1(T)}} p_2^{\frac{1}{\mu_2(T)}} \right)^{f_0(T)} \ge \left(n p_0 p_1^{3+\epsilon} p_2^{2+\epsilon} \right)^{f_0(T)} \xrightarrow[n \to \infty]{} \infty$$

By Theorem 1(B) of [CF2] stated in the previous section, S' is a.a.s. simplically embeddable into $Y \in (\Omega, \mathbb{P}_p)$, completing the proof of Theorem 3.1(1). The proof of Theorem 3.1(2) appears in Section 7. Corollary 3.2 easily follows from Theorem 3.1.

4. Domains in the open 2-simplex

In this section we focus our attention on one 2-simplex Δ of the original complex S. We will be interested in open planar surfaces of the form $\Delta - (\partial \Delta \cup T)$ where T is a pure subcomplex of $\mathfrak{s}^k \Delta$. By a pure subcomplex we mean a subcomplex which is the union of simplices of $\mathfrak{s}^k \Delta$ of fixed dimension (either 1 or 2).

Definition 4.1. Given a simplicial complex S and a subcomplex T of S, the open complex defined by S - T is the collection W of all open simplices in S - T. The underlying space of W, i.e. the union of open simplices in W, will be denoted |W|.

Definition 4.2. Let Δ be a fixed 2-simplex.

- (1) We denote by V_k the open complex defined by $\mathfrak{s}^k \Delta \partial(\mathfrak{s}^k \Delta)$, where $\partial(\mathfrak{s}^k \Delta)$ is the subcomplex of $\mathfrak{s}^k \Delta$ which is the topological boundary of Δ . So, $|V_k|$ is the interior of Δ , i.e. an open disc. The open simplices of dimensions 0,1,2 in V_k will be called vertices, edges and triangles, respectively.
- (2) Given a collection $L = \{E_1, \ldots, E_m\}$ of *closed* simplices in $\mathfrak{s}^k \Delta$ of fixed dimension, we denote by U_L the open complex defined by

$$\mathfrak{s}^k \Delta - \left(\partial(\mathfrak{s}^k \Delta) \cup \bigcup_{E_i \in L} E_i \right).$$

An open complex of the form U_L will be called a *domain* in V_k . The fixed dimension of the simplices in L will be called the "type" of U_L . Note that V_k itself is a domain with $L = \emptyset$.

(3) For a domain $U = U_L$ we define $f_i(U)$ to be the number of open simplices of dimension iin U, i = 0, 1, 2, and define $\mu_i(U) = \frac{f_0(U)}{f_i(U)}$, i = 1, 2, whenever $f_i(U) \neq 0$.

Lemma 4.3. Let $U = U_L$ be a domain in V_k , then $\mu_1(U) < \frac{1}{3}$ and $\mu_2(U) < \frac{1}{2}$ whenever they are defined.

Proof. If there are no vertices in U then $\mu_1(U) = \mu_2(U) = 0$ and we are done. Otherwise, let v be a vertex in U. Since U is open, all 6 edges surrounding v must be in U. On the other hand, given an edge e in U, at most 2 vertices in U lie at its endpoints. This gives $\mu_1(U) \leq \frac{1}{3}$, and to show strict inequality we must show that if $\mu_1(U)$ is defined, i.e. if there are edges in U, then at least one of them does not have both endpoints in U. Indeed, starting with an edge e in U, follow a path of edges in V_k to ∂V_k . Since U is open, this path must include an edge with the desired property.

Similarly, if v is a vertex in U then since U is open, all 6 triangles surrounding v are in U. On the other hand at most 3 vertices in U lie on the boundary of a given triangle in U, and triangles adjacent to the boundary of U have strictly less than 3 such vertices. This gives $\mu_2(U) < \frac{1}{2}$. \Box Regarding V_k itself we note the following.

Proposition 4.4. $\lim_{k\to\infty} \mu_1(V_k) = \frac{1}{3}$ and $\lim_{k\to\infty} \mu_2(V_k) = \frac{1}{2}$.

Proof. We claim that for $k \ge 1$, $f_0(V_k) = 3 \cdot 4^{k-1} - 3 \cdot 2^{k-1} + 1$, $f_1(V_k) = 9 \cdot 4^{k-1} - 3 \cdot 2^{k-1}$ and $f_2(V_k) = 6 \cdot 4^{k-1}$, from which the statement follows. This is verified by induction noting that $f_0(V_1) = 1$, $f_1(V_1) = 6$, $f_2(V_1) = 6$, and for $k \ge 1$, $f_0(V_{k+1}) = f_0(V_k) + f_1(V_k)$, $f_1(V_{k+1}) = 2f_1(V_k) + 3f_2(V_k)$ and $f_2(V_{k+1}) = 4f_2(V_k)$.

An open hexagon in V_k is a domain in V_k which is convex and has six sides, which are necessarily parallel to the six sides of V_k . Two examples appear in Figure 4. We denote the boundary of an open hexagon U by ∂U , though strictly speaking it is the boundary of the corresponding closed hexagon.

Lemma 4.5. For U an open hexagon in V_k , let $f_i = f_i(U)$ and b = the number of edges in ∂U , then $f_0 = \frac{f_2 - b + 2}{2}$ and $f_1 = \frac{3f_2 - b}{2}$.

Proof. Every triangle in U is surrounded by three edges. In this way each edge in U is counted twice, and each edge in ∂U is counted once, so $3f_2 = 2f_1 + b$. Also, since the boundary of the closed hexagon contributes 0 to its Euler characteristic, we have $f_0 - f_1 + f_2 = 1$. Our claim follows by combining these two equalities.

For a domain U in V_k , a "shifted image" of U in V_k is a domain in V_k obtained from U by a (perhaps trivial) translation. A "side" of an open hexagon U in V_k is one of the six sides of ∂U .

Lemma 4.6. Let U be an open hexagon in V_k with $U \neq V_k$. Then there is a shifted image of U in V_k whose longest side is not included in ∂V_k .

Proof. If I is one of the six sides of U, then we define its length $\ell(I)$ to be the number of edges in I. Let I be the longest side of U, let $t = \ell(I)$, and let s denote the side length of V_k . If $t \ge s + 1$ we are done since I cannot be included in ∂V_k . If $t \le s - 1$ then U is included in a regular hexagon of side s - 1 and so the claim is again clear. So assume t = s. It cannot be that all sides of U are of length s since then $U = V_k$. So the sides of U may be consecutively labeled I_1, \ldots, I_6 so that $\ell(I_1) = s$, $\ell(I_2) \le s - 1$, and $\ell(I_3) \le s$. If I_1 is not included in ∂V_k we are done, so assume I_1 is a side of V_k . Now let U' be the hexagon with corresponding sides J_1, \ldots, J_6 with $J_1 = I_1$ so $\ell(J_1) = s$, and having $\ell(J_2) = s - 1$ and $\ell(J_3) = \ell(J_4) = s$, as seen in Figure 2. It is easy to



FIGURE 2. The hexagon U' with $\ell(J_2) = s - 1$ and $\ell(J_1) = \ell(J_3) = \ell(J_4) = s$.

see that necessarily $U \subseteq U'$ and so U may be shifted by one to the right, pulling I_1 away from ∂V_k .

We conclude this section with the following trivial lemma that we will use repeatedly throughout this work. It will be used e.g. when say we have disjoint A, B with $\mu_i(A) > c$ and $\mu_i(B) > c$ to deduce $\mu_i(A \cup B) > c$. Another instance where it will be used is when say $f_0 = x$, $f_i = y$, and we are interested to know whether μ_i increases if we replace x, y by $x \pm a, y \pm b$.

Lemma 4.7.

(1) Let x, a ≥ 0, y, b > 0. Then x+a/y+b = tx/y + (1 - t)a/b with 0 < t < 1. That is, either x/y = a/b = x+a/y+b, or x+a/y+b is strictly between x/y and a/b. In particular if a/b > x/y then x+a/y+b > x/y.
(2) Let x ≥ a ≥ 0, y > b > 0. Then x/y = tx-a/y-b + (1 - t)a/b with 0 < t < 1. That is, either x/y = a/b = x-a/y-b, or x/y is strictly between x-a/y-b and a/b. In particular if a/b < x/y then x-a/y-b > x/y.

Proof. For (1) take $t = \frac{y}{y+b}$. For (2) take $t = \frac{y-b}{y}$.

5. Counting triangles

In this section we prove Proposition 3.3(2), where the main technical effort is the following.

Proposition 5.1. For sufficiently large k, every nonempty type-2 domain $U = U_L$ in V_k with $U \neq V_k$ has $\mu_2(U) < \mu_2(V_k)$.



FIGURE 3. Cyclically ordered 6-tuples of o's and x's for type-2 domain.

Proof. Throughout this proof, a type-2 domain will simply be called a domain. We establish our claim by showing that for every domain $U \neq V_k$ there is another domain U' with $\mu_2(U') > \mu_2(U)$. If $\mu_2(U) = 0$ then simply take $U' = V_k$, so we assume $\mu_2(U) > 0$, i.e. $f_0(U) > 0$, which implies $f_2(U) \ge 6$.

To each vertex v of V_k we assign a cyclically ordered 6-tuple of symbols x and o describing the six triangles surrounding v, where an x denotes a 2-simplex in L and an o denotes a 2-simplex which is not in L, and so its interior (and perhaps more of it) is included in U. To each vertex of ∂V_k we also attach such a 6-tuple. It will include x's and o's defined as before, and then completed with a sequence of x's to reach a total of six symbols. One may think of these additional "virtual" x's as corresponding to triangles outside V_k in a tiling of a neighborhood of V_k in the plane. See Figure 3, where the triangles in U are depicted in gray color.

If for some vertex v of V_k the corresponding 6-tuple includes one or two x's then we may remove the corresponding 2-simplices from L. This adds one or two triangles to U, and at least one vertex, namely v itself. So if a is the number of additional vertices and b is the number of additional triangles then $\frac{a}{b} \geq \frac{1}{2}$. From Lemmas 4.3 and 4.7(1) it then follows that μ_2 strictly increases. So we may assume that every 6-tuple includes either 0 or at least 3 x's. This is also true for the vertices of ∂V_k since each such vertex has 3 or 4 virtual x's, as seen in Figure 3.

If for some vertex v of V_k or ∂V_k the corresponding 6-tuple includes a fragment of the form xoxthen we may add the middle 2-simplex to L, changing the given fragment to xxx. This removes one triangle from U, and leaves the number of vertices unchanged, and so it strictly increases μ_2 .

We thus need to establish our claim when all 6-tuples have 0 or at least 3 x's, and a fragment of the form xox does not appear. It follows that all 6-tuples that include both x's and o's are either ooxxxxx or oooxxx. That is, all boundary angles of the connected components of U are either 120°



FIGURE 4. Adding a row of triangles along the longest side of an open hexagon.

or 180° . It follows that each connected component of U is an open hexagon. Indeed, note that a connected component of U cannot have more than one boundary component, since then it would have an interior boundary component, along which there would be angles greater than 180° . By Lemma 4.7(1) it is enough to prove our claim for each such hexagon separately, and so we assume U is a single hexagon.

Let *I* be the longest side of *U*, having length *t*. By Lemma 4.6 we may shift *U* in V_k so that *I* is not included in the boundary of V_k . We may now look at a larger hexagon *U'* in V_k obtained from *U* by adding a full row of triangles along *I* as in Figure 4. This process adds t - 1 vertices and 2t - 1 triangles, as seen in Figure 4 where t = 4. By Lemma 4.5 we have $\mu_2(U) = \frac{f_2 - b + 2}{2f_2}$ where *b* is the number of edges in ∂U . We would like to use Lemma 4.7(1) to show that $\mu_2(U') > \mu_2(U)$, so we would like to have $\frac{t-1}{2t-1} > \frac{f_2 - b + 2}{2f_2}$, i.e. $f_2 < (b-2)(2t-1)$.

We show that this indeed holds if $b \ge 12$. Since I is the longest side, we have $t \ge \frac{b}{6}$ and so it is enough to show that $f_2 < (b-2)(\frac{b}{3}-1) = \frac{1}{3}(b^2-5b+6)$. If we think of our triangles as equilateral triangles in the plane with sides of length 1, then the area A of U is $\frac{\sqrt{3}}{4}f_2$, the length of ∂U is b, and our required inequality may be rephrased as $A < \frac{1}{4\sqrt{3}}(b^2-5b+6)$. By the isoperimetric inequality in the plane we know $A \le \frac{b^2}{4\pi}$ and so it is enough to have $\frac{b^2}{4\pi} < \frac{1}{4\sqrt{3}}(b^2-5b)$, i.e. $\frac{\sqrt{3}}{\pi} < 1-\frac{5}{b}$, which holds for $b \ge 12$.

Finally, there are only finitely many possible hexagons with $b \leq 11$, and so by Lemmas 4.3 and 4.4, for sufficiently large k we have $\mu_2(V_k) > \mu_2(U)$ for every hexagon U with $b \leq 11$, and so for such k our statement is established.

We may now proceed to prove Proposition 3.3(2), namely, that given $\epsilon > 0$, then for sufficiently large k, every subcomplex T of $\mathfrak{s}^k S$ has $\mu_2(T) > \frac{1}{2} - \epsilon$. (Recall that T is a subcomplex in the usual sense, i.e. closed in S.) We first note that it is enough to prove the claim for pure 2-dimensional subcomplexes since given any subcomplex T we may replace it by the subcomplex which is the union of its (closed) 2-simplices. This does not change $f_2(T)$ and may only reduce $f_0(T)$ and so $\mu_2(T)$ may only be reduced. Now, given $\epsilon > 0$, by Proposition 4.4 we have $\mu_2(V_k) > \frac{1}{2} - \epsilon$ for sufficiently large k, and we show that every pure 2-dimensional subcomplex T of $\mathfrak{s}^k S$ satisfies $\mu_2(T) > \frac{1}{2} - \epsilon$. For a fixed 2-simplex Δ of S we look at the corresponding V_k . If $T \cap V_k \neq \emptyset$ we claim that $\mu_2(T \cap V_k) > \frac{1}{2} - \epsilon$ (where $\mu_2(T \cap V_k)$ is defined, as usual, by counting open simplices). Indeed this holds if $T \cap V_k = V_k$. Otherwise look at $U = V_k - (T \cap V_k)$, which is a nonempty type-2 domain in V_k . By Proposition 5.1 we have $\mu_2(U) < \mu_2(V_k)$, and so by Lemma 4.7(2) we have $\mu_2(T \cap V_k) = \mu_2(V_k - U) > \mu_2(V_k) > \frac{1}{2} - \epsilon$. If we look at the union of $T \cap V_k$ over all 2-simplices of S then by Lemma 4.7(1) we still have $\mu_2 > \frac{1}{2} - \epsilon$. All that is missing from the full subcomplex T is $T \cap \mathfrak{s}^k S^1$, where S^1 is the 1-skeleton of S. But $T \cap \mathfrak{s}^k S^1$ may only contribute additional vertices, and no additional 2-simplices, so μ_2 may only increase, and so we finally get $\mu_2(T) > \frac{1}{2} - \epsilon$, completing the proof of Proposition 3.3(2).

Discussion. At this point the reader may have noticed that all our computations up to and including the proof of Proposition 5.1 dealt with domains. Only in the previous paragraph have we finally returned to subcomplexes T of S, and domains made their appearance as the *complement* of $T \cap V_k$ in V_k . The reason for concentrating all our attention on the complement of $T \cap V_k$ in V_k , rather than on T itself, is the following.

Our aim was to show that $\mu_2(T)$ is sufficiently large. We have done this by separately considering the intersection of T with each 2-simplex of S. If we had considered $T \cap \Delta$ for the *closed* simplex Δ , then the vertices in $T \cap \mathfrak{s}^k S^1$ would have been multiply counted since each 1-simplex of S may be included in many 2-simplices of S. For this reason we were forced to consider $T \cap int\Delta = T \cap V_k$. This is a peculiar geometric object with only part of its boundary present. We were not able however to ignore this partial boundary since without it we would be left with a domain in V_k , for which we have seen above that in fact μ_2 tends to be *small*. On the other hand, the fact that the *complement* of $T \cap V_k$ in V_k is a domain, and thus has small μ_2 , was to our advantage since we could use it, via Lemma 4.7(2), to deduce that $T \cap V_k$ itself has large μ_2 .

The same approach will be used in the next section, for proving Proposition 3.3(1).

6. Counting edges

In this section we prove Proposition 3.3(1). Again, the main technical effort is the following.



FIGURE 5. Cyclically ordered 6-tuples of o's and x's for type-1 domain.

Proposition 6.1. For sufficiently large k, every type-1 domain $U = U_L$ in V_k with $U \neq V_k$ has $\mu_1(U) < \mu_1(V_k)$, if $\mu_1(U)$ is defined.

Proof. Throughout this proof, a type-1 domain will simply be called a domain. Again, we establish our claim by showing that for every domain $U \neq V_k$ there is another domain U' with $\mu_1(U') > \mu_1(U)$. If $\mu_1(U) = 0$ then simply take $U' = V_k$, so we assume $\mu_1(U) > 0$, i.e. $f_0(U) > 0$, which implies $f_1(U) \ge 6$. Note that since the simplices in L are of dimension 1, all triangles of V_k are in U.

To each vertex v of V_k we assign a cyclically ordered 6-tuple of symbols x and o, this time describing the six *edges* surrounding v, where an o denotes an edge of V_k that is in U, and an xdenotes an edge of V_k which is not in U. For each vertex of ∂V_k we also define such a 6-tuple, by adding, as before, a sequence of "virtual" x's, to reach a total of six symbols. See Figure 5, where the edges of U are depicted in white color (i.e. they are missing). Since all triangles of V_k are in U, and they are also white, the regions appearing in Figure 5 are in fact the connected components of U. Note that a 6-tuple corresponding to a vertex of ∂V_k includes at most 2 o's, and so at least 4 x's.

If for some edge e of U both endpoints of e are not in U (including the case of an endpoint in ∂V_k), then we can add e to L thus decreasing the number of edges in U and leaving the number of vertices in U unchanged, and so strictly increasing μ_1 . So we may assume that for every edge e of V_k , if both endpoints of e are not in U then e is not in U. If follows that for each triangle of V_k , if two of its sides are not in U (including the case of a side contained in ∂V_k), then also its third side is not in U. This implies that all boundary angles of a connected component of U which is not a single triangle, are at least 120°.

If for some vertex v of V_k the corresponding 6-tuple includes 1, 2, or 3 x's then we may remove the corresponding 1-simplices from L. This adds 1, 2, or 3 edges to U, and at least one vertex, namely v itself. So if a is the number of additional vertices and b is the number of additional edges then $\frac{a}{b} \geq \frac{1}{3}$. From Lemmas 4.3 and 4.7(1) it then follows that μ_1 strictly increases. We may thus assume that for each vertex of V_k the corresponding 6-tuple includes either 0 or at least 4 x's. This is also true for the vertices of ∂V_k , since as we have noted above, each such vertex has at least 4 x's. It follows that all boundary angles of the connected components of U are at most 180° .

We thus need to establish our claim when all boundary angles of the connected components of U which are not single triangles, are either 120° or 180°. It follows, as before, that each connected component of U is either a single triangle or a hexagon. The single triangles have no vertices and no edges and so they do not contribute to $\mu_1(U)$. Thus by Lemma 4.7(1) it is enough to prove our claim for each of the hexagons.

Look at one such hexagon, and denote it again by U. We construct a larger hexagon U' in V_k , as in the proof of Proposition 5.1, by shifting U in V_k if needed, using Lemma 4.6, and then adding a full row of triangles along the longest edge of U, whose length we denote t. This process adds t - 1 vertices and 3t - 2 edges, as seen in Figure 4 where t = 4. By Lemma 4.5 we have $\mu_1 = \frac{f_2 - b + 2}{3f_2 - b}$. To use Lemma 4.7(1) for showing that $\mu_1(U') > \mu_1(U)$ we need $\frac{t-1}{3t-2} > \frac{f_2 - b + 2}{3f_2 - b}$, i.e. $f_2 < (b-3)(2t-1) + 1$. The argument proceeds from here as in the proof of Proposition 5.1, this time leading to $\frac{\sqrt{3}}{\pi} < 1 - \frac{6}{b}$, which holds for $b \ge 14$.

We proceed to prove Proposition 3.3(1), namely, that given $\epsilon > 0$, then for sufficiently large k, every subcomplex T of $\mathfrak{s}^k S$ has $\mu_1(T) > \frac{1}{3} - \epsilon$. Since μ_1 does not involve 2-simplices, we may replace any T with its 1-skeleton, and so it is enough to prove our claim for 1-dimensional subcomplexes, i.e. graphs. Furthermore, it is enough to prove the claim for pure 1-dimensional subcomplexes, i.e. graphs with no isolated vertices, since given any graph T we may remove its isolated vertices, which may only reduce $\mu_1(T)$. Now, given $\epsilon > 0$, by Proposition 4.4 we have $\mu_1(V_k) > \frac{1}{3} - \epsilon$ for sufficiently large k, and we show that every subgraph T of $\mathfrak{s}^k S$ with no isolated vertices satisfies $\mu_1(T) > \frac{1}{3} - \epsilon$. For fixed 2-simplex Δ of S we look at the corresponding V_k . If $T \cap V_k \neq \emptyset$ we claim that $\mu_1(T \cap V_k) > \frac{1}{3} - \epsilon$. Indeed this holds if $T \cap V_k$ includes all edges of V_k ,

since then $\mu_1(T \cap V_k) = \mu_1(V_k)$. Otherwise look at $U = V_k - (T \cap V_k)$, which is a type-1 domain in V_k that includes edges and so $\mu_1(U)$ is defined. By Proposition 6.1 we have $\mu_1(U) < \mu_1(V_k)$, and so by Lemma 4.7(2) we have $\mu_1(T \cap V_k) = \mu_1(V_k - U) > \mu_1(V_k) > \frac{1}{3} - \epsilon$. If we look at the union of $T \cap V_k$ over all 2-simplices of S then by Lemma 4.7(1) we still have $\mu_1 > \frac{1}{3} - \epsilon$.

This completes the computation for the intersection of T with the interiors of the 2-simplices of S, but we are still missing $T \cap \mathfrak{s}^k S^1$ in case it is nonempty, (recall S^1 is the 1-skeleton of S). We claim that for $k \geq 1$, every subgraph of $\mathfrak{s}^k S^1$ has $\mu_1 > \frac{1}{2} > \frac{1}{3} - \epsilon$, which would complete our proof using Lemma 4.7(1). Denoting the graph $\mathfrak{s}^{k-1}S^1$ by P, we have $\mathfrak{s}^k S^1 = \mathfrak{s}^1 P$. Note that for every edge e of $\mathfrak{s}^1 P$, one endpoint of e is an original vertex of P, and the other endpoint is a vertex subdividing an edge of P, and so its degree in $\mathfrak{s}^1 P$ is 2. Now let $G \subseteq \mathfrak{s}^1 P$ be a subgraph, and we may assume as before that G has no isolated vertices. Let v_1, \ldots, v_m be those vertices of G which are vertices of P, and let $d_i > 0$ be the degree of v_i in G. Let H_i be v_i together with its d_i adjacent edges in G, and "half" of the vertex on the other side of each such edge. Then the H_i are disjoint, and H_i includes $1 + \frac{d_i}{2}$ vertices and d_i edges, and so $\mu_1(H_i) = \frac{1+\frac{d_i}{d_i}}{d_i} > \frac{1}{2}$. By Lemma 4.7(1) the combined contribution of all H_i produces $\mu_1 > \frac{1}{2}$. All that is now missing is perhaps some number of half vertices, which may only increase μ_1 , so we finally get $\mu_1(G) > \frac{1}{2}$, completing the proof of Proposition 3.3(1).

Now that both parts of Proposition 3.3 are proved, we have completed the proof of Theorem 3.1(1).

7. TOPOLOGICAL NON-EMBEDDABILITY

In this concluding section we prove Theorem 3.1(2) stating that for a probability triple \mathfrak{p} satisfying $\lim_{n\to\infty} np_0p_1^3p_2^2 = 0$, the torus is a.a.s. not topologically embeddable into $Y \in (\Omega_n, \mathbb{P}_{\mathfrak{p}})$. We will need the following lemma.

Lemma 7.1. Let F be a closed surface and Y a 2-dimensional simplicial complex. If $h : F \to Y$ is a topological embedding then h(F) is a subcomplex of Y. It follows that there is a triangulation S of F for which the embedding h is simplicial.

Proof. We first show that if V is the interior of a 2-simplex in Y, then either $h(F) \supseteq V$ or $h(F) \cap V = \emptyset$. This we do by showing that $h(F) \cap V$ is closed and open in V. Since h(F) is compact, it is closed in Y, and so $h(F) \cap V$ is closed in V. On the other hand, since Y is a

2-dimensional complex, V is open in Y and so $h^{-1}(V)$ is open in F. By invariance of domain (Corollary 19.9 in [B]), $h|_{h^{-1}(V)} : h^{-1}(V) \to V$ is an open mapping, and so its image, $h(F) \cap V$, is open in V.

Now let G be the union of those 2-simplices in Y whose interiors are contained in h(F). Then G is a subcomplex of Y and we show h(F) = G. Since h(F) is closed in Y and is disjoint from the interiors of all other 2-simplices of Y, we have $G \subseteq h(F) \subseteq G \cup Y^1$, where Y^1 is the 1-skeleton of Y. Let $U = F - h^{-1}(G)$ then U is open in F and $h(U) \subseteq Y^1$. Since Y^1 may be embedded into some surface as a subset with empty interior, by invariance of domain again we must have $U = \emptyset$ and so h(F) = G.

By Lemma 7.1, the torus is topologically embeddable into a given 2-dimensional simplicial complex Y iff there exists a triangulation S of the torus that is simplicially embeddable into Y. If S is a triangulation of the torus with f_i simplices of dimension i, then there are $n(n-1)\cdots(n-f_0+1) \leq n^{f_0}$ injections of its vertex set into [n]. Given such an injection, the probability that it determines a simplicial embedding of S into $Y \in (\Omega_n, \mathbb{P}_p)$ is precisely $p_0^{f_0} p_1^{f_1} p_2^{f_2}$. It follows that the probability that S is simplicially embeddable into Y is at most $n^{f_0} p_0^{f_0} p_1^{f_1} p_2^{f_2}$. Every triangulation of the torus satisfies $f_0 - f_1 + f_2 = 0$ and $3f_2 = 2f_1$, from which it follows that $f_1 = 3f_0$ and $f_2 = 2f_0$. Furthermore, it follows from [G] that there is a constant c such that the number of nonequivalent triangulations of the torus with f_0 vertices which is simplicially embeddable into Y is at most $(cnp_0p_1^3p_2^2)^{f_0} = u_n^{f_0}$ where $u_n = cnp_0p_1^3p_2^2 \longrightarrow 0$ by our assumption. Finally, summing over all possible f_0 , the probability that there is some triangulation of the torus which is simplicially embeddable into $Y \in (\Omega_n, \mathbb{P}_p)$ is bounded by

$$\sum_{f_0=1}^{\infty} u_n^{f_0} = \frac{u_n}{1-u_n} \xrightarrow[n \to \infty]{} 0.$$

References

- [B] G. E. Bredon: *Topology and Geometry*. Springer (1993).
- [CCFK] D. Cohen, A. Costa, M. Farber, T. Kappeler: Topology of Random 2-Complexes. Discrete & Computational Geometry (2012) 47:117–149
- [CF1] A. Costa, M. Farber: Large random simplicial complexes, I. Journal of Topology and Analysis Vol. 8, No. 3 (2016) 399–429.

- [CF2] A. Costa, M. Farber: Large random simplicial complexes, II; the fundamental group. Journal of Topology and Analysis Vol. 9, No. 3 (2017) 441–483.
- [CF3] A. Costa, M. Farber: Large random simplicial complexes, III the critical dimension. Journal of Knot Theory and Its Ramifications Vol. 26, No. 2 (2017).
- [FMN] M. Farber, L. Mead, T. Nowik: Random Simplicial Complexes, Duality and The Critical Dimension. Journal of Topology and Analysis to appear. arXiv:1901.09578
- [G] Z. Gao: The number of rooted triangular maps on a surface. Journal of Combinatorial Theory, Series B
 Vol. 52, No. 2 (1991) 236–249.
- [GW] A. Gundert, U. Wagner: On topological minors in random simplicial complexes. Proceedings of the American Mathematical Society Vol. 144, No. 4 (2016) 1815–1828.
- [LM] N. Linial, R. Meshulam: Homological connectivity of random 2-complexes. Combinatorica Vol. 26, No. 4 (2006) 475–487.
- [MW] R. Meshulam, N. Wallach: Homological connectivity of random k-dimensional complexes. Random Structures & Algorithms Vol. 34, No. 3 (2009) 408–417.

SCHOOL OF MATHEMATICAL SCIENCES, QUEEN MARY UNIVERSITY OF LONDON, LONDON E1 4NS, UK *E-mail address*: M.Farber@qmul.ac.uk *URL*: www.qmul.ac.uk/maths/profiles/farberm.html

DEPARTMENT OF MATHEMATICS, BAR-ILAN UNIVERSITY, RAMAT-GAN 5290002, ISRAEL E-mail address: tahl@math.biu.ac.il URL: www.math.biu.ac.il/~tahl