On some geometric aspects of the class of hv-convex swtching components

Paolo Dulio * Andrea Frosini †

December 16, 2020

Abstract

In the usual aim of discrete tomography, the reconstruction of an unknown discrete set is considered, by means of projection data collected along a set U of discrete directions. Possible ambiguous reconstructions can arise if and only if switching components occur, namely, if and only if non-empty images exist having null projections along all the directions in U. In order to lower the number of allowed reconstructions, one tries to incorporate possible extra geometric constraints in the tomographic problem, such as the request for connectedness, or some reconstruction satisfying special convexity constraints. In particular, the class $\mathbb P$ of horizontally and vertically convex connected sets (briefly, hv-convex polyominoes) has been largely considered.

In this paper we introduce the class of hv-convex switching components, and prove some preliminary results on their geometric structure. The class includes all switching components arising when the tomographic problem is considered in \mathbb{P} , which highly motivates the investigation of such configurations, also in view of possible uniqueness results for hv-convex polyominoes.

It turns out that the considered class can be partitioned in two disjointed subclasses of closed patterns, called windows and curls, respectively, according as the pattern can be travelled by turning always clockwise (or always counterclockwise), or points with different turning directions exist. It follows that all windows have a unique representation, while curls consist of interlaced sequences of sub-patterns, called Z-paths, which leads to the problem of understanding the combinatorial structure of such sequences

We provide explicit constructions of families of curls associated to some special sequences, and also give additional details on further allowed or forbidden configurations by means of a number of illustrative examples.

Keywords: curl, Discrete Tomography, *hv*-convex set, polyomino, projection, switching-component, window, X-ray.

Math subject classification: 52A30, 68R01, 52C30, 52C45

^{*}Dipartimento di Matematica "F. Brioschi", Politecnico di Milano, Piazza Leonardo da Vinci 32, I-20133 Milano, Italy

 $^{^\}dagger \text{Università}$ di Firenze, Dipartimento di Sistemi e Informatica, Viale Morgagni 65, 50134 Firenze, Italy

1 Introduction

Discrete Tomography is a part of the wider area of Computerized Tomography, which relates to a huge number of applications where image reconstruction from X-ray collected data is required. While Computerized Tomography involves analytical techniques and continuous mathematics (see, for instance [2, 1]), Discrete Tomography is mainly concerned with discrete and combinatorial structures, it works with a small number of density values, in particular with homogeneous objects, and usually allows very few X-rays directions to be considered (see [19, 20] for a general introduction to the main problems of Discrete Tomography).

The reconstruction problem is usually *ill-posed*, meaning that *ambiguous* reconstructions are expected. To limit the number of allowed configurations, further information is usually incorporated in the tomographic problem, which sometimes leads to a unique solution (see for instance [15]) in the case of convex reconstructions), or to the enumeration of the allowed solutions (an example with two projections is [3]).

In case different discrete sets Y_1 and Y_2 are tomographically equivalent with respect to a set U of directions, namely Y_1, Y_2 can be reconstructed by means of the same X-rays with respect to U, then there exist specific patterns, called switching components which turn Y_1 into Y_2 . Understanding the combinatorial and the geometric structure of the switching components is a main issue in discrete tomography (see, for instance [5, 6, 7, 9, 12, 13, 14, 15, 17, 16, 18]).

A largely investigated case concerns the class \mathbb{P} of hv-convex polyominoes, i.e., finite connected subsets of \mathbb{Z}^2 that are horizontally and vertically convex. Early results for two projections can be found in [6], where a uniqueness conjecture has been also stated, later disproved in [11]. On this regard, a main role is played by switching components with respect to the horizontal and to the vertical directions, respectively denoted by \overrightarrow{h} and \overrightarrow{v} . In [17], such switching components have been studied from an enumerative and an algorithmic point of view, which provided a very interesting and illustrative presentation of their connection with the complexity of the reconstruction problem. In this paper we also focus on such switching components, but we follow a different approach, based on a special geometrical condition (see Definition 2), which defines a class of patterns called hv-convex switching components. It includes the classes of regular and of irregular switching components considered in [17], that we redefine in terms of hv-convex windows and hv-convex curls, respectively.

The geometric condition in Definition 2 is always satisfied when the switching component is determined by a pair of sets Y_1, Y_2 both internal to the class \mathbb{P} . This motivates a deep investigation of the structure of hv-convex switching components, in view of possible uniqueness results for hv-convex polyominoes.

We give a geometric characterization of hv-convex windows (Theorem1), and a necessary condition for a curl to be a hv-convex switching component (Theorem 2). In general, the condition is not sufficient, but it provides a basic information concerning the geometric structure of hv-convex curls, which leads to the problem of understanding their geometric and combinatorial structure.

2 Notations and preliminaries

We first introduce some notations and basic definitions. As usual, \mathbb{R}^2 denotes the Euclidean two-dimensional space, and $\mathbb{Z}^2 \subset \mathbb{R}^2$ is the lattice of points having integer coordinates. If A is a subset of \mathbb{R}^2 , we denote by int(A) and by conv(A) the *interior* and the $convex\ hull$ of A, respectively. If A consists of two distinct points v and w, then conv(A) is a segment, denoted by s(v, w). If A is a finite set of \mathbb{Z}^2 , then A is said to be a $lattice\ set$, and |A| denotes the number of elements of A. A $convex\ lattice\ set$ is a lattice set $A \subset \mathbb{Z}^2$ such that $A = (conv(A)) \cap \mathbb{Z}^2$.

By \overrightarrow{h} , \overrightarrow{v} we mean the horizontal and the vertical directions, respectively. For any point $v \in \mathbb{R}^2$, we indicate by $L_h(v)$ and $L_v(v)$ the horizontal and the vertical line passing through v, respectively.

Finally, we define horizontal (resp. vertical) projection of a finite set $A \subset \mathbb{Z}^2$ to be the integer vector H(A) (resp. V(A)) counting the number of points of A that lie on each horizontal (resp. vertical) line passing through it. We underline that such a notion of projection can be defined for a generic set of discrete lines parallel to a given (discrete) direction.

In literature, the word *polyomino* indicates a connected finite discrete set of points. In particular, a polyomino is hv-convex if each one of its rows and columns is connected. As it is commonly assumed, a polyomino is composed by rows and columns due to the habit of representing it by a binary matrix whose dimensions are those of its minimal bounding rectangle. The class of all hv-convex polyominoes is denoted by \mathbb{P} .

Given a point $v = (i, j) \in \mathbb{Z}^2$, the four following closed regions are defined (with the same notations as in [8, 10]):

$$Z_0(v) = \{(i', j') \in \mathbb{R}^2 : i' \le i, \ j' \le j\}, \quad Z_1(v) = \{(i', j') \in \mathbb{R}^2 : i' \ge i, \ j' \le j\},$$

$$Z_2(v) = \{(i', j') \in \mathbb{R}^2 : i' \ge i, \ j' \ge j\}, \quad Z_3(v) = \{(i', j') \in \mathbb{R}^2 : i' \le i, \ j' \ge j\}.$$

A set of points A is said to be Q-convex (quadrant convex) along the horizontal and vertical directions if $Z_l(v) \cap A \neq \emptyset$ for all l = 0, 1, 2, 3 implies $v \in A$.

Lemma 1. Let P be a hv-convex polyomino, and consider a point $v \in \mathbb{Z}^2$. If $w_1, w_2, w_3 \in P$ exist such that $Z_i(v) \cap \{w_1, w_2, w_3\} \neq \emptyset$ for all i = 0, 1, 2, 3, then $v \in P$.

Proof. By [8, Proposition 2.3], a hv-convex set is also Q-convex with respect to the horizontal and to the vertical directions. The statement follows immediately by the hv-convex property of P.

2.1 Switching components and the uniqueness problem

Definition 1. A pair $S = (S^0, S^1)$ of sets of points is a hv-switching if:

- $S^0 \cap S^1 = \emptyset$ and $|S^0| = |S^1|$;
- $H(S^0)=H(S^1)$ and $V(S^0)=V(S^1)$, i.e., S^0 and S^1 have the same horizontal and vertical projections.

Each set S^0 and S^1 is indicated as hv-switching component. We underline that also the notion of switching can be extended to the projections along a

generic set of discrete directions (again refer to [19, 20] for these definitions and the related main results).

A discrete set A contains a hv-switching component if $S^0 \subseteq A$ and $S^1 \cap A = \emptyset$. In this case, we consider $A = Y \cup S^0$, with Y being a (possibly void) discrete set; we define the set $A' = Y \cup S^1$ as the dual of A, and we say that the switching S is associated to A and A'.

2.2 hv-convex switching

A classical result in [18] states that if A_1 and A_2 are two discrete sets sharing the same horizontal and vertical projections, then A_2 is the dual of A_1 with respect to a hv-switching. So, for any point $v \in S^0$ (resp. $v \in S^1$), there exist points $w_1, w_2 \in S^1$ (resp. $w_1, w_2 \in S^0$) such that $w_1 \in L_h(v)$ and $w_2 \in L_v(v)$.

If the sets A_1 and A_2 are hv-convex polyominoes, then, due to Lemma 1, for any $x \in S$ there exists one and only one $i \in \{0, 1, 2, 3\}$ such that $Z_i(x) \cap S$ consists of points all belonging to the same component of S as x. The quadrant $Z_i(x)$ is said to be the *free region* of x, or the S-free region of x in case we wish to emphasize that the free region relates to the switching S. We denote by F(x) (or by $F_S(x)$) the free region of $x \in S$. Also, $F_i(S)$ denotes the subset of S consisting of all points having free region $Z_i(x)$, namely $F_i(S) = \{x \in S, F_S(x) = Z_i(x)\}$, $i \in \{0, 1, 2, 3\}$.

We have the following

Lemma 2. Let $S = (S^0, S^1)$ be a hv-switching. Then, the following conditions are equivalent

$$\bigcup_{i=0}^{3} Z_i(S) = S. \tag{1}$$

$$v, w \in Z_i(S), i \in \{0, 1, 2, 3\}, v \in S^0, w \in S^1 \Rightarrow v \notin Z_j(w), w \notin Z_j(v), j = i + 2 \pmod{4}.$$

Proof. Let $v, w \in S$ such that $v, w \in Z_i(S)$, with $v \in S^0, w \in S^1$. Suppose that $v \in Z_j(w)$, with $j = i + 2 \pmod{4}$. Then $w \in Z_i(v)$, a contradiction. Analogously, if $w \in Z_j(v)$, with $j = i + 2 \pmod{4}$, then $v \in Z_i(w)$, a contradiction. Therefore, (2) holds. Conversely, assume that (2) holds. Let $v \in S$, and suppose $v \in S^0$. Since S is a hv-switching, then there exist three values of $k \in \{0, 1, 2, 3\}$ such that $Z_k(v) \cap S^1 \neq \emptyset$. Suppose that $w \in S^1$ exists such that $w \in Z_i(v)$ for $i \neq k$. Then $v \in Z_j(w)$, where $j = i + 2 \pmod{4}$, which contradicts (2). Therefore, $v \in S^1$ has a free region, namely $v \in S^1$ has a free region. Therefore, each point of $v \in S^1$ has a nonempty free region, and (1) follows.

Definition 2. Let $S = (S^0, S^1)$ be a hv-switching. Then, S is said to be a hv-convex switching if one of the equivalent conditions of Lemma 2 holds.

Remark 1. By the above discussion, if $S = (S^0, S^1)$ is a hv-switching associated to a pair of hv-convex polyominoes, then (1) holds, so S is a hv-convex switching. However, the converse is not necessarily true, namely it could exist two polyominoes P_1 and P_2 that are one the dual of the other with respect

to S and such that one or both of them are not hv-convex polyominoes. An interesting case is Figure 23 in [17], or Figure 2 below.

2.3 Squared spirals

A closed polygonal curve K in \mathbb{R}^2 is said to be a squared spiral if K consists of segments having, alternatively, horizontal and vertical direction. Their endpoints form the set of vertices of the polygonal, denoted by V(K). Two squared spirals are said to intersect in case some of (possibly all) their segments done. Assume to travel K according to a prescribed orientation. A vertex v of K is said to be a counterclockwise point if, crossing v, implies a counterclockwise change of direction. Differently, v is a clockwise point. Of course, by reversing the travelling orientation, clockwise and counterclockwise vertices mutually exchange. The bounding rectangle of K is the smallest rectangle R_K containing K.

2.4 Windows and curls

We now introduce two classes of special squared spirals that provide a geometric reformulation of the notions of regularity and of irregularity discussed in [17], which, in addition, constitute the main focus of our study. A squared spiral W is said to be a window if it can be traveled by turning always clockwise, or always counterclockwise. Differently, the squared spiral is said to be a curl. Therefore, travelling a curl needs changes of turning direction.

Obviously a rectangle is a particular case of window that coincides with its bounding rectangle.

Remark 2. Each window and each curl form a hv-switching $S = (S^0, S^1)$ by considering the corresponding vertices alternatively belonging to S^0 and S^1 .

2.5 Z-paths

A Z-path is a staircase shaped pattern consisting of a monotone sequence of horizontal and vertical segments, whose vertices alternate between clockwise and counterclockwise points. We say that the Z-path is of type SE-NW, or SW-NE, according as it can be travelled moving from South-East to North-West (or conversely), or from South-West to North-East (or conversely), respectively. A simple, or one-level, Z-path consists of just three segments, horizontal-verticalhorizontal, or vertical-horizontal-vertical, referred to as hvh, or vhv Z-path, respectively. Excluding its endpoints, a simple Z-path exhibits a pair of vertices having a specified orientation, clockwise-counterclockwise, or clockwisecounterclockwise, according to the considered type, and moving from south to north along the pattern. In general, for q > 0, we have a q-level Z-path if, excluding its endpoints, it consists of q+1 vertices having alternating orientations. Therefore, if q is odd, we have q horizontal and q-1 vertical segments, or conversely, and we refer to the corresponding Z-path with the notation $h(vh)_{q-1}$ and $v(hv)_{q-1}$, respectively. If q is even, then the Z-path consists of q horizontal and of q vertical segments, and we adopt the notation $(hv)_q$, or $(vh)_q$, according as the first segment is horizontal or vertical (see Figure 1). Any Z-path is a hvconvex set. In a SE-NW Z-path, any vertex v, different from an endpoint, has free region $Z_0(v)$ or $Z_2(v)$, while, in a SW-NE Z-path, the free region is $Z_1(v)$ or $Z_3(v)$. In any case, the elements of the sets of free regions $\{Z_0(v), Z_2(v)\}$, or $\{Z_1(v), Z_3(v)\}$ alternate along the Z-path. Since the vertices of a Z-path are, alternatively, clockwise and counterclockwise oriented, then no q-level Z-path, with q>0, can be found in a window, while any curl surely includes some Z-paths. Differently, if q=0, we have an L-shaped path, consisting of an horizontal and a vertical segment, with just one intermediate point. We refer to such a path as a degenerate Z-path. Note that a window can be considered as a consecutive sequence of degenerate Z-paths, while, in a curl, different Z-paths (possibly degenerate) can appear. In what follows, we provide a precise characterization of how these paths can be combined together.



Figure 1: Different types of Z-paths. From left to right: A simple hvh SE-NW Z-path, a $v(hv)_2$ SE-NW Z-path, a $(hv)_3$ SW-NE Z-path, and a $v(hv)_2$ SW-NE Z-path.

3 Characterization of hv-convex windows

We give a necessary and sufficient condition for a window to be hv-convex. This leads to a geometric characterization of the hv-convex switchings that have the structure of a hv-convex window.

Theorem 1. Let W be a window of size $n \geq 1$ and $\{w_1, w_2, ..., w_{4n}\}$ be the set of its vertices. Then W is a hv-convex switching if and only if a point $x \in \mathbb{R}^2$ exists such that $w_i \in Z_0(x) \cup Z_2(x)$ for all the odd indices, and $w_i \in Z_1(x) \cup Z_3(x)$ for all the even indices.

Proof. Assume that a point $x \in \mathbb{R}^2$ exists such that $w_i \in Z_0(x) \cup Z_2(x)$ for all the odd indices, and $w_i \in Z_1(x) \cup Z_3(x)$ for all the even indices. Then, by definition of window, W has the same number of vertices in each $Z_i(x)$, i = 0, 1, 2, 3, namely, $w_i \in Z_0(x)$, for $i = 1 \pmod{4}$, $w_i \in Z_1(x)$, for $i = 2 \pmod{4}$, $w_i \in Z_2(x)$, for $i = 3 \pmod{4}$, and $w_i \in Z_3(x)$, for $i = 0 \pmod{4}$. Therefore, if W^0 and W^1 are, respectively, the set of the even and of the odd labeled vertices of W, then each point of W^0 has a horizontal and a vertical corresponding in W^1 and conversely. This implies that the free regions of all points in W^0 are contained in $Z_1(x)$ or in $Z_3(x)$, and the free regions of all points in W^1 are contained in $Z_0(x)$ or in $Z_2(x)$. Therefore, W is hv-convex.

Conversely, suppose that W is a hv-convex switching. Without loss of generality we can assume that W is traveled counterclockwise, starting from w_1 . Also, up to a rotation (which does not change the argument) we can always assume that the free region of w_1 is $Z_0(w_1)$. Then the free region of w_i is $Z_j(w_i)$ where $i-j=1 \pmod 4$. For j=0,1,2,3, let H_j be the set

$$H_j = \bigcup_{i=j+1 \pmod{4}} Z_j(w_i).$$

Due to the hv-convexity of W, the sets H_i are mutually disjointed. Consider the strip bounded by the two horizontal lines supporting $H_0 \cup H_1$ and $H_2 \cup H_3$, and the strip bounded by the two vertical lines supporting $H_0 \cup H_3$ and $H_1 \cup H_2$ (see Figure 2).




Figure 2: The four regions H_i , $i \in \{0, 1, 2, 3\}$ related to a hv-convex window. The rectangle R_x contains all the points having the property stated in Theorem 1.

The intersection of such strips forms a rectangle R, having horizontal and vertical sides, and with no points of W belonging to the internal int(R) of R. Let x be any point such that $x \in int(R)$. Then $H_j \subseteq Z_j(x)$ for all j = 0, 1, 2, 3, and the statement follows.

Remark 3. The property stated in Theorem 1 is not restricted to a single point, but it involves all the points belonging to the interior of the rectangle R.

For any hv-convex window W, and point $x \in \mathbb{R}^2$ as in Theorem 1, all quadrants $Z_i(x)$, $i \in \{0, 1, 2, 3\}$ contain the same number of points of W, which is said the *size* of the window. Note that a window can be a switching component with respect to the horizontal and vertical directions without being hv-convex.

4 Characterization of hv-convex curls

Moving to curls, a deeper analysis is required, as it has been pointed out in [17] in terms of irregular switching components. Here we push the study a step ahead, by investigating the geometric nature and the main features of those curls that form hv-convex switching. As a first result, we prove a necessary condition for a curl to be a hv-convex switching, say hv-convex curl. In general, the given condition is not sufficient, but it spreads light on the geometric structure of the hv-convex curls, and leads to their characterization in terms of Z-paths. As a consequence, the class of hv-convex curls will be partitioned into two subclasses.

Theorem 2. Let C be a curl that forms a hv-switching, and let v and w be two points in V(C) with the same orientation. If precisely 2n > 0 consecutive vertices between v and w exist, and having their opposite orientation, then C is not a hv-convex curl.

Proof. Suppose that C is hv-convex. Without loss of generality, we can assume that travelling C from v to w the vertices v and w are counterclockwise oriented. Up to a rotation we can also assume that $Z_0(v)$ is the free region of v, so that a vertex $v_1 \in Z_2(v) \cap Z_3(v)$ exists, with v, v_1 in different components of C. Let

 $x_1, ..., x_{2n}$ be the 2n > 0 clockwise oriented vertices of C that are crossed when moving from v to w.

The segment $s(v,x_1)$ is horizontal. The same holds for the segment $s(x_{2n},w)$, and also for all segments $s(x_{2k},x_{2k+1})$, for $1 \leq k \leq n-1$. Analogously, all segments $s(x_{2k-1},x_{2k})$, for $1 \leq k \leq n$ are vertical. Then $Z_2(x_1)$ is the free region of x_1 , $Z_1(x_2)$ is the free region of x_2 , $Z_0(x_3)$ is the free region of x_3 , and, in general, the free region of x_i is the quadrant $Z_j(x_i)$ such that $i+j=3 \pmod 4$. Therefore, the free region of x_{2n} is $Z_1(x_{2n})$ if n is odd, and $Z_3(x_{2n})$ if n is even, which implies that the free region of w is, respectively, $Z_3(w)$ and $Z_1(w)$ (see Figure 3, where the case $F(w)=Z_1(w)$ is represented).




Figure 3: Positions of consecutive vertices having a same orientation in a curl.

Now, all the vertices x_h , with h odd, belong to a component different from that v, so they do not lie in $F(v) = Z_0(v)$. Since $x_1 \in Z_1(v) \cap Z_2(v)$, then $x_2 \in int(Z_1(v))$. If n = 1, then $w \in int(Z_1(v))$, and $F(w) = Z_3(w)$, so that $v \in F(w)$, a contradiction, since v and w belong to different components of C. So, the statement follows for n = 1. If n > 1, then $x_3 \in int(Z_1(v))$, so $x_4 \in Z_1(v) \cup Z_2(v)$. However, $x_4 \notin Z_1(v)$, since, otherwise $v_1 \in F(x_4) = Z_3(x_4)$, a contradiction, being v and v_1 in different components of C. By iterating the argument, we get that all the vertices of the form x_{2k} , with $k \leq n$ and k even must belong to $Z_2(v)$. Analogously, all the vertices of the form x_{2k} , with $k \leq n$ and k odd must belong to $Z_1(v)$, since, differently, x_{2k} would belong to $F(x_1) = Z_3(x_1)$, or conversely, $x_1 \in F(x_{2k}) = Z_1(x_{2k})$, a contradiction, since x_1 and x_{2k} belong to different components of C. This implies that the vertices x_{2k-1} with $1 \leq k \leq n$ and k odd belong to $Z_2(v)$. Consequently, also $w \in Z_1(v) \cup Z_2(v)$.

Suppose that $w \in Z_1(v)$. Since the segment $s(x_{2n}, w)$ is horizontal, then x_{2n} also belongs to $Z_1(v)$. As shown above, this implies that n is odd, so $F(w) = Z_3(w)$, and consequently $v \in F(w)$, a contradiction.

Hence $w \in Z_2(v)$, then x_{2n} also belongs to $Z_2(v)$, which implies that n is even, and consequently $F(x_{2n}) = Z_3(x_{2n})$, and $F(w) = Z_1(w)$. Therefore, w must belong to $Z_2(x_2)$, otherwise $w \in Z_3(x_2)$, and consequently $x_2 \in F(w) = Z_1(w)$, a contradiction. From $w \in Z_2(x_2)$, and $w \in Z_2(v)$, it follows that $w \in Z_2(x_1)$. Since C is a switching with respect to the vertical direction, then there exists a vertex $w_1 \in Z_2(w) \cap Z_3(w)$, with w and w_1 in different components, and consequently also x_1 and w_1 belong to different components of C (see Figure 3). Since $w \in Z_2(x_1)$, then also $w_1 \in F(x_1) = Z_2(x_1)$, a contradiction.

Consequently, the assumption that C is hv-convex always leads to a contradiction, and the statement follows. \Box

5 On some sequences associated to hv-convex switchings

Let S be a squared spiral. We associate to S an integer sequence $(k_1, k_2, ..., k_n)$, say hv-sequence, where each k_i represents the i-th maximal sequence of k_i vertices that can be travelled clockwise or counterclockwise, with i = 1, 2, ..., n. The starting vertex is not indicated, so the sequence can be considered up to circular shifts. If the sequence $(k_1, k_2, ..., k_n)$ is periodic, then we adopt the notation $(k_1, ..., k_{n'})_h$, to represent the h time repetition of the sequence $(k_1, ..., k_{n'})$, with $n = n' \cdot h$; if h = 1, we choose to omit it. We are interested in characterizing the hv-sequences that admit a hv-convex switching, say hv-convex sequences. Therefore, we are led to the following general problem.

Problem 1. For which $k_1,...,k_n,h \in \mathbb{N}$ there exists a hv-convex sequence $(k_1,...,k_n)_h$?

Concerning windows, Problem 1 has an easy solution.

Theorem 3. For each n > 0, (4n) is a hv-convex sequence if and only if the associated square spiral is a window.

Proof. By Theorem 2, the hv-sequence associated to a curl is of the form $(k_1, ..., k_{n'})_h$, where $k_1, ..., k_n$ are odd. Therefore, (4n) cannot be the hv-sequence associated to a curl. Let W be a window. By Theorem 1, $x \in \mathbb{R}^2$ exists such that all quadrants $Z_i(x)$, $i \in \{0, 1, 2, 3\}$ contain the same number of points of W. Then, W has 4n vertices, for some n > 0, which implies that the hv-sequence associated to W is (4n).

Differently, Problem 1 seems to require a deeper investigation of the geometrical and combinatorial structure of the set of vertices of a curl. In this view, we give here some preliminary remarks. First of all, note that, having a $(k_1, ..., k_n)_h$ curl, is in general not sufficient to get hv-convexity, since the conditions in Lemma 2 do not automatically hold.

A vertex v of a curl whose turning direction differs from that of the previous encountered point, is said to be a *changing point*. In order to improve our knowledge on the hv-sequences allowed for curls, it is worth focusing on the possible Z-paths that can be included in a hv-convex curl, which reflects in the understanding of how changing points can occur. As already observed, the simplest hv-convex curl is the $(3,3)_1$ curl shown in Figure 4 (a). Its vertices consists of six points x_1, \ldots, x_6 , where $F(x_1) = Z_1(x_1)$, $F(x_2) = Z_0(x_2)$, $F(x_3) = Z_3(x_3)$, $F(x_4) = Z_1(x_4), F(x_5) = Z_2(x_5), F(x_6) = Z_3(x_6).$ The $(3,3)_1$ curl can also be considered as the join of two simple hvh and vhv SW-NE Z-paths (see Section 2.5) having x_2 and x_5 in common. This means that x_1, x_4 are changing points (or x_3 , x_6 , depending on the starting choice for the walking direction). Figure 4 (b) shows a $(3,3)_2$ hv-convex curl, consisting of two different pairs of intersecting simple SW-NE Z-paths (analogous constructions can be performed by using SE-NW Z-paths). Analogously, for any integer number $h \geq 1$, a curl C can be constructed having h pairs of intersecting SW-NE Z-paths. These can be consecutively arranged, or, differently, connected by means of L-shaped paths, as described above. See Figure 4 (c) for an example where h = 4.

Of course, curl containing Z-paths of higher level can be also constructed.




Figure 4: (a) A $(3,3)_1$ hv-convex curl C, corresponding to a SW-NE vertexgluing of two rectangles. (b) A $(3,3)_2$ hv-convex curl with two pairs of intersecting simple Z-paths. (c) Example of curl associated to the hv-sequence $(3,3)_4$.

However, different Z-paths of a same curl are not necessarily consecutive. For instance, Figure 5 shows how to insert degenerate Z-paths (L-shaped paths) between the bottom-left endpoint of a Z-path and the upper-right endpoint of a different Z-path, so transforming a $(3,3)_1$ curl into a $(5,5)_1$ curl.



Figure 5: Including L-shaped paths in a given curl. (a) The starting curl. (b) The split of the two constituent simple Z-paths. (c) The connection of the two simple Z-paths by joining their extremal vertices with two degenerate Z-pats.

Further constructions also exist having associated hv-convex sequence of type $(k_1, k_2)_h$, with $k_1 \neq k_2$. Figure 6 shows a curl associated to the hv-convex sequence $(3, 5)_2$.




Figure 6: The hv-convex curl associated to the hv-convex sequence $(3,5)_2$.

6 Conclusion and remarks

We have introduced the class of hv-convex switching components, which includes all switching components associated to a pair of tomographically equivalent hvconvex polyominoes. We have separated the class in two disjointed subclasses of closed patterns, the windows and the curls, respectively. We have given geometrical results on both subclasses, which leads to the problem of characterizing them in terms of hv-convex sequences. While windows provide a complete and easy solution, deeper investigation is required for curls. We have discussed a few preliminary allowed or forbidden hv-sequences, which provide partial answers to Problem 1 in the case of curls. For a complete solution to Problem 1 it becomes relevant to understand how, in general, different Z-paths can be connected between them in a same curl. In particular, it would be worth exploring possible connections between the allowed levels of the Z-paths in a same curl, and the degree of convexity of L-convex sets [3]. We wish to investigate in these directions in separated further works

References

- [1] F. Natterer, The Mathematics of Computerized Tomography, Teubner, Stuttgart Ed. (1986).
- [2] C. Kak Avinash, M. Slaney, Principles of Computerized Tomographic Imaging (Classics in Applied Mathematics): Aninash C. Kak, Malcolm Society for Industrial and Applied Mathematics, (2001).
- [3] G. Castiglione, A. Frosini, A. Restivo, S. Rinaldi, *Enumeration of L-convex polyominoes by rows and columns*, Theoret. Comput. Sci., Vol. 347, 336-352 (2005).
- [4] R.J. Gardner, P. Gritzmann, P., Discrete tomography: determination of finite sets by X-rays Trans. Amer. Math. Soc. Vol. 349, 2271-2295 (1997).
- [5] A. Alpers, and R. Tijdeman, *The two-dimensional Prouhet-Tarry-Escott problem*, J. Number Theory, Vol. 123, 403-412 (2007).
- [6] E. Barcucci, A. Del Lungo, M. Nivat, R. Pinzani, X-rays characterizing some classes of discrete sets, Linear Algebra Appl., Vol. 339, 3-21 (2001).
- [7] E. Barcucci, P. Dulio, A. Frosini, S. Rinaldi, Ambiguity results in the characterization of hv-convex polyominoes from projections, SERIES = Lecture Notes in Comput. Sci., Vol. 10502, 147-158 (2017).
- [8] S. Brunetti, A. Daurat, A., An algorithm reconstructing convex lattice sets, Theoret. Comput. Sci., Vol. 304, 35-57 (2003).
- [9] S. Brunetti, P. Dulio, C. Peri, Discrete tomography determination of bounded lattice sets from four X-rays, Discrete Appl. Math., Vol. 161, 2281-2292 (2013).
- [10] P. Dulio, C. Peri, Discrete tomography for inscribable lattice sets, Discrete Appl. Math., Vol. 161, (1959-1974 (2013).
- [11] P. Dulio, A. Frosini, S.M.C. Pagani, S. Rinaldi, Ambiguous reconstructions of hv-convex polyominoes Discrete Math., Vol. 343-10, (2020).
- [12] P. Dulio, and S.M.C. Pagani, A rounding theorem for unique binary tomographic reconstruction, Discrete Applied Mathematics, Vol. 268, 54 - 69 (2019).

- [13] P. C. Fishburn, J. C. Lagarias, J. A. Reeds, L. A. Shepp, Sets uniquely determined by projections on axes Discrete Math., Vol. 91, 149-159 (1991).
- [14] A. Frosini, L. Vuillon, *Tomographic reconstruction of 2-convex polyominoes using dual Horn clauses*, Theoret. Comput. Sci. Vol. 777, 329 337, (2019).
- [15] R. J. Gardner and P. Gritzmann, Discrete tomography: determination of finite sets by X-rays, Trans. Amer. Math. Soc. 349, 2271-2295 (1997).
- [16] L. Hajdu, R. Tijdeman, Agebraic aspects of discrete tomography, J. Reine Angew. Math., Vol. 534, 119-128, (2001).
- [17] Y. Gerard, Regular switching components, Theoret. Comput. Sci., Vol. 777, 338-355 (2019).
- [18] H. J. Ryser, Combinatorial properties of matrices of zeros and ones, Canadian Journal of Mathematics, Vol. 9, 371-377 (1957).
- [19] G. T. Herman and A. Kuba, Discrete tomography: Foundations algorithms and applications, Birkhauser, Boston (1999).
- [20] G. T. Herman and A. Kuba, Advances in Discrete Tomography and Its Applications, Birkhauser, Boston (2007).