Embedding Kozen-Tiuryn Logic into Residuated One-Sorted Kleene Algebra with Tests^{*}

Igor Sedlár and Johann J. Wannenburg

The Czech Academy of Sciences, Institute of Computer Science Pod Vodárenskou věží 271/2, Prague, The Czech Republic

Abstract

Kozen and Tiuryn have introduced the substructural logic S for reasoning about correctness of while programs (ACM TOCL, 2003). The logic S distinguishes between tests and partial correctness assertions, representing the latter by special implicational formulas. Kozen and Tiuryn's logic extends Kleene altebra with tests, where partial correctness assertions are represented by equations, not terms. Kleene algebra with codomain, KAC, is a one-sorted alternative to Kleene algebra with tests that expands Kleene algebra with an operator that allows to construct a Boolean subalgebra of tests. In this paper we show that Kozen and Tiuryn's logic embeds into the equational theory of the expansion of KAC with residuals of Kleene algebra multiplication and the upper adjoint of the codomain operator.

Keywords: Hoare logic, Kleene algebra with codomain, Kleene algebra with tests, Partial correctness, Substructural logic

1 Introduction

Kleene algebra with tests [10], KAT, is a simple algebraic framework for verifying properties of propositional while programs. KAT is two-sorted, featuring a Boolean algebra of tests embedded into a Kleene algebra of programs. The equational theory of KAT is PSPACE-complete [2] and KAT subsumes Propositional Hoare logic, PHL [11]. A PHL partial correctness assertion $\{b\} p\{c\}$, meaning that c holds after each terminating execution of program p in a state satisfying b, is represented in KAT by the equation $bp\bar{c} = 0$ or, equivalently, bp = bpc. The universal Horn theory (or the quasi-equational theory) of KAT consisting of quasi-equations of the form $E \longrightarrow p = q$ where E is a finite set of equations of the form r = 0 embeds into the equational theory of KAT [13] and is therefore PSPACE-complete as well.

Kozen and Tiuryn [14] introduce a substructural logic S that extends KAT and represents partial correctness assertions as implicational formulas $bp \Rightarrow c$. They argue that the implicational rendering of partial correctness assertions has certain advantages over the equational one, e.g. it facilitates a better distinction between local and

^{*}Preprint of: I. Sedlár and J.J. Wannenburg: Embedding Kozen-Tiuryn Logic into Residuated One-Sorted Kleene Algebra with Tests. In: A. Ciabattoni, E. Pimentel, R. J. G. B. de Queiroz (Eds.): *Proc. 28th Int. Conference of Logic, Language, Information, and Computation (WoLLIC 2022)*, pp. 221-236. LNCS 13468. Springer, 2022. (Available online at https://link.springer.com/chapter/10.1007/978-3-031-15298-6_14.)

global properties. PHL embeds into S and S is PSPACE-complete [12]. Kozen and Tiuryn prove that S induces a left-handed KAT structure on programs. However, a more thorough discussion of the relation of S to residuated extensions of Kleene algebra, such as Pratt's action logic [18], or to substructural logics in general is not provided. We believe that a deeper investigation of these connections would shed more light on the landscape of program logics and algebras.

In this paper we prove an embedding result that contributes to this. We note that a simple extension of KAT with a residual \rightarrow of the Kleene algebra multiplication does not enable an obvious embedding of S: implication formulas of S are test-like in the sense that they entail the multiplicative unit 1, but terms $p \rightarrow b$ of residuated KAT are not test-like, even if b is. To avoid this problem, we work with Kleene algebra with codomain, KAC [3, 5], a one-sorted alternative to KAT featuring a test-forming codomain operator c. In particular, we introduce an extension of KAC with both residuals of Kleene algebra multiplication and the upper adjoint e of the codomain operator, which we call SKAT. We show that $c(p \rightarrow e(b))$ behaves like $p \Rightarrow b$ of S. Our main technical result is that S embeds into the equational theory of SKAT^{*}, the class of algebras in SKAT that are based on *-continuous Kleene algebras.

The paper is structured as follows. Section 2 recalls KAT and S. Section 3 introduces a residuated version of KAT. We show that a naive translation of the implication connective of S into the residual operation does not constitute an embedding. In Section 4 we recall KAC and in Section 5 we introduce SKAT and SKAT^{*}. Section 6 establishes the embedding result. Section 7 concludes the paper and lists some interesting open problems.

2 KAT and S

In this section we recall KAT (Sect. 2.1) and we outline S (Sect. 2.2).

2.1 Kleene algebra with tests

We assume that the reader is familiar with the notion of an *idempotent semiring*.

Definition 1. A Kleene algebra [8] is an idempotent semiring $(K, \cdot, +, 1, 0)$ expanded with an operation $*: K \to K$ such that

$$1 + xx^* \le x^* \tag{1}$$

$$1 + x^* x \le x^* \tag{2}$$

$$y + xz \le z \implies x^* y \le z \tag{3}$$

$$y + zx \le z \implies yx^* \le z \tag{4}$$

A Kleene algebra is *-continuous iff it satisfies the condition

$$xy^*z = \sum_{n \ge 0} xy^n z \tag{5}$$

where $y^0 = 1$ and $y^{n+1} = y^n y$. $\sum X$ is the supremum of the set $X \subseteq K$ with respect to the partial order induced by Kleene algebra addition +.

It follows from the definition that x^* is the least element z such that $1 \le z, xz \le z$ and $zx \le z$. A standard example of a Kleene algebra is a relational Kleene algebra where K is a set of binary relations over some set S, \cdot is relational composition, + is set union, * is reflexive transitive closure, 1 is identity on S and 0 is the empty set. A relational Kleene algebra where K is the power set of some $S \times S$ will be called *fully relational*. Another standard example is the Kleene algebra of regular languages over a finite alphabet, where * corresponds to finite iteration (Kleene star). Relational Kleene algebras and Kleene algebras of regular languages are both *-continuous. Part of the definition of *-continuous Kleene algebras is the requirement that suprema of sets of elements of the form xy^nz exist. Not each Kleene algebra is *-continuous [7], but the equational theories of Kleene algebras and *-continuous Kleene algebras coincide [8].

Definition 2. A Kleene algebra with tests [10] is a structure $(K, B, \cdot, +, *, 1, 0, \overline{})$ such that

- $(K, \cdot, +, *, 1, 0)$ is a Kleene algebra,
- $B \subseteq K$, and
- $(B, \cdot, +, \bar{}, 1, 0)$ is a Boolean algebra.

A Kleene algebra with tests is *-continuous if its underlying Kleene algebra is.

Every Kleene algebra is a Kleene algebra with tests; take $B = \{0, 1\}$ and define $\bar{0} = 1, \bar{1} = 0$ and $\bar{x} = x$ for $x \notin \{1, 0\}$.¹ A standard example of a (*-continuous) Kleene algebra with tests is a relational Kleene algebra expanded with a Boolean subalgebra of the *negative cone*, i.e. the elements $x \leq 1$, which in the relational case are subsets of the identity relation. The class of Kleene algebras with tests is denoted as KAT and the class of *-continuous Kleene algebras with tests is denoted as KAT*. As in the test-free case, the equational theory of KAT is identical to the equational theory of KAT* [13].

Kleene algebras with tests are able to represent while programs and facilitate equational reasoning about their partial correctness:

- $\mathbf{skip} = 1$
- p;q = pq
- if b then p else $q = (bp) + (\overline{b}q)$
- while $b \operatorname{do} p = (bp)^* \overline{b}$
- $\{b\} p \{c\}$ corresponds to $bp\bar{c} = 0$

2.2 Substructural logic of partial correctness

This section outlines the substructural logic S [14] that allows to represent a partial correctness assertion by a formula (term) instead of an equation as in KAT. As with KAT, the logic S is many-sorted. Let $B = \{b_i \mid i \in \omega\}$ be the set of test variables and let $P = \{p_i \mid i \in \omega\}$ be the set of program variables. We define the following sorts of syntactic objects:

¹In contrast to the standard formulation of Kleene algebra with tests [10], we consider $\bar{}$ to be defined on all elements of K.

tests	$b,c := b_i \mid 0 \mid b \Rightarrow c$
programs	$p,q := p_i \mid b \mid p \oplus q \mid p \otimes q \mid p^+$
formulas	$f,g := b \mid p \Rightarrow f$
environments	$\Gamma, \Delta := \epsilon \mid \Gamma, p \mid \Gamma, f$
sequents	$\Gamma \vdash f$

We define $1 := 0 \Rightarrow 0$, $\neg b := b \Rightarrow 0$ and $p^* := 1 \oplus p^+$. We will sometimes write pq instead of $p \otimes q$ and \bar{b} instead of $\neg b$. Let $\mathsf{E} = \mathsf{B} \cup \mathsf{P}$ and let Ex_{S} , the set of S-expressions, be the union of the sets of formulas, programs and environments.

Kozen and Tiuryn introduce three kinds of semantics for their language: semantics based on guarded strings, traces and binary relations, respectively. We will work only with binary relational semantics.

Definition 3. A Kozen–Tiuryn model is a pair M = (W, V), where W is a non-empty set and $V : \mathsf{E} \to 2^{W \times W}$ such that $V(\mathsf{b}) \subseteq \mathrm{id}_W$ for all $\mathsf{b} \in \mathsf{B}$.

For each Kozen-Tiuryn model M, we define the M-interpretation function $[]_M : Ex_S \to 2^{W \times W}$ as follows:

- $[b]_M = V(b)$
- $[p]_M = V(p)$
- $[0]_M = \emptyset$
- $[b \Rightarrow c]_M = \{(s,s) \mid (s,s) \notin [b]_M \text{ or } (s,s) \in [c]_M\}$
- $[p \oplus q]_M = [p]_M \cup [q]_M$
- $[p \otimes q]_M = [p]_M \circ [q]_M$
- $[p^+]_M = [p]_M^+$
- $[p \Rightarrow f]_M = \{(s,s) \mid \forall t.(s,t) \in [p]_M \implies (t,t) \in [f]_M\}$
- $[\epsilon]_M = \mathrm{id}_W$
- $[\Gamma, \Delta]_M = [\Gamma]_M \circ [\Delta]_M$

(Here + denotes transitive closure and \circ denotes relational composition.)

A sequent $\Gamma \vdash f$ is valid in M iff, for all $s, t \in W$, if $(s, t) \in [\Gamma]_M$, then $(t, t) \in [f]_M$.

Observe that $[f]_M \subseteq \operatorname{id}_W$ for all formulas f; if $(s,s) \in [f]_M$, then we may say that formula f is true in s. Note that $[bp \Rightarrow c]_M$ is the set of (s,s) such that, for all t, if $(s,s) \in [b]_M$ and $(s,t) \in [p]_M$, then $(t,t) \in [c]_M$. Hence, $bp \Rightarrow c$ represents a partial correctness assertion: the formula is true in s iff b is true in s and p connects s with a state t only if c is true in t.

Figure 1 shows the sequent proof system for S. A sequent $\Gamma \vdash f$ is *provable in* S iff there is a finite sequence of sequents that ends with $\Gamma \vdash f$ each of which is either of the form (Id) or (I0), or is derived from previous sequents using some of the inference rules.

Theorem 1 (Kozen and Tiuryn). $\Gamma \vdash f$ is provable in S iff $\Gamma \vdash f$ is valid in all models M.

Figure 1: The sequent proof system for S.

S can be seen as a substructural logic [19]. From that point of view, S is an extension of the Multiplicative-additive Lambek calculus with the transitive closure operator $^+$. However, S contains some rules not usual from the substructural logic perspective, namely, the sort-specific weakening rules (Wf) and (Wp), and the implication-formula rules (TC) and (I \Rightarrow). It is therefore interesting to inquire how S relates to mainstream substructural logics. The embedding result of Section 6 is helpful in this respect.

3 Residuated KAT

The crucial difference between KAT and S is that the latter contains an implication operator \Rightarrow that is used to formalize partial correctness assertions as formulas (terms) of the language. Therefore, one may wonder if S can be thought of as KAT expanded with a residual of Kleene algebra multiplication. In this section we introduce residuated KAT, RKAT, and we show that a naive mapping from S to RKAT is not an embedding.

We note that residuated Kleene algebras with tests were studied also in [6]. In that framework, however, it is assumed by definition that the negative cone of the underlying Kleene algebra forms a Boolean algebra of tests. Kozen [10] argues, however, that this approach is not entirely satisfactory: tests are usually easily decidable (unlike other propositions, such as halting assertions), and not each Kleene algebra has a Boolean negative cone. Our definition of residuated Kleene algebras with tests adds residuals to KAT defined as usual.

Definition 4. A residuated idempotent semiring is a structure of the form $(K, \cdot, +, \rightarrow$

Figure 2: A residuated Kleene algebra with tests where $x \to 0$ is not test complementation.

 $, \hookrightarrow, 1, 0)$ where

- $(K, \cdot, +, 1, 0)$ is an idempotent semiring, and
- \rightarrow , \hookrightarrow are binary operations on K satisfying the residuation laws

$$xy \le z \iff x \le y \to z \iff y \le x \hookrightarrow z$$
 (6)

In residuated idempotent semirings, \rightarrow is called the *right residual* of \cdot and \rightarrow is called the *left residual*. An idempotent semiring is called *right-residuated* (*left-residuated*) if it is a reduct of a residuated idempotent semiring with only the right (left) residual. Note that residuated idempotent semirings necessarily have a top element: $x0 \leq 0$ implies $x \leq 0 \rightarrow 0$ and $0x \leq 0$ implies $x \leq 0 \rightarrow 0$.

The study of residuated semirings has some precedent. Pratt adds residuals to Kleene algebras to obtain *action algebras* [18], and Kozen adds a lattice meet operation to action algebras to obtain *action lattices* [9]; see also [6]. Equational theories of both action algebras and action lattices are undecidable [15].

Definition 5. A right-residuated Kleene algebra with tests is a structure of the form $(K, B, \cdot, +, *, \rightarrow, -, 1, 0)$ where

- $(K, B, \cdot, +, *, \bar{}, 1, 0) \in KAT$, and
- $(K, \cdot, +, \rightarrow, 1, 0)$ is a right-residuated idempotent semiring.

We will usually call right-residuated Kleene algebras with tests just residuated. The class of residuated Kleene algebras with tests will be denoted as RKAT.

One may be tempted to define \bar{x} as $x \to 0$, but this temptation is better resisted. Note that $0 \in B$ in every Kleene algebra with tests, but $0 \to 0$ is the top element. Hence, if we have a residuated Kleene algebra with tests where B does not contain the top element, then B is not closed under the operation $x \to 0$. An example of such an algebra is given in Figure 2, where $B = \{0, 1\}$ and + is join behaving as indicated in the diagram on the left: $0 \to 0 = \top$ but $\bar{0} = 1$.

For a similar reason, formulas $p \Rightarrow b$ of **S** do not directly translate into $p \rightarrow b$; the former entail 1, but the latter do not, as we have seen. However, a consideration of the relational case sheds some light on the matter. Recall that for $R \subseteq S \times S$ and $B \subseteq id_S$,

$$R \to B = \{(s, u) \mid \forall v. (u, v) \in R \implies (s, v) \in B\}$$
$$R \Rightarrow B = \{(u, u) \mid \forall v. (u, v) \in R \implies (v, v) \in B\}$$

Now consider two operations $c, e: 2^{S \times S} \to 2^{S \times S}$ such that

$$c(R) = \{(u, u) \mid \exists s. (s, u) \in R\}$$
(7)

$$e(R) = \{(s, u) \mid (u, u) \in R\}$$
(8)

Note that c and e form a *Galois connection* in the sense that, for all binary relations Q and R on S,

$$c(R) \subseteq Q \iff R \subseteq e(Q). \tag{9}$$

We have the following observation.

Proposition 1. For all $R \subseteq S \times S$ and $B \subseteq id_S$,

$$c(R \to e(B)) = R \Rightarrow B \tag{10}$$

Proof. If $(u, u) \in R \Rightarrow B$, then $(u, u) \in R \to e(B)$ since $(v, v) \in B$ iff $(u, v) \in e(B)$ by (8). But then obviously $(u, u) \in c(R \to e(B))$. Hence, $R \Rightarrow B \subseteq c(R \to e(B))$. Conversely, assume that $(u, u) \in c(R \to e(B))$ and $(u, v) \in R$, then $(s, u) \in R \to e(B)$ by (7) and so $(s, v) \in e(B)$. It follows that $(v, v) \in B$. Hence, $(u, u) \in R \Rightarrow B$ and so $c(R \to e(B)) \subseteq R \Rightarrow B$.

Hence, it seems that expanding RKAT with abstractions of c and e would make it possible to express the implication operator of S. An abstraction of c was added to semirings and Kleene algebras in the works on Kleene algebra with (co)domain [5, 3]. We discuss Kleene algebra with codomain in the next section.

4 Kleene algebra with codomain

Kleene algebra with codomain, KAC, is a one-sorted alternative to KAT: instead of assuming the existence of a Boolean subalgebra of tests, KAC introduces a unary *codomain* operation c such that the set of c(x) for x in the underlying Kleene algebra forms a Boolean algebra. One of the motivations for studying one-sorted alternatives to Kleene algebra with tests is that they seem to be better suited to automated theorem proving (see [5], p. 194). Our presentation follows [5, 3], where a symmetric variant of KAC, namely, Kleene algebra with *domain*, was studied.

Kleene algebras with codomain add to Kleene algebras a unary operator that generalizes the *anticodomain* operator on binary relations:

$$a(R) = \{(u, u) \mid \neg \exists s.(s, u) \in R\}.$$
(11)

Note that, assuming (7), c(R) = a(a(R)).

Definition 6. A Kleene algebra with codomain *is an algebra of the form* $(K, \cdot, +, *, a, 1, 0)$ where $a : K \to K$ such that

$$x \cdot \mathsf{a}(x) \le 0 \tag{12}$$

$$\mathsf{a}(x \cdot y) \le \mathsf{a}\big(\mathsf{a}(\mathsf{a}(x)) \cdot y\big) \tag{13}$$

$$\mathsf{a}(x) + \mathsf{a}(\mathsf{a}(x)) = 1 \tag{14}$$

We define c(x) := a(a(x)).

Note that the relational anticodomain operator (11) satisfies the above equations.

Proposition 2. In every Kleene algebra with codomain, the operator c satisfies the following codomain equations:

$$x \le x \cdot \mathbf{c}(x) \tag{15}$$

$$c(xy) = c(c(x) \cdot y) \tag{16}$$

$$\mathsf{c}(x) \le 1 \tag{17}$$

$$\mathsf{c}(0) = 0 \tag{18}$$

$$c(x+y) = c(x) + c(y)$$
(19)

Moreover, c has the following properties:

$$\mathsf{c}(\mathsf{c}(x)) = \mathsf{c}(x) \tag{20}$$

$$\mathbf{c}(x) \cdot \mathbf{c}(x) = \mathbf{c}(x) \tag{21}$$

$$\mathsf{c}(xy) \le \mathsf{c}(y) \tag{22}$$

$$x \le x \mathbf{c}(y) \implies \mathbf{c}(x) \le \mathbf{c}(y) \tag{23}$$

Proof. For the codomain equations, see [5], Theorem 8.7, and the more detailed proof of the theorem in Appendix D of [4]. The arguments there are given for the domain and antidomain operators, but similar arguments work for codomain and anticodomain.

The fixpoint property (20) follows from $c(xy) = c(c(x) \cdot y)$ and idempotence (21) follows from the fixpoint property, $x \leq x \cdot c(x)$, and $c(x) \leq 1$. The codomain restriction property (22) is established by noting that $c(xy) \leq c(c(x)y) \leq c(1y) \leq c(y)$. The least preserver property (23) is established by noting that $x \leq xc(y)$ implies $c(x) \leq c(xc(y)) \leq c(c(y)) \leq c(y)$.

Proposition 3. Let $c(K) = \{x \mid x \in K \& \exists y.(x = c(y))\}$. In every Kleene algebra with codomain,

- 1. $(c(K), \cdot, +, 1, 0)$ is a subalgebra of $(K, \cdot, +, 1, 0)$;
- 2. $(c(K), \cdot, +, 1, 0)$ is a bounded distributive lattice;
- 3. c(x) + a(c(x)) = 1 and $c(x) \cdot a(c(x)) = 0$.

Hence, $c(\mathscr{K}) = (c(K), \cdot, +, a, 1, 0)$ is a Boolean algebra.

Proof. 1. follows from c(1) = 1, c(0) = 0, c(x) + c(y) = c(x + y), and $c(x) \cdot c(y) = c(c(x) \cdot c(y))$. We verified the latter using Prover 9 [16]. 2. follows from 1., together with $c(x) \le 1$ and c(x) = c(x)c(x) of Proposition 2. 3. follows from the axioms for a.

It can be shown that every full relational KAT can be seen as a KAC where a is the relational anticodomain operation, $a(R) := \{(u, u) \mid \neg \exists s. (s, u) \in R\}$. Moreover, Proposition 3 shows that every KAC gives rise to a KAT. A more detailed argument involving a translation from the language of KAT into the language of KAC then shows that the equational theory of KAT embeds into the equational theory of KAC. We omit the details.

5 SKAT: Residuated KAC with a Galois connection

In this section, we introduce SKAT, an extension of KAC with both residuals of Kleene algebra multiplication and the upper adjoint of the codomain operator. The *-continuous variant of SKAT will be denoted as SKAT*. The next section shows that S embeds into the equational theory of SKAT*.

Definition 7. An S-type Kleene algebra with codomain is a structure of the form $(K, \cdot, +, \rightarrow, \rightarrow *, \mathsf{a}, \mathsf{e}, 1, 0)$ where

- $(K, \cdot, +, \hookrightarrow, \rightarrow, *, 1, 0)$ is a residuated Kleene algebra,
- $(K, \cdot, +, *, a, 1, 0)$ is a Kleene algebra with codomain, and
- e is a unary operation on K that satisfies the following:

$$\mathsf{a}(\mathsf{a}(\mathsf{e}(x))) \le x \tag{24}$$

$$x \le \mathsf{e}(\mathsf{a}(\mathsf{a}(x))) \tag{25}$$

$$\mathsf{e}(x) \le \mathsf{e}(x+y) \tag{26}$$

An S-type Kleene algebra with codomain is *-continuous iff its underlying Kleene algebra is *-continuous. The class of S-type Kleene algebras with codomain will be denoted as SKAT and the class of the *-continuous ones as SKAT*.

We will call S-type Kleene algebras with codomain simply SKAT-algebras. We will usually omit bracketing between elements of $\{a, e\}$, writing ec(x) instead of e(c(x)), for instance.

Every SKAT-algebra is a Kleene algebra with codomain by definition, and so the test algebra $(c(\mathcal{K}), \cdot, +, a, 1, 0)$ of each SKAT-algebra \mathcal{K} is a Boolean algebra. It is well known that residuated Kleene algebras form a variety [18], and so SKAT forms a variety.

Proposition 4. The following hold in each SKAT-algebra:

$$\mathsf{c}(x) \le y \iff x \le \mathsf{e}(y) \tag{27}$$

$$\mathsf{c}(x \to y) \le x \to x\mathsf{c}(y) \tag{28}$$

$$\mathsf{a}(x) = \mathsf{c}(\mathsf{c}(x) \to 0) \tag{29}$$

Proof. (27) follows directly from the SKAT axioms concerning e. (28) is established as follows. Take any z such that c(z) = z; note that $c(x \to y)$ is of this form. We reason as follows:

$$z \leq \mathsf{c}(x \to y) \implies zx \leq \mathsf{c}(x \to y)x \stackrel{(16)}{\Longrightarrow} \mathsf{c}(zx) \leq \mathsf{c}((x \to y)x)$$
$$\implies \mathsf{c}(zx) \leq \mathsf{c}(y) \implies zx\mathsf{c}(zx) \leq \mathsf{c}(z)x\mathsf{c}(y) \implies zx \leq x\mathsf{c}(y)$$
$$\implies z \leq x \to x\mathsf{c}(y) \implies \mathsf{c}(z) \leq \mathsf{c}(x \to x\mathsf{c}(y)) \implies z \leq \mathsf{c}(x \to x\mathsf{c}(y)).$$

(29) is established as follows. Note that ac(x) = a(x) by (13) and c(x)a(x) = a(x)c(x) by Proposition 3. Hence, $a(x) \le c(x) \to 0$ by (12) and residuation. It follows that $ca(x) \le c(c(x) \to 0)$ and so $a(x) \le c(c(x) \to 0)$.

The converse inequality follows from

$$x\mathbf{c}(y) = 0 \implies \mathbf{c}(x) \le \mathbf{a}(y)$$
 (30)

Figure 3: A SKAT-algebra falsifying (31).

In particular, $c(c(x) \to 0)c(x) = 0$ by (28), and so $cc(c(x) \to 0) \le a(x)$ by (30), which means that $c(c(x) \to 0) \le a(x)$ by the fixpoint property (20). The quasi-equation (30) is established similarly as the quasi-equation (26) on p. 386 of [4]. If xc(y) = 0, then a(c(x)c(y)) = 1 and so c(x)c(y) = 0. But then

$$\mathsf{c}(x) = \mathsf{c}(x)(\mathsf{c}(y) + \mathsf{a}(y)) = \mathsf{c}(x)\mathsf{c}(y) + \mathsf{c}(x)\mathsf{a}(y) = \mathsf{c}(x)\mathsf{a}(y) \le \mathsf{a}(y) \,.$$

We note that an extension of Kleene algebra with domain with a right residual of Kleene algebra multiplication is considered in [4, 5]. However, the goal there is to induce a Heyting algebra of tests, and \rightarrow is assumed to be a partial operation, defined only on the test algebra. It is also assumed that the test algebra is closed under \rightarrow , which we may transpose to the setting that uses codomain instead of domain as follows:

$$c(c(x) \to c(y)) = c(x) \to c(y)$$
(31)

In our setting, \rightarrow is a total operation which is not included in the signature of (Boolean) test algebras. It can also be shown that (31) does not hold; see Fig. 3. Operations \cdot and * are defined as in Fig. 2, \hookrightarrow is identical to \rightarrow . It is easily verified that $c(0) \rightarrow c(0) = \top$, and $c(\top) = 1$.

6 The embedding result

In this section we prove that S embeds into the equational theory of SKAT*.

Let $X = \{x_i \mid i \in \omega\}$ be a countable set of variables. The set of terms Tm is the set of expressions formed using the following grammar:

$$p,q := \mathsf{x}_i \mid 1 \mid 0 \mid p \cdot q \mid p + q \mid p \to q \mid p^* \mid \mathsf{c}(p) \mid \mathsf{e}(p) \,.$$

(We abuse the notation by re-using the same variables that range over programs in S, but this will hopefully cause no confusion.) An equation is an ordered pair of terms, denoted as $p \approx q$. We use $p \preceq q$ as short for $p + q \approx q$. If $\mathscr{K} \in \mathsf{SKAT}$, then a \mathscr{K} -valuation is any homomorphism from Tm into \mathscr{K} . An equation $p \approx q$ is satisfied by a \mathscr{K} -valuation [] iff [p] = [q]. The equational theory of SKAT (SKAT^*) is the set of equations that are satisfied by all \mathscr{K} valuations where $\mathscr{K} \in \mathsf{SKAT}$ ($\mathscr{K} \in \mathsf{SKAT}^*$).

Definition 8. We define $Tr : Ex_S \to Tm$ as follows:

- $Tr(\mathbf{p}_n) = \mathbf{x}_{2n}$
- $Tr(\mathbf{b}_n) = \mathbf{c}(\mathbf{x}_{2n+1})$
- Tr(0) = c(0)

- $Tr(b \Rightarrow c) = c(Tr(b) \rightarrow e(Tr(c)))$
- $Tr(p \oplus q) = Tr(p) + Tr(q)$
- $Tr(p \otimes q) = Tr(p) \cdot Tr(q)$
- $Tr(p^+) = Tr(p) \cdot Tr(p)^*$
- $Tr(p \Rightarrow f) = c(Tr(p) \rightarrow e(Tr(f)))$
- $Tr(\epsilon) = 1$
- $Tr(\Gamma, \Delta) = Tr(\Gamma) \cdot Tr(\Delta)$

Theorem 2. A sequent $\Gamma \vdash f$ is provable in S iff $c(Tr(\Gamma)) \leq Tr(f)$ belongs to the equational theory of SKAT^{*}.

Theorem 2 is established in two steps. First we show that every model M of S with a valuation [] can be transformed into a $\mathcal{M} \in \mathsf{SKAT}^*$ and a \mathcal{M} -valuation [] such that $[\alpha] = \llbracket Tr(\alpha) \rrbracket$ for all $\alpha \in Ex_S$. Then we show by induction on the length of S-proofs that if $\Gamma \vdash f$ is provable in S, then $\mathsf{c}(Tr(\Gamma)) \preceq Tr(f)$ belongs to the equational theory of SKAT^* .

Lemma 1. Let M = (W, V) be a model of S and let [] be a *M*-interpretation. Then there is $\mathcal{M} \in \mathsf{SKAT}^*$ and a \mathcal{M} -valuation [] such that, for all $\alpha \in Ex_S$,

$$[\alpha] = \llbracket Tr(\alpha) \rrbracket \tag{32}$$

Proof. Fix M = (W, V). Let \mathscr{M} be the full relational residuated Kleene algebra of binary relations on W; define $a(R) = \{(u, u) \mid \neg \exists s. (s, u) \in R\}$ and $e(R) = \{(s, u) \mid (u, u) \in R\}$; note that $c(R) = \{(u, u) \mid \exists s. (s, u) \in R\}$.² It is easily checked that $\mathscr{M} \in \mathsf{SKAT}^*$. In order to prove (24), note that $(s, t) \in \mathsf{ce}(R)$ only if s = t and there is u such that $(u, t) \in \mathsf{e}(R)$, which holds only if s = t and $(t, t) \in R$. In order to prove (25), note that $(s, t) \in R$ implies $(t, t) \in \mathsf{c}(R)$, which implies $(s, t) \in \mathsf{ec}(R)$. Finally, e is obviously monotonic, which entails (26).

Define $\llbracket I]$ as the unique SKAT homomorphism from Tm to \mathscr{M} satisfying the following, for all $n \in \omega$:

- $\llbracket x_{2n} \rrbracket = \llbracket p_n \rrbracket;$
- $[x_{2n+1}] = [b_n].$

We will prove (32) by induction on α . First we prove that (32) holds for all tests b by induction on b. The base case is established as follows. We know by definition that $\llbracket Tr(\mathbf{b}_n) \rrbracket = \mathbf{c}\llbracket \mathbf{x}_{2n+1} \rrbracket = \mathbf{c}\llbracket \mathbf{b}_n \rrbracket$. The latter is $\{(u, u) \mid \exists s. (s, u) \in \llbracket \mathbf{b}_n \rrbracket \} = \llbracket \mathbf{b}_n \rrbracket$, since $\llbracket \mathbf{b}_n \rrbracket \subseteq \operatorname{id}_W$. The induction step is established as follows. $\llbracket Tr(0) \rrbracket = \mathbf{c}\llbracket 0 \rrbracket = \emptyset = \llbracket 0 \rrbracket$ and the other case uses (10) of Proposition 1 (we are working with a relational Kleene algebra):

$$\llbracket Tr(b \Rightarrow c) \rrbracket = \mathsf{c}(\llbracket Tr(b) \rrbracket \to \mathsf{e}\llbracket Tr(c) \rrbracket)$$
$$= \mathsf{c}(\llbracket b \rrbracket \to \mathsf{e}\llbracket c \rrbracket)$$
$$\stackrel{(10)}{=} \llbracket b \Rightarrow c \rrbracket$$

²Recall that $R \to Q = \{(s, u) \mid \forall v. (u, v) \in R \implies (s, u) \in Q\}$ and $R \hookrightarrow Q = \{(s, u) \mid \forall v. (v, s) \in R \implies (v, u) \in Q\}.$

This shows that (32) holds for all tests b. The fact that it holds for all programs is established by easy induction (Tr virtually commutes with Kleene algebra operations). Next, we show that (32) holds for all formulas. The only thing to prove is the induction step for $\alpha = p \Rightarrow f$. The claim can be established using (10) as before. The claim for environments is trivial.

Lemma 2. If $\Gamma \vdash f$ is provable in S, then $c(Tr(\Gamma)) \preceq Tr(f)$ belongs to the equational theory of SKAT^{*}.

Proof. We reason by induction on the length of proofs. We note that the proof of this lemma, namely the claim for the (I^+) rule, is the only point where the assumption of *-continuity seems to be required.

A useful fact to note is that Tr(f) is of the form c(p) for all formulas f. By a translation of a sequent $\Gamma \vdash f$ we mean the equation $c(Tr(\Gamma)) \preceq Tr(f)$. We will prove that translations of axioms of S are in the equational theory of SKAT^{*}, and that if translations of all premises of an inference rule of S are in the equational theory of SKAT^{*}, then so is the consequence of the rule. (If $p \preceq q$ is in the equational theory of SKAT^{*}, then we say that $p \preceq q$ is valid.)

(Id) $c(Tr(b)) \leq Tr(b)$ is valid since $cc(x) \leq c(x)$ by (20).

(I0) $\mathsf{c}(Tr(\Gamma) \cdot Tr(0) \cdot Tr(\Delta)) \preceq Tr(f)$ is valid since $Tr(0) = \mathsf{c}(0) = 0$.

(TC)
$$\frac{c(Tr(\Gamma)Tr(b)Tr(\Delta)) \leq Tr(f) \quad c(Tr(\Gamma)Tr(b \Rightarrow 0)Tr(\Delta)) \leq Tr(f)}{c(Tr(\Gamma)Tr(\Delta)) \leq Tr(f)}$$
preserves validity thanks to (19) and (29):

$$c(xc(y)z) \leq a \text{ and } c(xc(c(y) \to 0)z) \leq a$$

$$\stackrel{(19)}{\Longrightarrow} \quad c(xc(y)z + xc(c(y) \to 0)z) \leq a$$

$$\stackrel{(29)}{\Longrightarrow} \quad c(xc(y)z + xa(y)z) \leq a$$

$$\stackrel{(14)}{\Longrightarrow} \quad c(xz) \leq a$$
(cut)
$$\frac{c(Tr(\Gamma)) \leq Tr(g) \quad c(Tr(\Gamma)Tr(g)Tr(\Delta)) \leq Tr(f)}{c(Tr(\Gamma)Tr(\Delta)) \leq Tr(f)}$$
preserves validity thanks to (15) and (19):

$$c(x) \leq a_2 \text{ and } c(xa_2y) \leq a_1$$

$$\stackrel{(15)}{\Longrightarrow} xy \leq xa_2y \text{ and } c(xa_2y) \leq a_1$$

 $\stackrel{(19)}{\Longrightarrow}$ $\mathsf{c}(xy) \leq \mathsf{c}(xa_2y)$ and $\mathsf{c}(xa_2y) \leq a_1$

 $\implies \mathsf{c}(xy) \le a_1$

$$(\mathbf{R}\Rightarrow) \quad \frac{\mathsf{c}(Tr(\Gamma)Tr(p)) \preceq Tr(f)}{\mathsf{c}(Tr(\Gamma)) \preceq Tr(p \Rightarrow f)} \text{ preserves validity thanks to (27) and (19):}$$
$$\mathsf{c}(xy) \le z \quad \stackrel{(27)}{\Longrightarrow} \quad xy \le \mathsf{e}(z)$$
$$\implies x \le y \to \mathsf{e}(z) \quad \stackrel{(19)}{\Longrightarrow} \quad \mathsf{c}(x) \le \mathsf{c}(y \to \mathsf{e}(z))$$

(I \Rightarrow) $\frac{\mathsf{c}(Tr(\Gamma)Tr(p)Tr(f)Tr(\Delta)) \preceq Tr(g)}{\mathsf{c}(Tr(\Gamma)Tr(p \Rightarrow f)Tr(p)Tr(\Delta)) \preceq Tr(g)}$ preserves validity thanks to (28) and the fact that (27) entails $\mathsf{cec}(y) \le \mathsf{c}(y)$:

$$c(x \to ec(y))x \le (x \to x cec(y))x \le x cec(y) \le xc(y)$$

(Recall that $Tr(p \Rightarrow f) = c(q \rightarrow ec(r))$ for some terms q, r.)

(I \otimes) follows directly from the definition of Tr, and so do (E \otimes), (I \oplus), (E \oplus_1) and (E \oplus_2).

(I⁺)
$$\frac{\mathsf{c}(Tr(g)Tr(p)) \preceq Tr(f) \quad \mathsf{c}(Tr(g)Tr(p)) \preceq Tr(g)}{\mathsf{c}(Tr(g)Tr(p)Tr(p)^*) \preceq Tr(f)}.$$

It can be proven by induction on n, using (16), that if $c(yx) \leq c(z)$ and $c(yx) \leq y$, then $\forall n \geq 0. c(yxx^n) \leq c(z)$. By (27), $\forall n \geq 0. yxx^n \leq ec(z)$, and so by *-continuity $yxx^* \leq ec(z)$, which means that $c(yxx^*) \leq c(z)$. (Note that *-continuity entails that infinite suprema of elements of the form xy^*z exist.) In fact, this is the only point where we need to assume *-continuity.

(E⁺) follows from the fact that $x \le xx^*$ in Kleene algebras; and (CC⁺) follows from the fact that $x^*xx^* \le x^*$.

(Wf)
$$\frac{\mathsf{c}(Tr(\Gamma)Tr(\Delta)) \preceq Tr(g)}{\mathsf{c}(Tr(\Gamma)Tr(f)Tr(\Delta)) \preceq Tr(g)}$$
 preserves validity thanks to $\mathsf{c}(x) \le 1$.

(Wp)
$$\frac{\mathsf{c}(Tr(\Gamma)) \preceq Tr(f)}{\mathsf{c}(Tr(p)Tr(\Gamma)) \preceq Tr(f)}$$
 follows from (22).

This concludes the proof of the lemma.

Now we are in the position to prove Theorem 2.

Proof. If $\Gamma \vdash f$ is not provable in S, then there is a model M and an interpretation function [] such that $(w, u) \in [\Gamma]$ and $(u, u) \notin [f]$ by Kozen and Tiuryn's relational completeness theorem. By Lemma 1, there is $\mathscr{M} \in \mathsf{SKAT}^*$ and a \mathscr{M} -valuation $\llbracket \mathbb{I}$ such that $(u, u) \in \llbracket \mathsf{c}(Tr(\Gamma)) \rrbracket$ and $(u, u) \notin \llbracket Tr(f) \rrbracket$. Hence, $\mathsf{c}(Tr(\Gamma)) \preceq Tr(f)$ is not in the equational theory of SKAT*. Conversely, if $\Gamma \vdash f$ is provable in S, then $\mathsf{c}(Tr(\Gamma)) \preceq Tr(f)$ is in the equational theory of SKAT* by Lemma 2.

7 Conclusion

We have shown in this paper that Kozen and Tiuryn's substructural logic of partial correctness S embeds into the equational theory of an extension of *-continuous Kleene algebra with codomain with both residuals of the Kleene algebra multiplication and the upper adjoint of the codomain operator, SKAT*. We believe that this result sheds more light on the landscape of program logics and algebras.

A number of interesting problems remain open:

- 1. Does S embed into the equational theory of SKAT?
- 2. Which residuated Kleene algebras extend to algebras in SKAT or SKAT*? Is the equational theory of SKAT (SKAT*) a conservative extension of the equational theory of residuated (*-continuous) Kleene algebras?
- 3. Related to the previous problem, is the equational theory of SKAT^{*} identical to the equational theory of SKAT? The equational theory of *-continuous residuated Kleene algebras is not identical to the equational theory of residuated Kleene algebras [1].
- 4. What are the free algebras in SKAT and SKAT^{*}? McLean [17] shows that regular sets of labelled pointed trees form free Kleene algebras with domain, and a version of his construction may turn out to apply to our case as well.
- 5. What is the complexity of the equational theories of SKAT and SKAT^{*}? The equational theory of *-continuous residuated Kleene algebras is Π_1^0 -complete (hence not r.e.) and the equational theory of residuated Kleene algebras is Σ_1^0 -complete.
- 6. Consider the set of terms generated by the following restricted grammar:

$$p, q := \mathsf{x}_i \mid 1 \mid 0 \mid p \cdot q \mid p + q \mid p^* \mid \mathsf{c}(p) \mid \mathsf{c}(p \to \mathsf{ec}(q)).$$

Is the fragment of the equational theory of SKAT restricted to terms generated by this grammar decidable?

7. What are the Gentzen systems sound and (weakly) complete for SKAT, in the style of [6]?

Acknowledgement

We are grateful to two anonymous reviewers for valuable suggestions. The first author is grateful to Vít Punčochář for discussions that led to the work on this paper, and to the audience at the Proof Theory Seminar of the Steklov Mathematical Institute for illuminating discussion after his talk on 14 February 2022. The work of the second author was carried out within the project *Supporting the Internationalization of the Institute of Computer Science of the Czech Academy of Sciences* (no. CZ.02.2.69/0.0/0.0/18_053/0017594), funded by the Operational Programme Research, Development and Education of the Ministry of Education, Youth and Sports of the Czech Republic. The project is co-funded by the EU.

References

- W. Buszkowski. On Action Logic: Equational Theories of Action Algebras. Journal of Logic and Computation, 17(1):199–217, 10 2006.
- [2] E. Cohen, D. Kozen, and F. Smith. The complexity of Kleene algebra with tests. Technical Report TR96-1598, Computer Science Department, Cornell University, July 1996.
- [3] J. Desharnais, B. Möller, and G. Struth. Kleene algebra with domain. ACM Trans. Comput. Logic, 7(4):798–833, oct 2006.
- [4] J. Desharnais and G. Struth. Modal semirings revisited. In International Conference on Mathematics of Program Construction, pages 360–387. Springer, 2008.
- [5] J. Desharnais and G. Struth. Internal axioms for domain semirings. Science of Computer Programming, 76(3):181–203, 2011. Special issue on the Mathematics of Program Construction (MPC 2008).
- [6] P. Jipsen. From semirings to residuated Kleene lattices. Studia Logica, 76(2):291– 303, Mar 2004.
- [7] D. Kozen. On Kleene algebras and closed semirings. In B. Rovan, editor, International Symposium on Mathematical Foundations of Computer Science, pages 26–47. Springer, 1990.
- [8] D. Kozen. A completeness theorem for Kleene algebras and the algebra of regular events. *Information and Computation*, 110(2):366 390, 1994.
- [9] D. Kozen. On action algebras. In Logic and Information Flow, pages 78 88. MIT Press, 1994.
- [10] D. Kozen. Kleene algebra with tests. ACM Trans. Program. Lang. Syst., 19(3):427–443, May 1997.
- [11] D. Kozen. On Hoare logic and Kleene algebra with tests. ACM Trans. Comput. Logic, 1(1):60–76, July 2000.
- [12] D. Kozen. Automata on guarded strings and applications. Technical report, Cornell University, 2001.
- [13] D. Kozen and F. Smith. Kleene algebra with tests: Completeness and decidability. In D. van Dalen and M. Bezem, editors, *Computer Science Logic*, pages 244–259, Berlin, Heidelberg, 1997. Springer Berlin Heidelberg.
- [14] D. Kozen and J. Tiuryn. Substructural logic and partial correctness. ACM Trans. Computational Logic, 4(3):355–378, July 2003.
- [15] S. Kuznetsov. Action logic is undecidable. ACM Trans. Comput. Logic, 22(2):1– 26, May 2021.
- [16] W. McCune. Prover9 and Mace4. http://www.cs.unm.edu/~mccune/prover9/, 2005-2010.

- [17] B. McLean. Free kleene algebras with domain. Journal of Logical and Algebraic Methods in Programming, 117:100606, 2020.
- [18] V. Pratt. Action logic and pure induction. In Logics in AI. European workshop JELIA '90, Amsterdam, the Netherlands, September 10-14, 1990. Proceedings, pages 97–120. Berlin etc.: Springer-Verlag, 1991.
- [19] G. Restall. An Introduction to Substructural Logics. Routledge, London, 2000.