Monitoring edge-geodetic sets in graphs

Subhadeep R. Dev^{*}

Sanjana Dey[†] Florent Foucaud^{‡§} K Lekshmi Ramasubramony Sulochana $^{\parallel}$

Krishna Narayanan[¶]

October 31, 2023

Abstract

We introduce a new graph-theoretic concept in the area of network monitoring. In this area, one wishes to monitor the vertices and/or the edges of a network (viewed as a graph) in order to detect and prevent failures. Inspired by two notions studied in the literature (edge-geodetic sets and distance-edge-monitoring sets), we define the notion of a monitoring edge-geodetic set (MEG-set for short) of a graph G as an edge-geodetic set $S \subseteq V(G)$ of G (that is, every edge of G lies on some shortest path between two vertices of S) with the additional property that for every edge e of G, there is a vertex pair x, y of S such that e lies on all shortest paths between x and y. The motivation is that, if some edge e is removed from the network (for example if it ceases to function), the monitoring probes x and y will detect the failure since the distance between them will increase.

We explore the notion of MEG-sets by deriving the minimum size of a MEG-set for some basic graph classes (trees, cycles, unicyclic graphs, complete graphs, grids, hypercubes, corona products...) and we prove an upper bound using the feedback edge set of the graph.

We also show that determining the smallest size of an MEG-set of a graph is NP-hard, even for graphs of maximum degree at most 9.

1 Introduction

We introduce a new graph-theoretic concept, that is motivated by the problem of network monitoring, called *monitoring edge-geodetic sets*. In the area of network monitoring, one wishes to detect or repair faults in a network; in many applications, the monitoring process involves distance probes [2, 3, 4, 10]. Our networks are modeled by finite, undirected simple connected graphs, whose vertices represent systems and whose edges represent the connections between them. We wish to monitor a network such that when a connection (an edge) fails, we can detect the said failure by means of certain probes. To do this, we select a small subset of vertices (representing the probes) of the network such that all connections are covered by the shortest paths between pairs of vertices in the network. Moreover, any two probes are able to detect the current distance that separates them. The goal is that, when an edge of the network fails, some pair of probes detects a change in their distance value, and therefore the failure can be detected. Our inspiration comes from two areas: the concept of *geodetic sets* in graphs and its variants [12], and the concept of *distance edge-monitoring sets* [9, 10].

We now proceed with some necessary definitions. A geodesic is a shortest path between two vertices u, v of a graph G [20]. The length of a geodesic between two vertices u, v in G is the distance $d_G(u, v)$

^{*}Indian Statistical Institute, Kolkata, India

[†]National University of Singapore, Singapore

[‡]Université Clermont Auvergne, CNRS, Clermont Auvergne INP, Mines Saint-Etienne, LIMOS, 63000 Clermont-Ferrand, France.

 $^{^{\$}}$ Research financed by the French government IDEX-ISITE initiative 16-IDEX-0001 (CAP 20-25) and by the ANR project GRALMECO (ANR-21-CE48-0004).

[¶]Department of Mathematics, Simon Fraser University, Burnaby, Canada

PSG College of Technology, Coimbatore, India

between them. For an edge e of G, we denote by G-e the graph obtained by deleting e from G. An edge e in a graph G is a *bridge* if G-e has more connected components than G. The *open neighborhood* of a vertex $v \in V(G)$ is $N_G(v) = \{u \in V \mid uv \in E(G)\}$ and its *closed neighborhood* is the set $N_G[v] = N_G(v) \cup \{v\}$.

Monitoring edge-geodetic sets. We now formally define our main concept.

Definition 1.1. Two vertices x, y monitor an edge e in graph G if e belongs to all shortest paths between x and y. A set S of vertices of G is called a monitoring edge-geodetic set of G (MEG-set for short) if, for every edge e of G, there is a pair x, y of vertices of S that monitors e.

We denote by meg(G) the size of a smallest MEG-set of G. We note that V(G) is always an MEG-set of G, thus meg(G) is always well-defined.

Related notions. A set S of vertices of a graph G is a geodetic set if every vertex of G lies on some shortest path between two vertices of S [12]. An edge-geodetic set of G is a set $S \subseteq V(G)$ such that every edge of G is contained in a geodesic joining some pair of vertices in S [19]. A strong edge-geodetic set of G is a set S of vertices of G such that for each pair u, v of vertices of S, one can select a shortest u - vpath, in a way that the union of all these $\binom{|S|}{2}$ paths contains E(G) [16]. It follows from these definitions that any strong edge-geodetic set is an edge-geodetic set, and any edge-geodetic set is a geodetic set (if the graph has no isolated vertices). In fact, every MEG-set is a strong edge-geodetic set. Indeed, given an MEG-set S, one can choose any shortest path between each pair of vertices of S, and the set of these paths covers E(G). Indeed, every edge of G is contained in all shortest paths between some pair of S. Hence, MEG-sets can be seen as an especially strong form of strong edge-geodetic sets.

A set S of vertices of a graph G is a distance-edge monitoring set if, for every edge e, there is a vertex x of S and a vertex y of G such that e lies on all shortest paths between x and y [9, 10]. Thus, it follows immediately that any MEG-set of a graph G is also a distance-edge monitoring set of G.

Our results. We start by presenting some basic lemmas about the concept of MEG-sets in Section 2, that are helpful for understanding this concept. We then study in Section 3 the optimal value of meg(G) when G is a tree, cycle, unicyclic graph, complete (multipartite) graph, hypercubes, grids and corona products. In Section 4, we show that meg(G) is bounded above by a linear function of the *feedback edge set number* of G (the smallest number of edges of G needed to cover all cycles of G, also called *cyclomatic number*) and the number of leaves of G. This implies that meg(G) is bounded above by a function of the *max leaf number* of G (the maximum number of leaves in a spanning tree of G). These two parameters are popular in structural graph theory and in the design of algorithms. We refer to Figure 1 for the relations between parameter meg and other graph parameters. We show that determining meg(G) is an NP-complete problem, even in graphs of maximum degree at most 9, in Section 5. Finally, we conclude in Section 6.

Further related work on MEG-sets. This paper is the full version of a paper presented at the CALDAM 2023 conference [11], where the notion of MEG-sets was presented for the first time. It contains the full proofs of the results in the conference version (with a corrected version of Theorem 3.5), as well as the new result on corona products, and the new NP-completeness proof. In the meantime, Haslegrave [14] proved that the MEG-set problem is NP-complete on general graphs, and also studied MEG-sets for other graph products.

2 Preliminary lemmas

We now give some useful lemmas about the basic properties of MEG-sets. A vertex is *simplicial* if its neighborhood forms a clique.

Lemma 2.1. In a graph G with at least one edge, any simplicial vertex belongs to any edge-geodetic set and thus, to any MEG-set of G.



Figure 1: Relations between the parameter meg and other structural parameters in graphs (with no isolated vertices). For the relationships of distance edge-monitoring sets, see [9, 10]. Arcs between parameters indicate that the value of the bottom parameter is upper-bounded by a function of the top parameter.

Proof. Let us consider by contradiction an MEG-set of G that does not contain said simplicial vertex v. Any shortest path passing through its neighbors will not pass through v, because all the neighbors are adjacent, hence leaving the edges incident to v uncovered, a contradiction.

Two distinct vertices u and v of a graph G are open twins if N(u) = N(v) and closed twins if N[u] = N[v]. Further, u and v are twins in G if they are open twins or closed twins in G.

Lemma 2.2. If two vertices are twins of degree at least 1 in a graph G, then they must belong to any MEG-set of G.

Proof. For any pair u, v of open twins in G, for any shortest path passing through u, there is another one passing through v. Thus, if u, v were not part of the MEG-set, then the edges incident to u and v would remain unmonitored, a contradiction.

If u, v are closed twins, if some shortest path contains the edge uv, then it must be of length 1 and consist of the edge uv itself (otherwise there would be a shortcut). Thus, to monitor uv, both u, v must belong to any MEG-set.

The next two lemmas concern cut-vertices and subgraphs, and will be useful in some of our proofs.

Lemma 2.3. Let G be a graph with a cut-vertex v and C_1, C_2, \ldots, C_k be the k components obtained when removing v from G. If S_1, S_2, \ldots, S_k are MEG-sets of the induced subgraphs $G[C_1 \cup \{v\}], G[C_2 \cup \{v\}], \ldots, G[C_k \cup \{v\}],$ then $S = (S_1 \cup S_2, \ldots, \cup S_k) \setminus \{v\}$ is an MEG-set of G.

Proof. Consider any edge e of G, say in C_1 . Then, there are two vertices x, y of S_1 such that e belongs to all shortest paths between x and y in $G_1 = G[C_1 \cup \{v\}]$. Assume first that $v \notin \{x, y\}$. All shortest paths between x and y in G also exist in G_1 . Thus, e is monitored by $\{x, y\} \subseteq S$ in G. Assume next that $v \in \{x, y\}$: without loss of generality, v = x. At least one edge exists in $G[C_2 \cup \{v\}]$, which implies that $S_2 \setminus \{v\}$ is nonempty, say, it contains z. Then, e is monitored by y and z, since $z \in S$. Thus, S monitors all edges of G, as claimed.

Lemma 2.4. Let G be a graph and H an induced subgraph of G such that for all vertex pairs $\{x, y\}$ in H, no shortest path between them uses edges in G - H. Then, for any set S of vertices of G containing an MEG-set of H, the edges of H are monitored by S in G.

Proof. Let S be an MEG-set of G containing an MEG-set S' of H. Let e be an edge in H that lies on all shortest paths between some pair $\{x, y\}$ of vertices of S'. By our hypothesis, no shortest path between x

and y in G uses any edges of G - H. Thus, the shortest paths between x and y in H are the same as in G, and therefore in G, e is also monitored by $\{x, y\} \subseteq S$.

3 Basic graph classes and bounds

In this section, we study MEG-sets for some standard graph classes.

3.1 Trees

Theorem 3.1. For any tree T with at least one edge, the only optimal MEG-set of T consists of the set of leaves of T.

Proof. The fact that all leaves must be part of any MEG-set follows from Lemma 2.1, as they are simplicial. For the other side, let L be the set of leaves of T. Let e = xy be an edge of T and consider two leaves of T, l_x and l_y , such that l_x is closer to x than to y and that l_y is closer to y than to x. We note that e belongs to the unique (shortest) path between l_x and l_y , thus e is monitored by L. Hence, L is an MEG-set of T.

Corollary 3.2. For any path graph P_n , where $n \ge 2$, we have $meg(P_n) = 2$

This provides a lower bound which is tight for path graphs, which have order n and exactly 2 leaves.

Corollary 3.3. For any tree T of order $n \ge 3$, we have $2 \le meg(T) \le n-1$.

The upper bound is tight for star graphs, which have order n and n-1 leaves.

3.2 Cycle graphs

Theorem 3.4. Given an n-cycle graph C_n , for n = 3 and $n \ge 5$, $\operatorname{meg}(C_n) = 3$. Moreover, $\operatorname{meg}(C_4) = 4$.

Proof. Let us first prove that we need at least three vertices to monitor any cycle. By contradiction, let us assume that two vertices suffice. For any arbitrary vertex pair in the cycle graph, there are two paths joining them, but there is either one single shortest path or two equidistant shortest paths between them. Thus, the edges on at least one of the two paths between the pair will not be monitored by it. Hence, we need at least three vertices in any MEG-set of C_n $(n \ge 3)$.

We now prove the upper bound. Let $n \ge 5$ or n = 3, with the vertices of C_n from v_0 to v_{n-1} . Consider the set $S = \{v_0, v_{\lfloor \frac{n}{2} \rfloor}, v_{\lfloor \frac{2n}{2} \rfloor}\}$. We show that S is an MEG-set of C_n .

Consider every edge of C_n between a vertex pair v_x and v_y in S, then we note that they lie on every (unique) shortest path between these vertices, which has a length at least one for $n \leq 5$ and at least two otherwise, and at most $\lfloor \frac{n}{2} - 1 \rfloor$. Thus, $\operatorname{meg}(C_n) = 3$ when $n \geq 5$ or n = 3.

In the case of C_4 , the above construction does not work. Consider a set of three vertices, say v_0 , v_1 , v_2 without loss of generality due to the symmetries of C_4 . Notice that the edge v_0v_3 is unmonitored by this set. Thus, we have $meg(C_4) = 4$.

3.3 Unicyclic graphs

A unicyclic graph is a connected graph containing exactly one cycle [13]. We now determine the optimal size of an MEG-set of such graphs. Note that in the short version of this paper, published in the proceedings of CALDAM [11], the statement was slightly mistaken. Here, we correct it.

Theorem 3.5. Let G be a unicyclic graph where the only cycle C has length k and whose set of pendant vertices is L(G), |L(G)| = l. Let V_c^+ be the set of vertices of C with degree at least 3. Let p(G) = 1 if $G[V(C)\setminus V_c^+]$ contains a path whose length is at least $\left|\frac{k}{2}\right|$, and p(G) = 0 otherwise. Then, if $k \in \{3, 4\}$,

$$\operatorname{meg}(G) = l + k - |V_c^+|.$$

Otherwise $(k \ge 5)$, then

$$\operatorname{meg}(G) = \begin{cases} 3, \ if \ |V_c^+| = 0; \\ l+2, \ if \ |V_c^+| = 1; \\ l+p(G)+1, \ if \ |V_c^+| = 2, \ k \ is \ even, \ and \ the \ vertices \ in \ V_c^+ \ are \ adjacent \ or \ opposite \ on \ C; \\ l+p(G), \ in \ all \ other \ cases. \end{cases}$$

Proof. Let G be a unicyclic graph where the only cycle C has length k and whose set of pendant vertices is L(G). By Lemma 2.1, all leaves are part of any MEG-set of G. This implies that meg(G) is at least l. If $|V_C^+| = 0$ (i.e. l = 0), we are done by Theorem 3.4, so let us assume $|V_C^+| > 0$ and thus, l > 0.

Similarly as in the proof of Lemma 2.3, for every vertex v of V_C^+ , we know that at least one leaf will exist in the tree component T_v formed if we remove the neighbors of v in C from G. Informally speaking, towards the rest of the graph, this leaf simulates the fact that v is in the solution set.

If $k \in \{3, 4\}$, we consider S = L(G) and we add to S all vertices of C that are of degree 2 in G. One can easily check that this is an MEG-set. Moreover, one can see that adding these degree 2 vertices is necessary by using similar arguments as in the proof of Theorem 3.4 on cycles.

Next, we assume that $k \ge 5$. Let v_0, \ldots, v_{k-1} be the vertices of C.

When $|V_c^+| = 1$, without loss of generality, consider the vertex in V_c^+ to be v_0 . Then, the vertices $\{v_{\lfloor \frac{k}{3} \rfloor}, v_{\lfloor \frac{2k}{3} \rfloor}\}$ on the cycle are sufficient to monitor the graph, in the same way as in Theorem 3.4. Moreover, by the same arguments as in the proof of Theorem 3.4, one can see that if at most one vertex on C is chosen in the MEG-set, some edge of C will not be monitored.

If $|V_c^+| > 1$ and p(G) = 0, the *l* pendant vertices are sufficient to monitor *G*. Indeed, consider an edge *e*. If *e* is not on *C*, let *v* be the vertex of V_C^+ closest to *e*, and let $w \neq v$ be the vertex of V_C^+ closest to *v* (it exists because $|V_c^+| > 1$). Consider a leaf *f* of *G* such that *e* lies on some path from *v* to *f*. Since p(G) = 0, the path from *w* to *f* is a unique shortest path, and thus, *e* is monitored by *f* and some leaf whose closest vertex on *C* is *w*.

If e is an edge of C, e lies on a path between two vertices v, w of V_C^+ . Since p(G) = 0, this path is a shortest path, and e is monitored by two leaves, each of which has v and w as its closest vertex of C, respectively.

Consider the case where p(G) = 1, k is even and $|V_c^+| = 2$ such that the vertices in V_c^+ are adjacent. Then, $G[V(C)\setminus V_c^+]$ consists of a single path P with k-2 vertices, and the edges of P are not monitored by the set of leaves of G. We need to pick at least one vertex of P to monitor the edges of P. If we pick only one vertex of P, then at least one of the edges of P is not monitored (for example, the one that is, in the subgraph of G induced by P, farthest from the picked vertex). Thus, we need to pick at least two additional vertices. Picking the two middle vertices of P, for example, is sufficient and yields an MEG-set of the desired size, l + 2.

Similarly, if p(G) = 1, k is even and $|V_c^+| = 2$ such that the vertices in V_c^+ are opposite on C, there are exist paths of equal length between the two vertices of V_c^+ , and $G[V(C)\setminus V_c^+]$ consists of two paths P_1 and P_2 , each with k/2 - 1 vertices. To monitor the edges of these paths, we need to pick at least one vertex of each of P_1, P_2 . We can in fact pick any vertex of P_1 and any vertex of P_2 , and obtain an MEG-set of size l + 2, as desired.

Finally, consider all other cases where p(G) = 1 and $|V_c^+| > 1$. Now, there is only one path P of $G[V(C)\setminus V_c^+]$ of length at least $\lfloor \frac{k}{2} \rfloor$. As in the previous cases, we can see that the edges of P are not monitored by the set of leaves of G, which implies that $meg(G) \ge l + 1$. To show that $meg(G) \le l + 1$, we select as an MEG-set, the set of leaves together with the middle vertex of P (if P has odd length) or one of the middle vertices of P (if P has even length). One can see that this is an MEG-set by similar arguments as in the previous cases.

3.4 Complete graphs

The following follows immediately from Lemma 2.1, since every vertex of a complete graph is simplicial.

Theorem 3.6. For any $n \ge 2$, we have $meg(K_n) = n$.

3.5 Complete multipartite graphs

The complete k-partite graph K_{p_1,p_2,\ldots,p_k} consists of k disjoint sets of vertices of sizes p_1, p_2, \ldots, p_k , with an edge between any two vertices from distinct sets.

Theorem 3.7. We have $\operatorname{meg}(K_{p_1,p_2,\ldots,p_k}) = |V(K_{p_1,p_2,\ldots,p_k})|$, with the exceptional case of a bipartite graph $K_{1,p}$ with an independent set of size 1 (a star graph), for which $\operatorname{meg}(K_{1,p}) = p$.

Proof. In a complete k-partite graph, all vertices in a given partite set are twins. Therefore, by Lemma 2.2, all vertices of ant partite set of size at least 2 need to be a part of any MEG-set.

If we have several partite sets of size 1, then the vertices from these sets are closed twins, and again by Lemma 2.2 they all belong to any MEG-set.

Thus, we are done, unless there is a unique partite set of size 1, whose vertex we call v. If there are at least three partite sets, then note that v is never part of a unique shortest path, and thus the edges incident with v cannot be monitored if v is not part of the MEG-set.

On the other hand, if the graph is bipartite, it is a star $K_{1,p}$. Here, we know by Theorem 3.1 that meg(G) = p, as claimed.

3.6 Hypercubes

The hypercube of dimension n, denoted by Q_n , is the undirected graph consisting of $k = 2^n$ vertices labelled from 0 to $2^n - 1$ and such that there is an edge between any two vertices if and only if the binary representations of their labels differ by exactly one bit [18]. The Hamming distance H(A, B) between two vertices A, B of a hypercube is the number of bits where the two binary representations of its vertices differ.

We next show that not only C_4 has the whole vertex set as its only MEG-set (Theorem 3.4), but that this also holds for all hypercubes.

Theorem 3.8. For a hypercube graph Q_n with $n \ge 2$, we have $meg(Q_n) = 2^n$.

Proof. Assume by contradiction that there is an MEG-set M of size at most $2^n - 1$. Let $v \in V(G)$ be a vertex that is not in M. It is known that for every vertex pair $\{v_x, v\}$ with $H(v_x, v) \leq n$, there are $H(v_x, v)$ vertex-disjoint paths of length $H(v_x, v)$ between them [18]. Thus, there is no vertex pair in M with a unique shortest path going through the edges incident with v, and M is not an MEG-set, a contradiction.

3.7 Grid graphs

The graph $G \square H$ is the Cartesian product of graphs G and H and with vertex set $V(G \square H) = V(G)$ x V(H), and for which $\{(x, u), (y, v)\}$ is an edge if x = y and $\{u, v\} \in E(H)$ or $\{x, y\} \in E(G)$ and u = v. The grid graph G(m, n) is the Cartesian product $P_m \square P_n$ with vertex set $\{(i, j) \mid 1 \leq i \leq m, 1 \leq j \leq n\}$.

Theorem 3.9. For any $m, n \ge 2$, we have meg(G(m, n)) = 2(m + n - 2).

Proof. We claim that the set $S = \{(i, j) \in V(G(m, n)), i \in \{1, m\} \text{ and } 1 \leq j \leq n \text{ or } j \in \{1, n\} \text{ and } 1 \leq i \leq m\}$ of 2(m + n - 2) vertices of G(m, n) that form the boundary vertices of the grid, form the only optimal MEG-set.

For the necessity side, let us assume that some vertex v = (i, j) of S is not part of the MEG-set. If v is a corner vertex (without loss of generality say v = (1, 1), the two edges incident with v are not monitored, as for any shortest path going through them, there is another one going through vertex (2, 2). If v is not a corner vertex (without loss of generality say v = (1, j) with $2 \le j \le n-1$), then the edge e between v = (1, j) and (2, j) is not monitored, indeed for any shortest path containing e, there is another one avoiding it, either going through vertex (2, j-1) or through (2, j+1).

To see that S is an MEG-set, first see that each boundary edge is monitored by its endpoints. Next, consider an edge e that is not a boundary edge, without loss of generality, e is between (i, j) and (i+1, j). Then, it is monitored by (1, j) and (m, j), whose unique shortest path goes through e.

3.8 Corona products

The corona product $G \odot H$ of two graphs G and H is defined as the graph obtained by taking one copy of G and |V(G)| copies of H and joining the i^{th} vertex of G to every vertex in the i^{th} copy of H.

Let graph G be simple and connected and graph H be simple. Let $\{g_i\}$ and $\{h_i\}$ denote the vertex set of G and H respectively. Let $|V(G)| = n_1$ and $|V(H)| = n_2$.

Theorem 3.10. Given a simple, connected graph G and a simple graph H, we have $meg(G \odot H) = n_1 n_2$.

Proof. By definition of $G \odot H$, every vertex g_i is a cut-vertex in it. Therefore, the edge $e = (g_i, g_j)$ lies in the unique shortest path connecting the i^{th} copy H_i with the j^{th} copy H_j in $G \odot H$. i.e. every edge of G is in the unique shortest path of some H_i and H_j . Hence no vertex of G is in $meg(G \odot H)$.

If some vertex h of H_i is not in the MEG-set, consider an edge of H_i incident with h, if that edge is monitored, then there are two vertices other than h monitoring it; these two vertices must be in H_i and at distance 2 from each other, but then there are two edge-disjoint paths of length 2 joining them (one through h and one through g_i), a contradiction.

4 Relation to feedback edge set number

A feedback edge set of a graph G is a set of edges which when removed from G leaves a forest. The smallest size of such a feedback edge set of G is denoted by fes(G) and is sometimes called the *cyclomatic* number of G.

We next introduce the following terminology from [7]. A vertex is a *core vertex* if it has degree at least 3. A path with all internal vertices of degree 2 and whose end-vertices are core vertices is called a *core path*. Do note that we allow the two end-vertices to be equal, but that every other vertex must be distinct. A core path that is a cycle (that is, both end-vertices are equal) is a *core cycle*. For the sake of distinction, a core path that is not a core cycle is called a *proper core path*. We say that a (non-empty) path from a core vertex u to a leaf v is a *leg* of u if all internal vertices of the path have degree 2 (u is not considered to be a part of the leg). The *base graph* of a graph G is the graph of minimum degree 2 obtained from G by iteratively removing vertices of degree 1. A *hanging tree* is a connected subtree of G which is the union of some legs removed from G during the process of creating the base graph G_b of G. Thus, G can be decomposed into its base graph and a set of maximal hanging trees. The root of such a maximal hanging tree T is the vertex common to T and G.

See Figure 2 for a graph whose core vertices are in red. It has two hanging trees, three core cycles, three proper core paths of length 4, and five proper core paths of length 1.



Figure 2: Example of a graph G with its core vertices in red.

Based on the aforementioned, we have the following lemma.

Lemma 4.1 ([7, 15]). Let G be a graph with $fes(G) = k \ge 2$. The base graph of G has at most 2k - 2 core vertices, that are joined by at most 3k - 3 edge-disjoint core paths. Equivalently, G can be obtained from a multigraph H of order at most 2k - 2 and size at most 3k - 3 by subdividing its edges an arbitrary number of times and iteratively adding degree 1 vertices.

Lemma 4.2. Let S be an MEG-set of the base graph G_b of G and L(G) be the set of leaves in G. Then, $S \cup L(G)$ is an MEG-set of G.

Proof. Let G_b be a base graph of G. Consider all vertices that are roots of maximal hanging trees on G_b . By Theorem 3.1, the optimal MEG-set of each tree consists of all leaves. We repeatedly apply Lemma 2.3 to G, where for each application of Lemma 2.3, the cut-vertex is the root of a hanging tree in consideration.

Lemma 2.1, Theorem 3.1 and Lemma 4.2 together imply that if fes(G) = 0, then $meg(G) \leq fes(G) + |L(G)|$. Moreover, if fes(G) = 1, then $meg(G) \leq fes(G) + |L(G)| + 3$, where |L(G)| is the number of leaves of G. We next give a similar bound when $fes(G) \geq 2$.



Figure 3: Example of a graph G and its base graph G_b with four core cycles.

Theorem 4.3. If $fes(G) \ge 2$, then $meg(G) \le 9 fes(G) + |L(G)| - 8$ where |L(G)| is the number of leaves of G.

Proof. Let k = fes(G). We show how to construct a MEG-set M of G_b of order at most 9k - 8 and, by applying Lemma 4.2 to G, of order 9k - 8 + |L(G)| for G. If an edge e is part of a maximal hanging tree, then by Lemma 2.1 and Lemma 2.3, it is monitored by the leaves of G on the maximal hanging tree. M is constructed as follows.

- We let all core vertices of G_b be part of M.
- One or two internal vertices from each proper core path belongs to M, only if the length is at least 2, as explained below.
- Two or three internal vertices from each core cycle, as explained below.

Consider a proper core path P, with core vertex endpoints c and c', and the median vertex x_1 in the case of an odd-length path and x_1 , x_2 in the case of an even-length path, with d edges (on P) between the endpoints and the respective medians in P. Then, we choose the single median vertex x_1 or the two median vertices x_1 , x_2 from each of the core paths into M.

For each core cycle, in addition to the core vertex of that cycle, we add three vertices that are as equidistant as possible on the cycle, to be part of M (as in Theorem 3.4).

Let e by any edge of G. We now show that our construction M monitors any such edge in G_b . If e lies on a core cycle, assume an origin core vertex of v_0 . Then, based on Lemma 2.3 and Theorem 3.4, we deduce that in the worst case, four vertices together suffice to monitor the edges.

If the edge e lies on a proper core path P, then we have the following cases. Let c and c' be the core vertex endpoints of P, and the median vertex x_1 in the case of an odd-length path and x_1 , x_2 in the case of an even-length path and d edges of P between the end points and the respective medians in P. Without loss of generality, let us say that e lies on the path P such that its closest core vertex is c and closest median x_1 in the event of an even-length path. Suppose first that d is even. Given that the distance between c and x_1 is d in P, the length of any other path between them must be at least d + 2. Therefore, c and x_1 monitor e. We can similarly argue that if the closest core vertex to e was c' and the closest median vertex was x_2 , then c' and x_2 monitor e. If e lay in between the median vertices x_1 and x_2 , then we know that those vertices would monitor e because they are adjacent. If the path was of odd length, then depending on which of the core vertices c and c' was closest to e, the distance between the median vertex x_1 would monitor the edge apart from the core vertices. This justifies our construction of M for G_b .

By Lemma 4.1, the number of core vertices of G_b is at most 2k-2, and there are at most 3k-3 core paths.

If we have core cycles in our graph, then we must note that there can be at most k such cycles in the graph. Indeed, if there were k + 1 core cycles in the graph, since they are all edge-disjoint, we need at least k + 1 edges to be removed from G to obtain a forest, a contradiction to the fact that fes(G) = k.

Let n_c be the number of core cycles and n_p be the number of proper core paths. We have $|M| = 3n_c + 2n_p + 2k - 2$. Since $n_c \leq k$ and $n_c + n_p \leq 3k - 3$ by Lemma 4.1, we get $|M| \leq 3k + 2(2k - 3) + 2k - 2 = 9k - 8$.

Recall that the max leaf number of G, denoted mln(G), is the maximum number of leaves in a spanning tree of G. It can be seen as a refinement of the feedback edge set number of G [6]. We get the following corollary.

Corollary 4.4. For any graph G, we have $meg(G) \leq 10 mln(G)$, where mln(G) is the max leaf number of G.

Proof. It is known that $fes(G) \leq mln(G)$ [6], and clearly, $|L(G)| \leq mln(G)$, thus the bound follows from Theorem 4.3.

Proposition 4.5. For any integer $k \ge 2$, there exists a graph G with fes(G) = k and meg(G) = 3k + |L(G)|.

Proof. Consider G and its base graph G_b in Figure 3. We know that the leaves must be part of any MEG-set by Lemma 2.1. The MEG-set for G_b consists of all the vertices in each of the core cycles (each a C_4) in G_b , except the common core vertex. It is easy to check that no smaller set can work. The size of the optimal MEG-set in this example is 3k + |L(G)| and therefore, this is an instance where this proposition holds.

5 NP-completeness for graphs of small maximum degree

The MONITORING EDGE GEODETIC SET decision problem is defined as follows.

MONITORING EDGE GEODETIC SET Instance: A graph G = (V(G), E(G)) and an integer k. Question: Is there an *MEG-set* $S \subseteq V(G)$ of G of size at most k?

In this section, we show that the problem is NP-hard by a reduction from VERTEX COVER.

VERTEX COVER Instance: A graph G = (V(G), E(G)) and an integer k. Question: Is there a vertex cover $C \subseteq V(G)$ of size at most k such that every edge in E(G) is incident with some vertex in C?

Overview of the reduction. Consider a cubic planar graph. We subdivide each edge of this graph four times. Let this graph be G with vertex set V(G) and edge set E(G). Also, let |V(G)| = n. Observe that the vertices of G have degree 2 or 3 and its girth is at least 15. It is well know that VERTEX COVER is NP-hard when the input graph is cubic, see for example [17]. We show that it is also NP-hard when the input graph is restricted to resemble graphs such as G. (Note that such a result seems to be already known for INDEPENDENT SET [1], which has the same complexity as VERTEX COVER. We state the proof for VERTEX COVER for completeness, as [1] is in Russian.)

Then, given G, we construct a new graph H_G of maximum degree 8 and show that G has a vertex cover of size k if and only if H_G has an MEG-set of size 3n + k. This completes our proof.

Lemma 5.1 ([1]). VERTEX COVER is NP-complete, even for input graphs obtained from a cubic graph by subdividing every edge four times.

Proof. Let G' be the original (cubic planar) graph before subdivision and let |E(G')| = m'. To show that finding the minimum vertex cover in G is NP-hard it suffices to show that G' has a vertex cover of size k if and only if G has a vertex cover of size 2m' + k.

(The if part.) Let C be a vertex cover of G of size 2m' + k. We contruct a vertex cover C' of G' of size at most k as follows. Consider all maximal paths of G of length 5 whose end points are degree 3 vertices and whose internal points are degree 2 vertices. Let $\pi_{uv}, u, v \in V(G')$, represent one such path. Observe that π_{uv} which is in G, represents the edge uv in G'. Since C is a vertex cover, all edges in π_{uv} are covered by some vertex in C. We add to C' the vertex u and/or v if it was included in C. If both u and v were not in C then we arbitrarily choose one of these two vertices and add it to C'.

Since for each edge of G' we have included one of its end points, C' is a vertex cover of G. Also, π_{uv} needs at least 3 vertices to cover all its edges. Therefore, if one of u or v is inserted into C', two other vertices of π_{uv} , which are in C, are excluded. Again, if u and v both belong to C then π_{uv} needs at least 4 vertices to cover all its edges. Therefore, if both u and v are inserted into C', two other vertices of π_{uv} , which are in C, are excluded. See Figure 4 for reference. Since for every such maximal path we excluded at least 2 vertices from C from being inserted into C', the size of C' is at most k.



Figure 4: Choice of C when $u \in C'$ and $u, v \in C'$.

(The only if part.) Let C' be a vertex cover of G' of size k. We construct a vertex cover of G of size at most 2m' + k as follows. As before, consider all maximal paths of G and let $\pi_{uv}, u, v \in V(G')$, be one such path. Either u or v need to be in C'. We insert u or v or both into C if they belong to C' and

are not already inserted. We select two additional vertices of π_{uv} such that all its edges are covered and insert them into C'. As evident from Figure 4 this is always possible. C is a vertex cover of G. Since for each maximal path π_{uv} we insert two additional vertices into C, the size of C is at most 2m' + k. \Box

Now we show how to construct H_G from G.

Construction. For each degree 3 vertex $u \in V(G)$, we construct a vertex gadget H_G^u , with 16 vertices and 18 edges as shown in Figure 5(a). Each vertex $f_i^u, i \in \{1, 2, 3\}$, is associated with a unique side of the triangle $\triangle c_1^u c_2^u c_3^u$ as depicted in the figure. For example, f_1^u is associated with the side $\overline{c_1^u d_1^u c_2^u}$. Each f_i^u also has a unique associate vertex c_i^u . Note that c_i^u belongs to the side of $\triangle c_1^u c_2^u c_3^u$ associated with f_i^u and its distance from a^u is 2. For a degree 2 vertex $v \in V(G)$, we construct a vertex gadget H_G^v , with 14 vertices and 15 edges as shown in Figure 5(b). Again, each $f_i^v, i \in \{1, 2\}$, vertex is associated with a unique side of the triangle $\triangle c_1^v c_2^v c_3^v$ depicted in the figure. In this case however, the assosiate vertex of f_1^v is c_1^v and that of f_2^v is c_3^v . Again note that the associate vertex of f_i^v belongs to the side of $\triangle c_1^v c_2^v c_3^v$ associated with f_i^u and its distance from a^v is 2. Also, observe that the missing elements in H_G^v , corresponding to H_G^u , are the vertices b_2^v and f_3^v and their incident edges.



(a) Vertex gadget for a degree 3 vertex.

(b) Vertex gadget for a degree 2 vertex.

Figure 5: The vertex gadgets.

Let u be a degree 3 vertex in G and let $v \in V(G)$ be such that $uv \in E(G)$. According to our construction of G, v is always of degree 2. We connect the vertices of H_G^u and H_G^v using two vertices (g_1^{uv}) and $g_2^{uv})$ and five edges as shown in Figure 6. Observe that H_G^u and H_G^v are connected to each other via vertices f_1^u and f_2^v , thus g_1^{uv} and g_2^{uv} are connected to f_1^u 's and f_2^v 's associated sides. Since u is of degree 3 in G, there are also similar connections between H_G^u and two additional vertex gadgets which are shown in the figure. One of them is labelled H_G^t , the other is not labelled.

For a degree 2 vertex v of G, the connections are similar with the only exception being that, instead of three, the vertex gadget is connected to only two other vertex gadgets. This means that c_1^v and c_3^v are both adjacent to one vertex of type $g_i^{xv}, i \in \{1, 2\}, x \in V(G)$, while c_2^v , similar to the c_j^u 's (u is of degree 3 in G), is adjacent to two such vertices.

Next, consider any two vertices $u, w \in V(G)$ such that the distance between them in G is 2. Let us again assume u to be of degree 3 in G. By the structure of G, this implies w to be of degree 2. Let $v \in V(G)$ be the unique vertex adjacent to both u and w. Let $f_{i'}^u$ (resp. $f_{j'}^w$) be the vertex in H_G^u (resp. H_G^w) which is connected to H_G^v . Let $c_{i'}^u$ be the associate vertex of $f_{i'}^u$ and c_h^w the associate vertex of $f_{j'}^w$.

We connect $c_{i'}^u$ and c_h^w . This ensures that the length of the (unique) shortest path between a^u and a^w in H_G is 5 and it passes through the edge $c_{i'}^u c_h^w$.

Now consider two vertices $t, w \in V(G)$ such that the distance between them in G is 3. Let $u, v \in V(G)$ be the two vertices included in the unique shortest path between t and w in G. Also let u be adjacent to t and v to w in G. Let $f_{i'}^t$ (resp. $f_{j'}^w$) be the vertex in H_G^t (resp. H_G^w) which is connected to H_G^u (resp. H_G^v). Let c_g^t and c_h^w be the associate vertices of $f_{i'}^t$ and $f_{j'}^w$, respectively. We connect c_g^t and c_h^w , such that the length of the (unique) shortest path between a^t and a^w is 5 and it passes through the edge $c_g^t c_h^w$. See Figure 6 for reference.



Figure 6: Connections between vertex gadgets. Here, vertex u was of degree 3 in G, while the other vertices are of degree 2 in G.

The subgraphs H_G^u , $\forall u \in V(G)$, along with the connections between them constitute the graph H_G . Recall that the maximum degree of a vertex in G is 3 and any two degree 3 vertices of G are at least distance 5 apart. Therefore, for any vertex $u \in V(G)$, the maximum number of vertices at distance either exactly 2 or 3 in G is 3. If u is of degree 3 in G then any vertex c_i^u , $i \in \{1, 2, 3\}$, is connected to four vertices from $V(H_G^u)$, two vertices of type g_j^{ux} , $j \in \{1, 2\}$, $x \in V(G)$, and one vertex each of type c_h^y , where $h \in \{1, 3\}$ and $y \in V(G)$ is either at a distance 2 or 3 from u. This makes the total degree of c_i^u to be 8. If v is of degree 2 in G then a vertex c_i^v , $i \in \{1, 2, 3\}$, is connected to at most 4 vertices from $V(H_G^v)$, at most two vertices of type g_j^{vx} , $j \in \{1, 2\}$, $x \in V(G)$, and at most 2 vertex each of type $c_{h'}^y$, where $h' \in \{1, 2, 3\}$ and $y' \in V(G)$ is either at a distance 2 or 3 from v. This makes the total degree of c_i^v to be at most 10. A close inspection of H_G reveals that this upper bound is actually 9 and the maximum degree of H_G is determined by some c_i^v and hence is also 9. We prove the following result.

Lemma 5.2. G has a vertex cover of size k if and only if H_G has a MEG-set of size 3n + k.

Proof. (The if part.) Let $S \subseteq V(H_G)$ be an MEG-set of H_G of size 3n + k and let $S' \subset S$ be the set which remains after removing all degree 1 vertices from S. Since H_G has 3n degree 1 vertices, all of which have to be included in every MEG-set, the size of S' is k. Let $C \subseteq V(G)$ be the set of all vertices $v \in V(G)$ for which $V(H_G^v)$ has at least one element in S. It is easy to see that the size of C is at most k. We show that it is also a vertex cover of G.

Let us assume for the sake of contradiction that C is not a vertex cover of G. Then, there exists an edge $uv \in E(G), u, v \in V(G)$, such that neither u nor v belong to C. Let us assume u to be a degree 3 vertex of G and let H_G^u and H_G^v be connected as shown in Figure 6. The other case, when u is of degree 2, is similar. Now, consider the edge $f_1^u f_2^v \in E(H_G)$. Since S is still an MEG-set, there exists a pair of vertices $x, y \in S$ which monitors the edge $f_1^u f_2^v$ i.e. every shortest path from x to y passes through $f_1^u f_2^v$. Let π_{xy} be one such path. Without loss of generality we assume that f_1^u is encountered before f_2^v while traversing π_{xy} from x to y.

Let $V'(H_G^v) = V(H_G^v) \setminus \{e_1^v, e_2^v, e_3^v\}$. Let x' (resp. y') be the first vertex from $V'(H_G^u)$ (resp. $V'(H_G^v)$) encountered when traversing π_{xy} starting at x (resp. y). Since both u and v do not belong to C, from our contruction of C, no element of S belongs to $V'(H_G^u) \cup V'(H_G^v)$. This implies that neither x nor y belong to $V'(H_G^u) \cup V'(H_G^v)$. Since x does not belong to $V'(H_G^u)$, x' is either $c_i^u, i \in \{1, 2, 3\}$, or $f_i^u, i' \in \{2, 3\}$, and similarly since y does not belong to $V'(H_G^v)$, y' is either $c_j^v, j \in \{1, 2, 3\}$, or f_1^v . We have the following cases.

Case 1 $(x' \text{ is } c_i^u \text{ and } y' \text{ is } c_j^v)$. Note that in this case $i, j \in \{1, 2, 3\}$. It is easy to see that any shortest path between c_i^u and c_j^v passes through either the vertex g_1^{uv} or g_2^{uv} and does not include the edge $f_1^v f_2^v$. This means that x and y cannot monitor the edge $f_1^u f_2^v$, which implies a contradiction.

Case 2 $(x' \text{ is } c_i^u \text{ and } y' \text{ is } f_1^v)$. Note that in this case $i \in \{1, 2, 3\}$. Let $f_{j'}^w, j' \in \{1, 2\}$ and $w \in V(G)$, be the vertex encountered just before f_1^v when traversing π_{xy} starting at y. Since the distance between u and w in G is 2, by construction there exists an edge $c_i^u c_h^w, i' = 1$ and $h \in \{1, 3\}$, in G. Therefore, the length of the shortest path between c_i^u and $f_{j'}^w$ is at most 6 and passes through the edge $c_i^u c_h^w$ and not $f_1^u f_2^v$. This means that x and y cannot monitor the edge $f_1^u f_2^v$, which is a contradiction. The other case when x' is some $f_q^u, g \in \{2, 3\}$, and y' some $c_h^v, h \in \{1, 2, 3\}$, can be similarly disproved.

Case 3 $(x' \text{ is } f_i^u \text{ and } y' \text{ is } f_1^v)$. Note that in this case $i \in \{2,3\}$. Let $f_{i'}^t, i' \in \{1,2\}$ and $t \in V(G)$, be the vertex encountered just before f_i^u when traversing π_{xy} starting at x. Similarly, let $f_{j'}^w, j' \in \{1,2\}$ and $w \in V(G)$, be the vertex encountered just before f_1^v when traversing π_{xy} starting at y. Since u is of degree 3, both t and w are degree 2 vertices of G. Since the distance between t and w in G is 3, by construction there exists an edge $c_g^t c_h^w, g, h \in \{1,3\}$, in H_G . Therefore, there are at least two shortest path (of length 7) between $f_{i'}^t$ and $f_{j'}^w$, one which includes the edge $f_1^u f_2^v$ and the other which includes the edge $c_{g'}^t c_h^w$. This means that x and y cannot monitor the edge $f_1^u f_2^v$, a contradiction.

(The only if part.) Let $C \subseteq V(G)$ be a vertex cover of G of size k. We construct the set $S \subset V(H_G)$ as follows. For each vertex $v \in V(G)$, insert into S the vertices e_1^v, e_2^v and e_3^v of H_G . If $v \in C$ then we also add the vertex $a^v \in V(H_G^v)$ to S. Observe that the size of S is 3n + k. We show that it is also an MEG-set of H_G .

For every vertex $u \in V(G)$, the vertices $e_1^u, e_2^u, e_3^u \in S$ monitor all the edges in the cycle $c_1^u, d_1^u, c_2^u, d_2^u, c_3^u, d_3^u$ of H_G^u , as well as the edges $c_i^v e_i^v, \forall i \in \{1, 2, 3\}$. This is because for each pair of vertices from e_1^v, e_2^v and e_3^v , the shortest path between them is unique and hence all edges on the shortest path are monitored. For example, for the vertex pair e_1^v and e_2^v , the unique shortest path is e_1^v, c_2^v, e_2^v and all its edges are monitored by the vertex pair. Here we assume u to be a degree 3 vertex of G; the other case, when u is degree 2, is similar.

Next, with respect to an edge $uv \in E(G)$, let the subgraphs H_G^u and H_G^v be connected as shown in Figure 6. Again, the shortest path between e_1^u and e_3^v (which passes through the vertex g_1^{uv}), and between e_2^u and e_2^v (which passes through the vertex g_2^{uv}) are unique. Therefore, all edges of H_G incident on g_1^{uv} and g_2^{uv} are monitored.

If $u \in V(G)$ is part of the vertex cover C, then $a^u \in S$ and a^u, b^u_i, c^u_i, e^u_i is the unique shortest path between a^u and $e^u_i, \forall i \in \{1, 2, 3\}$. Therefore, all edges on these paths are monitored.

If $u \in V(G)$ is not part of the vertex cover C, then all three neighbors of u in G belong to C. Let

 $v \in V(G)$ be one such neighbor. Then $a^v \in V(H_G^v)$ is part of S. Observe that the only shortest path between e_3^u and a^v is the path $e_3^u, c_3^u, b_3^u, a^u, f_1^u, f_2^v, a^v$ of length 6. Therefore, all edges on this path are monitored. Any other path between e_3^u and a^v passes through either c_1^u or c_2^u and is of length at least 7. This is ensured in the construction phase by restricting c_3^u from being directly connected to any $c_{i'}^w$, $i' \in \{1, 3\}$. Similarly, we can show that edges $a^u b_i^u$ and $b_i^u c_i^u, \forall i \in \{1, 2\}$, are also monitored.

Now consider all edges $c_i^t c_j^w$, $i, j \in \{1, 2, 3\}$ and $t, w \in V(G)$, of H_G such that t and w are at distance 2 or 3 in G. The shortest path between e_i^t and e_j^w is of length 3, is unique, and passes through the edge $c_i^t c_j^w$, thereby monitoring it. Since we have shown all edges of H_G to be monitored by some vertex pair in S, therefore S is an MEG-set of H_G .

Since MONITORING EDGE GEODETIC SET is clearly in NP [14] and H_G has maximum degree 9, Lemma 5.1 and Lemma 5.2 imply the following theorem.

Theorem 5.3. MONITORING EDGE GEODETIC SET is NP-complete, even for graphs of maximum degree 9.

6 Conclusion

Inspired by a network monitoring application, we have defined the new concept of MEG-sets of a graph, which is a common refinement of the popular concept of a geodetic set and its variants, and of the previously studied distance-edge-monitoring sets.

We have studied the concept on basic graph classes. It is interesting to note that there are many graph classes which require the entire vertex set in any MEG-set: complete graphs, complete multipartite graphs, and hypercubes. More examples are provided in [14]. It would be an interesting question to characterize all such graphs, if they can be described in a meaningful way.

Our upper bound using the feedback edge set number is probably not tight. What is a tight bound on this regard?

Finally, it remains to investigate further computational aspects of the problem. Clearly, MONITORING EDGE GEODETIC SET is polynomial-time solvable on the graph classes studied in Section 3, such as trees, unicyclic graphs, etc. What about graphs of bounded tree-width? Also, VERTEX COVER remains NP-hard on planar cubic graphs [17]. However, our NP-hardness reduction does not preserve planarity. Is MONITORING EDGE GEODETIC SET NP-complete for planar graphs? For subcubic graphs? What about other standard graph classes like interval graphs? Also, the approximation complexity and the parameterized complexity of the problem could be investigated.

Acknowledgements

Florent Foucaud thanks Ralf Klasing and Tomasz Radzik for initial discussions in Bordeaux in 2019, which inspired the present study.

References

- V. E. Alekseev. The effect of local constraints on the complexity of determination of the graph independence number. *Combinatorial-algebraic methods in applied mathematics*, 3-13, Gorkiy University Press, Gorky, 1982 (in Russian).
- [2] E. Bampas, D. Bilò, G. Drovandi, L. Gualà, R. Klasing and G. Proietti. Network verification via routing table queries. *Journal of Computer and System Sciences* 81(1):234–248, 2015.
- [3] Z. Beerliova, F. Eberhard, T. Erlebach, A. Hall, M. Hoffmann, M. Mihalák and L. S. Ram. Network discovery and verification. *IEEE Journal on Selected Areas in Communications* 24(12):2168–2181, 2006.

- [4] Y. Bejerano and R. Rastogi. Robust monitoring of link delays and faults in IP networks. *IEEE/ACM Transactions on Networking* 14(5):1092–1103, 2006.
- [5] H. S. M. Coxeter. Self-dual configurations and regular graphs. Bulletin of the American Mathematical Society 56(5): 413–455, 1950.
- [6] D. Eppstein. Metric dimension parameterized by max leaf number. Journal of Graph Algorithms & Applications 19(1):313-323, 2015.
- [7] L. Epstein, A. Levin and G. J. Woeginger. The (weighted) metric dimension of graphs: hard and easy cases. *Algorithmica* 72(4), 1130-1171, 2015.
- [8] M. R. Fellows, D. Lokshtanov, N. Misra, M. Mnich, F. Rosamond, and S. Saurabh. The complexity ecology of parameters: an illustration using bounded max leaf number. *Theory of Computing Systems* 45(4):822–848, 2009.
- [9] F. Foucaud, R. Klasing, M. Miller and J. Ryan. Monitoring the edges of a graph using distances. Proceedings of the 6th International Conference on Algorithms and Discrete Applied Mathematics (CALDAM 2020), Lecture Notes in Computer Science 12016:28-40, 2020.
- [10] F. Foucaud, S. Kao, R. Klasing, M. Miller and J. Ryan. Monitoring the edges of a graph using distances. Discrete Applied Mathematics 319:424–438, 2022.
- [11] F. Foucaud, K. Narayanan and L. Ramasubramony Sulochana. Monitoring edge-geodetic sets in graphs. Proceedings of the 9th International Conference on Algorithms and Discrete Applied Mathematics (CALDAM 2023), *Lecture Notes in Computer Science* 13947:245-256, 2023.
- [12] F. Harary, E. Loukakis, and C. Tsouros. The geodetic number of a graph. Mathematical and Computational Modelling 17:89–95, 1993.
- [13] F. Harary. *Graph Theory*. Reading, MA: Addison-Wesley, 1994.
- [14] J. Haslegrave. Monitoring edge-geodetic sets: hardness and graph products. Discrete Applied Mathematics 340:79–84, 2023.
- [15] L. Kellerhals and T. Koana. Parameterized complexity of geodetic set. Proceedings of the 15th International Symposium on Parameterized and Exact Computation (IPEC 2020), *Leibniz International Proceedings in Informatics (LIPIcs)* 180, 20:1–20:14, 2020.
- [16] P. Manuel, S. Klavžar, A. Xavier, A. Arokiaraj and E. Thomas. Strong edge geodetic problem in networks. Open Mathematics 15(1):1225–1235, 2017.
- [17] B. Mohar. Face covers and the genus problem for apex graphs. Journal of Combinatorial Theory, Series B, 82(1), 102-117, 2001.
- [18] Y. Saad and M. Schultz. Topological properties of hypercubes. *IEEE Transactions on Computers*, 37(7), 867–872, 1988.
- [19] A. P Santhakumaran, and J. John. Edge geodetic number of a graph. Journal of Discrete Mathematical Sciences and Cryptography 10:415–432, 2007.
- [20] S. Skiena. Implementing Discrete Mathematics: Combinatorics and Graph Theory with Mathematica. Reading, MA: Addison-Wesley, 1990.