Sensitivity versus Certificate Complexity of Boolean Functions *

Andris Ambainis, Krišjānis Prūsis, and Jevgēnijs Vihrovs

Faculty of Computing, University of Latvia, Raina bulv. 19, Rīga, LV-1586, Latvia

Abstract Sensitivity, block sensitivity and certificate complexity are basic complexity measures of Boolean functions. The famous sensitivity conjecture claims that sensitivity is polynomially related to block sensitivity. However, it has been notoriously hard to obtain even exponential bounds. Since block sensitivity is known to be polynomially related to certificate complexity, an equivalent of proving this conjecture would be showing that the certificate complexity is polynomially related to sensitivity. Previously, it has been shown that $bs(f) \leq C(f) \leq 2^{s(f)-1}s(f) - (s(f)-1)$. In this work, we give a better upper bound of $bs(f) \leq C(f) \leq \max\left(2^{s(f)-1}\left(s(f)-\frac{1}{3}\right),s(f)\right)$ using a recent theorem limiting the structure of function graphs. We also examine relations between these measures for functions with 1-sensitivity $s_1(f)=2$ and arbitrary 0-sensitivity $s_0(f)$.

1 Introduction

Sensitivity and block sensitivity are two well-known combinatorial complexity measures of Boolean functions. The sensitivity of a Boolean function, s(f), is just the maximum number of variables x_i in an input assignment $x = (x_1, \ldots, x_n)$ with the property that changing x_i changes the value of f. Block sensitivity, bs(f), is a generalization of sensitivity to the case when we are allowed to change disjoint blocks of variables.

Sensitivity and block sensitivity are related to the complexity of computing f in several different computational models, from parallel random access machines or PRAMs [6] to decision tree complexity, where block sensitivity has been useful for showing the complexities of deterministic, probabilistic and quantum decision trees are all polynomially related [9,4,5].

A very well-known open problem is the sensitivity vs. block sensitivity conjecture which claims that the two quantities are polynomially related. This problem is very simple to formulate (so simple that it can be assigned as an undergraduate research project). At the same time, the conjecture appears quite difficult to solve. It has been known for over 25 years and the best upper and lower bounds

 $^{^\}star$ The research leading to these results has received funding from the European Union Seventh Framework Programme (FP7/2007-2013) under projects RAQUEL (Grant Agreement No. 323970) and ERC Advanced Grant MQC.

are still very far apart. We know that block sensitivity can be quadratically larger than sensitivity [10,12,2] but the best upper bounds on block sensitivity in terms of sensitivity are still exponential (of the form $bs(f) \leq c^{s(f)}$) [11,8,1].

Block sensitivity is polynomially related to a number of other complexity measures of Boolean functions: $certificate\ complexity$, $polynomial\ degree$ and the number of queries to compute f either deterministically, probabilistically or quantumly [5]. This gives a number of equivalent formulations for the sensitivity vs. block sensitivity conjecture: it is equivalent to asking whether sensitivity is polynomially related to any one of these complexity measures.

Which of those equivalent forms of the conjecture is the most promising one? We think that certificate complexity, C(f), is the combinatorially simplest among all of these complexity measures. Certificate complexity being at least c simply means that there is an input $x=(x_1,\ldots,x_n)$ that is not contained in an (n-(c-1))-dimensional subcube of the Boolean hypercube on which f is constant. Therefore, we now focus on the "sensitivity vs. certificate complexity" form of the conjecture.

1.1 Prior Work

The best upper bound on certificate complexity in terms of sensitivity is

$$C_0(f) \le 2^{s_1(f)-1} s_0(f) - (s_1(f)-1)$$
 (1)

due to Ambainis et al. $[1]^1$ The bounds for $C_0(f)$ also hold for $C_1(f)$ symmetrically (in this case, $C_1(f) \leq 2^{s_0(f)-1}s_1(f) - (s_0(f)-1)$), so it is sufficient to focus on the former.

Since $bs(f) \leq C(f)$, this immediately implies that

$$bs(f) \le C(f) \le 2^{s(f)-1}s(f) - (s(f)-1). \tag{2}$$

1.2 Our Results

In this work, we give improved upper bounds for the "sensitivity vs. certificate complexity" problem. Our main technical result is

Theorem 1. Let f be a Boolean function which is not constant. If $s_1(f) = 1$, then $C_0(f) = s_0(f)$. If $s_1(f) > 1$, then

$$C_0(f) \le 2^{s_1(f)-1} \left(s_0(f) - \frac{1}{3} \right).$$
 (3)

A similar bound for $C_1(f)$ follows by symmetry. This implies a new upper bound on block sensitivity and certificate complexity in terms of sensitivity:

Here, C_0 (C_1) and s_0 (s_1) stand for certificate complexity and sensitivity, restricted to inputs x with f(x) = 0 (f(x) = 1).

Corollary 1. Let f be a Boolean function. Then

$$bs(f) \le C(f) \le \max\left(2^{s(f)-1}\left(s(f) - \frac{1}{3}\right), s(f)\right). \tag{4}$$

On the other hand, the function of Ambainis and Sun [2] gives the separation of

$$C_0(f) = \left(\frac{2}{3} + o(1)\right) s_0(f) s_1(f) \tag{5}$$

for arbitrary values of $s_0(f)$ and $s_1(f)$. For $s_1(f) = 2$, we show an example of f that achieves

$$C_0(f) = \left| \frac{3}{2} s_0(f) \right| = \left| \frac{3}{4} s_0(f) s_1(f) \right|.$$
 (6)

We also study the relation between $C_0(f)$ and $s_0(f)$ for functions with low $s_1(f)$, as we think these cases may provide insights into the more general case.

If $s_1(f) = 1$, then $C_0(f) = s_0(f)$ follows from (1). So, the easiest non-trivial case is $s_1(f) = 2$, for which (1) becomes $C_0(f) \le 2s_0(f) - 1$.

For $s_1(f) = 2$, we prove a slightly better upper bound of $C_0(f) \leq \frac{9}{5}s_0(f)$. We also show that $C_0(f) \leq \frac{3}{2}s_0(f)$ for $s_1(f) = 2$ and $s_0(f) \leq 6$ and thus our example (6) is optimal in this case. We conjecture that $C_0(f) \leq \frac{3}{2}s_0(f)$ is a tight upper bound for $s_1(f) = 2$.

Our results rely on a recent "gap theorem" by Ambainis and Vihrovs [3] which says that any sensitivity-s induced subgraph G of the Boolean hypercube must be either of size 2^{n-s} or of size at least $\frac{3}{2}2^{n-s}$ and, in the first case, G can only be a subcube obtained by fixing s variables. Using this theorem allows refining earlier results which used Simon's lemma [11] – any sensitivity-s induced subgraph G must be of size at least 2^{n-s} – but did not use any more detailed information about the structure of such G.

We think that further research in this direction may uncover more interesting facts about the structure of low-sensitivity subsets of the Boolean hypercube, with implications for the "sensitivity vs. certificate complexity" conjecture.

2 Preliminaries

Let $f: \{0,1\}^n \to \{0,1\}$ be a Boolean function on n variables. The i-th variable of input x is denoted by x_i . For an index set $P \subseteq [n]$, let x^P be the input obtained from an input x by flipping every bit x_i , $i \in P$.

We briefly define the notions of sensitivity, block sensitivity and certificate complexity. For more information on them and their relations to other complexity measures (such as deterministic, probabilistic and quantum decision tree complexities), we refer the reader to the surveys by Buhrman and de Wolf [5] and Hatami et al. [7].

Definition 1. The sensitivity complexity s(f,x) of f on an input x is defined as $\left|\left\{i \mid f(x) \neq f\left(x^{\{i\}}\right)\right\}\right|$. The b-sensitivity $s_b(f)$ of f, where $b \in \{0,1\}$, is defined as $\max(s(f,x) \mid x \in \{0,1\}^n, f(x) = b)$. The sensitivity s(f) of f is defined as $\max(s_0(f), s_1(f))$.

Definition 2. The block sensitivity bs(f, x) of f on input x is defined as the maximum number t such that there are t pairwise disjoint subsets B_1, \ldots, B_t of [n] for which $f(x) \neq f(x^{B_i})$. We call each B_i a block. The b-block sensitivity $bs_b(f)$ of f, where $b \in \{0, 1\}$, is defined as $\max(bs(f, x) \mid x \in \{0, 1\}^n, f(x) = b)$. The block sensitivity bs(f) of f is defined as $\max(bs_0(f), bs_1(f))$.

Definition 3. A certificate c of f on input x is defined as a partial assignment $c: P \to \{0,1\}, P \subseteq [n]$ of x such that f is constant on this restriction. We call |P| the length of c. If f is always 0 on this restriction, the certificate is a 0-certificate. If f is always 1, the certificate is a 1-certificate.

Definition 4. The certificate complexity C(f,x) of f on input x is defined as the minimum length of a certificate that x satisfies. The b-certificate complexity $C_b(f)$ of f, where $b \in \{0,1\}$, is defined as $\max(C(f,x) \mid x \in \{0,1\}^n, f(x) = b)$. The certificate complexity C(f) of f is defined as $\max(C_0(f), C_1(f))$.

In this work we look at $\{0,1\}^n$ as a set of vertices for a graph Q_n (called the *n*-dimensional Boolean cube or hypercube) in which we have an edge (x,y) whenever $x=(x_1,\ldots,x_n)$ and $y=(y_1,\ldots,y_n)$ differ in exactly one position. We look at subsets $S\subseteq\{0,1\}^n$ as subgraphs (induced by the subset of vertices S) in this graph.

Definition 5. Let c be a partial assignment $c: P \to \{0,1\}, P \subseteq [n]$. An (n-|P|)-dimensional subcube of Q_n is a subgraph G induced on a vertex set $\{x \mid \forall i \in P(x_i = c(i))\}$. It is isomorphic to $Q_{n-|P|}$. We call the value $\dim(G) = n - |P|$ the dimension and the value |P| the co-dimension of G.

Note that each certificate of length l corresponds to one subcube of Q_n with co-dimension l.

Definition 6. Let G be a subcube defined by a partial assignment $c: P \to \{0,1\}, P \subseteq [n]$. Let $c': P \to \{0,1\}$ where $c'(i) \neq c(i)$ for exactly one $i \in P$. Then we call the subcube defined by c' a neighbour subcube of G.

Definition 7. Let G and H be induced subgraphs of Q_n . By $G \cap H$ denote the intersection of G and H that is the graph induced on $V(G) \cap V(H)$. By $G \cup H$ denote the union of G and H that is the graph induced on $V(G) \cup V(H)$. By $G \setminus H$ denote the complement of G in H that is the graph induced by $V(G) \setminus V(H)$.

Definition 8. Let G and H be induced subgraphs of Q_n . By R(G, H) denote the relative size of G in H:

$$R(G,H) = \frac{|G \cap H|}{|H|}. (7)$$

We extend the notion of sensitivity to the induced subgraphs of Q_n :

Definition 9. Let G be a non-empty induced subgraph of Q_n . The sensitivity $s(G, Q_n, x)$ of a vertex $x \in Q_n$ is defined as $|\{i \mid x^{\{i\}} \notin G\}|$, if $x \in G$, and $|\{i \mid x^{\{i\}} \in G\}|$, if $x \notin G$. Then the sensitivity of G is defined as $s(G, Q_n) = \max(s(G, Q_n, x) \mid x \in G)$.

Our results rely on the following generalization of Simon's lemma [11], proved by Ambainis and Vihrovs [3]:

Theorem 2. Let G be a non-empty induced subgraph of Q_n with sensitivity at most s. Then either $R(G,Q_n)=\frac{1}{2^s}$ and G is an (n-s)-dimensional subcube or $R(G,Q_n)\geq \frac{3}{2}\cdot \frac{1}{2^s}$.

3 Upper Bound on Certificate Complexity in Terms of Sensitivity

In this section we prove Corollary 1. In fact, we prove a slightly more specific result.

Theorem 1. Let f be a Boolean function which is not constant. If $s_1(f) = 1$, then $C_0(f) = s_0(f)$. If $s_1(f) > 1$, then

$$C_0(f) \le 2^{s_1(f)-1} \left(s_0(f) - \frac{1}{3} \right).$$
 (8)

Note that a similar bound for $C_1(f)$ follows by symmetry. For the proof, we require the following lemma.

Lemma 1. Let $H_1, H_2, ..., H_k$ be distinct subcubes of Q_n such that the Hamming distance between any two of them is at least 2. Take

$$T = \bigcup_{i=1}^{k} H_i, \qquad T' = \left\{ x \, \middle| \, \exists i \, \left(x^{\{i\}} \in T \right) \right\} \setminus T. \tag{9}$$

If $T \neq Q_n$, then $|T'| \geq |T|$.

Proof. If k=1, then the co-dimension of H_1 is at least 1. Hence H_1 has a neighbour cube, so $|T'| \ge |T| = |H_1|$.

Assume $k \geq 2$. Then $n \geq 2$, since there must be at least 2 bit positions for cubes to differ in. We use an induction on n.

Base case. n=2. Then we must have that H_1 and H_2 are two opposite vertices. Then the other two vertices are in T', hence |T'|=|T|=2.

Inductive step. Divide Q_n into two adjacent (n-1)-dimensional subcubes Q_0 and Q_1 by the value of x_1 . We will prove that the conditions of the lemma hold for each $T \cap Q_b$, $b \in \{0,1\}$. Let $H_u^b = H_u \cap Q_b$. Assume $H_u^b \neq \emptyset$ for some $u \in [k]$. Then either $x_1 = b$ or x_1 is not fixed in H_u . Thus, if there are two non-empty subcubes H_u^b and H_v^b , they differ in the same bit positions as H_u and H_v . Thus the Hamming distance between H_u^b and H_v^b is also at least 2. On the other hand, $Q_b \not\subseteq T$, since then k would be at most 1.

Let $T_b = T \cap Q_b$ and $T_b' = \{x \mid x \in Q_b, \exists i \ (x^{\{i\}} \in T_b)\} \setminus T_b$. Then by induction we have that $|T_b'| \geq |T_b|$. On the other hand, $T_0 \cup T_1 = T$ and $T_0' \cup T_1' \subseteq T'$. Thus

$$|T'| \ge |T_0'| + |T_1'| \ge |T_0| + |T_1| = |T|. \tag{10}$$

Proof of Theorem 1. Let z be a vertex such that f(z) = 0 and $C(f, z) = C_0(f)$. Pick a 0-certificate S_0 of length $C_0(f)$ and $z \in S_0$. It has $m = C_0(f)$ neighbour subcubes which we denote by S_1, S_2, \ldots, S_m .

We work with a graph G induced on a vertex set $\{x \mid f(x) = 1\}$. Since S_0 is a minimum certificate for $z, S_i \cap G \neq \emptyset$ for $i \in [m]$.

As S_0 is a 0-certificate, it gives 1 sensitive bit to each vertex in $G \cap S_i$. Then $s(G \cap S_i, S_i) \leq s_1(f) - 1$.

Suppose $s_1(f) = 1$, then for each $i \in [m]$ we must have that $G \cap S_i$ equals to the whole S_i . But then each vertex in S_0 is sensitive to its neighbour in $G \cap S_i$, so $m \leq s_0(f)$. Hence $C_0(f) = s_0(f)$.

Otherwise $s_1(f) \geq 2$. By Theorem 2, either $R(G, S_i) = \frac{1}{2^{s_1(f)-1}}$ or $R(G, S_i) \geq \frac{3}{2^{s_1(f)}}$ for each $i \in [m]$. We call the cube S_i either light or heavy respectively. We denote the number of light cubes by l, then the number of heavy cubes is m-l. We can assume that the light cubes are S_1, \ldots, S_l .

Let the average sensitivity of $x \in S_0$ be $as(S_0)$. Since each vertex of G in any S_i gives sensitivity 1 to some vertex in S_0 , $\sum_{i=1}^m R(G, S_i) \leq as(S_0)$. Clearly $as(S_0) \leq s_0(f)$. We have that

$$l\frac{1}{2^{s_1(f)-1}} + (m-l)\frac{3}{2^{s_1(f)}} \le as(S_0) \le s_0(f)$$
(11)

$$m\frac{3}{2^{s_1(f)}} - l\frac{1}{2^{s_1(f)}} \le as(S_0) \le s_0(f).$$
(12)

Then we examine two possible cases.

Case 1. $l \leq (s_0(f) - 1)2^{s_1(f)-1}$. Then we have

$$m\frac{3}{2^{s_1(f)}} - (s_0(f) - 1)\frac{2^{s_1(f)-1}}{2^{s_1(f)}} \le as(S_0) \le s_0(f)$$
(13)

$$m\frac{3}{2^{s_1(f)}} \le s_0(f) + \frac{1}{2}(s_0(f) - 1)$$
 (14)

$$m\frac{3}{2^{s_1(f)}} \le \frac{3}{2}s_0(f) - \frac{1}{2} \tag{15}$$

$$m \le 2^{s_1(f)-1} \left(s_0(f) - \frac{1}{3} \right).$$
 (16)

Case 2. $l = (s_0(f) - 1)2^{s_1(f)-1} + \epsilon$ for some $\epsilon > 0$. Since $s_1(f) \ge 2$, the number of light cubes is at least $2(s_0(f) - 1) + \epsilon$, which in turn is at least $s_0(f)$.

Let $\mathcal{F} = \{F \mid F \subseteq [l], |F| = s_0(f)\}$. Denote its elements by $F_1, F_2, \ldots, F_{|\mathcal{F}|}$. We examine $H_1, H_2, \ldots, H_{|\mathcal{F}|}$ – subgraphs of S_0 , where H_i is the set of vertices whose neighbours in S_j are in G for each $j \in F_i$. By Theorem 2, $G \cap S_i$ are subcubes for $i \leq l$. Then so are the intersections of their neighbours in S_0 , including each H_i .

Let $N_{i,j}$ be the common neighbour cube of S_i and S_j that is not S_0 . Suppose $v \in S_0$. Then by v_i denote the neighbour of v in S_i . Let $v_{i,j}$ be the common neighbour of v_i and v_j that is in $N_{i,j}$.

We will show that the Hamming distance between any two subcubes H_i and H_i , $i \neq j$ is at least 2.

Assume there is an edge (u,v) such that $u \in H_i$ and $v \in H_j$. Then $u_k \in G$ for each $k \in F_i$. Since $i \neq j$, there is an index $t \in F_j$ such that $t \notin F_i$. The vertex u is sensitive to S_k for each $k \in F_i$ and, since $|F_i| = s_0(f)$, has full sensitivity. Thus $u_t \notin G$. On the other hand, since each S_k is light, u_k has full 1-sensitivity, hence $u_{k,t} \in G$ for all $k \in F_i$. This gives full 0-sensitivity to u_t . Hence $v_t \notin G$, a contradiction, since $v \in H_j$ and $t \in F_j$.

Thus there are no such edges and the Hamming distance between H_i and H_j is not equal to 1. That leaves two possibilities: either the Hamming distance between H_i and H_j is at least 2 (in which case we are done), or both H_i and H_j are equal to a single vertex v, which is not possible, as then v would have a 0-sensitivity of at least $s_0(f) + 1$.

Let $T = \bigcup_{i=1}^{|\mathcal{F}|} H_i$. We will prove that $T \neq S_0$. Since $s_1(f) \geq 2$, by Theorem 2 it follows that $\dim(G \cap S_i) = \dim(S_i) - s_1(f) + 1 \leq \dim(S_0) - 1$ for each $i \in [l]$. Thus $\dim(H_1) \leq \dim(S_0) - 1$, and $H_1 \neq S_0$. Then it has a neighbour subcube H'_1 in S_0 . But since the Hamming distance between H_1 and any other H_i is at least 2, we have that $H'_1 \cap H_i = \emptyset$, thus T is not equal to S_0 .

Therefore, $H_1, H_2, \ldots, H_{|\mathcal{F}|}$ satisfy all the conditions of Lemma 1. Let T' be the set of vertices in $S_0 \setminus T$ with a neighbour in T. Then, by Lemma 1, $|T'| \geq |T|$ or, equivalently, $R(T', S_0) \geq R(T, S_0)$.

Then note that $R(T', S_0) \geq R(T, S_0) \geq \frac{\epsilon}{2^{s_1(f)-1}}$, since $R(G, S_i) = \frac{1}{2^{s_1(f)-1}}$ for all $i \in [l]$, there are a total of $(s_0(f) - 1)2^{s_1(f)-1} + \epsilon$ light cubes and each vertex in S_0 can have at most $s_0(f)$ neighbours in G.

Let S_h be a heavy cube, and $i \in [|\mathcal{F}|]$. The neighbours of H_i in S_h must not be in G, or the corresponding vertex in H_i would have sensitivity $s_0(f) + 1$.

Let $k \in F_i$. As S_k is light, all the vertices in $G \cap S_k$ are fully sensitive, therefore all their neighbours in $N_{k,h}$ are in G. Therefore all the neighbours of H_i in S_h already have full 0-sensitivity. Then all their neighbours must also not be in G.

This means that vertices in T' can only have neighbours in G in light cubes. But they can have at most $s_0(f)-1$ such neighbours each, otherwise they would be in T, not in T'. As $R(T', S_0) \geq \frac{\epsilon}{2^{s_1(f)-1}}$, the average sensitivity of vertices in S_0 is at most

$$as(S_0) \le s_0(f)R(S_0 \setminus T', S_0) + (s_0(f) - 1)R(T', S_0) \le$$
(17)

$$\leq s_0(f) \left(1 - \frac{\epsilon}{2^{s_1(f)-1}}\right) + (s_0(f) - 1) \frac{\epsilon}{2^{s_1(f)-1}} =$$
 (18)

$$= s_0(f) - \frac{\epsilon}{2^{s_1(f)-1}}. (19)$$

Then by inequality (12) we have

$$m\frac{3}{2^{s_1(f)}} - \left((s_0(f) - 1)2^{s_1(f) - 1} + \epsilon \right) \frac{1}{2^{s_1(f)}} \le s_0(f) - \frac{\epsilon}{2^{s_1(f) - 1}}. \tag{20}$$

Rearranging the terms, we get

$$m\frac{3}{2^{s_1(f)}} \le \left((s_0(f) - 1)2^{s_1(f) - 1} + \epsilon \right) \frac{1}{2^{s_1(f)}} + s_0(f) - \frac{\epsilon}{2^{s_1(f) - 1}} \tag{21}$$

$$m\frac{3}{2^{s_1(f)}} \le s_0(f) + \frac{1}{2}(s_0(f) - 1) - \frac{\epsilon}{2^{s_1(f)}}$$
 (22)

$$m\frac{3}{2^{s_1(f)}} \le \frac{3}{2}s_0(f) - \frac{1}{2} - \frac{\epsilon}{2^{s_1(f)}}$$
(23)

$$m \le 2^{s_1(f)-1} \left(s_0(f) - \frac{1}{3} \right) - \frac{\epsilon}{3}.$$
 (24)

Theorem 1 immediately implies Corollary 1:

Proof of Corollary 1. If f is constant, then C(f) = s(f) = 0 and the statement is true. Otherwise by Theorem 1

$$C(f) = \max(C_0(f), C_1(f)) \le$$
 (25)

$$\leq \max_{b \in \{0,1\}} \left(\max \left(2^{s_{1-b}(f)-1} \left(s_b(f) - \frac{1}{3} \right), s_b(f) \right) \right) \leq \tag{26}$$

$$\leq \max\left(2^{s(f)-1}\left(s(f) - \frac{1}{3}\right), s(f)\right) \tag{27}$$

On the other hand, $bs(f) \leq C(f)$ is a well-known fact.

4 Relation between $C_0(f)$ and $s_0(f)$ for $s_1(f) = 2$

Ambainis and Sun exhibited a class of functions that achieves the best known separation between sensitivity and block sensitivity, which is quadratic in terms of s(f) [2]. This function also produces the best known separation between 0-certificate complexity and 0/1-sensitivity:

Theorem 3. For arbitrary $s_0(f)$ and $s_1(f)$, there exists a function f such that

$$C_0(f) = \left(\frac{2}{3} + o(1)\right) s_0(f) s_1(f). \tag{28}$$

Thus it is possible to achieve a quadratic gap between the two measures. As $bs_0(f) \leq C_0(f)$, it would be tempting to conjecture that quadratic separation is the largest possible. Therefore we are interested both in improved upper bounds and in functions that achieve quadratic separation with a larger constant factor.

In this section, we examine how $C_0(f)$ and $s_0(f)$ relate to each other for small $s_1(f)$. If $s_1(f) = 1$, it follows by Theorem 1 that $C_0(f) = s_0(f)$. Therefore we consider the case $s_1(f) = 2$.

Here we are able to construct a separation that is better than (28) by a constant factor.

Theorem 4. There is a function f with $s_1(f) = 2$ and arbitrary $s_0(f)$ such that

$$C_0(f) = \left| \frac{3}{4} s_0(f) s_1(f) \right| = \left| \frac{3}{2} s_0(f) \right|.$$
 (29)

Proof. Consider the function that takes value 1 iff its 4 input bits are in either ascending or descending sorted order. Formally,

$$SORT_4(x) = 1 \Leftrightarrow (x_1 \le x_2 \le x_3 \le x_4) \lor (x_1 \ge x_2 \ge x_3 \ge x_4).$$
 (30)

One easily sees that $C_0(SORT_4) = 3$, $s_0(SORT_4) = 2$ and $s_1(SORT_4) = 2$.

Denote the 2-bit logical AND function by AND₂. We have $C_0(\text{AND}_2) = s_0(\text{AND}_2) = 1$ and $s_1(\text{AND}_2) = 2$.

To construct the examples for larger $s_0(f)$ values, we use the following fact (it is easy to show, and a similar lemma was proved in [2]):

Fact 1. Let f and g be Boolean functions. By composing them with OR to $f \lor g$ we get

$$C_0(f \vee g) = C_0(f) + C_0(g), \tag{31}$$

$$s_0(f \vee g) = s_0(f) + s_0(g), \tag{32}$$

$$s_1(f \vee g) = \max(s_1(f), s_1(g)).$$
 (33)

Suppose we need a function with $k = s_0(f)$. Assume k is even. Then by Fact 1 for $g = \bigvee_{i=1}^{\frac{k}{2}} \operatorname{SORT}_4$ we have $C_0(g) = \frac{3}{2}k$. If k is odd, consider the function $g = \left(\bigvee_{i=1}^{\frac{k-1}{2}} \operatorname{SORT}_4\right) \vee \operatorname{AND}_2$. Then by Fact 1 we have $C_0(g) = 3 \cdot \frac{k-1}{2} + 1 = \left\lfloor \frac{3}{2}k \right\rfloor$. \square

A curious fact is that both examples of (28) and Theorem 4 are obtained by composing some primitives using OR. The same fact holds for the best examples of separation between bs(f) and s(f) that preceded the [2] construction [10,12].

We are also able to prove a slightly better upper bound in case $s_1(f) = 2$.

Theorem 5. Let f be a Boolean function with $s_1(f) = 2$. Then

$$C_0(f) \le \frac{9}{5}s_0(f).$$
 (34)

Proof. Let z be a vertex such that f(z) = 0 and $C(f, z) = C_0(f)$. Pick a 0-certificate S_0 of length $m = C_0(f)$ and $z \in S_0$. It has m neighbour subcubes which we denote by S_1, S_2, \ldots, S_m . Let $n' = n - m = \dim(S_i)$ for each S_i .

We work with a graph G induced on a vertex set $\{x \mid f(x) = 1\}$. Let $G_i = G \cap S_i$. As S_0 is a minimal certificate for z, we have $G_i \neq \emptyset$ for each $i \in [m]$. Since any $v \in G_i$ is sensitive to S_0 , we have $s(G_i, S_i) \leq 1$. Thus by Theorem 2 either G_i is an (n'-1)-subcube of S_i with $R(G_i : S_i) = \frac{1}{2}$ or $R(G_i : S_i) \geq \frac{3}{4}$. We call S_i light or heavy, respectively.

Let $N_{i,j}$ be the common neighbour cube of S_i , S_j that is not S_0 . Let $G_{i,j} = G \cap N_{i,j}$. Suppose $v \in S_0$. Let v_i be the neighbour of v in S_i . Let $v_{i,j}$ be the neighbour of v_i and v_j in $N_{i,j}$.

Let S_i, S_j be light. By G_i^0, G_j^0 denote the neighbour cubes of G_i, G_j in S_0 . We call $\{S_i, S_j\}$ a pair, iff $G_i^0 \cup G_j^0 = S_0$. In other words, a pair is defined by a single dimension. Also we have either $z_i \notin G$ or $z_j \notin G$: we call the corresponding cube the representative of this pair.

Proposition 1. Let \mathcal{P} be a set of mutually disjoint pairs of the neighbour cubes of S_0 . Then there exists a 0-certificate S_0' such that $z \in S_0'$, $\dim(S_0') = \dim(S_0)$ and S_0' has at least $|\mathcal{P}|$ heavy neighbour cubes.

Proof. Let \mathcal{R} be a set of mutually disjoint pairs of the neighbour cubes of S_0 . W.l.o.g. let $S_1, \ldots, S_{|\mathcal{R}|}$ be the representatives of \mathcal{R} . Let F_i be the neighbour cube of $S_i \setminus G$ in S_0 . Let $B_{\mathcal{R}} = \bigcap_{i=1}^{|\mathcal{R}|} F_i$. Suppose $S_0 + x$ is a coset of S_0 and $x_t = 0$ if the t-th dimension is not fixed in S_0 : let $B_{\mathcal{R}}(S_0 + x)$ be $B_{\mathcal{R}} + x$.

Pick $\mathcal{R} \subseteq \mathcal{P}$ with the largest size, such that for each two representatives S_i , S_j of \mathcal{R} , $B_{\mathcal{R}}(N_{i,j})$ is a 0-certificate.

We will prove that the subcube S'_0 spanned by $B_{\mathcal{R}}, B_{\mathcal{R}}(S_1), \ldots, B_{\mathcal{R}}\left(S_{|\mathcal{R}|}\right)$ is a 0-certificate. It corresponds to an $|\mathcal{R}|$ -dimensional hypercube $Q_{|\mathcal{R}|}$ where $B_{\mathcal{R}}(S_0 + x)$ corresponds to a single vertex for each coset $S_0 + x$ of S_0 .

Let $T \subseteq Q_{|\mathcal{R}|}$ be the graph induced on $\{v \mid B_{\mathcal{R}}(H) \text{ corresponds to } v, B_{\mathcal{R}}(H) \text{ is not a 0-certificate}\}$. Then we have $s(T, Q_{|\mathcal{R}|}) \leq 2$. Suppose $B_{\mathcal{R}}$ corresponds to $0^{|\mathcal{R}|}$. Let L_d be the set of $Q_{|\mathcal{R}|}$ vertices that are at distance d from $0^{|\mathcal{R}|}$. We prove by induction that $L_d \cap T = \emptyset$ for each d.

Proof. Base case. $d \leq 2$. The required holds since all $B_{\mathcal{R}}, B_{\mathcal{R}}(S_i), B_{\mathcal{R}}(N_{i,j})$ are 0-certificates.

Inductive step. $d \geq 3$. Examine $v \in L_d$. As v has d neighbours in L_{d-1} , $L_{d-1} \cap T = \emptyset$ and $s(T, Q_{|\mathcal{R}|}) \leq 2$, we have that $v \notin T$.

Let k be the number of distinct dimensions that define the pairs of \mathcal{R} , then $k \leq |\mathcal{R}|$. Hence $\dim(S'_0) = |\mathcal{R}| + \dim(B_{\mathcal{R}}) = |\mathcal{R}| + (\dim(S_0) - k) \geq \dim(S_0)$. But S_0 is a minimal 0-certificate for z, therefore $\dim(S'_0) = \dim(S_0)$.

Note that a light neighbour S_i of S_0 is separated into a 0-certificate and a 1-certificate by a single dimension, hence we have $s(G, S_i, v) = 1$ for every $v \in S_i$. As S_i neighbours S_0 , every vertex in its 1-certificate is fully sensitive. The same holds for any light neighbour S_i' of S_0' .

Now we will prove that each pair in \mathcal{P} provides a heavy neighbour for S'_0 . Let $\{S_a, S_b\} \in \mathcal{P}$, where S_a is the representative. We distinguish two cases:

- $-B_{\mathcal{R}}(S_b)$ is a 1-certificate. Since S_b is light, it has full 1-sensitivity. Therefore, $v \in G$ for all $v \in B_R(N_{i,b})$, for each $i \in [|\mathcal{R}|]$. Let S_b' be the neighbour of S_0' that contains $B_{\mathcal{R}}(S_b)$ as a subcube. Then for each $v \in B_{\mathcal{R}}(S_b)$ we have $s(G, S_b', v) = 0$. Hence S_b' is heavy.
- Otherwise, $\{S_a, S_b\}$ is defined by a different dimension than any of the pairs in \mathcal{R} . Let $\mathcal{R}' = \mathcal{R} \cup \{S_a, S_b\}$. Examine the subcube $B_{\mathcal{R}'}$. By definition of \mathcal{R} , there is a representative S_i of \mathcal{R} such that $B_{\mathcal{R}'}(N_{i,a})$ is not a 0-certificate. Let S'_a be the neighbour of S'_0 that contains $B_{\mathcal{R}}(S_a)$ as a subcube. Then there is a vertex $v \in B_{\mathcal{R}'}(S_a)$ such that $s(G, S'_a, v) \geq 2$. Hence S'_a is heavy.

Let \mathcal{P} be the largest such set. Let l and h=m-l be the number of light and heavy neighbours of S_0 , respectively. Each pair in \mathcal{P} gives one neighbour in G to each vertex in S_0 . Now examine the remaining $l-2|\mathcal{P}|$ light cubes. As they are not in \mathcal{P} , no two of them form a pair. Hence there is a vertex $v \in S_0$ that is sensitive to each of them. Then $s_0(f) \geq s_0(f,v) \geq |\mathcal{P}| + (l-2|\mathcal{P}|) = l - |\mathcal{P}|$. Therefore $|\mathcal{P}| \geq l - s_0(f)$.

Let q be such that $m = qs_0(f)$. Then there are $qs_0(f) - l$ heavy neighbours of S_0 . On the other hand, by Proposition 1, there exists a minimal certificate S'_0 of z with at least $l - s_0(f)$ heavy neighbours. Then z has a minimal certificate with at least $\frac{(qs_0(f)-l)+(l-s_0(f))}{2} = \frac{q-1}{2} \cdot s_0(f)$ heavy neighbour cubes.

W.l.o.g. let S_0 be this certificate. Then $l=qs_0(f)-h\leq (q-\frac{q-1}{2})s_0(f)=\frac{q+1}{2}\cdot s_0(f)$. As each $v\in G_i$ for $i\in [m]$ gives sensitivity 1 to its neighbour in S_0 ,

$$l\frac{1}{2} + h\frac{3}{4} \le s_0(f). \tag{35}$$

Since the constant factor at l is less than at h, we have

$$\frac{q+1}{2} \cdot s_0(f) \cdot \frac{1}{2} + \frac{q-1}{2} \cdot s_0(f) \cdot \frac{3}{4} \le s_0(f) \tag{36}$$

By dividing both sides by $s_0(f)$ and simplifying terms, we get $q \leq \frac{9}{5}$.

This result shows that the bound of Theorem 1 can be improved. However, it is still not tight. For some special cases, through extensive casework we can also prove the following results:

Theorem 6. Let f be a Boolean function with $s_1(f) = 2$ and $s_0(f) \ge 3$. Then

$$C_0(f) \le 2s_0(f) - 2.$$
 (37)

Theorem 7. Let f be a Boolean function with $s_1(f) = 2$ and $s_0(f) \ge 5$. Then

$$C_0(f) \le 2s_0(f) - 3.$$
 (38)

These theorems imply that for $s_1(f) = 2$, $s_0(f) \le 6$ we have $C_0(f) \le \frac{3}{2}s_0(f)$, which is the same separation as achieved by the example of Theorem 4. This leads us to the following conjecture:

Conjecture 1. Let f be a Boolean function with $s_1(f) = 2$. Then

$$C_0(f) \le \frac{3}{2}s_0(f).$$
 (39)

We consider $s_1(f) = 2$ to be the simplest case where we don't know the actual tight upper bound on $C_0(f)$ in terms of $s_0(f)$, $s_1(f)$. Proving Conjecture 1 may provide insights into relations between C(f) and s(f) for the general case.

References

- A. Ambainis, M. Bavarian, Y. Gao, J. Mao, X. Sun, and S. Zuo. Tighter relations between sensitivity and other complexity measures. In J. Esparza, P. Fraigniaud, T. Husfeldt, and E. Koutsoupias, editors, *International Colloquium on Automata*, *Languages, and Programming*, volume 8572 of *Lecture Notes in Computer Science*, pages 101–113. Springer Berlin Heidelberg, 2014.
- 2. A. Ambainis and X. Sun. New separation between s(f) and bs(f). CoRR, abs/1108.3494, 2011.
- 3. A. Ambainis and J. Vihrovs. Size of sets with small sensitivity: a generalization of Simon's lemma. In R. Jain, S. Jain, and F. Stephan, editors, *Theory and Applications of Models of Computation*, volume 9076 of *Lecture Notes in Computer Science*, pages 122–133. Springer International Publishing, 2015.
- 4. R. Beals, H. Buhrman, R. Cleve, M. Mosca, and R. de Wolf. Quantum lower bounds by polynomials. *J. ACM*, 48(4):778–797, 2001.
- 5. H. Buhrman and R. de Wolf. Complexity measures and decision tree complexity: a survey. *Theoretical Computer Science*, 288(1):21 43, 2002.
- S. Cook, C. Dwork, and R. Reischuk. Upper and lower time bounds for parallel random access machines without simultaneous writes. SIAM Journal on Computing, 15:87–97, 1986.
- 7. P. Hatami, R. Kulkarni, and D. Pankratov. Variations on the Sensitivity Conjecture. Number 4 in Graduate Surveys. Theory of Computing Library, 2011.
- C. Kenyon and S. Kutin. Sensitivity, block sensitivity, and ℓ-block sensitivity of Boolean functions. Information and Computation, 189(1):43 − 53, 2004.
- 9. N. Nisan. CREW PRAMS and decision trees. In *Proceedings of the Twenty-first Annual ACM Symposium on Theory of Computing*, STOC '89, pages 327–335, New York, NY, USA, 1989. ACM.
- 10. D. Rubinstein. Sensitivity vs. block sensitivity of Boolean functions. *Combinatorica*, 15(2):297–299, 1995.
- 11. H.-U. Simon. A tight $\Omega(\log \log N)$ -bound on the time for parallel RAM's to compute nondegenerated Boolean functions. In *Proceedings of the 1983 International FCT-Conference on Fundamentals of Computation Theory*, pages 439–444, London, UK, 1983. Springer-Verlag.
- 12. M. Virza. Sensitivity versus block sensitivity of Boolean functions. *Information Processing Letters*, 111(9):433 435, 2011.

A Proof of Theorem 6

Proof. Let G be a graph induced on a vertex set $\{x \mid f(x) = 1\}$. Suppose z is a vertex such that f(z) = 0 and $C(f, z) = C_0(f)$. Pick a 0-certificate S_0 of length $C_0(f)$ and $z \in S_0$. It has $m = C_0(f)$ neighbour subcubes which we denote by S_1, S_2, \ldots, S_m . Since S_0 is a minimum certificate for z, we have that $S_i \cap G \neq \emptyset$ for $i \in [m]$. Let the dimension of S_0 be n'.

Let $N_{i,j}$ be the common neighbour cube of S_i and S_j that is not S_0 . Suppose $v \in S_0$. Then by v_i denote the neighbour of v in S_i . Let $v_{i,j}$ be the common neighbour of v_i and v_j that is in $N_{i,j}$.

As S_0 is a 0-certificate, each vertex of G in any S_i is sensitive to its neighbour in S_0 . Then $s(G \cap S_i, S_i) \leq 1$. By Theorem 2, it follows that either $R(G, S_i) = \frac{1}{2}$ and $G \cap S_i$ is a (n'-1)-dimensional subcube or $R(G, S_i) \geq \frac{3}{4}$. We call such subcubes light or heavy respectively. Let the number of light cubes be l, then the number of heavy cubes is m-l. Assume the light cubes are S_1, S_2, \ldots, S_l .

Each vertex of G in any S_i gives sensitivity 1 to its neighbour in S_0 , thus

$$\sum_{i=1}^{m} R(G, S_i) \le s_0(f). \tag{40}$$

Hence

$$l\frac{1}{2} + (m-l)\frac{3}{4} \le s_0(f),\tag{41}$$

$$3m - 4s_0(f) \le l. \tag{42}$$

Assume on the contrary that $C_0(f) > 2s_0(f) - 2$ or equivalently $m \ge 2s_0(f) - 1$. In that case $l \ge 3(2s_0(f) - 1) - 4s_0(f) = 2s_0(f) - 3$.

Let $s_0(f) = 3$, then $l \ge 3$. First assume l = 3. Then S_1, S_2, S_3 are light and S_4, S_5 are heavy. This implies

$$3 = s_0(f) \ge \sum_{i=1}^m R(G, S_i) \ge 3 \cdot \frac{1}{2} + 2 \cdot \frac{3}{4} = 3.$$
 (43)

Thus for each $j \in [4; 5]$ we have $R(G, S_j) = \frac{3}{4}$, which means that $R(S_j \setminus G, S_j) = \frac{1}{4}$. By Theorem 2, $s(S_j \setminus G, S_j) \ge 2$. Let $v_j \notin G$ be a vertex with $s(S_j \setminus G, S_j, v_j) = 2$. Assume $v_i \in G$ for some $i \in [3]$. As v_i has full sensitivity, its neighbour $v_{i,j} \in G$. Hence there is at most one such $i \in [3]$ that $v_i \in G$. But then $v \in S_0$ can only have neighbours in G in one light and one heavy cube. Hence v does not have full sensitivity: a contradiction, since $\sum_{i=1}^m R(G, S_i) = 3$. Thus we have $l \ge 4$ for $s_0(f) = 3$.

Now let $s_0(f) \geq 3$. Assume there is a vertex $v \in S_0$ that has $s_0(f)$ neighbours in G among light cubes. Let S_i be a light cube with $v_i \in G$. If $s_0(f) \geq 4$, then $l \geq 2s_0(f) - 3 > s_0(f)$; if $s_0(f) = 3$, then $l \geq 4 > 3$. Thus $l > s_0(f)$. Hence there is a light cube S_j such that $v_j \notin G$. Since v_i has full sensitivity, $v_{i,j} \in G$, and there are $s_0(f)$ such i. But v_j is also sensitive to one neighbour in S_j ; hence v_j has sensitivity $s_0(f) + 1$, a contradiction.

Thus any vertex $v \in S_0$ has at most $s_0(f) - 1$ neighbours in G in light cubes. Then $l \leq 2(s_0(f) - 1)$, otherwise we would have a contradiction by the pigeonhole principle. If there are no heavy cubes, then $m = l \leq 2s_0(f) - 2$ and we are done. Otherwise there is a heavy cube S_h . Let T be the subset of vertices in S_0 that each has exactly $s_0(f) - 1$ neighbours in G in light cubes. Since $l \geq 2s_0(f) - 3$, we have $R(T, S_0) \geq \frac{1}{2}$.

Pick a vertex $v \in S'$. Let S_i, S_j be light cubes with $v_i \in G$ and $v_j \notin G$. If $s_0(f) \geq 4$, then by $l \geq 2s_0(f) - 3$ we have that the number of choices for j is at least $(2s_0(f) - 3) - (s_0(f) - 1) = 2$; if $s_0(f) = 3$, then since $l \geq 4$, this number is also at least 2. Since v_i has full sensitivity, $v_{i,j} \in G$, and there are $s_0(f) - 1$ choices for i. On the other hand, as S_j is a light cube, v_j is sensitive to a neighbour in S_j . Hence v_j has full sensitivity, so its neighbour $v_{j,h} \notin G$. But then v_h has at least 3 neighbours not in G, and, as $s_1(f) = 2$, we have $v_h \notin G$.

This shows that for a vertex $v \in S'$, its neighbour in S_h does not belong to G. Let S'_h be the set of S' neighbours in S_h . Then $R(S_h \setminus G, S_h) \ge R(S'_h, S_h) = R(S', S_0) \ge \frac{1}{2}$: a contradiction, since S_h is a heavy cube.

B Proof of Theorem 7

Lemma 2. Let G be a non-empty subgraph of Q_n induced on the vertex set $\{x \mid f(x) = 1\}$ of a function f with $s_1(f) \leq 1$. Then either $G = Q_n$, or G is an (n-1)-dimensional subcube, or the number of fully sensitive vertices in G is at least $2|Q_n \setminus G|$. Furthermore, in the last case each vertex in $Q_n \setminus G$ has a sensitivity of at least 2.

Proof. We examine the induced graph $Q_n \setminus G$. Each connected component in this graph must be a subcube, otherwise some of the vertices of G in the smallest subcube containing the component would have sensitivity at least 2. Furthermore, the Hamming distance between any two of these subcubes is at least 3, otherwise the vertices between them would have sensitivity at least 2.

If there are no such subcubes, $G = Q_n$. If there is such an (n-1)-subcube, it must be the only one and G is its opposite (n-1)-subcube. Otherwise each of these subcubes is an (n-2)-subcube or smaller. Therefore each vertex in them has at least 2 neighbours in G. Since $s_1(f) \leq 1$, these 2 neighbours are fully sensitive and must be different for each such vertex, thus there are at least $2|Q_n \setminus G|$ of them.

We denote a subcube that can be obtained by fixing some continuous sequence b of starting bits by Q_b . For example, Q_0 and Q_1 can be obtained by fixing the first bit and Q_{01} can be obtained by fixing the first two bits to 01. We use a wildcard * symbol to indicate that the bit in the corresponding position is not fixed. For example, by Q_{*10} we denote a cube obtained by fixing the second and the third bit to 10.

Proof of Theorem 7. Assume on the contrary that such a function exists. Denote $m = C_0(f) = 2s_0(f) - 2$ and $k = s_0(f) \ge 5$.

We work with a graph G induced on a vertex set $\{x \mid f(x) = 1\}$. W.l.o.g. let z be a vertex such that f(z) = 0 and $C(f, z) = C_0(f)$. Pick a 0-certificate S_0 of length $m = C_0(f)$ and $z \in S_0$. It has m neighbour subcubes which we denote by S_1, S_2, \ldots, S_m . Let n' = n - m, the number of dimensions in each of S_i .

Each vertex of G in any S_i gives sensitivity 1 to some vertex in S_0 , therefore $\sum_{i=1}^m R(G, S_i) \leq s_0(f) = k$.

Let $G_i = G \cap S_i$. Each G_i is nonempty, otherwise we would obtain a shorter certificate for z.

Since any 1-vertex in each of S_i is sensitive to S_0 , we have $s(G, S_i) \le s_1(f) - 1 = 1$. Thus by Theorem 2 G_i can be either an (n' - 1)-subcube of S_i with $R(G_i, S_i) = \frac{1}{2}$ or $R(G_i, S_i) \ge \frac{3}{4}$. We will call these cubes *light* or *heavy*, respectively.

Let F_i be the set of fully sensitive vertices in G_i . Note that $F_i = G_i$ for light cubes and $|F_i| = 2|S_i \setminus G_i|$ for heavy cubes by Lemma 2.

By $N_{i,j}$ denote the common neighbour cube of S_i , S_j that is not S_0 . For a vertex $v \in S_0$, denote by v_i its neighbour in S_i . For $v_i, v_j, i \neq j$, by $v_{i,j}$ denote their common neighbour in $N_{i,j}$.

We will first show that no vertex in S_0 has k fully sensitive neighbours. Assume that there exists such a vertex v. W.l.o.g. assume that its k fully sensitive neighbours are in S_1, \ldots, S_k . Then examine v_m . As v is already fully sensitive, $v_m \notin G$. As $v_i \in G$ and fully sensitive for each $i \in [k]$, we have that each $v_{i,m}$ is also in G.

But then it follows by induction that no vertex in S_m can be in G. As a basis, we have that v_m is also fully sensitive, therefore all of its neighbours in S_m must also not be in G. Now assume that all vertices in S_m with a distance to v_m no more that $i \geq 1$ are not in G, then examine a vertex u_m at distance i + 1. u_m differs from v_m in at least $i + 1 \geq 2$ bits, therefore it has at least 2 neighbours closer to v_m which are in $S_m \setminus G$. But we have that G_m has sensitivity at most $s_1(f) - 1 = 1$ inside S_m , therefore $u_m \notin G$.

But then $G_m = \emptyset$, a contradiction.

Since $\sum_{i=1}^{m} R(G, S_i) \leq k$, we have that there are at most 4 heavy cubes. We will now examine each possible number of heavy cubes separately.

First we examine the case where there are m-4 light cubes and 4 heavy cubes. Since $\sum_{i=1}^{m} R(G, S_i) \leq k$, for each heavy cube S_i it holds that $R(G_i : S_i) = \frac{3}{4}$ exactly.

Then by Lemma 2, $R(F_i, S_i) \ge \frac{1}{2}$ for all i. Then $\sum_{i=1}^m R(F_i, S_i) \ge k-1$. Since no vertex in S_0 can have k fully sensitive neighbours, we have that each vertex in S_0 has exactly k-1 fully sensitive neighbours.

Now examine a vertex $v \in S_0$ which has at least k-3 neighbours in G in light cubes, such a vertex must exist as there are a total of 2k-6 light cubes in this case. W.l.o.g. assume that its k-1 fully sensitive neighbours are in S_1, \ldots, S_{k-1} . As v already has k-3 neighbours in G in light cubes and can have at most k neighbours in G, it must have a neighbour not in G in a heavy cube, W.l.o.g.

let that heavy cube be S_m . Then examine $v_m \notin G$. As $v_i \in G$ and fully sensitive for each $i \in [k-1]$, we have that each $v_{i,m}$ is also in G. In addition, v_m has at least 2 neighbours in G in S_m by Lemma 2. But then it has sensitivity at least k+1, a contradiction.

Now we examine the case where there are no heavy cubes. $R(F_i, S_i) = \frac{1}{2}$ for all i and $\sum_{i=1}^{m} R(F_i, S_i) = k - 1$. Since no vertex in S_0 can have k fully sensitive neighbours, we have that each vertex in S_0 has exactly k - 1 fully sensitive neighbours in G.

If no two of G_i were opposite subcubes in $Q_{n'}$, they would all overlap in $Q_{n'}$, giving a vertex in S_0 with sensitivity m > k. Therefore at least 2 of them are opposite, W.l.o.g. let $G_1 = Q_0 \cap S_1$ and $G_2 = Q_1 \cap S_2$.

Now W.l.o.g. examine the case where $z \in Q_0$. Then we examine the 0-certificate $S_0' = Q_0 \cap (S_0 \cup S_2)$. It is also a minimal certificate, but has a heavy neighbour $Q_0 \cap (S_1 \cup N_{1,2})$, therefore we have reduced this case to the cases where at least one of S_i is heavy.

We now examine the case where there are m-1 light cubes and 1 heavy cube, let them be S_1, \ldots, S_{m-1} and S_m respectively. Note that $m-1 \ge k+1$, as $k \ge 5$. Then $\sum_{i=1}^{m-1} R(F_i, S_i) = \frac{1}{2}(2k-3) = k-\frac{3}{2}$. Since no vertex in S_0 can have k fully sensitive neighbours, we have that half of all vertices in S_0 have exactly k-1 fully sensitive neighbours in G in light cubes.

We now examine one such vertex $v \in S_0$. W.l.o.g. assume that its k-1 fully sensitive neighbours are in S_1, \ldots, S_{k-1} . Then for all $i \in [k-1]$ we have that $v_{i,k}, v_{i,k+1}, v_{i,m}$ are all in G. Since S_k and S_{k+1} are light, we have that v_k and v_{k+1} are fully sensitive. Therefore $v_{k,m}$ and $v_{k+1,m}$ are not in G. Therefore v_m has at least 3 neighbours not in G and is also not in G. But then v_m has at least 2 additional neighbours in G by Lemma 2. Therefore it has sensitivity at least k+1, a contradiction.

We now examine the case where there are m-2 light cubes and 2 heavy cubes, let them be S_1, \ldots, S_{m-2} and S_{m-1}, S_m respectively. Then $\sum_{i=1}^{m-2} R(F_i, S_i) = k-2$. If there exists a vertex $v \in S_0$ with k-1 neighbours in G in light cubes, we can derive a contradiction similarly to the case with 1 heavy cube (since $m-2 \ge k+1$ as well). Therefore no such vertex exists and each vertex in S_0 has exactly k-2 neighbours in G in light cubes.

We will next show that $G_{m-1} = S_{m-1}$ and $G_m = S_m$. Assume otherwise, that there exists a vertex in S_{m-1}, S_m not in G. W.l.o.g. let $v_m \notin G$. By Lemma 2, it has a fully sensitive neighbour $u_m \in G_m$. We have that exactly k-2 of u_1, \ldots, u_{m-2} are in G, W.l.o.g. let them be u_1, \ldots, u_{k-2} . Now examine a vertex $u_i \notin G$, $i \in [k-1, m-2]$. There are k-2 such u_i . We have that $u_{j,i}$ belongs to G for all $j \in [k-2]$. Additionally so does $u_{i,m}$. Finally, u_i has sensitivity 1 in S_i . Then it is fully sensitive, therefore $u_{i,m-1} \notin G$. Then u_{m-1} has at least k-1 neighbours not in G and it is also not in G. But $u_{j,m-1}$ belongs to G for all $j \in [k-2]$ and so does $u_{m-1,m}$. By Lemma 2, u_{m-1} also has sensitivity 2 in S_{m-1} , giving it a total sensitivity of k+1, a contradiction.

If no two light cubes were opposite subcubes in $Q_{n'}$, they would all overlap in $Q_{n'}$, giving a vertex in S_0 with m-2 neighbours in G in light cubes. Therefore at least 2 of them are opposite, W.l.o.g. let them be G_1 and G_2 . Then we have that each vertex in S_0 has exactly k-3 neighbours in G in G_3, \ldots, G_{m-2} and again we have that at least 2 of them must be opposite, W.l.o.g. let them be G_3, G_4 . Similarly we obtain that G_{2i-1}, G_{2i} are opposite for all $i \in [\frac{m-2}{2}]$. Now we examine 2 cases:

- 1. All pairs of light cubes consist of the same subcubes. W.l.o.g. we have that $G_{2i-1} = Q_0$ and $G_{2i} = Q_1$ for all $i \in \left[\frac{m-2}{2}\right]$. Now w.l.o.g. examine the case where $z \in Q_0$. Then we examine the 0-certificate $S_0' = Q_0 \cap (S_0 \cup S_2)$. It is also a minimal certificate, but has heavy neighbours $Q_0 \cap (S_{2i-1} \cup N_{2i-1,2})$ for all $i \in \left[\frac{m-2}{2}\right]$. Therefore we have reduced this case to the cases where at least 3 of S_i are heavy.
- 2. At least 2 of the pairs of light cubes consist of different subcubes. W.l.o.g. we have that $G_1 = Q_0, G_2 = Q_1, G_3 = Q_{*0}, G_4 = Q_{*1}$. W.l.o.g. assume that $z \in Q_{00}$.

Then we examine the 0-certificate $S_0' = Q_0 \cap (S_0 \cup S_2)$. It is also a minimal certificate, and has a heavy neighbour $Q_0 \cap (S_1 \cup N_{1,2})$. If it has 2 more heavy neighbours we have reduced this case to the cases where at least 3 of S_i are heavy. Otherwise at least one of $Q_0 \cap (S_{m-1} \cup N_{2,m-1})$, $Q_0 \cap (S_m \cup N_{2,m})$ is light. W.l.o.g. assume it is $Q_0 \cap (S_{m-1} \cup N_{2,m-1})$. Since $G_{m-1} = S_{m-1}$, this means that $Q_0 \cap N_{2,m-1}$ does not have any vertices in G. Then $Q_0 \cap S_{m-1}$ is fully sensitive.

Now examine the 0-certificate $S_0'' = Q_{*0} \cap (S_0 \cup S_4)$. It is also a minimal certificate, and has a heavy neighbour $Q_{*0} \cap (S_3 \cup N_{3,4})$. If it has 2 more heavy neighbours we have reduced this case to the cases where at least 3 of S_i are heavy. Otherwise at least one of $Q_{*0} \cap (S_{m-1} \cup N_{4,m-1})$, $Q_{*0} \cap (S_m \cup N_{4,m})$ is light. As $Q_0 \cap S_{m-1}$ is fully sensitive, we have that $Q_0 \cap N_{4,m-1} \subset G$ and it must be the case that $Q_{*0} \cap (S_m \cup N_{4,m})$ is light. Since $G_m = S_m$, this means that $Q_{*0} \cap N_{4,m}$ does not have any vertices in G. Then $Q_{*0} \cap S_m$ is fully sensitive.

Now note that, as G_5 and G_6 are opposite subcubes, one of z_5 , z_6 must not belong to G. W.l.o.g. let it be z_5 . As each vertex in S_0 has exactly k-2 neighbours in G in light cubes, z_5 has at least k-2 neighbours in G in $N_{5,j}$ for $j \in [m-2]$. Since $Q_0 \cap S_{m-1}$ and $Q_{*0} \cap S_m$ are fully sensitive, $z_{5,m-1}$ and $z_{5,m}$ are also in G. As z_5 also has a neighbour in G in S_5 , it has a sensitivity of at least k+1, a contradiction.

We are left with the case where there are m-3 light cubes and 3 heavy cubes, let them be S_1, \ldots, S_{m-3} and S_{m-2}, S_{m-1}, S_m respectively.

We will first show that no vertex in S_0 has k-1 neighbours in G in light cubes. Assume on the contrary that such a vertex $v \in S_0$ exists. W.l.o.g. assume that its k-1 fully sensitive neighbours are in S_1, \ldots, S_{k-1} . Then for all $i \in [k-1], j \in [m-2, m]$ we have that $v_{i,j} \in G$. Then, if $v_j \notin G$, it would have

sensitivity k+1 $(k-1 \text{ in } v_{i,j}, 2 \text{ in } S_j \text{ by Lemma 2})$. But if $v_j \in G$ for all $j \in [m-2, m]$, v would have sensitivity k+2. Therefore no such v exists.

Now note that $\sum_{i=1}^{m-3} R(F_i, S_i) = k - \frac{5}{2}$. Therefore at least half the vertices of S_0 have k-2 neighbours in G in light cubes. Examine one such vertex v. W.l.o.g. assume that v_1, \ldots, v_{k-2} are in G. We will now show that at most one of v_{m-2}, v_{m-1}, v_m is in G. Assume that on the contrary 2 of them are (if all 3 were, the sensitivity of v would be k+1). W.l.o.g. let $v_{m-2} \notin G$, $v_{m-1}, v_m \in G$. As $v_{i,m-2} \in G$ for $i \in [k-2]$ and v_{m-2} has sensitivity 2 in S_{m-2} by Lemma 2, we have that v_{m-2} is fully sensitive. Therefore $v_{m-2,m-1}$ and $v_{m-2,m}$ are both not in G. Then v_{m-1} and v_m are both fully sensitive and $v_{k-1,m-1}, v_{k-1,m} \in G$. But $v_{i,k-1} \in G$ for $i \in [k-2]$ and v_{k-1} has sensitivity 1 in S_{k-1} , giving it a total sensitivity of k+1, a contradiction.

As $R(G, S_i) \geq \frac{3}{4}$ for $i \in [m-2, m]$, we have that $\sum_{i=m-2}^{m} R(G, S_i) \geq \frac{9}{4}$. However, by the previous paragraph, $\frac{1}{2}$ of the vertices in S_0 have at most one neighbour in a heavy cube. Denote by G' the vertices in G neighbouring the other half of S_0 . We now have that $\sum_{i=m-2}^{m} R(G', S_i) \geq \frac{7}{4}$. But then by the pigeonhole principle at least one vertex in S_0 would need to have 4 neighbours in G in heavy cubes, which is impossible with only 3 heavy cubes.