Automated Complexity Analysis Based on the Dependency Pair Method^{*}

Nao Hirokawa

School of Information Science, Japan Advanced Institute of Science and Technology, Japan, hirokawa@jaist.ac.jp

> Georg Moser Institute of Computer Science, University of Innsbruck, Austria georg.moser@uibk.ac.at

> > June 2011

This article is concerned with automated complexity analysis of term rewrite systems. Since these systems underlie much of declarative programming, time complexity of functions defined by rewrite systems is of particular interest. Among other results, we present a variant of the dependency pair method for analysing runtime complexities of term rewrite systems automatically. The established results significantly extent previously known techniques: we give examples of rewrite systems subject to our methods that could previously not been analysed automatically. Furthermore, the techniques have been implemented in the Tyrolean Complexity Tool. We provide ample numerical data for assessing the viability of the method.

Key words: Term rewriting, Termination, Complexity Analysis, Automation, Dependency Pair Method

^{*} This research is partly supported by FWF (Austrian Science Fund) project P20133, the Grant-in-Aid for Young Scientists Nos. 20800022 and 22700009 of the Japan Society for the Promotion of Science, and Leading Project e-Society (MEXT of Japan), and STARC.

Contents

1	Introduction	2
2	Preliminaries 2.1 Rewriting 2.2 Matrix Interpretations	3 3 5
3	Runtime Complexity	6
4	Usable Replacement Maps	8
5	Weak Dependency Pairs	12
6	The Weight Gap Principle	17
7	Weak Dependency Graphs	23
8	Experiments	31
9	Conclusion	33

1 Introduction

This article is concerned with automated complexity analysis of term rewrite systems (TRSs for short). Since these systems underlie much of declarative programming, time complexity of functions defined by TRSs is of particular interest.

Several notions to assess the complexity of a terminating TRS have been proposed in the literature, compare [1, 2, 3, 4]. The conceptually simplest one was suggested by Hofbauer and Lautemann in [2]: the complexity of a given TRS is measured as the maximal length of derivation sequences. More precisely, the *derivational complexity function* with respect to a terminating TRS relates the maximal derivation height to the size of the initial term. However, when analysing complexity of a function, it is natural to refine derivational complexity so that only terms whose arguments are constructor terms are employed. Conclusively the runtime complexity function with respect to a TRS relates the length of the longest derivation sequence to the size of the initial term, where the arguments are supposed to be in normal form. This terminology was suggested in [4]. A related notion has been studied in [1], where it is augmented by an *average case* analysis. Finally [3] studies the complexity of the functions *computed* by a given TRS. This latter notion is extensively studied within implicit computational complexity theory (ICC for short), see [5] for an overview. A conceptual difference from runtime complexity is that polynomial computability addresses the number of steps by means of (deterministic) Turing machines, while runtime complexity measures the number of rewrite steps which is closely related to operational semantics of programs. For instance, a statement like a quadratic complexity of sort algorithm is in the latter sense.

This article presents methods for (over-)estimating runtime complexity automatically. We establish the following results:

- 1) We extend the applicability of direct techniques for complexity results by showing how the monotonicity constraints can be significantly weakened through the employ of *usable replacement maps*.
- 2) We revisit the *dependency pair method* in the context of complexity analysis. The dependency pair method is originally developed for proving termination [6], and known as one of the most successful methods in automated termination analysis.
- 3) We introduce the *weight gap principle* which allows the estimation of the complexity of a TRS in a modular way.
- 4) We revisit the dependency graph analysis of the dependency pair method in the context of complexity analysis. For that we introduce a suitable notion of *path analysis* that allows to modularise complexity analysis further.

Note that while we have taken seminal ideas from termination analysis as starting points, often the underlying principles are crucially different from those used in termination analysis.

A preliminary version of this article appeared in [4, 7]. Apart from the correction of some shortcomings, we extend our earlier work in the following way: First, all results on usable replacement maps are new (see Section 4). Second, the side condition for the weight gap principle [4, Theorem 24] is corrected in Section 6. Thirdly, the weight gap principle is extended by exploiting the initial term conditions and is generalised by means of matrix interpretations (see Section 6). Finally, the applicability of the path analysis is strengthened in comparison to the conference version [7] (see Section 7).

The remainder of this article is organised as follows. In the next section we recall basic notions. We define runtime complexity and a subclass of matrix interpretations for its analysis in Section 3. In Section 4 we relate context-sensitive rewriting to runtime complexity. In the next sections several ingredients in the dependency pair method are recapitulated for complexity analysis: dependency pairs and usable rules (Section 5), reduction pairs via the weight gap principle (Section 6), and dependency graphs (Section 7). In order to access viability of the presented techniques all techniques have been implemented in the *Tyrolean Complexity Tool*¹ (T_CT for short) and its empirical data is provided in Section 8. Finally we conclude the article by mentioning related works in Section 9.

2 Preliminaries

We assume familiarity with term rewriting [8, 9] but briefly review basic concepts and notations from term rewriting, relative rewriting, and context-sensitive rewriting. Moreover, we recall matrix interpretations.

2.1 Rewriting

Let \mathcal{V} denote a countably infinite set of variables and \mathcal{F} a signature, such that \mathcal{F} contains at least one constant. The set of terms over \mathcal{F} and \mathcal{V} is denoted by $\mathcal{T}(\mathcal{F}, \mathcal{V})$. The root symbol of a term t, denoted as $\operatorname{root}(t)$, is either t itself, if $t \in \mathcal{V}$, or the symbol f, if $t = f(t_1, \ldots, t_n)$. The set of position $\mathcal{P}os(t)$ of a term t is defined as usual. We write $\mathcal{P}os_{\mathcal{G}}(t) \subseteq \mathcal{P}os(t)$ for

¹ http://cl-informatik.uibk.ac.at/software/tct/.

the set of positions of subterms, whose root symbol is contained in $\mathcal{G} \subseteq \mathcal{F}$. The subterm of t at position p is denoted as $t|_p$, and $t[u]_p$ denotes the term that is obtained from t by replacing the subterm at p by u. The subterm relation is denoted as \leq . $\mathcal{V}ar(t)$ denotes the set of variables occurring in a term t. The *size* |t| of a term is defined as the number of symbols in t:

$$|t| := \begin{cases} 1 & \text{if } t \text{ is a variable }, \\ 1 + \sum_{1 \leq i \leq n} |t_i| & \text{if } t = f(t_1, \dots, t_n) \;. \end{cases}$$

A term rewrite system (TRS) \mathcal{R} over $\mathcal{T}(\mathcal{F}, \mathcal{V})$ is a finite set of rewrite rules $l \to r$, such that $l \notin \mathcal{V}$ and $\mathcal{V}ar(l) \supseteq \mathcal{V}ar(r)$. The smallest rewrite relation that contains \mathcal{R} is denoted by $\rightarrow_{\mathcal{R}}$. The transitive closure of $\rightarrow_{\mathcal{R}}$ is denoted by $\rightarrow_{\mathcal{R}}^+$, and its transitive and reflexive closure by $\rightarrow_{\mathcal{R}}^*$. We simply write \rightarrow for $\rightarrow_{\mathcal{R}}$ if \mathcal{R} is clear from context. Let s and t be terms. If exactly n steps are performed to rewrite s to t we write $s \to^n t$. Sometimes a derivation $s = s_0 \rightarrow s_1 \rightarrow \cdots \rightarrow s_n = t$ is denoted as $A: s \rightarrow^* t$ and its length n is referred to as |A|. A term $s \in \mathcal{T}(\mathcal{F}, \mathcal{V})$ is called a normal form if there is no $t \in \mathcal{T}(\mathcal{F}, \mathcal{V})$ such that $s \to t$. With $NF(\mathcal{R})$ we denote the set of all normal forms of a term rewrite system \mathcal{R} . The *innermost* rewrite relation $\xrightarrow{i}_{\mathcal{R}}$ of a TRS \mathcal{R} is defined on terms as follows: $s \xrightarrow{i}_{\mathcal{R}} t$ if there exist a rewrite rule $l \to r \in \mathcal{R}$, a context C, and a substitution σ such that $s = C[l\sigma], t = C[r\sigma]$. and all proper subterms of $l\sigma$ are normal forms of \mathcal{R} . Defined symbols of \mathcal{R} are symbols appearing at root in left-hand sides of \mathcal{R} . The set of defined function symbols is denoted as \mathcal{D} , while the constructor symbols $\mathcal{F} \setminus \mathcal{D}$ are collected in \mathcal{C} . We call a term $t = f(t_1, \ldots, t_n)$ *basic* or constructor based if $f \in \mathcal{D}$ and $t_i \in \mathcal{T}(\mathcal{C}, \mathcal{V})$ for all $1 \leq i \leq n$. The set of all basic terms are denoted by $\mathcal{T}_{\mathbf{b}}$. A TRS \mathcal{R} is called *duplicating* if there exists a rule $l \to r \in \mathcal{R}$ such that a variable occurs more often in r than in l. We call a TRS (innermost) terminating if no infinite (innermost) rewrite sequence exists.

We recall the notion of relative rewriting, cf. [10, 9]. Let \mathcal{R} and \mathcal{S} be TRSs. The relative TRS \mathcal{R}/\mathcal{S} is the pair $(\mathcal{R}, \mathcal{S})$. We define $s \to_{\mathcal{R}/\mathcal{S}} t := s \to_{\mathcal{S}}^* \cdot \to_{\mathcal{R}} \cdot \to_{\mathcal{S}}^* t$ and we call $\to_{\mathcal{R}/\mathcal{S}}$ the relative rewrite relation of \mathcal{R} over \mathcal{S} . Note that $\to_{\mathcal{R}/\mathcal{S}} = \to_{\mathcal{R}}$, if $\mathcal{S} = \emptyset$. \mathcal{R}/\mathcal{S} is called terminating if $\to_{\mathcal{R}/\mathcal{S}}$ is well-founded. In order to generalise the innermost rewriting relation to relative rewriting, we introduce the slightly technical construction of the restricted rewrite relation, compare [11]. The restricted rewrite relation $\xrightarrow{\mathcal{Q}}_{\mathcal{R}}$ is the restriction of $\to_{\mathcal{R}}$ where all arguments of the redex are in normal form with respect to the TRS \mathcal{Q} . We define the innermost relative rewriting relation (denoted as $\xrightarrow{i}_{\mathcal{R}/\mathcal{S}}$) as follows:

$$\xrightarrow{i}_{\mathcal{R}/\mathcal{S}} := \xrightarrow{\mathcal{R}\cup\mathcal{S}} \xrightarrow{*}_{\mathcal{S}} \cdot \xrightarrow{\mathcal{R}\cup\mathcal{S}}_{\mathcal{R}} \cdot \xrightarrow{\mathcal{R}\cup\mathcal{S}} \xrightarrow{*}_{\mathcal{S}},$$

We briefly recall context-sensitive rewriting. A replacement map μ is a function with $\mu(f) \subseteq \{1, \ldots, n\}$ for all *n*-ary functions with $n \ge 1$. The set $\mathcal{P}os_{\mu}(t)$ of μ -replacing positions in t is defined as follows:

$$\mathcal{P}os_{\mu}(t) = \begin{cases} \{\epsilon\} & \text{if } t \text{ is a variable }, \\ \{\epsilon\} \cup \{ip \mid i \in \mu(f) \text{ and } p \in \mathcal{P}os_{\mu}(t_i)\} & \text{if } t = f(t_1, \dots, t_n) \end{cases}$$

A μ -step $s \xrightarrow{\mu} t$ is a rewrite step $s \to t$ whose rewrite position is in $\mathcal{P}os_{\mu}(s)$. The set of all non- μ -replacing positions in t is denoted by $\overline{\mathcal{P}os}_{\mu}(t)$; namely, $\overline{\mathcal{P}os}_{\mu}(t) := \mathcal{P}os(t) \setminus \mathcal{P}os_{\mu}(t)$.

2.2 Matrix Interpretations

One of the most powerful and popular techniques for analysing derivational complexities is use of orders induced from matrix interpretations [12]. In order to define it first we define (weakly) monotone algebras.

A proper order is a transitive and irreflexive relation and a preorder (or quasi-order) is a transitive and reflexive relation. A proper order \succ is well-founded if there is no infinite decreasing sequence $t_1 \succ t_2 \succ t_3 \cdots$. We say a proper order \succ and a TRS \mathcal{R} are compatible if $\mathcal{R} \subseteq \succ$.

An \mathcal{F} -algebra \mathcal{A} consists of a carrier set A and a collection of interpretations $f_{\mathcal{A}}$ for each function symbol in \mathcal{F} . By $[\alpha]_{\mathcal{A}}(\cdot)$ we denote the usual evaluation function of \mathcal{A} according to an assignment α which maps variables to values in A. A monotone \mathcal{F} -algebra is a pair (\mathcal{A}, \succ) where \mathcal{A} is an \mathcal{F} -algebra and \succ is a proper order such that for every function symbol $f \in \mathcal{F}$, $f_{\mathcal{A}}$ is strictly monotone in all coordinates with respect to \succ . A weakly monotone \mathcal{F} -algebra $(\mathcal{A}, \succcurlyeq)$ is defined similarly, but for every function symbol $f \in \mathcal{F}$, it suffices that $f_{\mathcal{A}}$ is weakly monotone in all coordinates (with respect to the quasi-order \succcurlyeq). A monotone \mathcal{F} -algebra (\mathcal{A}, \succ) is called *well-founded* if \succ is well-founded. We write WMA instead of well-founded monotone algebra.

Any (weakly) monotone \mathcal{F} -algebra (\mathcal{A}, R) induces a binary relation $R_{\mathcal{A}}$ on terms: define $s \ R_{\mathcal{A}} \ t$ if $[\alpha]_{\mathcal{A}}(s) \ R \ [\alpha]_{\mathcal{A}}(t)$ for all assignments α . Clearly if R is a proper order (quasiorder), then $R_{\mathcal{A}}$ is a proper order (quasi-order) on terms and if R is a well-founded, then $R_{\mathcal{A}}$ is well-founded on terms. We say \mathcal{A} is *compatible* with a TRS \mathcal{R} if $\mathcal{R} \subseteq R_{\mathcal{A}}$. Let $\succeq_{\mathcal{A}}$ denote the quasi-order induced by a weakly monotone algebra (\mathcal{A}, \succeq) , then $=_{\mathcal{A}}$ denotes the equivalence (on terms) induced by $\succeq_{\mathcal{A}}$. Let μ denote a replacement map. Then we call a well-founded algebra (\mathcal{A}, \succ) μ -monotone if for every function symbol $f \in \mathcal{F}$, $f_{\mathcal{A}}$ is strictly monotone on $\mu(f)$, i.e., $f_{\mathcal{A}}$ is strictly monotone with respect to every argument position in $\mu(f)$. Similarly a (strict) relation R is called μ -monotone if (strictly) monotone on $\mu(f)$ for all $f \in \mathcal{F}$. Let \mathcal{R} be a TRS compatible with a μ -monotone relation R. Then clearly any μ -step $s \xrightarrow{\mu} t$ implies $s \ R t$.

We recall the concept of *matrix interpretations* on natural numbers (see [12] but compare also [13]). Let \mathcal{F} denote a signature. We fix a dimension $d \in \mathbb{N}$ and use the set \mathbb{N}^d as the carrier of an algebra \mathcal{A} , together with the following extension of the natural order > on \mathbb{N} :

$$(x_1, x_2, \dots, x_d) > (y_1, y_2, \dots, y_d) : \iff x_1 > y_1 \land x_2 \ge y_2 \land \dots \land x_d \ge y_d .$$

Let μ be a replacement map. For each *n*-ary function symbol f, we choose as an interpretation a linear function of the following shape:

$$f_{\mathcal{A}}: (\vec{v}_1, \dots, \vec{v}_n) \mapsto F_1 \vec{v}_1 + \dots + F_n \vec{v}_n + \vec{f},$$

where $\vec{v}_1, \ldots, \vec{v}_n$ are (column) vectors of variables, F_1, \ldots, F_n are matrices (each of size $d \times d$), and \vec{f} is a vector over \mathbb{N} . Moreover, suppose for any $i \in \mu(f)$ the top left entry $(F_i)_{1,1}$ is positive. Then it is easy to see that the algebra \mathcal{A} forms a μ -monotone WMA. Let \mathcal{A} be a matrix interpretation, let α_0 denotes the assignment mapping any variable to $\vec{0}$, i.e., $\alpha_0(x) = \vec{0}$ for all $x \in \mathcal{V}$, and let t be a term. In the following we write $[t], [t]_j$ as an abbreviation for $[\alpha_0]_{\mathcal{A}}(t)$, or $([\alpha_0]_{\mathcal{A}}(t))_j$ $(1 \leq j \leq d)$, respectively, if the algebra \mathcal{A} is clear from the context.

3 Runtime Complexity

In this section we formalise runtime complexity and then define a subclass of matrix interpretations that give polynomial upper-bounds.

The derivation height of a term s with respect to a well-founded, finitely branching relation \rightarrow is defined as: $dh(s, \rightarrow) = \max\{n \mid \exists t \ s \rightarrow^n t\}$. Let \mathcal{R} be a TRS and T be a set of terms. The complexity function with respect to a relation \rightarrow on T is defined as follows:

$$\operatorname{comp}(n, T, \rightarrow) = \max\{\operatorname{dh}(t, \rightarrow) \mid t \in T \text{ and } |t| \leq n\}.$$

In particular we are interested in the (innermost) complexity with respect to $\rightarrow_{\mathcal{R}} (\stackrel{i}{\rightarrow}_{\mathcal{R}})$ on the set \mathcal{T}_{b} of all *basic* terms.

Definition 3.1. Let \mathcal{R} be a TRS. We define the *runtime complexity function* $\mathsf{rc}_{\mathcal{R}}(n)$, the *innermost runtime complexity function* $\mathsf{rc}_{\mathcal{R}}^{i}(n)$, and the *derivational complexity function* $\mathsf{dc}_{\mathcal{R}}(n)$ as $\mathsf{comp}(n, \mathcal{T}_{\mathsf{b}}, \rightarrow_{\mathcal{R}})$, $\mathsf{comp}(n, \mathcal{T}_{\mathsf{b}}, \stackrel{i}{\rightarrow}_{\mathcal{R}})$, and $\mathsf{comp}(n, \mathcal{T}(\mathcal{F}, \mathcal{V}), \rightarrow_{\mathcal{R}})$, respectively.

Note that the above complexity functions need not be defined, as the rewrite relation $\rightarrow_{\mathcal{R}}$ is not always well-founded *and* finitely branching. We sometimes say the (innermost) runtime complexity of \mathcal{R} is *linear*, *quadratic*, or *polynomial* if there exists a (linear, quadratic) polynomial p(n) such that $\mathsf{rc}_{\mathcal{R}}^{(i)}(n) \leq p(n)$ for sufficiently large n. The (innermost) runtime complexity of \mathcal{R} is called *exponential* if there exist constants c, d with $c, d \geq 2$ such that $\mathsf{c}_{\mathcal{R}}^{(i)}(n) \leq \mathsf{d}^n$ for sufficiently large n.

The next example illustrates a difference between derivational complexity and runtime complexity.

Example 3.2. Consider the following TRS \mathcal{R}_{div}^2

1:
$$x - 0 \rightarrow x$$
 3: $0 \div \mathbf{s}(y) \rightarrow 0$
2: $\mathbf{s}(x) - \mathbf{s}(y) \rightarrow x - y$ 4: $\mathbf{s}(x) \div \mathbf{s}(y) \rightarrow \mathbf{s}((x - y) \div \mathbf{s}(y))$

Although the functions computed by \mathcal{R}_{div} are obviously feasible this is not reflected in the derivational complexity of \mathcal{R}_{div} . Consider rule 4, which we abbreviate as $C[x] \to D[x, x]$. Since the maximal derivation height starting with $C^n[x]$ equals 2^{n-1} for all n > 0, \mathcal{R}_{div} admits (at least) exponential derivational complexity. In general any duplicating TRS admits (at least) exponential derivational complexity.

In general it is not possible to bound $dc_{\mathcal{R}}$ polynomially in $rc_{\mathcal{R}}$, as witnessed by Example 3.2 and the observation that the runtime complexity of \mathcal{R} is linear (see Example 4.10, below). We will use Example 3.2 as our running example.

Below we define classes of orders whose compatibility with a TRS \mathcal{R} bounds its runtime complexity from the above. Note that $dh(t, \succ)$ is undefined, if the relation \succ is not well-founded or not finitely branching. In fact compatibility of a constructor TRS with the polynomial path order $>_{pop*}$ ([15]) induces polynomial innermost runtime complexity, whereas $f(x) >_{pop*} \cdots >_{pop*} g^2(x) >_{pop*} g(x) >_{pop*} x$ holds when precedence f > g is used. Hence $dh(t, >_{pop*})$ is undefined, while the order $>_{pop*}$ can be employed in complexity analysis.

 $^{^{2}}$ This is Example 3.1 in Arts and Giesl's collection of TRSs [14].

Definition 3.3. Let R be a binary relation over terms, let \succ be a proper order on terms, and let G denote a mapping associating a term with a natural number. Then \succ is G-collapsible on R if G(s) > G(t), whenever s R t and $s \succ t$ holds. An order \succ is collapsible (on R), if there is a mapping G such that \succ is G-collapsible (on R).

Lemma 3.4. Let R be a finitely branching and well-founded relation. Further, let \succ be a G-collapsible order with $R \subseteq \succ$. Then $dh(t, R) \leq G(t)$ holds for all terms t.

The alert reader will have noticed that any proper order \succ is collapsible on a finitely branching and well-founded relation R: simply set G(t) := dh(t, R). However, this observation is of limited use if we wish to bound the derivation height of t in independence of R.

If a TRS \mathcal{R} and a μ -monotone matrix interpretation \mathcal{A} are compatible, $\mathsf{G}(t)$ can be given by $[t]_1$. In order to estimate derivational or runtime complexity, one needs to associate $[t]_1$ to |t|. For this sake we define degrees of matrix interpretations.

Definition 3.5. A matrix interpretation is of *(basic)* degree d if there is a constant c such that $[t]_i \leq c \cdot |t|^d$ for all (basic) terms t and i, respectively.

An upper triangular complexity matrix is a matrix M in $\mathbb{N}^{d \times d}$ such that we have $M_{j,k} = 0$ for all $1 \leq k < j \leq d$, and $M_{j,j} \leq 1$ for all $1 \leq j \leq d$. We say that a WMA \mathcal{A} is a triangular matrix interpretation (TMI for short) if \mathcal{A} is a matrix interpretation (over \mathbb{N}) and all matrices employed are of upper triangular complexity form. It is easy to define triangular matrix interpretations, such that an algebra \mathcal{A} based on such an interpretation, forms a wellfounded weakly monotone algebra. To simplify notation we will also refer to \mathcal{A} as a TMI, if no confusion can arise from this. A TMI \mathcal{A} of dimension 1, that is a linear polynomial, is called a strongly linear interpretation (SLI for short) if all interpretation functions $f_{\mathcal{A}}$ are strongly linear. Here a polynomial $P(x_1, \ldots, x_n)$ is strong linear if $P(x_1, \ldots, x_n) = x_1 + \cdots + x_n + c$.

Lemma 3.6. Let \mathcal{A} be a TMI and let M denote the component-wise maximum of all matrices occurring in \mathcal{A} . Further, let d denote the number of ones occurring along the diagonal of M. Then for all $1 \leq i, j \leq d$ we have $(M^n)_{i,j} = O(n^{d-1})$.

Proof. The lemma is a direct consequence of Lemma 4 in [16] together with the observation that for any triangular complexity matrix, the diagonal entries denote the multiset of eigenvalues. \Box

Lemma 3.7. Let \mathcal{A} and d be defined as in Lemma 3.6. Then \mathcal{A} is of degree d.

Proof. For any (triangular) matrix interpretation \mathcal{A} , there exist vectors \vec{v}_i and a vector \vec{w} such that the evaluation [t] of t can be written as follows:

$$[t] = \sum_{i=1}^{\ell} \vec{v}_i + \vec{w} ,$$

where each vector \vec{v}_i is the product of those matrices employed in the interpretation of function symbols in \mathcal{A} and a vector representing the constant part of a function interpretation. It is not difficult to see that there is a one-to-one correspondence between the number of vectors $\vec{v}_1, \ldots, \vec{v}_\ell$ and the number of subterms of t and thus $\ell = |t|$. Moreover for each \vec{v}_i the number of products is less than the depth of t and thus bounded by |t|. In addition, due to Lemma 3.6 the entries of the vectors \vec{v}_i and \vec{w} are bounded by a polynomial of degree at most d-1. Thus for all $1 \leq j \leq d$, there exists $k \leq d$ such that $([t])_j = O(|t|^k)$.

Theorem 3.8. [16, Theorem 9], [17] Let \mathcal{A} and d be defined as in Lemma 3.6. Then, $\succ_{\mathcal{A}}$ is $O(n^d)$ -collapsible.

Proof. The theorem is a direct consequence of Lemmas 3.6 and 3.7.

In order to cope with runtime complexity, a similar idea to restricted polynomial interpretations (see [18]) can be integrated to triangle matrix interpretations. We call \mathcal{A} a *restricted matrix interpretation* (*RMI* for short) if \mathcal{A} is a matrix interpretation, but for each constructor symbol $f \in \mathcal{F}$, the interpretation $f_{\mathcal{A}}$ of f employs upper triangular complexity matrices, only. The next theorem is a direct consequence of the definitions in conjunction with Lemma 3.7.

Theorem 3.9. Let \mathcal{A} be an RMI and let t be a basic term. Further, let M denote the component-wise maximum of all matrices used for the interpretation of constructor symbol, and let d denote the number of ones occurring along the diagonal of M. Then \mathcal{A} is of basic degree d. Furthermore, if M is the unit matrix then \mathcal{A} is of basic degree 1.

4 Usable Replacement Maps

Unfortunately, there is no RMI compatible with the TRS of our running example. The reason is that the monotonicity requirement of TMI is too severe for complexity analysis. Inspired by the idea of Fernández [19], we show how context-sensitive rewriting is used in complexity analysis. Here we briefly explain our idea. Let **n** denote the numeral $s^n(0)$. Consider the derivation from $\mathbf{4} \div \mathbf{2}$:

$$\underline{4\div 2} \to \mathsf{s}((3-1)\div 2) \to \mathsf{s}((\underline{2-0})\div 2) \to \mathsf{s}(\underline{2\div 2}) \to \cdots$$

where redexes are underlined. Observe that e.g. any second argument of \div is never rewritten. More precisely, any derivation from a basic term consists of only μ -steps with the replacement map μ : $\mu(\mathbf{s}) = \mu(\div) = \{1\}$ and $\mu(-) = \emptyset$.

We present a simple method based on a variant of ICAP in [20] to estimate a suitable replacement map. Let μ be a replacement map. Clearly the function μ is representable as set of ordered pairs (f, i). Below we often confuse the notation of μ as a function or as a set. Recall that $\mathcal{P}os_{\mu}(t)$ denotes the set of μ -replacing positions in t and $\overline{\mathcal{P}os}_{\mu}(t) =$ $\mathcal{P}os(t) \setminus \mathcal{P}os_{\mu}(t)$. Further, a term t is a μ -replacing term with respect to a TRS \mathcal{R} if $t|_{p} \notin \mathsf{NF}(\mathcal{R})$ implies that $p \in Pos_{\mu}(t)$. The set of all μ -replacing terms is denoted by $\mathcal{T}(\mu)$. In the following \mathcal{R} will always denote a TRS.

Definition 4.1. Let \mathcal{R} be a TRS and let μ be a replacement map. We defined the operator $\Upsilon^{\mathcal{R}}$ as follows:

$$\Upsilon^{\mathcal{R}}(\mu) := \{ (f,i) \mid l \to C[f(r_1,\ldots,r_n)] \in \mathcal{R} \text{ and } \mathsf{CAP}^l_{\mu}(r_i) \neq r_i \} .$$

Here $\mathsf{CAP}^{s}_{\mu}(t)$ is inductively defined on t as follows:

$$\mathsf{CAP}^{s}_{\mu}(t) = \begin{cases} t & t = s|_{p} \text{ for some } p \in \overline{\mathcal{P}\mathsf{os}}_{\mu}(s) ,\\ u & \text{if } t = f(t_{1}, \dots, t_{n}) \text{ and } u \text{ and } l \text{ unify for no } l \to r \in \mathcal{R} ,\\ y & \text{otherwise }, \end{cases}$$

where, $u = f(\mathsf{CAP}^s_{\mu}(t_1), \dots, \mathsf{CAP}^s_{\mu}(t_n))$, y is a fresh variable, and $\mathcal{V}ar(l) \cap \mathcal{V}ar(u) = \emptyset$ is assumed.

We define the *innermost usable replacement map* $\mu_i^{\mathcal{R}}$ as follows $\mu_i^{\mathcal{R}} := \Upsilon^{\mathcal{R}}(\emptyset)$ and let the *usable replacement map* $\mu_f^{\mathcal{R}}$ denote the least fixed point of $\Upsilon^{\mathcal{R}}$. The existence of $\Upsilon^{\mathcal{R}}$ follows from the monotonicity of $\Upsilon^{\mathcal{R}}$. If \mathcal{R} is clear from context, we simple write μ_i , μ_f , and Υ , respectively. Usable replacement maps satisfy a desired property for runtime complexity analysis. In order to see it several preliminary lemmas are necessary.

First we take a look at $\mathsf{CAP}^s_{\mu}(t)$. Suppose $s \in \mathcal{T}(\mu)$: observe that the function $\mathsf{CAP}^s_{\mu}(t)$ replaces a subterm u of t by a fresh variable if $u\sigma$ is a redex for some $s\sigma \in \mathcal{T}(\mu)$. This is exemplified below.

Example 4.2. Consider the TRS \mathcal{R}_{div} . Let $l \to r$ be rule 4, namely, $l = \mathbf{s}(x) \div \mathbf{s}(y)$ and $r = \mathbf{s}((x - y) \div \mathbf{s}(y))$. Suppose $\mu(f) = \emptyset$ for all functions f and let w and z be fresh variables. The next table summarises $\mathsf{CAP}^l_{\mu}(t)$ for each proper subterm t in r. To see the computation process, we also indicate the term u in Definition 4.1.

t	x	y	x - y	s(y)	$(x-y) \div s(y)$
u	—	_	x - y	s(y)	$w \div s(y)$
$CAP^l_\mu(t)$	x	y	w	s(y)	z

By underlining proper subterms t in r such that $\mathsf{CAP}^l_{\mu}(t) \neq t$, we have

$$\mathsf{s}((x-y) \div \mathsf{s}(y))$$

which indicates $(s, 1), (\div, 1) \in \Upsilon(\mu)$.

The next lemma states a role of $\mathsf{CAP}^s_{\mu}(t)$.

Lemma 4.3. If $s\sigma \in \mathcal{T}(\mu)$ and $\mathsf{CAP}^s_{\mu}(t) = t$ then $t\sigma \in \mathsf{NF}(\mathcal{R})$.

Proof. We use induction on t. Suppose $s\sigma \in \mathcal{T}(\mu)$ and $\mathsf{CAP}^s_{\mu}(t) = t$. If $t = s|_p$ for some $p \in \overline{\mathcal{Pos}}_{\mu}(s)$ then $t\sigma = (s\sigma)|_p \in \mathsf{NF}$ follows by definition of $\mathcal{T}(\mu)$.

We can assume that $t = f(t_1, \ldots, t_n)$. Assume otherwise that $t = x \in \mathcal{V}$, then $\mathsf{CAP}^s_{\mu}(x) = x$ entails that $x\sigma$ occurs at a non- μ -replacing position in $s\sigma$. Hence $x\sigma \in \mathsf{NF}$ follows from $s\sigma \in \mathcal{T}(\mu)$. Moreover, by assumption we have:

- 1) $\mathsf{CAP}^{s}_{\mu}(t_{i}) = t_{i}$ for each i, and
- 2) there is no rule $l \to r \in \mathcal{R}$ such that t and l unify.

Due to 2) $l\sigma$ is not reducible at the root, and the induction hypothesis yields $t_i\sigma \in \mathsf{NF}$ because of 1). Therefore, we obtain $t\sigma \in \mathsf{NF}$.

For a smooth inductive proof of our key lemma we prepare a characterisation of the set of μ -replacing terms $\mathcal{T}(\mu)$.

Definition 4.4. The set $\{(f,i) \mid f(t_1,\ldots,t_n) \leq t \text{ and } t_i \notin \mathsf{NF}(\mathcal{R})\}$ is denoted by v(t).

Lemma 4.5. $\mathcal{T}(\mu) = \{t \mid v(t) \subseteq \mu\}.$

Proof. The inclusion from left to right essentially follows from the definitions. Let $t \in \mathcal{T}(\mu)$ and let $(f, i) \in v(t)$. We show $(f, i) \in \mu$. By Definition 4.4 there is a position $p \in \mathcal{P}os(t)$ with $t|_p = f(t_1, \ldots, t_n)$ and $t|_{pi} \notin \mathsf{NF}$. Thus $pi \in \mathcal{P}os_{\mu}(t)$ and $i \in \mathcal{P}os_{\mu}(t|_p)$. Hence $(f, i) \in \mu$ is concluded.

Next we consider the reverse direction $\{t \mid v(t) \subseteq \mu\} \subseteq \mathcal{T}(\mu)$. Let t be a minimal term such that $v(t) \subseteq \mu$ and $t \notin \mathcal{T}(\mu)$. One can write $t = f(t_1, \ldots, t_n)$. Then, there exists a position $p \in \overline{\mathcal{Pos}}_{\mu}(t)$ such that $t|_p \notin \mathsf{NF}$. Because $\epsilon \notin \overline{\mathcal{Pos}}_{\mu}(t)$ holds in general, p is of the form iq with $i \in \mathbb{N}$. As $iq \in \overline{\mathcal{Pos}}_{\mu}(t)$ one of $(f, i) \notin \mu$ or $q \in \overline{\mathcal{Pos}}_{\mu}(t|_i)$ must hold. As t is minimal and $t|_{iq} \notin \mathsf{NF}$ implies that $t|_i \notin \mathsf{NF}$, we have $(f, i) \notin \mu$. However, by Definition 4.4, $(f, i) \in v(t) \subseteq \mu$. Contradiction.

The next lemma about the operator Υ is a key for the main theorem. Note that every subterm of a μ -replacing term is a μ -replacing term.

Lemma 4.6. If $l \to r \in \mathcal{R}$ and $l\sigma \in \mathcal{T}(\mu)$ then $r\sigma \in \mathcal{T}(\mu \cup \Upsilon(\mu))$.

Proof. Let $l \to r \in \mathcal{R}$ and suppose $l\sigma \in \mathcal{T}(\mu)$. By Lemma 4.5 we have

$$\mathcal{T}(\mu) = \{t \mid v(t) \subseteq \mu\} \qquad \mathcal{T}(\mu \cup \Upsilon(\mu)) = \{t \mid v(t) \subseteq \mu \cup \Upsilon(\mu)\}.$$

Hence it is sufficient to show $v(r\sigma) \subseteq \mu \cup \Upsilon(\mu)$. Let $(f,i) \in v(r\sigma)$. There is $p \in \mathcal{P}os(r\sigma)$ with $r\sigma|_p = f(t_1, \ldots, t_n)$ and $t_i \notin \mathsf{NF}$. If p is below some variable position of r, $r\sigma|_p$ is a subterm of $l\sigma$, and thus $v(r\sigma|_p) \subseteq v(l\sigma) \subseteq \mu$. Otherwise, p is a non-variable position of r. We may write $r|_p = f(r_1, \ldots, r_n)$ and $r_i\sigma = t_i \notin \mathsf{NF}$. Due to Lemma 4.3 we obtain $\mathsf{CAP}^l_\mu(r_i) \neq r_i$. Therefore, $(f, i) \in \Upsilon(\mu)$.

Remark that if $s, t \in \mathcal{T}(\mu)$ and $p \in \mathcal{P}os_{\mu}(s)$ then $s[t]_p \in \mathcal{T}(\mu)$.

Lemma 4.7. The following implications hold.

- 1) If $s \in \mathcal{T}(\mu_i)$ and $s \xrightarrow{i} t$ then $t \in \mathcal{T}(\mu_i)$.
- 2) If $s \in \mathcal{T}(\mu_{f})$ and $s \to t$ then $t \in \mathcal{T}(\mu_{f})$.

Proof. We show property 1). Suppose $s \in \mathcal{T}(\mu_i)$ and $s \stackrel{i}{\to} t$ is a rewrite step at p. Due to the definition of innermost rewriting, we have $s|_p \in \mathcal{T}(\emptyset)$. Hence, $t|_p \in \mathcal{T}(\mu_i)$ is obtained by Lemma 4.6. Because $s \in \mathcal{T}(\mu_i)$ we have $p \in \mathcal{Pos}_{\mu_i}(s)$. Hence due to $t|_p \in \mathcal{T}(\mu_i)$ we conclude $t = s[t|_p]_p \in \mathcal{T}(\mu_i)$ due to the above remark. The proof of 2) proceeds along the same pattern and is left to the reader.

We arrive at the main result of this section.

Theorem 4.8. Let \mathcal{R} be a TRS, and let $\rightarrow^*(L)$ denote the descendants of the set of terms L. Then $\stackrel{i}{\rightarrow}^*_{\mathcal{R}}(\mathcal{T}(\emptyset)) \subseteq \mathcal{T}(\mu_i)$ and $\rightarrow^*_{\mathcal{R}}(\mathcal{T}(\emptyset)) \subseteq \mathcal{T}(\mu_f)$. *Proof.* Recall that $\to^*(L) := \{t \mid \exists s \in L \text{ such that } s \to^* t\}$. We focus on the second part of the theorem, where we have to prove that $t \in \mathcal{T}(\mu_f)$, whenever there exists $s \in \mathcal{T}(\emptyset)$ such that $s \to^*_{\mathcal{R}} t$. As $\mathcal{T}(\emptyset) \subseteq \mathcal{T}(\mu_f)$ this follows directly from Lemma 4.7.

Note that $\mathcal{T}(\emptyset)$ is the set of all argument normalised terms. Therefore, $\mathcal{T}_{b} \subseteq \mathcal{T}(\emptyset)$. The following corollary to Theorem 4.8 is immediate.

Corollary 4.9. Let \mathcal{R} be a TRS and let $\xrightarrow{\mu_i}$, $\xrightarrow{\mu_f}$ denote the μ_i -step and μ_f -step relation, respectively. Then for all terminating terms $t \in \mathcal{T}_b$ we have $dh(t, \xrightarrow{i} \mathcal{R}) \leq dh(t, \xrightarrow{\mu_i})$ and $dh(t, \rightarrow_{\mathcal{R}}) \leq dh(t, \xrightarrow{\mu_f})$.

An advantage of the use of context-sensitive rewriting is that the compatibility requirement of monotone algebra in termination or complexity analysis is relaxed to μ -monotone algebra. We illustrate its use in the next example.

Example 4.10. Recall the TRS \mathcal{R}_{div} given in Example 3.2 above. The usable argument positions are as follows:

$$\mu_{i}(-) = \varnothing \quad \mu_{i}(s) = \mu_{i}(\div) = \{1\} \qquad \mu_{f}(s) = \mu_{f}(-) = \mu_{f}(\div) = \{1\} \ .$$

Consider the 1-dimensional RMI \mathcal{A} (i.e., linear polynomial interpretations) with

$$\mathbf{0}_{\mathcal{A}} = 1 \qquad \qquad \mathbf{s}_{\mathcal{A}}(x) = x + 2 \qquad \qquad -_{\mathcal{A}}(x, y) = x + 1 \qquad \qquad \div_{\mathcal{A}}(x, y) = 3x \ .$$

which is strictly μ_i -monotone and μ_f -monotone. The rules in \mathcal{R}_{div} are interpreted and ordered as follows.

1:
$$x+1 > x$$
 3: $3 > 1$
2: $x+3 > x+2$ 4: $3x+6 > 3x+5$

Therefore, $\mathcal{R}_{div} \subseteq >_{\mathcal{A}}$ holds. By an application of Theorem 3.9 we conclude that the (innermost) runtime complexity is *linear*, which is optimal.

We cast the observations in the example into another corollary to Theorem 4.8.

Corollary 4.11. Let \mathcal{R} be a TRS and let \mathcal{A} be a d-degree μ_i -monotone (or μ_f -monotone) RMI compatible with \mathcal{R} . Then the (innermost) runtime complexity function $\mathsf{rc}_{\mathcal{R}}^{(i)}$ with respect to \mathcal{R} is bounded by a d-degree polynomial.

Proof. It suffices to consider the case for full rewriting. Let s, t be terms such that $s \to_{\mathcal{R}} t$. By the theorem, we have $s \xrightarrow{\mu_f} t$. Furthermore, by assumption $\mathcal{R} \subseteq \succ_{\mathcal{A}}$ and for any $f \in \mathcal{F}$, $f_{\mathcal{A}}$ is strictly monotone on all $\mu_f(f)$. Thus $s \succ_{\mathcal{A}} t$ follows. Finally, the corollary follows by application of Theorem 3.9.

We link Theorem 4.8 to related work by Fernández [19]. In [19] it is shown how contextsensitive rewriting is used for proving innermost termination.

Proposition 4.12 ([19]). A TRS \mathcal{R} is innermost terminating if $\xrightarrow{\mu_i}$ is terminating.

Proof. We show the contraposition. If \mathcal{R} is not innermost terminating, there is an infinite sequence $t_0 \xrightarrow{i} t_1 \xrightarrow{i} t_2 \xrightarrow{i} \cdots$, where $t_0 \in \mathcal{T}(\emptyset)$. From Theorem 4.8 and Lemma 4.7 we obtain $t_0 \xrightarrow{\mu_i} t_1 \xrightarrow{\mu_i} t_2 \xrightarrow{\mu_i} \cdots$. Hence, $\xrightarrow{\mu_i}$ is not terminating.

One might think that a similar claim holds for full termination if one uses μ_{f} . The next examples clarifies that this is not the case.

Example 4.13. Consider the famous Toyama's example \mathcal{R}

$$f(a, b, x) \rightarrow f(x, x, x)$$
 $g(x, y) \rightarrow x$ $g(x, y) \rightarrow y$.

The replacement map μ_{f} is empty. Thus, the algebra \mathcal{A} over \mathbb{N}

$$\mathsf{f}_{\mathcal{A}}(x,y,z) = \max\{x-y,0\} \qquad \mathsf{g}_{\mathcal{A}}(x,y) = x+y+1 \qquad \mathsf{a}_{\mathcal{A}} = 1 \qquad \mathsf{b}_{\mathcal{A}} = 0 \; .$$

is μ_{f} -monotone and we have $\mathcal{R} \subseteq >_{\mathcal{A}}$. However, we should not conclude termination of \mathcal{R} , because f(a, b, g(a, b)) is non-terminating.

5 Weak Dependency Pairs

In Section 4 we investigated argument positions of rewrite steps. This section is concerned about contexts surrounding rewrite steps. Recall the derivation:

$$\begin{array}{c|c} \underline{\mathbf{4} \div \mathbf{2}} & \rightarrow_{\mathcal{R}_{\text{div}}} \mathsf{s}(\boxed{(\mathbf{3} - \mathbf{1}) \div \mathbf{2}}) & \rightarrow^2_{\mathcal{R}_{\text{div}}} \mathsf{s}(\boxed{\mathbf{2} \div \mathbf{2}}) \\ & \rightarrow_{\mathcal{R}_{\text{div}}} \mathsf{s}(\mathsf{s}(\boxed{(\mathbf{1} - \mathbf{1}) \div \mathbf{2}})) & \rightarrow^2_{\mathcal{R}_{\text{div}}} \mathsf{s}(\mathsf{s}(\boxed{\mathbf{0} \div \mathbf{2}})) \\ & \rightarrow_{\mathcal{R}_{\text{div}}} \mathsf{s}(\mathsf{s}(\mathbf{0})) \ , \end{array}$$

where we boxed outermost occurrences of defined symbols. Obviously, their surrounding contexts are not rewritten. Here an idea is to simulate rewrite steps from basic terms with new rewrite rules, obtained by dropping unnecessary contexts. In termination analysis this method is known as the dependency pair method [6]. We recast its main ingredient called dependency pairs.

Let X be a set of symbols. We write $C\langle t_1, \ldots, t_n \rangle_X$ to denote $C[t_1, \ldots, t_n]$, whenever $\operatorname{root}(t_i) \in X$ for all $1 \leq i \leq n$ and C is an n-hole context containing no X-symbols. (Note that the context C may be degenerate and doesn't contain a hole \Box or it may be that C is a hole.) Then, every term t can be uniquely written in the form $C\langle t_1, \ldots, t_n \rangle_X$.

Lemma 5.1. Let t be a terminating term, and let σ be a substitution. Then $dh(t\sigma, \rightarrow_{\mathcal{R}}) = \sum_{1 \leq i \leq n} dh(t_i\sigma, \rightarrow_{\mathcal{R}})$, whenever $t = C\langle t_1, \ldots, t_n \rangle_{\mathcal{D} \cup \mathcal{V}}$.

The idea is to replace such a *n*-hole context with a fresh *n*-ary function symbol. We define the function COM as a mapping from tuples of terms to terms as follows: $COM(t_1, \ldots, t_n)$ is t_1 if n = 1, and $c(t_1, \ldots, t_n)$ otherwise. Here *c* is a fresh *n*-ary function symbol called *compound symbol*. The above lemma motivates the next definition of *weak dependency pairs*.

Definition 5.2. Let t be a term. We set $t^{\sharp} := t$ if $t \in \mathcal{V}$, and $t^{\sharp} := f^{\sharp}(t_1, \ldots, t_n)$ if $t = f(t_1, \ldots, t_n)$. Here f^{\sharp} is a new n-ary function symbol called *dependency pair symbol*. For a signature \mathcal{F} , we define $\mathcal{F}^{\sharp} = \mathcal{F} \cup \{f^{\sharp} \mid f \in \mathcal{F}\}$. Let \mathcal{R} be a TRS. If $l \to r \in \mathcal{R}$ and $r = C\langle u_1, \ldots, u_n \rangle_{\mathcal{D} \cup \mathcal{V}}$ then the rewrite rule $l^{\sharp} \to \operatorname{COM}(u_1^{\sharp}, \ldots, u_n^{\sharp})$ is called a *weak dependency pair* of \mathcal{R} . The set of all weak dependency pairs is denoted by WDP(\mathcal{R}).

While dependency pair symbols are defined with respect to $WDP(\mathcal{R})$, these symbols are not defined with respect to the original system \mathcal{R} . In the sequel defined symbols refer to the defined function symbols of \mathcal{R} .

Example 5.3 (continued from Example 3.2). The set $WDP(\mathcal{R}_{div})$ consists of the next four weak dependency pairs:

5:
$$x \to 0 \to x$$
 7: $0 \div \sharp \mathbf{s}(y) \to \mathbf{c}$
6: $\mathbf{s}(x) \to \xi(y) \to x \to y$ 8: $\mathbf{s}(x) \div \sharp \mathbf{s}(y) \to (x-y) \div \sharp \mathbf{s}(y)$

Here c denotes a fresh compound symbols of arity 0.

The derivation on page 12 corresponds to the derivation of $WDP(\mathcal{R}_{div}) \cup \mathcal{R}_{div}$:

$$\begin{array}{ll} \mathbf{4} \div^{\sharp} \mathbf{2} & \rightarrow_{\mathsf{WDP}(\mathcal{R}_{\mathsf{div}})} & (\mathbf{3} - \mathbf{1}) \div^{\sharp} \mathbf{2} & \rightarrow^{2}_{\mathcal{R}_{\mathsf{div}}} \mathbf{2} \div^{\sharp} \mathbf{2} \\ & \rightarrow_{\mathsf{WDP}(\mathcal{R}_{\mathsf{div}})} & (\mathbf{1} - \mathbf{1}) \div^{\sharp} \mathbf{2} & \rightarrow^{2}_{\mathcal{R}_{\mathsf{div}}} \mathbf{0} \div^{\sharp} \mathbf{2} \\ & \rightarrow_{\mathsf{WDP}(\mathcal{R}_{\mathsf{div}})} & \mathsf{c} \ , \end{array}$$

which preserves the length. The next lemma states that this is generally true.

Lemma 5.4. Let $t \in \mathcal{T}(\mathcal{F}, \mathcal{V})$ be a terminating term with defined root. Then we obtain: $dh(t, \rightarrow_{\mathcal{R}}) = dh(t^{\sharp}, \rightarrow_{WDP(\mathcal{R})\cup\mathcal{R}}).$

Proof. We show $dh(t, \to_{\mathcal{R}}) \leq dh(t^{\sharp}, \to_{\mathsf{WDP}(\mathcal{R})\cup\mathcal{R}})$ by induction on $dh(t, \to_{\mathcal{R}})$. Let $\ell = dh(t, \to_{\mathcal{R}})$. If $\ell = 0$, the inequality is trivial. Suppose $\ell > 0$. Then there exists a term u such that $t \to_{\mathcal{R}} u$ and $dh(u, \to_{\mathcal{R}}) = \ell - 1$. We distinguish two cases depending on the rewrite position p.

- 1) If p is a position below the root, then clearly $\operatorname{root}(u) = \operatorname{root}(t) \in \mathcal{D}$ and $t^{\sharp} \to_{\mathcal{R}} u^{\sharp}$. Induction hypothesis yields $\operatorname{dh}(u, \to_{\mathcal{R}}) \leq \operatorname{dh}(u^{\sharp}, \to_{\operatorname{WDP}(\mathcal{R})\cup\mathcal{R}})$, and we obtain $\ell \leq \operatorname{dh}(t^{\sharp}, \to_{\operatorname{WDP}(\mathcal{R})\cup\mathcal{R}})$.
- 2) If p is a root position, then there exist a rewrite rule $l \to r \in \mathcal{R}$ and a substitution σ such that $t = l\sigma$ and $u = r\sigma$. There exists a context C such that $r = C\langle u_1, \ldots, u_n \rangle_{\mathcal{D} \cup \mathcal{V}}$ and thus by definition $l^{\sharp} \to \operatorname{COM}(u_1^{\sharp}, \ldots, u_n^{\sharp}) \in \mathsf{WDP}(\mathcal{R})$ such that $t^{\sharp} = l^{\sharp}\sigma$. Now, either $u_i \in \mathcal{V}$ or $\operatorname{root}(u_i) \in \mathcal{D}$ for every $1 \leq i \leq n$. Suppose $u_i \in \mathcal{V}$. Then $u_i^{\sharp}\sigma = u_i\sigma$ and clearly no dependency pair symbol can occur and thus,

$$\mathsf{dh}(u_i\sigma,\to_{\mathcal{R}}) = \mathsf{dh}(u_i^{\sharp}\sigma,\to_{\mathcal{R}}) = \mathsf{dh}(u_i^{\sharp}\sigma,\to_{\mathsf{WDP}(\mathcal{R})\cup\mathcal{R}})$$

Otherwise, if $\operatorname{root}(u_i) \in \mathcal{D}$ then $u_i^{\sharp} \sigma = (u_i \sigma)^{\sharp}$. Hence $\operatorname{dh}(u_i \sigma, \to_{\mathcal{R}}) \leq \operatorname{dh}(u, \to_{\mathcal{R}}) < \ell$, and we conclude $\operatorname{dh}(u_i \sigma, \to_{\mathcal{R}}) \leq \operatorname{dh}(u_i^{\sharp} \sigma, \to_{\mathsf{WDP}(\mathcal{R})\cup\mathcal{R}})$ from the induction hypothesis. Therefore,

$$\begin{split} \ell &= \mathsf{dh}(u, \to_{\mathcal{R}}) + 1 \\ &= \sum_{1 \leqslant i \leqslant n} \mathsf{dh}(u_i \sigma, \to_{\mathcal{R}}) + 1 \leqslant \sum_{1 \leqslant i \leqslant n} \mathsf{dh}(u_i^{\sharp} \sigma, \to_{\mathsf{WDP}(\mathcal{R}) \cup \mathcal{R}}) + 1 \\ &= \mathsf{dh}(\mathsf{COM}(u_1^{\sharp}, \dots, u_n^{\sharp}) \sigma, \to_{\mathsf{WDP}(\mathcal{R}) \cup \mathcal{R}}) + 1 \leqslant \mathsf{dh}(t^{\sharp}, \to_{\mathsf{WDP}(\mathcal{R}) \cup \mathcal{R}}) \,. \end{split}$$

Here we used Lemma 5.1 for the second equality.

Note that t is \mathcal{R} -reducible if and only if t^{\sharp} is $\mathsf{WDP}(\mathcal{R}) \cup \mathcal{R}$ -reducible. Hence as t is terminating, t^{\sharp} is terminating on $\rightarrow_{\mathsf{WDP}(\mathcal{R})\cup\mathcal{R}}$. Thus, similarly, $\mathsf{dh}(t, \rightarrow_{\mathcal{R}}) \ge \mathsf{dh}(t^{\sharp}, \rightarrow_{\mathsf{WDP}(\mathcal{R})\cup\mathcal{R}})$ is shown by induction on $\mathsf{dh}(t^{\sharp}, \rightarrow_{\mathsf{WDP}(\mathcal{R})\cup\mathcal{R}})$.

In the case of innermost rewriting we need not include collapsing dependency pairs as in Definition 5.2. This is guaranteed by the next lemma.

Lemma 5.5. Let t be a terminating term and σ a substitution such that $x\sigma$ is a normal form of \mathcal{R} for all $x \in \mathcal{V}ar(t)$. Then $dh(t\sigma, \rightarrow_{\mathcal{R}}) = \sum_{1 \leq i \leq n} dh(t_i\sigma, \rightarrow_{\mathcal{R}})$, whenever $t = C\langle t_1, \ldots, t_n \rangle_{\mathcal{D}}$.

Definition 5.6. Let \mathcal{R} be a TRS. If $l \to r \in \mathcal{R}$ and $r = C\langle u_1, \ldots, u_n \rangle_{\mathcal{D}}$ then the rewrite rule $l^{\sharp} \to \text{COM}(u_1^{\sharp}, \ldots, u_n^{\sharp})$ is called a *weak innermost dependency pair* of \mathcal{R} . The set of all weak innermost dependency pairs is denoted by WIDP(\mathcal{R}).

Example 5.7 (continued from Example 3.2). The set $WIDP(\mathcal{R}_{div})$ consists of the next three weak innermost dependency pairs (with respect to \xrightarrow{i}):

$$\begin{split} \mathsf{s}(x) &-^{\sharp} \mathsf{s}(y) \to x -^{\sharp} y & \mathsf{0} \div^{\sharp} \mathsf{s}(y) \to \mathsf{c} \\ \mathsf{s}(x) &\div^{\sharp} \mathsf{s}(y) \to (x - y) \div^{\sharp} \mathsf{s}(y) \,. \end{split}$$

The next lemma adapts Lemma 5.4 to innermost rewriting.

Lemma 5.8. Let t be an innermost terminating term in $\mathcal{T}(\mathcal{F}, \mathcal{V})$ with $\operatorname{root}(t) \in \mathcal{D}$. We have $\operatorname{dh}(t, \xrightarrow{i}_{\mathcal{R}}) = \operatorname{dh}(t^{\sharp}, \xrightarrow{i}_{\mathcal{W}\mathsf{IDP}(\mathcal{R})\cup\mathcal{R}})$.

Looking at the simulated version of the derivation on page 12, rules 1 and 2 are used, but neither rule 3 nor 4 is used in the \mathcal{R} -steps. In general we can approximate a subsystem of a TRS that can be used in derivations from basic terms, by employing the notion of usable rules in the dependency pair method (cf. [6, 21, 22]).

Definition 5.9. We write $f \rhd_d g$ if there exists a rewrite rule $l \to r \in \mathcal{R}$ such that $f = \operatorname{root}(l)$ and g is a defined function symbol in $\mathcal{F}un(r)$. For a set \mathcal{G} of defined function symbols we denote by $\mathcal{R} \upharpoonright \mathcal{G}$ the set of rewrite rules $l \to r \in \mathcal{R}$ with $\operatorname{root}(l) \in \mathcal{G}$. The set $\mathcal{U}(t)$ of usable rules of a term t is defined as $\mathcal{R} \upharpoonright \{g \mid f \rhd_d^* g \text{ for some } f \in \mathcal{F}un(t)\}$. Finally, if \mathcal{P} is a set of (weak) dependency pairs then $\mathcal{U}(\mathcal{P}) = \bigcup_{l \to r \in \mathcal{P}} \mathcal{U}(r)$.

Example 5.10 (continued from Examples 5.3 and 5.7). The set $\mathcal{U}(WDP(\mathcal{R}_{div}))$ of usable rules for the weak dependency pairs consists of the two rules:

1: $x - \mathbf{0} \to x$ 2: $\mathbf{s}(x) - \mathbf{s}(y) \to x - y$.

Note that we have that $\mathcal{U}(\mathsf{WDP}(\mathcal{R}_{\mathsf{div}})) = \mathcal{U}(\mathsf{WIDP}(\mathcal{R}_{\mathsf{div}})).$

We show a usable rule criterion for complexity analysis by exploiting the property that the starting terms are basic. Recall that \mathcal{T}_{b} denotes the set of basic terms; we set $\mathcal{T}_{b}^{\sharp} = \{t^{\sharp} \mid t \in \mathcal{T}_{b}\}$.

Lemma 5.11. Let \mathcal{P} be a set of weak dependency pairs and let $(t_i)_{i=0,1,\dots}$ be a (finite or infinite) derivation of $\mathcal{P} \cup \mathcal{R}$. If $t_0 \in \mathcal{T}_b^{\sharp}$ then $(t_i)_{i=0,1,\dots}$ is a derivation of $\mathcal{P} \cup \mathcal{U}(\mathcal{P})$.

Proof. Let \mathcal{G} be the set of all non-usable symbols with respect to \mathcal{P} . We write P(t) if $t|_q \in \mathsf{NF}(\mathcal{R})$ for all $q \in \mathcal{P}\mathsf{os}_{\mathcal{G}}(t)$. First we prove by induction on i that $P(t_i)$ holds for all i.

- 1) Assume i = 0. Since $t_0 \in \mathcal{T}_b^{\sharp}$, we have $t_0 \in \mathsf{NF}(\mathcal{R})$ and thus $t|_p \in \mathsf{NF}(\mathcal{R})$ for all positions p. The assertion P follows trivially.
- 2) Suppose i > 0. By induction hypothesis, $P(t_{i-1})$ holds, i.e., there exist $p \in \mathcal{P}os(t_{i-1})$, a substitution σ , and $l \to r \in \mathcal{U}(\mathcal{P}) \cup \mathcal{P}$, such that $t_{i-1}|_p = l\sigma$ and $t_i|_p = r\sigma$. In order to show property P for t_i , we fix a position $q \in \mathcal{P}os_{\mathcal{G}}(t)$. We have to show $t_i|_q \in \mathsf{NF}(\mathcal{R})$. We distinguish three subcases:
 - Suppose that q is above p. Then $t_{i-1}|_q$ is reducible, but this contradicts the induction hypothesis $P(t_{i-1})$.
 - Suppose p and q are parallel but distinct. Since $t_{i-1}|_q = t_i|_q \in \mathsf{NF}(\mathcal{R})$ holds, we obtain $P(t_i)$.
 - Otherwise, q is below p. Then, $t_i|_q$ is a subterm of $r\sigma$. Because r contains no \mathcal{G} -symbols by the definition of usable symbols, $t_i|_q$ is a subterm of $x\sigma$ for some $x \in \mathcal{V}ar(r) \subseteq \mathcal{V}ar(l)$. Therefore, $t_i|_q$ is also a subterm of $t_{i-1}|_q$, from which $t_i|_q \in \mathsf{NF}(\mathcal{R})$ follows. We obtain $P(t_i)$.

Hence property P holds for all t_i in the assumed derivation. Thus any reduction step $t_i \to_{\mathcal{R}\cup\mathcal{P}} t_{i+1}$ can be simulated by a step $t_i \to_{\mathcal{U}(\mathcal{P})\cup\mathcal{P}} t_{i+1}$. From this the lemma follows. \Box

Note that the proof technique adopted for termination analysis [21, 22] cannot be directly used in this context. The technique transforms terms in a derivation to exclude non-usable rules. However, since the size of the initial term increases, this technique does not suit to our use. On the other hand, the transformation employed in [22] is adaptable to a complexity analysis in the large, cf. [23].

The next theorem follows from Lemmas 5.4 and 5.8 in conjunction with the above Lemma 5.11. It adapts the usable rule criteria to complexity analysis.

Theorem 5.12. Let \mathcal{R} be a TRS and let $t \in \mathcal{T}_b$. If t is terminating with respect to \rightarrow then $dh(t, \rightarrow) = dh(t^{\sharp}, \rightarrow_{\mathcal{P} \cup \mathcal{U}(\mathcal{P})})$, where \rightarrow denotes $\rightarrow_{\mathcal{R}}$ or $\stackrel{i}{\rightarrow}_{\mathcal{R}}$ depending on whether $\mathcal{P} = WDP(\mathcal{R})$ or $\mathcal{P} = WIDP(\mathcal{R})$.

To clarify the applicability of the theorem in complexity analysis, we instantiate the theorem by considering RMIs.

Corollary 5.13. Let \mathcal{R} be a TRS, let μ be the (innermost) usable replacement map and let $\mathcal{P} = \mathsf{WDP}(\mathcal{R})$ (or $\mathcal{P} = \mathsf{WIDP}(\mathcal{R})$). If $\mathcal{P} \cup \mathcal{U}(\mathcal{P})$ is compatible with a d-degree μ -monotone RMI \mathcal{A} , then the (innermost) runtime complexity function $\mathrm{rc}_{\mathcal{R}}^{(i)}$ with respect to \mathcal{R} is bounded by a d-degree polynomial.

Proof. For simplicity we suppose $\mathcal{P} = \mathsf{WDP}(\mathcal{R})$ and let \mathcal{A} be a μ -monotone RMI of degree d. Compatibility of \mathcal{A} with $\mathcal{P} \cup \mathcal{U}(\mathcal{P})$ implies the well-foundedness of the relation $\rightarrow_{\mathcal{P} \cup \mathcal{U}(\mathcal{P})}$ on the set of terms $\mathcal{T}_{\mathsf{b}}^{\sharp}$, cf. Theorem 4.8. This in turn implies the well-foundedness of $\rightarrow_{\mathcal{R}}$, cf. Lemma 5.11. Hence Theorem 5.12 is applicable and we conclude $\mathsf{dh}(t, \rightarrow_{\mathcal{R}}) = \mathsf{dh}(t^{\sharp}, \rightarrow_{\mathcal{P} \cup \mathcal{U}(\mathcal{P})})$. On the other hand, due to Theorem 3.9 compatibility with \mathcal{A} implies that $\mathsf{dh}(t^{\sharp}, \rightarrow_{\mathcal{P} \cup \mathcal{U}(\mathcal{P})}) = \mathsf{O}(|t^{\sharp}|^d)$. As $|t^{\sharp}| = |t|$, we can combine these equalities to conclude polynomial runtime complexity of \mathcal{R} .

The below given example applies Corollary 5.13 to the motivating Example 3.2 introduced in Section 1.

Example 5.14 (continued from Example 5.10). Consider the TRS \mathcal{R}_{div} for division used as running example; the weak dependency pairs $\mathcal{P} := \mathsf{WDP}(\mathcal{R}_{div})$ are given in Example 5.3. We have $\mathcal{U}(\mathcal{P}) = \{1,2\}$ and let $\mathcal{S} = \mathcal{P} \cup \mathcal{U}(\mathcal{P})$. The usable replacement map $\mu := \mu_{\mathsf{f}}^{\mathcal{S}}$ is defined as follows:

$$\mu(s) = \mu(-) = \mu(-^{\sharp}) = \emptyset \qquad \mu(\div^{\sharp}) = \{1\}$$

Note that μ_{f}^{S} is smaller than $\mu_{f}^{\mathcal{R}}$ on \mathcal{F} (see Example 4.10). Consider the 1-dimensional RMI \mathcal{A} with $\mathbf{0}_{\mathcal{A}} = \mathbf{c}_{\mathcal{A}} = \mathbf{d}_{\mathcal{A}} = 0$, $\mathbf{s}_{\mathcal{A}}(x) = x + 2$, $-\mathcal{A}(x,y) = -\overset{\sharp}{\mathcal{A}}(x,y) = x + 1$, and $\div^{\sharp}_{\mathcal{A}}(x,y) = x + 1$. The algebra \mathcal{A} is strictly monotone on all usable argument positions and the rules in \mathcal{S} are interpreted and ordered as follows:

1:
$$x+1 > x$$
 5: $1 > 0$ 7: $1 > 0$
2: $x+3 > x+1$ 6: $x+3 > x+1$ 8: $x+3 > x+2$

Therefore, S is compatible with A and the runtime complexity function $\mathsf{rc}_{\mathcal{R}}$ is linear. Remark that by looking at the coefficients of the interpretations more precise bound can be inferred. Since all coefficients are at most one, we obtain $\mathsf{rc}_{\mathcal{R}}(n) \leq n + c$ for some $c \in \mathbb{N}$.

It is worth stressing that it is (often) easier to analyse the complexity of $\mathcal{P} \cup \mathcal{U}(\mathcal{P})$ than the complexity of \mathcal{R} . This is exemplified by the next example.

Example 5.15. Consider the TRS \mathcal{R}_D

$$\begin{split} \mathsf{D}(\mathsf{c}) &\to \mathbf{0} \qquad \mathsf{D}(x+y) \to \mathsf{D}(x) + \mathsf{D}(y) \qquad \mathsf{D}(x \times y) \to (y \times \mathsf{D}(x)) + (x \times \mathsf{D}(y)) \\ \mathsf{D}(\mathsf{t}) \to \mathbf{1} \qquad \mathsf{D}(x-y) \to \mathsf{D}(x) - \mathsf{D}(y) \; . \end{split}$$

There is no 1-dimensional $\mu_{\rm f}$ -monotone RMI compatible with $\mathcal{R}_{\rm D}$. On the other hand $WDP(\mathcal{R}_{\rm D})$ consists of the five pairs

$$\begin{aligned} \mathsf{D}^{\sharp}(\mathsf{c}) &\to \mathsf{c}_{1} \qquad \mathsf{D}^{\sharp}(x+y) \to \mathsf{c}_{3}(\mathsf{D}^{\sharp}(x),\mathsf{D}^{\sharp}(y)) \qquad \mathsf{D}^{\sharp}(x\times y) \to \mathsf{c}_{5}(y,\mathsf{D}^{\sharp}(x),x,\mathsf{D}^{\sharp}(y)) \\ \mathsf{D}^{\sharp}(\mathsf{t}) \to \mathsf{c}_{2} \qquad \mathsf{D}^{\sharp}(x-y) \to \mathsf{c}_{4}(\mathsf{D}^{\sharp}(x),\mathsf{D}^{\sharp}(y)) \;, \end{aligned}$$

and $\mathcal{U}(\mathsf{WDP}(\mathcal{R}_{\mathsf{D}})) = \emptyset$. The usable replacement map μ_{f} for $\mathsf{WDP}(\mathcal{R}_{\mathsf{D}}) \cup \mathcal{U}(\mathcal{R}_{\mathsf{D}})$ is defined as $\mu_{\mathsf{f}}(\mathsf{c}_3) = \mu_{\mathsf{f}}(\mathsf{c}_4) = \{1, 2\}, \ \mu_{\mathsf{f}}(\mathsf{c}_5) = \{2, 4\}, \ \text{and} \ \mu_{\mathsf{f}}(f) = \emptyset$ for all other symbols f. Since the 1-dimensional μ_{f} -monotone RMI \mathcal{A} with

$$\begin{aligned} \mathsf{D}^{\sharp}_{\mathcal{A}}(x) &= 2x \qquad \mathsf{c}_{\mathcal{A}} = \mathsf{t}_{\mathcal{A}} = 1 \qquad +_{\mathcal{A}}(x,y) = -_{\mathcal{A}}(x,y) = \times_{\mathcal{A}}(x,y) = x + y + 1 \\ \mathsf{c}_{1\mathcal{A}} &= \mathsf{c}_{2\mathcal{A}} = 0 \qquad \mathsf{c}_{3\mathcal{A}}(x,y) = \mathsf{c}_{4\mathcal{A}}(x,y) = x + y \qquad \mathsf{c}_{5\mathcal{A}}(x,y,z,w) = y + w \,, \end{aligned}$$

is compatible with \mathcal{R}_D , linear runtime complexity of \mathcal{R}_D is concluded. Remark that this bound is optimal.

We conclude this section by discussing the (in-)applicability of standard dependency pairs (see [6]) in complexity analysis. For that we recall the definition of standard dependency pairs.

Definition 5.16 ([6]). The set $\mathsf{DP}(\mathcal{R})$ of (standard) dependency pairs of a TRS \mathcal{R} is defined as $\{l^{\sharp} \to u^{\sharp} \mid l \to r \in \mathcal{R}, u \leq r, \operatorname{root}(u) \text{ is defined, and } u \not \lhd l\}.$

The next example shows that Lemma 5.4 (Lemma 5.8) does not hold if we replace weak (innermost) dependency pairs with standard dependency pairs.

Example 5.17. Consider the one-rule TRS \mathcal{R} : $f(\mathbf{s}(x)) \to \mathbf{g}(f(x), f(x))$. $\mathsf{DP}(\mathcal{R})$ is the singleton of $f^{\sharp}(\mathbf{s}(x)) \to f^{\sharp}(x)$. Let $t_n = f(\mathbf{s}^n(x))$ for each $n \ge 0$. Since $t_{n+1} \to_{\mathcal{R}} \mathbf{g}(t_n, t_n)$ holds for all $n \ge 0$, it is easy to see $\mathsf{dh}(t_{n+1}, \to_{\mathcal{R}}) \ge 2^n$, while $\mathsf{dh}(t_{n+1}^{\sharp}, \to_{\mathsf{DP}(\mathcal{R})\cup\mathcal{R}}) = n$.

6 The Weight Gap Principle

Let $\mathcal{P} = \mathsf{WDP}(\mathcal{R}_{\mathsf{div}})$ and recall the derivation over $\mathcal{P} \cup \mathcal{R}_{\mathsf{div}}$ on page 13. This derivation can be represented as derivation of \mathcal{P} modulo $\mathcal{U}(\mathcal{P})$:

$$4 \div^{\sharp} 2 \rightarrow_{\mathcal{P}/\mathcal{U}(\mathcal{P})} 2 \div^{\sharp} 2 \rightarrow_{\mathcal{P}/\mathcal{U}(\mathcal{P})} 0 \div^{\sharp} 2 \rightarrow_{\mathcal{P}/\mathcal{U}(\mathcal{P})} \mathsf{c}$$

As we see later linear runtime complexity of $\mathcal{U}(\mathcal{P})$ and $\mathcal{P}/\mathcal{U}(\mathcal{P})$ can be easily obtained. If linear runtime complexity of $\mathcal{P} \cup \mathcal{U}(\mathcal{P})$ would follow from them, linear runtime complexity of \mathcal{R} could be established in a modular way.

In order to bound complexity of relative TRSs we define a variant of a reduction pair [6]. Note that G is associated to a given collapsible order.

Definition 6.1. A μ -complexity pair for a relative TRS \mathcal{R}/\mathcal{S} is a pair (\gtrsim, \succ) such that \gtrsim is a μ -monotone proper order and \succ is a strict order. Moreover \gtrsim and \succ are compatible, that is, $\gtrsim \cdot \succ \subseteq \succ$ or $\succ \cdot \gtrsim \subseteq \succ$. Finally \succ is collapsible on $\rightarrow_{\mathcal{R}/\mathcal{S}}$ and all compound symbols are μ -monotone with respect to \succ .

Lemma 6.2. Let $\mathcal{P} = \mathsf{WDP}(\mathcal{R})$ and (\gtrsim,\succ) a $\mu_{\mathsf{f}}^{\mathcal{P} \cup \mathcal{U}(\mathcal{P})}$ -complexity pair for $\mathcal{P}/\mathcal{U}(\mathcal{P})$. If $\mathcal{P} \subseteq \succ$ and $\mathcal{U}(\mathcal{P}) \subseteq \gtrsim$ then $\mathsf{dh}(t, \rightarrow_{\mathcal{P}/\mathcal{U}(\mathcal{P})}) \leqslant \mathsf{G}(t)$ for any $t \in \mathcal{T}_{\mathsf{b}}^{\sharp}$.

Example 6.3 (continued from Example 5.14). Consider the 1-dimensional RMI \mathcal{A} with

$$\mathbf{0}_{\mathcal{A}} = \mathbf{c}_{\mathcal{A}} = \mathbf{d}_{\mathcal{A}} = 0 \qquad \mathbf{s}_{\mathcal{A}}(x) = x + 1 \qquad -\mathcal{A}(x, y) = -\overset{\sharp}{\mathcal{A}}(x, y) = \div \overset{\sharp}{\mathcal{A}}(x, y) = x + 1$$

which yields the complexity pair $(\geq_{\mathcal{A}}, \geq_{\mathcal{A}})$ for $\mathcal{P}/\mathcal{U}(\mathcal{P})$. Since $\mathcal{P} \subseteq \geq_{\mathcal{A}}$ and $\mathcal{U}(\mathcal{P}) \subseteq \geq_{\mathcal{A}}$ hold, $\operatorname{comp}(n, \mathcal{T}_{\mathsf{b}}^{\sharp}, \rightarrow_{\mathcal{P}/\mathcal{U}(\mathcal{P})}) = \mathsf{O}(n)$.

First we show the main theorem of this section.

Definition 6.4. Let \mathcal{A} be a matrix interpretation and let \mathcal{R}/\mathcal{S} be a relative TRS. A *weight* gap on a set T of terms is a number $\Delta \in \mathbb{N}$ such that $s \in \mathcal{A}^*_{\mathcal{R}\cup\mathcal{S}}(T)$ and $s \to_{\mathcal{R}} t$ implies $[t]_1 - [s]_1 \leq \Delta$.

Let T be a set of terms and let \mathcal{R}/\mathcal{S} be a relative TRS.

Theorem 6.5. If \mathcal{R}/\mathcal{S} is terminating, \mathcal{A} admits a weight gap Δ on T, and \mathcal{A} is a matrix interpretation of degree d such that \mathcal{S} is compatible with \mathcal{A} , then there exists $c \in \mathbb{N}$ such that $dh(t, \rightarrow_{\mathcal{R}\cup\mathcal{S}}) \leq (1+\Delta) \cdot dh(t, \rightarrow_{\mathcal{R}/\mathcal{S}}) + c \cdot |t|^d$ for all $t \in T$. Consequently, $comp(n, T, \rightarrow_{\mathcal{R}\cup\mathcal{S}}) = O(comp(n, T, \rightarrow_{\mathcal{R}/\mathcal{S}}) + n^d)$ holds.

Proof. Let $m = dh(s, \rightarrow_{\mathcal{R}/\mathcal{S}})$ and n = |s|. Any derivation of $\rightarrow_{\mathcal{R}\cup\mathcal{S}}$ is representable as follows:

$$s = s_0 \to_{\mathcal{S}}^{k_0} t_0 \to_{\mathcal{R}} s_1 \to_{\mathcal{S}}^{k_1} t_1 \to_{\mathcal{R}} \cdots \to_{\mathcal{S}}^{k_m} t_m .$$

Without loss of generality we may assume that the derivation is maximal and ground. We observe:

- 1) $k_i \leq [s_i]_1 [t_i]_1$ holds for all $0 \leq i \leq m$. This is because $[s]_1 > [t]_1$, whenever $s \to_S t$ by the assumption S is compatible with A. By definition of >, we conclude $[s]_1 \geq [t]_1 + 1$ whenever $s \to_S t$. From the fact that $s_i \to_S^{k_i} t_i$ we thus obtain $k_i \leq [s_i]_1 [t_i]_1$.
- 2) $([s_{i+1}])_1 \leq ([t_i])_1 + \Delta$ holds for all $0 \leq i < m$ by the assumption.
- 3) There exists a number c such that for any term $s \in T$, $[s]_1 \leq c \cdot |s|^d$. This follows by the degree of \mathcal{A} .

We obtain the following inequalities:

$$\begin{aligned} \mathsf{dh}(s_0, \to_{\mathcal{R}\cup\mathcal{S}}) &= m + k_0 + \dots + k_m \\ &\leqslant m + ([s_0]_1 - [t_0]_1) + \dots + ([s_m]_1 - [t_m]_1) \\ &= m + [s_0]_1 + ([s_1]_1 - [t_0]_1) + \dots + ([s_m]_1 - [t_{m-1}]_1) - [t_m]_1 \\ &\leqslant m + [s_0]_1 + ([t_0]_1 + \Delta - [t_0]_1) + \dots - [t_m]_1 \\ &\leqslant m + [s_0]_1 + m\Delta - [t_m]_1 \\ &\leqslant m + [s_0]_1 + m\Delta \\ &\leqslant (1 + \Delta)m + c \cdot |s_0|^d . \end{aligned}$$

Here we use property 1) *m*-times in the second line. We used property 2) in the third line and property 3) in the last line. \Box

A question is when a weight gap is admitted. We present two conditions. We start with a simple version for derivational complexity, and then we adapt it for runtime complexity.

We employ a very restrictive form of TMIs. Every $f \in \mathcal{F}$ is interpreted by the following restricted linear function:

$$f_{\mathcal{A}}: (\vec{v}_1, \ldots, \vec{v}_n) \mapsto \mathbf{1}\vec{v}_1 + \ldots + \mathbf{1}\vec{v}_n + \vec{f}.$$

I.e., the only matrix employed in this interpretation is the unit matrix **1**. Such a matrix interpretation is called *strongly linear* (*SLMI* for short).

Lemma 6.6. If \mathcal{R} is non-duplicating and \mathcal{A} is an SLMI, then \mathcal{R}/\mathcal{S} and \mathcal{A} admit a weight gap on all terms.

Proof. Let $\Delta := \max\{[r]_1 \doteq [l]_1 \mid l \rightarrow r \in \mathcal{R}\}$. We show that Δ gives a weight gap. In proof, we first show the following equality.

$$\Delta = \max\{([\alpha]_{\mathcal{A}}(r))_1 \div ([\alpha]_{\mathcal{A}}(l))_1 \mid l \to r \in \mathcal{R}, \alpha \colon \mathcal{V} \to \mathcal{A}\}.$$
 (1)

Although the proof is not difficult, we give the full account in order to utilise it later. Observe that for any matrix interpretation \mathcal{A} and rule $l \to r \in \mathcal{R}$, there exist matrices (over \mathbb{N}) $L_1, \ldots, L_k, R_1, \ldots, R_k$ and vectors \vec{l}, \vec{r} such that:

$$[\alpha]_{\mathcal{A}}(l) = \sum_{i=1}^{k} L_i \cdot \alpha(x_i) + \vec{l} \qquad [\alpha]_{\mathcal{A}}(r) = \sum_{i=1}^{k} R_i \cdot \alpha(x_i) + \vec{r},$$

where k denotes the cardinality of $Var(l) \supseteq Var(r)$. Conclusively, we obtain:

$$[\alpha]_{\mathcal{A}}(r) \doteq [\alpha]_{\mathcal{A}}(l) = \sum_{i=1}^{k} (R_i \doteq L_i)\alpha(x_i) + (\vec{r} \doteq \vec{l}) .$$
⁽²⁾

Here \div denotes the natural component-wise extension of the modified minus to vectors.

As \mathcal{A} is an SLMI the matrices L_i , R_i are obtained by multiplying or adding unit matrices, where the latter case can only happen if (at least one) of the variables x_i occurs multiple times in l or r. Due to the fact that $l \to r$ is non-duplicating, this effect is canceled out. Thus the right-hand side of (2) is independent on the assignment α and we conclude:

$$[r]_1 \div [l]_1 = ([\alpha]_{\mathcal{A}}(r) \div [\alpha]_{\mathcal{A}}(l))_1 = (\vec{r} \div \vec{l})_1$$

By definition $\Delta = \max\{[r]_1 \doteq [l]_1 \mid l \rightarrow r \in \mathcal{R}\}$ and thus (1) follows.

Let $C[\Box]$ denote a (possible empty) context such that $s = C[l\sigma] \rightarrow_{\mathcal{R}} C[r\sigma] = t$, where $l \rightarrow r \in \mathcal{R}$ and σ a substitution. We prove the lemma by induction on C.

- 1) Suppose $C[\Box] = \Box$, that is, $s = l\sigma$ and $t = r\sigma$. There exists an assignment α_1 such that $[l\sigma] = [\alpha_1]_{\mathcal{A}}(l)$ and $[r\sigma] = [\alpha_1]_{\mathcal{A}}(r)$. By (1) we conclude for the assignment α_1 : $([\alpha_1]_{\mathcal{A}}(l))_1 + \Delta \ge ([\alpha_1]_{\mathcal{A}}(r))_1$. Therefore in sum we obtain $[s]_1 + \Delta \ge [t]_1$.
- 2) Suppose $C[\Box] = f(t_1, \ldots, t_{i-1}, C'[\Box], t_{i+1}, \ldots, t_n)$. Hence, we obtain:

$$[f(t_1, \dots, C'[l\sigma], \dots, t_n)]_1 + \Delta$$

= $[t_1]_1 + \dots + ([C'[l\sigma]]_1 + \Delta) + \dots + [t_n]_1 + (\vec{f})_1$
$$\geq [t_1]_1 + \dots + [C'[r\sigma]]_1 + \dots + [t_n]_1 + (\vec{f})_1$$

= $[f(t_1, \dots, C'[r\sigma], \dots, t_n)]_1$,

for some vector $\vec{f} \in \mathbb{N}^d$. In the first and last line, we employ the fact that \mathcal{A} is strongly linear. In the second line the induction hypothesis is applied together with the (trivial) fact that \mathcal{A} is strictly monotone on all arguments of f by definition.

Note that the combination of Theorem 6.5 and Lemma 6.6 corresponds to (the corrected version of) Theorem 24 in [4]. In [4] 1-dimensional SLMIs are called *strongly linear interpretations* (*SLIs* for short).

Example 6.7. Consider the TRS \mathcal{R}

$$1: \ \mathsf{f}(\mathsf{s}(x)) \to \mathsf{f}(x-\mathsf{s}(\mathbf{0})) \qquad 2: \ x-\mathsf{0} \to x \qquad 3: \ \mathsf{s}(x)-\mathsf{s}(y) \to x-y \; .$$

 $\mathcal{P} := \mathsf{WDP}(\mathcal{R})$ consists of the three pairs

$$\mathsf{f}^{\sharp}(\mathsf{s}(x)) \to \mathsf{f}^{\sharp}(x - \mathsf{s}(0)) \qquad \qquad x - {}^{\sharp} \, \mathsf{0} \to x \qquad \qquad \mathsf{s}(x) - {}^{\sharp} \, \mathsf{s}(y) \to x - {}^{\sharp} \, y \; ,$$

and $\mathcal{U}(\mathcal{P}) = \{2, 3\}$. Obviously \mathcal{P} is non-duplicating and there exists an SLI \mathcal{A} with $\mathcal{U}(\mathcal{P}) \subseteq \mathcal{F}_{\mathcal{A}}$. Thus, Lemma 6.6 yields a weight gap for $\mathcal{P}/\mathcal{U}(\mathcal{P})$. By taking the 1-dimensional RMI \mathcal{B} with

$$\begin{aligned} \mathsf{s}_{\mathcal{B}}(x) &= x + 1 & -_{\mathcal{B}}(x, y) = x & \mathsf{f}_{\mathcal{B}}(x) = \mathsf{f}_{\mathcal{B}}^{\sharp}(x) = x \\ \mathsf{0}_{\mathcal{B}} &= 0 & -^{\sharp}_{\mathcal{B}}(x, y) = x + 1 , \end{aligned}$$

we obtain $\mathcal{P} \subseteq \succ_{\mathcal{B}}$ and $\mathcal{U}(\mathcal{P}) \subseteq \succeq_{\mathcal{B}}$. Therefore, $\operatorname{comp}(n, \mathcal{T}_{\mathsf{b}}^{\sharp}, \rightarrow_{\mathcal{P}/\mathcal{U}(\mathcal{P})}) = \mathsf{O}(n)$. Hence, $\operatorname{rc}_{\mathcal{R}}(n) = \operatorname{comp}(n, \mathcal{T}_{\mathsf{b}}^{\sharp}, \rightarrow_{\mathcal{P}\cup\mathcal{U}(\mathcal{P})}) = \mathsf{O}(n)$ is concluded by Theorem 6.5.

The next lemma shows that there is no advantage to consider SLMIs of dimension $k \ge 2$. Lemma 6.8. If S is compatible with some SLMI A then S is compatible with some SLI B.

Proof. Let \mathcal{A} be an SLMI of dimension k. Further, let $\alpha : \mathcal{V} \to \mathbb{N}$ denote an arbitrary assignment. We define $\hat{\alpha} : \mathcal{V} \to \mathbb{N}^k$ as $\hat{\alpha}(x) = (\alpha(x), 0, \dots, 0)^{\top}$ for each variable x. We define the SLI \mathcal{B} by $f_{\mathcal{B}}(x_1, \dots, x_n) = x_1 + \dots + x_n + \vec{f_1}$. Then,

$$f_{\mathcal{B}}(x_1, \dots, x_n) = \left((x_1, 0, \dots, 0)^\top + \dots + (x_n, 0, \dots, 0)^\top + \vec{f} \right)_1$$
$$= \left(f_{\mathcal{A}}((x_1, 0, \dots, 0)^\top, \dots, (x_n, 0, \dots, 0)^\top)) \right)_1$$

Therefore, easy structural induction shows that $[\alpha]_{\mathcal{B}}(t) = ([\widehat{\alpha}]_{\mathcal{A}}(t))_1$ for all terms t. Hence, $\mathcal{S} \subseteq \succ_{\mathcal{B}}$ whenever $\mathcal{S} \subseteq \succ_{\mathcal{A}}$.

The next example shows that in Lemma 6.6 SLMIs cannot be simply replaced by RMIs. Example 6.9. Consider the TRSs \mathcal{R}_{exp}

$$\begin{aligned} \exp(0) &\to \mathsf{s}(0) & \mathsf{d}(0) \to 0 \\ \exp(\mathsf{r}(x)) &\to \mathsf{d}(\exp(x)) & \mathsf{d}(\mathsf{s}(x)) \to \mathsf{s}(\mathsf{s}(\mathsf{d}(x))) \end{aligned}$$

This TRS formalises the exponentiation function. Setting $t_n = \exp(\mathbf{r}^n(\mathbf{0}))$ we obtain $dh(t_n, \rightarrow_{\mathcal{R}_{exp}}) \ge 2^n$ for each $n \ge 0$. Thus the runtime complexity of \mathcal{R}_{exp} is exponential.

In order to show the claim, we split \mathcal{R}_{exp} into two TRSs $\mathcal{R} = \{exp(0) \rightarrow s(0), exp(r(x)) \rightarrow d(exp(x))\}$ and $\mathcal{S} = \{d(0) \rightarrow 0, d(s(x)) \rightarrow s(s(d(x)))\}$. Then it is easy to verify that the next 1-dimensional RMI \mathcal{A} is compatible with \mathcal{S} :

$$\mathbf{0}_{\mathcal{A}} = 0$$
 $\mathbf{d}_{\mathcal{A}}(x) = 3x$ $\mathbf{s}_{\mathcal{A}}(x) = x + 1$.

Moreover an upper-bound of $dh(t_n, \rightarrow_{\mathcal{R}/S})$ can be estimated by using the following 1dimensional TMI \mathcal{B} :

$$\mathbf{0}_{\mathcal{B}} = 0$$
 $\mathbf{d}_{\mathcal{B}}(x) = \mathbf{s}_{\mathcal{B}}(x) = x$ $\exp_{\mathcal{B}}(x) = \mathbf{r}_{\mathcal{B}}(x) = x + 1$.

Since $\rightarrow_{\mathcal{R}} \subseteq >_{\mathcal{B}}$ and $\rightarrow_{\mathcal{S}}^* \subseteq \geq_{\mathcal{B}}$ hold, we have $\rightarrow_{\mathcal{R}/\mathcal{S}} \subseteq >_{\mathcal{B}}$. Hence $\mathsf{dh}(t_n, \rightarrow_{\mathcal{R}/\mathcal{S}}) \leq [\alpha_0]_{\mathcal{B}}(t_n) = n + 2$. But clearly from this we cannot conclude a polynomial bound on the derivation length of $\mathcal{R} \cup \mathcal{S} = \mathcal{R}_{\mathsf{exp}}$, as the runtime complexity of $\mathcal{R}_{\mathsf{exp}}$ is exponential.

Furthermore, non-duplication of \mathcal{R} is also essential for Lemma 6.6.³

Example 6.10. Consider the following $\mathcal{R} \cup \mathcal{S}$

1:
$$f(s(x), y) \rightarrow f(x, d(y, y, y))$$

3: $d(0, 0, x) \rightarrow x$
3: $d(s(x), s(y), z) \rightarrow d(x, y, s(z))$

Let $\mathcal{R} = \{1\}$ and let $\mathcal{S} = \{2, 3\}$. The following SLI \mathcal{A} is compatible with \mathcal{S} :

$$\mathsf{d}_{\mathcal{A}}(x,y,z) = x + y + z + 1 \qquad \mathsf{s}_{\mathcal{A}}(x) = x + 1 \qquad \mathsf{0}_{\mathcal{A}} = 0 \; .$$

Furthermore, the following $\mu_{f}^{\mathcal{R}\cup\mathcal{S}}$ -monotone 1-dimensional RMI \mathcal{B} orients the rule in \mathcal{R} strictly, while the rules in \mathcal{S} are weakly oriented.

$$f_{\mathcal{B}}(x,y) = x$$
 $d_{\mathcal{B}}(x,y,z) = x + y + z$ $s_{\mathcal{B}}(x) = x + 1$ $0_{\mathcal{B}} = 0$

Thus, $\operatorname{comp}(n, \mathcal{T}_{\mathsf{b}}, \to_{\mathcal{R}/\mathcal{S}}) = \mathsf{O}(n)$ is obtained. If the restriction that \mathcal{R} is non-duplicating could be dropped from Lemma 6.6, we would conclude $\operatorname{rc}_{\mathcal{R}\cup\mathcal{S}}(n) = \mathsf{O}(n)$. However, it is easy to see that $\operatorname{rc}_{\mathcal{R}\cup\mathcal{S}}$ is at least exponential. Setting $t_n := \mathsf{f}(\mathsf{s}^n(\mathsf{0}),\mathsf{s}(\mathsf{0}))$, we obtain $\operatorname{dh}(t_n, \to_{\mathcal{R}\cup\mathcal{S}}) \geq 2^n$ for any $n \geq 1$.

We present a weight gap condition for runtime complexity analysis. When considering the derivation in the beginning of this section (on page 17), every step by a weak dependency pair only takes place as an outermost step. Exploiting this fact we can relax the restriction that was imposed in the above examples. To this end, we introduce a generalised notion of non-duplicating TRSs.

Below max { $([\alpha]_{\mathcal{A}}(r))_1 \div ([\alpha]_{\mathcal{A}}(l))_1 | l \rightarrow r \in \mathcal{P} \text{ and } \alpha : \mathcal{V} \rightarrow \mathcal{A}$ } is referred to as $\Delta(\mathcal{A}, \mathcal{P})$. We say that a μ -monotone RMI is *adequate* if all compound symbols are interpreted as μ -monotone SLMI.

Lemma 6.11. Let $\mathcal{P} = \mathsf{WDP}(\mathcal{R})$ and let \mathcal{A} be an adequate $\mu_{\mathsf{f}}^{\mathcal{P} \cup \mathcal{U}(\mathcal{P})}$ -monotone RMI. Suppose $\Delta(\mathcal{A}, \mathcal{P})$ is well-defined on \mathbb{N} . Then, $\mathcal{P}/\mathcal{U}(\mathcal{P})$ and \mathcal{A} admit a weight gap on $\mathcal{T}_{\mathsf{h}}^{\sharp}$.

Proof. The proof follows the proof of Lemma 6.6. We set $\Delta = \Delta(\mathcal{A}, \mathcal{P})$. Let $s \to_{\mathcal{P}} t$ with $s \in \to_{\mathcal{P}\cup\mathcal{U}(\mathcal{P})}(\mathcal{T}_{\mathsf{b}}^{\sharp})$. One may write $s = C[l\sigma]$ and $t = C[r\sigma]$ with $l \to r \in \mathcal{P}$, where C denotes a context. Note that due to $s \in \to_{\mathcal{P}\cup\mathcal{U}(\mathcal{P})}(\mathcal{T}_{\mathsf{b}}^{\sharp})$ all function symbols above the hole in C are compound symbols. We perform induction on C.

- 1) If $C = \Box$ then $[t]_1 [s]_1 \leq \Delta$ by the definition of $\Delta(\mathcal{A}, \mathcal{P})$.
- 2) For inductive step, C must be of the form $c(u_1, \ldots, u_{i-1}, C', u_{i+1}, \ldots, u_n)$ with $i \in \mu(c)$. Since \mathcal{A} is adequate, $c_{\mathcal{A}}$ is a SLMI. The rest of reasoning is same with 2) in the proof of Lemma 6.6.

³ This example is due to Dieter Hofbauer and Andreas Schnabl.

Example 6.12 (continued from Example 6.3). Consider the following adequate $\mu_{f}^{\mathcal{P}\cup\mathcal{U}(\mathcal{P})}$ -monotone 1-dimensional RMI \mathcal{B} :

$$\mathbf{0}_{\mathcal{B}} = \mathbf{c}_{\mathcal{B}} = \mathbf{d}_{\mathcal{B}} = 0 \qquad \mathbf{s}_{\mathcal{B}}(x) = x + 2 \qquad -_{\mathcal{B}}(x, y) = -_{\mathcal{B}}^{\sharp}(x, y) = \div_{\mathcal{B}}^{\sharp}(x, y) = x + 1$$

Since $\Delta(\mathcal{B}, \mathcal{P})$ is well-defined (indeed 1), \mathcal{B} admits the weight gap of Lemma 6.11. Moreover, $\mathcal{U}(\mathcal{P})$ is compatible with $\succ_{\mathcal{B}}$. As $\mathsf{comp}(n, \mathcal{T}_{\mathsf{b}}^{\sharp}, \rightarrow_{\mathcal{P}/\mathcal{U}(\mathcal{P})}) = \mathsf{O}(n)$ was shown in Example 6.3, Theorem 6.5 deduces linear runtime complexity for $\mathcal{R}_{\mathsf{div}}$.

In Lemma 6.11 $\Delta(\mathcal{A}, \mathcal{P})$ must be well-defined.

Example 6.13. Consider the following TRS \mathcal{R}

$$\begin{aligned} 1: \quad \mathsf{f}([]) \to [] & 3: \quad \mathsf{g}([], z) \to z \\ 2: \quad \mathsf{f}(x:y) \to x: \mathsf{f}(\mathsf{g}(y, [])) & 4: \quad \mathsf{g}(x:y, z) \to \mathsf{g}(y, x:z) \end{aligned}$$

whose optimal innermost runtime complexity is quadratic. The weak innermost dependency pairs $\mathcal{P} := \mathsf{WIDP}(\mathcal{R})$ are

$$\begin{aligned} 5: \quad \mathsf{f}^{\sharp}([]) \to \mathsf{c} & 7: \quad \mathsf{g}^{\sharp}([], z) \to \mathsf{d} \\ 6: \quad \mathsf{f}^{\sharp}(x:y) \to \mathsf{f}^{\sharp}(\mathsf{g}(y, [])) & 8: \quad \mathsf{g}^{\sharp}(x:y, z) \to \mathsf{g}^{\sharp}(y, x:z) \end{aligned}$$

and $\mathcal{U}(\mathcal{P}) = \{3,4\}$. It is not difficult to show $\operatorname{comp}(n, \mathcal{T}_{\mathsf{b}}^{\sharp}, \stackrel{i}{\to}_{\mathcal{P}/\mathcal{U}(\mathcal{P})}) = \mathsf{O}(n)$ with a 1-dimensional RMI. Moreover, the $\mu_{i}^{\mathcal{P}\cup\mathcal{U}(\mathcal{P})}$ -monotone 1-dimensional RMI \mathcal{A} with

$$\begin{bmatrix} \mathbf{a}_{\mathcal{A}} = 0 & \mathbf{a}_{\mathcal{A}}(x, y) = y + 1 \\ \mathbf{b}_{\mathcal{A}}(x) = \mathbf{f}_{\mathcal{A}}^{\sharp}(x) = x & \mathbf{g}_{\mathcal{A}}^{\sharp}(x, y) = 0 \\ \mathbf{c}_{\mathcal{A}} = \mathbf{d}_{\mathcal{A}} = 0 \\ \mathbf{c}_{\mathcal{A}} = \mathbf{d}_{\mathcal{A}} = 0 \end{bmatrix}$$

is compatible with $\mathcal{U}(\mathcal{P})$. If Lemma 6.11 would be applicable without its well-definedness, linear innermost runtime complexity of \mathcal{R} would be concluded falsely. Note that $\Delta(\mathcal{A}, \mathcal{P})$ is *not* well-defined on \mathbb{N} due to pair 6.

Corollary 6.14. Let \mathcal{R} be a TRS, \mathcal{P} the set of weak (innermost) dependency pairs, and μ be the (innermost) usable replacement map. Suppose \mathcal{B} is a RMI such that $(\succeq_{\mathcal{B}}, \succ_{\mathcal{B}})$ forms a μ -complexity pair with $\mathcal{U}(\mathcal{P}) \subseteq \succeq_{\mathcal{B}}$ and $\mathcal{P} \subseteq \succ_{\mathcal{B}}$. Further, suppose \mathcal{A} is an adequate μ -monotone RMI such that $\Delta(\mathcal{A}, \mathcal{P})$ is well-defined on \mathbb{N} and \mathcal{P} is compatible with $\mathcal{U}(\mathcal{P})$.

Then the (innermost) runtime complexity function $\operatorname{rc}_{\mathcal{R}}^{(i)}$ with respect to \mathcal{R} is polynomial. Here the degree of the polynomial is given by the maximum of the degrees of the used RMIs.

Let \mathcal{A} be an RMI as in the corollary. In order to verify that $\Delta(\mathcal{A}, \mathcal{P})$ is well-defined, we use the following simple trick in the implementation. Let $l \to r \in \mathcal{P}$ and let k denotes the cardinality of $\operatorname{Var}(l) \supseteq \operatorname{Var}(r)$. Recall the existence of matrices (over \mathbb{N}) L_1, \ldots, L_k , R_1, \ldots, R_k and vectors \vec{l}, \vec{r} such that $[\alpha]_{\mathcal{A}}(l) \doteq [\alpha]_{\mathcal{A}}(r) = \sum_{i=1}^k (R_i \doteq L_i)\alpha(x_i) + (\vec{r} \doteq \vec{l})$. Then $\Delta(\mathcal{A}, \mathcal{P})$ is well-defined if $(R_i \doteq L_i) \leq \mathbf{0}$.

7 Weak Dependency Graphs

In this section we extend the above refinements by revisiting dependency graphs in the context of complexity analysis. Let $\mathcal{P} = \mathsf{WDP}(\mathcal{R}_{\mathsf{div}})$ and recall the derivation over $\mathcal{P} \cup \mathcal{U}(\mathcal{P})$ on page 17. Looking more closely at this derivation we observe that we do not make use of all weak dependency pairs in \mathcal{P} , but we only employ the pairs 7 and 8:

$$4 \div^{\sharp} 2 \rightarrow_{\{8\}/\mathcal{U}(\mathcal{P})} 2 \div^{\sharp} 2 \rightarrow_{\{8\}/\mathcal{U}(\mathcal{P})} 0 \div^{\sharp} 2 \rightarrow_{\{7\}/\mathcal{U}(\mathcal{P})} \mathsf{c}.$$

Therefore it is a natural idea to modularise our complexity analysis and apply the previously obtained techniques only to those pairs that are relevant. Dependencies among weak dependency pairs are formulated by the notion of weak dependency graphs, which is an easy variant of *dependency graphs* [6].

Definition 7.1. Let \mathcal{R} be a TRS over a signature \mathcal{F} and let \mathcal{P} be the set of weak, weak innermost, or (standard) dependency pairs. The nodes of the *weak dependency graph* $\mathsf{WDG}(\mathcal{R})$, *weak innermost dependency graph* $\mathsf{WIDG}(\mathcal{R})$, or *dependency graph* $\mathsf{DG}(\mathcal{R})$ are the elements of \mathcal{P} and there is an arrow from $s \to t$ to $u \to v$ if and only if there exist a context C and substitutions $\sigma, \tau \colon \mathcal{V} \to \mathcal{T}(\mathcal{F}, \mathcal{V})$ such that $t\sigma \to^* C[u\tau]$, where \to denotes $\to_{\mathcal{R}}$ or $\overset{i}{\to}_{\mathcal{R}}$ depending on whether $\mathcal{P} = \mathsf{WDP}(\mathcal{R}), \mathcal{P} = \mathsf{DP}(\mathcal{R})$, or $\mathcal{P} = \mathsf{WIDP}(\mathcal{R})$, respectively.

Example 7.2 (continued from Example 5.3). The weak dependency graph $WDG(\mathcal{R}_{div})$ has the following form.



Since weak dependency graphs represent call graphs of functions, grouping mutual parts helps analysis. A graph is called *strongly connected* if any node is connected with every other node by a (possibly empty) path. A *strongly connected component* (*SCC* for short) is a maximal strongly connected subgraph.⁴

Definition 7.3. Let \mathcal{G} be a graph, let \equiv denote the equivalence relation induced by SCCs, and let \mathcal{P} be a SCC in \mathcal{G} . Consider the *congruence graph* \mathcal{G}_{\equiv} induced by the equivalence relation \equiv . The set of all source nodes in \mathcal{G}_{\equiv} is denoted by $Src(\mathcal{G}_{\equiv})$. Let $\mathcal{K} \in \mathcal{G}_{\equiv}$ and let \mathcal{C} denote the SCC represented by \mathcal{K} . Then we write $l \to r \in \mathcal{K}$ if $l \to r \in \mathcal{C}$. For nodes \mathcal{K} and \mathcal{L} in \mathcal{G}_{\equiv} we write $\mathcal{K} \rightsquigarrow \mathcal{L}$, if \mathcal{K} and \mathcal{L} are connected by an edge. The reflexive (transitive, reflexive-transitive) closure of \rightsquigarrow is denoted as $\rightsquigarrow^{=} (\rightsquigarrow^{+}, \rightsquigarrow^{*})$.

Example 7.4 (continued from Example 7.2). Let \mathcal{G} denote $WDG(\mathcal{R}_{div})$. There are 4 SCCs in \mathcal{G} : {5}, {6}, {7}, and {8}. Thus the congruence graph \mathcal{G}_{\equiv} has the following form:

$$6 \longrightarrow 5 \qquad 8 \longrightarrow 7$$

Here $Src(\mathcal{G}_{\equiv}) = \{\{6\}, \{8\}\}.$

⁴ We use SCCs in the standard graph theoretic sense, while in the literature SCCs are sometimes defined as *maximal cycles* (e.g. [24, 25, 11]). This alternative definition is of limited use in our context.

Example 7.5. Consider the TRS \mathcal{R}_{gcd} which computes the greatest common divisor.⁵

The set $WDP(\mathcal{R}_{gcd})$ consists of the next ten weak dependency pairs:

The congruence graph \mathcal{G}_{\equiv} of $\mathcal{G} := \mathsf{WDG}(\mathcal{R}_{\mathsf{gcd}})$ has the following form:

$$11 \leftarrow 13 \longrightarrow 12$$
 $15 \longrightarrow 14$ $\{18, 19, 20\} \longrightarrow 16$ 17

Here $\mathsf{Src}(\mathcal{G}_{\equiv}) = \{\{13\}, \{15\}, \{17\}, \{18, 19, 20\}\}.$

The main result in this section is stated as follows: Let \mathcal{R} be a TRS, $\mathcal{P} = \mathsf{WDP}(\mathcal{R})$, $\mathcal{G} = \mathsf{WDG}(\mathcal{R})$, and furthermore

 $\mathsf{L}(t) := \max\{\mathsf{dh}(t, \stackrel{(i)}{\longrightarrow}_{\mathcal{Q} \cup \mathcal{U}(\mathcal{Q})}) \mid (\mathcal{P}_1, \dots, \mathcal{P}_k) \text{ is a path in } \mathcal{G}_{\equiv} \text{ and } \mathcal{P}_1 \in \mathsf{Src}(\mathcal{G}_{\equiv})\} ,$

where $\mathcal{Q} = \bigcup_{i=1}^{k} \mathcal{P}_i$. Then, $dh(t, \to_{\mathcal{R}}) = O(L(t))$ holds for all basic term t. This means that one may decompose $\mathcal{P} \cup \mathcal{U}(\mathcal{P})$ into several smaller fragments and analyse these fragments separately.

Reconsider the derivation on page 23. The only dependency pairs are from the set $\{7, 8\}$. Observe that the order these pairs are applied is representable by the path ($\{8\}, \{7\}$) in the congruence graph. This observation is cast into the following definition.

Definition 7.6. Let \mathcal{P} be the set of weak (innermost) dependency pairs and let \mathcal{G} denote the weak (innermost) dependency graph. Suppose $A: s \xrightarrow{(i)}_{\mathcal{P}/\mathcal{U}(\mathcal{P})} t$ denote a derivation, such that $s \in \mathcal{T}_{b}^{\sharp}$. If A can be written in the following form:

$$s \xrightarrow{(i)} {}^{*}_{\mathcal{P}_{1}/\mathcal{U}(\mathcal{P})} \cdots \xrightarrow{(i)} {}^{*}_{\mathcal{P}_{k}/\mathcal{U}(\mathcal{P})} t$$

then A is based on the sequence of nodes $(\mathcal{P}_1, \ldots, \mathcal{P}_k)$ (in \mathcal{G}_{\equiv}).

The next lemma is an easy generalisation of the above example.

⁵ This is Example 3.6a in Arts and Giesl's collection of TRSs [14].

Lemma 7.7. Let \mathcal{R} be a TRS, let \mathcal{P} be the set of weak (innermost) dependency pairs and let \mathcal{G} denote the weak (innermost) dependency graph. Suppose that all compound symbols are nullary. Then any derivation $A: s \xrightarrow{(i)}_{\mathcal{P}/\mathcal{U}(\mathcal{P})} t$ such that $s \in \mathcal{T}_{b}^{\sharp}$ is based on a path in \mathcal{G}_{\equiv} .

From Lemma 7.7 we see that the above mentioned modularity result easily follows as long as the arity of the compound symbols is restricted. We lift the assumption that all compound symbols are nullary. Perhaps surprisingly this generalisation complicates the matter. As exemplified by the next example, Lemma 7.7 fails if there exist non-nullary compound symbols.

Example 7.8. Consider the TRS $\mathcal{R} = \{f(0) \to a, f(s(x)) \to b(f(x), f(x))\}$. The set WDP(\mathcal{R}) consists of the two weak dependency pairs: 1: $f^{\sharp}(0) \to c$ and 2: $f^{\sharp}(s(x)) \to d(f^{\sharp}(x), f^{\sharp}(x))$. The corresponding congruence graph only contains the single edge from $\{2\}$ to $\{1\}$. Writing t_n for $f^{\sharp}(s^n(0))$, we have the sequence

$$t_{2} \rightarrow^{2}_{\{2\}} \mathsf{d}(\mathsf{d}(t_{0}, t_{0}), t_{1}) \rightarrow_{\{1\}} \mathsf{d}(\mathsf{d}(\mathsf{c}, t_{0}), t_{1}) \\ \rightarrow_{\{2\}} \mathsf{d}(\mathsf{c}(\mathsf{c}, t_{0}), \mathsf{d}(t_{0}, t_{0})) \rightarrow^{3}_{\{1\}} \mathsf{d}(\mathsf{d}(\mathsf{c}, \mathsf{c}), \mathsf{d}(\mathsf{c}, \mathsf{c})) \ .$$

whereas $(\{2\}, \{1\}, \{2\}, \{1\})$ is not a path in the graph.

Note that the derivation in Example 7.8 can be reordered (without affecting its length) such that the derivation becomes based on the path ($\{2\}, \{1\}$). More generally, we observe that a weak (innermost) dependency pair containing an *m*-ary (m > 1) compound symbol can induce *m* independent derivations. This allows us to reorder (sub-)derivations. We show this via the following sequence of lemmas.

Let \mathcal{R} be a TRS, let \mathcal{P} denote the set of weak (innermost) dependency pairs, and let \mathcal{G} denote the weak (innermost) dependency graph. The set $\mathcal{T}_{\mathsf{c}}^{\sharp}$ is inductively defined as follows (i) $\mathcal{T}^{\sharp} \cup \mathcal{T} \subseteq \mathcal{T}_{\mathsf{c}}^{\sharp}$, where $\mathcal{T}^{\sharp} = \{t^{\sharp} \mid t \in \mathcal{T}\}$ and (ii) $c(t_1, \ldots, t_n) \in \mathcal{T}_{\mathsf{c}}^{\sharp}$, whenever $t_1, \ldots, t_n \in \mathcal{T}_{\mathsf{c}}^{\sharp}$ and c a compound symbol. The next lemma formalises an easy observation.

Lemma 7.9. Let C be a set of nodes in G and let $A: t = t_0 \xrightarrow{(i)}_{C/U(\mathcal{P})} t_n$ denote a derivation based on C with $t \in \mathcal{T}_c^{\sharp}$. Then A has the following form: $t = t_0 \xrightarrow{(i)}_{C/U(\mathcal{P})} t_1 \xrightarrow{(i)}_{C/U(\mathcal{P})} \cdots \xrightarrow{(i)}_{C/U(\mathcal{P})} t_n$ where each $t_i \in \mathcal{T}_c^{\sharp}$.

A key is that consecutive two weak dependency pairs may be swappable.

Lemma 7.10. Let \mathcal{K} and \mathcal{L} denote two different nodes in \mathcal{G}_{\equiv} such that there is no edge from \mathcal{K} to \mathcal{L} . Let $s \in \mathcal{T}_{c}^{\sharp}$ and suppose the existence of a derivation A of the following form:

$$s \xrightarrow{(i)} \mathcal{K}/\mathcal{U}(\mathcal{P}) \cdot \xrightarrow{(i)} \mathcal{L}/\mathcal{U}(\mathcal{P}) t$$
.

Then there exists a derivation B

$$s \xrightarrow{(i)} \mathcal{L}/\mathcal{U}(\mathcal{P}) \cdot \xrightarrow{(i)} \mathcal{K}/\mathcal{U}(\mathcal{P}) t$$
,

such that |A| = |B|.

Proof. We only show the full rewriting case since the innermost case is analogous. According to Lemma 7.9 an arbitrary terms u reachable from s belongs to \mathcal{T}_{c}^{\sharp} . Writing $C\langle u_{1}, \ldots, u_{i}, \ldots, u_{m} \rangle_{\mathcal{F} \cup \mathcal{F}^{\sharp}}$ for u, the *m*-hole context C consists of compound symbols and variables, $u_{1}, \ldots, u_{m} \in \mathcal{T} \cup \mathcal{T}^{\sharp}$. Therefore, A can be written in the following form:

$$s \rightarrow_{\mathcal{U}(\mathcal{P})}^{n_1} C\langle u_1, \dots, u_i, \dots, u_m \rangle_{\mathcal{F} \cup \mathcal{F}^{\sharp}} =: u$$

$$\rightarrow_{\mathcal{L}} C[u_1, \dots, u'_i, \dots, u_m]$$

$$\rightarrow_{\mathcal{U}(\mathcal{P})}^{n_2} C[v_1, \dots, v_i, \dots, v_j, \dots, v_m]$$

$$\rightarrow_{\mathcal{K}} C[v_1, \dots, v_i, \dots, v'_j, \dots, v_m] \rightarrow_{\mathcal{U}(\mathcal{P})}^{n_3} t$$

with $u'_i \to^k_{\mathcal{U}(\mathcal{P})} v_i$. Here $i \neq j$ holds, because i = j induces $\mathcal{L} \rightsquigarrow \mathcal{K}$. Easy induction on n_2 shows

$$s \rightarrow_{\mathcal{U}(\mathcal{P})}^{n_1} u = C[u_1, \dots, u_i, \dots, u_j, \dots, u_m]$$

$$\rightarrow_{\mathcal{U}(\mathcal{P})}^{n_2-k} C[v_1, \dots, u_i, \dots, v_j, \dots, v_m]$$

$$\rightarrow_{\mathcal{K}} C[v_1, \dots, u_i, \dots, v'_j, \dots, v_m]$$

$$\rightarrow_{\mathcal{L}} C[v_1, \dots, u'_i, \dots, v'_j, \dots, v_m]$$

$$\rightarrow_{\mathcal{U}(\mathcal{P})}^k C[v_1, \dots, v_i, \dots, v'_j, \dots, v_m] \rightarrow_{\mathcal{U}(\mathcal{P})}^{n_3} t ,$$

which is the desired derivation B.

The next lemma states that reordering is partly possible.

Lemma 7.11. Let $s \in \mathcal{T}_{c}^{\sharp}$, and let $A: s \xrightarrow{(i)}_{\mathcal{P}/\mathcal{U}(\mathcal{P})} t$ be a derivation based on a sequence of nodes $(\mathcal{P}_{1}, \ldots, \mathcal{P}_{k})$ such that $\mathcal{P}_{1} \in \operatorname{Src}(\mathcal{G}_{\equiv})$, and let $(\mathcal{Q}_{1}, \ldots, \mathcal{Q}_{\ell})$ be a path in \mathcal{G}_{\equiv} with $\{\mathcal{P}_{1}, \ldots, \mathcal{P}_{k}\} = \{\mathcal{Q}_{1}, \ldots, \mathcal{Q}_{\ell}\}$. Then there exists a derivation $B: s \xrightarrow{(i)}_{\mathcal{P}/\mathcal{U}(\mathcal{P})} t$ based on $(\mathcal{Q}_{1}, \ldots, \mathcal{Q}_{\ell})$ such that |A| = |B| and $\mathcal{P}_{1} = \mathcal{Q}_{1}$.

Proof. According to Lemma 7.9, for any derivation A

$$s \xrightarrow{(i)} {}^{*}_{\mathcal{P}_{1}/\mathcal{U}(\mathcal{P})} \cdots \xrightarrow{(i)} {}^{*}_{\mathcal{P}_{n}/\mathcal{U}(\mathcal{P})} t$$

if $\mathcal{P}_i \rightsquigarrow \mathcal{P}_{i+1}$ does not hold, there is a derivation B

$$s \xrightarrow{(i)} \stackrel{*}{\mathcal{P}_{1}/\mathcal{U}(\mathcal{P})} \cdots \xrightarrow{(i)} \stackrel{*}{\mathcal{P}_{i+1}/\mathcal{U}(\mathcal{P})} \cdot \xrightarrow{(i)} \stackrel{*}{\mathcal{P}_{i}/\mathcal{U}(\mathcal{P})} \cdots \xrightarrow{(i)} \stackrel{*}{\mathcal{P}_{n}/\mathcal{U}(\mathcal{P})} t ,$$

with |A| = |B|. By assumption (Q_1, \ldots, Q_ℓ) is a path, whence we obtain $Q_1 \rightsquigarrow \cdots \rightsquigarrow Q_\ell$. By performing bubble sort with respect to \rightsquigarrow^+ , A is transformed into the derivation B:

$$s \xrightarrow{(i)} {}^{*}_{\mathcal{Q}_1/\mathcal{U}(\mathcal{P})} \cdots \xrightarrow{(i)} {}^{*}_{\mathcal{Q}_m/\mathcal{U}(\mathcal{P})} t$$
,

such that |A| = |B|.

The next example shows that there is a derivation that cannot be transformed into a derivation based on a path.

Example 7.12. Consider the TRS $\mathcal{R} = \{f \to b(g,h), g \to a, h \to a\}$. Thus $WDP(\mathcal{R})$ consists of three dependency pairs: $1: f^{\sharp} \to c(g^{\sharp}, h^{\sharp}), 2: g^{\sharp} \to d$, and $3: h^{\sharp} \to e$. Let $\mathcal{P} := WDP(\mathcal{R})$ and let $\mathcal{G} := WDG(\mathcal{R})$. Note that \mathcal{G}_{\equiv} are identical to \mathcal{G} . We witness that the derivation

$$f^{\sharp} \rightarrow_{\mathcal{P}} c(g^{\sharp}, h^{\sharp}) \rightarrow_{\mathcal{P}} c(d, h^{\sharp}) \rightarrow_{\mathcal{P}} c(d, e)$$

is based neither on the path $(\{1\}, \{2\})$, nor on the path $(\{1\}, \{3\})$.

Lemma 7.11 shows that we can reorder a given derivation A that is based on a sequence of nodes that would in principle form a path in the congruence graph \mathcal{G}_{\equiv} . The next lemma shows that we can guarantee that any derivation is based on sequence of different paths.

Lemma 7.13. Let $s \in \mathcal{T}_{\mathsf{c}}^{\sharp}$ and let $A: s \xrightarrow[\mathcal{P}/\mathcal{U}(\mathcal{P})]{}^{\mathsf{v}} t$ be a derivation based on $(\mathcal{P}_1, \ldots, \mathcal{P}_k, \mathcal{Q}_1, \ldots, \mathcal{Q}_\ell)$, such that $(\mathcal{P}_1, \ldots, \mathcal{P}_k)$ and $(\mathcal{Q}_1, \ldots, \mathcal{Q}_\ell)$ form two disjoint paths in \mathcal{G} . Then there exists a derivation $B: s \xrightarrow[\mathcal{P}/\mathcal{U}(\mathcal{P})]{}^{\mathsf{v}} t$ based on the sequence of nodes $(\mathcal{Q}_1, \ldots, \mathcal{Q}_\ell, \mathcal{P}_1, \ldots, \mathcal{P}_k)$ such that |A| = |B|.

Proof. The lemma follows by an adaptation of the technique in the proof of Lemma 7.11. \Box

Lemma 7.13 shows that the maximal length of any derivation only differs from the maximal length of any derivation based on a path by a linear factor, depending on the size of the congruence graph \mathcal{G}_{\equiv} . We arrive at the main result of this section. Recall the definition of $L(\cdot)$ on page 24.

Theorem 7.14. Let \mathcal{R} be a TRS and \mathcal{P} the set of weak (innermost) dependency pairs. Then, $dh(t, \stackrel{(i)}{\longrightarrow}_{\mathcal{R}}) = O(L(t))$ holds for all $t \in \mathcal{T}_{h}^{\sharp}$.

Proof. Let a denotes the maximum arity of compound symbols and K denotes the number of SCCs in the weak (innermost) dependency graph \mathcal{G} . We show $\mathsf{dh}(s, \stackrel{(i)}{\to}_{\mathcal{R}}) \leq a^{K} \cdot \mathsf{L}(s)$ holds for all $s \in \mathcal{T}_{\mathsf{b}}^{\sharp}$. Theorem 5.12 yields that $\mathsf{dh}(s, \stackrel{(i)}{\to}_{\mathcal{R}}) = \mathsf{dh}(s, \to)$, where \to either denotes $\to_{\mathcal{P} \cup \mathcal{U}(\mathcal{P})}$ or $\stackrel{i}{\to}_{\mathcal{P} \cup \mathcal{U}(\mathcal{P})}$.

Let $A: s \to^* t$ be a derivation over $\mathcal{P} \cup \mathcal{U}(\mathcal{P})$ such that $s \in \mathcal{T}_b^{\sharp}$. Then A is based on a sequence of nodes in the congruence graph \mathcal{G}_{\equiv} such that there exists a maximal (with respect to subset inclusion) components of \mathcal{G}_{\equiv} that includes all these nodes. Let T denote this maximal component. T forms a directed acyclic graph. In order to (over-)estimate the number of nodes in this graph we can assume without loss of generality that T is a tree with root in $\operatorname{Src}(\mathcal{G}_{\equiv})$. Note that K bounds the height of this tree. Thus the number of nodes in the component T is less than

$$\frac{a^K - 1}{a - 1} \leqslant a^K \,.$$

Due to Lemma 7.13 the derivation A is conceivable as a sequence of subderivations based on paths in \mathcal{G}_{\equiv} . As the number of nodes in T is bounded from above by a^K , there exist at most be a^K different paths through T.

Hence in order to estimate |A|, it suffices to estimate the length of any subderivation B of A, based on a specific path. Let $(\mathcal{P}_1, \ldots, \mathcal{P}_k)$ be a path in \mathcal{P}_{\equiv} such that $\mathcal{P}_1 \in \operatorname{Src}(\mathcal{G}_{\equiv})$ and let $B: u \to^n v$, denote a derivation based on this path. Let $\mathcal{Q} := \bigcup_{i=1}^k \mathcal{P}_i$. By Definition 7.6 and the definition of usable rules, the derivation B can be written as:

$$u = u_0 \xrightarrow{(i)}_{\mathcal{P}_1/\mathcal{U}(\mathcal{Q})} u_{n_1} \xrightarrow{(i)}_{\mathcal{P}_2/\mathcal{U}(\mathcal{Q})} \cdots \xrightarrow{(i)}_{\mathcal{P}_k/\mathcal{U}(\mathcal{Q})} u_n = v ,$$

where $u \in \mathcal{T}_{\mathsf{b}}^{\sharp}$ each $u_i \in \mathcal{T}_{\mathsf{c}}^{\sharp}$. Hence *B* is contained in $u \xrightarrow{(i)} \mathcal{Q}_{\cup \mathcal{U}(\mathcal{Q})}^* v$ and thus $|B| \leq \mathsf{L}(u)$ by definition.

As the length of a derivation B based on a specific path can be estimated by L(s), we obtain that the length of an arbitrary derivation is less than $a^K \cdot L(s)$. This completes the proof of the theorem.

Corollary 7.15. Let \mathcal{R} be a TRS and let \mathcal{G} denote the weak (innermost) dependency graph. For every path $\overline{P} := (\mathcal{P}_1, \ldots, \mathcal{P}_k)$ in \mathcal{G}_{\equiv} such that $\mathcal{P}_1 \in \text{Src}(\mathcal{G}_{\equiv})$, we set $\mathcal{Q} := \bigcup_{i=1}^k \mathcal{P}_i$ and suppose

- 1) there exist a $\mu_{f}^{\mathcal{Q}\cup\mathcal{U}(\mathcal{Q})}$ -monotone ($\mu_{i}^{\mathcal{Q}\cup\mathcal{U}(\mathcal{Q})}$ -monotone) and adequate RMI $\mathcal{A}_{\bar{P}}$ that admits the weight gap $\Delta(\mathcal{A}_{\bar{P}}, \mathcal{Q})$ on \mathcal{T}_{b}^{\sharp} and $\mathcal{A}_{\bar{P}}$ is compatible with the usable rules $\mathcal{U}(\mathcal{Q})$,
- 2) there exists a $\mu_{f}^{\mathcal{Q}\cup\mathcal{U}(\mathcal{Q})}$ -monotone ($\mu_{i}^{\mathcal{Q}\cup\mathcal{U}(\mathcal{Q})}$ -monotone) RMI $\mathcal{B}_{\bar{P}}$ such that ($\succeq_{\mathcal{B}_{\bar{P}}}, \succeq_{\mathcal{B}_{\bar{P}}}$) forms a complexity pair for $\mathcal{P}_{k}/\mathcal{P}_{1}\cup\cdots\cup\mathcal{P}_{k-1}\cup\mathcal{U}(\mathcal{Q})$, and

Then the (innermost) runtime complexity of a TRS \mathcal{R} is polynomial. Here the degree of the polynomial is given by the maximum of the degrees of the used RMIs.

Proof. We restrict our attention to weak dependency pairs and full rewriting. First observe that the assumptions imply that any basic term $t \in \mathcal{T}_{\mathsf{b}}$ is terminating with respect to \mathcal{R} . Let \mathcal{P} be the set of weak dependency pairs. (Note that $\mathcal{P} \supseteq \mathcal{Q}$.) By Lemma 5.11 any infinite derivation with respect to \mathcal{R} starting in t can be translated into an infinite derivation with respect to $\mathcal{U}(\mathcal{P}) \cup \mathcal{P}$. Moreover, as the number of paths in \mathcal{G}_{\equiv} is finite, there exist a path $(\mathcal{P}_1, \ldots, \mathcal{P}_k)$ in \mathcal{G}_{\equiv} and an infinite rewrite sequence based on this path. This is a contradiction. Hence we can employ Theorem 6.5 in the following.

Let $(\mathcal{P}_1, \ldots, \mathcal{P}_k)$ be an arbitrary, but fixed path in the congruence graph \mathcal{G}_{\equiv} , let $\mathcal{Q} = \bigcup_{i=1}^k \mathcal{P}_i$, and let *d* denote the maximum of the degrees of the used RMIs. Due to Theorem 6.5 there exists $c \in \mathbb{N}$ such that:

$$\mathsf{dh}(t^{\sharp}, \to_{\mathcal{Q}\cup\mathcal{U}(\mathcal{Q})}) \leqslant (1 + \Delta(\mathcal{A}_{\bar{P}}, \mathcal{Q})) \cdot \mathsf{dh}(t^{\sharp}, \to_{\mathcal{Q}/\mathcal{U}(\mathcal{Q})}) + c \cdot |t|^{d} .$$

Due to Theorem 7.14 it suffices to consider a derivation A based on the path $(\mathcal{P}_1, \ldots, \mathcal{P}_k)$. Suppose $A: s \to_{\mathcal{Q}/\mathcal{U}(\mathcal{Q})}^n t$. Then A can be represented as follows:

$$s = s_0 \to_{\mathcal{P}_1/\mathcal{U}(\mathcal{P}_1)}^{n_1} s_{n_1} \to_{\mathcal{P}_2/\mathcal{U}(\mathcal{P}_1)\cup\mathcal{U}(\mathcal{P}_2)}^{n_2} \cdots \to_{\mathcal{P}_k/\mathcal{U}(\mathcal{P}_1)\cup\cdots\cup\mathcal{U}(\mathcal{P}_k)}^{n_k} s_n = t ,$$

such that $n = \sum_{i=1}^{k} n_i$. It is sufficient to bound each n_i from the above. Fix $i \in \{1, \ldots, k\}$. Consider the subderivation

$$A': s = s_0 \to_{\mathcal{P}_1/\mathcal{U}(\mathcal{P}_1)}^{n_1} s_{n_1} \cdots \to_{\mathcal{P}_k/\mathcal{U}(\mathcal{P}_1) \cup \cdots \cup \mathcal{U}(\mathcal{P}_i)}^{n_i} s_{n_i}$$

Then A' is contained in $A'': s \to_{\mathcal{P}_1 \cup \cdots \cup \mathcal{P}_{i-1} \cup \mathcal{U}(\mathcal{P}_1) \cup \cdots \cup \mathcal{U}(\mathcal{P}_i)}^{n_i} \cdot \to_{\mathcal{P}_k/\mathcal{U}(\mathcal{P}_1) \cup \cdots \cup \mathcal{U}(\mathcal{P}_i)}^{n_i} s_{n_i}$. Let $\hat{P}_i := (\mathcal{P}_1, \ldots, \mathcal{P}_i)$. By assumption there exists a μ -monotone complexity pair $(\geq_{\mathcal{B}_{\hat{P}_i}}, \succ_{\mathcal{B}_{\hat{P}_i}})$ such that $\mathcal{P}_1 \cup \cdots \cup \mathcal{P}_{i-1} \cup \mathcal{U}(\mathcal{P}_1 \cup \cdots \cup \mathcal{P}_i) \subseteq \geq_{\mathcal{B}_{\hat{P}_i}}$ and $\mathcal{P}_i \subseteq \succ_{\mathcal{B}_{\hat{P}_i}}$. Hence, we obtain $n_i \leq ([\alpha_0]_{\mathcal{B}_{\hat{P}_i}}(s))_1$ and in sum $n \leq k \cdot |s|^d$. Finally, defining the polynomial p as follows:

$$p(x) := (1 + \Delta(\mathcal{A}_{\bar{P}}, \mathcal{Q})) \cdot k \cdot x^d + c \cdot x^d ,$$

we conclude $\mathsf{dh}(t^{\sharp}, \to_{\mathcal{Q}\cup\mathcal{U}(\mathcal{Q})}) \leq p(|t|)$. Note that the polynomial p depends only on the algebras $\mathcal{A}_{\bar{P}}$ and $\mathcal{B}_{\hat{P}_1}, \ldots, \mathcal{B}_{\bar{P}_k}$.

As the path $(\mathcal{P}_1, \ldots, \mathcal{P}_k)$ was chosen arbitrarily, there exists a polynomial q, depending only on the employed RMIs such that $\mathsf{L}(t) \leq q(|t|)$. Thus the corollary follows due to Theorem 7.14.

Let t be an arbitrary term. By definition the set in L(t) may consider $2^{O(n)}$ -many paths, where n denotes the number of nodes in \mathcal{G}_{\equiv} . However, it suffices to restrict the definition on page 24 to maximal paths. For this refinement L(t) contains at most n^2 paths. This fact we employ in implementing the WDG method.

Example 7.16 (continued from Example 7.5). For $WDG(\mathcal{R}_{gcd})_{\equiv}$ the above set consists of 8 paths: ({13}), ({13}, {11}), ({13}, {12}), ({15}), ({15}, {14}), ({17}), ({18, 19, 20}), and ({18, 19, 20}, {16}). In the following we only consider the last three paths, since all other paths are similarly handled.

- Consider ({17}). Note $\mathcal{U}(\{17\}) = \emptyset$. By taking an arbitrary SLI \mathcal{A} and the linear restricted interpretation \mathcal{B} with $gcd_{\mathcal{B}}^{\sharp}(x, y) = x$ and $s_{\mathcal{B}}(x) = x + 1$, we have $\emptyset \subseteq >_{\mathcal{A}}$, $\emptyset \subseteq \geq_{\mathcal{B}}$, and $\{17\} \subseteq >_{\mathcal{B}}$.
- Consider ({18, 19, 20}). Note $\mathcal{U}(\{18, 19, 20\}) = \{1, \ldots, 5\}$. The following RMI \mathcal{A} is adequate for ({18, 19, 20}) and strictly monotone on $\mu_{f}^{\mathcal{P} \cup \mathcal{U}(\mathcal{P})}$. The presentation of \mathcal{A} is succinct as only the signature of the usable rules {1, ..., 5} is of interest.

$$\operatorname{true}_{\mathcal{A}} = \operatorname{false}_{\mathcal{A}} = \mathbf{0}_{\mathcal{A}} = \vec{0} \qquad \qquad \mathbf{s}_{\mathcal{A}}(\vec{x}) = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \vec{x} + \begin{pmatrix} 3 \\ 1 \end{pmatrix}$$
$$\leqslant_{\mathcal{A}}(\vec{x}, \vec{y}) = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \vec{y} + \begin{pmatrix} 1 \\ 3 \end{pmatrix} \qquad \qquad -_{\mathcal{A}}(\vec{x}, \vec{y}) = \vec{x} + \begin{pmatrix} 2 \\ 3 \end{pmatrix} .$$

Further, consider the RMI \mathcal{B} giving rise to the complexity pair $(\succeq_{\mathcal{B}}, \succeq_{\mathcal{B}})$.

$$\begin{split} \mathbf{0}_{\mathcal{B}} &= \mathsf{true}_{\mathcal{B}} = \mathsf{false}_{\mathcal{B}} = \leqslant_{\mathcal{B}}(\vec{x}, \vec{y}) = 0\\ \mathbf{s}_{\mathcal{B}}(\vec{x}) &= \begin{pmatrix} 1 & 3 \\ 0 & 0 \end{pmatrix} \vec{x} + \begin{pmatrix} 3 \\ 0 \end{pmatrix} \qquad -_{\mathcal{B}}(\vec{x}, \vec{y}) \qquad = \begin{pmatrix} 1 & 0 \\ 2 & 2 \end{pmatrix} \vec{x} + \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} \\ \mathsf{if}_{\mathsf{gcd}}_{\mathcal{B}}^{\sharp}(x, y, z) &= \begin{pmatrix} 3 & 0 \\ 0 & 0 \end{pmatrix} \vec{y} + \begin{pmatrix} 3 & 0 \\ 0 & 0 \end{pmatrix} \vec{z} \\ \mathsf{gcd}_{\mathcal{B}}^{\sharp}(x, y) &= \begin{pmatrix} 3 & 0 \\ 0 & 0 \end{pmatrix} \vec{x} + \begin{pmatrix} 3 & 0 \\ 0 & 0 \end{pmatrix} \vec{y} + \begin{pmatrix} 2 \\ 0 \end{pmatrix} \ . \end{split}$$

We obtain $\{1, \ldots, 5\} \subseteq \succ_{\mathcal{A}}, \{1, \ldots, 5\} \subseteq \succcurlyeq_{\mathcal{B}}, \text{ and } \{18, 19, 20\} \subseteq \succ_{\mathcal{B}}.$

• Consider ({18, 19, 20}, {16}). Note $\mathcal{U}(\{16\}) = \emptyset$. By taking the same \mathcal{A} and also \mathcal{B} as above, we have $\{1, \ldots, 5\} \subseteq \succ_{\mathcal{A}}, \{1, \ldots, 5, 18, 19, 20\} \subseteq \succcurlyeq_{\mathcal{B}}, \text{ and } \{16\} \subseteq \succ_{\mathcal{B}}.$

Thus, all path constraints are handled by suitably defined RMIs of dimension 2. Hence, the runtime complexity function of \mathcal{R}_{gcd} is at most quadratic, which is unfortunately not optimal, as $rc_{\mathcal{R}_{gcd}}$ is linear.

Corollary 7.15 is more powerful than Corollary 6.14. We illustrate it with a small example.

Example 7.17. Consider the TRS \mathcal{R}

$$\mathsf{f}(\mathsf{a},\mathsf{s}(x),y)\to\mathsf{f}(\mathsf{a},x,\mathsf{s}(y))\qquad\qquad \mathsf{f}(\mathsf{b},x,\mathsf{s}(y))\to\mathsf{f}(\mathsf{b},\mathsf{s}(x),y)\;.$$

Its weak dependency pairs $\mathsf{WDP}(\mathcal{R})$ are

1:
$$f^{\sharp}(\mathsf{a},\mathsf{s}(x),y) \to f^{\sharp}(\mathsf{a},x,\mathsf{s}(y))$$
 2: $f^{\sharp}(\mathsf{b},x,\mathsf{s}(y)) \to f^{\sharp}(\mathsf{b},\mathsf{s}(x),y)$.

The corresponding congruence graph consists of the two isolated nodes $\{1\}$ and $\{2\}$. It is not difficult to find suitable 1-dimensional RMIs for the nodes, and therefore $\mathsf{rc}_{\mathcal{R}}(n) = \mathsf{O}(n)$ is concluded. On the other hand, it can be verified that the linear runtime complexity cannot be obtained by Corollary 6.14 with a 1-dimensional RMI.

We conclude this section with a brief comparison of the path analysis developed here and the use of the dependency graph refinement in termination analysis. First we recall a theorem on the dependency graph refinement in conjunction with usable rules and innermost rewriting (see [24], but also [25]). Similar results hold in the context of full rewriting, see [21, 22].

Theorem 7.18 ([24]). A TRS \mathcal{R} is innermost terminating if for every maximal cycle \mathcal{C} in the dependency graph $\mathsf{DG}(\mathcal{R})$ there exists a reduction pair (\gtrsim, \succ) such that $\mathcal{U}(\mathcal{C}) \subseteq \gtrsim$ and $\mathcal{C} \subseteq \succ$.

The following example shows that in the context of complexity analysis it is *not* sufficient to consider each cycle individually.

Example 7.19 (continued from Example 6.9). Consider the TRS \mathcal{R}_{exp} introduced in Example 6.9.

$$\begin{split} \exp(\mathbf{0}) &\to \mathbf{s}(\mathbf{0}) & & \mathbf{d}(\mathbf{0}) \to \mathbf{0} \\ \exp(\mathbf{r}(x)) &\to \mathbf{d}(\exp(x)) & & \mathbf{d}(\mathbf{s}(x)) \to \mathbf{s}(\mathbf{s}(\mathbf{d}(x))) \;. \end{split}$$

Recall that the (innermost) runtime complexity of \mathcal{R}_{exp} is exponential. Let \mathcal{P} denote the (standard) dependency pairs with respect to \mathcal{R}_{exp} . Then \mathcal{P} consists of three pairs: 1: $\exp^{\sharp}(\mathbf{r}(x)) \rightarrow d^{\sharp}(\exp(x))$, 2: $\exp^{\sharp}(\mathbf{r}(x)) \rightarrow \exp^{\sharp}(x)$, and 3: $d^{\sharp}(\mathbf{s}(x)) \rightarrow d^{\sharp}(x)$. Hence the dependency graph $\mathsf{DG}(\mathcal{R}_{exp})$ contains two maximal cycles: {2} and {3}.

We define two reduction pairs $(\succeq_{\mathcal{A}}, \succeq_{\mathcal{A}})$ and $(\succeq_{\mathcal{B}}, \succeq_{\mathcal{B}})$ such that the conditions of the theorem are fulfilled. Let \mathcal{A} and \mathcal{B} be SLIs such that $\exp_{\mathcal{A}}^{\sharp}(x) = x$, $\mathsf{r}_{\mathcal{A}}(x) = x + 1$ and $\mathsf{d}_{\mathcal{B}}^{\sharp}(x) = x$, $\mathsf{s}_{\mathcal{A}}(x) = x + 1$. Hence for any term $t \in \mathcal{T}_{\mathsf{b}}$, we have that the derivation heights $\mathsf{dh}(t^{\sharp}, \stackrel{i}{\to}_{\{2\}/\mathcal{U}(\mathcal{P})})$ and $\mathsf{dh}(t^{\sharp}, \stackrel{i}{\to}_{\{3\}/\mathcal{U}(\mathcal{P})})$ are linear in |t|, while $\mathsf{dh}(t, \stackrel{i}{\to}_{\mathcal{R}})$ is (at least) exponential in |t|.

Observe that the problem exemplified by Example 7.19 cannot be circumvented by replacing the dependency graph employed in Theorem 7.18 with weak (innermost) dependency graphs. The exponential derivation height of terms t_n in Example 7.19 is not controlled by the cycles $\{2\}$ or $\{3\}$, but achieved through the non-cyclic pair 1 and its usable rules. Example 7.19 shows an exponential speed-up between the maximal number of dependency pair steps within a cycle in the dependency graph and the runtime complexity of the initial TRS. In the context of derivational complexity this speed-up may even increase to a primitive recursive function, cf. [23].

While Example 7.19 shows that the usable rules need to be taken into account fully for any complexity analysis, it is perhaps tempting to think that it should suffice to demand that at least one weak (innermost) dependency pair in each cycle decreases strictly. However this intuition is deceiving as shown by the next example.

Example 7.20. Consider the TRS \mathcal{R} of $f(s(x), 0) \to f(x, s(0))$ and $f(x, s(y)) \to f(x, y)$. WDP(\mathcal{R}) consists of 1: $f^{\sharp}(s(x), 0) \to f^{\sharp}(x, s(x))$ and 2: $f^{\sharp}(x, s(y)) \to f^{\sharp}(x, y)$, and the weak dependency graph WDG(\mathcal{R}) contains two cycles {1,2} and {2}. There are two linear restricted interpretations \mathcal{A} and \mathcal{B} such that $\{1, 2\} \subseteq \geq_{\mathcal{A}} \cup \geq_{\mathcal{A}}, \{1\} \subseteq \geq_{\mathcal{A}}, \text{ and } \{2\} \subseteq \geq_{\mathcal{B}}$. Here, however, we must not conclude linear runtime complexity, because the runtime complexity of \mathcal{R} is at least quadratic.

8 Experiments

All described techniques have been incorporated into the *Tyrolean Complexity Tool* $T_{C}T$, an open source complexity analyser⁶. The testbed is based on version 8.0.2 of the *Termination Problems Database (TPDB* for short). We consider TRSs without theory annotation, where the runtime complexity analysis is non-trivial, that is the set of basic terms is infinite. This testbed comprises 1695 TRSs. All experiments were conducted on a machine that is identical to the official competition server (8 AMD Opteron[®] 885 dual-core processors with 2.8GHz, 8x8 GB memory). As timeout we use 60 seconds. The complete experimental data can be found at http://cl-informatik.uibk.ac.at/software/tct/experiments, where also the testbed employed is detailed.

Table 1 summarises the experimental results of the here presented techniques for full runtime complexity analysis in a restricted setting. The tests are based on the use of oneand two-dimensional RMIs with coefficients over $\{0, 1, \ldots, 7\}$ as direct technique (compare Theorem 3.9) as well as in combination with the WDP method (compare Corollaries 5.13 and 6.14) and the WDG method (compare Corollary 7.15). Weak dependency graphs are estimated by the TCAP-based technique ([20]). The tests indicate the power of the transformation techniques introduced. Note that for linear and quadratic runtime complexity the latter techniques are more powerful than the direct approach. Furthermore note that the WDG method provides overall better bounds than the WDP method.

However if we consider RMIs upto dimension 3 the picture becomes less clear, cf. Table 2. Again we compare the direct approach, the WDP and WDG method and restrict to coefficients over $\{0, 1, \ldots, 7\}$. Consider for example the test results for cubic runtime complexity with respect to full rewriting. While the transformation techniques are still more powerful than the direct approach, the difference is less significant than in Table 1. On one hand this is due to the fact that RMIs employing matrices of dimension k may have a degree strictly smaller than k, compare Theorem 3.9 and on the other hand note the increase in timeouts for the more advanced techniques.

⁶ Available at http://cl-informatik.uibk.ac.at/software/tct.

		full						
result	direct (1)	direct (2)	WDP (1)	WDP (2)	WDG(1)	WDG(2)		
O(1)	16	18	0	0	10	10		
O(n)	106	113	123	70	130	67		
$O(n^2)$	106	148	123	157	130	158		
timeout $(60s)$	20	88	55	127	103	261		

Table 1: Experiment results I (one- and two-dimensional RMIs separated)

Moreover note the seemingly strange behaviour of the WDG method for innermost rewriting: already for quadratic runtime the WDP method performs better, if we only consider the number of yes-instances. This seems to contradict the fact that the WDG method is in theory more powerful than the WDP method. However, the explanation is simple: first the sets of yes-instances are incomparable and second the more advanced technique requires more computation power. If we would use (much) longer timeout the set of yes-instances for WDP would become a *proper* subset of the set of yes-instances for WDG. For example the WDG method can prove cubic runtime complexity of the TRS AProVE_04/Liveness 6.2 from the TPDB, while the WDP method fails to give its bound.

		fu	11	innermost			
result	direct	WDP	WDG	direct	WDP	WDG	
O(1)	18	0	10	20	0	10	
O(n)	135	141	140	135	142	145	
$O(n^2)$	161	163	162	173	181	172	
$O(n^3)$	163	167	169	179	185	178	
timeout (60s)) 310	459	715	311	458	718	

Table 2: Experiment results II (1–3-dimensional RMIs combined)

In order to assess the advances of this paper in contrast to the conference versions (see [4, 7]), we present in Table 3 a comparison between RMIs with/without the use of usable arguments and a comparison of the WDP or WDG method with/without the use of the extended weight gap principle. Again we restrict our attention to full rewriting, as the case for innermost rewriting provides a similar picture (see http://cl-informatik.uibk.ac.at/software/tct/experime for the full data).

Finally, in Table 4 we present the overall power obtained for the automated runtime complexity analysis. Here we test the version of T_CT that run for the international annual termination competition (TERMCOMP)⁷ in 2010 in comparison to the most recent version of T_CT incorporating all techniques developed in this paper. In addition we compare with a recent version of C_aT .⁸

⁷ http://termcomp.uibk.ac.at/termcomp/.

⁸ http://cl-informatik.uibk.ac.at/software/cat/.

	full						
result	$\operatorname{direct}\left(-\right)$	$\operatorname{direct}\left(+\right)$	$\mathrm{WDP}\left(-\right)$	$\mathrm{WDP}\left(+\right)$	WDG(-)	WDG(+)	
O (1)	4	18	5	0	10	10	
O(n)	105	135	102	141	105	140	
$O(n^2)$	127	161	118	163	119	162	
$O(n^3)$	130	163	120	167	122	169	
timeout (60s)) 306	310	505	459	655	715	

Table 3: Experiment results III (1–3-dimensional RMIs combined)

		full	innermost			
result	$T_{C}T({\rm old})$	$T_{C}T(\mathrm{new})$	CaT	$T_{C}T({\rm old})$	$T_{C}T(\mathrm{new})$	CaT
O (1)	10	3	0	10	3	0
O(n)	393	486	439	401	488	439
$O(n^2)$	394	493	452	403	502	452
$O(n^3)$	397	495	453	407	505	453
$O(n^4)$	397	495	454	407	505	454

Table 4: Experiment results IV (1–3-dimensional RMIs combined)

The results in Table 4 clearly show the increase in power in T_{CT} , which is due to the fact that the techniques developed in this paper have been incorporated.

9 Conclusion

In this article we are concerned with automated complexity analysis of TRSs. More precisely, we establish new and powerful results that allow the assessment of polynomial runtime complexity of TRSs fully automatically. We established the following results: Adapting techniques from context-sensitive rewriting, we introduced *usable replacement maps* that allow to increase the applicability of direct methods. Furthermore we established the *weak dependency pair method* as a suitable analog of the dependency pair method in the context of (runtime) complexity analysis. Refinements of this method have been presented by the use of the *weight gap principle* and *weak dependency graphs*. In the experiments of Section 8 we assessed the viability of these techniques. It is perhaps worthy of note to mention that our motivating examples (Examples 3.2, 5.15, and 7.5) could not be handled by any known technique prior to our results.

To conclude, we briefly mention related work. Based on earlier work by Arai and the second author (see [26]) Avanzini and the second author introduced POP^{*} a restriction of the recursive path order (RPO) that induces polynomial innermost runtime complexity (see [27, 15]). With respect to derivational complexity, Zankl and Korp generalised a simple variant of our weight gap principle to achieve a modular derivational complexity analysis

(see [28, 29]). Neurauter et al. refined in [16] matrix interpretations in the context of derivational complexity derivational complexity (see also [30]). Furthermore, Waldmann studied in [17] the use of weighted automata in this setting. Based on [4, 7] Noschinski et al. incorporated a variant of weak dependency pairs (not yet published) into the termination prover AProVE.⁹ Currently this method is restricted to innermost runtime complexity, but allows for a complexity analysis in the spirit of the dependency pair framework. Preliminary evidence suggests that this technique is orthogonal to the methods presented here. While all mentioned results are concerned with *polynomial* upper bounds on the derivational or runtime complexity of a rewrite system, Schnabl and the second author provided in [31, 23, 32] an analysis of the dependency pair method and its framework from a complexity point of view. The upshot of this work is that the dependency pair framework may induce multiple recursive derivational complexity, even if only simple processors are considered.

Investigations into the complexity of TRSs are strongly influenced by research in the field of ICC, which contributed the use of restricted forms of polynomial interpretations to estimate the complexity, cf. [18]. Related results have also been provided in the study of term rewriting characterisations of complexity classes (compare [33]). Inspired by Bellantoni and Cook's recursion theoretic characterisation of the class of all polynomial time computable functions in [34], Marion [35] defined LMPO, a variant of RPO whose compatibility with a TRS implies that the functions computed by the TRS is polytime computable (compare [3]). A remarkable milestone on this line is the quasi-interpretation method by Bonfante et al. [36]. The method makes use of standard termination methods in conjunction with special polynomial interpretations to characterise the class of polytime computable functions. In conjunction with *sup-interpretations* this method is even capable of making use of *standard* dependency pairs (see [37]).

In principle we cannot directly compare our result on *polynomial* runtime complexity of TRSs with the results provided in the setting of ICC: the notion of complexity studied is different. However, due to a recent result by Avanzini and the second author (see [38], but compare also [39, 40]) we know that the runtime complexity of a TRS is an *invariant* cost model. Whenever we have polynomial runtime complexity of a TRS \mathcal{R} , the functions computed by this \mathcal{R} can be implemented on a Turing machine that runs in polynomial time. In this context, our results provide automated techniques that can be (almost directly) employed in the context of ICC. The qualification only refers to the fact that our results are presented for an abstract form of programs, viz. rewrite systems.

References

- C. Choppy, S. Kaplan, M. Soria, Complexity analysis of term-rewriting systems, Theor. Comput. Sci. 67 (2–3) (1989) 261–282.
- [2] D. Hofbauer, C. Lautemann, Termination proofs and the length of derivations, in: Proc. 3rd International Conference on Rewriting Techniques and Applications, no. 355 in LNCS, Springer Verlag, 1989, pp. 167–177.

⁹ This novel version of AProVE (see http://aprove.informatik.rwth-aachen.de/) for (innermost) runtime complexity took part in TERMCOMP in 2010.

- [3] E.-A. Cichon, P. Lescanne, Polynomial interpretations and the complexity of algorithms, in: Proc. 11th International Conference on Automated Deduction, Vol. 607 of LNCS, 1992, pp. 139–147.
- [4] N. Hirokawa, G. Moser, Automated complexity analysis based on the dependency pair method, in: Proc. 4th International Joint Conference on Automated Reasoning, no. 5195 in LNAI, Springer Verlag, 2008, pp. 364–380.
- [5] P. Baillot, J.-Y. Marion, S. R. D. Rocca, Guest editorial: Special issue on implicit computational complexity, ACM Trans. Comput. Log. 10 (4).
- [6] T. Arts, J. Giesl, Termination of term rewriting using dependency pairs, Theor. Comput. Sci. 236 (2000) 133–178.
- [7] N. Hirokawa, G. Moser, Complexity, graphs, and the dependency pair method, in: Proc. 15th International Conference on Logic for Programming Artificial Intelligence and Reasoning, no. 5330 in LNCS, Springer Verlag, 2008, pp. 652–666.
- [8] F. Baader, T. Nipkow, Term Rewriting and All That, Cambridge University Press, 1998.
- [9] TeReSe, Term Rewriting Systems, Vol. 55 of Cambridge Tracks in Theoretical Computer Science, Cambridge University Press, 2003.
- [10] A. Geser, Relative termination, Ph.D. thesis, Universität Passau (1990).
- [11] R. Thiemann, The DP framework for proving termination of term rewriting, Ph.D. thesis, University of Aachen, Department of Computer Science (2007).
- [12] J. Endrullis, J. Waldmann, H. Zantema, Matrix interpretations for proving termination of term rewriting, J. Automated Reasoning 40 (3) (2008) 195–220.
- [13] D. Hofbauer, J. Waldmann, Termination of string rewriting with matrix interpretations, in: Proc. 17th International Conference on Rewriting Techniques and Applications, Vol. 4098 of LNCS, 2006, pp. 328–342.
- [14] T. Arts, J. Giesl, A collection of examples for termination of term rewriting using dependency pairs, Tech. Rep. AIB-2001-09, RWTH Aachen (2001).
- [15] M. Avanzini, G. Moser, Dependency pairs and polynomial path orders, in: Proc. 20th International Conference on Rewriting Techniques and Applications, Vol. 5595 of LNCS, 2009, pp. 48–62.
- [16] F. Neurauter, H. Zankl, A. Middeldorp, Revisiting matrix interpretations for polynomial derivational complexity of term rewriting, in: Proc. 17th International Conference on Logic for Programming Artificial Intelligence and Reasoning, Vol. 6397 of LNCS (ARCoSS), 2010, pp. 550–564.
- [17] J. Waldmann, Polynomially bounded matrix interpretations, in: Proc. 21st International Conference on Rewriting Techniques and Applications, Vol. 6 of LIPIcs, 2010, pp. 357–372.

- [18] G. Bonfante, A. Cichon, J.-Y. Marion, H. Touzet, Algorithms with polynomial interpretation termination proof, J. Funct. Program. 11 (1) (2001) 33–53.
- [19] M. L. Fernández, Relaxing monotonicity for innermost termination, Inform. Proc. Lett. 93 (1) (2005) 117–123.
- [20] J. Giesl, R. Thiemann, P. Schneider-Kamp, Proving and disproving termination of higher-order functions, in: Proc. 5th International Workshop on Frontiers of Combining Systems, 5th International Workshop, Vol. 3717 of LNAI, 2005, pp. 216–231.
- [21] J. Giesl, R. Thiemann, P. Schneider-Kamp, S. Falke, Mechanizing and improving dependency pairs, J. Automated Reasoning 37 (3) (2006) 155–203.
- [22] N. Hirokawa, A. Middeldorp, Tyrolean termination tool: Techniques and features, Inform. and Comput. 205 (2007) 474–511.
- [23] G. Moser, A. Schnabl, The derivational complexity induced by the dependency pair method, Logical Methods in Computer ScienceAccepted for publication.
- [24] J. Giesl, T. Arts, E. Ohlebusch, Modular termination proofs for rewriting using dependency pairs, J. Symbolic Comput. 34 (2002) 21–58.
- [25] N. Hirokawa, A. Middeldorp, Automating the dependency pair method, Inform. and Comput. 199 (1,2) (2005) 172–199.
- [26] T. Arai, G. Moser, Proofs of termination of rewrite systems for polytime functions, in: Proc. 25th Conference on Foundations of Software Technology and Theoretical Computer Science, no. 3821 in LNCS, Springer Verlag, 2005, pp. 529–540.
- [27] M. Avanzini, G. Moser, Complexity analysis by rewriting, in: Proc. 9th International Symposium on Functional and Logic Programming, no. 4989 in LNCS, Springer Verlag, 2008, pp. 130–146.
- [28] H. Zankl, M. Korp, Modular complexity analysis via relative complexity, in: Proc. 21st International Conference on Rewriting Techniques and Applications, Vol. 6 of LIPIcs, 2010, pp. 385–400.
- [29] H. Zankl, M. Korp, Modular complexity analysis via relative complexity, Logical Methods in Computer ScienceSubmitted.
- [30] G. Moser, A. Schnabl, J. Waldmann, Complexity analysis of term rewriting based on matrix and context dependent interpretations, in: Proc. 28th Conference on Foundations of Software Technology and Theoretical Computer Science, LIPIcs, 2008, pp. 304–315.
- [31] G. Moser, A. Schnabl, The derivational complexity induced by the dependency pair method, in: Proc. 20th International Conference on Rewriting Techniques and Applications, Vol. 5595 of LNCS, 2009, pp. 255–269.
- [32] G. Moser, A. Schnabl, Termination proofs in the dependency pair framework may induce multiply recursive derivational complexities, in: Proc. 22nd International Conference on Rewriting Techniques and Applications, Vol. 10 of LIPIcs, 2011, pp. 235–250.

- [33] E.-A. Cichon, A. Weiermann, Term rewriting theory for the primitive recursive functions., Ann. Pure Appl. Logic 83 (3) (1997) 199–223.
- [34] S. Bellantoni, S. Cook, A new recursion-theoretic characterization of the polytime functions, Comput. Complexity 2 (2) (1992) 97–110.
- [35] J.-Y. Marion, Analysing the implicit complexity of programs, Inform. and Comput. 183 (2003) 2–18.
- [36] G. Bonfante, J.-Y. Marion, J.-Y. Moyen, Quasi-interpretations: A way to control resources, Theor. Comput. Sci. To appear.
- [37] J.-Y. Marion, R. Péchoux, Sup-interpretations, a semantic method for static analysis of program resources, ACM Trans. Comput. Log. 10 (4).
- [38] M. Avanzini, G. Moser, Closing the gap between runtime complexity and polytime computability, in: Proc. 21st International Conference on Rewriting Techniques and Applications, Vol. 6 of LIPIcs, 2010, pp. 33–48.
- [39] U. Dal Lago, S. Martini, On constructor rewrite systems and the lambda-calculus, in: Proc. 36th ICALP, Vol. 5556 of LNCS, Springer Verlag, 2009, pp. 163–174.
- [40] U. Dal Lago, S. Martini, Derivational Complexity is an Invariant Cost Model, in: Proc. 1st FOPARA, 2009.