

On the optimality of gluing over scales

Alexander Jaffe*

James R. Lee*

Mohammad Moharrami*

Abstract

We show that for every $\alpha > 0$, there exist n -point metric spaces (X, d) where every “scale” admits a Euclidean embedding with distortion at most α , but the whole space requires distortion at least $\Omega(\sqrt{\alpha \log n})$. This shows that the scale-gluing lemma [Lee, SODA 2005] is tight, and disproves a conjecture stated there. This matching upper bound was known to be tight at both endpoints, i.e. when $\alpha = \Theta(1)$ and $\alpha = \Theta(\log n)$, but nowhere in between.

More specifically, we exhibit n -point spaces with doubling constant λ requiring Euclidean distortion $\Omega(\sqrt{\log \lambda \log n})$, which also shows that the technique of “measured descent” [Krauthgamer, et. al., *Geometric and Functional Analysis*] is optimal. We extend this to L_p spaces with $p > 1$, where one requires distortion at least $\Omega((\log n)^{1/q}(\log \lambda)^{1-1/q})$ when $q = \max\{p, 2\}$, a result which is tight for every $p > 1$.

1 Introduction

Suppose one is given a collection of mappings from some finite metric space (X, d) into a Euclidean space, each of which reflects the geometry at some “scale” of X . Is there a non-trivial way of gluing these mappings together to form a global mapping which reflects the entire geometry of X ? The answers to such questions have played a fundamental role in the best-known approximation algorithms for Sparsest Cut [7, 10, 4, 1] and Graph Bandwidth [17, 7, 11], and have found applications in approximate multi-commodity max-flow/min-cut theorems in graphs [17, 7]. In the present paper, we show that the approaches of [7] and [10] are optimal, disproving a conjecture stated in [10].

Let (X, d) be an n -point metric space, and suppose that for every $k \in \mathbb{Z}$, we are given a non-expansive mapping $\phi_k : X \rightarrow L_2$ which satisfies the following. For every $x, y \in X$ with $d(x, y) \geq 2^k$, we have

$$\|\phi_k(x) - \phi_k(y)\| \geq \frac{2^k}{\alpha}.$$

The Gluing Lemma of [10] (generalizing the approach of [7]) shows that the existence of such a collection $\{\phi_k\}$ yields a Euclidean embedding of (X, d) with distortion $O(\sqrt{\alpha \log n})$. (See Section 1.1 for the relevant definitions on embeddings and distortion.) This is known to be tight when $\alpha = \Theta(1)$ [16] and also when $\alpha = \Theta(\log n)$ [13, 2], but nowhere in between. In fact, in [10], the second named author conjectured that one could achieve $O(\alpha + \sqrt{\log n})$ (this is indeed stronger, since one can always construct $\{\phi_k\}$ with $\alpha = O(\log n)$).

In the present paper, we give a family of examples which shows that the $\sqrt{\alpha \log n}$ bound is tight for any dependence $\alpha(n) = O(\log n)$. In fact, we show more. Let $\lambda(X)$ denote the *doubling*

*Research partially supported by NSF CCF-0644037 and a Sloan Research Fellowship.

constant of X , i.e. the smallest number λ so that every open ball in X can be covered by λ balls of half the radius. In [7], using the method of “measure descent,” the authors show that (X, d) admits a Euclidean embedding with distortion $O(\sqrt{\log \lambda(X) \log n})$. (This is a special case of the Gluing Lemma since one can always find $\{\phi_k\}$ with $\alpha = O(\log \lambda(X))$ [5]). Again, this bound was known to be tight for $\lambda(X) = \Theta(1)$ [8, 9, 5] and $\lambda(X) = n^{\Theta(1)}$ [13, 2], but nowhere in between. We provide the matching lower bound for any dependence of $\lambda(X)$ on n . We also generalize our method to give tight lower bounds on L_p distortion for every fixed $p > 1$.

Construction and analysis. In some sense, our lower bound examples are an interpolation between the multi-scale method of [16] and [8], and the expander Poincaré inequalities of [13, 2, 14]. We start with a vertex-transitive expander graph G on m nodes. If D is the diameter of G , then we create $D + 1$ copies G^1, G^2, \dots, G^{D+1} of G where $u \in G^i$ is connected to $v \in G^{i+1}$ if (u, v) is an edge in G , or if $u = v$. We then connect a vertex s to every node in G^1 and a vertex t to every node in G^{D+1} by edges of length D . This yields the graph \vec{G} described in Section 2.2.

In Section 3, we show that whenever there is a non-contracting embedding f of \vec{G} into L_2 , the following holds. If $\gamma = \frac{\|f(s) - f(t)\|}{d_{\vec{G}}(s, t)}$, then some edge of \vec{G} gets stretched by at least $\sqrt{\gamma^2 + \Omega(\log m)^2}$, i.e. there is a “stretch increase.” This is proved by combining the uniform convexity of L_2 (i.e. the Pythagorean theorem), with the well-known contraction property of expander graphs mapped into Hilbert space. To convert the “average” nature of this contraction to information about a specific edge, we symmetrize the embedding over all automorphisms of G (which was chosen to be vertex-transitive).

To exploit this stretch increase recursively, we construct a graph $\vec{G}^{\otimes k}$ inductively as follows: $\vec{G}^{\otimes k}$ is formed by replacing every edge of $\vec{G}^{\otimes k-1}$ by a copy of \vec{G} (see Section 2.1 for the formal definitions). Now a simple induction shows that in a non-contracting embedding of $\vec{G}^{\otimes k}$, there must be an edge stretched by at least $\Omega(\sqrt{k} \log m)$. In Section 3.1, a similar argument is made for L_p distortion, for $p > 1$, but here we have to argue about “quadrilaterals” instead of “triangles” (in order to apply the uniform convexity inequality in L_p), and it requires slightly more effort to find a good quadrilateral.

Finally, we observe that if \tilde{G} is the graph formed by adding two tails of length $3D$ hanging off s and t in \vec{G} , then (following the analysis of [8, 9]), one has $\log \lambda(\tilde{G}^{\otimes k}) \lesssim \log m$. The same lower bound analysis also works for $\tilde{G}^{\otimes k}$, so since $n = |V(\tilde{G}^{\otimes k})| = 2^{\Theta(k \log m)}$, the lower bound is

$$\sqrt{k} \log m \approx \sqrt{\log m \log n} \gtrsim \sqrt{\log \lambda(\tilde{G}^{\otimes k}) \log n},$$

completing the proof.

1.1 Preliminaries

For a graph G , we will use $V(G), E(G)$ to denote the sets of vertices and edges of G , respectively. Sometimes we will equip G with a non-negative length function $\text{len} : E(G) \rightarrow \mathbb{R}_+$, and we let d_{len} denote the shortest-path (semi-)metric on G . We refer to the pair (G, len) as a *metric graph*, and often len will be implicit, in which case we use d_G to denote the path metric. We use $\text{Aut}(G)$ to denote the group of automorphisms of G .

Given two expressions E and E' (possibly depending on a number of parameters), we write $E = O(E')$ to mean that $E \leq CE'$ for some constant $C > 0$ which is independent of the parameters. Similarly, $E = \Omega(E')$ implies that $E \geq CE'$ for some $C > 0$. We also write $E \lesssim E'$ as a synonym for $E = O(E')$. Finally, we write $E \approx E'$ to denote the conjunction of $E \lesssim E'$ and $E \gtrsim E'$.




Figure 1: A single edge H , $H \odot K_{2,3}$, and $H \odot K_{2,3} \odot K_{2,2}$.

Embeddings and distortion. If $(X, d_X), (Y, d_Y)$ are metric spaces, and $f : X \rightarrow Y$, then we write

$$\|f\|_{\text{Lip}} = \sup_{x \neq y \in X} \frac{d_Y(f(x), f(y))}{d_X(x, y)}.$$

If f is injective, then the *distortion* of f is defined by $\text{dist}(f) = \|f\|_{\text{Lip}} \cdot \|f^{-1}\|_{\text{Lip}}$. A map with distortion D will sometimes be referred to as *D-bi-lipschitz*. If $d_Y(f(x), f(y)) \leq d_X(x, y)$ for every $x, y \in X$, we say that f is *non-expansive*. If $d_Y(f(x), f(y)) \geq d_X(x, y)$ for every $x, y \in X$, we say that f is *non-contracting*. For a metric space X , we use $c_p(X)$ to denote the least distortion required to embed X into some L_p space.

Finally, for $x \in X$, $r \in \mathbb{R}_+$, we define the open ball $B(x, r) = \{y \in X : d(x, y) < r\}$. Recall that the *doubling constant* of a metric space (X, d) is the infimum over all values λ such that every ball in X can be covered by λ balls of half the radius. We use $\lambda(X, d)$ to denote this value.

We now state the main theorem of the paper.

Theorem 1.1. *For any positive nondecreasing function $\lambda(n)$, there exists a family of n -vertex metric graphs $\tilde{G}^{\odot k}$ such that $\lambda(\tilde{G}^{\odot k}) \lesssim \lambda(n)$, and for every fixed $p > 1$,*

$$c_p(\tilde{G}^{\odot k}) \gtrsim (\log n)^{1/q} (\log \lambda(n))^{1-1/q},$$

where $q = \max\{p, 2\}$.

2 Metric construction

2.1 \odot -products

An *s-t* graph G is a graph which has two distinguished vertices $s, t \in V(G)$. For an *s-t* graph, we use $s(G)$ and $t(G)$ to denote the vertices labeled s and t , respectively. We define the length of an *s-t* graph G as $\text{len}(G) = d_{\text{len}}(s, t)$.

Definition 2.1 (Composition of *s-t* graphs). *Given two s-t graphs H and G , define $H \odot G$ to be the s-t graph obtained by replacing each edge $(u, v) \in E(H)$ by a copy of G (see Figure 1). Formally,*

- $V(H \odot G) = V(H) \cup (E(H) \times (V(G) \setminus \{s(G), t(G)\}))$.
- For every edge $e = (u, v) \in E(H)$, there are $|E(G)|$ edges,

$$\left\{ \left((e, v_1), (e, v_2) \right) \mid (v_1, v_2) \in E(G) \text{ and } v_1, v_2 \notin \{s(G), t(G)\} \right\} \cup \\ \left\{ \left((u, (e, w)), (s(G), w) \right) \mid (s(G), w) \in E(G) \right\} \cup \left\{ \left((e, w), v \right) \mid (w, t(G)) \in E(G) \right\}$$

- $s(H \odot G) = s(H)$ and $t(H \odot G) = t(H)$.

If H and G are equipped with length functions $\text{len}_H, \text{len}_G$, respectively, we define $\text{len} = \text{len}_{H \odot G}$ as follows. Using the preceding notation, for every edge $e = (u, v) \in E(H)$,

$$\begin{aligned} \text{len}((e, v_1), (e, v_2)) &= \frac{\text{len}_H(e)}{d_{\text{len}_G}(s(G), t(G))} \text{len}_G(v_1, v_2) \\ \text{len}(u, (e, w)) &= \frac{\text{len}_H(e)}{d_{\text{len}_G}(s(G), t(G))} \text{len}_G(s(G), w) \\ \text{len}((e, w), v) &= \frac{\text{len}_H(e)}{d_{\text{len}_G}(s(G), t(G))} \text{len}_G(w, t(G)). \end{aligned}$$

This choice implies that $H \odot G$ contains an isometric copy of $(V(H), d_{\text{len}_H})$.

Observe that there is some ambiguity in the definition above, as there are two ways to substitute an edge of H with a copy of G , thus we assume that there exists some arbitrary orientation of the edges of H . However, for our purposes the graph G will be symmetric, and thus the orientations are irrelevant.

Definition 2.2 (Recursive composition). For an s - t graph G and a number $k \in \mathbb{N}$, we define $G^{\odot k}$ inductively by letting $G^{\odot 0}$ be a single edge of unit length, and setting $G^{\odot k} = G^{\odot k-1} \odot G$.

The following result is straightforward.

Lemma 2.3 (Associativity of \odot). For any three graphs A, B, C , we have $(A \odot B) \odot C = A \odot (B \odot C)$, both graph-theoretically and as metric spaces.

Definition 2.4. For two graphs G, H , a subset of vertices $X \subseteq V(H)$ is said to be a copy of G if there exists a bijection $f : V(G) \rightarrow X$ with distortion 1, i.e. $d_H(f(u), f(v)) = C \cdot d_G(u, v)$ for some constant $C > 0$.

Now we make the following two simple observations about copies of H and G in $H \odot G$.

Observation 2.5. The graph $H \odot G$ contains $|E(H)|$ distinguished copies of the graph G , one copy corresponding to each edge in H .

Observation 2.6. The subset of vertices $V(H) \subseteq V(H \odot G)$ form an isometric copy of H .

2.2 A stretched \vec{G}

Let $G = (V, E)$ be an unweighted graph, and put $D = \text{diam}(G)$. We define a metric s - t graph \vec{G} which has $D + 1$ layers isomorphic to G , with edges between the layers, and a pair of endpoints s, t . Formally,

$$\begin{aligned} V(\vec{G}) &= \{s, t\} \cup \{v^{(i)} : v \in V, i \in [D + 1]\} \\ E(\vec{G}) &= \{(s, v^{(1)}), (v^{(D+1)}, t) : v \in V\} \\ &\quad \cup \left\{ (u^{(i)}, v^{(i+1)}), (u^{(j)}, v^{(j)}) : (u, v) \in E, i \in [D], j \in [D + 1] \right\} \\ &\quad \cup \{ (v^{(i)}, v^{(i+1)}) : v \in V, i \in [D] \}. \end{aligned}$$

We put $\text{len}(s, v^{(1)}) = \text{len}(v^{(D+1)}, t) = D$ for $v \in V$, $\text{len}(u^{(i)}, v^{(i+1)}) = \text{len}(u^{(j)}, v^{(j)}) = 1$ for $(u, v) \in E$, $i \in [D]$, $j \in [D+1]$ and $\text{len}(v^{(i)}, v^{(i+1)}) = 1$ for $v \in V, i \in [D]$. We refer to edges of the form $(u^{(i)}, v^{(i)})$ as *vertical edges*. All other edges are called *horizontal edges*. In particular, there are $D+1$ copies $G^{(1)}, \dots, G^{(D+1)}$ of G in \tilde{G} which are isometric to G itself, and their edges are all vertical.

A doubling version, following Laakso. Let \vec{G} be a stretched graph as in Section 2.2, with $D = \text{diam}(G)$, and let $s' = s(\vec{G}), t' = t(\vec{G})$. Consider a new metric s - t graph \tilde{G} , which has two new vertices s, t and two new edges $(s, s'), (t', t)$ with $\text{len}(s, s') = \text{len}(t', t) = 3D$.

Claim 2.7. *For any graph G with $|V(G)| = m$, and any $k \in \mathbb{N}$, we have $\log \lambda(\tilde{G}^{\otimes k}) \lesssim \log m$.*

The proof of the claim is similar to [8, 9], and follows from the following three results.

We define $\text{tri}(G) = \max_{v \in V(G)} (d_{\text{len}}(s, v) + d_{\text{len}}(v, t))$. For any graph G , we have $\text{len}(\tilde{G}) = d(s, t) = 9D$, and it is not hard to verify that $\text{tri}(\tilde{G}^{\otimes k}) \leq \text{len}(\tilde{G}^{\otimes k}) (1 + \frac{1}{9D-1})$. For convenience, let G_0 be the top-level copy of \tilde{G} in $\tilde{G}^{\otimes k}$, and H be the graph $\tilde{G}^{\otimes k-1}$. Then for any $e \in E(G_0)$, we refer to the copy of H along edge e as H_e .

Observation 2.8. *If $r > \frac{\text{tri}(\tilde{G}^{\otimes k})}{3}$, then the ball $B(x, r)$ in $\tilde{G}^{\otimes k}$ may be covered by at most $|V(\tilde{G})|$ balls of radius $r/2$.*

Proof. For any $e \in E(G_0)$, we have $r > \frac{\text{len}(e)}{\text{len}(H)} \text{tri}(H)$, so every point in H_e is less than $r/2$ from an endpoint of e . Thus all of $\tilde{G}^{\otimes k}$ is covered by placing balls of radius $\frac{\text{tri}(\tilde{G}^{\otimes k})}{6}$ around each vertex of G_0 . \square

Lemma 2.9. *If $s \in B(x, r)$, then one can cover the ball $B(x, r)$ in $\tilde{G}^{\otimes k}$ with at most $|E(\tilde{G})||V(\tilde{G})|$ balls of radius $r/2$.*

Proof. First consider the case in which $r > \frac{\text{len}(\tilde{G}^{\otimes k})}{6}$. Then for any edge e in $\tilde{G}^{\otimes k}$, we have $r > \frac{\text{len}(e)}{\text{len}(H)} \cdot \frac{\text{tri}(H)}{3}$. Thus by Observation 2.8, we may cover H_e by $|V(\tilde{G})|$ balls of radius $r/2$. This gives a covering of all of $\tilde{G}^{\otimes k}$ by at most $|E(\tilde{G})||V(\tilde{G})|$ balls of radius $r/2$.

Otherwise, assume $\frac{\text{len}(\tilde{G}^{\otimes k})}{6} \geq r$. Since $s \in B(x, r)$, but $2r \leq \frac{\text{len}(\tilde{G}^{\otimes k})}{3}$, the ball must be completely contained inside $H_{(s, s')}$. By induction, we can find a sufficient cover of this smaller graph. \square

Lemma 2.10. *We can cover any ball $B(x, r)$ in $\tilde{G}^{\otimes k}$ with at most $2|V(\tilde{G})||E(\tilde{G})|^2$ balls of radius $r/2$.*

Proof. We prove this lemma using induction. For $\tilde{G}^{\otimes 0}$, the claim holds trivially. Next, if any H_e contains all of $B(x, r)$, then by induction we are done. Otherwise, for each H_e containing x , $B(x, r)$ contains an endpoint of e . Then by Lemma 2.9, we may cover H_e by at most $|E(\tilde{G})||V(\tilde{G})|$ balls of radius $r/2$. For all other edges $e' = (u, v)$, $x \notin H_{e'}$, so we have:

$$V(H_{e'}) \cap B(x, r) \subseteq B(v, \max(0, r - d(x, v))) \cup B(u, \max(0, r - d(x, u))).$$

Thus, using Lemma 2.9 on both of the above balls, we may cover $V(H_{e'}) \cap B(x, r)$ by at most $2|E(\tilde{G})||V(\tilde{G})|$ balls of radius $r/2$. Hence, in total, we need at most $2|V(\tilde{G})||E(\tilde{G})|^2$ balls of radius $r/2$ to cover all of $B(x, r)$. \square

Proof of Claim 2.7. First note that $|V(\tilde{G})| = m(D+1) + 2 \lesssim m^2$. By Lemma 2.10, we have

$$\lambda(\tilde{G}^{\otimes k}) \leq 2|V(\tilde{G})||E(\tilde{G})|^2 \leq 2|V(\tilde{G})|^5 \lesssim m^{10}.$$

Hence $\log \lambda(\tilde{G}^{\otimes k}) \lesssim \log m$. □

3 Lower bound

For any $\pi \in \text{Aut}(G)$, we define a corresponding automorphism $\tilde{\pi}$ of \tilde{G} by $\tilde{\pi}(s) = s$, $\tilde{\pi}(t) = t$, $\tilde{\pi}(s') = s'$, $\tilde{\pi}(t') = t'$, and $\tilde{\pi}(v^{(i)}) = \pi(v)^{(i)}$ for $v \in V, i \in [D+1]$.

Lemma 3.1. *Let G be a vertex transitive graph. Let $f : V(\tilde{G}) \rightarrow L_2$ be an injective mapping and define $\bar{f} : V(\tilde{G}) \rightarrow L_2$ by*

$$\bar{f}(x) = \frac{1}{\sqrt{|\text{Aut}(G)|}} \left(f(\tilde{\pi}x) \right)_{\pi \in \text{Aut}(G)}.$$

Let β be such that for every $i \in [D+1]$ there exists a vertical edge $(u^{(i)}, v^{(i)})$ with $\|\bar{f}(u^{(i)}) - \bar{f}(v^{(i)})\| \geq \beta$. Then there exists a horizontal edge $(x, y) \in E(\tilde{G})$ such that

$$\frac{\|\bar{f}(x) - \bar{f}(y)\|^2}{d_{\tilde{G}}(x, y)^2} \geq \frac{\|f(s) - f(t)\|^2}{d_{\tilde{G}}(s, t)^2} + \frac{\beta^2}{36} \quad (1)$$

Proof. Let $D = \text{diam}(G)$. We first observe three facts about \bar{f} , which rely on the fact that when $\text{Aut}(G)$ is transitive, for every $x \in V$, the orbits $\{\pi(x)\}_{\pi \in \text{Aut}(G)}$ all have the same cardinality.

$$(F1) \quad \|\bar{f}(s) - \bar{f}(t)\| = \|f(s) - f(t)\|$$

$$(F2) \quad \text{For all } u, v \in V,$$

$$\begin{aligned} \|\bar{f}(s) - \bar{f}(v^{(1)})\| &= \|\bar{f}(s) - \bar{f}(u^{(1)})\|, \\ \|\bar{f}(t) - \bar{f}(v^{(D+1)})\| &= \|\bar{f}(t) - \bar{f}(u^{(D+1)})\|. \end{aligned}$$

$$(F3) \quad \text{For every } u, v \in V, i \in [D],$$

$$\|\bar{f}(v^{(i)}) - \bar{f}(v^{(i+1)})\| = \|\bar{f}(u^{(i)}) - \bar{f}(u^{(i+1)})\|.$$

$$(F4) \quad \text{For every pair of vertices } u, v \in V \text{ and } i \in [D+1],$$

$$\langle \bar{f}(s) - \bar{f}(t), \bar{f}(u^{(i)}) - \bar{f}(v^{(i)}) \rangle = 0.$$

Let $z = \frac{\bar{f}(s) - \bar{f}(t)}{\|\bar{f}(s) - \bar{f}(t)\|}$. Fix some $r \in V$ and let $\rho_0 = |\langle z, \bar{f}(s) - \bar{f}(r^{(1)}) \rangle|$, $\rho_i = |\langle z, \bar{f}(r^{(i)}) - \bar{f}(r^{(i+1)}) \rangle|$ for $i = 1, 2, \dots, D$ and $\rho_{D+1} = |\langle z, \bar{f}(t) - \bar{f}(r^{(D+1)}) \rangle|$. Note that, by (F2) and (F3) above, the values $\{\rho_i\}$ do not depend on the representative $r \in V$. In this case, we have

$$\sum_{i=0}^{D+1} \rho_i \geq \|\bar{f}(s) - \bar{f}(t)\| = 9\gamma D, \quad (2)$$

where we put $\gamma = \frac{\|\bar{f}(s) - \bar{f}(t)\|}{d_{\tilde{G}}(s, t)}$. Note that $\gamma > 0$ since f is injective.

Recalling that $d_{\tilde{G}}(s, t) = 9D$ and $d_{\tilde{G}}(s, r^{(1)}) = 4D$, observe that if $\rho_0^2 \geq \left(1 + \frac{\beta^2}{36\gamma^2}\right)(4\gamma D)^2$, then

$$\max\left(\frac{\|\bar{f}(s) - \bar{f}(s')\|^2}{d_{\tilde{G}}(s, s')^2}, \frac{\|\bar{f}(s') - \bar{f}(r^{(1)})\|^2}{d_{\tilde{G}}(s', r^{(1)})^2}\right) \geq \gamma^2 + \frac{\beta^2}{36},$$

verifying (1). The symmetric argument holds for ρ_{D+1} , thus we may assume that

$$\rho_0, \rho_{D+1} \leq 4\gamma D \sqrt{1 + \frac{\beta^2}{36\gamma^2}} \leq 4\gamma D \left(1 + \frac{\beta^2}{72\gamma^2}\right).$$

In this case, by (2), there must exist an index $j \in [D]$ such that

$$\rho_j \geq \left(1 - \frac{8\beta^2}{72\gamma^2}\right)\gamma = \left(1 - \frac{\beta^2}{9\gamma^2}\right)\gamma.$$

Now, consider a vertical edge $(u^{(j+1)}, v^{(j+1)})$ with $\|\bar{f}(u^{(j)}) - \bar{f}(v^{(j)})\| \geq \beta$, and let

$$u' = \bar{f}(u^{(j)}) + \langle z, \bar{f}(u^{(j)}) - \bar{f}(u^{(j+1)}) \rangle z.$$

From (F4), and the Pythagorean inequality we have,

$$\begin{aligned} \max(\|\bar{f}(u^{(j)}) - \bar{f}(u^{(j+1)})\|^2, \|\bar{f}(u^{(j)}) - \bar{f}(v^{(j+1)})\|^2) &= \\ \|\bar{f}(u^{(j)}) - u'\|^2 + \max(\|u' - \bar{f}(u^{(j+1)})\|^2, \|u' - \bar{f}(v^{(j+1)})\|^2) & \\ \geq \rho_j^2 + \frac{\beta^2}{4} & \\ \geq \left(1 - \frac{2\beta^2}{9\gamma^2}\right)\gamma^2 + \frac{\beta^2}{4} & \\ \geq \gamma^2 + \frac{\beta^2}{36}, & \end{aligned}$$

again verifying (1) for one of the two edges $(u^{(j)}, v^{(j+1)})$ or $(u^{(j)}, u^{(j+1)})$. \square

The following lemma is well-known, and follows from the variational characterization of eigenvalues (see, e.g. [15, Ch. 15]).

Lemma 3.2. *If $G = (V, E)$ is a d -regular graph with second Laplacian eigenvalue $\mu_2(G)$, then for any mapping $f : V \rightarrow L_2$, we have*

$$\mathbb{E}_{x, y \in V} \|f(x) - f(y)\|^2 \lesssim \frac{d}{\mu_2(G)} \mathbb{E}_{(x, y) \in E} \|f(x) - f(y)\|^2 \quad (3)$$

The next lemma shows that when we use an expander graph, we get a significant increase in stretch for edges of \tilde{G} .

Lemma 3.3. *Let $G = (V, E)$ be a d -regular vertex-transitive graph with $m = |V|$ and $\mu_2 = \mu_2(G)$. If $f : V(\tilde{G}) \rightarrow L_2$ is any non-contractive mapping, then there exists a horizontal edge $(x, y) \in E(\tilde{G})$ with*

$$\frac{\|f(x) - f(y)\|^2}{d_{\tilde{G}}(x, y)^2} \geq \frac{\|f(s) - f(t)\|^2}{d_{\tilde{G}}(s, t)^2} + \Omega\left(\frac{\mu_2}{d}(\log_d m)^2\right). \quad (4)$$

Proof. We need only prove the existence of an $(x, y) \in E(\tilde{G})$ such that (4) is satisfied for \bar{f} (as defined in Lemma 3.1), as this implies it is also satisfied for f (possibly for some other edge (x, y)).

Consider any layer $G^{(i)}$ in \tilde{G} , for $i \in [D + 1]$. Applying (3) and using the fact that f is non-contracting, we have

$$\begin{aligned} \mathbb{E}_{(u,v) \in E} \|\bar{f}(u^{(i)}) - \bar{f}(v^{(i)})\|^2 &= \mathbb{E}_{(u,v) \in E} \|f(u^{(i)}) - f(v^{(i)})\|^2 \\ &\gtrsim \frac{\mu_2}{d} \mathbb{E}_{u,v \in V} \|f(u^{(i)}) - f(v^{(i)})\|^2 \\ &\geq \frac{\mu_2}{d} \mathbb{E}_{u,v \in V} d_G(u, v)^2 \\ &\gtrsim \frac{\mu_2}{d} (\log_d m)^2. \end{aligned}$$

In particular, in every layer $i \in [D + 1]$, at least one vertical edge $(u^{(i)}, v^{(i)})$ has $\|\bar{f}(u^{(i)}) - \bar{f}(v^{(i)})\| \gtrsim \sqrt{\frac{\mu_2}{d}} \log_d m$. Therefore the desired result follows from Lemma 3.1. \square

We now to come our main theorem.

Theorem 3.4. *If $G = (V, E)$ is a d -regular, m -vertex, vertex-transitive graph with $\mu_2 = \mu_2(G)$, then*

$$c_2(\tilde{G}^{\otimes k}) \gtrsim \sqrt{\frac{\mu_2 k}{d}} \log_d m.$$

Proof. Let $f : V(\tilde{G}^{\otimes k}) \rightarrow L_2$ be any non-contracting embedding. The theorem follows almost immediately by induction: Consider the top level copy of \tilde{G} in $\tilde{G}^{\otimes k}$, and call it G_0 . Let $(x, y) \in E(G_0)$ be the horizontal edge for which $\|f(x) - f(y)\|$ is longest. Clearly this edge spans a copy of $\tilde{G}^{\otimes k-1}$, which we call G_1 . By induction and an application of Lemma 3.3, there exists a (universal) constant $c > 0$ and an edge $(u, v) \in E(G_1)$ such that

$$\begin{aligned} \frac{\|f(u) - f(v)\|^2}{d_{\tilde{G}^{\otimes k}}(u, v)^2} &\geq \frac{c\mu_2(k-1)}{d} (\log_d m)^2 + \frac{\|f(x) - f(y)\|^2}{d_{\tilde{G}^{\otimes k}}(x, y)^2} \\ &\geq \frac{c\mu_2(k-1)}{d} (\log_d m)^2 + \frac{c\mu_2}{d} (\log_d m)^2 + \frac{\|f(s) - f(t)\|^2}{d_{\tilde{G}^{\otimes k}}(s, t)^2}, \end{aligned}$$

completing the proof. \square

Corollary 3.5. *If $G = (V, E)$ is an $O(1)$ -regular m -vertex, vertex-transitive graph with $\mu_2 = \Omega(1)$, then*

$$c_2(\tilde{G}^{\otimes k}) \gtrsim \sqrt{k} \log m \approx \sqrt{\log m \log N},$$

where $N = |V(\tilde{G}^{\otimes k})| = 2^{\Theta(k \log m)}$.

We remark that infinite families of $O(1)$ -regular vertex-transitive graphs with $\mu_2 \geq \Omega(1)$ are well-known. In particular, one can take any construction coming from the Cayley graphs of finitely generated groups. We refer to the survey [6]; see, in particular, Margulis' construction in Section 8.

3.1 Extension to other L_p spaces

Our previous lower bound dealt only with L_2 . We now prove the following.

Theorem 3.6. *If $G = (V, E)$ is an $O(1)$ -regular m -vertex, vertex-transitive graph with $\mu_2 = \Omega(1)$, for any $p > 1$, there exists a constant $C(p)$ such that*

$$c_p(\tilde{G}^{\otimes k}) \gtrsim C(p)k^{1/q} \log m \approx C(p)(\log m)^{1-1/q}(\log N)^{1/q}$$

where $N = |V(\tilde{G}^{\otimes k})|$ and $q = \max\{p, 2\}$.

The only changes required are to Lemma 3.2 and Lemma 3.1 (which uses orthogonality). The first can be replaced by Matoušek's [14] Poincaré inequality: If $G = (V, E)$ is an $O(1)$ -regular expander graph with $\mu_2 = \Omega(1)$, then for any $p \in [1, \infty)$ and $f : V \rightarrow L_p$,

$$\mathbb{E}_{x,y \in V} \|f(x) - f(y)\|_p^p \leq O(2p)^p \mathbb{E}_{(x,y) \in E} \|f(x) - f(y)\|_p^p.$$

Generalizing Lemma 3.1 is more involved. We need the following well-known 4-point inequalities for L_p spaces.

Lemma 3.7. *Consider any $p \geq 1$ and $u, v, w, x \in L_p$. If $1 \leq p \leq 2$, then*

$$\|u - w\|_p^2 + (p-1)\|x - v\|_p^2 \leq \|u - v\|_p^2 + \|v - w\|_p^2 + \|x - w\|_p^2 + \|u - x\|_p^2. \quad (5)$$

If $p \geq 2$, then

$$\|u - w\|_p^p + \|x - v\|_p^p \leq 2^{p-2} (\|u - v\|_p^p + \|v - w\|_p^p + \|x - w\|_p^p + \|u - x\|_p^p). \quad (6)$$

Proof. The following inequalities are known for $a, b \in L_p$ (see, e.g. [3]). If $1 \leq p \leq 2$, then

$$\left\| \frac{a+b}{2} \right\|_p^2 + (p-1) \left\| \frac{a-b}{2} \right\|_p^2 \leq \frac{\|a\|_p^2 + \|b\|_p^2}{2}.$$

On the other hand, if $p \geq 2$, then

$$\left\| \frac{a+b}{2} \right\|_p^p + \left\| \frac{a-b}{2} \right\|_p^p \leq \frac{\|a\|_p^p + \|b\|_p^p}{2}.$$

In both cases, the desired 4-point inequalities are obtained by averaging two incarnations of one of the above inequalities with $a = u - v, b = v - w$ and then $a = u - x, b = x - w$ and using convexity of the L_p norm (see, e.g. [12, Lem. 2.1]). \square

Lemma 3.8. *Let G be a vertex transitive graph, and suppose $p > 1$. If $q = \max\{p, 2\}$, then there exists a constant $K(p) > 0$ such that the following holds. Let $f : V(\tilde{G}) \rightarrow L_p$ be an injective mapping and define $\bar{f} : V(\tilde{G}) \rightarrow L_p$ by*

$$\bar{f}(x) = \frac{1}{|\text{Aut}(G)|^{1/p}} \left(f(\tilde{\pi}x) \right)_{\pi \in \text{Aut}(G)}.$$

Suppose that β is such that for every $i \in [D+1]$, there exists a vertical edge $(u^{(i)}, v^{(i)})$ which satisfies $\|\bar{f}(u^{(i)}) - \bar{f}(v^{(i)})\|_p \geq \beta$. Then there exists a horizontal edge $(x, y) \in E(\tilde{G})$ such that

$$\frac{\|\bar{f}(x) - \bar{f}(y)\|_p^q}{d_{\tilde{G}}(x, y)^q} \geq \frac{\|f(s) - f(t)\|_p^q}{d_{\tilde{G}}(s, t)^q} + K(p)\beta^q. \quad (7)$$

Proof. Let $D = \text{diam}(G)$. For simplicity, we assume that D is even in what follows.

$$(F1) \quad \|\bar{f}(s) - \bar{f}(t)\|_p = \|f(s) - f(t)\|_p$$

$$(F2) \quad \text{For all } u, v \in V,$$

$$\begin{aligned} \|\bar{f}(s) - \bar{f}(v^{(1)})\|_p &= \|\bar{f}(s) - \bar{f}(u^{(1)})\|_p, \\ \|\bar{f}(t) - \bar{f}(v^{(D+1)})\|_p &= \|\bar{f}(t) - \bar{f}(u^{(D+1)})\|_p. \end{aligned}$$

$$(F3) \quad \text{For every } u, v \in V, i \in [D],$$

$$\|\bar{f}(v^{(i)}) - \bar{f}(v^{(i+1)})\|_p = \|\bar{f}(u^{(i)}) - \bar{f}(u^{(i+1)})\|_p.$$

Fix some $r \in V$ and let $\rho_0 = \|\bar{f}(s) - \bar{f}(r^{(1)})\|_p$, $\rho_i = \|\bar{f}(r^{(2i-1)}) - \bar{f}(r^{(2i+1)})\|_p$ for $i = 1, \dots, D/2$, $\rho_{D/2+1} = \|\bar{f}(t) - \bar{f}(r^{(D+1)})\|_p$. Also let $\rho_{i,1} = \|\bar{f}(r^{(2i-1)}) - \bar{f}(r^{(2i)})\|_p$ and $\rho_{i,2} = \|\bar{f}(r^{(2i)}) - \bar{f}(r^{(2i+1)})\|_p$ for $i = 1, \dots, D/2$.

Note that, by (F2) and (F3) above, the values $\{\rho_i\}$ do not depend on the representative $r \in V$. In this case, we have

$$\sum_{i=0}^{D/2+1} \rho_i \geq \|\bar{f}(s) - \bar{f}(t)\|_p = 9\gamma D, \quad (8)$$

where we put $\gamma = \frac{\|f(s) - f(t)\|_p}{d_{\bar{G}}(s, t)}$. Note that $\gamma > 0$ since f is injective.

Let $\delta = \delta(p)$ be a constant to be chosen shortly. Recalling that $d_{\bar{G}}(s, t) = 9D$ and $d_{\bar{G}}(s, r^{(1)}) = 4D$, observe that if $\rho_0^q \geq \left(1 + \delta \frac{\beta^q}{\gamma^q}\right) (4\gamma D)^q$, then

$$\max \left(\frac{\|\bar{f}(s) - \bar{f}(s')\|_p^q}{d_{\bar{G}}(s, s')^q}, \frac{\|\bar{f}(s') - \bar{f}(r^{(1)})\|_p^q}{d_{\bar{G}}(s', r^{(1)})^q} \right) \geq \gamma^q + \delta \beta^q,$$

verifying (7). The symmetric argument holds for $\rho_{D/2+1}$, thus we may assume that

$$\rho_0, \rho_{D/2+1} \leq 4\gamma D \left(1 + \delta \frac{\beta^q}{\gamma^q}\right)^{1/q} \leq 4\gamma D \left(1 + \delta \frac{\beta^q}{\gamma^q}\right).$$

Similarly, we may assume that $\rho_{i,1}, \rho_{i,2} \leq \gamma \left(1 + \delta \frac{\beta^q}{\gamma^q}\right)^{1/q}$ for every $i \in [D/2]$.

In this case, by (8), there must exist an index $j \in \{1, 2, \dots, D/2\}$ such that

$$\rho_j \geq \left(1 - 8\delta \frac{\beta^q}{\gamma^q}\right) 2\gamma.$$

Now, consider a vertical edge $(u^{(2j)}, v^{(2j)})$ with $\|f(u^{(2j)}) - f(v^{(2j)})\|_p \geq \beta$. Also consider the vertices $v^{(2j-1)}$ and $v^{(2j+1)}$. We now replace the use of orthogonality ((F4) in Lemma 3.1) with Lemma 3.7.

We apply one of (5) or (6) of these two inequalities with $x = f(u^{(2j)}), v = f(v^{(2j)}), u = f(v^{(2j-1)}), w = f(v^{(2j+1)})$. In the case $p \geq 2$, we use (5) to conclude that

$$\begin{aligned} \|f(u^{(2j)}) - f(v^{(2j-1)})\|_p^p + \|f(u^{(2j)}) - f(v^{(2j+1)})\|_p^p &\geq 2^{-p+2} \rho_j^p + 2^{-q+2} \beta^p - \rho_{j,1}^p - \rho_{j,2}^p \\ &\geq 2\gamma^p + 2^{-p+2} \beta^p - 34\delta p \beta^p. \end{aligned}$$

Thus choosing $\delta = \frac{2^{1-p}}{34p}$ yields the desired result for one of $(u^{(2j)}, v^{(2j-1)})$ or $(u^{(2j)}, v^{(2j+1)})$.

In the case $1 \leq p \leq 2$, we use (6) to conclude that

$$\|f(u^{(2j)}) - f(v^{(2j-1)})\|_p^2 + \|f(u^{(2j)}) - f(v^{(2j+1)})\|_p^2 \geq \rho_j^2 + (p-1)\beta^2 - \rho_{j,1}^2 - \rho_{j,2}^2.$$

A similar choice of δ again yields the desired result. \square

References

- [1] S. Arora, J. R. Lee, and A. Naor. Euclidean distortion and the Sparsest Cut. *J. Amer. Math. Soc.*, 21(1):1–21, 2008.
- [2] Y. Aumann and Y. Rabani. An $O(\log k)$ approximate min-cut max-flow theorem and approximation algorithm. *SIAM J. Comput.*, 27(1):291–301 (electronic), 1998.
- [3] K. Ball, E. A. Carlen, and E. H. Lieb. Sharp uniform convexity and smoothness inequalities for trace norms. *Invent. Math.*, 115(3):463–482, 1994.
- [4] S. Chawla, A. Gupta, and H. Räcke. An improved approximation to sparsest cut. In *Proceedings of the 16th Annual ACM-SIAM Symposium on Discrete Algorithms*, Vancouver, 2005. ACM.
- [5] A. Gupta, R. Krauthgamer, and J. R. Lee. Bounded geometries, fractals, and low-distortion embeddings. In *44th Symposium on Foundations of Computer Science*, pages 534–543, 2003.
- [6] S. Hoory, N. Linial, and A. Wigderson. Expander graphs and their applications. *Bull. Amer. Math. Soc. (N.S.)*, 43(4):439–561 (electronic), 2006.
- [7] R. Krauthgamer, J. R. Lee, M. Mendel, and A. Naor. Measured descent: A new embedding method for finite metrics. *Geom. Funct. Anal.*, 15(4):839–858, 2005.
- [8] T. J. Laakso. Plane with A_∞ -weighted metric not bi-Lipschitz embeddable to \mathbb{R}^N . *Bull. London Math. Soc.*, 34(6):667–676, 2002.
- [9] U. Lang and C. Plaut. Bilipschitz embeddings of metric spaces into space forms. *Geom. Dedicata*, 87(1-3):285–307, 2001.
- [10] J. R. Lee. On distance scales, embeddings, and efficient relaxations of the cut cone. In *SODA '05: Proceedings of the sixteenth annual ACM-SIAM symposium on Discrete algorithms*, pages 92–101, Philadelphia, PA, USA, 2005. Society for Industrial and Applied Mathematics.
- [11] J. R. Lee. Volume distortion for subsets of Euclidean spaces. *Discrete Comput. Geom.*, 41(4):590–615, 2009.
- [12] J. R. Lee and A. Naor. Embedding the diamond graph in L_p and dimension reduction in L_1 . *Geom. Funct. Anal.*, 14(4):745–747, 2004.
- [13] N. Linial, E. London, and Y. Rabinovich. The geometry of graphs and some of its algorithmic applications. *Combinatorica*, 15(2):215–245, 1995.
- [14] J. Matoušek. On embedding expanders into l_p spaces. *Israel J. Math.*, 102:189–197, 1997.

- [15] J. Matoušek. *Lectures on discrete geometry*, volume 212 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 2002.
- [16] I. Newman and Y. Rabinovich. A lower bound on the distortion of embedding planar metrics into Euclidean space. *Discrete Comput. Geom.*, 29(1):77–81, 2003.
- [17] S. Rao. Small distortion and volume preserving embeddings for planar and Euclidean metrics. In *Proceedings of the 15th Annual Symposium on Computational Geometry*, pages 300–306, New York, 1999. ACM.