

The Clique Problem in Ray Intersection Graphs*

Sergio Cabello[†]

Jean Cardinal[‡]

Stefan Langerman[§]

October 31, 2018

Abstract

Ray intersection graphs are intersection graphs of rays, or halflines, in the plane. We show that any planar graph has an even subdivision whose complement is a ray intersection graph. The construction can be done in polynomial time and implies that finding a maximum clique in a segment intersection graph is NP-hard. This solves a 21-year old open problem posed by Kratochvíl and Nešetřil.

arXiv:1111.5986v1 [cs.CG] 25 Nov 2011

*Research was supported in part by the Slovenian Research Agency, program P1-0297 and projects J1-4106, L7-4119, BI-BE/11-12-F-002, and within the EUROCORES Programme EUROGIGA (projects GReGAS and ComPoSe) of the European Science Foundation. Large part of the work was done while Sergio was visiting ULB.

[†]Department of Mathematics, IMFM, and Department of Mathematics, FMF, University of Ljubljana, Slovenia. email: sergio.cabello@fmf.uni-lj.si

[‡]Université Libre de Bruxelles (ULB), Brussels, Belgium. Email: jcardin@ulb.ac.be

[§]Maitre de Recherches, Fonds de la Recherche Scientifique (F.R.S-FNRS), Université Libre de Bruxelles (ULB), Brussels, Belgium. Email: stefan.langerman@ulb.ac.be

1 Introduction

The intersection graph of a collection of sets has one vertex for each set, and an edge between two vertices whenever the corresponding sets intersect. Of particular interest are families of intersection graphs corresponding to geometric sets in the plane. In this contribution, we will focus on *segment intersection graphs*, intersection graphs of line segments in the plane.

In a seminal paper, Kratochvíl and Nešetřil [11] proposed to study the complexity of two classical combinatorial optimization problems, the maximum independent set and the maximum clique, in geometric intersection graphs. While those problems are known to be hard to approximate in general graphs (see for instance [5, 13]), their restriction to geometric intersection graphs may be more tractable. They proved that the maximum independent set problem remains NP-hard for segment intersection graphs, even if those segments have only two distinct directions. It was also shown that in that case, the maximum clique problem can be solved in polynomial time. The complexity of the maximum clique problem in general segment intersection graphs was left as an open problem, and remained so until now. In their survey paper “On six problems posed by Jarik Nešetřil” [3], Bang-Jensen *et al.* describe this problem as being “among the most tantalizing unsolved problem in the area”.

Some progress has been made in the meanwhile. In 1992, Middendorf and Pfeiffer [12] showed, with a simple proof, that the maximum clique problem was NP-hard for intersection graphs of 1-intersecting curve segments that are either line segments or curves made of two orthogonal line segments. They also give a polynomial time dynamic programming algorithm for the special case of line segments with endpoints of the form $(x, 0), (y, i)$, with $i \in \{1, \dots, k\}$ for some fixed k . Another step was made by Ambühl and Wagner [2] in 2005, who showed that the maximum clique problem was NP-hard for intersection graphs of ellipses of fixed, arbitrary, aspect ratio. Unfortunately, this ratio must be bounded, which excludes the case of segments.

Our results. We prove that the maximum clique problem in segment intersection graphs is NP-hard. In fact, we prove the stronger result that the problem is NP-hard even in *ray intersection graphs*, defined as intersection graphs of rays, or halflines, in the plane. This complexity result is a consequence of the following structural lemma: every planar graph has an even subdivision whose complement is a ray intersection graph. Furthermore, the corresponding set of rays has a natural polynomial size representation. Hence solving the maximum clique problem in this graph allows to recover the maximum independent set in the original planar graph, a task well known to be NP-hard [8]. The construction is detailed in Section 2.3.

Related work. We prove that the complement of some subdivision of any planar graph can be represented as a segment intersection graph. Whether the complement of every planar graph is a segment intersection graph remains an open question. In 1998, Kratochvíl and Kuběna [9] showed that the complement of any planar graph is the intersection graph of a set of convex polygons. More recently, Francis, Kratochvíl, and Vyskočil [7] proved that the complement of any partial 2-tree is a segment intersection graph. Partial 2-trees are planar, and in particular every outerplanar graph is a partial 2-tree. The representability of planar graphs by segment intersection graphs, formerly known as Scheinerman’s conjecture, was proved recently by Chalopin and Gonçalves [4].

The maximum independent set problem in intersection graphs has been studied by Agarwal and Mustafa [1]. In particular, they proved that it could be approximated within a factor $n^{1/2+o(1)}$ in polynomial time for segment intersection graphs. This has been recently improved by Fox and Pach [6], who described, for any $\epsilon > 0$, a n^ϵ -approximation algorithm. In fact, their technique also

applies to the maximum clique problem, and therefore n^ϵ is the best known approximation factor for this problem too.

In 1994, Kratochvíl and Matoušek [10] proved that the recognition problem for segment intersection graphs was in PSPACE, and was also NP-hard. It is still not clear whether it is NP-complete.

Notation. For any natural number m we use $[m] = \{1, \dots, m\}$. In a graph G , a *rotation system* is a list $\pi = (\pi_v)_{v \in V(G)}$, where each π_v fixes the clockwise order of the edges of $E(G)$ incident to v . When G is an embedded planar graph, the embedding uniquely defines a rotation system, which is often called a combinatorial embedding. For the rest of the paper we use *ray* to refer to an *open ray*, that is, a ray does not contain its origin. Therefore, whenever two rays intersect they do so in the relative interior of both. Since our construction does not use degeneracies, we are not imposing any restriction by considering only open rays. A subdivision of a graph G is said to be *even* if each edge of G is subdivided an even number of times.

2 Construction

Let us start providing an overview of the approach. We first construct a set of curves that will form the *reference frame*. This construction is quite generic and depends only on a parameter $k \in \mathbb{N}$. We then show that the complement of any tree has a special type of representation, called *snooker representation*, which is constructed iteratively over the levels of the tree. The number of levels of the tree is closely related to k , the parameter used for the reference frame. We then argue that if G is a planar graph that consists of a tree T and a few, special paths of length two and three, then the complement of G can be represented as an intersection graph of rays by extending a snooker representation of T . Finally, we argue that any planar graph has an even subdivision that can be decomposed into a tree and a set of paths of length two and three with the required properties.

We first describe the construction using real coordinates. The construction does not rely on degeneracies, and thus we can slightly perturb the coordinates used in the description. This perturbation is enough to argue that a representation can be computed in polynomial time. Then, using a relation between the independence number of a graph G and an even subdivision of G , we obtain that computing a maximum clique in a ray intersection graph is NP-hard.

2.1 Reference frame

Let k be an *odd* number to be chosen later. We set $\theta = \frac{k-1}{k}\pi$, and define for $i = 0, \dots, k-1$ the points

$$p_i = (\cos(i \cdot \theta), \sin(i \cdot \theta)).$$

The points p_i lie on a unit circle centered at the origin; see Figure 1. For each $i \in [k-2]$ we construct a rectangle R_i as follows. Let q_i be the point $p_i + (p_i - p_{i+1})$, symmetric of p_{i+1} with respect to p_i , let m_i be the midpoint between p_i and q_i , and let t_i be, among the two points along the line through q_{i+1} and m_i with the property $|m_i t_i| = |m_i p_i|$, the one that is furthest from q_{i+1} . We define R_i to be the rectangle with vertices p_i , t_i , and q_i . The fourth vertex of R_i is implicit and is denoted by r_i . Any two rectangles R_i and R_j are congruent with respect to a rotation around the origin. We have constructed the rectangles R_i in such way that, for any $i \in [k-2]$, the line supporting the diagonal $p_i q_i$ of R_i contains p_{i+1} and the line supporting the diagonal $r_i t_i$ of R_i contains q_{i+1} .

For each $i \in [k-2]$, let α_i be the arc of circle that is tangent to both diagonals of R_i and has endpoints t_i and r_i (see Figure 2). Note that the curves α_i and the rectangles R_i have been chosen




Figure 1: The points p_0, \dots, p_k and the rectangles R_1, \dots, R_{k-2} .

so that any line that intersects α_i twice or is tangent to the curve α_i must intersect the curve α_{i+1} . For any $i \in [k-2]$, let Γ_i be the set of rays that intersect α_i twice or are tangent to α_i and have its origin on α_{i+1} . We also define Γ_0 as the set of rays with origin on α_1 and passing through p_0 . The rays of Γ_i that are tangent to α_i will play a special role. In fact, we will only use rays of Γ_i that are “near-tangent” to α_i .

Lemma 2.1. *When $|j-i| > 1$, any ray from Γ_i intersects any ray from Γ_j .*

Lemma 2.2. *Any ray tangent to α_{i+1} at the point $x \in \alpha_{i+1}$ intersects any ray from Γ_i , except those having their origin at x .*

Note that the whole construction depends only on the parameter k . We will refer to it as *reference frame*.

2.2 Complement of Trees

Let T be a graph with a rotation system π_T , let r vertex in T and let $rs \in E(T)$ be an arbitrary edge incident to r . The triple (π_T, r, rs) induces a natural linear order $\tau = \tau(\pi_T, r, rs)$ on the vertices of T . This order τ corresponds to the order followed by a breadth-first traversal of T from r with the following additional restrictions:

- (i) s is the second vertex;




Figure 2: The circular arcs $\alpha_1, \alpha_2, \dots$

- (ii) the children of any vertex v are visited according to the clockwise order π_v ;
- (iii) if $v \neq r$ has parent v' , the first child of u of v is such that vu is the successor of vv' in the clockwise order π_v .

We say that vertices v and v' at the same level are *consecutive* when they are consecutive in τ . See Figure 3. The linear order will be fixed through our discussion, so we will generally drop it from the notation. Henceforth, whenever we talk about a tree T and a linear order τ on $V(T)$, we assume that τ is the natural linear order induced by a triple (π_T, r, rs) . In fact, the triple (π_T, r, rs) is implicit in τ . For any vertex v we use v^+ for its successor and v^- for its predecessor.

A *snooker representation* of the complement of an embedded tree T with linear order τ is a representation of T with rays that satisfies the following properties:

- (a) Each vertex v at level i in T is represented by a ray γ_v from Γ_i . Thus, the origin of γ_v , denoted by a_v , is on α_{i+1} . Note that this implies that k is larger than the depth of T .
- (b) If a vertex u has parent v , then γ_u passes through the origin a_v of γ_v . (Here it is relevant that we consider all rays to be open, as otherwise γ_u and γ_v would intersect.) In particular, all rays corresponding to the children of v pass through the point a_v .
- (c) The origins of rays corresponding to consecutive vertices u and v of level i are consecutive along α_{i+1} . That is, no other ray in the representation has its origin on α_{i+1} and between the origins γ_u and γ_v .




Figure 3: In this example, assuming that π is as drawn, the linear order τ is $r, s, v_1^1, \dots, v_1^5, v_2^1, \dots$

Lemma 2.3. *The complement of any embedded tree with a linear order τ has a snooker representation.*

Proof. Consider a reference frame with k larger than the depth of T . The construction we provide is iterative over the levels of T . Note that, since we provide a snooker representation, it is enough to tell for each vertex $v \neq r$ the origin a_v of the ray γ_v . Property (b) of the snooker representation provides another point on the ray γ_v , and thus γ_v is uniquely defined. The ray γ_r for the root r is the ray of Γ_0 with origin a_r in the center of α_1 .

Consider any level $i > 1$ and assume that we have a representation of the vertices at level $i - 1$. Consider a vertex v at level $i - 1$ and let u_1, \dots, u_d denote its d children. If the successor v^+ of v is also at level $i - 1$, we take $a_v^+ = a_{v^+}$, and else we take a_v^+ to be an endpoint of α_i such that no other origin is between the endpoint and a_v . See Figure 4. Similarly, if the predecessor v^- of v is at level $i - 1$, we take $a_v^- = a_{v^-}$, and else we take a_v^- to be an endpoint of α_i such that no other origin is between the endpoint and a_v . (If v is the only one vertex at level i , we also make sure that $a_v^- \neq a_v^+$.) Let ℓ_v^+ be the line through a_v and a_v^+ . Similarly, let ℓ_v^- be the line through a_v and a_v^- . We then choose the points a_{u_1}, \dots, a_{u_d} on the portion of α_{i+1} contained between ℓ_v^+ and ℓ_v^- such that the $d + 2$ points $\ell_v^- \cap \alpha_{i+1}, a_{u_1}, \dots, a_{u_d}, \ell_v^+ \cap \alpha_{i+1}$ are regularly spaced. Since the ray γ_{u_j} has origin a_{u_j} and passes through a_v , this finishes the description of the procedure. Because a_{u_j} lies between ℓ_v^+ and ℓ_v^- , the ray γ_{u_j} either intersects α_i twice or is tangent to α_i , and thus $\gamma_{u_j} \in \Gamma_i$.

Recall that any ray from Γ_i intersects any ray from Γ_j when $|j - i| > 1$. Therefore, vertices from levels i and j , where $|i - j| > 1$, intersect. For vertices u and v at levels $i - 1$ and i , respectively, the convexity of the curve α_i and the choices for a_v imply that γ_v intersects γ_u if and only if u is not the parent of v in T . For vertices u and v at the same level i , the rays γ_u and γ_v intersect: if they have the same parent w , then they intersect on a_w , and if they have different parents, the order of their origins a_u and a_v on α_{i+1} and the order of their intersections with α_i are reversed. \square

2.3 A tree with a few short paths

Let T be an embedded tree with a linear order τ . An *admissible extension* of T is a graph P with the following properties

- P is the union of vertex-disjoint paths (i.e., two paths don't share internal vertices but they are allowed to share endpoints);




Figure 4: An example showing the construction of a snooker representation when v has three children and v^+ and v^- are at the same level as v .

- each maximal path in P has 3 or 4 vertices;
- the endpoints of each maximal path in P are leaves of T that are consecutive and at the same level;
- the internal vertices of any path in P are not vertices of $V(T)$.

Note that $T + P$ is a planar graph because we only add paths between consecutive leaves.

Lemma 2.4. *Let T be an embedded tree and let P be an admissible extension of T . The complement of $T + P$ is a ray intersection graph.*

Proof. We construct a snooker representation of T using Lemma 2.3 where k , the number of levels, is the depth of T plus 2 or 3, whichever is odd. We will use a local argument to represent each maximal path of P , and each maximal path of P is treated independently. It will be obvious from the construction that rays corresponding to vertices in different paths intersect. We distinguish the case where the maximal path has one internal vertex or two.

Consider first the case of a maximal path in P with one internal vertex. Thus, the path is uvw where u and v are consecutive leaves in T and $w \notin V(T)$ is not yet represented by a ray. The origins a_u and a_v of the rays γ_u and γ_v , respectively, are distinct and consecutive along α_{i+1} because we have a snooker representation. We thus have the situation depicted in Figure 5. We can then just take the γ_w to be the line through a_u and a_v . (This line can also be a ray with an origin sufficiently far away.) This line intersects the ray of any other vertex, different that γ_u and γ_v .

Consider now the case of a maximal path in P with two internal vertices. Thus, the path is $uww'v$ where u and v are consecutive leaves in T and $w, w' \notin V(T)$. In this case, the construction depends on the relative position of the origins a_u and a_v , and we distinguish two scenarios: (i) shifting the origin a_u of ray γ_u towards a_v while maintaining the slope introduces an intersection between γ_u and the ray for the parent of u or (ii) shifting the origin a_v of ray γ_v towards a_u while maintaining the slope introduces an intersection between γ_v and the ray for the parent of v . Note that exactly one of the two options must occur.

Let us consider only scenario (i), since the other one is symmetric; see Figure 6. We choose a point b_w on α_{i+1} between a_u and a_v very near a_u and represent w with a ray γ_w parallel to γ_u with origin b_w . Thus γ_w does not intersect γ_u but intersects any other ray because we are in scenario




Figure 5: Case 1 in the proof of Lemma 2.4: adding a path with one internal vertex.

(i). Finally, we represent w' with the line $\gamma_{w'}$ through points b_w and a_v . With this, $\gamma_{w'}$ intersects the interior of any other ray but γ_w and γ_v , as desired. Note that $\gamma_{w'}$ also intersects the rays for vertices in the same level because those rays are near-tangent to α_{i+2} , which is intersected by γ_w .

Note that in every case, since the rays γ_w and $\gamma_{w'}$, respectively, are actually lines, no new ray has its origin on α_{i+2} . Hence rays having consecutive origins on α_{i+2} remain so after the inclusion of a path in level $i + 1$. \square

Lemma 2.5. *Any embedded planar graph G has an even subdivision $T + P$, where T is an embedded tree and P is an admissible extension of T . Furthermore, such T and P can be computed in polynomial time.*

Proof. Let r be an arbitrary vertex in the outer face f of G . Let B be the set of edges in a BFS tree of G from r . With a slight abuse of notation, we also use B for the BFS tree itself. Let $C = E(G) \setminus B$. In the graph G^* dual to G , the edges $C^* = \{c^* \mid c \in C\}$ are a spanning tree of G^* , which with a slight abuse of notation we also denote by C^* . We root C^* at the node f^* , corresponding to the outer face. This is illustrated on Figure 7.

We define for each edge $e \in C$ the number k_e of subdivisions it will undertake using a bottom-up approach. Any edge $e \in C$ that is incident to a leaf of C^* gets assigned $k_e = 4$. For any other edge $e \in C$, we define k_e as 2 plus the maximum $k_{e'}$ over all descendants $(e')^* \in C^*$ of e^* in C^* . Let H be the resulting subdivision. For any edge $e \in C$, let Q_e be the path in H that corresponds to the subdivision of e . We use in H the combinatorial embedding induced by G .

We can now compute the tree T and the paths P . Since B is a BFS tree, every edge $e \in C$ connects two vertices that are either at the same level in B , or at two successive levels. For every edge $e \in C$ that connects two vertices at the same level in B , let P_e be the length-two subpath in the middle of Q_e . For every edge $uv \in C$ that connects vertex u at level i to vertex v at level $i + 1$ in B , let P_e be the length-three subpath obtained from the length-two subpath in the middle of $Q_e - u$. We then take $P = \bigcup_{e \in C} P_e$ and take T to be the graph $H - P$, after removing isolated vertices. In T we use the rotation system inherited from H and use the edge rs to define the linear order, where rs is an edge in f .

It is clear from the construction that $T + P = H$ is an even subdivision of G . We have to check that P is indeed an admissible extension of T . The maximal paths of P are vertex disjoint and




Figure 6: Case 2(i) in the proof of Lemma 2.4: adding a path with two internal vertices.




Figure 7: Figure for the proof of Lemma 2.5. (a) A planar graph with a BFS from r in bold. (b) The spanning tree C^* of the dual graph. (c) The subdivision of the edges from C . (d) The resulting subdivided graph with P marked in red and thinner lines. (e) The resulting graph T after the removal of P , drawn such that the height of the vertices corresponds to their level.

connect leaves of P because the paths Q_e , $e \in C$, are edge-disjoint and each P_e is strictly contained in the interior of Q_e . Since in $H - P$ we removed isolated vertices, it is clear that no internal vertex of a path of P is in T . The graph T is indeed a tree because, for every edge $e \in C$, we have removed some edges from its subdivision Q_e . Since B is a BFS tree and P_{uv} is centered within Q_{uv} , when u and v are at the same level, or within $Q_{uv} - u$, when u is one level below v , the maximal paths of P connect vertices that are equidistant from r in H .

It remains to show that the endpoints of any maximal path in P are consecutive vertices in T . This is so because of the inductive definition of k_e . The base case is when $e \in C$ is incident to a leaf of C^* , and is subdivided $k_e = 4$ times. The edge e either connects two vertices at the same level in B , or at two successive levels. In both cases it can be checked that P_e connects two consecutive vertices in T . The inductive case is as follows. We let k_e be equal to 2 plus the maximum $k_{e'}$ over all descendants $(e')^* \in C^*$ of e^* in C^* . By induction, all the corresponding $P_{e'}$ connect two consecutive vertices of T , say at level i in T . By definition, P_e will connect two consecutive vertices of T at level $i + 1$.

The construction of T and P only involves computing the BFS tree B , the spanning tree C^* , and the values k_e , which can clearly be done in polynomial time. \square

Combining lemmas 2.4 and 2.5 directly yields the following.

Theorem 2.6. *Any planar graph has an even subdivision whose complement is a ray intersection graph. Furthermore, this subdivision can be computed in polynomial time.*

2.4 Polynomial-time construction

The construction of the ray intersection graph in Lemmas 2.3 and 2.4, uses real coordinates. We wish to prove that the maximum clique problem is NP-hard even when a geometric description of the ray intersection graph is given as input. Hence we need to argue how to carry out this construction using integer coordinates, each using a polynomial number of bits. In what follows, we let $n = |V(T + P)|$.

Lemma 2.7. *In the construction of Lemma 2.4,*

- *any two points are at distance at least $n^{-O(n)}$ and at most $O(n)$;*
- *the distance between any line through two origins and any other point is at least $n^{-O(n)}$, unless the three points are collinear.*

Proof. Recall that the circle containing points p_0, p_1, \dots has radius 1. The rectangles R_i are all congruent and have two diagonals of length $|p_0 p_1| = \Theta(1)$. The small side of rectangle R_i has size $\Theta(1/n)$ and both diagonals of R_i form an angle of $\Theta(1/n)$. It follows that the center of the circles supporting α_i have coordinates $\Theta(n)$. For points from different curves α_i there is at least a separation of $\Theta(1/n)$.

We first bound the distance between the origins for the rays representing vertices of T . Let us refer to the origins of rays lying on α_i and the extremes of α_i as *features* on the curve α_i . Let δ_i be the minimum separation between any two features on the curve α_i . On the curve α_1 there are three features: the two extremes of α_1 and the origin a_r of γ_r , which is in the middle of α_1 . Since α_1 has length $\Omega(1)$, it follows that $\delta_1 = \Omega(1)$.

We will bound the ratio δ_{i+1}/δ_i for $i \geq 1$. Consider the construction of Lemma 2.3 to determine the features on α_{i+1} . By induction, any two consecutive features along α_i are separated at least by δ_i . Since α_i is supported by a circle of radius $\Theta(n)$, the lines ℓ_v^+ and ℓ_v^- form an angle of at

least $\Omega(\delta_i/n)$. This implies that the points $\ell_v^- \cap \alpha_{i+1}, a_{u_1}, \dots, a_{u_d}, \ell_v^+ \cap \alpha_{i+1}$ have a separation of $\Omega(\delta_i/(nd))$. It follows that $\delta_{i+1} = \Omega(\delta_i/(nd)) = \Omega(\delta_i/n^2)$, and thus $\delta_{i+1}/\delta_i = \Omega(1/n^2)$. Since T has depth at most n , all features are at distance at least $n^{-O(n)}$.

We can now argue that the origins of the rays used in the construction of Lemma 2.4 also have a separation of $n^{-O(n)}$. In Case 1, we just add a line through previous features. In Case 2, it is enough to place b_w at distance $|a_u a_v|/n$ from a_u , and thus the features keep a separation of at least $n^{-O(n)}$.

The second item is a consequence of the fact that if a point (a, b) is not on the line through (x, y) and (x', y') then its distance is at least $|y + \frac{y'-y}{x'-x}(a-x) - b|$. \square

We can now give the following algorithmic version of Lemma 2.4.

Lemma 2.8. *Let T be an embedded tree and let P be an admissible extension of T . We can find in polynomial time a family of rays described with integer coordinates whose intersection graph is isomorphic to the complement of $T + P$.*

Proof. Recall that the construction in Lemma 2.4 consists of first constructing a snooker representation of T , as described in Lemma 2.3, then adding the rays corresponding to the paths in the extension. In the first part, each new point is created by: (a) computing the two intersections between two lines ℓ_v^- and ℓ_v^+ through two existing points and a curve α_{i+1} , then by (b) equally spacing $O(n)$ points between those two points on α_{i+1} (see Figure 4). In the second part, the only new points of the construction are the points b_w added on α_{i+1} , between a_u and a_v (see Figure 6). We refer to this latter case as (c).

Consider that each point is moved by a distance $< \varepsilon$ after it is constructed. This may cause any point further constructed from those to move as well. In cases (b) and (c), the new points would be moved by no more than ε as well.

In case (a) however, the error could be amplified. Let a and b be two previously constructed points, and suppose they are moved to a' and b' , within a radius of ε . By the previous lemma, the distance between a and b is at least $n^{-O(n)}$ and so the angle between the line ab and $a'b'$ is at most $\varepsilon n^{O(n)}$. Because the radius of the supporting circle of α_i is $\Theta(n)$, the distance between $ab \cap \alpha_i$ and $a'b' \cap \alpha_i$ is at most $O(n)\varepsilon n^{O(n)} = \varepsilon n^{O(n)}$.

Therefore, an error in one construction step expands by a factor $n^{O(n)}$. Now observe that each point of type (a) is constructed on the next level, and a point of type (b) is always constructed from points of type (a). Therefore, as there are $O(n)$ levels, an error is propagated at most $O(n)$ times, and the total error propagated from moving each constructed point by a distance $< \varepsilon$ is at most $\varepsilon n^{O(n^2)}$.

By the previous lemma, there is a constant c such that any origin in the construction of Lemma 2.4 is separated at least by a distance $A = 1/n^{cn}$ from any other origin and any ray not incident to it. Therefore, by choosing $\varepsilon = n^{-c'n^3}$ for c' large enough, the total propagation will never exceed A , and therefore perturbing the basic construction points of the reference frame and each further constructed point by a distance $< \varepsilon$ will not change the intersection graph of the modified rays.

Therefore, to construct the required set of rays in polynomial time, we multiply every coordinate by the smallest power of 2 larger than $1/\varepsilon$ and snap every constructed point to the nearest integer while following the construction of Lemma 2.4. Each coordinate can then be represented by $O(n^3)$ bits. \square

Let $\alpha(G)$ be the size of the largest independent set in a graph G . The following simple lemma can be deduced from the observation that subdividing an edge twice increases the independence number by exactly one.

Lemma 2.9. *If G' is an even subdivision of G where each edge $e \in E(G)$ is subdivided $2k_e$ times, then $\alpha(G') = \alpha(G) + \sum_e k_e$*

By combining Lemmas 2.5, 2.8, and 2.9, we obtain:

Theorem 2.10. *Finding a maximum clique in a ray intersection graph is NP-hard, even when the input is given by a geometric representation as a set of rays.*

References

- [1] Pankaj K. Agarwal and Nabil H. Mustafa. Independent set of intersection graphs of convex objects in 2D. *Computational Geometry*, 34(2):83–95, 2006.
- [2] Christoph Ambühl and Uli Wagner. The clique problem in intersection graphs of ellipses and triangles. *Theory Comput. Syst.*, 38(3):279–292, 2005.
- [3] Jorgen Bang-Jensen, Bruce Reed, Mathias Schacht, Robert Sámal, Bjarne Toft, and Uli Wagner. On six problems posed by Jarik Nešetřil. In Martin Klazar, Jan Kratochvíl, Martin Loeb, Jiří Matoušek, and Robin Thomas, editors, *Topics in Discrete Mathematics*, volume 26 of *Algorithms and Combinatorics*, pages 613–627. Springer-Verlag, 2006.
- [4] Jérémie Chalopin and Daniel Gonçalves. Every planar graph is the intersection graph of segments in the plane. In *STOC*, pages 631–638, 2009.
- [5] Lars Engebretsen and Jonas Holmerin. Clique is hard to approximate within $n^{1-o(1)}$. In *Proceedings of the 27th International Colloquium on Automata, Languages and Programming (ICALP'00)*, pages 2–12, 2000.
- [6] Jacob Fox and János Pach. Computing the independence number of intersection graphs. In *SODA*, pages 1161–1165, 2011.
- [7] Mathew C. Francis, Jan Kratochvíl, and Tomáš Vyskočil. Segment representation of a subclass of co-planar graphs. ArXiv e-print:1011.1332, November 2010.
- [8] M. R. Garey and David S. Johnson. The rectilinear steiner tree problem is NP-complete. *SIAM J. Appl. Math.*, 32:826–834, 1977.
- [9] Jan Kratochvíl and Aleš Kuběna. On intersection representations of co-planar graphs. *Discrete Math.*, 178(1-3):251–255, 1998.
- [10] Jan Kratochvíl and Jirí Matoušek. Intersection graphs of segments. *J. Comb. Theory, Ser. B*, 62(2):289–315, 1994.
- [11] Jan Kratochvíl and Jaroslav Nešetřil. Independent set and clique problems in intersection-defined classes of graphs. *Commentationes Mathematicae Universitatis Carolinae*, 31(1):85–93, 1990.
- [12] Matthias Middendorf and Frank Pfeiffer. The max clique problem in classes of string-graphs. *Discrete Math.*, 108(1-3):365–372, 1992.
- [13] Johan Håstad. Clique is hard to approximate within $n^{1-\epsilon}$. In *Proceedings of the 37th Annual Symposium on Foundations of Computer Science (FOCS'96)*, page 627, 1996.