ON THE FUNCTOR ℓ^2

CHRIS HEUNEN

ABSTRACT. We study the functor ℓ^2 from the category of partial injections to the category of Hilbert spaces. The former category is finitely accessible, and its homsets are algebraic domains; the latter category has conditionally algebraic domains for homsets. The functor preserves daggers, monoidal structures, enrichment, and various (co)limits, but has no adjoints. Up to unitaries, its direct image consists precisely of the partial isometries, but its essential image consists of all continuous linear maps between Hilbert spaces.

I am delighted to dedicate this paper to Samson Abramsky, on the occasion of his 60th birthday. Among all the wisdom he has imparted on me is this contradictory gem: "Never solve a problem completely, or noone will have a reason to cite you". My better nature gladly took some time off to let this paper follow his advice.

1. Introduction

The rich theory of Hilbert spaces underpins much of modern functional analysis and therefore quantum physics [25, 20], yet important parts of it have resisted categorical treatment. In any categorical analysis of a species of mathematical objects, free objects of that kind play a significant role. The important ℓ^2 -construction is in many ways the closest thing there is to a free Hilbert space: if X is a set, then

$$\ell^2(X) = \left\{ \varphi \colon X \to \mathbb{C} \; \middle| \; \sum_{x \in X} |\varphi(x)|^2 < \infty \right\}$$

is a Hilbert space, in fact the only one of its dimension up to isomorphism. The ℓ^2 construction can be made into a functor, if we take partial injections as morphisms
between the sets X, as first observed by Barr [6]. Outside functional analysis, it also
plays a historically important role in the geometry of interaction (which has been
noticed by many authors; an incomplete list of references includes [9, 1, 12, 13, 17]).

Explicit categorical properties of the ℓ^2 -construction are few and far between in the literature. These notes gather and augment them in a systematic study. Section 2 starts with the category of Hilbert spaces: it is self-dual, has two monoidal structures, and its homsets are algebraic domains, but its enrichment and limit behaviour is wanting. Section 3 discusses the category of partial injections, which is more well-behaved: it is also self-dual, has two monoidal structures, and is enriched over algebraic domains; moreover, it is finitely accessible. Section 4 introduces and studies the functor ℓ^2 itself. It preserves the self-dualities, monoidal structures, and enrichment. It also preserves (co)kernels and finite (co)products, but not general

Date: September 19, 2018.

The author was supported by U.S. Office of Naval Research Grant Number N000141010357, and would like to thank Samson Abramsky, Rasmus Bentmann, John Bourke, Bart Jacobs, Peter Hines, Robin Kaarsgaard, Martti Karvonen, Prakash Panangaden, Jiří Rosicky, Norbert Schuch, and Jamie Vicary for encouragement and pointers.

(co)limits. Therefore it has no adjoints, and in that sense does not provide free Hilbert spaces. It is faithful and essentially surjective on objects. Section 5 studies the image of the functor ℓ^2 . Up to unitaries, its direct image consists precisely of partial isometries. Remarkably, it is essentially full, that is, its essential image is the whole category of Hilbert spaces.

Choice issues are lurking closely beneath the surface of these results. In fact, $\ell^2(X)$ is not just a Hilbert space; it carries a priviledged orthonormal basis. The functor ℓ^2 is an equivalence between the category of partial injections, and the category of Hilbert spaces with a chosen orthonormal basis and morphisms preserving it. But the latter class of morphisms is too restrictive: all interesting applications of Hilbert spaces require a change of basis. Following the guiding thought "a gentleman does not choose a basis", Section 6 suggests directions for further research.

2. The codomain

Definition 2.1. We are interested in the category **Hilb**, whose objects are complex Hilbert spaces, and whose morphisms are continuous linear functions.

- **2.2.** The category **Hilb** has a *dagger*, that is, a contravariant involutive functor \dagger : **Hilb**^{op} \to **Hilb** that acts as the identity on objects. On a morphism $f: H \to K$ it is given by the unique adjoint $f^{\dagger}: K \to H$ satisfying $\langle f(x) | y \rangle = \langle x | f^{\dagger}(y) \rangle$. For example, an isomorphism u is unitary when $u^{-1} = u^{\dagger}$.
- **2.3.** Furthermore, the usual tensor product of Hilbert spaces provides the category **Hilb** with symmetric monoidal structure. The monoidal unit is the 1-dimensional Hilbert space \mathbb{C} . In fact, **Hilb** has dagger symmetric monoidal structure, i.e. $(f \otimes q)^{\dagger} = f^{\dagger} \otimes q^{\dagger}$, and all coherence isomorphisms are unitaries.
- **2.4.** Direct sums of Hilbert spaces provide the category **Hilb** with (finite) dagger biproducts. That is, $H \oplus K$ is simultaneously a product and a coproduct, the projections are the daggers of the corresponding coprojections, and $(f \oplus g)^{\dagger} = f^{\dagger} \oplus g^{\dagger}$. Similarly, the 0-dimensional Hilbert space is a zero object, i.e. simultaneously initial and terminal.
- **2.5.** Let us emphasize that we take continuous linear maps as morphisms between Hilbert spaces, rather than linear contractions. The category of Hilbert spaces with the latter morphisms is rather well-behaved, see *e.g.* [5]. However, it is the former choice of morphisms that is of interest in functional analysis and quantum physics. Unfortunately it also reduces the limit behaviour of the category **Hilb**, as the following lemma shows.

Lemma 2.6. The category Hilb:

- (i) has (co)equalizers;
- (ii) does not have infinite (co)products;
- (iii) does not have directed (co)limits.

Proof. Part (i) holds because **Hilb** is enriched over abelian groups and has kernels [16]. For (ii), consider the following counterexample. Define an N-indexed family $H_n = \mathbb{C}$ of objects of **Hilb**. Suppose the family (H_n) had a coproduct H with coprojections $\kappa_n \colon H_n \to H$. Define $f_n \colon H_n \to \mathbb{C}$ by $f_n(z) = n \cdot ||\kappa_n|| \cdot z$. These are bounded maps, since $||f_n|| = n \cdot ||\kappa_n||$. Then for all $n \in \mathbb{N}$ the norm of the cotuple $f \colon H \to \mathbb{C}$ of (f_n) must satisfy

$$n \cdot ||\kappa_n|| = ||f_n|| = ||f \circ \kappa_n|| \le ||f|| \cdot ||\kappa_n||,$$

so that $n \leq ||f||$. This contradicts the boundedness and hence continuity of f. Finally, part (iii) follows from (ii) and [24, IX.1.1]

2.7. Despite the previous lemma, **Hilb** is *conditionally* (co) *complete*, in the sense that it does have objects that partially obey the universal property of infinite (co) products: for a family H_i of Hilbert spaces,

$$H = \{(x_i) \in \prod_i H_i \mid \sum_i ||x_i||^2 < \infty \}.$$

is a well-defined Hilbert space under the inner product $\langle (x_i) \mid (y_i) \rangle = \sum_i \langle x_i \mid y_i \rangle$ [20]. The evident morphisms $\pi_i \colon H \to H_i$ satisfy $\pi_i \circ \pi_i^{\dagger} = \operatorname{id}$ and $\pi_i \circ \pi_j^{\dagger} = 0$ when $i \neq j$. A cone $f_i \colon K \to H_i$ allows a unique well-defined morphism $f \colon K \to H$ satisfying $\pi_i \circ f = f_i$ if and only if $\sum_i \|f_i\|^2 < \infty$. Note, however, that the cone (π_i) itself does not satisfy this condition. In this sense, $\ell^2(X)$ is the conditional coproduct of X many copies of $\mathbb C$.

- **2.8.** A similar phenomenon occurs for simpler types of (co)limits. Monomorphisms in **Hilb** are precisely the injective morphisms, and epimorphisms are precisely those morphisms with dense range [15, A.3]. Not every monic epimorphism is an isomorphism. For example, the morphism $f: \ell^2(\mathbb{N}) \to \ell^2(\mathbb{N})$ defined by $f(\varphi)(n) = \frac{1}{n}\varphi(n)$ is injective, self-adjoint, and hence also has dense image. But it is not surjective, as the vector $\varphi \in \ell^2(\mathbb{N})$ determined by $\varphi(n) = \frac{1}{n}$ is not in its range. Monic epimorphisms are called *bimorphisms*.
- **2.9.** If $f,g: H \to K$ are morphisms in **Hilb**, then so are f+g and zf for $z \in \mathbb{C}$. Because composition respects these operations, **Hilb** is enriched over complex vector spaces. In general, the homsets are not Hilbert spaces themselves [2], so **Hilb** is not enriched over itself, and hence not Cartesian closed. At any rate, there is another way to structure the homsets of **Hilb**, which is of more interest here. Say $f \leq g$ when $\ker(f)^{\perp} \subseteq \ker(g)^{\perp}$ and f(x) = g(x) for $x \in \ker(f)^{\perp}$. The following proposition shows that this makes all homsets into conditional algebraic domains, but that this is not respected by composition. Here, by conditional algebraic domain we mean an algebraic domain [4], except that it does not need to have all directed suprema, but only bounded ones, in the same sense as 2.7: more precisely, every is a directed family of parallel morphisms f_i in **Hilb** for which $||f_i||$ converges has a supremum. This is closely related to [8, 2.1.4], but **Hilb** is not a restriction category in the sense of that paper: setting \overline{f} to be the projection onto $\ker(f)^{\perp}$ does not satisfy $\overline{fg} = \overline{gf}$.

Proposition 2.10. All homsets in the category **Hilb** are conditional algebraic domains, but composition is not monotone.

Proof. The least upper bound of a directed family f_i is given by continuous extension to the closure of $\bigcup_i \ker(f_i)^{\perp}$; this makes all homsets into directed-complete partially ordered sets. If $f \leq \bigvee_i f_i$ always implies $f \leq f_i$ for some i, then $\ker(f)^{\perp}$ must have been finite-dimensional; thus morphisms f with $\dim(\ker(f)^{\perp}) < \infty$ are the compact elements. It is now easy to see that any morphism is the directed supremum of compact ones below it, making all homsets into algebraic domains.

Now consider composition. First suppose that $f \leq f'$ and $g \leq g'$. If $x \in \ker(f)$, then clearly gf(x) = 0. If $x \in \ker(f)^{\perp}$, then f(x) = f'(x), so g'f'(x) = 0 implies $f(x) \in \ker(g') \subseteq \ker(g)$. Because we may write $\operatorname{dom}(gf) = \ker(f) \oplus \ker(f)^{\perp}$, we

conclude $\ker(gf)^{\perp} \subseteq \ker(g'f')^{\perp}$. But unless $f(\ker(gf)^{\perp}) \subseteq \ker(g)^{\perp}$, it need not be the case that gf equals g'f' on $\ker(gf)^{\perp}$. For an explicit counterexample, let

$$f = f' = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, \qquad g = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \qquad g' = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}.$$

Then $f \leq f'$ and $g \leq g'$. But $\ker(gf)^{\perp} = \{ \begin{pmatrix} x \\ -x \end{pmatrix} \mid x \in \mathbb{C} \}^{\perp} = \{ \begin{pmatrix} x \\ x \end{pmatrix} \mid x \in \mathbb{C} \}$, and $gf\begin{pmatrix} x \\ x \end{pmatrix} = \begin{pmatrix} 2x \\ 0 \end{pmatrix} \neq \begin{pmatrix} 2x \\ x \end{pmatrix} = g'f'\begin{pmatrix} x \\ x \end{pmatrix}$, so $gf \not\leq g'f'$.

3. The domain

Definition 3.1. A partial injection is a partial function that is injective, wherever it is defined. More precisely, it(s graph) is a relation $R \subseteq X \times Y$ such that for each x there is at most one y with $(x, y) \in R$, and for each y there is at most one x with $(y, x) \in R$. Sets and partial injections form a category **PInj** under composition of relations $S \circ R = \{(x, z) \mid \exists y \colon (x, y) \in R, (y, z) \in S\}$.

3.2. Notationally, a partial injection $f: X \to Y$ can be conveniently represented as a span $(X \leftarrow f_1 \frown F \succ f_2 \rightarrow Y)$ of monics in **Set**. Here, f_1 is (the inclusion of) the domain of definition of f, and f_2 is its (injective) action on that domain. Composition in this representation is by pullback. We will also write $Dom(f) = f_1(F)$ for the domain of definition, and $Im(f) = f_2(F)$ for the range of f.

If it wasn't already, the span notation immediately makes it clear that **PInj** is a dagger category: $(X \leftarrow f_1 \rightarrow F_2 \rightarrow Y)^{\dagger} = (Y \leftarrow f_2 \rightarrow F_1 \rightarrow X)$.

3.3. The category **PInj** has two dagger *symmetric monoidal* structures. The first one, that we denote by \otimes , acts as the Cartesian product on objects. Because the Cartesian product of injections is again injective, \otimes is well-defined on morphisms of **PInj** as well. The monoidal unit is a singleton set 1. Notice that \otimes is not a product, and hence not a coproduct either.

The second dagger symmetric monoidal structure on **PInj**, denoted by \oplus , is given by disjoint union on objects. It is easy to see that a disjoint union of injections is again injective, making \oplus well-defined on morphisms of **PInj**. The monoidal unit is the empty set. Notice that \oplus is not a coproduct, and hence not a product either.

Lemma 3.4. The category PInj:

- (i) has (co)equalizers;
- (ii) has a zero object;
- (iii) does not have finite (co)products;

Proof. The equalizer of $f, g: X \to Y$ is the inclusion of

$$\{x \in X \mid x \notin (\mathrm{Dom}(f) \cup \mathrm{Dom}(g)) \lor (x \in (\mathrm{Dom}(f) \cap \mathrm{Dom}(g)) \land f(x) = g(x))\}$$
 into X. The empty set is a zero object in **PInj**.

Towards (iii), notice that if $(X^{-\kappa_X} \to X + Y \leftarrow \kappa_Y - Y)$ were a coproduct in $\operatorname{\mathbf{PInj}}$, then one must have $\operatorname{Dom}(\kappa_X) = X$, $\operatorname{Dom}(\kappa_Y) = Y$ and $\operatorname{Im}(\kappa_X) \cap \operatorname{Im}(\kappa_Y) = \emptyset$, because otherwise unique existence of mediating morphisms is violated. Hence any coproduct must contain the disjoint union of X and Y. Let $f: X \to Z$ and $g: Y \to Z$ be any morphisms. Then a mediating morphism $m: X + Y \to Z$ has to satisfy m(x) = f(x) for $x \in \operatorname{Dom}(f)$ and m(y) = g(y) for $y \in \operatorname{Dom}(g)$. But such an m is not unique, unless $\operatorname{Dom}(f) = X$ and $\operatorname{Dom}(g) = Y$. In fact, it is not even a partial injection unless $\operatorname{Im}(f) \cap \operatorname{Im}(g) = \emptyset$. We conclude that $\operatorname{\mathbf{PInj}}$ does not have binary (co)products.

3.5. In fact, part (ii) of the previous lemma follows from the existence of directed colimits, which we now work towards. Recall that a category has directed colimits if and only if it has colimits of chains, *i.e.* colimits of well-ordered diagrams [5, Corollary 1.7]. Observe that for a chain $D: I \to \mathbf{PInj}$, if $c_i: D(i) \to X$ is a cocone on D, then $\mathrm{Dom}(c_i) \subseteq \mathrm{Dom}(D(i \le j))$ for all $j \ge i$. To see this, notice that $c_i = c_j \circ D(i \le j)$ since c_i is a cocone, and therefore

$$Dom(c_i) = Dom(c_j \circ D(i \le j)) \subseteq Dom(D(i \le j)).$$

This observation suggests that the colimit of a well-ordered diagram in **PInj** should consist of all 'infinite paths'. The following proposition shows that this is indeed a colimit.

Proposition 3.6. The category PInj has directed colimits.

Proof. Let $D: I \to \mathbf{PInj}$ be a chain. Define

$$X = \{x \in \coprod_{i} D(i) \mid \forall_{j \ge i} [x \in \text{Dom}(D(i \le j))]\} / \sim,$$

where the coproduct is taken in **Set**, and the equivalence relation \sim is generated by $x \sim D(i \leq j)(x)$ for all $i \leq j$ in I and $x \in \text{Dom}(D(i \leq j))$. For $i \in I$, define $c_i \colon D(i) \to X$ by

$$Dom(c_i) = \{x \in D(i) \mid \forall_{j>i} [x \in Dom(D(i \le j))]\},\$$

and $c_i(x) = [x]$.

First of all, let us show that the c_i form a cocone. One has:

$$Dom(c_i \circ D(i \leq j))$$

$$= \{x \in D(i) \mid x \in \text{Dom}(D(i \le j)) \land D(i \le j)(x) \in \text{Dom}(c_j)\}\$$

$$= \{x \in D(i) \mid x \in \text{Dom}(D(i \le j)) \land \forall_{k > j} [D(i \le j)(x) \in \text{Dom}(D(j \le k))] \}.$$

The well-orderedness of I implies that

$$\forall_{k>i}[P(k)] \Leftrightarrow \forall_{k>j}[P(k)] \wedge P(j)$$

for any property P on the objects of I, whence

$$Dom(c_i \circ D(i \le j)) = \{x \in D(i) \mid \forall_{k \ge i} [x \in Dom(D(i \le k))]\} = Dom(c_i).$$

Moreover $c_j \circ D(i \leq j)(x) = [D(i \leq j)(x)] = [x] = c_i(x)$ for $x \in \text{Dom}(c_i)$, by definition of the equivalence relation.

Next, we show that c_i is universal. Let $d_i : D(i) \to Y$ be any cocone, and define $m : X \to Y$ by

$$Dom(m) = \{ [x] \mid x \in Dom(d_i) \}$$


and $m([x]) = d_i(x)$ for $x \in D(i)$; this is well-defined since d_i is a cocone. Then

$$dom(m \circ c_i) = \{x \in D(i) \mid x \in Dom(c_i) \land m_i(x) \in Dom(m)\}$$
$$= \{x \in D(i) \mid \forall_{j \ge i} [x \in Dom(D(i \le j))] \land x \in Dom(d_i)\}$$
$$= Dom(d_i)$$

by 3.5, and $m \circ c_i(x) = m([x]) = d_i(x)$ for $x \in D(i)$. Thus $m \circ c_i = d_i$, so m is indeed a mediating morphism.

Finally, if $m': X \to Y$ satisfies $m \circ c_i = d_i$, then it follows from the above considerations that Dom(m') = Dom(m) and m'(x) = m(x) for $x \in Dom(m)$. Hence m is the unique mediating morphism.

3.7. Recall that an object X in a category \mathbf{C} is called *finitely presentable* when the hom-functor $\mathbf{C}(X,-)\colon \mathbf{C}\to \mathbf{Set}$ preserves directed colimits. Explicitly, this means that for any directed poset $D\colon I\to \mathbf{C}$, any colimit cocone $d_i\colon D(i)\to Y$ and any morphism $f\colon X\to Y$, there are $j\in I$ and a morphism $g\colon X\to D(j)$ such that $f=d_j\circ g$. Moreover, this morphism g is essentially unique, in the sense that if $f=d_i\circ g=d_i\circ g'$, then $D(i\to i')\circ g=D(i\to i')\circ g'$ for some $i'\in I$.



A category is called *finitely accessible* [5] when it has directed colimits and every object is a directed colimit of finitely presentable objects.

Lemma 3.8. A set is finitely presentable in **PInj** if and only if it is finite.

Proof. The only thing, in the situation of 3.7 with X finite, is to notice that if a partial injection g is to exist, we must have Dom(g) = Dom(f). The rest follows from [5, 1.2.1].

Theorem 3.9. The category PInj is finitely accessible.

Proof. It suffices to prove that every set in **PInj** is a directed colimit of finite ones. But that is easy: X is the colimit of the directed diagram consisting of its finite subsets.

Definition 3.10. An inverse category is a category \mathbf{C} in which every morphism $f \colon X \to Y$ allows a unique morphism $f^{\dagger} \colon Y \to X$ satisfying $f = ff^{\dagger}f$ and $f^{\dagger} = f^{\dagger}ff^{\dagger}$. Equivalently, it is a dagger category satisfying $f = ff^{\dagger}f$ and pq = qp for idempotents $p, q \colon X \to X$. The proof of equivalence of these two statements is the same as for inverse semigroups (see [23, Theorem 1.1.3] or [8, Theorem 2.20]). Inverse categories are a special case of restriction categories [8].

The category **PInj** is an inverse category under its dagger (see 3.2). The following categorification of the Wagner–Preston theorem [23, Theorem 1.5.1] shows that it is in fact a representative one. See also [8, 3.4].

Proposition 3.11. [21] Any locally small inverse category C allows a faithful embedding $F: C \to PInj$ that preserves daggers.

3.12. Like any inverse category, the homsets of **PInj** carry a natural partial order: $f \leq g$ when $f = gf^{\dagger}f$. Concretely, $f \leq g$ means $Dom(f) \subseteq Dom(g)$ and f(x) = g(x) for $x \in Dom(f)$. It is easy to see that this makes homsets into directed-complete partially ordered sets, with $Dom(\bigvee_i f_i) = \bigcup_i Dom(f_i)$ for a directed family of morphisms $f_i \colon X \to Y$. In fact, as in Proposition 2.10, homsets are algebraic domains: any partial injection is the supremum of compact ones below it, which are those partial injections with finite domain. Moreover, composition respects these operations. Thus **PInj** is enriched in algebraic domains. This is a satisfying reflection of Theorem 3.9 on the level of homsets.

4. The functor

Definition 4.1. There is a functor $\ell^2 \colon \mathbf{PInj} \to \mathbf{Hilb}$, acting on a set X as

$$\ell^2(X) = \{\varphi \colon X \to \mathbb{C} \mid \sum_{x \in X} |\varphi(x)|^2 < \infty\},\,$$

which is a well-defined Hilbert space under the inner product $\langle \varphi \mid \psi \rangle = \sum_{x \in X} \overline{\varphi(x)} \psi(x)$. The action on morphisms sends a partial injection $(X \leftarrow f_1 \leftarrow F_2 \rightarrow Y)$ to the linear function $\ell^2 f \colon \ell^2(X) \to \ell^2(Y)$ determined informally by $\ell^2 f = (_) \circ f^{\dagger}$. Explicitly,

$$(\ell^2 f)(\varphi)(y) = \sum_{x \in f_2^{-1}(y)} \varphi(f_1(x)).$$

4.2. In verifying that $\ell^2 f$ is indeed a well-defined morphism of **Hilb**, it is essential that f is a (partial) injection.

$$\sum_{y \in Y} |(\ell^2 f)(\varphi)(y)|^2 = \sum_{y \in Y} |\sum_{x \in f_2^{-1}(y)} \varphi(f_1(x))|^2 \le \sum_{y \in Y} \sum_{x \in f_2^{-1}(y)} |\varphi(f_1(x))|^2$$
$$= \sum_{x \in F} |\varphi(f_1(x))|^2 \le \sum_{x \in X} |\varphi(x)|^2 < \infty.$$

That this breaks down for functions f in general, instead of (partial) injections, was first noticed in [6], and further studied in [13]. That is, ℓ^2 is well-defined on the category of sets and partial injections; on the category of finite sets and functions; but not on the category of sets and functions; nor on the category of finite sets and relations. Functoriality of ℓ^2 is easy to verify.

4.3. The following calculation shows that the ℓ^2 functor preserves daggers. For a partial injection $(X \leftarrow f_1 \rightarrow F_2 \rightarrow Y)$, $\varphi \in \ell^2(X)$ and $\psi \in \ell^2(Y)$:

$$\begin{split} \langle (\ell^2 f)(\varphi) \, | \, \psi \rangle_{\ell^2(Y)} &= \sum_{y \in Y} \overline{(\ell^2 f)(\varphi)(y)} \cdot \psi(y) = \sum_{y \in Y} \sum_{x \in f_2^{-1}(y)} \overline{\varphi(f_1(x))} \cdot \psi(y) \\ &= \sum_{x \in F} \overline{\varphi(f_1(x))} \cdot \psi(f_2(x)) = \sum_{x \in X} \sum_{x' \in f_1^{-1}(x)} \overline{\varphi(x)} \cdot \psi(f_2(x')) \\ &= \sum_{x \in X} \overline{\varphi(x)} \cdot (\sum_{x' \in f_1^{-1}(x)} \psi(f_2(x'))) = \langle \varphi \, | \, \ell^2(f^\dagger)(\psi) \rangle_{\ell^2(X)}. \end{split}$$

- **4.4.** The functor ℓ^2 preserves the tensor product \otimes , *i.e.* it is symmetric (strong) monoidal. There is a canonical isomorphism $\mathbb{C} \cong \ell^2(\mathbf{1})$. The required natural morphisms $\ell^2(X) \otimes \ell^2(Y) \to \ell^2(X \otimes Y)$ are given by mapping (φ, ψ) to the function $(x,y) \mapsto \varphi(x)\psi(y)$. That there are inverses is seen when one realizes that $\ell^2(X \otimes Y)$ is the Cauchy-completion of the set of functions $X \times Y \to \mathbb{C}$ with finite support. The required coherence diagrams follow easily.
- **4.5.** Also, the ℓ^2 functor is symmetric (strong) monoidal with respect to \oplus . There is a canonical isomorphism between the 0-dimensional Hilbert space and the set $\ell^2(\emptyset)$ consisting only of the empty function. The natural morphisms $\ell^2(X) \oplus \ell^2(Y) \to \ell^2(X \oplus Y)$ map (φ, ψ) to the cotuple $[\varphi, \psi] \colon X \oplus Y \to \mathbb{C}$. One sees that these are isomorphisms by recalling that $\ell^2(X \oplus Y)$ is the closure of the span of the Kronecker functions δ_x and δ_y for $x \in X$ and $y \in Y$, on which the inverse acts as the appropriate coprojection. Coherence properties readily follow.

- **4.6.** From the description of the structure of homsets in **PInj** and **Hilb** as algebraic domains in 3.12 and 2.9, respectively, it is clear that the functor ℓ^2 preserves this enrichment: $\ell^2(\bigvee_i f_i) = \bigvee_i \ell^2 f_i$ if $f_i \colon X \to Y$ is a directed family of morphisms in **PInj**. See also [18, Theorem 13].
- **4.7.** The functor ℓ^2 preserves (co)kernels and finite (co)products (because **PInj** has very few of the latter). But it follows from Lemma 2.6(iii) and Proposition 3.6 that ℓ^2 cannot preserve arbitrary (co)limits. For an explicit counterexample to preservation of equalizers, take $X = \{0, 1\}$, $Y = \{a\}$, and let $f, g: X \to Y$ be the partial injections $f = \{(0, a)\}$ and $g = \{(1, a)\}$. Their equaliser in **PInj** is \emptyset . But

$$\begin{split} \operatorname{eq}(\ell^{2}(f),\ell^{2}(g)) &= \{ \varphi \in \ell^{2}(X) \mid \ell^{2}(f)(\varphi) = \ell^{2}(g)(\varphi) \} \\ &= \left\{ \varphi \in \ell^{2}(X) \mid \forall_{y \in Y}. \sum_{u \in f_{2}^{-1}(y)} \varphi(f_{1}(u)) = \sum_{v \in g_{2}^{-1}(y)} \varphi(g_{1}(v)) \right\} \\ &= \{ \varphi \colon \{0,1\} \to \mathbb{C} \mid \varphi(0) = \varphi(1) \} \cong \mathbb{C}. \end{split}$$

Hence $eq(\ell^2(f), \ell^2(g)) \cong \mathbb{C} \ncong \{\emptyset\} = \ell^2(eq(f,g)).$

Corollary 4.8. The functor ℓ^2 : PInj \rightarrow Hilb has no adjoints.

Proof. If ℓ^2 had an adjoint, it would preserve (co)limits, contradicting 4.7.

- **4.9.** The functor ℓ^2 is clearly faithful. It is also essentially surjective on objects: every Hilbert space H has an orthonormal basis X, so $H \cong \ell^2(X)$. It cannot be full because of 4.8, but it does reflect isomorphisms: if $\ell^2 f$ is invertible, so is f.
- **4.10.** If X is a set, $\ell^2(X)$ is not just a Hilbert space; it comes equipped with a chosen orthonormal basis (given by the Kronecker functions $\delta_x \in \ell^2(X)$ for $x \in X$). Hence we could think of ℓ^2 as a functor to a category of Hilbert spaces H with a priviledged orthonormal basis $X \subseteq H$. If we choose as morphisms $(H, X) \to (K, Y)$ those continuous linear $f: H \to K$ satisfying $f(X) \subseteq Y$ and $ff^{\dagger}f = f$, then the functor ℓ^2 in fact becomes (half of) an equivalence of categories [3, 4.3].
- **4.11.** Lemma 4.8 showed that $\ell^2(X)$ is not the free Hilbert space on X, at least not in the categorically accepted meaning. It also makes precise the intuition that 'choosing bases is unnatural': the functor $\ell^2 \colon \mathbf{PInj} \to \mathbf{Hilb}$ cannot have a (functorial) converse, even though one can choose an orthonormal basis for every Hilbert space.

It is perhaps also worth mentioning that ℓ^2 is not a fibration in the technical sense of the word, not even a nonsplit or noncloven one, as the reader might perhaps think; Cartesian liftings in general do not exist because 'choosing bases is unnatural'.

5. The image

5.1. The choice of morphisms in 4.10 is quite strong, and does not capture all morphisms of interest to quantum physics. From that point of view, one would at least like to relax to partial isometries: morphisms i of Hilbert spaces that satisfy $ii^{\dagger}i = i$. Equivalently, the restriction of i to the orthogonal complement of its kernel is an isometry. The following proposition proves that, up to isomorphisms, the direct image of the functor ℓ^2 consists precisely of partial isometries.

Definition 5.2. For a category C, denote by C_{\cong} the category with the same objects as C whose morphisms are the bimorphisms of C.

The category \mathbf{Hilb}_{\cong} carries two dagger symmetric monoidal structures: \oplus and \otimes . Because having (co)limits only depends on a skeleton of the specifying diagram, \mathbf{Hilb}_{\cong} does not have (co)equalizers, nor (finite) (co)products, but does have directed (co)limits.

Proposition 5.3. A morphism in **Hilb** is a partial isometry if and only if it is of the form $v \circ \ell^2 f \circ u$ for morphisms f in **PInj** and unitaries u, v in **Hilb** \cong .

Proof. Clearly a map of the form $v \circ \ell^2 f \circ u$ is a partial isometry. Conversely, suppose that $i \colon H \to K$ is a partial isometry. Choose an orthonormal basis $X \subseteq H$ for its initial space $\ker(i)^{\perp}$, and choose an orthonormal basis $X' \subseteq H$ for $\ker(i)$, giving a unitary $u \colon H \to \ell^2(X \oplus X')$. Let $Y = i(X) \subseteq K$. Then Y will be an orthonormal basis for the final space $\ker(i^{\dagger})^{\perp}$ because i acts isometrically on X. Choose an orthonormal basis $Y' \subseteq K$ for $\ker(i^{\dagger})$, giving a unitary $v \colon \ell^2(Y \oplus Y') \to K$. Now, if we define $f = (X \oplus X' \longleftarrow X) \xrightarrow{i \to Y} Y \oplus Y'$, then $i = v \circ \ell^2 f \circ u$.

- **5.4.** However, partial isometries are not closed under composition. To see this, consider the partial isometries $\binom{1}{0}: \mathbb{C} \to \mathbb{C}^2$ and $\left(\sin(\theta)\cos(\theta)\right): \mathbb{C}^2 \to \mathbb{C}$ for a fixed real number θ . Their composition is $\left(\sin(\theta)\right): \mathbb{C} \to \mathbb{C}$, which is not a partial isometry unless θ is a multiple of $\pi/2$. There are other compositions that do make partial isometries into a category [19], but these are not of interest here. Instead, we shall extend the previous proposition to highlight one of the most remarkable features of the functor ℓ^2 .
- **5.5.** The example in 5.4 shows that any linear function $\mathbb{C} \to \mathbb{C}$ between -1 and 1 is a composition of partial isometries. Note that the projections $\pi_i \colon \mathbb{C}^m \to \mathbb{C}$ and coprojections $\pi_i^{\dagger} \colon \mathbb{C} \to \mathbb{C}^n$ are partial isometries, as are the weighted diagonal $\Delta/\sqrt{n} \colon \mathbb{C} \to \mathbb{C}^n$ given by $\Delta(x) = (x, \dots, x)$ and the weighted codiagonal $\Delta^{\dagger}/\sqrt{m} \colon \mathbb{C}^m \to \mathbb{C}$ given by $\Delta^{\dagger}(x_1, \dots, x_m) = \sum_i x_i$. Moreover, it is easy to see that if f and g are (compositions of) partial isometries, then so is $f \oplus g$. Finally, any linear map $f \colon \mathbb{C}^m \to \mathbb{C}^n$ has a matrix expansion, and can hence be written in terms of biproduct structure as $f = \Delta^{\dagger} \circ (\bigoplus_{i=1}^m \bigoplus_{j=1}^n \pi_j^{\dagger} \circ \pi_j \circ f \circ \pi_i^{\dagger} \circ \pi_i) \circ \Delta$. Thus any $f \colon \mathbb{C}^m \to \mathbb{C}^n$ with $\|f\| \le 1/\sqrt{mn}$ is a composition of partial isometries.
- **5.6.** The essential image of a functor $F : \mathbf{C} \to \mathbf{D}$ is the smallest subcategory of \mathbf{D} that contains all morphisms F(f) for f in \mathbf{C} , and that is closed under composition with bimorphisms of \mathbf{D} .

It follows from 5.5 that the essential image of the functor ℓ^2 contains at least all morphisms of **Hilb** of finite rank. For infinite rank that strategy fails because Δ is then no longer a valid morphism (see 2.7). Nevertheless, Theorem 5.11 below will prove that the essential image of ℓ^2 is all of **Hilb**. In preparation we accommodate an intermezzo on *polar decomposition*.

A morphism $p: H \to H$ in **Hilb** is nonnegative when $\langle px | x \rangle \geq 0$ for all $x \in H$, and positive when $\langle px | x \rangle > 0$. Nonnegative maps are precisely those of the form $p = f^{\dagger}f$ for some morphism f.

Proposition 5.7. For every morphism $f: H \to K$ between Hilbert spaces, there exist a unique nonnegative map $p: H \to H$ and partial isometry $i: H \to K$ satisfying f = ip and $\ker(p) = \ker(i)$.

Proof. See [14, problem 134].

5.8. The previous proposition stated the usual formulation of polar decomposition, but the unicity condition $\ker(p) = \ker(i)$ is something of a red herring. It should be understood as saying that both i and p are uniquely determined on the orthogonal complement of $\ker(f) = \ker(p) = \ker(i)$. On each point of $\ker(f)$, one of i and p must be zero, but the other's behaviour has no restrictions apart from being a partial isometry or positive map, respectively. Dropping the unicity condition, we may take p to be a positive map, by altering i to be zero on $\ker(f)$, and p to be nonzero on $\ker(f)$. More precisely, define p' = p on $\ker(f)^{\perp}$ and $p' = \mathrm{id}$ on $\ker(f)$; since $\ker(f)$ is a closed subspace, $H \cong \ker(f) \oplus \ker(f)^{\perp}$, and this gives a well-defined positive operator $p' \colon H \to H$. Similarly, setting i' = i on $\ker(f)^{\perp}$ and i' = 0 on $\ker(f)$ gives a well-defined partial isometry $i' \colon H \to K$, satisfying f = i'p'.

In the following we have to consider bimorphisms rather than isomorphism because of the issues in 2.8. In the finite-dimensional case, bimorphisms are isomorphisms.

Lemma 5.9. Positive operators on Hilbert spaces are bimorphisms.

Proof. Let $p: H \to H$ be a positive operator in **Hilb**. If p(x) = 0 then certainly $\langle p(x) | x \rangle = 0$ which contradicts positivity. Hence $\ker(p) = 0$, and so p is monic.

To see that p is epic, suppose that $p \circ f = p \circ g$ for parallel morphisms f, g. Then $\langle p \circ (f-g)(x) | x \rangle = 0$ for all x. By positivity, For each x there is $p_x > 0$ such that $p \circ (f-g)x = p_x \cdot (f-g)(x)$. Hence $\langle (f-g)(x) | x \rangle = 0$ for all x, that is, f=g and p is epic.

Definition 5.10. A functor $F: \mathbf{C} \to \mathbf{D}$ is essentially full when for each morphism g in \mathbf{D} there exist f in \mathbf{C} and u, v in \mathbf{C}_{\cong} such that $g = v \circ Ff \circ u$.

It follows that the essential image of such a functor is all of **D**.

Theorem 5.11. The functor $\ell^2 \colon \mathbf{PInj} \to \mathbf{Hilb}$ is essentially full.

Proof. Let g be a morphism in **Hilb**. By Proposition 5.7 and 5.8, we can write g = pi for a positive morphism p and a partial isometry i. Use Proposition 5.3 to decompose $i = v' \circ \ell^2 f \circ u$ for f in **PInj** and unitaries v', u. Finally, Lemma 5.9 shows that $v = p \circ v'$ in \mathbf{Hilb}_{\cong} satisfies $g = v \circ \ell^2 f \circ u$.

5.12. Writing **2** for the ordinal $2=(0 \le 1)$ regarded as a category, the category ${\bf C^2}$ is the arrow category of ${\bf C}$: its objects are morphisms of ${\bf C}$, and its morphisms are pairs of morphisms of ${\bf C}$ making the square commute. A functor $F: {\bf C} \to {\bf D}$ is essentially full when $F^2: {\bf C^2} \to {\bf D^2}$ is essentially surjective on objects. From this point of view Definition 5.10 is quite natural. Nonetheless we might consider weakening it to take $u={\bf id}$ or $v={\bf id}$. But this would break the previous theorem. For example, if $g: \ell^2(X) \to \ell^2(Y)$ is a morphism in **Hilb**, there need not be $f: X \to Y$ in **PInj** and v in **Hilb** \cong with $g=v\circ\ell^2 f$. For a counterexample, take $X=Y=\{a,b\}$, and g(a)=g(b)=a; if $g=v\circ\ell^2 f$, then $(v\circ\ell^2 f)(a)=(v\circ\ell^2 f)(b)$, so $(\ell^2 f)(a)=(\ell^2 f)(b)$, so f(a)=f(b), whence f cannot be a partial injection. Similarly, because of the dagger, if $g:\ell^2(X)\to\ell^2(Z)$ is a morphism in **Hilb**, there need not be $f:Y\to Z$ in **PInj** and u in ${\bf Hilb}_\cong$ with $g=\ell^2 f\circ u$.

6. The future

6.1. Theorem 5.11 naturally raises a coherence question: is there any regularity to the bimorphisms u and v that enable us to write an arbitrary morphism of

Hilb in the form $v \circ \ell^2 f \circ u$? How do they behave under composition? Curiously enough, essentially full functors do not seem to have been studied in the categorical literature at all. The results in this article suggest such a study.

It would be very interesting to reconstruct **Hilb** (up to equivalence) from $\mathbf{Hilb} \cong$ and \mathbf{PInj} via the ℓ^2 functor. The objects are easily recovered, because they are the same as those of \mathbf{Hilb}_{\cong} . Theorem 5.11 also lets us recover the homsets and identities, as soon as we can identify when two morphisms in \mathbf{Hilb} of the form $v \circ \ell^2 f \circ u$ are equal. The main problem is how to recover composition, which requires a way to turn $\ell^2 g \circ v \circ \ell^2 f$ into $w \circ \ell^2 h \circ u$. (Note that turning $\ell^2 g \circ v$ into $w \circ \ell^2 h$ would be sufficient, because we could then use functoriality of ℓ^2 and composition in \mathbf{PInj} . But 5.12 obstructs this; the bimorphism v in the middle is crucial.) This will likely lead into bicategorical territory.

6.2. The ℓ^2 -construction has a continuous counterpart, that turns a measure space (X,μ) into a Hilbert space $L^2(X,\mu)$ of square integrable complex functions on X. The L^2 -construction is quite fundamental and well-studied, but surprisingly enough functorial aspects seem not to have been considered before. One possibility is to mimic Definition 4.1, and endow the category of measure spaces with essential injections $(X,\mu) \to (Y,\nu)$ as morphisms, *i.e.* subsets $R \subseteq X \times Y$ such that $\nu(\{y \mid xRy\}) = 0$ for all $x \in X$ and $\mu(\{x \mid xRy\}) = 0$ for all $y \in Y$.

The importance of L^2 -spaces lies in the following formulation of the spectral theorem: every normal operator $f \colon H \to H$ is of the form $f = u^{-1} \circ g \circ u$ for a unitary $u \colon H \to L^2(X,\mu)$ and an operator g induced by multiplication with a measurable function $X \to \mathbb{C}$. This perspective warrants choosing complex measurable functions as (endo)morphisms on measure spaces, with multiplication for composition. With 5.8 in mind, we could even restrict to a groupoid of positive maps. A solution to 6.1 could then be regarded as reconstructing quantum mechanics (as embodied by **Hilb**) from its continuous, quantitative aspects (encoded by the L^2 functor), and its discrete, qualitative aspects (encoded by the ℓ^2 functor).

At any rate, the continuous cousin L^2 of ℓ^2 poses an interesting research topic.

- **6.3.** Letting \mathcal{L} be the class of positive morphisms, and \mathcal{R} the class of partial isometries in **Hilb**:
 - (1) every morphism f can be factored as f = rl with $l \in \mathcal{L}$ and $r \in \mathcal{R}$;
 - (2) every commutative square as below with $l \in \mathcal{L}$ and $r \in \mathcal{R}$ allows a unique diagonal fill-in d making both triangles commute.



The second property would follows from Lemma 5.9 for isomorphisms l. The established notion of *orthogonal factorization system* additionally demands that (3) both \mathcal{L} and \mathcal{R} are closed under composition, and (4) all isomorphisms are in both \mathcal{L} and \mathcal{R} . But (3) is not satisfied by 5.4, and the map $-1: H \to H$ is a counterexample to (4).

Write 3 for the ordinal $3 = (0 \le 1 \le 2)$, regarded as a category. Then objects of \mathbb{C}^3 are composable pairs of morphisms. Recall that a functorial factorization is a functor $F \colon \mathbb{C}^2 \to \mathbb{C}^3$ that splits the composition functor. Lemma 5.9 ensures that polar decomposition at least provides a functorial factorization system. It is usual

to require extra conditions on top of a functorial factorization, such as in a *natural* weak factorization system. For details we refer to [11]. It leads too far afield here, but polar decomposition does not satisfy the axioms of a natural weak factorization system.

In short, polar decomposition unquestionably provides a notion of factorization. But it does not fit existing categorical notions, despite the fact that factorization has been a topic of quite intense study in category theory [10, 7, 11, 22, 26]. This is an interesting topic for further investigation.

References

- [1] Samson Abramsky. Retracing some paths in process algebra. In 7th International Conference on Concurrency Theory, pages 1–17. Springer, 1996.
- [2] Samson Abramsky, Richard Blute, and Prakhash Panangaden. Nuclear and trace ideals in tensored *-categories. Journal of Pure and Applied Algebra, 143:3–47, 1999.
- [3] Samson Abramsky and Chris Heunen. H*-algebras and nonunital Frobenius algebras: first steps in infinite-dimensional categorical quantum mechanics. In Samson Abramsky and Michael Mislove, editors, *Clifford Lectures*, volume 71 of *Proceedings of Symposia in Applied Mathematics*, pages 1–24. American Mathematical Society, 2012.
- [4] Samson Abramsky and Achim Jung. Domain theory. In Handbook of Logic in Computer Science Volume 3, pages 1–168. Oxford University Press, 1994.
- [5] Jiří Adámek and Jiří Rosicky. Locally Presentable and Accessible Categories. Number 189 in London Mathematical Society Lecture Note Series. Cambridge University Press, 1994.
- [6] Michael Barr. Algebraically compact functors. Journal of Pure and Applied Algebra, 82:211–231, 1992.
- [7] Aldridge K. Bousfield. Constructions of factorization systems in categories. *Journal of Pure and Applied Algebra*, 9(2–3):207–220, 1977.
- [8] J. Robin B. Cockett and Stephen Lack. Restriction categories I: categories of partial maps. Theoretical Computer Science, 270(1-2):223-259, 2002.
- [9] Vincent Danos and Laurent Regnier. Proof-nets and the Hilbert space. In Advances in Linear Logic, pages 307–328. Cambridge University Press, 1995.
- [10] Peter Freyd and Max Kelly. Categories of continuous functors I. Journal of Pure and Applied Algebra, 2, 1972.
- [11] Marco Grandis and Walther Tholen. Natural weak factorization systems. Archivum Mathematicum, 42:397–408, 2006.
- [12] Esfandiar Haghverdi. A categorical approach to linear logic, geometry of proofs and full completeness. PhD thesis, University of Ottawa, 2000.
- [13] Esfandiar Haghverdi and Phil Scott. A categorical model for the geometry of interaction. Theoretical Computer Science, 350:252–274, 2006.
- [14] Paul Halmos. A Hilbert space problem book. Springer, 2nd edition, 1982.
- [15] Chris Heunen. An embedding theorem for Hilbert categories. Theory and Applications of Categories, 22(13):321–344, 2009.
- [16] Chris Heunen and Bart Jacobs. Quantum logic in dagger kernel categories. Order, 27(2):177–212, 2010.
- [17] Peter Hines. The algebra of self-similarity and its applications. PhD thesis, University of Wales, 1997.
- [18] Peter Hines. Quantum circuit oracles for abstract machine computations. Theoretical Computer Science, 411(11-13):1501-1520, 2010.
- [19] Peter Hines and Samuel L. Braunstein. Semantic techniques in quantum computation, chapter The structure of partial isometries, pages 361–389. Cambridge University Press, 2009.
- [20] Richard V. Kadison and John R. Ringrose. Fundamentals of the theory of operator algebras. Academic Press, 1983.
- [21] J. Kastl. Algebraische Modelle, Kategorien und Gruppoide, volume 7 of Studien zur Algebra und ihre Anwendungen, chapter Inverse categories, pages 51–60. Akademie-Verlag Berlin, 1979.
- [22] Mareli Korostenski and Walter Tholen. Factorization systems as Eilenberg-Moore algebras. Journal of Pure and Applied Algebra, 85(1):57–72, 1993.

- [23] Mark V. Lawson. Inverse semigroups: the theory of partial symmetries. World Scientific, 1998.
- $[24] \ \ Saunders \ Mac \ Lane. \ \textit{Categories for the Working Mathematician}. \ Springer, \ 2nd \ edition, \ 1971.$
- [25] Michael Reed and Barry Simon. Methods of Modern Mathematical Physics, Vol I: Functional Analysis. Academic Press, 1972.
- [26] Jiří Rosicky and Walter Tholen. Lax factorization algebras. Journal of Pure and Applied Algebra, 175:355–382, 2002.

Department of Computer Science, University of Oxford, Wolfson Building, Parks Road, OX1 3QD, Oxford, UK

 $E\text{-}mail\ address: \verb|heunen@cs.ox.ac.uk||$