QCSP on partially reflexive cycles - the wavy line of tractability

Florent Madelaine¹ and Barnaby Martin^{2*}

¹ Clermont Université, Université d'Auvergne, LIMOS, BP 10448, F-63000 Clermont-Ferrand, France.

² CNRS / LIX UMR 7161, École Polytechnique, Palaiseau, France.

Abstract. We study the (non-uniform) quantified constraint satisfaction problem $QCSP(\mathcal{H})$ as \mathcal{H} ranges over partially reflexive cycles. We obtain a complexity-theoretic dichotomy: $QCSP(\mathcal{H})$ is either in NL or is NP-hard. The separating conditions are somewhat esoteric hence the epithet "wavy line of tractability" (see Figure 5 at end).

1 Introduction

The quantified constraint satisfaction problem QCSP(\mathcal{B}), for a fixed template (structure) \mathcal{B} , is a popular generalisation of the constraint satisfaction problem CSP(\mathcal{B}). In the latter, one asks if a primitive positive sentence (the existential quantification of a conjunction of atoms) Φ is true on \mathcal{B} , while in the former this sentence may be positive Horn (where universal quantification is also permitted). Much of the theoretical research into CSPs is in respect of a large complexity classification project – it is conjectured that CSP(\mathcal{B}) is always either in P or NP-complete [11]. This dichotomy conjecture remains unsettled, although dichotomy is now known on substantial classes (e.g. structures of size ≤ 3 [20, 3] and smooth digraphs [12, 1]). Various methods, combinatorial (graph-theoretic), logical and universal-algebraic have been brought to bear on this classification project, with many remarkable consequences. A conjectured delineation for the dichotomy was given in the algebraic language in [4].

Complexity classifications for QCSPs appear to be harder than for CSPs. Indeed, a classification for QCSPs will give a fortiori a classification for CSPs (if $\mathcal{B} \uplus \mathcal{K}_1$ is the disjoint union of \mathcal{B} with an isolated element, then QCSP($\mathcal{B} \uplus \mathcal{K}_1$) and CSP(\mathcal{B}) are polynomially equivalent). Just as CSP(\mathcal{B}) is always in NP, so QCSP(\mathcal{B}) is always in Pspace. However, no overarching polychotomy has been conjectured for the complexities of QCSP(\mathcal{B}), as \mathcal{B} ranges over finite structures, but the only known complexities are P, NP-complete and Pspace-complete (see [2, 17] for some trichotomies). It seems plausible that these complexities are the only ones that can be so obtained (for more in this see [6]).

In this paper we study the complexity of $QCSP(\mathcal{H})$, where \mathcal{H} is a partially reflexive cycle. In this respect, our paper is a companion to the similar classification for partially reflexive forests in [16]. We derive a classification between those

^{*} The author is supported by ANR Blanc International ALCOCLAN.

cases that are in NL and those that are NP-hard. For some of the NP-hard cases we are able to demonstrate Pspace-completeness. The dichotomy, as depicted in Figure 5 at the end, is quite esoteric and deviates somewhat from similar classifications (e.g. for retraction as given in [10]). To our minds, this makes it interesting in its own right. Some of our hardness proofs come from judicious amendments to the techniques used in [16]. Several others use different elaborate encodings of retraction problems, known to be hard from [10]. All but one of our NL-membership results follow from a majority polymorphism in an equivalent template (indeed – the so-called *Q-core* of [14]), as they did in [10]. However, \mathcal{C}_{0111} is special. It has no QCSP-equivalent that admits a majority (indeed, it omits majority and is a Q-core), so we have to give a specialised algorithm, based on ideas from [13], to demonstrate that $QCSP(\mathcal{C}_{0111})$ is in L. Indeed, and in light of the observations in [14], this is the principal news from the partially reflexive cycles classification that removes it from being simply a sequel to partially reflexive forests: for a partially reflexive forest \mathcal{H} , either the Q-core of \mathcal{H} admits a majority polymorphism and $QCSP(\mathcal{H})$ is in NL, or $QCSP(\mathcal{C})$ is NP-hard. The same can not be said for partially reflexive cycles, due to the odd case of \mathcal{C}_{0111} .

This paper is organised as follows. After the preliminaries, we address small cycles in Section 3. Then we deal with reflexive cycles, cycles whose loops induce a path and cycles with disconnected loops in Sections 4, 5 and 6, respectively. Finally we give our classification in Section 7 and our conclusions in Section 8. For reasons of space, many proofs are deferred to the appendix.

2 Definitions and preliminaries

Let $[n] := \{1, \ldots, n\}$. A graph \mathcal{G} has vertex set G, of cardinality |G|, and edge set $E(\mathcal{G})$. For a sequence $\alpha \in \{0, 1\}^*$, of length $|\alpha|$, let \mathcal{P}_{α} be the undirected path on $|\alpha|$ vertices such that the *i*th vertex has a loop iff the *i*th entry of α is 1 (we may say that the path \mathcal{P} is of the form α). We will usually envisage the domain of a path with n vertices to be [n], where the vertices appear in the natural order. Similarly, for $\alpha \in \{0, 1\}^*$, let \mathcal{C}_{α} be the $|\alpha|$ cycle with domain [n] and edge set $\{(i, j) : |j - i| = 1 \mod n\} \cup \{(i, i) : \alpha[i] = 1\}$ (note $|n - 1| = |1 - n| = 1 \mod n$). If α and α' are sequences in $\{0, 1\}^n$ such that $\alpha[i] = \alpha'[i+1 \mod n]$ then \mathcal{C}_{α} and $\mathcal{C}_{\alpha'}$ are isomorphic.

A partially reflexive cycle is one that may include some self-loops. For such an *m*-cycle C, whose vertices we will imagine to be v_1, \ldots, v_m in their natural mod *m* adjacencies, let $[v_i \Rightarrow v_j]$ be shorthand for a conjunction specifying a path, whichever is the fastest way mod *m*, from v_i to v_j . For example, if m = 5, then 1.) $[v_1 \Rightarrow v_3]$ is $E(v_1, v_2) \wedge E(v_2, v_3), 2$.) $[v_3 \Rightarrow v_1]$ is $E(v_3, v_2) \wedge E(v_2, v_1)$, and 3.) $[v_4 \Rightarrow v_1]$ is $E(v_4, v_5) \wedge E(v_5, v_1)$. We ask the reader to endure the following relaxation of this notation; $[v_i, v_{i+1} \Rightarrow v_j]$ indicates an edge from v_i to v_{i+1} then a path to v_j (which may not be the same as $[v_i \Rightarrow v_j]$ as the latter may take the other path around the cycle). Finally, let $\text{Ref}(v_i, \ldots, v_j)$ indicate $E(v_i, v_i) \wedge \ldots \wedge E(v_j, v_j)$, whichever is the quickest way around the cycle mod *m*. All graphs in this paper are undirected, so edge statements of the form E(x, y) should be read as asserting $E(x, y) \wedge E(y, x)$.

The problems $\operatorname{CSP}(\mathcal{T})$ and $\operatorname{QCSP}(\mathcal{T})$ each take as input a sentence Φ , and ask whether this sentence is true on \mathcal{T} . For the former, the sentence involves the existential quantification of a conjunction of atoms – *primitive positive* logic. For the latter, the sentence involves the arbitrary quantification of a conjunction of atoms – *positive Horn* logic. By convention equalities are permitted in both of these, but these may be propagated out by substitution in all but trivial degenerate cases. The *retraction problem* $\operatorname{RET}(\mathcal{B})$ takes as input some \mathcal{G} , with \mathcal{H} an induced substructure of \mathcal{G} , and asks whether there is a homomorphism $h: \mathcal{G} \to \mathcal{H}$ such that h is the identity on \mathcal{H} . It is important that the copy of \mathcal{H} is specified in \mathcal{G} ; it can be that \mathcal{H} appears twice as an induced substructure and there is a retraction from one of these instances but not to the other. The problem $\operatorname{Ret}(\mathcal{H})$ is easily seen to be logspace equivalent with the problem $\operatorname{CSP}(\mathcal{H}^c)$, where \mathcal{H}^c is \mathcal{H} expanded with all constants (one identifies all elements assigned to the same constant and enforces the structure \mathcal{H} on those constants).

The direct product $\mathcal{G} \times \mathcal{H}$ of two graphs \mathcal{G} and \mathcal{H} has vertex set $\{(x, y) : x \in G, y \in H\}$ and edge set $\{((x, u), (y, v)) : x, y \in G, u, v \in H, (x, y) \in E(\mathcal{G}), (u, v) \in E(\mathcal{H})\}$. Direct products are (up to isomorphism) associative and commutative. The kth power \mathcal{G}^k of a graph \mathcal{G} is $\mathcal{G} \times \ldots \times \mathcal{G}$ (k times). A homomorphism from a graph \mathcal{G} to a graph \mathcal{H} is a function $h: G \to H$ such that, if $(x, y) \in E(\mathcal{G})$, then $(h(x), h(y)) \in E(\mathcal{G})$. A k-ary polymorphism of a graph is a homomorphism from \mathcal{G}^k to \mathcal{G} . A ternary function $f: G^3 \to G$ is designated a majority operation if f(x, x, y) = f(x, y, x) = f(y, x, x) = x, for all $x, y \in G$.

A positive Horn sentence Φ in the language of graphs induces naturally a graph \mathcal{G}_{Φ} whose vertices are the variables of Φ and whose edges are the atoms of Φ . In the case of primitive positive Φ one would call \mathcal{G}_{Φ} the canonical database and Φ its canonical query. With positive Horn Φ there is additional extra structure and one may talk of a vertex-variable as being existential/ universal and as coming before/ after (earlier/ later), in line with the quantifier **block** and its order in Φ . Variables in the same quantifier block will not need their orders considered (there is commutativity within a block anyway). A typical reduction from a retraction problem $\operatorname{Ret}(\mathcal{C})$, where $|\mathcal{C}| = m$, builds a positive Horn Φ sentence involving variables v_1, \ldots, v_m where we want \mathcal{G}_{Φ} restricted to $\{v_1, \ldots, v_m\}$ (itself a copy of \mathcal{C}) to map automorphically to \mathcal{C} . Typically, we can force this with some evaluation of the variables (some of which might be universally quantified). The other valuations are degenerate and we must ensure at least that they map \mathcal{G}_{Φ} restricted to $\{v_1, \ldots, v_m\}$ homomorphically to \mathcal{C} .

3 Small cycles

The classification for QCSP for cycles of length ≤ 4 is slightly esoteric, although it does match the analogous classification for Retraction (the former is a dichotomy between NL and Pspace-complete; the latter is a dichotomy between P and NP-complete). The following has appeared in [15], where it was given as an application of counting quantifiers in QCSPs. We review it now because we will need to generalise it later in Section 4.

Proposition 1 ([15]). $QCSP(C_{1111})$ is Pspace-complete.

Proof. We will reduce from the problem $QCSP(\mathcal{K}_4)$ (known to be Pspace-complete from, e.g., [2]). We will borrow heavily from the reduction of $CSP(\mathcal{K}_4)$ to $Ret(\mathcal{C}_{1111})$ in [9]. We introduce the following shorthands (x', x'') must appear nowhere else in φ , which may contain more free variables than just x).

$$\begin{split} \exists^{\geq 1} x \ \varphi(x) &:= \exists x \ \varphi(x) \\ \exists^{\geq 2} x \ \varphi(x) &:= \forall x' \exists x \ E(x', x) \land \varphi(x) \\ \exists^{\geq 3} x \ \varphi(x) &:= \forall x'' \forall x' \exists x \ E(x'', x) \land E(x', x) \land \varphi(x) \end{split}$$

On \mathcal{C}_{1111} , it is easy to verify that, for each $i \in [4], \exists^{\geq i} x \varphi(x)$ holds iff there exist at least *i* elements *x* satisfying φ . Thus our borrowing the notation of counting quantifiers is justified.

We now reduce an instance Φ of QCSP(\mathcal{K}_4) to an instance Ψ of QCSP(\mathcal{C}_{1111}).

We begin with a cycle C_{1111} on vertices 1, 2, 3 and 4, which we realise through their canonical query (without quantification) as $\theta(v_1, v_2, v_3, v_4) := E(v_1, v_2) \wedge E(v_2, v_3) \wedge E(v_3, v_4) \wedge E(v_4, v_1)$ (recall that the canonical query is in fact the reflexive closure of this, but this will not be important in this case or many future cases – when it is important it will be stated explicitly). If Φ contains an atom E(x, y), then this gives rise to a series of atoms in Ψ as dictated by the gadget in Figure 1 (for each atom we add many new vertex-variables, corresponding to the vertices in the gadget that are not x, y, 1, 2, 3, 4). These atoms can be seen



Fig. 1. Edge gadget in reduction from $QCSP(\mathcal{K}_4)$ to $QCSP(\mathcal{C}_{1111})$

to join up with the atoms of θ as in the right end of the figure. Build Ψ''' from Φ by this process and conjunction with θ . Then make Ψ'' from Ψ''' by existentially quantifying all of the variables other than those associated to atoms of Φ (x, y in the figure) and v_1, v_2, v_3, v_4 ($1, \ldots, 4$ in the figure). Now, we build Ψ' from Ψ'' , by copying the quantifier order of Φ on the outside of the existential quantifiers that we already have. Thus $\Psi'(v_1, v_2, v_3, v_4)$ is a positive Horn formula with precisely four free variables.

It is not hard to see that when v_1, v_2, v_3, v_4 are evaluated as (an automorphism of) 1, 2, 3, 4, then we have a faithful simulation of $QCSP(\mathcal{K}_4)$). This is because

x and y, as in the gadget drawn, may evaluate precisely to distinct vertices on \mathcal{C}_{1111} . Finally, we build $\Psi := \exists v_1 \exists^{\geq 2} v_2 \exists^{\geq 3} v_3 \exists^{\geq 2} v_4 \Psi'(v_1, v_2, v_3, v_4)$. It is not hard to see that Ψ forces on some evaluation of v_1, v_2, v_3, v_4 that these map isomorphically to 1, 2, 3, 4 in \mathcal{C}_{1111} . Further, a rudimentary case analysis shows us that when they do not, we can still evaluate the remainder of Ψ' , if we could have done when they did. In fact, if v_1, v_2, v_3, v_4 are not mapped isomorphically (but still homomorphically, of course) to 1, 2, 3, 4 then we can extend each of the edge gadgets to homomorphism under **all maps** of vertices x and y to 1, 2, 3, 4 (not just ones in which x and y are evaluated differently).

Proposition 2. $QCSP(C_{0111})$ is in L.

Proof. Recall C_{0111} has vertices $\{1, 2, 3, 4\}$ in cyclic order and 1 is the only nonloop. Consider an input Φ for QCSP(C_{0111}), w.l.o.g. without any equalities, and its evaluation on C_{0111} as a game on Φ between *Prover*, playing (evaluating on C_{0111}) existential variables, and *Adversary*, playing universal variables. Adversary never gains by playing 3, as any existential edge witness to anything from $\{4, 1, 2\}$ is already an edge-witness to 3. That is, if E(x, 3) then already each of E(x, 4), E(x, 1) and E(x, 2). Similarly, Prover never gains by playing 1. Thus, Φ is true on C_{0111} iff it is true with all universal variables relativised to $\{4, 1, 2\}$ and all existential variables relativised to $\{2, 3, 4\}$. (This intuition is formalised in the notion of U-X-surjective hyper-endomorphism in [13]. What we are saying is that $\frac{2|2}{|4|}$ is a surjective hyper-endomorphism of C_{0111} .) Henceforth, we will

make this assumption of relativisation in our inputs.

Given an input Φ we will describe a procedure to establish whether it is true on C_{0111} based around a list of forbidden subinstances.

- (i.) An edge E(x, y) in \mathcal{G}_{Φ} with the later of x and y being universal.
- (*ii.*) A 3-star $E(x_1, y), E(x_2, y), E(x_3, y)$ where x_1, x_2, x_3 are universal variables coming before y existential.
- (*iii.*) A path y_1, \ldots, y_m of existential variables, where: both y_1 and y_m have edges to **two** earlier universal variables, and y_2, \ldots, y_{m-1} have edges each to **one** earlier universal variable.
- (iv.) A path y_1, \ldots, y_m of existential variables, where y_1 comes before y_m , and y_1 has an edge to an earlier universal variable. y_m has edges to two earlier universal variables at least one of which comes after y_1 . Finally, y_2, \ldots, y_{m-1} each have edges to an earlier universal variable.

These cases are illustrated in Figure 2. Using the celebrated result of [19] it can be seen that one may recognise in logspace whether or not Φ contains any of these forbidden subinstances. It is not hard to see that if Φ contains such a subinstance then Φ is false on C_{0111} (the universal variables adjacent and before y_m can be played as either 1,2 or 1,4 to force y_m to be either 2 or 4). We now claim all other instances Φ are true on C_{0111} and we demonstrate this by giving a winning strategy for Prover on such an instance. Owing to Case (*i*) being omitted, Adversary has no trivial win. Prover will now *always play* 3 if she can. Owing to the omission of Case (ii), Prover never has to answer a variable adjacent to more than two elements. It can be seen that there are few circumstances in which she can not play 3. Indeed, the only one is if she is forced at some point to play 2 or 4 as a neighbour to Adversary's having played 1. In this case, Adversary can force this response to be propagated as in the chain of cases (iii) and (iv), but because these cases are forbidden, Adversary will never succeed here in winning the game.



Fig. 2. Cases from proof of Proposition 2.

4 The reflexive cycles



Fig. 3. Left ends of the edge gadgets in reductions from $QCSP(\mathcal{K}_5)$ to $QCSP(\mathcal{C}_{1^5})$ and $QCSP(\mathcal{K}_6)$ to $QCSP(\mathcal{C}_{1^6})$. In the former case, the full gadget contains a chain of five copies of \mathcal{C}_{1^5} ; in the latter case it is a chain of six copies of \mathcal{C}_{1^6}

We will use similar edge gadgets to those of Figure 1 to prove NP-hardness of $QCSP(\mathcal{C}_{1^m})$, for $m \ge 4$. If $m \ge 4$ is even, then the edge gadget \mathcal{E}_m consists of m copies of \mathcal{C}_{1^m} where each copy – with vertices $1, \ldots, m$, is connected to its successor by edges joining vertex k with vertices k and $k + 1 \pmod{m}$. In the first of the copies, the vertex $\frac{m}{2} + 1$ is labelled y and a reflexive path of length $\frac{m}{2} - 1$ is added to the vertex labelled 1, which culminates in a vertex labelled x. The last of the copies of \mathcal{C}_{1^m} retains the vertex labelling $1, \ldots, m$ – we consider the other vertices (except for x and y) to become unlabelled. Of course, \mathcal{E}_4 is already drawn in Figure 1. The left end of \mathcal{E}_6 is drawn in Figure 3 (to the right). If $m \geq 4$ is odd, then the edge gadget \mathcal{E}_m is drawn in a similar manner, except vertex $\frac{m+1}{2} + 1$ becomes y and the reflexive path of length $\frac{m-3}{2}$ that culminates in x at one end and at the other end a vertex that makes a triangle with vertices 1 and 2, respectively. The left end of \mathcal{E}_5 is drawn in Figure 3 (to the left). These gadgets are borrowed from [9] and have the property that when the right-hand cycle \mathcal{C}_{1^m} is evaluated automorphically to itself then the rest of the cycles are also evaluated automorphically (but may twist $\frac{1}{m}$ th each turn – this is why we have m copies; m is a minimum number, more copies would still work). Finally, in the left-hand cycle it can be seen that x can be evaluated anywhere except y.

Just as in Proposition 1, we want to try to force vertex-variables v_1, \ldots, v_m , corresponding to $1, \ldots, m$, to be evaluated (up to isomorphism) around C_{1^m} .

Proposition 3. $QCSP(\mathcal{C}_{1^m})$, for any $m \ge 4$ is NP-hard.

5 Cycles whose loops induce a path

We begin by recalling the result for $QCSP(\mathcal{P}_{101})$ from [16], on which our proof for Propositions 5 and Corollary 1 will be based. In this proof we introduce the notions of *pattern* and \forall -selector that will recur in the sequel.



Fig. 4. Variable and clause gadgets in reduction to $QCSP(\mathcal{P}_{101})$.

Proposition 4. $QCSP(\mathcal{P}_{101})$ is Pspace-complete.

Proof. For hardness, we reduce from *quantified not-all-equal 3-satisfiability*, whose Pspace-completeness is well known [18], where we will ask for the extra condition that no clause has three universal variables (of course, any such instance would be trivially false). From an instance Φ of QNAESAT we will build an instance Ψ of QCSP(\mathcal{P}_{101}) such that Φ is in QNAE3SAT iff Ψ in QCSP(\mathcal{P}_{101}). We will consider the quantifier-free part of Ψ , itself a conjunction of atoms, as a graph, and use the language of homomorphisms.

We begin by describing a graph \mathcal{G}_{Φ} , whose vertices will give rise to the variables of Ψ , and whose edges will give rise to the atoms listed in the quantifier-free part of Ψ . Most of these variables will be existentially quantified, but a small handful will be universally quantified. \mathcal{G}_{Φ} consists of two reflexive paths, labelled \top and \perp which contain inbetween them gadgets for the clauses and variables of Φ . We begin by assuming that the paths \top and \perp are evaluated to vertices 1 and 3 in P_{101} , respectively (the two ends of P_{101}); later on we will show how we can effectively enforce this. Of course, once one vertex of one of the paths is evaluated to, say, 1, then that whole path must also be so evaluated – as the only looped neighbour of 1 in \mathcal{P}_{101} is 1. The gadgets are drawn in Figure 4. The *pattern* is the path \mathcal{P}_{101} , that forms the edges of the diamonds in the clause gadgets as well as the tops and bottoms of the variable gadgets. The diamonds are braced by two horizontal edges, one joining the centres of the top patterns and the other joining the centres of the bottom patterns. The \forall -selector is the path \mathcal{P}_{10} , which travels between the universal variable node v_2 and the labelled vertex \forall . (The remainder of this proof is deferred to the appendix.)

Proposition 5. Let $C_{0^d1^e}$ be a cycle in which e > d+3 (d odd) or e > d+2 (d even). Then $QCSP(C_{0^d1^e})$ is Pspace-complete.

Proof. The reduction is similar to that employed for Proposition 4. We use the pattern $\mathcal{P}_{10^{d_1}}$ and \forall -selector $\mathcal{P}_{0^{\lfloor \frac{c}{2} \rfloor_1}}$. The key part to the reduction is how we get v_{\top} and v_{\perp} to evaluate suitably. Let $x_1, \ldots, y_1, \ldots, z_1, \ldots$ be variables not appearing in $\Psi'(v_{\top}, v_{\perp})$ (cf. proof of Proposition 4). In the following, interpret $\lfloor \frac{e-d-5}{2} \rfloor$ to be 0, if $\frac{e-d-5}{2} < 0$. For d odd, set $\Psi := \forall x_1 \exists x_2, \ldots, x_{\frac{d+1}{2}}$

$$\begin{split} &\exists x_{\frac{d+3}{2}}, \dots, x_{\frac{d+3}{2}+\lfloor\frac{e-d-5}{2}\rfloor}, \exists v_{\top} \forall y_{1} \exists y_{2}, \dots, y_{\frac{d+1}{2}} \exists y_{\frac{d+3}{2}}, \dots, y_{\frac{d+3}{2}+\lfloor\frac{e-d-5}{2}\rfloor}, \exists v_{\perp} \\ &\exists z_{1}, \dots, z_{d} \\ & [x_{1} \Rightarrow x_{\frac{d+3}{2}+\lfloor\frac{e-d-5}{2}\rfloor}, v_{\top}] \land [y_{1} \Rightarrow y_{\frac{d+3}{2}+\lfloor\frac{e-d-5}{2}\rfloor}, v_{\perp}] \land [v_{\top}, z_{1}, \dots, z_{d}, v_{\perp}] \land \\ & \operatorname{Ref}(x_{\frac{d+3}{2}}, \dots, x_{\frac{d+3}{2}+\lfloor\frac{e-d-5}{2}\rfloor}) \land \operatorname{Ref}(y_{\frac{d+3}{2}}, \dots, y_{\frac{d+3}{2}+\lfloor\frac{e-d-5}{2}\rfloor}) \land \Psi'(v_{\top}, v_{\perp}) \end{split}$$

For d even, set $\Psi := \forall x_1 \exists x_2, \dots, x_{\frac{d}{2}}$

$$\begin{split} &\exists x_{\frac{d+2}{2}}, \dots, x_{\frac{d}{2}+\lfloor\frac{e-d-5}{2}\rfloor}, \exists v_{\top} \forall y_{1} \exists y_{2}, \dots, y_{\frac{d}{2}} \exists y_{\frac{d+2}{2}}, \dots, y_{\frac{d}{2}+\lfloor\frac{e-d-5}{2}\rfloor}, \exists v_{\perp} \\ &\exists z_{1}, \dots, z_{d} \\ & [x_{1} \Rightarrow x_{\frac{d}{2}+\lfloor\frac{e-d-5}{2}\rfloor}, v_{\top}] \land [y_{1} \Rightarrow y_{\frac{d}{2}+\lfloor\frac{e-d-5}{2}\rfloor}, v_{\perp}] \land [v_{\top}, z_{1}, \dots, z_{d}, v_{\perp}] \land \\ & \operatorname{Ref}(x_{\frac{d+2}{2}}, \dots, x_{\frac{d}{d}+\lfloor\frac{e-d-5}{2}\rfloor}) \land \operatorname{Ref}(y_{\frac{d+2}{2}}, \dots, y_{\frac{d}{d}+\lfloor\frac{e-d-5}{2}\rfloor}) \land \Psi'(v_{\top}, v_{\perp}) \end{split}$$

Note how the previous proof breaks down in boundary cases, for example on the cycle $C_{0^21^3}$.

The following proofs make use of reductions from $\operatorname{Ret}(\mathcal{C})$, where $|\mathcal{C}| = m$. It is ultimately intended that the variables v_1, \ldots, v_m in the created instance map automorphically to the cycle. The cycle will be found when the universally quantified v_1 is mapped to a non-looped vertex at maximal distance from the looped vertices (sometimes this is unique, other times there are two). We then require that the universally quantified x_1 be mapped to a neighbour of v_1 at maximal distance from the loops (given v_1 's evaluation, this will either be unique or there will be two). All other maps of v_1 and x_1 lead to degenerate cases.

Proposition 6. Let C be an odd m-cycle which contains an induced \mathcal{P}_{11100} (or is $\mathcal{C}_{0^21^3}$). Then QCSP(C) is NP-hard.

Proposition 7. Let C be an even m-cycle which contains an induced \mathcal{P}_{11100} . Then QCSP(C) is NP-hard.

We note that the previous two propositions do not quite use the same techniques as one another. All cases of Proposition 5 involving more than one non-loop are weakly subsumed by Propositions 6 and 7 in the sense that Pspace-completeness only becomes NP-hardness.

It is interesting to note that the Proposition 7 breaks down on even cycles with two consecutive loops only. It is no longer possible to ensure to encircle the cycle. For these cases we will need yet another specialised construction.

Proposition 8. For $m \ge 6$, let C be an even m-cycle which contains only two consecutive loops. Then QCSP(C) is NP-hard.

6 Cycles in which the loops induce a disconnected graph

Let $D_{\mathcal{C}} := \{\{\lceil \frac{d_1}{2} \rceil, \ldots, \lceil \frac{d_m}{2} \rceil\}\}$ be the multiset (of two or more elements), where d_1, \ldots, d_m are the maximal non-looped induced sections (paths) of a cycle \mathcal{C} in which the loops induce a disconnected graph. E.g. a single non-loop between two loops contributes a value $\lceil 1/2 \rceil = 1$ to $D_{\mathcal{C}}$. We need to split into three cases.

Proposition 9. Let C be a partially reflexive m-cycle in which the loops induce a disconnected graph. If D_C contains a unique maximal element $\lceil \frac{d}{2} \rceil$, then QCSP(C) is Pspace-complete.

Proposition 10. Let C be a partially reflexive m-cycle in which the loops induce a disconnected graph. If D_C contains only one value, then QCSP(C) is Pspace-complete.

Corollary 1. Let C be a partially reflexive m-cycle in which the loops induce a disconnected graph. Then QCSP(C) is Pspace-complete.

7 Classification

Theorem 1. Let $m = d + e \ge 5$. Then $QCSP(\mathcal{C}_{0^d1^e})$ is in NL if I.) m is odd and e = 1 or 2, or II.) m is even and e = 0 or 1. Otherwise, $QCSP(\mathcal{C}_{0^d1^e})$ is NP-hard.

Proof. Pspace-hardness for irreflexive odd cycles is well-known (see [17]). Hardness for cycles with disconnected loops follows from Corollary 1. Otherwise, for most cycles hardness follows from Propositions 6 and 7. For reflexive cycles see

Proposition 3 and for cycles with a single non-loop see Proposition 5. Finally, the outstanding cases of even cycles with two loops are taken care of in Proposition 8.

Now we address the NL cases. For even cycles with no loops, we are equivalent to $QCSP(\mathcal{K}_2)$ (in NL – see [17]). For even cycles $\mathcal{C}_{0^{2i+1}1}$, $QCSP(\mathcal{C}_{0^{2i+1}1})$ is equivalent to $QCSP(\mathcal{P}_{0^{i+1}1})$ (in NL – see [16]). This is because there are surjective homomorphisms from both $(\mathcal{P}_{0^{i+1}1})^2$ to $\mathcal{C}_{0^{2i+1}1}$ and $\mathcal{C}_{0^{2i+1}1}$ to $\mathcal{P}_{0^{i+1}1}$ (see [7]). For odd cycles, there are surjective homomorphisms from $(\mathcal{P}_{0^{i}1})^2$ to $\mathcal{C}_{0^{2i-1}11}$. Thus, $QCSP(\mathcal{C}_{0^{2i}1})$ is equivalent to both $QCSP(\mathcal{C}_{0^{2i-1}11})$ and $QCSP(\mathcal{P}_{0^{i}1})$ and the result follows from [16].

Theorem 2. For C a partially reflexive cycle, either QCSP(C) is in NL or it is NP-hard.

Proof. Owing to Theorem 1, it remains only to consider partially reflexive cycles on four or fewer vertices.

Firstly, we consider NL-membership. Each of $C := C_{001}, C_{011}, C_{111}, C_{0000}, C_{0001}$ and C_{0011} admits a majority polymorphism. It follows form [5] that QCSP(C) reduces to the verification of a polynomial number of instances of CSP(C^c), each of which is in NL by [8]. Finally, the case C_{0111} is taken care of in Proposition 2.

For hardness, it is well-known that quantified 3-colouring $QCSP(\mathcal{K}_3)$ is Pspacecomplete [18]. And the like result for C_{1111} appears as Proposition 1.



Fig. 5. The wavy line of tractability.

8 Conclusion

We have given a systematic classification for the QCSP of partially reflexive cycles. Many of the tractable cases can be explained by the notion of Q-core – a minimal equivalent QCSP template (see [14]) – and this is done implicitly in Theorem 1. All NP-hard cases we have seen in this paper have templates that are already Q-cores, with the sole exception of C_{0101} , whose Q-core is \mathcal{P}_{101} . By contrast, all of the tractable cases are not Q-cores, except C_{0011} and C_{0111} .

Since finding an algorithm for $\text{QCSP}(\mathcal{C}_{0111})$ we became aware of a polymorphism enjoyed by this structure. \mathcal{C}_{0111} has a ternary polymorphism f so that there is $c \in C_{0111}$ such that each of the three binary functions f(c, u, v), f(u, c, v) and f(u, v, c) is surjective. The code from Figure 6 will verify such a polymorphism and is intended to run on the excellent program of Miklós Maróti³. The naming of the vertices has been altered according to the bijection $\frac{\frac{13}{20}}{\frac{31}{44}}$ (vertices are numbered from 0 for the computer). If follows from [5] that \mathcal{C}_{0111} is 2-collapsible,

```
arity 3; size 4; idempotent; preserves 01 10 12 21 23 32 30 03 00 11 22; value 302=0; value 320=2; value 311=1; value 123=1; value 223=2; value 003=0; value 032=0; value 030=1; value 230=2
```

Fig. 6. Code for Maróti's program (semicolons indicate new line).

and hence $QCSP(\mathcal{C}_{0111})$ may be placed in NL by other means.

We would like to improve the lower bound from NP-hardness to Pspacehardness in the cases of Propositions 3, 6, 7 and 8. This might be quite messy in the last three cases, involving careful consideration of the hardness proof for the retraction problem. For the reflexive cycles, though, it just requires more careful analysis of the degenerate cases. This is because we may only have a homomorphic image under f of the cycle for v_1, \ldots, v_m , but the universal variables may be evaluated outside of the image of $f(\{v_1, \ldots, v_m\})$. We will require something of the following form (\mathcal{E}_m is defined at the beginning of Section 4).

Conjecture 1. Let f be a function from $\{1, \ldots, m\}$ in \mathcal{E}_m to \mathcal{C}_{1^m} that is a nonsurjective homomorphism. Let f_{ij} , for some $i, j \in \{1, \ldots, m\}$, be the partial function that extends f from $\{1, \ldots, m, x, y\}$ in \mathcal{E}_m to \mathcal{C}_{1^m} , by mapping $x \mapsto i$ and $y \mapsto j$. Then f_{ij} can be extended to a homomorphism from \mathcal{E}_m to \mathcal{C}_{1^m} .

The proof of this seems to be rather technical. For those who doubt it, let us remind ourselves that the length of the chain in \mathcal{E}_m may be any fixed function of m, and the reduction of Propositions 1 and 3 will still work. The conjecture is surely easier to prove if we make the chains much longer (say exponential in m).

³ See: http://www.math.u-szeged.hu~maroti/applets/GraphPoly.html

Finally, we conjecture that none of the NL cases are NL-hard, and that most likely our dichotomy can be perfected to L/ Pspace-complete.

Acknowledgements. The authors thank the referees and are grateful to St. Catherine.

References

- 1. BARTO, L., KOZIK, M., AND NIVEN, T. The CSP dichotomy holds for digraphs with no sources and no sinks (a positive answer to a conjecture of Bang-Jensen and Hell). *SIAM Journal on Computing* 38, 5 (2009), 1782–1802.
- BÖRNER, F., BULATOV, A. A., CHEN, H., JEAVONS, P., AND KROKHIN, A. A. The complexity of constraint satisfaction games and qcsp. *Inf. Comput. 207*, 9 (2009), 923–944.
- BULATOV, A. A dichotomy theorem for constraint satisfaction problems on a 3-element set. J. ACM 53, 1 (2006), 66–120.
- BULATOV, A., KROKHIN, A., AND JEAVONS, P. G. Classifying the complexity of constraints using finite algebras. SIAM Journal on Computing 34 (2005), 720–742.
- 5. CHEN, H. The complexity of quantified constraint satisfaction: Collapsibility, sink algebras, and the three-element case. SIAM J. Comput. 37, 5 (2008), 1674–1701.
- CHEN, H. Meditations on quantified constraint satisfaction. CoRR abs/1201.6306 (2012). Appeared in Festschrift for Dexter Kozen 60th.
- CHEN, H., MADELAINE, F., AND MARTIN, B. Quantified constraints and containment problems. In 23rd Annual IEEE Symposium on Logic in Computer Science (2008), pp. 317–328.
- DALMAU, V., AND KROKHIN, A. A. Majority constraints have bounded pathwidth duality. *Eur. J. Comb.* 29, 4 (2008), 821–837.
- FEDER, T., AND HELL, P. List homomorphisms to reflexive graphs. J. Comb. Theory, Ser. B 72, 2 (1998), 236–250.
- FEDER, T., HELL, P., JONSSON, P., KROKHIN, A. A., AND NORDH, G. Retractions to pseudoforests. SIAM J. Discrete Math. 24, 1 (2010), 101–112.
- FEDER, T., AND VARDI, M. The computational structure of monotone monadic SNP and constraint satisfaction: A study through Datalog and group theory. *SIAM Journal on Computing 28* (1999), 57–104.
- 12. HELL, P., AND NEŠETŘIL, J. On the complexity of H-coloring. Journal of Combinatorial Theory, Series B 48 (1990), 92–110.
- MADELAINE, F., AND MARTIN, B. A tetrachotomy for positive equality-free logic. In Proceedings of the 26th Annual IEEE Symposium on Logic in Computer Science, LICS 2011 (2011).
- MADELAINE, F. R., AND MARTIN, B. Containment, equivalence and coreness from CSP to QCSP and beyond. In 18th CP 2012 (2012), pp. 480–495.
- 15. MADELAINE, F. R., MARTIN, B., AND STACHO, J. Constraint satisfaction with counting quantifiers. In 7th International CSR 2012 (2012), pp. 253–265.
- 16. MARTIN, B. QCSP on partially reflexive forests. In 17th CP 2011 (2011).
- MARTIN, B., AND MADELAINE, F. Towards a trichotomy for quantified H-coloring. In 2nd Conf. on Computatibility in Europe, LNCS 3988 (2006), pp. 342–352.
- 18. PAPADIMITRIOU, C. H. Computational Complexity. Addison-Wesley, 1994.
- 19. REINGOLD, O. Undirected connectivity in log-space. J. ACM 55, 4 (2008), 1–24.
- SCHAEFER, T. J. The complexity of satisfiability problems. In Proceedings of STOC'78 (1978), pp. 216–226.

Appendix

8.1 Proposition 3 in full

Proposition 3. $QCSP(\mathcal{C}_{1^m})$, for any $m \ge 4$ is NP-hard.

Proof. We will reduce from the problem $\text{CSP}(\mathcal{K}_m)$ in essentially the same manner, for m > 4, as we did for m := 4 and $\text{QCSP}(\mathcal{K}_m)$ in Proposition 1. The salient points remain unchanged, except that we use the gadgets \mathcal{E}_m for edges in the graph input of $\text{QCSP}(\mathcal{K}_m)$, instead of \mathcal{E}_4 , and we may assume that all variables in our reduction instance are existential. So it is that we build a formula $\Psi'(v_1, \ldots, v_m)$ in place of the formula $\Psi'(v_1, \ldots, v_4)$ that we built in Proposition 1. For m > 4, let us define the shorthands (the variables x' and $x_1, \ldots, x_{\frac{m}{2}-2}$ are new and must not appear in φ).

$$\Diamond x \ \varphi(x) := \forall x' \exists x_1, \dots, x_{\lceil \frac{m}{2} \rceil - 2} \exists x \ E(x', x_1) \land \dots \land E(x_{\lceil \frac{m}{2} \rceil - 2}, x) \land \varphi(x)$$

Recalling that C_{1^m} is reflexive, it is not hard to verify that $\Diamond x$ may be satisfied on C_{1^m} iff x takes at least 2 values (m even) or iff x takes at least two non-adjacent values (m odd). To elucidate, let us consider the two examples, when m := 6, 5.

$$\begin{aligned} & \Diamond x \ \varphi(x) := \forall x' \exists x_1 \exists x \ E(x', x_1) \land E(x_1, x) \land \varphi(x) \text{ if } m := 6 \\ & \Diamond x \ \varphi(x) := \forall x' \exists x \ E(x', x) \land \varphi(x) & \text{ if } m := 5 \end{aligned}$$

For m even, we set $\Psi :=$

$$\exists v_1 \forall v_{\frac{m}{2}+1} \exists v_2, \dots, v_{\frac{m}{2}} \quad [v_1 \Rightarrow v_{\frac{m}{2}+1}] \land \Diamond v_{\frac{m}{2}+2} \exists v_{\frac{m}{2}+3}, \dots, v_m [v_{\frac{m}{2}+1}, v_{\frac{m}{2}+2} \Rightarrow v_1] \land \\ \Psi'(v_1, \dots, v_m)$$

For m odd, note that $\exists x \Diamond y \ E(x, y)$ selects adjacent pairs (but x and y distinct!). For m odd, we set $\Psi :=$

$$\exists v_1 \diamondsuit v_2 \ E(v_1, v_2) \land \forall v_{\frac{m+3}{2}} \exists v_3, \dots, v_{\frac{m+1}{2}}, v_{\frac{m+5}{2}}, \dots, v_m \ [v_2 \Rightarrow v_{\frac{m+3}{2}}] \land [v_{\frac{m+3}{2}} \Rightarrow v_1] \land$$
$$\Psi'(v_1, \dots, v_m)$$

It is not hard to see that Ψ forces on some evaluation of v_1, \ldots, v_m that these map isomorphically to $1, \ldots, m$ in \mathcal{C}_{1^m} . If they do not, then they instead map according to a homomorphism f, and we are in a degenerate case. If they had mapped according to the identity, then a yes-instance (all other variables are existential) extends to homomorphism (from the graph associated with Ψ) by, say, g. But now $f \circ g$ extends to homomorphism in the degenerate case, and we are done.

8.2 Conclusion of proof of Proposition 4

For each existential variable v_1 in Φ we add the gadget on the far left, and for each universal variable v_2 we add the gadget immediately to its right. There is a single vertex in that gadget that will eventually give rise to a variable in Ψ that is universally quantified, and it is labelled \forall (however, it is the vertex at the other end, labelled v_2 in Figure 4, that actually corresponds to the universally quantified variable). For each clause of Φ we introduce a copy of the clause gadget drawn on the right. We then introduce an edge between a variable v and literal l_i ($i \in \{1, 2, 3\}$) if $v = l_i$ (note that all literals in QNAE3SAT are positive). We reorder the literals in each clause, if necessary, to ensure that literal l_2 of any clause is never a variable in Φ that is universally quantified. It is not hard to verify that homomorphisms from \mathcal{G}_{Φ} to \mathcal{P}_{101} (such that the paths \top and \bot are evaluated to 1 and 3, respectively) correspond exactly to satisfying not-all-equal assignments of Φ . The looped vertices must map to either 1 or $3 - \top$ or \bot – and the clause gadgets forbid exactly the all-equal assignments. Now we will consider the graph \mathcal{G}_{Φ} realised as a formula Ψ'' , in which we will existentially quantify innermost all of the variables of Ψ'' except:

- one variable each, v_{\top} and v_{\perp} , corresponding respectively to some vertex from the paths \top and \perp , and
- all variables corresponding to the centre vertex of an existential variable gadget, and
- all variables corresponding to the \forall -selector (the centre vertex of a universal variable gadget and the vertex labelled \forall).

We now build Ψ' by quantifying, adding outermost and in the order of the quantifiers of Φ :

- existentially, the variable corresponding to the centre vertex of an existential variable gadget,
- universally, the variable corresponding to the extra vertex labelled \forall of a universal variable gadget, and then existentially, the remaining vertices of the \forall -selector.

The reason we do not directly universally quantify the vertex associated with a universal variable is because we want it to be forced to range over only the looped vertices 1 and 3 (which it does as its unlooped neighbour \forall is forced to range over all $\{1, 2, 3\}$). $\Psi'(v_{\top}, v_{\perp})$ is therefore a positive Horn formula with two free variables, v_{\top} and v_{\perp} , such that, Φ is QNAE3SAT iff $\mathcal{P}_{101} \models \Psi'(1, 3)$. Finally, we construct Ψ from Ψ' with the help of two \forall -selectors, adding new variables v'_{\top} and v'_{\perp} , and setting

$$\Psi := \forall v'_{\top}, v'_{\perp} \exists v_{\top}, v_{\perp} \ E(v'_{\top}, v_{\top}) \land E(v'_{\perp}, v_{\perp}) \land \Psi'(v_{\top}, v_{\perp}).$$

The purpose of universally quantifying the new variables v'_{\top} and v'_{\perp} , instead of directly quantifying v_{\top} and v_{\perp} , is to force v'_{\top} and v'_{\perp} to range over $\{1, 3\}$ (recall that $E(v_{\top}, v_{\top})$ and $E(v_{\perp}, v_{\perp})$ are both atoms of Ψ). This is the same reason we add the vertex \forall to the universal variable gadget.

We claim that $\mathcal{P}_{101} \models \Psi'(1,3)$ iff $\mathcal{P}_{101} \models \Psi$. It suffices to prove that $\mathcal{P}_{101} \models \Psi'(1,3)$ implies $\mathcal{P}_{101} \models \Psi'(3,1), \Psi'(1,1), \Psi'(3,3)$. The first of these follows by symmetry. The second two are easy to verify, and follow because the second literal in any clause is forbidden to be universally quantified in Φ . If both paths \top and \bot are w.l.o.g. evaluated to 1, then, even if some l_1 - or l_3 -literals are forced to evaluate to 3, we can still extend this to a homomorphism from \mathcal{G}_{Φ} to \mathcal{P}_{101} .

8.3 Latter part of Section 5 in full

The following proofs make use of reductions from $\operatorname{Ret}(\mathcal{C})$, where |C| = m. It is ultimately intended that the variables v_1, \ldots, v_m in the created instance map automorphically to the cycle. The cycle will be found when the universally quantified v_1 is mapped to a non-looped vertex at maximal distance from the looped vertices (sometimes this is unique, other times there are two). We then require that the universally quantified x_1 be mapped to a neighbour of v_1 at maximal distance from the loops (given v_1 's evaluation, this will either be unique or there will be two). All other maps of v_1 and x_1 lead to degenerate cases.

Proposition 6. Let C be an odd *m*-cycle which contains an induced \mathcal{P}_{11100} (or is $\mathcal{C}_{0^21^3}$). Then QCSP(C) is NP-hard.

Proof. We reduce from the problem of $\operatorname{Ret}(\mathcal{C})$, known to be NP-complete from [10]. Let \mathcal{C} be of the form $\mathcal{C}_{0^d 1^e}$. We find a copy of \mathcal{C} with the variables v_1, \ldots, v_m in the following fashion.

$$\begin{split} \forall v_1 \exists v_2, \dots, v_{\frac{m-1}{2}} \\ \forall x_1 \exists x_2, \dots, x_{\frac{m+1}{2}} \exists v_{\frac{m+1}{2}} [v_1 \Rightarrow v_{\frac{m+1}{2}}] \wedge [x_1 \Rightarrow x_{\frac{m+1}{2}}] \wedge \operatorname{Ref}(v_{\lceil \frac{d+2}{2} \rceil}, \dots, v_{\frac{m+1}{2}}) \wedge \\ \operatorname{Ref}(x_{\lceil \frac{d+2}{2} \rceil}, \dots, x_{\frac{m+1}{2}}) \wedge x_{\frac{m+1}{2}} = v_{\frac{m+1}{2}} \wedge \\ \exists v_m, \dots, v_{\frac{m+3}{2}} x_{\frac{m-1}{2}} = v_{\frac{m+3}{2}} \wedge [v_1, v_m \Rightarrow v_{\frac{m+1}{2}}] \wedge \operatorname{Ref}(v_{\frac{m+3}{2}}, \dots, v_{\lceil \frac{m+2}{2} \rceil}) \wedge \dots \end{split}$$

It is interesting to see where the previous proof breaks down cycles in which there are less than three loops or a single non-loop. Let us consider the sentence obtained in the case for C_{10^4} .

$$\forall v_1 \exists v_2, v_3 \forall x_1 \exists x_2, x_3 \ [v_1 \Rightarrow v_3] \land [x_1 \Rightarrow x_3] \land E(v_3, v_3) \land x_3 = v_3 \land \\ \exists v_5, v_4 \ x_2 = v_4 \land [v_1, v_5 \Rightarrow v_3] \land E(v_3, v_3) \land \dots$$

The problem with C_{10^4} occurs when v_1 and x_1 are both evaluated as, say, vertex 3 (there can be no v_5 between v_1 and x_2).

Proposition 7. Let C be an even *m*-cycle which contains an induced \mathcal{P}_{11100} . Then QCSP(C) is NP-hard.

Proof. We reduce from the problem $\operatorname{Ret}(\mathcal{C})$, known to be NP-complete from [10]. Let \mathcal{C} be of the form $\mathcal{C}_{0^d 1^e}$. We find a copy of \mathcal{C} with the variables v_1, \ldots, v_m in the following fashion.

 $\begin{array}{l} \forall v_1 \exists v_2, \dots, v_{\lceil \frac{m-d+2}{2} \rceil} \\ \forall x_1 \exists x_2, \dots, x_{\lceil \frac{m-d+2}{2} \rceil} [v_1 \Rightarrow v_{\lceil \frac{m-d+2}{2} \rceil}] \land [x_1 \Rightarrow x_{\lceil \frac{m-d+2}{2} \rceil}] \land \operatorname{Ref}(v_{\lceil \frac{m-d+2}{2} \rceil}, x_{\lceil \frac{m-d+2}{2} \rceil}) \land \\ \exists v_{\lceil \frac{m-d+2}{2} \rceil+1}, \dots, v_{\lceil \frac{m-d+2}{2} \rceil+d-1} \\ v_{\lceil \frac{m-d+2}{2} \rceil+d-1} = x_{\lceil \frac{m-d+2}{2} \rceil} \land [v_{\lceil \frac{m-d+2}{2} \rceil+1} \Rightarrow v_{\lceil \frac{m-d+2}{2} \rceil+d-1}] \land \operatorname{Ref}(v_{\lceil \frac{m-d+2}{2} \rceil+1}, v_{\lceil \frac{m-d+2}{2} \rceil+d-1}) \land \\ \exists v_{\lceil \frac{m-d+2}{2} \rceil+d}, \dots, v_m [v_{\lceil \frac{m-d+2}{2} \rceil+d-1} \Rightarrow v_m] \end{array}$

We note that the previous two propositions do not quite use the same techniques as one another. All cases of Proposition 5 involving more than one non-loop are weakly subsumed by Propositions 6 and 7 in the sense that Pspace-completeness only becomes NP-hardness.

It is interesting to see where the Proposition 7 breaks down on even cycles with two consecutive loops only. It is no longer possible to ensure to encircle the cycle. For these cases we will need yet another specialised construction.

Proposition 8. For $m \ge 6$, let \mathcal{C} be an even *m*-cycle which contains only two consecutive loops. Then $QCSP(\mathcal{C})$ is NP-hard.

Proof. Let us label the vertices of the cycle 1 to m s.t. the loops are at positions $\frac{m}{2}$ and $\frac{m}{2} + 1$. We introduce the following shorthand $(u' \text{ does not appear in } \varphi)$.

$$\heartsuit u \varphi(u) := \forall u' \exists u \ E(u', u) \land \varphi(u)$$

The \heartsuit quantifier is some kind of weak universal quantifier, with the following important property: if $\varphi(2)$ and $\varphi(m-1)$ are false then $\varphi(1)$ and $\varphi(m)$ are true.

Unlike the previous proofs, which used just v_1, \ldots, v_m , we will need also w_1, \ldots, w_n and we will be able to say that at least one of them maps automorphically to C. We begin with

$$\forall v_1 \exists v_2, \dots, v_{\frac{m}{2}} [v_1 \Rightarrow v_{\frac{m}{2}}] \land E(v_{\frac{m}{2}}, v_{\frac{m}{2}}) \land \\ \forall w_1 \exists w_2, \dots, w_{\frac{m}{2}} [w_1 \Rightarrow w_{\frac{m}{2}}] \land E(w_{\frac{m}{2}}, w_{\frac{m}{2}}) \land$$

If v_1 and w_1 are evaluated on 1 and m, respectively, then $v_{\frac{m}{2}}$ and $w_{\frac{m}{2}}$ will be on $\frac{m}{2}$ and $\frac{m}{2} + 1$, respectively. In the past, we did not care about degenerate cases, so long as they extend to homomorphism, but here we will use the fact that some degenerate cases *could not* extend to homomorphism, if we had used a universal quantifier. Assuming v_1, w_1 is mapped to 1, m, these degenerate cases would come about with z mapped to either 2 or m - 1. Thus, using the \heartsuit quantifier we can force z to be mapped to both 1 and m - 1. We complete with

$$\begin{aligned} & \forall z \; \exists x_2, \dots, x_{\frac{m}{2}} \exists v_{\frac{m}{2}+1}, v_{\frac{m}{2}+2} \quad E(v_{\frac{m}{2}+1}, v_{\frac{m}{2}+1}) \land E(v_{\frac{m}{2}+1}, v_{\frac{m}{2}+2}) \land \\ & \quad [z, x_2 \Rightarrow x_{\frac{m}{2}}, v_{\frac{m}{2}+1}, v_{\frac{m}{2}+2}] \land v_{\frac{m}{2}+1} = x_{\frac{m}{2}} \land v_{\frac{m}{2}+2} = x_{\frac{m}{2}-1} \land \\ & \quad \exists y_2, \dots, y_{\frac{m}{2}} \exists w_{\frac{m}{2}+1}, w_{\frac{m}{2}+2} \in E(w_{\frac{m}{2}+1}, w_{\frac{m}{2}+1}) \land E(w_{\frac{m}{2}+1}, w_{\frac{m}{2}+2}) \land \\ & \quad [z, y_2 \Rightarrow y_{\frac{m}{2}}, w_{\frac{m}{2}+1}, w_{\frac{m}{2}+2}] \land w_{\frac{m}{2}+1} = y_{\frac{m}{2}} \land w_{\frac{m}{2}+2} = y_{\frac{m}{2}-1} \land \\ & \quad \exists v_{\frac{m}{2}+3}, \dots, v_m [v_{\frac{m}{2}+2} \Rightarrow v_m, v_1] \land \\ & \quad \exists w_{\frac{m}{2}+3}, \dots, w_m [w_{\frac{m}{2}+2} \Rightarrow w_m, w_1]. \end{aligned}$$

The key observation is that the statements $v_{\frac{m}{2}+2} = x_{\frac{m}{2}-1}$ and $w_{\frac{m}{2}+2} = y_{\frac{m}{2}-1}$ are impossible to satisfy if v_1, w_1, z is mapped to 1, m, 2 or 1, m, m - 1. This is why a universal quantifier can not be used for z and why we can not add $v_{\frac{m}{2}+2} = x_{\frac{m}{2}-1}$ to the construction from the previous proposition.

8.4 Section 6 in full

Let $D_{\mathcal{C}} := \{\{\lceil \frac{d_1}{2} \rceil, \dots, \lceil \frac{d_m}{2} \rceil\}\}$ be the multiset (of two or more elements), where d_1, \dots, d_m are the maximal non-looped induced sections (paths) of a cycle \mathcal{C} in

which the loops induce a disconnected graph. E.g. a single non-loop between two loops contributes a value $\lceil 1/2 \rceil = 1$ to $D_{\mathcal{C}}$. We need to split into three cases. In the following, as in [16], we refer to paths involving loops and non-loops according to words $\{0, 1\}^*$ – where 0 indicates a non-loop and 1 a loop.

Proposition 9 Let C be a partially reflexive *m*-cycle in which the loops induce a disconnected graph. If $D_{\mathcal{C}}$ contains a unique maximal element $\lceil \frac{d}{2} \rceil$, then QCSP(C) is Pspace-complete.

Proof. Let \mathcal{P} be the partially reflexive path, obtained by removing the middle (or middle two) vertices in the unique section of non-loops of maximal length in \mathcal{C} . It follows from inspection of the proofs from [16] that QCSP(\mathcal{P}) is Pspace-complete even when the universal variables are restricted to range over some subset of P that contains at least each of the two end-most loops. We reduce from this problem. The ∀-selector will be $\mathcal{P}_{0^{\lceil \frac{d}{2} \rceil_1}}$. It remains to show how to relativise the existential variables so they range over precisely $P \subseteq C$. In order to do this, we ensure that each existential variable is at distance $\lceil \frac{d}{2} \rceil - 1$ from its nearest loop, by adding a path of length $\lceil \frac{d}{2} \rceil - 1$ to a loop. The reduction is clear and the result follows.

Proposition 10 Let C be a partially reflexive *m*-cycle in which the loops induce a disconnected graph. If $D_{\mathcal{C}}$ contains only one value, then $QCSP(\mathcal{C})$ is Pspace-complete.

Proof. Let \mathcal{C} be over n vertices, and suppose $D_{\mathcal{C}}$ contains k instances of its maximal element f. These may be produced by sections of non-loops of length 2f or 2f - 1. Let g^+ be the maximal such length of non-loops. The reduction is similar to that employed in Proposition 4. The pattern will be \mathcal{P}_{10g^+1} , while the \forall -selector will be \mathcal{P}_{0f_1} . It remains to explain how to evaluate v_{\top} and v_{\perp} . We begin with a universal variable followed by a path of f non-loops and then a loop $(0^f 1)$ - just as in the \forall -selector. We then have a sequence of alternations of $1^n 0^{2f}$ followed finally by 1^n . The number of these alternations is $\lceil (\lfloor \frac{k}{2} \rfloor - 1)/2 \rceil$ from v_{\top} and $\lfloor (\lfloor \frac{k}{2} \rfloor - 1)/2 \rfloor$ from v_{\perp} . The point is that when the initial universal variables are evaluated at opposite sides of \mathcal{C} , with respect to the k large gaps, then some evaluation forces v_{\top} and v_{\perp} to be at distance exactly distance g^+ , across non-loops.

Corollary 1 Let C be a partially reflexive *m*-cycle in which the loops induce a disconnected graph. Then QCSP(C) is Pspace-complete.

Proof. Taking into consideration the results of Propositions 9 and 10, the only remaining case is where $D_{\mathcal{C}}$ has multiple -k – copies of its maximal element, and some other smaller elements also. But, this case can be handled very similarly to Proposition 10. Simply amend the alternating motif $1^n 0^{2f}$ to $1^n (0^{2f-2} 1^n)^n 0^{2f}$ – thereby allowing any section of 2f - 2 or fewer non-loops to be crossed for free.