ALGEBRAIC METHODS FOR COMPUTING SMALLEST ENCLOSING AND CIRCUMSCRIBING CYLINDERS OF SIMPLICES

RENÉ BRANDENBERG AND THORSTEN THEOBALD

ABSTRACT. We provide an algebraic framework to compute smallest enclosing and smallest circumscribing cylinders of simplices in Euclidean space \mathbb{E}^n . Explicitly, the computation of a smallest enclosing cylinder in \mathbb{E}^3 is reduced to the computation of a smallest circumscribing cylinder. We improve existing polynomial formulations to compute the locally extreme circumscribing cylinders in \mathbb{E}^3 and exhibit subclasses of simplices where the algebraic degrees can be further reduced. Moreover, we generalize these efficient formulations to the *n*-dimensional case and provide bounds on the number of local extrema. Using elementary invariant theory, we prove structural results on the direction vectors of any locally extreme circumscribing cylinder for regular simplices.

1. INTRODUCTION

Radii (of various types) belong to the most important functionals of polytopes and general convex bodies in Euclidean space \mathbb{E}^n [3, 14, 16], and they are related to applications in computer vision, robotics, computational biology, functional analysis, and statistics (see [15]). Following the notation in [3], the *outer j-radius* $R_j(\mathcal{P})$ of a convex body $\mathcal{C} \subset \mathbb{E}^n$ is the radius of a smallest enclosing *j*-dimensional sphere in the optimal orthogonal projection of \mathcal{C} onto a *j*-dimensional linear subspace. Studying these radii, mainly for regular simplices and regular polytopes, is a classical topic of convex geometry (see [2, 4, 11, 14]).

From the computational point of view, most of the existing algorithms for computing these radii focus on approximation [5, 17]. A major reason is that exact computations lead to algebraic problems of high degree, even for computing, say, the outer (n-1)-radius in \mathbb{E}^n (already if n = 3). However, since some approaches for computing radii of general polytopes consider the computation of a smallest enclosing cylinder of a simplex as a black box within a larger computation [1, 23], these core problems on simplices are of fundamental importance.

Recently, the authors of [9] demonstrated that using their state-of-the-art numerical polynomial solvers, various problems related to cylinders in \mathbb{E}^3 can be solved rather efficiently. In particular, the authors give a polynomial formulation for the smallest circumscribing cylinder of a simplex in \mathbb{E}^3 , whose Bézout number – the product of the degrees of the polynomial equations – is 60. However, these equations contain certain undesired solutions with multiplicity 4, and as a result of these multiplicities the computation times

Date: November 19, 2018.

²⁰⁰⁰ Mathematics Subject Classification. 51N20, 52B55, 68U05, 68W30, 90C90.

(using state-of-the-art numerical techniques) are about a factor 100 larger than those of similar problems in which all solutions occur with multiplicity 1.

Here, we provide a general algebraic framework for computing smallest enclosing and circumscribing cylinders of simplices in \mathbb{E}^n . First we reduce the computation of a smallest enclosing cylinder in \mathbb{E}^3 to the computation of a smallest circumscribing cylinder, thus combining these two problems. Then we investigate smallest circumscribing cylinders of simplices in \mathbb{E}^3 . We improve the results of [9] by providing a polynomial formulation for the locally extreme cylinders, whose Bézout bound is 36 and whose solutions generically have multiplicity one. Our formulations use techniques from the paper [21] which studies the lines simultaneously tangent to four unit spheres. These techniques also enable us to present classes of simplices for which the algebraic degrees in computing the smallest circumscribing cylinder can be considerably reduced.

Then, in Section 4, we give a generalization of our approach to smallest circumscribing cylinders of a simplex in \mathbb{E}^n . Based on this formulation we give bounds on the number of locally extreme cylinders based on the Bézout number. Since this bound is not tight, we provide better bounds for small dimensions; these bounds are based on mixed volume computations and Bernstein's Theorem. Moreover, we study in detail the locally extreme circumscribing cylinders of a regular simplex in \mathbb{E}^n . To exploit many symmetries in the analysis, we provide a formulation based on symmetric polynomials. Using elementary invariant theory we show that the direction vector of every locally extreme circumscribing cylinder has at most three distinct values in its components. With this result we can illustrate our combinatorial results on the number of solutions for general simplices.

As a byproduct of our computational studies, we discovered a subtle but severe mistake in the paper [31] on the explicit determination of the outer (n-1)-radius for a regular simplex in \mathbb{E}^n , thus completely invalidating the proof given there. In the appendix we give a description of that flaw, including some computer-algebraic calculations illustrating it.

2. Preliminaries and background

2.1. *j*-radii and cylinders. Throughout the paper we work in Euclidean space \mathbb{E}^n , i.e., \mathbb{R}^n with the usual scalar product $x \cdot y = \sum_{i=1}^3 x_i y_i$ and norm $||x|| = (x \cdot x)^{1/2}$. We write x^2 for $x \cdot x$.

A *j*-flat is an affine subspace of dimension j. For a convex polytope $\mathcal{P} \subset \mathbb{E}^n$ (or a finite point set $\mathcal{P} \subset \mathbb{E}^n$) and a *j*-flat E, we consider

$$\mathcal{RD}(\mathcal{P}, E) := \max_{p \in \mathcal{P}} \operatorname{dist}(p, E),$$

where dist(p, E) denotes the Euclidean distance from p to E. The outer *j*-radius of \mathcal{P} is

$$R_j(\mathcal{P}) := \min_{E \text{ is an } (n-j)-\text{flat}} \mathcal{RD}(\mathcal{P}, E).$$

The choice of the indexing in the *j*-radius stems from the fact that it measures the radius of an enclosing *j*-dimensional sphere in the optimal orthogonal projection of \mathcal{P} onto a *j*-dimensional linear subspace (cf. [3, 14]).

One of the most natural representatives of this class is the one with j = 2, n = 3, i.e., the smallest enclosing (circular) cylinder of a polytope. In \mathbb{E}^n , we define a cylinder to be a set of the form

$$\operatorname{bd}(\ell + \rho \mathbb{B}^n),$$

where ℓ is a line in \mathbb{E}^n , \mathbb{B}^n denotes the unit ball, $\rho > 0$, the addition denotes the Minkowski sum, and bd(·) denotes the boundary of a set. We say that *P* can be enclosed in a cylinder \mathcal{C} if *P* is contained in the convex hull of \mathcal{C} . Thus the outer (n-1)-radius gives the radius of the smallest enclosing cylinder of a polytope.

A simplex in \mathbb{E}^n is the convex hull of n + 1 affinely independent points. An enclosing cylinder \mathcal{C} of a simplex \mathcal{P} is called a *circumscribing* cylinder of \mathcal{P} if all the vertices of \mathcal{P} are contained in (the hypersurface) \mathcal{C} .

2.2. Smallest circumscribing cylinders and smallest enclosing cylinders. The following statement connects the computation of a smallest enclosing cylinder of a polytope with the computation of a smallest circumscribing cylinder of a simplex.¹

Theorem 1. Let $\mathcal{P} = \{p_1, \ldots, p_m\}$ be a set of $m \ge 4$ points in \mathbb{E}^3 , not all collinear. If \mathcal{P} can be enclosed in a circular cylinder \mathcal{C} of radius r, then there exists a circular cylinder \mathcal{C}' of radius r enclosing all elements of \mathcal{P} such that the surface \mathcal{C}' passes through

- (i) at least four non-collinear points of \mathcal{P} , or
- (ii) three non-collinear points of \mathcal{P} , and the axis ℓ of \mathcal{C}' is contained in
 - (a) the cylinder naturally defined by spheres of radius r centered at two of these points;
 - (b) the double cone naturally defined by spheres of radius r centered at two of these points (and these spheres are disjoint);
 - (c) or the set of lines which are tangent to the two spheres of radius r centered at two these points and which are contained in the plane equidistant from these points (and the spheres are non-disjoint).

Moreover, C can be transformed into C' by a continuous motion.

Figures 1 and 2 visualize the three geometric properties in the second possibility.

Since the second possibility in Theorem 1 characterizes the possible special cases, this lemma in particular reduces the computation of a smallest enclosing cylinder of a simplex in \mathbb{E}^3 to the computation of a smallest circumscribing cylinder of a simplex. Namely, it suffices to compute the smallest circumscribing cylinder (corresponding to case (i)) as well as the smallest enclosing cylinders whose axes satisfies one of the condition in (ii); the latter case gives a constant number of problems of smaller algebraic degree (since the positions of the axes are very restricted).

Remark 2. Before we start with the proof, we remark that Theorem 1 and its different cases show a quite similar behaviour as the well known statement that the (unique) circumsphere of a simplex touches all its vertices, or one of its great (n-1)-circles is the circumsphere of one of the (n-1)-faces of the simplex (see [2, p. 54]).

¹We remark that a similar statement has already been used in [23], but the manuscript referenced there does not contain a complete proof.



FIGURE 1. Extreme situations of the set of hyperboloids for disjoint spheres



(a) Hyperboloid for $0 < x_h < 2r^2/a$ (b) Degenerated hyperboloid for $x_h = a/2$

FIGURE 2. The left figure shows a general situation for disjoint spheres; the right figure shows an extreme situation for non-disjoint spheres

In the proof we will apply the following geometric equivalence. A point $x \in \mathbb{E}^3$ is enclosed in a cylinder with axis ℓ if and only if ℓ is a transversal of the sphere with radius r centered at x (i.e., ℓ is a line intersecting the sphere).

Proof of Theorem 1. Let \mathcal{C} be a cylinder with axis ℓ and radius r enclosing \mathcal{P} . Then, denoting by $S_i := S(p_i, r)$ the sphere with radius r centered at p_i , ℓ is a common transversal to S_1, \ldots, S_m . By continuously translating and rotating ℓ , we can assume that ℓ is tangent to two of the spheres, say S_1 and S_2 . Further, by changing coordinates, we can assume that S_1 and S_2 have the form $S_1 = S((0,0,0)^T, r), S_2 = S((a,0,0)^T, r)$ for some a > 0.

The set of lines tangent to two spheres of radius r constitutes a set of hyperboloids (see, e.g., [8, 18]). Moreover, any of these hyperboloids touches the sphere S_1 on a circle lying in a hyperplane parallel to the yz-plane. Hence, the set of hyperboloids can be parametrized by the x-coordinate of this hyperplane which we denote by x_h .

If $S_1 \cap S_2 = \emptyset$ then the boundary values are $x_h = 0$ and $x_h = 2r^2/a$. These two extreme situations yield a cylinder and a double cone with apex $(a/2, 0, 0)^T$, respectively (see Figure 1). For $0 < x_h < 2r^2/a$ we obtain a hyperboloid of one sheet (see Figure 2(a)). If $S_1 \cap S_2 \neq \emptyset$ then the boundary values are $x_h = 0$ and $x_h = a/2$. Here, for $0 < x_h < a/2$ we obtain hyperboloids of one sheet, too. For $x_h = a/2$ the hyperboloid degenerates to a set of tangents which are tangents to the circle with radius $r_c = \sqrt{4r^2 - a^2}$ in the hyperplane x = a/2 (see Figure 2(b)).

Let $x_{h,0}$ be the parameter value of the hyperboloid containing the line ℓ . The tangent to S_1 and S_2 is contained in the hyperboloid with some parameter value $x_{h,0}$. By decreasing the parameter x_h starting from $x_{h,0}$ the hyperboloid changes its shape towards the cylinder around S_1 and S_2 . Let $x_{h,1}$ be the infimum of all $0 \leq x_h < x_{h,0}$ such that the hyperboloid does not contain a generating line tangent to some other sphere $S(p_i, r)$ for some $3 \leq i \leq m$. If $x_{h,1} = 0$, then by choosing any point of \mathcal{P} not collinear to p_1 and p_2 we are in case (ii) (a).

If $x_{h,1} > 0$ then let p_3 be the corresponding point. Let $T(S_1, S_2, S_3)$ denote the set of lines simultaneously tangent to S_1 , S_2 , and S_3 . Now let $x_{h,2}$ be the infimum of all $0 \le x_h < x_{h,0}$ such that there exists a continuous function $\ell : (x_{h,2}, x_{h,1}) \to T(\{S_1, S_2, S_3\})$ with $\ell(x_h)$ lying on the hyperboloid with parameter x_h . Since the spheres are compact, the infimum is a minimum. If $x_{h,2} > 0$ then one of three hyperboloids involved by the three pairs of spheres must be one of the extreme hyperboloids in that situation and we are in cases (ii) (a), (b), or (c). If $x_{h,2} = 0$ then we distinguish between two possibilities. Either during this process we also reached a tangent to some other sphere $S(p_i, r)$ for some $4 \le i \le m$; in this case we are in case (i). Or during the transformation all the points p_4, \ldots, p_m are enclosed in the cylinder with axis ℓ and radius r, but none of them is contained in it. Then we arrive at situation (ii) (a).

3. Computing the smallest circumscribing cylinders of a simplex in \mathbb{E}^3

So far, we have seen how to reduce the computation of a smallest enclosing cylinder of a simplex in \mathbb{E}^3 to the computation of a smallest circumscribing cylinder. In order to apply algebraic methods to compute a smallest circumscribing cylinder, there are many different ways to formulate that problem in terms of polynomial equations. It is well-known that the computational costs of solving a system of polynomial equations are mainly dominated by the Bézout number (= product of the degrees) and the mixed volume (the latter one is discussed in Section 4). See [6, 7, 26] for comprehensive introductions and the state-of-the-art. Hence, it is an essential task to find the right formulations. Moreover, we are interested in simplex classes for which the degrees can be further reduced.

3.1. General simplices in \mathbb{E}^3 . In the proof of [9, Theorem 6], a polynomial formulation is given to compute the smallest enclosing cylinder of a simplex in \mathbb{E}^3 . This formulation describes the problem by three equations in the direction vector $v = (v_1, v_2, v_3)^T$ of the line, one of them normalizing the direction vector v by

(3.1)
$$v_1^2 + v_2^2 + v_3^2 = 1.$$

The equations are of degree 10, 3, and 2, respectively, thus giving a Bézout number of 60. However, as pointed out in that paper, some of the solutions to that system are artificially introduced by the formulation and occur with higher multiplicity, and there are only 18 really different solutions. Even more severely, in the experiments in that paper

(using SYNAPS, a state-of-the-art software for numerical polynomial computations), the numerical treatment of these multiple solutions needs much time, roughly a factor 100 compared to similar systems without multiple solutions.

Here, we present an approach, which reflects the true algebraic bound of 18. Namely, we give a polynomial formulation with Bézout bound 36 in which every solution generically has multiplicity one. The additional factor 2 just results from the fact that due to the normalization condition (3.1) every solution v also implies that -v is a solution as well.

Our framework is based on [21] in which the lines simultaneously tangent to four unit spheres are studied. A line in \mathbb{E}^3 is represented by a point $u \in \mathbb{E}^3$ lying on the line and a direction vector $v \in \mathbb{E}^3$ with $v^2 = 1$. We can make u unique by requiring that $u \cdot v = 0$. A line $\ell = (u, v)$ has Euclidean distance r from a point $p \in \mathbb{E}^3$ if and only if the quadratic equation $(u + tv - p)^2 = r^2$ has a solution of multiplicity two. This gives the condition

$$\frac{\left(v\cdot(u-p)\right)^2}{v^2} - (u-p)^2 + r^2 = 0.$$

Expanding this equation yields

(3.2)
$$v^2 u^2 - 2v^2 u \cdot p + v^2 p^2 - (v \cdot p)^2 - r^2 v^2 = 0.$$

Rather than using $v^2 = 1$ to further simplify this equation, we prefer to keep the homogenous form, in which all terms are of degree 4.

Now let p_1, \ldots, p_4 be the affinely independent vertices of the given simplex. Without loss of generality we can choose p_4 to be the located in the origin. Then the remaining points span \mathbb{E}^3 . Subtracting the equation for the point in the origin from the equations for p_1, p_2, p_3 gives the following program to compute the square of the radius of the minimal circumscribing cylinder.

(3.3)
$$\begin{array}{rcl} \min u^2 \\ \text{s.t.} & u \cdot v &= 0, \\ & 2v^2 u \cdot p_i &= v^2 p_i^2 - (v \cdot p_i)^2, \\ & v^2 &= 1. \end{array}$$

We remark that the set of admissible solutions is nonempty; a proof of that statement (for general dimension) is contained in Section 4.

Since the points p_1, p_2, p_3 are linearly independent, the matrix $M := (p_1, p_2, p_3)^T$ is invertible, and we can solve the equations in the bottom line of (3.3) for u:

(3.4)
$$u = \frac{1}{2v^2} M^{-1} \begin{pmatrix} v^2 p_1^2 - (v \cdot p_1)^2 \\ v^2 p_2^2 - (v \cdot p_2)^2 \\ v^2 p_3^2 - (v \cdot p_3)^2 \end{pmatrix}.$$

Now substitute this expression for u into the the objective function and into the first constraint of the system (3.3). After setting $v^2 = 1$ in the denominator of the first constraint, this gives a homogeneous cubic equation which we denote by $g_1(v_1, v_2, v_3) = 0$. Hence, we arrive at the following polynomial optimization formulation in terms of the variables v_1 , v_2 , and v_3 .

(3.5)
$$\min \left(\frac{1}{2} M^{-1} \begin{pmatrix} v^2 p_1^2 - (v \cdot p_1)^2 \\ v^2 p_2^2 - (v \cdot p_2)^2 \\ v^2 p_3^2 - (v \cdot p_3)^2 \end{pmatrix} \right)^2$$
s.t.
$$g_1(v_1, v_2, v_3) = 0,$$
$$g_2(v_1, v_2, v_3) := v^2 - 1 = 0.$$

Note that the objective function is a homogeneous polynomial of degree 4. We denote this polynomial by f.

Using Lagrange multipliers λ_1 and λ_2 , a necessary local optimality condition is

(3.6)
$$\operatorname{grad} f = \lambda_1 \operatorname{grad} g_1 + \lambda_2 \operatorname{grad} g_2$$

By thinking of an additional factor λ_0 before grad f and considering (3.6) as a system of linear equations in λ_0 , λ_1 , λ_2 , we see that if (3.6) is satisfied for some vector v then the determinant

$$(3.7) \qquad \det \begin{pmatrix} -\frac{\partial f}{\partial v_1} & \frac{\partial g_1}{\partial v_1} & \frac{\partial g_2}{\partial v_1} \\ -\frac{\partial f}{\partial v_2} & \frac{\partial g_2}{\partial v_2} & \frac{\partial g_2}{\partial v_2} \\ -\frac{\partial f}{\partial v_3} & \frac{\partial g_1}{\partial v_3} & \frac{\partial g_2}{\partial v_3} \end{pmatrix}$$

vanishes.

Lemma 3. (a) Any direction vector $(v_1, v_2, v_3)^T \in \mathbb{E}^3$ of the axis of a locally extreme circumscribing cylinder satisfies the polynomial system (3.5). If this system has only finitely many solutions then this number is bounded by 36.

(b) For a generic simplex the number of solutions is indeed finite, and all solutions have multiplicity one.

Proof. Let v be the direction vector of an axis of a locally extreme circumscribing cylinder. Then v satisfies the first constraint of (3.5), and the determinant (3.7) vanishes. Since these are homogeneous equations of degree 3 and 6, respectively, Bézout's Theorem implies that in connection with $v^2 = 1$ we obtain at most 36 isolated solutions.

For the second statement it suffices to check that for one specific simplex there are only finitely many solutions and that all solutions are pairwise distinct. \Box

3.2. Special simplex classes in \mathbb{E}^3 . In this section, we investigate conditions under which the degree of the resulting equations is reduced. Moreover, we show that for the equifacial simplex, the minimal circumscribing radius can be computed quite easily.

We use the following classification from [21, 22].

Proposition 4. Let T be a simplex in \mathbb{E}^3 with vertices p_1, \ldots, p_4 . The polynomial g_1 in the cubic equation factors into a linear polynomial and an irreducible quadratic polynomial if and only if the four faces of T can be partitioned into two pairs of faces $\{F_1, F_2\}, \{F_3, F_4\}$ with $\operatorname{area}(F_1) = \operatorname{area}(F_2) \neq \operatorname{area}(F_3) = \operatorname{area}(F_4)$. Moreover, g_1 factors into three linear terms if and only if the areas of all four faces of T are equal.

R. BRANDENBERG AND T. THEOBALD

First let us consider the case where g_1 decomposes into a linear polynomial and an irreducible quadratic polynomial. By optimizing separately over the linear and the quadratic constraint, the degrees of our equations are smaller than for the general case. Namely, analogously to the derivation in Section 3.1, for the quadratic constraint we obtain a Bézout bound of

$$(3+1+1) \cdot 2 \cdot 2 = 20,$$

and for the linear constraint we obtain

$$(3+0+1) \cdot 1 \cdot 2 = 8$$
.

Thus, we can conclude:

Lemma 5. If the four faces of the simplex can be partitioned into two pairs of faces $\{F_1, F_2\}, \{F_3, F_4\}$ with $area(F_1) = area(F_2) \neq area(F_3) = area(F_4)$ then there are at most 28 isolated local extrema for the minimal circumscribing cylinder. They can be computed from two polynomial systems with Bézout numbers 20 and 8, respectively.

Equifacial simplices. A simplex in \mathbb{E}^3 is called *equifacial* if all four faces have the same area. By Proposition 4, for an equifacial simplex the cubic polynomial g_1 factors into three linear terms. Hence, we obtain at most $3 \cdot 8 = 24$ local extrema. Somewhat surprisingly, using a characterization from [28], it is even possible to compute smallest circumscribing cylinder of an equifacial simplex esentially without any algebraic computation.

Namely, it is well-known that the vertices of an equifacial simplex T can be regarded as four pairwise non-adjacent vertices of a rectangular box (see, e.g., [19]). Hence, there exists a representation $p_1 = (w_1, w_2, w_3)^T$, $p_2 = (w_1, -w_2, -w_3)^T$, $p_3 = (-w_1, w_2, -w_3)^T$, $p_4 = (-w_1, -w_2, w_3)^T$ with $w_1, w_2, w_3 > 0$.

Assuming without loss of generality $v^2 = 1$, (3.2) gives

(3.8)
$$(v \cdot p_i)^2 + 2u \cdot p_i = \sum_{j=1}^3 w_j^2 + u^2 - r^2, \qquad 1 \le i \le 4.$$

Subtracting these equations pairwise gives

$$4(w_2u_2 + w_3u_3) = -4(w_1w_3v_1v_3 + w_1w_2v_1v_2)$$

(for indices 1, 2) and analogous equations, so that

$$w_1u_1 = -w_2w_3v_2v_3, \quad w_2u_2 = -w_1w_3v_1v_3, \quad w_3u_3 = -w_1w_2v_1v_2.$$

Since $u \cdot v = 0$, this yields $v_1 v_2 v_3 = 0$. Without loss of generality we can assume $v_1 = 0$. In this case,

$$u = \left(-\frac{w_2w_3}{w_1}v_2v_3, 0, 0\right)^T.$$

So we can express (3.8) in terms of the direction vector v,

$$w_2^2 v_2^2 + w_3^2 v_3^2 = \sum_{j=1}^3 w_j^2 + \left(-\frac{w_2 w_3}{w_1} v_2 v_3\right)^2 - r^2,$$

8

which, by using $v_2^2 + v_3^2 = 1$, gives

(3.9)
$$r^{2} = -\frac{w_{2}^{2}w_{3}^{2}}{w_{1}^{2}}v_{2}^{4} - \left(w_{2}^{2} - w_{3}^{2} - \frac{w_{2}^{2}w_{3}^{2}}{w_{1}^{2}}\right)v_{2}^{2} + w_{1}^{2} + w_{2}^{2} + w_{1}^{2} + w_{1}^{2} + w_{2}^{2} + w_{1}^{2} + w_{1}$$

Thus, by computing the derivative of this expression $r^2 = r^2(v_2)$ and taking into account the three cases $v_i = 0$, we can reduce the computation of the minimal circumscribing cylinder to solving three univariate equations of degree 3. However, we can still do better. Substitute $z_2 := v_2^2$, and let ρ be the expression for r^2 in terms of z_2 ,

$$\rho(z_2) = -\frac{w_2^2 w_3^2}{w_1^2} z_2^2 - \left(w_2^2 - w_3^2 - \frac{w_2^2 w_3^2}{w_1^2}\right) z_2 + w_1^2 + w_2^2.$$

Since the second derivative of that quadratic function is negative, $\rho(z_2)$ is a concave function. Hence, within the interval $z_2 \in [0, 1]$, the minimum is attained at one of the boundary values $z_2 \in \{0, 1\}$. Consequently, two of the components of $(v_1, v_2, v_3)^T$ must be zero and therefore v is perpendicular to two opposite edges. Since the latter geometric characterization is independent of our specific choice of coordinates, we can conclude:

Lemma 6. If all four faces of the simplex T have the same area then the axis of a minimum circumscribing cylinder is perpendicular to two opposite edges.

Hence, for an equifacial simplex it suffices to investigate the cross products of the three pairs of opposite edges (equipped with an orientation), and we do not need to solve a system of polynomial equations at all.

In order to illustrate how these three solutions relate to the 18 solutions of the general approach above, we consider the regular simplex in \mathbb{E}^3 . In the general approach, as already pointed out in [9], the six edge directions $p_i p_j$ $(1 \leq i < j \leq 4)$ all have multiplicity 1, and each of the three directions in Lemma 6, $p_1 p_2 \times p_3 p_4$, $p_1 p_3 \times p_2 p_4$, $p_1 p_4 \times p_2 p_3$, have multiplicity 4.

4. Smallest circumscribing cylinders in higher dimensions

In Section 3 we have given polynomial formulations with small Bézout number for computing smallest circumscribing cylinders of a simplex in \mathbb{E}^3 . Using the characterization in [25] of lines simultaneously tangent to 2n-2 spheres in \mathbb{E}^n , we generalize these formulations to smallest circumscribing cylinders of a simplex in \mathbb{E}^n , $n \ge 2$. Analogous to the three-dimensional case let p_1, \ldots, p_{n+1} be the affinely independent vertices of the simplex in \mathbb{E}^n , and let p_{n+1} be located in the origin.

First note that (3.3) also holds in general dimension n if we replace the index 3 by the index n. Since the points p_1, \ldots, p_n are linearly independent, the matrix $M := (p_1, \ldots, p_n)^T$ is invertible, and we can solve for u:

(4.1)
$$u = \frac{1}{2v^2} M^{-1} \begin{pmatrix} v^2 p_1^2 - (v \cdot p_1)^2 \\ \vdots \\ v^2 p_n^2 - (v \cdot p_n)^2 \end{pmatrix}.$$

Hence, by generalizing the formulation for the three-dimensional case, we obtain the program

(4.2)
$$\min \left(\frac{1}{2} M^{-1} \begin{pmatrix} v^2 p_1^2 - (v \cdot p_1)^2 \\ \vdots \\ v^2 p_n^2 - (v \cdot p_n)^2 \end{pmatrix} \right)^2$$
s.t.
$$g_1(v_1, \dots, v_n) = 0,$$
$$g_2(v_1, \dots, v_n) := v^2 - 1 = 0,$$

where g_1 denotes the cubic equation as before. In order to show that set of admissible solutions for our optimization problem is nonempty, we record the following result.

Lemma 7. For any simplex in \mathbb{E}^n the $\binom{n+1}{2}$ edge directions of the simplex are direction vectors of circumscribing cylinders.

Proof. Since the edge directions $p_i - p_j$ have a simple description in the basis p_1, \ldots, p_n , we express the cubic equation $g_1(v) = 0$ in that basis. Let v be an arbitrary direction vector, and let the representation of v in the basis p_1, \ldots, p_n be

$$v = \sum_{i=1}^n t_i p_i \, .$$

Further, let p'_1, \ldots, p'_n be a dual basis to p_1, \ldots, p_n ; i.e., let p'_1, \ldots, p'_n be defined by $p'_i \cdot p_j = \delta_{ij}$, where δ_{ij} denotes Kronecker's delta function. By elementary linear algebra, we have $t_i = p'_i \cdot v$.

When expressing u in this dual basis, $u = \sum u'_i p'_i$, the second constraint of (3.3) gives

$$u_i' = \frac{1}{2v^2} \left(v^2 p_i^2 - (v \cdot p_i)^2 \right) \,.$$

Substituting this representation of u into the equation $g_1(v) = 0$ gives

$$0 = g_1(v) = v^2(u \cdot v) = v^2 \left(\sum_{i=1}^n u'_i p'_i\right) \cdot v = v^2 \sum_{i=1}^n u'_i t_i,$$

where the last step uses the duality of the bases. Hence, we obtain the cubic equation

$$\frac{1}{2}\sum_{i=1}^{n}(v^{2}p_{i}^{2}-(v\cdot p_{i})^{2})t_{i} = 0.$$

Expressing v in terms of the *t*-variables yields

$$\frac{1}{2} \sum_{1 \le i \ne j \le n} \alpha_{ij} t_i^2 t_j + \sum_{1 \le i < j < k \le n} \beta_{ijk} t_i t_j t_k = 0,$$

10

where

$$\alpha_{ij} = (\operatorname{vol}_2(p_i, p_j))^2 = \det \begin{pmatrix} p_i \cdot p_i & p_i \cdot p_j \\ p_j \cdot p_i & p_j \cdot p_j \end{pmatrix},$$

$$\beta_{ijk} = \det \begin{pmatrix} p_i \cdot p_j & p_i \cdot p_k \\ p_k \cdot p_j & p_k \cdot p_k \end{pmatrix} + \det \begin{pmatrix} p_i \cdot p_k & p_i \cdot p_j \\ p_j \cdot p_k & p_j \cdot p_j \end{pmatrix},$$

$$+ \det \begin{pmatrix} p_j \cdot p_k & p_j \cdot p_i \\ p_i \cdot p_k & p_i \cdot p_i \end{pmatrix},$$

and $\operatorname{vol}_2(p_i, p_j)$ denotes the oriented area of the parallelogram spanned by p_i and p_j . In terms of the *t*-coordinates, the $\binom{n+1}{2}$ edges of the simplex are $t = e_i$, $1 \leq i \leq n$, and $t = e_i - e_j$, $1 \leq i < j \leq n$, where e_i denotes the *i*-th standard unit vector. For all these edges, the cubic equation is satisfied.

Considering Lagrange multipliers λ_1 and λ_2 yields the following necessary optimality condition.

(4.3)
$$\begin{array}{rcl} \operatorname{grad} f &=& \lambda_1 \operatorname{grad} g_1 + \lambda_2 \operatorname{grad} g_2 \,, \\ g_1(v_1, \dots, v_n) &=& 0 \,, \\ g_2(v_1, \dots, v_n) &=& 0 \,. \end{array}$$

Since the Bézout bound of this system is $3^n \cdot 3 \cdot 2 = 2 \cdot 3^{n+1}$, we have:

Lemma 8. For $n \ge 2$, the number of isolated local extrema for the minimal circumscribing cylinder is bounded by $2 \cdot 3^{n+1}$.

This bound is not tight. Trying to reduce this upper bound of isolated solutions like in the three-dimensional case, we can eliminate the linear occurrences of the Lagrange variables λ_1 and λ_2 . Generalizing (3.7), we have to consider the vanishing of all 3×3 subdeterminants of the matrix

(4.4)
$$\begin{pmatrix} -\frac{\partial f}{\partial v_1} & \frac{\partial g_1}{\partial v_1} & \frac{\partial g_2}{\partial v_1} \\ -\frac{\partial f}{\partial v_2} & \frac{\partial g_1}{\partial v_2} & \frac{\partial g_2}{\partial v_2} \\ \vdots & \vdots & \vdots \\ -\frac{\partial f}{\partial v_n} & \frac{\partial g_1}{\partial v_n} & \frac{\partial g_2}{\partial v_n} \end{pmatrix}.$$

Thus, for $n \ge 4$ we arrive at a non-complete intersection of equations where we have more equations than variables. Hence, we cannot apply our Bézout bound on these systems.

However, for small dimensions we can improve Lemma 8 by directly working on the formulation (4.3). In order to provide better bounds, we use well-known characterizations of the number of zeroes of a polynomial equation by the mixed volume of a Minkowski sum of polytopes (for an easily accessible introduction into this topic we refer to [7]). Here, let $\mathbb{C}^* := \mathbb{C} \setminus \{0\}$.

Lemma 9. For $2 \le n \le 7$, the number of solutions of the system (4.3) in $(v_1, \ldots, v_n, \lambda_1, \lambda_2) \in (\mathbb{C}^*)^{n+2}$ is bounded by

$$6\binom{n+1}{3},$$

where $\binom{n}{k}$ denotes the Stirling number of the second kind (see, e.g., [20, 24]).

The sequence $6\binom{n+1}{3}$ starts as follows

n	2	3	4	5	6	7
$6\binom{n+1}{3}$	6	36	150	540	1806	5796

Proof. For a polynomial $h = \sum_{\alpha \in \mathbb{N}_0^n} c_{\alpha} x^{\alpha} \in \mathbb{C}[x_1, \ldots, x_n]$, let

 $NP(h) := \operatorname{conv}\{\alpha \in \mathbb{N}_0^n : c_\alpha \neq 0\}$

denote the Newton polytope of h (see, e.g., $[7, \S7.1]$). Let h_1, \ldots, h_n be the polynomials of the gradient equation in (4.3). Further let $P_1, \ldots, P_n, Q_1, Q_2$ be the Newton polytopes of $h_1, \ldots, h_n, g_1, g_2$ for generic instances of these equations.

Recall that the mixed volume $MV(P_1, \ldots, P_n, Q_1, Q_2)$ is the coefficient of the monomial $\lambda_1 \cdot \lambda_2 \cdots \lambda_n \cdot \mu_1 \cdot \mu_2$ in the (n+2)-dimensional volume $Vol_{n+2}(\lambda_1 P_1 + \ldots + \lambda_n P_n + \mu_1 Q_1 + \mu_2 Q_2)$ (which is a polynomial expression in $\lambda_1, \ldots, \lambda_n, \mu_1, \mu_2$). By Bernstein's Theorem, the number of isolated common zeroes in $(\mathbb{C}^*)^{n+2}$ of the set of polynomials $h_1, \ldots, h_n, g_1, g_2$ is bounded aboved by

 $MV(P_1,\ldots,P_n,Q_1,Q_2)$

(see [7, Chapter 8, Theorem 5.4]). For every given n this volume can be computed using software for computing mixed volumes (see, e.g, [12, 29]).

We conjecture that for any $n \ge 2$, the number of isolated solutions in $(\mathbb{C}^*)^{n+2}$ is bounded by $6\binom{n+1}{3}$.

4.1. The regular simplex in \mathbb{E}^n . Here, we analyze the local extrema of circumscribing cylinders for the regular simplex. Our aim is both to illustrate the algebraic formulations given before and to relate our investigations to classical investigations on the regular simplex in convex geometry. In order to achieve many symmetries in the algebraic formulation, we use a slightly modified coordinate system that is particularly suited for the regular simplex; these coordinates have also been used in [30, 4].

The equation $x_1 + \ldots + x_{n+1} = 1$ defines an *n*-dimensional affine subspace in \mathbb{E}^{n+1} . Now let the regular simplex in this *n*-dimensional subspace be given by the n+1 vertices $p_i = e_i$, where e_i denotes the *i*-th standard unit vector, $1 \le i \le n+1$. We consider the tangency equation (3.2) for the point p_{n+1} ,

$$v^2 u^2 - 2v^2 u_{n+1} + v^2 - v_{n+1}^2 - r^2 v^2 = 0.$$

Subtracting this equation from the equation for p_i , $1 \le i \le n$, yields

$$2v^{2}(u_{i} - u_{n+1}) = -(v_{i}^{2} - v_{n+1}^{2}), \quad 1 \le i \le n.$$

Moreover, the embedding into the hyperplane $\sum_{i=1}^{n+1} x_i = 1$ implies $\sum_{i=1}^{n+1} u_i = 1$. In order to solve these n+1 equations for u, let M be the $(n+1) \times (n+1)$ -matrix whose *i*-th row

contains the vector $e_i^T - e_{n+1}^T$ and whose *n*-th row is (1, 1, ..., 1). Since *M* is invertible, we obtain

(4.5)
$$u = \frac{1}{2v^2} M^{-1} \begin{pmatrix} -(v_1^2 - v_{n+1}^2) \\ \vdots \\ -(v_n^2 - v_{n+1}^2) \\ 2v^2 \end{pmatrix}.$$

As before, substituting this expression into $u \cdot v = 0$ and setting $v^2 = 1$ in the denominator gives a cubic equation $g_1(v) = 0$. Hence, we obtain the following optimization problem. Here, the objective function f stems from the condition for the vertex p_{n+1} , and the condition $\sum_{i=1}^{n+1} v_i = 0$ comes from the embedding.

(4.6)
$$\min u^2 - 2u_{n+1} + 1 - v_{n+1}^2$$
$$\text{s.t.} \quad g_1(v_1, \dots, v_{n+1}) = 0,$$
$$\sum_{i=1}^{n+1} v_i = 0,$$
$$v^2 = 1.$$

First we record that the functions f and g_1 are symmetric polynomials in the variables v_1, \ldots, v_{n+1} . In order to show this, let $\sigma_1, \ldots, \sigma_{n+1}$ be the elementary symmetric functions in v_1, \ldots, v_{n+1} ,

$$\sigma_1 = v_1 + \ldots + v_{n+1},$$

$$\vdots$$

$$\sigma_k = \sum_{1 \le i_1 < \ldots < i_k \le n+1} v_{i_1} v_{i_2} \cdots v_{i_k},$$

$$\vdots$$

$$\sigma_{n+1} = v_1 v_2 \cdots v_{n+1}$$

(see, e.g., [6, 27]). By providing explicit expressions for f and g_1 as polynomials in the elementary symmetric polynomials $\sigma_1, \ldots, \sigma_{n+1}$, the symmetry of f and g_1 follows. More precisely, we obtain:

Lemma 10. The quartic polynomial $f(v_1, \ldots, v_{n+1})$ and the cubic polynomial $g_1(v_1, \ldots, v_{n+1})$ are symmetric polynomials in the variables v_1, \ldots, v_{n+1} . In terms of the elementary symmetric functions, f results in

$$f = \frac{1}{4(n+1)} \left(n\sigma_1^4 - 4n\sigma_1^2\sigma_2 + 2(n-1)\sigma_2^2 - 4\sigma_1^2 + 8\sigma_2 + 4n \right) + \sigma_1\sigma_3 - \sigma_4,$$

and the homogeneous polynomial g_1 results in

$$g_1 = \frac{1}{2(n+1)} \left(-(n-2)\sigma_1^3 + 3(n-1)\sigma_1\sigma_2 \right) - \frac{3}{2}\sigma_3.$$

Since $\sigma_1 = 0$ and $\sum_{i=1}^{n+1} v_i^2 = \sigma_1^2 - 2\sigma_2$, we can also deduce the following formulation of our optimization problem:

Corollary 11. Finding the critical values of the minimization problem (4.6) is equivalent to finding the critical values $(v_1, \ldots, v_{n+1})^T$ of the maximization problem

(4.7)
$$\begin{aligned} \max & \sigma_4 \\ \text{s.t.} & \sigma_1 &= 0, \\ \sigma_2 &= -\frac{1}{2} \\ \sigma_3 &= 0, \end{aligned}$$

where σ_i are the elementary symmetric functions in v_1, \ldots, v_{n+1} .

Theorem 12. The direction vector $(v_1, \ldots, v_{n+1})^T$ of any locally extreme circumscribing cylinder satisfies $|\{v_1, \ldots, v_{n+1}\}| \leq 3$, i.e., for each solution vector the components take at most three distinct values.

Proof. For $n \leq 2$, the statement is trivial, so we can assume $n \geq 3$. Let v be the direction vector of a locally extreme circumscribing cylinder with $v^2 = 1$. Using Corollary 11, let $f(v) := -\sigma_4(v), g_1(v) := \sigma_3(v), g_2(v) := \sigma_2(v) - 1/2$, and $g_3(v) := \sigma_1(v)$. As a necessary condition for a local extremum, for any pairwise different indices $a, b, c, d \in \{1, \ldots, n+1\}$ the determinant

(4.8)
$$\det \begin{pmatrix} -\frac{\partial f}{\partial v_a} & \frac{\partial g_1}{\partial v_a} & \frac{\partial g_2}{\partial v_a} & \frac{\partial g_3}{\partial v_a} \\ -\frac{\partial f}{\partial v_b} & \frac{\partial g_1}{\partial v_b} & \frac{\partial g_2}{\partial v_b} & \frac{\partial g_3}{\partial v_b} \\ -\frac{\partial f}{\partial v_c} & \frac{\partial g_1}{\partial v_c} & \frac{\partial g_2}{\partial v_c} & \frac{\partial g_3}{\partial v_c} \\ -\frac{\partial f}{\partial v_d} & \frac{\partial g_1}{\partial v_d} & \frac{\partial g_2}{\partial v_d} & \frac{\partial g_3}{\partial v_d} \end{pmatrix}$$

vanishes. Since f, g_1 , g_2 , and g_3 are symmetric functions in the variables v_1, \ldots, v_{n+1} , we can assume without loss of generality a = 1, b = 2, c = 3, and d = 4. Setting $\alpha_n := \sum_{i=5}^{n+1} v_i$ and $\beta_n = \sum_{i=5}^{n+1} v_i^2$, we can write

$$\frac{\partial g_3}{\partial v_i} = 1,$$

$$\frac{\partial g_2}{\partial v_i} = \sum_{\substack{j=1\\j\neq i}}^4 v_j + \alpha_n,$$

$$\frac{\partial g_1}{\partial v_i} = \sum_{\substack{1 \le j < k \le 4\\j,k \ne i}} v_j v_k + \alpha_n \sum_{\substack{j=1\\j\neq i}}^4 v_j + \frac{1}{2} \left(\alpha_n^2 - \beta_n\right)$$

 $(1 \leq i \leq 4)$. Moreover, since $\sigma_3(v) = 0$, we can consider $\sigma_3 + \frac{\partial f}{\partial v_i}$ instead of $\frac{\partial f}{\partial v_i}$. This allows to express the resulting expression easily in terms of α_n and β_n . More precisely, we

obtain

$$\sigma_3 + \frac{\partial f}{\partial v_i} = v_i \left(\sum_{\substack{1 \le j < k \le 4\\ j, k \ne i}} v_j v_k + \alpha_n \sum_{\substack{j=1\\ j \ne i}}^4 v_j + \frac{1}{2} (\alpha_n^2 - \beta_n) \right) \,.$$

Thus we can consider the determinant (4.8) as a polynomial in $v_1, v_2, v_3, v_4, \alpha_n, \beta_n$. Evaluating this 4×4 -determinant Δ shows that it is independent of α_n , β_n and that it factors as

$$\Delta = (v_1 - v_2)(v_1 - v_3)(v_1 - v_4)(v_2 - v_3)(v_2 - v_4)(v_3 - v_4).$$

Hence, $|\{v_1, v_2, v_3, v_4\}| \leq 3$, and this holds true for any quadruple (a, b, c, d) of indices. \Box

Using this result, we illustrate the occurrence of the Stirling numbers in Lemma 9 for the case of a regular simplex. There are $\binom{n+1}{3}$ ways to partition the set $V := \{v_1, \ldots, v_{n+1}\}$ into three nonempty subsets V_1, V_2, V_3 . We assume that $v_i \in V_i, 1 \le i \le 3$, and that all variables within the same set take the same value. Setting $k := |V_1|$ and $l := |V_2|$, the formulation in Corollary 11 yields the system of equations

(4.9)
$$kv_{1} + lv_{2} + (n+1-k-l)v_{3} = 0,$$
$$kv_{1}^{2} + lv_{2}^{2} + (n+1-k-l)v_{3}^{2} = 1,$$
$$\sum_{\substack{0 \le i_{1} \le i_{2} \le i_{3} \le 3\\i_{1}+i_{2}+i_{3}=3}} \binom{k}{i_{1}} \binom{l}{i_{2}} \binom{n+1-k-l}{i_{3}} v_{1}^{i_{1}}v_{2}^{i_{2}}v_{3}^{i_{3}} = 0.$$

If one of the indices k, l, or n + 1 - k - l is zero then this system consists of three equations in two variables, so we do not expect any solutions. For every choice of k, lcorresponding to a partition into nonempty subsets, we obtain a system of equations with Bézout number 6. Thus, whenever the values of v_1 , v_2 , and v_3 in the solutions to (4.9) are distinct, then this reflects the bound in Lemma 9.

In particular, in the case n = 4 we obtain the following 150 solutions.

$$k = 1, l = 1$$
: The six solutions for $(v_1, v_2, v_3)^T$ of the system (4.9) are

$$\left(\frac{1}{\sqrt{2}}, -\frac{1}{\sqrt{2}}, 0\right)^T$$
, $\left(\frac{1}{20}\sqrt{110 - 30i\sqrt{15}}, \frac{1}{20}\sqrt{110 + 30i\sqrt{15}}, -\frac{1}{10}\sqrt{15}\right)^T$,

and the solutions obtained by permuting the first two components of the first solution and by changing the signs and/or permuting the first two components in the second solution.

For the program (4.7) in the variables $(v_1, \ldots, v_5)^T$, this gives $\binom{5}{2}\binom{2}{1} = 20$ critical positions of the form (i.e., up to variable permutations)

$$\left(\frac{1}{\sqrt{2}}, -\frac{1}{\sqrt{2}}, 0, 0, 0\right)^T$$
,

20 complex solutions of the form

$$\left(-\frac{1}{20}\sqrt{110-30i\sqrt{15}},-\frac{1}{20}\sqrt{110+30i\sqrt{15}},\frac{1}{10}\sqrt{15},\frac{1}{10}\sqrt{15},\frac{1}{10}\sqrt{15}\right)^{T},$$

and 20 complex solutions of the form

$$\left(\frac{1}{20}\sqrt{110 - 30i\sqrt{15}}, \frac{1}{20}\sqrt{110 + 30i\sqrt{15}}, -\frac{1}{10}\sqrt{15}, -\frac{1}{10}\sqrt{15}, -\frac{1}{10}\sqrt{15}\right)^T.$$

k = 1, l = 2: Here, we obtain 30 solutions of the form

$$\left(0, \frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}\right)^T$$
,

30 solutions of the form

$$\left(\frac{1}{5}\sqrt{10}, \frac{1}{4}\sqrt{2} - \frac{1}{20}\sqrt{10}, \frac{1}{4}\sqrt{2} - \frac{1}{20}\sqrt{10}, -\frac{1}{4}\sqrt{2} - \frac{1}{20}\sqrt{10}, -\frac{1}{4}\sqrt{2} - \frac{1}{20}\sqrt{10}\right)^{T},$$

and 30 solutions of the form

$$\left(-\frac{1}{5}\sqrt{10},\frac{1}{4}\sqrt{2}+\frac{1}{20}\sqrt{10},\frac{1}{4}\sqrt{2}+\frac{1}{20}\sqrt{10},-\frac{1}{4}\sqrt{2}+\frac{1}{20}\sqrt{10},-\frac{1}{4}\sqrt{2}+\frac{1}{20}\sqrt{10}\right)^{T}.$$

The global minimum is attained for the vector $(0, \frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, -\frac{1}{2})^T$, and the objective value of the global optimum is 49/80. Hence, the radius of the smallest circumscribing cylinder for a regular simplex in \mathbb{E}^4 with edge length $\sqrt{2}$ is $\sqrt{49/80} = 7\sqrt{5}/20 \approx 0.7826$.

APPENDIX: AN ERROR IN THE RESULTS OF WEISSBACH

In the course of our investigations, we discovered a subtle but severe mistake in the paper [31] on the explicit determination of the outer (n-1)-radius of a regular simplex in \mathbb{E}^n . Since this error completely invalidates the proof given there², we give a description of that flaw, including some computer-algebraic calculations illustrating it.

In that paper, the computation of the outer (n-1)-radius of a regular simplex is reduced to the analysis of the following optimization problem.

(4.10)
$$\min \sum_{i=1}^{n+1} u_i^4$$
$$s.t. \qquad \sum_{i=1}^{n+1} u_i^2 = 1,$$
$$\sum_{i=1}^{n+1} u_i = 0.$$

²In a personal communication this has been confirmed by B. Weißbach.

For any local optimum $(u_1, \ldots, u_{n+1})^T$ there exist Lagrange multipliers $\lambda_1, \lambda_2 \in \mathbb{R}$ such that

,

(4.11)
$$4u_i^3 + 2\lambda_1 u_i + \lambda_2 = 0, \qquad 1 \le i \le n+1$$
$$\sum_{i=1}^{n+1} u_i^2 = 1,$$
$$\sum_{i=1}^{n+1} u_i = 0.$$

Erroneously, in [31] it is argued that symmetry arguments imply that $\lambda_2 = 0$ in any solution. The following calculation in the computer algebra system SINGULAR [13] shows that for n = 3 this system has 26 solutions (counting multiplicity) over \mathbb{C} .

```
ring R = 0, (u1,u2,u3,u4,la1,la2), (dp);
```

```
ideal I =
    4*u1^3 + 2*la1*u1 + la2,
    4*u2^3 + 2*la1*u2 + la2,
    4*u3^3 + 2*la1*u3 + la2,
    4*u4^3 + 2*la1*u4 + la2,
    u1^2 + u2^2 + u3^2 + u4^2 - 1,
    u1 + u2 + u3 + u4;
```

degree(std(I));

This program first defines a polynomial ring in the variables $u_1, \ldots, u_4, \lambda_1, \lambda_2$ over a field of characteristic zero. We then use the **degree** command to compute the dimension and the degree of the ideal defined by our equations. The output of that command is

// codimension = 6
// dimension = 0
// degree = 26

Hence, there are finitely many solutions (since the dimension of the ideal is zero), and the degree of the ideal (the sum of the multiplicities of the solutions) is 26.

18 of these solutions refer to the case $\lambda_2 = 0$ (and those were the ones computed in [31]). Namely, if $\lambda_2 = 0$ then the first row of (4.11) simplifies to

$$u_i(2u_i^2 + \lambda_1) = 0, \qquad 1 \le i \le n+1.$$

If we are only interested in the real solutions to this system, then setting $\lambda_1 = -2\lambda^2$ for some $\lambda \ge 0$ gives

$$u_i(u_i^2 - \lambda^2) = 0, \qquad 1 \le i \le n+1.$$

Since the vector $(u_1, \ldots, u_{n+1})^T = (0, \ldots, 0)^T$ does not satisfy the second row in (4.11), the solutions with $\lambda_2 = 0$ are

$$u_{i} = \lambda, \quad i \in \{i_{1}, \dots, i_{h}\}, u_{i} = -\lambda, \quad i \in \{i_{h+1}, \dots, i_{2h}\}, u_{i} = 0, \quad i \in \{1, \dots, n+1\} \setminus \{i_{1}, \dots, i_{2h}\}$$

for some $h \ge 1$, some set $\{i_1, \ldots, i_{2h}\}$ of pairwise different indices, and $\lambda = (2h)^{-1/2}$. In the case n = 3, there are 12 possibilities to choose the indices and the signs for |h| = 1 and 6 possibilities to choose the indices and the signs for |h| = 2, giving 18 solutions to (4.11).

However, there are 8 additional solutions, which in fact are also real! Namely, these are the solutions

$$(u_1, \dots, u_4)^T = \frac{1}{2\sqrt{3}} (1, -3, 1, 1)^T, \quad \lambda_1 = -\frac{7}{6}, \quad \lambda_2 = \frac{1}{\sqrt{3}}, \\ (u_1, \dots, u_4)^T = \frac{1}{2\sqrt{3}} (-1, 3, -1, -1)^T, \quad \lambda_1 = -\frac{7}{6}, \quad \lambda_2 = -\frac{1}{\sqrt{3}},$$

as well as the six distinct solutions obtained from these two by permuting the variables u_1, \ldots, u_4 . These additional solutions invalidate the subsequent arguments in [31].

The omissions get even worse in the higher-dimensional case. E.g., for n = 4, besides the $\binom{5}{2}\binom{2}{1} + \binom{5}{4}\binom{4}{2} = 20 + 30 = 50$ solutions described in [31], we obtain the following solutions:

$$(u_1, \dots, u_5)^T = \frac{1}{\sqrt{30}} (-2, -2, -2, 3, 3)^T, \quad \lambda_1 = -\frac{7}{15}, \quad \lambda_2 = -\frac{2}{75}\sqrt{30},$$

$$(u_1, \dots, u_5)^T = \frac{1}{\sqrt{30}} (2, 2, 2, -3, -3)^T, \quad \lambda_1 = -\frac{7}{15}, \quad \lambda_2 = \frac{2}{75}\sqrt{30},$$

$$(u_1, \dots, u_5)^T = \frac{1}{2\sqrt{5}} (1, -4, 1, 1, 1)^T, \quad \lambda_1 = -\frac{13}{10}, \quad \lambda_2 = \frac{6}{25}\sqrt{5},$$

$$(u_1, \dots, u_5)^T = \frac{1}{2\sqrt{5}} (-1, 4, -1, -1, -1)^T, \quad \lambda_1 = -\frac{13}{10}, \quad \lambda_2 = -\frac{6}{25}\sqrt{5},$$

as well as those solutions obtained by permuting the variables. Altogether, we have 10 + 10 + 5 + 5 = 30 solutions with $\lambda_2 \neq 0$, and thus a total number of 80 solutions.

Finally, we remark that the paper [30], which computes the outer (n-1)-radius of a regular simplex in *odd* dimension n, is correct (cf. also [4]).

References

- P.K. Agarwal, B. Aronov, and M Sharir. Line transversals of balls and smallest enclosing cylinders in three dimensions. *Discrete Comput. Geom.*, 21:373–388, 1999.
- [2] T. Bonnesen and W. Fenchel, Theorie der konvexen Körper. Springer-Verlag, Berlin, 1934.
- [3] R. Brandenberg. Radii of Convex Polytopes. Ph.D. thesis, Dept. of Mathematics, Technische Universität München, 2002.
- [4] R. Brandenberg. Radii of regular polytopes. Preprint, 2002.

- [5] T.M. Chan. Approximating the diameter, width, smallest enclosing cylinder, and minimimum-width annulus. To appear in *Internat. J. of Comp. Geom. and Applications*.
- [6] D. Cox, J. Little, and D. O'Shea. Ideals, Varieties, and Algorithms: An Introduction to Computational Algebraic Geometry and Commutative Algebra. UTM, Springer-Verlag, New York, 1996, second edition.
- [7] D. Cox, J. Little, and D. O'Shea. Using Algebraic Geometry. Graduate Texts in Mathematics, vol. 185, Springer-Verlag, New York, 1998.
- [8] H.S.M. Coxeter, Introduction to Geometry. John Wiley & Sons, 1961.
- [9] O. Devillers, B. Mourrain, F.P. Preparata, and Ph. Trébuchet. On circular cylinders by four or five points in space. Technical Report, INRIA, no. 4195, 2001.
- [10] G. Dos Reis, B. Mourrain, F. Rouillier, and Ph. Trébuchet. SYNAPS: An environment for symbolic and numeric computation, 2002. http://www-sop.inria.fr/galaad/logiciels/synaps/.
- [11] H.G. Eggleston. Notes on Minkowski geometry (I): Relations between the circumradius, diameter, inradius, and minimal width of a convex set. J. London Math. Soc. 33:76–81 (1958).
- [12] I. Emiris and J. Canny. Efficient incremental algorithms for the sparse resultant and the mixed volume. J. Symb. Comp. 20:117–149, 1995.
- [13] G.-M. Greuel, G. Pfister, and H. Schönemann. SINGULAR 2.0. A computer algebra system for polynomial computations. Centre for Computer Algebra, University of Kaiserslautern, 2001. http://www.singular.uni-kl.de.
- [14] P. Gritzmann and V. Klee. Inner and outer *j*-radii of convex bodies in finite-dimensional normed spaces. *Discrete Comput. Geom.* 7:255-280, 1992.
- [15] P. Gritzmann and V. Klee. Computational complexity of inner and outer *j*-radii of polytopes in finite-dimensional normed spaces. *Math. Program.* 59A:163-213, 1993.
- [16] P. Gritzmann and V. Klee. Computational convexity. In Handbook of Discrete and Computational Geometry (J.E. Goodman, J. O'Rourke, eds.), 491–515, CRC Press, Boca Raton, 1997.
- [17] S. Har-Peled and K. Varadarajan. Projective clustering in high dimensions using core sets. Proc. ACM Symposium on Computational Geometry '02 (Barcelona), 312–318, 2002.
- [18] D. Hilbert and S. Cohn-Vossen. Anschauliche Geometrie. Springer-Verlag, Berlin, 1932. Translation: Geometry and the Imagination, Chelsea Publ., New York, 1952.
- [19] Y.S. Kupitz and H. Martini. Equifacial tetrahedra and a famous location problem. Math. Gazette, 83:464–467, 1999.
- [20] R.L. Graham, D.E. Knuth, and O. Patashnik. *Concrete mathematics*. Addison-Wesley, 1989.
- [21] I.G. Macdonald, J. Pach, and T. Theobald. Common tangents to four unit balls in ℝ³. Discrete Comput. Geom. 26:1–17, 2001.
- [22] H. Schaal. Ein geometrisches Problem der metrischen Getriebesynthese. In Sitzungsber., Abt. II, Österr. Akad. Wiss. 194:39–53, 1985.
- [23] E. Schömer, J. Sellen, M. Teichmann, and C. Yap. Smallest enclosing cylinders. Algorithmica, 27:170– 186, 2000.
- [24] R.P. Stanley. Enumerative Combinatorics. Cambridge University Press, 1997.
- [25] F. Sottile and T. Theobald. Lines tangent to 2n-2 spheres in \mathbb{R}^n . Trans. Amer. Math. Soc. 354:4815–4829, 2002.
- [26] B. Sturmfels. Solving Systems of Polynomial Equations. CBMS series, vol. 97, AMS, Providence, 2002.
- [27] B. Sturmfels. Algorithms in Invariant Theory. RISC Series in Symbolic Computation, Springer-Verlag, Wien, 1993.
- [28] T. Theobald. Visibility computations: From discrete algorithms to real algebraic geometry. To appear in Proc. DIMACS workshop on Algorithmic and Quantitative Aspects of Real Algebraic Geometry, 2001, AMS DIMACS series.
- [29] J. Verschelde. PHCpack: A general-purpose solver for polynomial systems by homotopy continuation. ACM Trans. Math. Software 25:251–276, 1999.

- [30] B. Weißbach. Über die senkrechten Projektionen regulärer Simplexe. Beitr. Algebra Geom. 15:35–41, 1983.
- [31] B. Weißbach. Über Umkugeln von Projektionen regulärer Simplexe. Beitr. Algebra Geom. 16:127– 137, 1983.

René Brandenberg, Zentrum Mathematik, Technische Universität München, Boltzmannstr. 3, D–85747 Garching bei München

Current address: on leave at Technische Universität Wien, Institut für Analysis und Technische Mathematik, Wiedner Hauptstr. 8–10, A–1040 Wien

E-mail address: brandenb@mathematik.tu-muenchen.de *URL*: http://www-m9.mathematik.tu-muenchen.de/~brandenb/

THORSTEN THEOBALD, ZENTRUM MATHEMATIK, TECHNISCHE UNIVERSITÄT MÜNCHEN, BOLTZ-MANNSTR. 3, D-85747 GARCHING BEI MÜNCHEN

E-mail address: theobald@mathematik.tu-muenchen.de *URL*: http://www-m9.mathematik.tu-muenchen.de/~theobald/

20