



## Correction to: Fast random field generation with $H$ -matrices

Michael Feischl<sup>1</sup> · Frances Y. Kuo<sup>2</sup> · Ian H. Sloan<sup>2</sup>

Published online: 21 March 2019

© Springer-Verlag GmbH Germany, part of Springer Nature 2019

### Correction to: Numer. Math. (2018) 140:639–676

<https://doi.org/10.1007/s00211-018-0974-2>

On page 3 in line -4, the correct statement is

...  $\mu \in (0, \infty]$ ,  $p \in 2\mathbb{N}$  are parameters.

The corrected version states the parameter range as  $p \in 2\mathbb{N}$  instead of  $p \in \mathbb{N}$ . The effect is to disallow non-smooth norms such as the  $\ell_1$ -norm for the distance measure. This, however, does not impact the usual definition of the Matérn class of covariance functions using the  $\ell_2$ -norm for all values of  $\mu \in (0, \infty]$  (particularly the exponential case for  $\mu = 1/2$  and the Gaussian case for  $\mu = \infty$ ). Moreover, all the numerical examples and particularly the non-stationary covariance functions on page 642 are still covered by the theory.

This minor change closes the small gap in the proof of Lemma 1 by ensuring the existence of all the derivatives which are shown to be bounded in the sense of asymptotic smoothness (1).

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

---

The original article can be found online at <https://doi.org/10.1007/s00211-018-0974-2>.

✉ Michael Feischl  
michael.feischl@kit.edu

Frances Y. Kuo  
f.kuo@unsw.edu.au

Ian H. Sloan  
i.sloan@unsw.edu.au

<sup>1</sup> Karlsruhe Institute of Technology, 76131 Karlsruhe, Germany

<sup>2</sup> School of Mathematics and Statistics, UNSW Sydney, Sydney, NSW 2052, Australia