

Approximating node connectivity problems via set covers*

Guy Kortsarz

Zeev Nutov

Open University of Israel, Klausner 16 Str.

Open University of Israel, Klausner 16 Str.

Ramat-Aviv, Israel

Ramat-Aviv, 61392, Israel

`guyk@oumail.openu.ac.il`

`nutov@oumail.openu.ac.il`

Abstract

Given a graph (directed or undirected) with costs on the edges, and an integer k , we consider the problem of finding a k -node connected spanning subgraph of minimum cost. For the general instance of the problem (directed or undirected), there is a simple $2k$ -approximation algorithm. Better algorithms are known for various ranges of n, k . For undirected graphs with metric costs Khuller and Raghavachari gave a $\left(2 + \frac{2(k-1)}{n}\right)$ -approximation algorithm. We obtain the following results.

- (i) For arbitrary costs, a k -approximation algorithm for undirected graphs and a $(k + 1)$ -approximation algorithm for directed graphs.
- (ii) For metric costs, a $\left(2 + \frac{k-1}{n}\right)$ -approximation algorithm for undirected graphs and a $\left(2 + \frac{k}{n}\right)$ -approximation algorithm for directed graphs.

For undirected graphs and $k = 6, 7$, we further improve the approximation ratio from k to $\lceil (k + 1)/2 \rceil = 4$; previously, $\lceil (k + 1)/2 \rceil$ -approximation algorithms were known only for $k \leq 5$. We also give a fast 3-approximation algorithm for $k = 4$.

The multiroot problem generalizes the min-cost k -connected subgraph problem. In the multiroot problem, requirements k_u for every node u are given, and the aim is to find a minimum-cost subgraph that contains $\max\{k_u, k_v\}$ internally disjoint paths between every pair of nodes u, v . For the general instance of the problem, the best known algorithm has approximation ratio $2k$, where $k = \max k_u$. For metric costs there is a 3-approximation algorithm. We consider the case of metric costs, and, using our techniques, improve for $k \leq 7$ the approximation guarantee from 3 to $2 + \frac{\lceil (k-1)/2 \rceil}{k} < 2.5$.

*This research was done at Max-Planck-Institute für Informatik, Saarbrücken, Germany, and at the Open University of Israel, Ramat-Aviv, Israel.

1 Introduction

A basic problem in network design is given a graph \mathcal{G} to find its minimum cost subgraph that satisfies given connectivity requirements (see [14, 8] for surveys). A fundamental problem in this area is the *survivable network design problem*: find a cheapest spanning subgraph such that for every pair of nodes (u, v) , there are at least k_{uv} internally disjoint paths from u to v , where k_{uv} is a nonnegative integer (requirement) associated with the pair (u, v) ; two paths are *internally disjoint* if they do not have any internal node in common. No efficient approximation algorithm for this problem is known. However, for undirected graphs, when the paths are required only to be *edge disjoint*, an approximation algorithm that produces a solution at most twice the value of an optimal was given by Jain [12]. Henceforth, unless stated otherwise, we consider node connectivity only.

A ρ -*approximation algorithm* for a minimization problem is a polynomial time algorithm that produces a solution of value no more than ρ times the value of an optimal solution; ρ is called the *approximation ratio* of the algorithm. A particular important case of the survivable network design problem is the problem of finding a cheapest k -node connected spanning subgraph, that is the case when $k_{uv} = k$ for every node pair (u, v) . For undirected graphs this problem is NP-hard for $k = 2$ (for $k = 1$ it is the minimum spanning tree problem) and for directed graphs it is NP-hard for $k = 1$. For both directed and undirected graphs, there is a simple $2k$ -approximation algorithm, see for example [3].

For undirected graphs, the following results were known. Ravi and Williamson [21] claimed a $2H(k)$ -approximation algorithm, where $H(k) = 1 + \frac{1}{2} + \dots + \frac{1}{k}$, but the proof was found to contain an error, see [22]. A $\lceil (k+1)/2 \rceil$ -approximation algorithms are known for $k \leq 5$; see [16] for $k = 2$, [2] for $k = 2, 3$, and [7] for $k = 4, 5$. For metric costs and k arbitrary, Khuller and Raghavachari [16] gave a $(2 + \frac{2(k-1)}{n})$ -approximation algorithm (see also a 3-approximation algorithm in [3]).

We extend and generalize some of these algorithms, and unify ideas from [16], [2, 7], [3], and [13] to show further improvements. Among our results are:

- (i) For arbitrary costs, a k -approximation algorithm for undirected graphs and a $(k+1)$ -approximation algorithm for directed graphs;
- (ii) For metric costs, a $(2 + \frac{k-1}{n})$ -approximation algorithm for undirected graphs and a $(2 + \frac{k}{n})$ -approximation algorithm for directed graphs;

For undirected graphs and $k = 6, 7$, we further improve the approximation ratio from k to $\lceil (k+1)/2 \rceil = 4$, and give a fast 3-approximation algorithm for $k = 4$.

Recently, Cheriyan et. al. [4] gave a $6H(k)$ -approximation algorithm for undirected graphs with $n \geq 6k^2$, where n is the number of vertices of the input graph. In [5] the same authors suggest an iterative rounding $O(\frac{n}{\sqrt{n-k}})$ -approximation algorithm for both directed and undirected graphs. For a combinatorial $O(\sqrt{n} \ln k)$ -approximation algorithm see [15].

Another particular case of the survivable network design problem is the (undirected) *multiroot problem*, where pairwise node requirements are defined by single node requirements; that is, requirements k_u for every node u are given, and the aim is to find a minimum-cost subgraph that contains $\max\{k_u, k_v\}$ internally disjoint paths between every pair of nodes u, v . A graph (directed or undirected) is said to be *k-outconnected from a node r* if it contains k internally disjoint paths from r to any other node; such node r is usually referred as the *root*. It is easy to see that a subgraph is a feasible solution to the multiroot problem if and only if it is k_u -outconnected from every node u . Given an instance of the multiroot problem, we use q to denote the number of nodes u with $k_u > 0$, and $k = \max k_u$ is the maximum requirement. Observe that the (undirected) min-cost k -connected subgraph problem is a special case of the multiroot problem when $k_u = k$ for every node u .

Directed and undirected one root problems were considered long ago. For *directed* graphs, Frank and Tardos [10] showed that the problem of finding a k -outconnected spanning subgraph of minimum cost is solvable in polynomial time; a faster algorithm is due to Gabow [11]. As was observed by Khuller and Raghavachari in [16], this implies a 2-approximation algorithm for the (undirected) one root problem, as follows. First, replace every undirected edge e of G by the two antiparallel directed edges with the same ends and of the same cost as e . Then compute an optimal k -outconnected from r subdigraph and output its underlying (undirected) simple graph. The algorithm can be implemented in $O(k^2 n^2 m)$ time using the algorithm of [11].

For the multiroot problem, a $2q$ -approximation algorithm follows by applying the above algorithm for each root and taking the union of the resulting q subgraphs. The approximation guarantee $2q$ of this algorithm is tight for $q \leq k$, see [3]. For metric costs and k arbitrary, Cheriyan et. al. [3] gave a 3-approximation algorithm. For metric costs and $k = 2$, it can be shown that the problem is equivalent to that of finding a 2-connected subgraph. For the latter, there is a 3/2-approximation algorithm, see [9]. We consider the case of metric costs, and improve for $3 \leq k \leq 7$ the approximation ratio from 3 to $2 + \frac{\lfloor (k-1)/2 \rfloor}{k} < 2.5$.

This paper is organized as follows. Section 2 contains preliminary results and definitions. Sections 3 and 4 present algorithms for arbitrary and metric costs, respectively. Section 5 shows a 4-approximation algorithm for $k \in \{6, 7\}$, and Section 6 shows a fast 3-approximation algorithm for $k = 4$. Section 7 considers the metric multiroot problem with $k \leq 7$.

2 Definitions and preliminary results

All the graphs (directed or undirected) in the paper are assumed to be simple (i.e., without loops and parallel edges). An edge from u to v is denoted by uv . For an arbitrary graph H , $V(H)$ denotes the node set of H , and $E(H)$ denotes the edge set of H . Let $G = (V, E)$ be a graph. For any set of edges and nodes $U = E' \cup V'$ we denote by $G - U$ (resp., $G + U$) the graph obtained from G by deleting U (resp., adding U), where deletion of a node implies also deletion of all the edges incident to it. For a nonnegative cost function c on the edges of G and a subgraph $G' = (V', E')$ of G we use the notation $c(G') = c(E') = \sum\{c(e) : e \in E'\}$.

For $S, T \subseteq V$ let $\delta(S, T) = \delta_G(S, T)$ denote the set of edges in G going from S to T . For $X \subseteq V$ we denote by $\Gamma(X) = \Gamma_G(X)$ the set $\{v \in V \setminus X : uv \in E \text{ for some } u \in X\}$ of *neighbors* of X . Let $X^* = X_G^* = V \setminus (X \cup \Gamma(X))$ denote the “node complement” of X in G . It is well known that the function $|\Gamma(\cdot)|$ is submodular, that is for any $X, Y \subseteq V$ holds:

$$|\Gamma(X)| + |\Gamma(Y)| \geq |\Gamma(X \cap Y)| + |\Gamma(X \cup Y)|. \quad (1)$$

Two sets $X, Y \subset V$ *cross* (or X *crosses* Y) if $X \cap Y \neq \emptyset$ and neither $X \subseteq Y$ nor $Y \subseteq X$. We say that $U \subseteq V$ *covers* a collection \mathcal{C} of subsets of V if $X \cap U \neq \emptyset$ for every $X \in \mathcal{C}$.

We say that $X \subset V$ is *l -tight* if $|\Gamma(X)| = l$ and $X^* \neq \emptyset$ (i.e., if $|\Gamma(X)| = l$ and $|X| \leq |V| - l - 1$); such X is an *l -core* if it does not contain any other l -tight set. A graph G is *k -(node)-connected* if for any pair of its nodes there are k internally disjoint paths from one node to the other. By Menger’s Theorem, G is k -connected if and only if $|V(G)| \geq k + 1$ and there are no l -tight sets with $l \leq k - 1$ in G .

For an undirected graph G , we say that $U \subseteq V$ is an *l -cover* if U covers all the l' -cores with $l' \leq l$. Note that if U is an l -cover, then for any l' -tight set X with $l' \leq l$ holds: $X \cap U \neq \emptyset$ and $X^* \cap U \neq \emptyset$. Thus if $|V(G)| \geq l + 2$, then by adding to G the edge set $E' = \{uv : u \neq v \in U\}$ of a complete graph on U we obtain an $(l + 1)$ -connected graph.

An edge e of a graph G is said to be *critical w.r.t. property P* if G satisfies property P , but $G - e$ does not. The following theorem is due to Mader.

Theorem 2.1 (Mader,[17]) *In a k -connected undirected graph, any cycle in which every edge is critical w.r.t. k -connectivity contains a node of degree k .*

Theorem 2.1 implies that if $|\Gamma(v)| \geq k - 1$ for every $v \in V(G)$, and if F is an inclusion minimal edge set such that $G + F$ is k -connected, then F is a forest (if not, then F contains a cycle C of critical edges, but every node of this cycle is incident to 2 edges of C and to at least $k - 1$ edges of G , contradicting Mader’s Theorem). This implies:

Corollary 2.2 *Let U be a $(k - 1)$ -cover in an undirected graph G , and let $E' = \{uv : u \neq v \in U\}$. Then $G + E'$ is k -connected. Moreover, if $|\Gamma(v)| \geq k - 1$ for every $v \in V$, and if $F \subseteq E'$ is an inclusion minimal edge set such that $G + F$ is k -connected, then $|F| \leq |U| - 1$.*

The following property of k -outconnected undirected graphs is from [2].

Lemma 2.3 ([2]) *Let G be an undirected graph which is k -outconnected from r , and let S be an l -tight set in G . Then $|S \cap \Gamma(r)| \geq k - l + 1$, and if $l \leq k - 1$ then $|S \cap \Gamma(r)| \geq 2$ and $r \in \Gamma(S)$. Thus G is $(k - \lfloor \frac{|\Gamma(r)|}{2} \rfloor + 1)$ -connected.*

Corollary 2.4 *Let G be an undirected graph which is k -outconnected from r . Then $\Gamma(r) - v$ is a $(k - 1)$ -cover in G for any $v \in \Gamma(r)$.*

Throughout the paper, for an instance of a problem, we will denote by \mathcal{G} the input graph, and by opt the value of an optimal solution; n denotes the number of nodes in \mathcal{G} , and m the number of edges in \mathcal{G} . We assume that \mathcal{G} contains a feasible solution; otherwise our algorithms can be easily modified to output an error.

For the min-cost k -connected subgraph problem, we can assume that \mathcal{G} is a complete graph, and that $c(e) \leq opt$ for every edge e of \mathcal{G} . Indeed, let $G = (V, E)$ be a k -connected spanning subgraph of \mathcal{G} and let $st \in E$. Let F_{st} be the edge set of cheapest k internally disjoint paths from s to t in \mathcal{G} . Then $(G - st) + F_{st}$ is k -connected and, clearly, $c(F_{st}) \leq opt$. Note that F_{st} as above can be found in $O(n \log n(m + n \log n))$ time by a min-cost k -flow algorithm of [20] (the node version), and flow decomposition.

The main idea of most of our algorithms is to find a certain subgraph of \mathcal{G} of low cost and with a small cardinality $(k - 1)$ -cover or augmenting edge set. For undirected graphs, such a subgraph is found by using the following two modifications of the 2-approximation algorithm for the one root problem. Each one of these modifications outputs a subgraph of \mathcal{G} of cost $\leq 2opt$ (here opt is the optimal cost of a k -connected spanning subgraph of \mathcal{G}) and a $(k - 1)$ -cover U of the subgraph.

The first modification is from [16], and we use it for the case of metric costs. Let \mathcal{G}_r be a graph constructed from \mathcal{G} by adding an external node r and connecting it by edges of cost 0 to an arbitrary set R of at least k nodes in \mathcal{G} . We compute a k -outconnected from r subgraph G_r of \mathcal{G}_r using the 2-approximation algorithm above, and output $G = G_r - r$. As was shown in [16], $c(G) \leq 2opt$. By Lemma 2.3 R is a $(k - 1)$ -cover of G . We shall refer to this modification as the *External Outconnected Subgraph Algorithm* (EOCSA). It can be implemented in $O(k^2 n^2 m)$ time using the algorithm of [11].

The second modification is from [2, 7]. It finds a subgraph G and a node r such that: G is

k -outconnected from r , $|\Gamma_G(r)| = k$, and $c(G) \leq 2opt$. The time complexity of the algorithm is $O(k^2n^3m)$ for the deterministic version, and $O(k^2n^2m \log n)$ for the randomized one. We shall refer to the deterministic version as the *Outconnected Subgraph Algorithm* (OCSA), and for the randomized version as the *Randomized Outconnected Subgraph Algorithm* (ROCSA).

3 Min-cost k -connected subgraphs

3.1 Undirected graphs with arbitrary costs

This section deals with undirected graphs only. It is not hard to get a k -approximation algorithm for the min-cost k -connected subgraph problem as follows. We execute OCSA (or ROCSA) to compute a corresponding root r and a subgraph G of \mathcal{G} . Let $v \in \Gamma_G(r)$ be arbitrary, and let $R = \Gamma_G(r) \setminus v$. Recall that, by Corollary 2.4, R is a $(k - 1)$ -cover in G . We then find an edge set F as in Corollary 2.2, so $G + F$ is k -connected and F is a forest on R . Finally, we replace every edge $st \in F$ by a cheapest set F_{st} of k internally disjoint paths between s and t in \mathcal{G} . By [2], $c(G) \leq 2opt$. Since $|R| = k - 1$ then $|F| \leq k - 2$. Thus the cost of the output subgraph is at most $2opt + (k - 2)opt = kopt$.

We can get a slightly better approximation ratio by executing OCSA and then iteratively increasing the connectivity by 1 until it reaches k .

Let G be an l -connected graph, $|V(G)| \geq l + 2$. We say that an l -tight set X is *small* if $|X| \leq \lfloor \frac{n-l}{2} \rfloor$. Clearly, if X is l -tight, then at least one of X, X^* is small. Thus G is $(l + 1)$ -connected if and only if it has no small l -tight sets. The following lemma is well known, e.g., see [13, Lemma 1.2].

Lemma 3.1 *Let X, Y be two intersecting small l -tight sets in an l connected graph G . Then*

- (i) $X \cap Y$ is a small l -tight set;
- (ii) $X \cup Y, (X \cup Y)^*$ are both l -tight, and at least one of them is small.

Corollary 3.2 *In an l -connected graph G , no small l -tight set crosses a small l -core. Thus any two distinct small l -cores are disjoint.*

Let $\hat{\nu}_l(G)$ denote the number of small l -cores in G . Note that G is $(l + 1)$ -connected if and only if $\hat{\nu}_l(G) = 0$. Let us call an edge e *reducing* for G if $\hat{\nu}_l(G + e) \leq \hat{\nu}_l(G) - 1$.

Lemma 3.3 *Let R be a cover of all small l -cores in an l -connected graph G . If R is not an l -cover, then there is a reducing edge for G .*

Proof: Let R be a cover of all small l -cores in G . If R is not an l -cover, then there is an l -core T such that $T \cap R = \emptyset$. Note that T cannot be small, thus T^* is small. Let $S \subseteq T^*$ be an arbitrary l -core. Consider the collection \mathcal{D} of all (inclusion) maximal small l -tight sets containing S . Note that $T^* \in \mathcal{D}$. By Lemma 3.1 (ii) and the maximality of the sets in \mathcal{D} , exactly one of the following holds: (i) $|\mathcal{D}| = 1$ (so $\mathcal{D} = \{T^*\}$), or (ii) $|\mathcal{D}| \geq 2$, and the union of any two sets from \mathcal{D} is an l -tight set which is not small.

If case (i) holds, then any edge $e = st$ where $s \in S$ and $t \in T$ is reducing for G , since in $G + st$ there cannot be a small l -tight set containing S . Assume therefore that case (ii) holds. Let L be a set in \mathcal{D} crossing with T^* . Then, by Lemma 3.1 (i), $L^* \cap T$ is tight and small, implying $L^* \cap T \cap R \neq \emptyset$. This contradicts our assumption that $T \cap R = \emptyset$. \square

Corollary 3.4 *Any l -connected graph can be made $(l+1)$ -connected by adding $\hat{\nu}_l(G)$ edges.*

Proof: If G has no reducing edge, we find an l -cover R of size $\hat{\nu}_l(G)$ by picking a node from every small l -core. By Lemma 3.3, R is an l -cover, and, by Corollary 2.2, we can find a forest F on R such that $G + F$ is $(l+1)$ -connected. Else, we find and add a reducing edge, and recursively apply the same process on the resulting graph. \square

Theorem 3.5 *For the problem of making a $(k-1)$ -connected graph G k -connected by adding a min-cost edge set there exists a $(2 + \lfloor \frac{k}{2} \rfloor)$ -approximation algorithm with time complexity $O(k^2 n^3 m)$ deterministic (using OCSA) and $O(k^2 n^2 m \log n)$ randomized (using ROCSA).*

Proof: At the first phase we reset the edge cost of edges of G to zero, and execute OCSA: let H be the output graph, r the corresponding root, and $R = \Gamma_H(r)$. Now, consider the graph $J = H + G$, and let $l = k - 1$. Note that $\hat{\nu}_l(J) \leq \lfloor k/2 \rfloor$, since every l -tight set in H , and thus in J , contains at least two nodes from R , and $|R| = k$. At the second phase we make J k -connected by adding an edge set F as in Lemma 3.3, with $l = k - 1$. Now, $c(J) + c(F) \leq 2opt + \lfloor k/2 \rfloor opt$. The analysis of the time complexity is straightforward. \square

One can get an approximation ratio slightly better than k by sequentially applying augmentation steps as above. That is, we execute OCSA, and from $l = \lfloor k/2 \rfloor + 1$ to $k - 1$ increase the connectivity by 1. At every iteration, $\hat{\nu}_l(G) \leq \lfloor \frac{k}{k-l+1} \rfloor$, where G denotes the current graph. By Corollary 3.4, G can be made $(l+1)$ -connected by adding $\hat{\nu}_l(G)$ edges. The following lemma implies that increasing the number of internally disjoint paths between s and t from l to $l+1$ costs at most $\frac{opt}{k-l}$.

Lemma 3.6 *Let G be a subgraph of a graph \mathcal{G} containing l internally disjoint paths from s to t , $s, t \in V(\mathcal{G})$. For an integer p let $F^p \subseteq I = E(\mathcal{G}) - E(G)$ be an optimal edge set such that $G + F^p$ contains $l + p$ internally disjoint paths from s to t . Then $c(F^1) \leq \frac{1}{p}c(F^p)$.*

Proof: One can view \mathcal{G} as a min-cost flow network with source s and sink t where all edges and nodes have unit capacity (the costs are determined by the costs of the edges in I , while the edges in E have cost zero). Apply the following standard two stage reduction. First, replace every undirected edge e by two opposite directed edges with the same ends and the same capacity and cost as e , to get a directed network. Second, apply a standard conversion of node capacities to edge capacities: replace every node $v \in V - \{s, t\}$ by the two nodes v^-, v^+ connected by the edge v^-v^+ having the same capacity as v and cost zero, and redirect the heads of the edges entering v to v^- and the tails of the edges leaving v to v^+ .

In the new network, let \vec{F}^p be a min-cost $(l + p)$ -flow. Using flow decomposition, it is not hard to see that $c(\vec{F}^p) = c(F^p)$. In particular, $c(\vec{F}^0) = c(F^0) = 0$. Now consider the (fractional) $(l + 1)$ -flow $\frac{1}{p}\vec{F}^p + \left(1 - \frac{1}{p}\right)\vec{F}^0$ which has cost $\frac{1}{p}c(\vec{F}^p) = \frac{1}{p}c(F^p)$. Since the capacities are integral, there must be an integral $(l + 1)$ -flow \vec{F}^1 of at most the same cost, which proves the lemma. \square

Lemma 3.6 implies that the approximation ratio of our algorithm is:

$$I(k) = 2 + \sum_{l=\lceil \frac{k}{2} \rceil + 1}^{k-1} \left\lfloor \frac{k}{k-l+1} \right\rfloor \frac{1}{k-l} = 2 + \sum_{j=1}^{\lfloor \frac{k}{2} \rfloor - 1} \frac{1}{j} \left\lfloor \frac{k}{j+1} \right\rfloor.$$

It is easy to check that $I(k) < k$ for $k \geq 7$, but $\lim_{k \rightarrow \infty} \frac{I(k)}{k} = 1$.

Theorem 3.7 *For the problem of making a k_0 -connected graph k -connected there exists an $I(k - k_0)$ -approximation algorithm with time complexity $O(k^2 n^3 m)$ deterministic (using OCSA) and $O(k^2 n^2 m \log n)$ randomized (using ROCSA).*

3.2 Directed graphs with arbitrary costs

Let us say that a directed graph is k -inconnected to r if it contains k internally disjoint paths from any its node to r . Our algorithm is as follows.

1. Choose an arbitrary set $R = \{r_1, \dots, r_k\} \subseteq V$ of k nodes, and for $i = 1, \dots, k$, compute a min-cost k -outconnected from r_i subgraph $G_i = (V, F_i)$ of \mathcal{G} .
2. Construct a graph \mathcal{G}_r by adding to \mathcal{G} an external node r , and edges $r_i r$ of cost 0, $i = 1, \dots, k$.
Compute a minimum cost k -inconnected to r spanning subgraph G_r of \mathcal{G}_r .
3. Output $H = (G_r + F) - r$, where $F = \bigcup_{i=1}^k F_i$.

Theorem 3.8 *There exists a $(k+1)$ -approximation algorithm with time complexity $O(k^2n^2m)$ for the directed min-cost k -connected subgraph problem.*

Proof: We need to show that the output graph H is k -connected and that $c(H) \leq (k+1)opt$.

If H is not k -connected, then, H has an l -tight set S with $l < k$. Since H is k -outconnected from any node that belongs to R , we must have $S \cap R = \emptyset$. Thus, S is also l -tight in $G_r \cup F$. We obtain a contradiction since then G_r cannot contain k internally disjoint paths from any node $s \in S$ to r .

We now prove the approximation ratio. Clearly, $c(F_i) \leq opt$, $i = 1, \dots, k$; thus $c(F) \leq kopt$. It remains to show that $c(G_r) \leq opt$. Let G^* be an optimal k -connected spanning subgraph of \mathcal{G} . Extend G^* to a spanning subgraph G_r^* of \mathcal{G}_r by adding to G^* the node r and the edges $r_i r$ of cost 0, $i = 1, \dots, k$. It is easy to see that G_r^* is k -inconnected to r . Therefore, $c(G_r) \leq c(G_r^*) = c(G^*) = opt$. \square

4 Metric k -connected subgraph problem

4.1 Undirected graphs with metric costs

In this section we consider the metric min-cost k -connected subgraph problem. We present a modification of the $(2 + \frac{2(k-1)}{n})$ -approximation algorithm of Khuller and Raghavachari [16] to achieve a slightly better approximation guarantee of $(2 + \frac{k-1}{n})$.

Here is a short description of the algorithm of [16]. An l -star is a tree with l nodes and $l - 1$ leaves; a node s is a *center* of the star if all the other nodes in the star are leaves. Note that a min-cost subgraph of \mathcal{G} which is l -star with center v can be computed in $O(ln)$ time, and the overall cheapest l -star in $O(ln^2)$ time. The algorithm of [16] finds the node set R of a cheapest k -star, executes EOCSA, and adds to the graph G calculated the edge set E' as in Corollary 2.2 (that is, all the edges with both endnodes in R that are not in G). In [16] it is shown that $c(E') \leq \frac{2(k-1)}{n}$.

In our algorithm, we make a slightly different choice of R , and add an extra phase of removing from E' the noncritical edges (that is we add an edge set F as in Corollary 2.2). We show that for our choice of R , $c(F) \leq \frac{k-1}{n}$. We use the following lemma:

Lemma 4.1 *Let J be a complete graph on a node set R with node weights $w(v) \geq 0$, $v \in R$, and edge weights $w(uv) = w(u) + w(v)$, $u, v \in R$. If F is a forest on R then*

$$w(F) \leq (|R| - 2) \max\{w(v) : v \in R\} + \sum\{w(v) : v \in R\}.$$

Proof: Let $s \in R$ be a node satisfying $w(s) = \max\{w(v) : v \in R\}$. Among all forests F on R for which $w(F)$ is maximal, let F^* be one with the maximum number of edges incident to s . We claim that F^* is a star centered at s and thus for any forest F on R holds:

$$w(F) \leq w(F^*) = \sum\{w(s) + w(v) : v \in R - s\} = (|R| - 2)w(s) + \sum\{w(v) : v \in R\}.$$

If not, then there is a node $v \neq s$, such that v is either an isolated node of F^* , or v is a leaf of F^* with $uv \in F^*$ and $u \neq s$. In both cases, $(F^* - uv) + sv$ is a forest of the weight at least $c(F^*)$, but with more edges incident to s than F^* ; this contradicts our choice of F^* . \square

In our algorithm, we start by choosing the cheapest $(k + 1)$ -star J_{k+1} . Let v_0 be its center, and let its leaves be v_1, \dots, v_k . Denote $w_0 = w(v_0) = 0$ and $w_i = w(v_i) = c(v_0v_i)$, $i = 1, \dots, k$. Without loss of generality, assume that $w_1 \leq w_2 \leq \dots \leq w_k$. Since the costs are metric, $c(v_iv_j) \leq w(v_iv_j) = w_i + w_j$, $0 \leq i \neq j \leq k$. Let us delete v_k from the star. This results in a k -star J_k , and let R be its node set. For such R , let G be the subgraph of \mathcal{G} calculated by EOCSA. Recall that R is a $(k - 1)$ -cover in G . Let F be an edge set as in Corollary 2.2, so $G + F$ is k -connected, and F is a forest. The algorithm will output $G + F$. All this can be implemented in $O(k^2n^2m)$ time.

Let us analyze the approximation ratio. By [16], $c(H) \leq 2opt$. We claim that $c(F) \leq \frac{k-1}{n}opt$. Indeed, similarly to [16], using the metric cost assumption it is not hard to show that $c(J_{k+1}) = \sum\{w(v) : v \in R\} + w_k \leq \frac{2}{n}opt$. Thus, by our choice of J_k , $w_{k-1} = \max\{w(v) : v \in R\} \leq \frac{1}{n}opt$. Using this, the metric costs assumption, and Lemma 4.1 we get:

$$\begin{aligned} c(F) &= \sum\{c(v_iv_j) : v_iv_j \in F\} \leq \sum\{w_i + w_j : v_iv_j \in F\} \leq \\ &\leq (k - 2)w_{k-1} + \sum\{w(v) : v \in R\} \leq (k - 2)w_{k-1} + \left(\frac{2}{n}opt - w_k\right) \leq \\ &\leq (k - 3)w_{k-1} + \frac{2}{n}opt \leq \frac{k - 3}{n}opt + \frac{2}{n}opt = \frac{k - 1}{n}opt. \end{aligned}$$

Theorem 4.2 *There exists a $(2 + \frac{k-1}{n})$ -approximation algorithm with time complexity $O(k^2n^2m)$ for the undirected metric min-cost k -connected subgraph problem.*

4.2 Directed graphs with metric costs

In this section we consider directed graphs only. We say that a pair (R^-, R^+) is an l -cover in a directed graph G if R^- covers all the l' -tight sets in G and R^+ covers all the l' -tight sets in the graph obtained from G by reversal of its arcs, for any $l' \leq l$. It is easy to see that if (R^-, R^+) is a $(k - 1)$ -cover in G , and $E' = \{uv : u \in R^-, v \in R^+\}$, then $G + E'$ is k -connected.

A $v \rightarrow l$ -star is a directed tree rooted at v , with l nodes and $l - 1$ leaves; a $v \leftarrow l$ -star is a graph which reversal of its edges results in a $v \rightarrow l$ -star. Let v be a node of \mathcal{G} . Among all subdigraphs of \mathcal{G} which are $v \rightarrow l$ -stars (resp., $v \leftarrow l$ -stars), let $X_l^-(v)$ (resp., $X_l^+(v)$) be a cheapest one. Our algorithm for directed graphs is as follows:

1. Find a node v_0 for which $c(X_{k+1}^-(v)) + c(X_{k+1}^+(v))$ is minimal, and set $u_0 = v_0$.
 Let $R^- = \{v_1, \dots, v_k\}$ be the leaves of $J_{k+1}^- = X_{k+1}^-(v_0)$, and $R^+ = \{u_1, \dots, u_k\}$ be the leaves of $J_{k+1}^+ = X_{k+1}^+(u_0)$, where $c(v_0v_i) \leq c(v_0v_{i+1})$ and $c(u_iu_0) \leq c(u_{i+1}u_0)$, $i = 1, \dots, k - 1$.
 Set $J_k^- = X_{k+1}^-(v_0) - v_k$, $J_k^+ = X_{k+1}^+(v_0) - u_k$.
2. Add a node r to \mathcal{G} and edges $v_i r, r u_i$ of the cost 0, $i = 0, \dots, k - 1$, obtaining a graph \mathcal{G}_r . Compute two spanning subgraphs of \mathcal{G}_r : an optimal k -outconnected from r , say G_r^- , and an optimal k -inconnected to r , say G_r^+ ;
3. The graph $G + E'$ is k -connected, where $G = (G_r^- + G_r^+) - r$ and $E' = \{uv : u \in R^-, v \in R^+\}$.
 Output $H = G + F$, where $F \subseteq E'$, and all the edges in F are critical w.r.t. k -connectivity in H .

The following directed counterpart of Lemma 2.3 implies that the pair (R^-, R^+) is a $(k - 1)$ -cover in G , and thus the algorithm correctly outputs a k -connected graph H .

Lemma 4.3 *Let G_r be k -inconnected to r , let $R = \{v \in V : r \in \Gamma(v)\}$, and let S be an l -tight set in G_r such that $r \notin S$. If $r \in \Gamma(S)$ then $|S \cap R| \geq k - l + 1$, and if $r \notin \Gamma(S)$ then $l \geq k$. Thus R covers all the l -tight sets in $G_r - r$, $l \leq k - 1$.*

Proof: Let $s \in S$, and consider a set of k internally disjoint paths from s to r . Let $R' = \{v_1, \dots, v_k\} \subseteq R$ be the nodes of these paths preceding r . If $r \in \Gamma(S)$, then at most $l - 1$ nodes from R' may not belong to S ; this implies $|R \cap S| \geq |R' \cap S| \geq k - (l - 1)$. Clearly, if $r \notin \Gamma(S)$ and $l < k$ there cannot be k internally disjoint paths from s to r , by Menger's Theorem. The last statement is obvious. \square

Let us analyze the approximation ratio, using the notation as in the algorithm. Similarly to the proof of Theorem 3.8, one can show that $c(G) \leq c(G_r^-) + c(G_r^+) \leq 2opt$.

We claim that $c(F) \leq \frac{k}{n}opt$. Construct a bipartite graph $J = (A, B, E(J))$ with weights on the nodes as follows. The node parts are $A = \{u_0, \dots, u_{k-1}\}$ and $B = \{v_0, \dots, v_{k-1}\}$. The node weights are $w(u_i) = c(u_0u_i)$, $w(v_j) = c(v_0v_j)$ and $w(u_0) = w(v_0) = 0$. To every directed edge $e = u_i v_j$ with $u_i \in R^-, v_j \in R^+$ naturally corresponds an undirected edge $e' = u_i v_j$

with $u_i \in A, v_j \in B$. Moreover, since the costs are metric, for any $u_i \in R^-$ and $v_j \in R^+$ we have $c(u_i v_j) \leq w(v_i v_j) = w(u_i) + w(v_j)$.

We need some definitions and facts to continue. An even length sequence of edges $C = (v_1 v_2, v_3 v_2, v_3 v_4, \dots, v_{2q-1} v_{2q}, v_1 v_{2q})$ of a directed graph G is called an *alternating cycle*; the nodes $v_1, v_3, \dots, v_{2q-1}$ are *C-out nodes*, and v_2, v_4, \dots, v_{2q} are *C-in nodes*.

Theorem 4.4 (Mader,[18]) *In a k -connected directed graph, any cycle C in which every edge is critical w.r.t. k -connectivity contains a C -in node of indegree k , or a C -out node of outdegree k .*

Theorem 4.4 implies that if the indegree and the outdegree of every node in $V(G)$ is at least $k - 1$, and if F is an inclusion minimal edge set such that $G + F$ is k -connected, then F contains no alternating cycle. Note that $F \subseteq \{uv : u \in R^-, v \in R^+\}$ has no alternating cycle if and only if the corresponding edge set F' in J is a forest. We also need the following directed counterpart of Lemma 4.1 (the proof is omitted):

Lemma 4.5 *Let $J = (A, B, E(J))$ be a complete bipartite directed graph with nonnegative node weights $w(v) \geq 0, v \in A \cup B$, and edge weights $w(ab) = w(a) + w(b), a \in A, b \in B$. If $F \subseteq E(J)$ is a forest, then*

$$w(F) \leq (|B| - 1) \max\{w(a) : a \in A\} + (|A| - 1) \max\{w(b) : b \in B\} + \sum\{w(v) : v \in A \cup B\}.$$

Let us set $w_i = w(u_i) + w(v_i), i = 0, \dots, k$. Similarly to [16], one can show that $c(J_{k+1}^- + J_{k+1}^+) \leq \frac{2}{n} \text{opt}$. Thus, $w_{k-1} \leq \frac{1}{n} \text{opt}$, by our choice of J_k^-, J_k^+ . Now, similarly to the undirected case we get:

$$\begin{aligned} c(F) &= \sum\{c(v_i v_j) : v_i v_j \in F\} \leq \sum\{w_i + w_j : v_i v_j \in F\} \leq \\ &\leq (k-1)w_{k-1} + \sum_{i=0}^{k-1} w_i \leq (k-1)w_{k-1} + \left(\frac{2}{n} \text{opt} - w_k\right) \leq \\ &\leq (k-2)w_{k-1} + \frac{2}{n} \text{opt} \leq \frac{k-2}{n} \text{opt} + \frac{2}{n} \text{opt} = \frac{k}{n} \text{opt}. \end{aligned}$$

Theorem 4.6 *There exists a $\left(2 + \frac{k}{n}\right)$ -approximation algorithm with time complexity $O(k^2 n^2 m)$ for the directed metric min-cost k -connected subgraph problem.*

5 Min-cost 6,7-connected subgraphs

This section presents our algorithms for the min-cost 6,7-connected (undirected) subgraph problems. The algorithm itself is simple, and the main difficulty is to show that for $k = 6, 7$

we can make the output graph of OCSA k -connected by adding an edge set F with $|F| \leq 2$. A similar approach was used previously in [7] for $k = 4, 5$ with $|F| \leq 1$:

Lemma 5.1 ([7], Lemma 4.5) *Let G be a graph which is k -outconnected from r , $k \in \{4, 5\}$. If $|\Gamma_G(r)| = k$, then there exists a pair of nodes $s, t \in \Gamma_G(r)$ such that $G + st$ is k -connected.*

In fact, Lemma 5.1 can be deduced from Lemma 2.3 and the following lemma:

Lemma 5.2 ([13], Lemma 3.2) *Let G be an l -connected graph such that the maximum number of pairwise disjoint l -cores in G is exactly 2. Then the family of l -cores of G consists of two disjoint sets $S, T \subset V(G)$, and for any l -tight set Z of G either $S \subseteq Z$ and $T \subseteq Z^*$ or $T \subseteq Z$ and $S \subseteq Z^*$.*

Our algorithm for $k = 6, 7$ is based on the following Lemma:

Lemma 5.3 *Let G be k -outconnected from r , $k \in \{6, 7\}$. If $|\Gamma_G(r)| \in \{6, 7\}$ then there exists two pairs of nodes $\{s_1, t_1\}, \{s_2, t_2\} \subset \Gamma_G(R)$ such that $G + \{s_1t_1, s_2t_2\}$ is k -connected.*

Proof: Let G be as in the lemma, and $k \in \{6, 7\}$. In the proof, let the default subscript of the functions Γ be G . For convenience, let us denote $R = \Gamma(r)$. Note that, by Lemma 2.3, G is $(k-2)$ -connected, and that if S is $(k-2)$ -tight and X is $(k-1)$ -tight then $|S \cap R| \geq 3$, $|X \cap R| \geq 2$, and $r \in \Gamma(S) \cap \Gamma(X)$. In particular, since $|R| \leq 7$, we have:

Proposition 5.4 *If S and T are two disjoint $(k-2)$ -tight sets then any $(k-1)$ - or $(k-2)$ tight set intersects at least one of S, T .*

In what follows, note that in any graph $G = (V, E)$ for any two sets $X, Y \subset V$ holds:

$$|\Gamma(X)| + |\Gamma(Y)| \geq |\Gamma(X^* \cap Y)| + |\Gamma(X \cap Y^*)| \quad (2)$$

$$|\Gamma(X \cap Y)| \leq |\Gamma(X) - Y^*| + |\Gamma(Y) \cap X|. \quad (3)$$

We now establish several properties of $(k-1)$ - and $(k-2)$ -cores for a graph G as in Lemma 5.3 using inequalities (1), (2), and (3).

Lemma 5.5 *Let S be a $(k-2)$ -core and let X be an arbitrary $(k-1)$ -tight set crossing S . Then at least one of the following holds:*

- $X \cap S$ is $(k-1)$ -tight and $X^* \cap S^*$ is $(k-2)$ -tight; or
- $X \cap S^*$ is $(k-2)$ -tight and $X^* \cap S$ is $(k-1)$ -tight.

Proof: If $X^* \cap S^* = (X \cup S)^* \neq \emptyset$ then $|\Gamma(X \cup S)| \geq k - 2$. By the minimality of S , $|\Gamma(X \cap S)| \geq k - 1$. Using inequality (1) we obtain

$$(k - 1) + (k - 2) = |\Gamma(X)| + |\Gamma(S)| \geq |\Gamma(X \cap S)| + |\Gamma(X \cup S)| \geq (k - 1) + (k - 2).$$

If $X \cap S^*, X^* \cap S \neq \emptyset$ then $|\Gamma(X \cap S^*)| \geq k - 2$. By the minimality of S , $|\Gamma(X^* \cap S)| \geq k - 1$. Then using (2) we obtain

$$(k - 1) + (k - 2) = |\Gamma(X)| + |\Gamma(S)| \geq |\Gamma(X \cap S^*)| + |\Gamma(X^* \cap S)| \geq (k - 2) + (k - 1).$$

In both cases, equality holds everywhere, and the claim of the lemma holds.

Assume now that $X^* \cap S^* = \emptyset$. Then $X^* \cap S \neq \emptyset$, since otherwise X^* is a $(k - 1)$ -tight set disjoint to both S, S^* , contradicting Proposition 5.4. Thus we must have $X^* \cap S \neq \emptyset$ and $X \cap S^* = \emptyset$. Then

$$|\Gamma(X) - S^*| = |\Gamma(X)| - |S^*| \leq |\Gamma(X)| - |S^* \cap R| \leq (k - 1) - 3.$$

Since $|\Gamma(S)| = k - 2 \leq 5$, then $|\Gamma(S) \cap X| \leq 2$ or $|\Gamma(S) \cap X^*| \leq 2$. If $|\Gamma(S) \cap X| \leq 2$ then by (3)

$$|\Gamma(X \cap S)| \leq |\Gamma(X) - S^*| + |\Gamma(S) \cap X| \leq (k - 4) + 2 = k - 2.$$

This contradicts the minimality of S . The contradiction for the case $|X^* \cap \Gamma(S)| \leq 2$ is obtained similarly. \square

Combining the last lemma with Proposition 5.4 we obtain:

Corollary 5.6 *If G is not $(k - 1)$ -connected, then any $(k - 1)$ -core either contains exactly one $(k - 2)$ -core, or is contained in such a core.*

Lemma 5.7 *Let X, Y be $(k - 1)$ -cores that cross. Then exactly one of the following holds:*

- (i) *at least one of the sets $X \cap Y, X \cap Y^*, X^* \cap Y, X \cup Y$ is $(k - 2)$ -tight, or*
- (ii) *G is $(k - 1)$ -connected, $X \cap Y$ is k -tight, and the only $(k - 1)$ -cores in G are X, Y, X^*, Y^* .*

Proof: Assume $X^* \cap Y^* = (X \cup Y)^* \neq \emptyset$ (see Fig. 1(a)). Then $|\Gamma(X \cup Y)| \geq k - 2$, and, by the minimality of X , $|\Gamma(X \cap Y)| \neq k - 1$. Now, by (1):

$$|\Gamma(X \cap Y)| + |\Gamma(X \cup Y)| \leq |\Gamma(X)| + |\Gamma(Y)| = 2k - 2,$$

which implies that $|\Gamma(X \cap Y)| = k - 2$ or $|\Gamma(X \cup Y)| = k - 2$.

Figure 1: Illustration to the proof of Lemma 5.7

Similar argument applies with (2) for the case when both $X \cap Y^*$, $X^* \cap Y$ are nonempty and gives for this case that $|\Gamma(X \cap Y^*)| = k - 2$ or that $|\Gamma(X^* \cap Y)| = k - 2$.

Assume therefore that $X^* \cap Y^* = \emptyset$, and that at least one of $X^* \cap Y$, $X \cap Y^*$ is also empty. Without loss of generality let us consider the case $X \cap Y^* = \emptyset$ (see Fig. 1(b)). Then $Y^* \subset \Gamma(X)$. Since $|Y^* \cap R| \geq 2$, we must have $|\Gamma(X) - Y^*| \leq k - 3$.

Now, assume that $X \cap Y$ is not $(k - 2)$ -tight. Then, by the minimality of Y , we must have $|\Gamma(X \cap Y)| \geq k$. Applying inequality (3) we get

$$k \leq |\Gamma(X \cap Y)| \leq |\Gamma(X - Y^*)| + |\Gamma(Y) \cap X| \leq (k - 3) + |\Gamma(Y) \cap X|,$$

so $|\Gamma(Y) \cap X| \geq 3$. This implies

$$|\Gamma(Y) \cap X^*| = (k - 1) - |\Gamma(Y) \cap X| - |\Gamma(Y) \cap \Gamma(X)| \leq (k - 1) - 3 - 1 \leq 2.$$

Now, if $X^* \cap Y$ is not $(k - 2)$ -tight, then $X^* \cap Y = \emptyset$. Otherwise, applying (3) on X^* and Y we get a contradiction to the minimality of Y :

$$|\Gamma(X^* \cap Y)| \leq |\Gamma(X^*) - Y^*| + |\Gamma(Y) \cap X^*| \leq |\Gamma(X) - Y^*| + |\Gamma(Y) \cap X^*| \leq (k - 3) + 2.$$

From the previous discussion we conclude, that if the first case of the lemma does not hold, then the following holds (see fig 1(c)): all the three sets $X \cap Y^*$, $X^* \cap Y$, $X^* \cap Y^*$ are empty; $|X^*| = |Y^*| = 2$, and thus $X^*, Y^* \subseteq R$ and X^*, Y^* are $(k - 1)$ -cores; and $|\Gamma(Y) \cap X| = |\Gamma(X) \cap Y| = 3$ and thus $|X| \geq 4$ and $|Y| \geq 4$. (Note that then also $k = 7$ and $\Gamma(Y) \cap \Gamma(X) = \{r\}$.) From that it is easy to see that $\Gamma(X^* \cup Y^*) = \Gamma(X \cap Y)$, so $X^* \cup Y^*$ is k -tight. We now prove that then the second case of the lemma must hold.

First, let us show that G is $(k - 1)$ -connected. If not, then by Corollary 5.6, there is a $(k - 2)$ -core S containing X^* . Using Lemma 5.5 and Proposition 5.4, it is not hard to see that we must have $S = X^* \cup Y^*$. This is a contradiction, since $|\Gamma(X^* \cup Y^*)| = k$.

Second, we prove that if Z is a $(k-1)$ -core in G then Z is one of X, Y, X^*, Y^* . Otherwise, Z crosses at least one of X, Y, X^*, Y^* . Since G is $(k-1)$ -connected, case (i) of the lemma does not hold, and we conclude that $|Z^*| = 2$. But then Z^* crosses at least one of X, Y, X^*, Y^* , and, by the previous discussion, we must have $|Z^*| \geq 4$, which is a contradiction. \square

We now are ready to finish the proof of Lemma 5.3.

Assume first that G is $(k-1)$ -connected. We will show that then there is a $(k-1)$ -cover $U \subset R$ with $|U| \leq 3$. Then, the statement is a straightforward consequence from Corollary 2.2. Recall that the maximum number of pairwise disjoint cores in G is at most 3. Thus, if no two $(k-1)$ -cores cross, then picking one node in R from every $(k-1)$ -core gives a $(k-1)$ -cover as required. If there exists a pair X, Y of $(k-1)$ -cores that cross, then we are in case (ii) of Lemma 5.7. In particular, $X \cap Y$ is k -tight, thus by Lemma 2.3 $X \cap Y \cap R \neq \emptyset$. Then $U = \{x, y, z\}$, where $x \in X^* \cap R$, $y \in Y^* \cap R$, and $z \in X \cap Y \cap R$ is a $(k-1)$ -cover as required.

Assume now that G is not $(k-1)$ -connected. Let S, T be the $(k-2)$ -cores in G (as in Lemma 5.2). Let \mathcal{S} (resp., \mathcal{T}) denote all the $(k-1)$ -cores contained in S (resp., in T). Note that there are at most two disjoint sets in \mathcal{S} , and that, by Lemma 5.7, for any two sets in \mathcal{S} that cross, their union is S . A similar statement holds for \mathcal{T} .

Lemma 5.8 *Let \mathcal{C} be a collection of subsets of S containing at most two disjoint subsets, and let U cover \mathcal{C} . If for any $X, Y \in \mathcal{C}$ that cross holds $X \cup Y = S$, then there is $U' \subseteq U$ with $|U'| \leq 2$ that covers \mathcal{C} .*

Proof: It is sufficient to prove the statement under the assumption that any two sets in \mathcal{C} either disjoint or cross. The proof is by induction on $|\mathcal{C}|$. For $|\mathcal{C}| \leq 3$ the statement is clear.

Assume now that $|\mathcal{C}| \geq 4$. Let $X_1, X_2, X_3 \in \mathcal{C}$ be arbitrary. Then any two of X_1, X_2, X_3 cross. Let $Z = X_1 \cap X_2 \cap X_3$, and let $X \in \mathcal{C} \setminus \{X_1, X_2, X_3\}$. By the assumption of the lemma, $(X_i \cap X_j) \setminus Z \subset X$ for $i \neq j = 1, 2, 3$, implying $S \setminus Z \subseteq X$. Now, if $U \setminus Z \neq \emptyset$, let $u \in U \setminus Z$. Then u covers all the sets in \mathcal{C} except of exactly one of X_1, X_2, X_3 . Let $v \in U$ be a node that covers the set not covered by u . Then $\{u, v\}$ is a cover as required. If $U \subseteq Z$, then let $\mathcal{C}' = \mathcal{C} \setminus \{X_1, X_2, X_3\}$. Note that \mathcal{C}' satisfies the conditions of the lemma. By the induction hypothesis, \mathcal{C}' has a cover U' as in the lemma. But then U' also covers \mathcal{C} , and the proof is complete. \square

By Lemma 5.8, there is a pair $\{s_1, s_2\} \in R$ that covers \mathcal{S} , and there is a pair $\{t_1, t_2\} \in R$ that covers \mathcal{T} .

Lemma 5.9 *The graph $G + \{s_1 t_1, s_2 t_2\}$ is k -connected.*

Proof: It is straightforward to see (via Lemma 5.2) that adding the edges s_1t_1, s_2t_2 adds at least 2 neighbors to any $(k - 2)$ -tight set. We will show that adding these edges also adds at least 1 neighbor to any $(k - 1)$ -tight set Z . If Z contains one of S, T and Z^* contains the other, then the claim is straightforward. Else, by Corollary 5.6, Z or Z^* is contained in one of S, T , say $Z \subset S$. Then $T \subset Z^*$, and the claim again follows. \square

The proof of Lemma 5.3 is done. \square

Two pairs $\{s_1, t_1\}, \{s_2, t_2\}$ as in Lemma 5.3 can be found in $O(m)$ time, e.g., by exhaustive search. Combining this and Lemma 5.3 we obtain:

Theorem 5.10 *For $k = 6, 7$, there exists a 4-approximation algorithm for the min-cost k -connected subgraph problem. The time complexity of the algorithm is $O(n^3m)$ deterministic (using OCSA) and $O(n^2m \log n)$ randomized (using ROCSA).*

6 Fast algorithm for $k = 4$

In this section we present a 3-approximation algorithm for $k = 4$ with complexity $O(n^4)$. This improves the previously best known time complexity $O(n^5)$ [7]. Let us call a subset R of nodes of a graph G k -connected if for every $u, v \in R$ there are k internally disjoint paths between u and v in G . The following Theorem is due to Mader.

Theorem 6.1 ([19]) *Any graph on $n \geq 5$ nodes with minimal degree at least k , $k \geq 2$, contains a k -connected subset R with $|R| = 4$.*

It is known that the problem of finding a min-cost spanning subgraph with minimal degree at least k is reduced to the weighted b -matching problem. Using the algorithm of Anstee [1] for the latter problem, such a subgraph can be found in $O(n^2m)$ time. We use these observations to obtain a 3-approximation algorithm for $k = 4$ as follows. The algorithm has two phases. At phase 1, among the subgraphs of \mathcal{G} with minimal degree 4, we find an optimal one, say G . Then, we find in G a 4-connected subset R with $|R| = 4$. At phase 2, we execute EOCSA on R , and let F be its output. Finally, the algorithm will output $G + F$.

Theorem 6.2 *There exists a 3-approximation algorithm for the min-cost 4-connected subgraph problem, with time complexity $O(n^2m + nT(n)) = O(n^4)$, where $T(n)$ is the time required for multiplying two $n \times n$ -matrices.*

Proof: The correctness follows from Theorem 6.1, Lemma 2.3 (i), and Corollary 2.2. To see the approximation ratio, recall that $c(F) \leq 2opt$, and note that $c(G) \leq opt$.

We now prove the time complexity. The complexity of each step, except of finding a 4-connected subset in G is $O(n^2m)$. Let us show that finding a 4-connected subset can be done in $O(n^2m + n(T(n)))$ time. Using the Ford-Fulkerson max-flow algorithm, we construct in $O(n^2m)$ time the graph $J = (V, E')$, where $(s, t) \in E'$ if and only if there are 4 internally disjoint paths between s and t in G . Now, R is a 4-connected subset in G if and only if the subgraph induced by R in J is a complete graph. Thus, finding R as above is reduced to finding a complete subgraph on 4 nodes in J . This can be implemented as follows. Observe that $R = \{s, u, v, w\}$ induces a complete subgraph in J if and only if $\{u, v, w\}$ form a triangle in the subgraph induced by $\Gamma_J(s)$ in J . It is known that finding a triangle in a graph is reduced to computing the square of the incidence matrix of the graph. The best known time bound for that is $O(n^{2.376})$ [6], and the time complexity follows. \square

7 Metric multiroot problem: cases $k \leq 7$

In this section we consider the metric-cost multiroot problem. Note that here \mathcal{G} is a complete graph, and every edge in \mathcal{G} has cost at most opt/k . This is since any feasible solution contains at least k edge disjoint paths between any two nodes s and t , and, by the metric cost assumption, each one of these paths has cost $\geq c(st)$. For $k \leq 7$, we give an algorithm with approximation ratio $2 + \frac{\lfloor (k-1)/2 \rfloor}{k} < 2.5$. This improves the previously best known approximation ratio 3 [3]. Our algorithm combines some ideas from [3], [2, 7], and some results from the previous section.

Splitting off two edges ru, rv means deleting ru and rv and adding a new edge uv .

Theorem 7.1 ([3], Theorem 17) *Let $G = (V, E)$ be a graph which is k -outconnected from a root node $r \in V$, and suppose that $|\Gamma_G(r)| \geq k + 2$ and every edge incident to r is critical w.r.t. k -outconnectivity from r . If G is not k -connected, then there exists a pair of edges incident to r that can be split off preserving k -outconnectivity from r .*

Consider now an instance of a metric cost multiroot problem, and let r be a node with the maximum requirement k . As was pointed in [3], Theorem 7.1 implies that we can produce a spanning subgraph G of \mathcal{G} , such that G is k -outconnected from r , $c(G) \leq 2opt$, and: G is k -connected, or $|\Gamma_G(r)| \in \{k, k + 1\}$. To handle the cases $k = 5, 7$, we show that by adding one edge, we can reduce the case $|\Gamma(r)| = k + 1$ to the already familiar case $|\Gamma(r)| = k$.

Lemma 7.2 *Let $G = (V, E)$ be k -outconnected from a root node $r \in V$, let $R = \Gamma_G(r)$, and let rx be critical w.r.t. k -outconnectivity from r . If $|R| \geq k + 1$, then there exists a node $y \in R$ such that $(G - rx) + xy$ is k -outconnected from r .*

Proof: Let $G = (V, E)$ be a graph which is k -outconnected from a root node $r \in V$. Following [3], for $X \subseteq V - r$ let $g(X) = |\Gamma_{G-r}(X)| + |X \cap R|$. It is easy to see that G is k -outconnected from r if and only if $g(X) \geq k$ for every $X \subseteq V - r$. Let us say that a set $X \subseteq V - r$ is *critical* if $g(X) = k$. Thus, rx is critical w.r.t. k -outconnectivity from r if and only if there is a critical set containing x . In [3, Lemma 6] was shown that:

The intersection and union of two intersecting critical sets are both critical. Thus for every critical edge rx there is unique maximal critical set containing x .

Now, assume that rx is critical w.r.t. k -outconnectivity from r , and let X be the maximal critical set containing x . We claim that if $R \cap X^* \neq \emptyset$ then for any $y \in R \cap X^*$ holds: $(G - rx) + xy$ is k -outconnected from r . Indeed, if $(G - rx) + xy$ is not k -outconnected from r , then there is a critical set X' with $x \in X'$, $y \in \Gamma(X')$. But then we must have $X' \subseteq X$. As a consequence, we must have $y \in X + \Gamma(X)$, contradicting that $y \in X^*$.

Now, suppose $|R| \geq k + 1$. We claim that then $R \cap X^* \neq \emptyset$. Else, $R \subseteq X \cup \Gamma(X)$. But then we must have $g(X) \geq |R| \geq k + 1$, contradicting that $g(X) = k$. \square

Lemma 7.3 *Let G be a graph which is k -outconnected from r , $3 \leq k \leq 7$, and suppose that $|\Gamma_G(r)| \in \{k, k + 1\}$. Then there is an edge set $F \subseteq \{uv : u \neq v \in \Gamma_G(r)\}$ such that: $G + F$ is k -connected and $|F| \leq \lfloor (k - 1)/2 \rfloor$.*

Proof: For $k \leq 4$, this is a straightforward consequence from Lemmas 2.3 and 5.2. For $k = 6$ this is a consequence from Lemma 5.3. For $k = 5, 7$, it can be easily deduced using Lemma 7.2 and: Lemma 5.1 for $k = 5$, or Lemma 5.3 for $k = 7$. \square

Using Lemma 7.3 and the fact that for every $s, t \in V$ holds $c(st) \leq \text{opt}/k$, we deduce:

Theorem 7.4 *For the metric cost multiroot problem with $3 \leq k \leq 7$, there exists a $(2 + \frac{\lfloor (k-1)/2 \rfloor}{k})$ -approximation algorithm with time complexity $O(n^3m)$.*

References

- [1] R. P. Anstee: A polynomial time algorithm for b -matchings: an alternative approach, *Information Processing Letters* 24, 1987, 153–157.
- [2] V. Auletta, Y. Dinitz, Z. Nutov, and D. Parente, A 2-approximation algorithm for finding an optimum 3-vertex connected spanning subgraph, *Journal of Algorithms* 32, 1999, 21–30.

- [3] J. Cheriyan, T. Jordán, and Z. Nutov, On rooted node-connectivity problems, *Algorithmica* 30 (special issue on APPROX'98), 2001, 353–375.
- [4] J. Cheriyan, S. Vempala, and A. Vetta, An approximation algorithm for the minimum cost k -vertex connected subgraph, manuscript, July 2001.
- [5] J. Cheriyan, S. Vempala, and A. Vetta, Network design via iterative rounding of setpair relaxations, manuscript, October 2001.
- [6] D. Coppersmith and S. Winograd: Matrix multiplication via arithmetic progressions, *J. Symbolic Comp.* 9, 1990, 251–280.
- [7] Y. Dinitz and Z. Nutov, A 3-approximation algorithm for finding optimum 4,5-vertex-connected spanning subgraphs, *Journal of Algorithms* 32, 1999, 31–40.
- [8] A. Frank, Connectivity augmentation problems in network design, *Mathematical Programming*, State of the Art, J. R. Birge and K. G. Murty eds., 1994, 34–63.
- [9] G. .N. Frederickson and J. Jájá, On the relationship between the biconnectivity augmentation and traveling salesman problems, *Theoretical Computer Science* 19(2), 1982, 189–201.
- [10] A. Frank and É. Tardos, An application of submodular flows, *Linear Algebra and its Applications* 114/115, 1989, 329–348.
- [11] H. N. Gabow, A representation for crossing set families with application to submodular flow problems, *Proc. 4th Annual ACM-SIAM Symp. on Discrete Algorithms* 1993, 202–211.
- [12] K. Jain, A factor 2 approximation algorithm for the generalized Steiner network problem, *Combinatorica*, 21(1), 2001, 39–60.
- [13] T. Jordán, On the optimal vertex-connectivity augmentation, *J. Combinatorial Theory Series B* 63, 1995, 8–20.
- [14] S. Khuller, Approximation algorithms for finding highly connected subgraphs, In *Approximation algorithms for NP-hard problems*, Ed. D. S. Hochbaum, PWS publishing co., Boston, 1996.
- [15] G. Kortzars and Z. Nutov, Combinatorial approximation algorithm for k -node connected subgraphs, manuscript.

- [16] S. Khuller and B. Raghavachari, Improved approximation algorithms for uniform connectivity problems, *J. of Algorithms* 21, 1996, 434–450.
- [17] W. Mader, Ecken vom Grad n in minimalen n -fach zusammenhängenden Graphen, *Archive der Mathematik* 23, 1972, 219–224.
- [18] W. Mader, Minimal n -fach in minimalen n -fach zusammenhängenden Digraphen, *J. Combinatorial Theory Series B*, 38, 1985, 102–117.
- [19] W. Mader, Degree and local connectivity in finite graphs, *Recent advances in graph theory (Proc. Second Czechoslovak Sympos., Prague, 1974)*, Academia, Prague, 1975, 341–344.
- [20] J. B. Orlin, A faster strongly polynomial minimum cost flow algorithm, *Operations Research* 41, 1993, 338–350.
- [21] R. Ravi and D. P. Williamson, An approximation algorithm for minimum-cost vertex-connectivity problems, *Algorithmica* 18, 1997, 21–43.
- [22] R. Ravi and D. P. Williamson, Erratum: An approximation algorithm for minimum-cost vertex-connectivity problems, manuscript, July 2001.