Maximum cut on interval graphs of interval count four is NP-complete

Celina M. H. de Figueiredo ⊠©

Federal University of Rio de Janeiro, Rio de Janeiro, Brazil

Alexsander A. de Melo 🖂 💿 Federal University of Rio de Janeiro, Rio de Janeiro, Brazil

Fabiano S. Oliveira ⊠ <a>D Rio de Janeiro State University, Rio de Janeiro, Brazil

Ana Silva ⊠[®] Federal University of Ceará, Ceará, Brazil

— Abstract -

The computational complexity of the MAXCUT problem restricted to interval graphs has been open since the 80's, being one of the problems proposed by Johnson in his Ongoing Guide to NP-completeness, and has been settled as NP-complete only recently by Adhikary, Bose, Mukherjee and Roy. On the other hand, many flawed proofs of polynomiality for MAXCUT on the more restrictive class of unit/proper interval graphs (or graphs with interval count 1) have been presented along the years, and the classification of the problem is still unknown. In this paper, we present the first NP-completeness proof for MAXCUT when restricted to interval graphs with bounded interval count, namely graphs with interval count 4.

1 Introduction

A cut is a partition of the vertex set of a graph into two disjoint parts and the maximum cut problem (denoted MAXCUT for short) aims to determine a cut with the maximum number of edges for which each endpoint is in a distinct part. The decision problem MAXCUT is known to be NP-complete since the seventies [18], and only recently its restriction to interval graphs has been announced to be hard [1], settling a long-standing open problem that appeared in a summary table in the 1985 column of the Ongoing Guide to NP-completeness by David S. Johnson [20]. We refer the reader to a revised version of Johnson's summary table in [15], where one can also find a parameterized complexity version of the said table.

An interval model is a family of closed intervals of the real line. A graph is an interval graph if there exists an interval model, for which each interval corresponds to a vertex of the graph, such that distinct vertices are adjacent in the graph if and only if the corresponding intervals intersect. Ronald L. Graham proposed in the 80's the study of the *interval count* of an interval graph as the smallest number of interval lengths used by an interval model of the graph. Interval graphs having interval count 1 are called *unit interval graphs* (these are also called proper interval graphs, or indifference graphs). Understanding the interval count, besides being an interesting and challenging problem by itself, can be also of value for the investigation of problems that are hard for general interval graphs, and easy for unit interval graphs (e.g. geodetic number [9, 16], optimal linear arrangement [10, 19], sum coloring [23, 24]). The positive results for unit interval graphs usually take advantage of the fact that a representation for these graphs can be found in linear time [11, 14]. Surprisingly, the recognition of interval graphs with interval count k is open, even for k = 2 [7]. Nevertheless, another generalization of unit interval graphs has been recently introduced which might be more promising in this aspect. These graphs are called k-nested interval graphs, for which an efficient recognition algorithm has firstly appeared in [8]. Recently, a linear time algorithm has been devised in [21].

In the same way that MAXCUT on interval graphs has evaded being solved for so long, the community has been puzzled by the restriction to unit interval graphs. Indeed, two attempts at solving it in polynomial time were proposed in [4,6] just to be disproved closely after [3,22]. In this paper, we give the first classification that bounds the interval count, namely, we prove that MAXCUT is NP-complete when restricted to interval graphs of interval count 4. This also implies



© Celina M. H. de Figueiredo, Alexsander A. de Melo, Fabiano S. Oliveira, and Ana Silva; licensed under Creative Commons License CC-BY 4.0

Leibniz International Proceedings in Informatics

LIPICS Schloss Dagstuhl – Leibniz-Zentrum für Informatik, Dagstuhl Publishing, Germany

XX:2 Maximum cut on interval graphs of interval count four is NP-complete

NP-completeness for the newly generalized class of 4-nested graphs, and opens the search for a full polynomial/NP-complete dichotomy classification in terms of the interval count. It can still happen that the problem is hard even on graphs of interval count 1. We contribute towards filling the complexity gap between interval and unit interval graphs. We have communicated the result at the MFCS 2021 conference [13], and previous versions of the full proof appeared in the ArXiv [12]. The present paper contains the improved and much shorter full proof.

Next, we establish basic definitions and notation. Section 2 describes our reduction and Section 3 discusses the interval count of the interval graph constructed in [1].

1.1 Preliminaries

In this work, all graphs considered are simple. For missing definitions and notation of graph theory, we refer to [5]. For a comprehensive study of interval graphs, we refer to [17].

Let G be a graph. Let X and Y be two disjoint subsets of V(G). We let $E_G(X, Y)$ be the set of edges of G with an endpoint in X and the other endpoint in Y. A cut of G is a partition of V(G)into two parts $A, B \subseteq V(G)$, denoted by [A, B]; the edge set $E_G(A, B)$ is called the *cut-set* of G associated with [A, B]. The size of a cut-set is defined as its cardinality. The size of a cut is the size of its associated cut-set. For each two vertices $u, v \in V(G)$, we say that u and v are in a same part of [A, B] if either $\{u, v\} \subseteq A$ or $\{u, v\} \subseteq B$; otherwise, we say that u and v are in opposite parts of [A, B]. Denote by $\mathsf{mc}(G)$ the maximum size of a cut-set of G. The MAXCUT problem has as input a graph G and a positive integer k, and it asks whether $\mathsf{mc}(G) \ge k$.

Let $I \subseteq \mathbb{R}$ be a closed interval of the real line. We let $\ell(I)$ and r(I) denote respectively the minimum and maximum points of I, which we call the *left* and the *right endpoints* of I, respectively. For every non-empty collection of intervals \mathcal{H} , we define the *left endpoint* of \mathcal{H} as $\ell(\mathcal{H}) = \min_{I \in \mathcal{H}} \ell(I)$ and the *right endpoint* of \mathcal{H} as $r(\mathcal{H}) = \max_{I \in \mathcal{H}} r(I)$. We denote a closed interval I by $[\ell(I), r(I)]$. Distinction from the cut notation will be clear from the context. For every two intersecting intervals I and I', we say that I covers I' if $\ell(I) \leq \ell(I')$ and $r(I) \geq r(I')$, that I intersects I' to the *left* if $\ell(I) < \ell(I') < r(I) < r(I')$, and that I intersects I' to the *right* if $\ell(I) < \ell(I) < r(I) < r(I)$. We say that an interval I precedes an interval I' if $r(I) < \ell(I')$; and more generally, we say that a collection of intervals \mathcal{H} occurs to the *left* of a collection \mathcal{H}' if every interval in \mathcal{H} precedes every interval in \mathcal{H}' . The *length* of an interval I is defined as $|I| = r(I) - \ell(I)$.

An interval model is a finite multiset \mathcal{M} of intervals. The interval count of an interval model \mathcal{M} , denoted by $ic(\mathcal{M})$, is defined as the number of distinct lengths of the intervals in \mathcal{M} . Let G be a graph and \mathcal{M} be an interval model. An \mathcal{M} -representation of G is a bijection $\phi: V(G) \to \mathcal{M}$ such that, for every two distinct vertices $u, v \in V(G)$, we have that $uv \in E(G)$ if and only if $\phi(u) \cap \phi(v) \neq \emptyset$. If such an \mathcal{M} -representation exists, we say that \mathcal{M} is an interval model of G. We note that a graph may have either no interval model or arbitrarily many distinct interval models. A graph is called an interval graph if it has an interval model. The interval count of an interval graph G, denoted by ic(G), is defined as $ic(G) = \min\{ic(\mathcal{M}): \mathcal{M} \text{ is an interval model of } G\}$. An interval graph is called a unit interval graph if its interval count is equal to 1.

Note that, for every interval model \mathcal{M} , there exists a unique (up to isomorphism) graph that admits an \mathcal{M} -representation. Thus, for every interval model $\mathcal{M} = \{I_1, \ldots, I_n\}$, we let $\mathbb{G}_{\mathcal{M}}$ be the graph with vertex set $V(\mathbb{G}_{\mathcal{M}}) = \{1, \ldots, n\}$ and edge set $E(\mathbb{G}_{\mathcal{M}}) = \{ij: I_i, I_j \in \mathcal{M}, I_i \cap I_j \neq \emptyset, i \neq j\}$. Since $\mathbb{G}_{\mathcal{M}}$ is uniquely determined (up to isomorphism) from \mathcal{M} , in what follows we may make an abuse of language and use graph terminologies to describe properties related to the intervals in \mathcal{M} . Two intervals $I_i, I_j \in \mathcal{M}$ are said to be *true twins* in $\mathbb{G}_{\mathcal{M}}$ if they have the same closed neighborhood in $\mathbb{G}_{\mathcal{M}}$.

2 Our reduction

The following theorem is the main contribution of this work:

▶ **Theorem 1.** MAXCUT is NP-complete on interval graphs of interval count 4.

This result is a stronger version of that of Adhikary et al. [1]. To prove Theorem 1, we present a polynomial-time reduction from MAXCUT on cubic graphs, which is known to be NP-complete [2]. Since our proof is based on that of Adhikary et al., we start by presenting some important properties of their key gadget.

2.1 Grained gadget

The interval graph constructed in the reduction of [1] is strongly based on two types of gadgets, called *V-gadgets* and *E-gadgets*. In fact, these gadgets have the same structure except for the number of intervals of certain kinds contained in each of them. In this subsection, we present a generalization of such gadgets, rewriting their key properties to suit our purposes. In order to discuss the interval count of the reduction of [1], we describe it in detail in Section 3.

Let x and y be two positive integers. An (x, y)-grained gadget (see Figure 1 to follow) is an interval model \mathcal{H} formed by 2y long intervals, y of which called *left long* and y called *right long* intervals, together with 2x pairwise disjoint short intervals, x of which called *left short* and x of which called *right short*. The y left long intervals all have the same right endpoint, which also is the left endpoint of each of the y right long intervals. The x left (resp. right) short intervals are all pairwise disjoint and intersect each left (resp. right) long interval, but intersect no right (resp. left) long interval. We write $\mathcal{LS}(\mathcal{H}), \mathcal{LL}(\mathcal{H}), \mathcal{RS}(\mathcal{H})$ and $\mathcal{RL}(\mathcal{H})$ to denote the left short, left long, right short and right long intervals of \mathcal{H} , respectively. And we omit \mathcal{H} when it is clear from the context.

Note that, if \mathcal{H} is an (x, y)-grained gadget, then $\mathbb{G}_{\mathcal{H}}$ is a split graph such that $\mathcal{LS} \cup \mathcal{RS}$ is an independent set of size 2x, $\mathcal{LL} \cup \mathcal{RL}$ is a clique of size 2y, and, for every vertex $u \in \mathcal{LS}$, $N_{\mathbb{G}_{\mathcal{H}}}(u) = \mathcal{LL}$ and, for every vertex $u \in \mathcal{RS}$, $N_{\mathbb{G}_{\mathcal{H}}}(u) = \mathcal{RL}$. Moreover, the intervals in \mathcal{LL} are true twins in $\mathbb{G}_{\mathcal{H}}$; similarly, the intervals in \mathcal{RL} are true twins in $\mathbb{G}_{\mathcal{H}}$.



Figure 1 General structure of an (x, y)-grained gadget.

Let \mathcal{M} be an interval model containing an (x, y)-grained gadget \mathcal{H} . We say that an interval of $\mathcal{M} \setminus \mathcal{H}$ intersects \mathcal{H} if it intersects at least one interval of \mathcal{H} . Otherwise, we say that the interval does not intersect \mathcal{H} . The possible types of intersections between an interval $I \in \mathcal{M} \setminus \mathcal{H}$ and \mathcal{H} in our construction are depicted in Figure 2, with the used nomenclature. More specifically, the intersection between I and \mathcal{H} is a cover intersection if I intersects all the intervals of \mathcal{H} (Figure 2(a)), a weak intersection to the left (right) if I intersects exactly the left (right) long intervals of \mathcal{H} (Figure 2(b-c)), and a strong intersection to the left (right) if I intersects exactly the left (right) long and short intervals of \mathcal{H} (Figure 2(d-e)). We say that \mathcal{M} respects the structure of \mathcal{H} if, for every interval $I \in \mathcal{M} \setminus \mathcal{H}$, we have that I either does not intersect \mathcal{H} , or the intersection between I and \mathcal{H} is of one of the types described above.

The advantage of this gadget is that, by manipulating the values of x and y, we can ensure that, in a maximum cut, the left long and right short intervals are placed in the same part, opposite to the part containing the left short and right long intervals, as proved in Lemma 3, presented shortly. Note that if \mathcal{M} is an interval model containing a grained gadget \mathcal{H} and \mathcal{M} respects the structure of \mathcal{H} , then every left (resp. right) short interval of \mathcal{H} intersects exactly the same set of intervals in \mathcal{M} . The following remark will be useful throughout the text.

▶ Remark 2. Let [A, B] be a maximum cut of a graph G. For any vertex $u \in V(G)$, if more than half of the neighbours of u are in one part of [A, B], say A, then $u \notin A$, or in other words $u \in B$.

XX:4 Maximum cut on interval graphs of interval count four is NP-complete



Figure 2 Interval $I \in \mathcal{M} \setminus \mathcal{H}$ (a) covering \mathcal{H} , (b-c) weakly intersecting \mathcal{H} to the left and to the right, and (d-e) strongly intersecting \mathcal{H} to the left and to the right.

Proof. Suppose that $u \in A$, and let [A', B'] be the cut of G such that $A' = A \setminus \{u\}$ and $B' = B \cup \{u\}$. Note that, if $e \in E_G(A, B) \setminus E_G(A', B')$, then e is incident to u. Thus, since u has more than half of its neighbours in A, the size of [A', B'] is strictly greater than the size of [A, B], contradicting the maximality of [A, B].

▶ Lemma 3. Let x and y be positive integers and \mathcal{M} an interval model containing an (x, y)-grained gadget \mathcal{H} . Suppose that \mathcal{M} respects the structure of \mathcal{H} . Let [A, B] be a maximum cut of $\mathbb{G}_{\mathcal{M}}$. Also, let t be the number of intervals in $\mathcal{M} \setminus \mathcal{H}$ intersecting \mathcal{H} , ℓ be the number of intervals in $\mathcal{M} \setminus \mathcal{LS}$ intersecting the left short intervals of \mathcal{H} , and r be the number of intervals in $\mathcal{M} \setminus \mathcal{RS}$ intersecting the right short intervals of \mathcal{H} . If ℓ and r are odd, y > t(x/y - 1) and x > t + 2y, then the following hold:

1. $\mathcal{LS}(\mathcal{H}) \subseteq A$ and $\mathcal{LL}(\mathcal{H}) \subseteq B$, or vice versa;

2. $\mathcal{RS}(\mathcal{H}) \subseteq A$ and $\mathcal{RL}(\mathcal{H}) \subseteq B$, or vice versa; and

3. $\mathcal{LL}(\mathcal{H}) \subseteq A$ and $\mathcal{RL}(\mathcal{H}) \subseteq B$, or vice versa.

Proof. First, we prove that all the left short intervals are in the same part of [A, B]. Denote by \mathcal{N} the set of intervals in $\mathcal{M} \setminus \mathcal{LS}$ that intersect the left short intervals.

Suppose, without loss of generality, that B contains more than half of the intervals in \mathcal{N} (it must occur for either A or B since ℓ is odd). Consider any $u \in \mathcal{LS}$. Then \mathcal{N} is the set of neighbours of u, and since more than half of the intervals of \mathcal{N} are in B, it follows that $u \in A$. This shows that $\mathcal{LS} \subseteq A$. Thus all the left short intervals are in the same part of [A, B]. Because r is also odd, a similar argument shows that all the right short intervals are in the same part of [A, B].

Now consider the left long intervals and suppose, without loss of generality, that all the left short intervals are contained in A. Observe that the number of intervals in $\mathcal{M} \setminus \mathcal{LS}$ intersecting a left long interval is less than t + 2y < x. Thus every left long interval has more than half of its neighbours from \mathcal{LS} , which are all in one part of [A, B]. It now follows that every left long interval is in the part of [A, B] opposite to that of the left short intervals, namely B. An analogous argument holds for the right long intervals. This proves Claims (1) and (2) in the statement of the lemma.

Finally, let \mathcal{L} denote the set of long intervals of \mathcal{H} and suppose by contradiction that $\mathcal{L} \subseteq A$. Let T be the set of intervals in $\mathcal{M} \setminus \mathcal{H}$ that intersect \mathcal{H} ; then t = |T|. Let $t_A = |T \cap A|$ and $t_B = |T \cap B|$. Now by switching the intervals in \mathcal{RL} to B and \mathcal{RS} to A, we gain at least $y^2 + yt_A + xt_B$ cut-edges and lose at most $xt_A + yt_B$ cut-edges. Since y > t(x/y - 1), we have $y^2 > t(x - y) = xt - yt$ or in other words, $y^2 > xt_A + xt_B - yt_A - yt_B$. So we get $y^2 + yt_A + xt_B > xt_A + yt_B + 2t_B(x - y)$. As x > y, we can conclude that $y^2 + yt_A + xt_B > xt_A + yt_B$, which means that we have more cut-edges in the new cut than in the cut [A, B], a contradiction.

We say that $(\mathcal{H}, \mathcal{M})$ is well-valued if the conditions of Lemma 3 are satisfied. Moreover, we say that the constructed model \mathcal{M} is well-valued if all its grained gadgets \mathcal{H} are well-valued with respect to the model \mathcal{M} . Finally, we say that \mathcal{H} is A-partitioned by [A, B] if $\mathcal{LS}(\mathcal{H}) \cup \mathcal{RL}(\mathcal{H}) \subseteq A$ and $\mathcal{RS}(\mathcal{H}) \cup \mathcal{LL}(\mathcal{H}) \subseteq B$. Define B-partitioned analogously.

2.2 Reduction graph

In this subsection, we formally present our construction. We will make a reduction from MAXCUT on cubic graphs. So, let G be a cubic graph on n vertices and m edges. Intuitively, we consider an ordering of the edges of G, and we divide the real line into m regions, with the j-th region holding the information about whether the j-th edge is in the cut-set. For this, each vertex u will be related to a subset of intervals traversing all the m regions, bringing the information about which part of the cut contains u. Let $\pi_V = (v_1, \ldots, v_n)$ be an ordering of V(G), $\pi_E = (e_1, \ldots, e_m)$ be an ordering of E(G), and $\mathfrak{G} = (G, \pi_V, \pi_E)$.



Figure 3 General structure of a region of an (n, m)-escalator. The shaded rectangles represent the (p, q)-grained gadgets \mathcal{H}_i^j .

We first describe the gadgets related to the vertices. Please refer to Figure 3 to follow the construction. The values of p, q used next will be defined later. An (n, m)-escalator is an interval model \mathcal{D} formed by m+1 (p, q)-grained gadgets for each v_i , denoted by $\mathcal{H}_i^1, \ldots, \mathcal{H}_i^{m+1}$, together with 2m link intervals, denoted by L_i^1, \ldots, L_i^{2m} , such that L_i^{2j-1} and L_i^{2j} weakly intersect \mathcal{H}_i^j to the right and weakly intersect \mathcal{H}_i^{j+1} to the left. Additionally, all the grained gadgets are mutually disjoint. More specifically, given $j \in \{1, \ldots, m+1\}$ and $i, i' \in \{1, \ldots, n\}$ with i < i', the grained gadget \mathcal{H}_i^j occurs to the left of $\mathcal{H}_{i'}^{j+1}$ for $j \in \{1, \ldots, m\}$.



Figure 4 General structure of the constructed interval model $\mathcal{M}(\mathfrak{G})$ highlighting the intersections between the intervals of the (n, m)-escalator \mathcal{D} , the intervals of the (p', q')-grained gadget \mathcal{E}_j , and the intervals $C_j^1, C_j^2, C_j^3, C_j^4$.

Now, we add the gadgets related to the edges. Please refer to Figure 4 to follow the construction.

XX:6 Maximum cut on interval graphs of interval count four is NP-complete

The values of p', q' used next will be defined later. For each edge $e_j = v_i v_{i'} \in E(G)$, with i < i', create a (p',q')-grained gadget \mathcal{E}_j and intervals $C_j^1, C_j^2, C_j^3, C_j^4$ in such a way that \mathcal{E}_j is entirely contained in the *j*-th region (i.e., in the open interval between the right endpoint of \mathcal{H}_n^j and the left endpoint of \mathcal{H}_1^{j+1}), C_j^1 and C_j^2 weakly intersect \mathcal{H}_i^j to the right and weakly intersect \mathcal{E}_j to the left, and C_j^3 and C_j^4 weakly intersect $\mathcal{H}_{i'}^j$ to the right and strongly intersect \mathcal{E}_j to the left. We call the intervals in $\{C_j^i \mid i \in \{1, \ldots, 4\}, j \in \{1, \ldots, m\}\}$ intervals of type C. Denote the constructed model by $\mathcal{M}(\mathfrak{G})$ (or simply by \mathcal{M} when \mathfrak{G} is clear from the context), which defines the reduction graph $\mathbb{G}_{\mathcal{M}(\mathfrak{G})}$.

The following straightforward lemma will be useful in the next section.

▶ Lemma 4. Let G be a graph, $\pi_V = (v_1, \ldots, v_n)$ and $\pi_E = (e_1, \ldots, e_m)$ be orderings of V(G) and E(G), respectively, and \mathcal{M} be the model constructed as before. The following holds for every grained gadget \mathcal{H} :

- $\blacksquare \mathcal{M} respects the structure of \mathcal{H};$
- \blacksquare The number of intervals covering \mathcal{H} is even; and
- The number of intervals strongly intersecting H to the left is either zero or two, and the number of intervals strongly intersecting H to the right is always zero.

Observe that Lemma 4 implies that, in order for the values ℓ and r in Lemma 3 to be odd, it suffices to choose odd values for q and q'.

2.3 Proof of Theorem 1: Maximum cut of the reduction graph

Consider a cubic graph G on n vertices and m = 3n/2 edges, and let $\pi_V = (v_1, \ldots, v_n)$ be an ordering of V(G), $\pi_E = (e_1, \ldots, e_m)$ be an ordering of E(G) and $\mathfrak{G} = (G, \pi_V, \pi_E)$. We now prove that $\mathsf{mc}(G) \geq k$ if and only if $\mathsf{mc}(\mathbb{G}_{\mathcal{M}(\mathfrak{G})}) \geq f(G, k)$, where f is a polynomial-time computable function defined at the end of this subsection. As it is usually the case in this kind of reduction, given a cut of G, constructing an appropriate cut of the reduction graph $\mathbb{G}_{\mathcal{M}(\mathfrak{G})}$ is an easy task. On the other hand, constructing an appropriate cut [X, Y] of G, from a given a cut [A, B] of the reduction graph $\mathbb{G}_{\mathcal{M}(\mathfrak{G})}$, requires that the intervals in $\mathcal{M}(\mathfrak{G})$ behave in a way with respect to [A, B] so that [X, Y] can be inferred, a task achieved by appropriately manipulating the values of p, q, p', q', as done in Lemma 3. We start by giving conditions on these values that ensure that the partitioning of the edge gadget related to an edge $e_j = v_i v_{i'}$, with i < i', depends solely on the partitioning of $\mathcal{H}^j_{i'}$.

▶ Lemma 5. Let G be a cubic graph, $\pi_V = (v_1, \ldots, v_n)$ and $\pi_E = (e_1, \ldots, e_m)$ be orderings of V(G)and E(G), respectively, and $\mathcal{M}(\mathfrak{G})$ be the model constructed as before, where $\mathfrak{G} = (G, \pi_V, \pi_E)$. Also, let [A, B] be a maximum cut of $\mathbb{G}_{\mathcal{M}(\mathfrak{G})}$, and consider $e_j = v_i v_{i'}$, i < i'. If $\mathcal{M}(\mathfrak{G})$ is well-valued, and q > 4n + p' + q' + 3, then

- 1. If \mathcal{H}_i^j is A-partitioned by [A, B], then $\{C_j^1, C_j^2\} \subseteq B$; otherwise, $\{C_j^1, C_j^2\} \subseteq A$; and
- **2.** If $\mathcal{H}_{i'}^j$ is A-partitioned by [A, B], then $\{C_j^3, C_j^4\} \subseteq B$ and \mathcal{E}_j is A-partitioned by [A, B]; otherwise, $\{C_j^3, C_j^4\} \subseteq A$ and \mathcal{E}_j is B-partitioned by [A, B].

Proof. Denote $\mathcal{M}(\mathfrak{G})$ by \mathcal{M} for simplicity. Since \mathcal{M} is well-valued, by Lemma 3, we may assume that \mathcal{H}_i^j is A-partitioned by [A, B], i.e., that $\mathcal{LS} \cup \mathcal{RL} \subseteq A$ and $\mathcal{LL} \cup \mathcal{RS} \subseteq B$. We make the arguments for C_j^1 and it will be clear that they also hold for C_j^2 . Observe first that all the grained gadgets covered by C_j^1 have a balanced number of intervals in A and in B. More formally, from the intervals within the gadgets \mathcal{H}_ℓ^j , $i + 1 \leq \ell \leq n$, which are all the grained gadgets covered by C_j^1 , there are exactly (n-i)(p+q) intervals in A, and (n-i)(p+q) intervals in B. Additionally, there are at most 2(n-i) link intervals intersecting C_j^1 to the left (these are the link intervals related to $v_{i''}$ for i'' > i in the (j-1)-th region, if j > 1), exactly 2(n-i) link intervals intersecting C_j^1 to the right (these are the link intervals related to $v_{i''}$ for i'' > i in the j-th region). This is a total of at most 2(n-i) + 2(n-i) + 2i = 4n - 2i < 4n link intervals. Adding finally C_i^2, C_i^3, C_i^4

and the q' right long intervals of \mathcal{E}_j , we get that the number of neighbors of C_j^1 that might be in B is at most (n-i)(p+q) + 4n + q' + 3, while the number of neighbors of C_j^1 that are in A is at least (n-i)(p+q) + q. Since $q > 4n + p' + q' + 3 \ge 4n + q' + 3$, we can conclude that there are more neighbours of C_j^1 in A than in B. From Remark 2, it follows that $C_j^1 \in B$.

Observe that a similar argument can be applied to C_j^3, C_j^4 , except that we gain also p' new edges from the left short intervals of \mathcal{E}_j . That is, supposing $\mathcal{H}_{i'}^j$ is A-partitioned by [A, B], then the number of neighbors of C_j^3 that might be in B is at most (n - i')(p + q) + 4n + p' + q' + 3, while the number of neighbors of C_j^3 that are in A is at least (n - i')(p + q) + q. It follows again by Remark 2 that C_j^3, C_j^4 are in B, since q > 4n + p' + q' + 3.

Finally, suppose that $\mathcal{H}_{i'}^j$ is A-partitioned by [A, B], in which case, from the previous paragraph, we get that $\{C_j^3, C_j^4\} \subseteq B$. Suppose that \mathcal{E}_j is B-partitioned. Then consider the cut [A', B'] obtained by switching the intervals of \mathcal{E}_j of side; formally, in which every interval $I \in \mathcal{M} \setminus \mathcal{E}_j$ is in A' if and only if $I \in A$ and every interval $I \in \mathcal{E}_j$ is in A' if and only if $I \in B$. Clearly, the number of cut-edges having both endpoints in \mathcal{E}_j is the same in both the cuts [A, B] and [A', B']. Since $|A \cap \mathcal{E}_j|$ $= |B \cap \mathcal{E}_j| = |A' \cap \mathcal{E}_j| = |B' \cap \mathcal{E}_j|$, and every interval other than C_j^1, C_j^2, C_j^3 and C_j^4 that intersects the gadget \mathcal{E}_j has a cover intersection with it, the number of cut-edges in [A, B] differ from that of [A', B'] only by the number of cut-edges between $C_j^1, C_j^2, C_j^3, C_j^4$ and \mathcal{E}_j . Since $C_j^3, C_j^4 \in B$, the 2p'edges between these two intervals and the intervals in $\mathcal{LS}(\mathcal{E}_j)$ are cut-edges in [A, B] but not in [A, B]. Meanwhile, the edges between $C_j^1, C_j^2, C_j^3, C_j^4$ and $\mathcal{LL}(\mathcal{E}_j)$. Thus [A', B'] has at least 2p' - 4q' edges more than [A, B]. Since \mathcal{M} is well-valued, we have 2p' > 4q', implying that [A', B'] is a cut of size larger than [A, B], which is a contradiction.

After ensuring that each grained gadget behaves well individually, we also need to ensure that \mathcal{H}_i^1 can be used to decide in which part of [X, Y] we should put v_i , and for this it is necessary that all gadgets related to v_i agree with one another. In other words, for each v_i , we want that the behaviour of the first gadget \mathcal{H}_i^1 influence the behaviour of the subsequent gadgets $\mathcal{H}_i^2, \ldots, \mathcal{H}_i^{m+1}$, as well as the behaviour of the gadgets related to edges incident to v_i . Given $v_i \in V(G)$ and a cut [A, B] of $\mathbb{G}_{\mathcal{M}(\mathfrak{G})}$, we say that the gadgets of v_i alternate in [A, B] if, for every $j \in \{1, \ldots, m\}$, we get that \mathcal{H}_i^j is A-partitioned if and only if \mathcal{H}_i^{j+1} is B-partitioned, while L_i^{2j-1}, L_i^{2j} are opposite to the right long intervals of \mathcal{H}_i^j . Also, we say that [A, B] is alternating partitioned if the gadgets of v_i alternate in [A, B], for every $v_i \in V(G)$. We add a further condition on the values of p, q, p', q' in order to ensure that every maximum cut is alternating partitioned. After this, we use the good behaviour of the constructed model in order to relate the sizes of the maximum cuts in G and in $\mathbb{G}_{\mathcal{M}(\mathfrak{G})}$.

▶ Lemma 6. Let G be a cubic graph, $\pi_V = (v_1, \ldots, v_n)$ and $\pi_E = (e_1, \ldots, e_m)$ be orderings of V(G) and E(G), respectively, and $\mathcal{M}(\mathfrak{G})$ be the model constructed as before, where $\mathfrak{G} = (G, \pi_V, \pi_E)$. Also, let [A, B] be a maximum cut of $\mathbb{G}_{\mathcal{M}(\mathfrak{G})}$. If $\mathcal{M}(\mathfrak{G})$ is well-valued, q > 4n + p' + q' + 3, and $q > 3(2n^2 + 2n + q' + 2)$, then [A, B] is alternating partitioned.

Proof. By hypothesis, the conditions of Lemmas 3 and 5 are satisfied. Thus, we can suppose that the obtained properties of those lemmas hold. Denote $\mathcal{M}(\mathfrak{G})$ by \mathcal{M} for simplicity, and let \mathcal{M}_i be the family of all the intervals related to vertex v_i ; more specifically, it contains every interval in some grained gadget \mathcal{H}_i^j , $j \in \{1, \ldots, m+1\}$, every link interval L_i^j , $j \in \{1, \ldots, 2m\}$, every interval of type C that intersects \mathcal{H}_i^j to the right (this happens if e_j has v_i as endpoint), and every interval in \mathcal{E}_j for e_j incident to v_i . In what follows, we count the number f_i of edges of the cut incident to some interval in \mathcal{M}_i and argue that, if the gadgets of v_i do not alternate in [A, B], then we can obtain a bigger cut by switching the side of some intervals, thus getting a contradiction.

Denote by $\overline{\mathcal{M}}_i$ the set of intervals $\mathcal{M} \setminus \mathcal{M}_i$, and by Λ the set of all link intervals.

In what follows, there are some values that must be added to f_i but remain the same in every maximum cut of $\mathbb{G}_{\mathcal{M}(\mathfrak{G})}$, independently of how \mathcal{M}_i is partitioned; we call these values *irrelevant* and do not add them to f_i . For instance, recall that every (x, y)-grained gadget has exactly x + yintervals in A and x + y in B. Thus, because of Lemmas 3 and 5, the number of edges of the cut

XX:8 Maximum cut on interval graphs of interval count four is NP-complete

between grained gadgets and intervals that cover them is irrelevant. In what follows, we count the other possible edges.

First, consider $j \in \{1, \ldots, m\}$; we want to count the maximum number of edges of the cut incident to L_i^{2j} (which holds analogously for L_i^{2j-1}). Denote by ℓ_A^j the number of intervals in $\overline{\mathcal{M}}_i \cap \Lambda \cap A$ that intersect L_i^{2j} ; define ℓ_B^j similarly. Observe that $\ell_A^j + \ell_B^j < 4n$ since it includes at most 2(n-i) link intervals in the *j*-th region, plus at most 2(n-i) link intervals of the (j-1)-th region, and at most 2(i-1) link intervals of the (j+1)-th region. Additionally, let a_j be equal to 1 if L_i^{2j} is opposite to the right long intervals of \mathcal{H}_i^{j+1} , and 0 otherwise; similarly, let b_j be equal to 1 if L_i^{2j} is opposite to the left long intervals of \mathcal{H}_i^{j+1} , and 0 otherwise. Because L_i^{2j} might also be opposite to C_j^1, \ldots, C_j^4 and it is possible that the edge between L_i^{2j} is at most $q(a_j + b_j) + \ell_A^j + \ell_B^j + 5$. Note that L_i^{2j} covers the gadgets \mathcal{E}_j and also every $\mathcal{H}_{i'}^{j'}$ with which it has an intersection except \mathcal{H}_i^j and \mathcal{H}_i^{j+1} , and hence the number of cut-edges between L_i^{2j} and intervals in these gadgets is irrelevant.

Now, let e_j be an edge incident to v_i and let $v_{i'}$ be the other endpoint of e_j (here i' might be smaller than i). We apply Lemma 5 in order to count the edges incident to $\mathcal{E}_j \cup \{C_j^1, \ldots, C_j^4\}$; observe that all these intervals are in \mathcal{M}_i . First observe that, since \mathcal{E}_j is always partitioned according to C_j^3, C_j^4 , we have an irrelevant value of 2p', namely the edges between C_j^3, C_j^4 and the left short intervals of \mathcal{E}_j . Now, suppose, without loss of generality, that $\{C_j^1, C_j^2\} \subseteq A$. If $\{C_j^3, C_j^4\} \subseteq A$, then there are no relevant edges to be added; otherwise, we get 2q' + 4 edges, those between C_j^1, C_j^2 and C_j^3, C_j^4 , and between C_j^1, C_j^2 and the left long intervals of \mathcal{E}_j . Finally, observe that the edges between $\{C_j^1, \ldots, C_j^4\}$ and \mathcal{H}_i^j are irrelevant because of Lemma 5 and the fact that C_j^1, C_j^2 cover $\mathcal{H}_{i'}^j$ (where e_j is the edge $v_i v_{i'}$), and that the edges between $\{C_j^1, \ldots, C_j^4\}$ and the link intervals have been counted previously. Note that the number of cut edges between two intervals in \mathcal{E}_j and the number of cut edges between intervals in \mathcal{E}_j and link intervals are both irrelevant. Note also that every gadget \mathcal{H}_i^j , for each $j \in \{1, 2, \ldots, m+1\}$, is covered by every interval from $\overline{\mathcal{M}}_i$ that it intersects, and hence the number of cut edges between vertices in \mathcal{H}_i^j is irrelevant. Also, the number of cut edges between vertices in \mathcal{H}_i^j have already been counted.

In order to put everything together, let e_{j_1} , e_{j_2} , e_{j_3} be all the edges incident to v_i , and for each $h \in \{1, 2, 3\}$, write $e_{j_h} = v_i v_{i_h}$ (note that here *i* is not necessarily smaller than i_h). For each $h \in \{1, 2, 3\}$, let c_h be equal to 1 if \mathcal{H}_i^j and $\mathcal{H}_{i_h}^j$ are partitioned differently, and 0 otherwise. We then get that:

$$f_i \le 2\sum_{j=1}^m (q(a_j + b_j) + \ell_A^j + \ell_B^j + 5) + \sum_{h=1}^3 c_h (2q' + 4).$$
(1)

If L_i^{2j} is on the same side as the right long intervals of \mathcal{H}_i^j and the left long intervals of \mathcal{H}_i^{j+1} , we can increase f_i simply by switching the side of L_i^{2j} . Indeed, in this case we would lose at most $\max\{\ell_A^j, \ell_B^j\} + 5 < 4n + 5$ edges, while gaining 2q, a positive exchange since 2q > 8n > 4n + 5considering n > 1. Observe that this implies $a_j + b_j \ge 1$. Note also that this type of argument can be always applied, and that it can be applied also for L_i^{2j-1} . Hence, whenever in what follows we switch side of the intervals in some vertex gadget, we can suppose that this property still holds, i.e. that L_i^{2j} and L_i^{2j-1} are always opposite to the left long intervals of \mathcal{H}_i^j .

Consider now j to be minimum such that \mathcal{H}_i^j and \mathcal{H}_i^{j+1} are partitioned in the same way, say they are both A-partitioned. Note that this implies that $a_j + b_j = 1$, since the right long intervals of \mathcal{H}_i^j are in A, while the left long intervals of \mathcal{H}_i^{j+1} are in B. We want to switch sides of \mathcal{H}_i^{j+1} , but in order to ensure an increase in the size of the cut, we need to also switch subsequent grained gadgets in case they were alternating. For this, let j' > j be minimum such that $\mathcal{H}_i^{j'+1}$ and $\mathcal{H}_i^{j'}$ are either both A-partitioned or both B-partitioned; if it does not exist, let j' = m + 1. For each $h \in \{j + 1, \ldots, j'\}$, we switch sides of \mathcal{H}_i^h , and put L_i^{2h-1}, L_i^{2h} in the side opposite to the right long intervals of \mathcal{H}_i^h . Also switch the intervals of type C and intervals in edge gadgets appropriately; i.e., in a way that Lemma 5 continues to hold. We prove that we gain at least 2q edges, while losing at

most $8m(n+1) + 6(q'+2) = 6(2n^2 + 2n + q' + 2)$ cut edges (recall that m = 3n/2); the result thus follows since $q > 3(2n^2 + 2n + q' + 2)$.

Observe that, by previous arguments, we have that, for every $h \in \{j, \ldots, j'\}$, the link intervals L_i^{2h-1}, L_i^{2h} are in B if and only if \mathcal{H}_i^h is A-partitioned. In particular, since \mathcal{H}_i^j is A-partitioned, L_i^{2j-1} and L_i^{2j} are in B. Additionally, because of the switch we now know that the left long intervals of \mathcal{H}_i^{j+1} are in A. This implies that we gain at least 2q edges. Now, we count our losses. Concerning intervals L_i^{2j-1} and L_i^{2j} , we lose at most $2(\ell_B^j + 4) \leq 8n + 8$ cut edges, namely the edges between these intervals and link intervals or intervals of type C. As for the intervals L_i^{2h-1}, L_i^{2h} for $h \in \{j+1,\ldots,j'\}$, by the definition of j' we know that we lose at most $2(\max\{\ell_A^h, \ell_B^h\} + 4) \leq 8n + 8$ cut edges, while the number of edges of the cut between them and the vertex grained gadgets can only increase. Hence, concerning the link intervals in \mathcal{M}_i , in total we lose at most $8m(n+1) = 12(n^2+n)$ cut edges. Additionally, observe the upper bound given by (1) to see that, in the worst case scenario, we have $\{j_1, j_2, j_3\} \subseteq \{j + 1, \ldots, j'\}$ and all the values c_h were previously equal to 1 and are now equal to 0; this leads to a possible loss of at most 6(q'+2) edges, as we wanted to show.

Now, if [A, B] is an alternating partitioned maximum cut of $\mathbb{G}_{\mathcal{M}(\mathfrak{G})}$, and $\mathcal{M}(\mathfrak{G})$ obeys the conditions in the statement of Lemma 5, we let $\Phi(A, B) = [X, Y]$ be the cut of G such that, for each vertex $v_i \in V(G)$, we have $v_i \in X$ if and only if \mathcal{H}_i^1 is A-partitioned by [A, B]. Note that [X, Y] is well-defined (i.e., Φ is a function). Additionally, given a cut [X, Y] of G, there is a unique alternating partitioned cut [A, B] of $\mathbb{G}_{\mathcal{M}(\mathfrak{G})}$ obeying the conditions of Lemma 5, such that $[X, Y] = \Phi(A, B)$ (i.e., Φ is one-to-one and onto). Therefore, it remains to relate the sizes of these cut-sets. Basically we can use the good behaviour of the maximum cuts in $\mathbb{G}_{\mathcal{M}(\mathfrak{G})}$ to prove that the size of [A, B] grows as a function of the size of $\Phi(A, B)$.

▶ Lemma 7. Suppose that all the conditions in Lemmas 3-6 hold, and that $q' \ge 13n^2$. Let $\Phi(A, B) = [X, Y]$, and k be a positive integer. Then (below, G denotes $\mathbb{G}_{\mathcal{M}(\mathfrak{G})})$

$$|E_G(X,Y)| \ge k$$
 if and only if $|E_{\mathbb{G}}(A,B)| \ge \gamma + (2q'+4)k_s$

where γ is a well-defined polynomial-time computable function on G, π_V, p, q, p', q' (i.e., does not depend on [A, B]).

Proof. We use the same notation as before and count the number of edges in $E_{\mathbb{G}}(A, B)$. We will count the number of edges of the cut-set separately in the following groups:

- among intervals of a vertex/edge grained-gadget;
- between intervals of a vertex grained-gadget and link intervals;
- between intervals of an edge grained-gadget and other intervals;
- \blacksquare among intervals of type C;
- among link intervals;
- \blacksquare between link intervals and intervals of type C; and
- \blacksquare between intervals of a vertex grained-gadget and intervals of type C.

First, we compute the number of edges of the cut-set within a given (x, y)-grained gadget. By Lemma 3, we get that this is exactly $y^2 + 2xy$. Since there are (m + 1)n (p, q)-grained gadgets (the ones related to the vertices), and m (p', q')-grained gadgets (the edge ones), we get a total of:

$$\beta_1 = n(m+1)(q^2 + 2pq) + m((q')^2 + 2p'q').$$

Now, we count the number of edges of the cut-set between a given vertex grained gadget $\mathcal{H} = \mathcal{H}_i^j$ and link intervals; again, denote the set of link intervals by Λ . If an interval I covers \mathcal{H} , then there are exactly p + q edges between I and \mathcal{H} , since there are these many intervals of \mathcal{H} in each of A and B. And if I intersects \mathcal{H} either to the left or to the right, then there are exactly q edges between Iand \mathcal{H} , since \mathcal{M} is alternating partitioned (i.e., I is opposite to the corresponding long intervals of

XX:10 Maximum cut on interval graphs of interval count four is NP-complete

 \mathcal{H}). It remains to count how many of each type of intervals there are. If $j \in \{2, \ldots, m\}$, then there are exactly 2n - 2 intervals covering \mathcal{H} , as well as 2 intervals intersecting \mathcal{H} to the left, and 2 to the right; this gives a total of (2n - 2)(p + q) + 4q = 2n(p + q) + 2(q - p) edges between \mathcal{H} and Λ . If j = 1, then there are 2(i - 1) intervals covering \mathcal{H} , and 2 intervals intersecting \mathcal{H} to the right, thus giving a total of 2(i - 1)(p + q) + 2q. Finally, if j = m + 1, then there are 2(n - i) intervals covering \mathcal{H} , and 2 intervals intersecting \mathcal{H} to the left, giving a total of 2(n - i)(p + q) + 2q. Summing up, we get:

$$\begin{array}{rcl} \beta_2 = & \sum_{j=2}^m \sum_{i=1}^n [2n(p+q) + 2(q-p)] + \\ & \sum_{i=1}^n [2(i-1)(p+q) + 2q + 2(n-i)(p+q) + 2q] \\ = & 2(m-1)n[n(p+q) + (q-p)] + 2n[(n-1)(p+q) + 2q] \\ = & 2n[(m-1)n(p+q) + (m-1)(q-p) + (n-1)(p+q) + 2q] \\ = & 2mn[n(p+q) + q-p]. \end{array}$$

We count now the number of edges of the cut-set between a given edge gadget \mathcal{E}_j and an interval I intersecting it, and among intervals of type C. As before, if I covers \mathcal{E}_j , then there are exactly (p'+q') edges between I and \mathcal{E}_j in the cut. If I strongly intersects \mathcal{E}_j to the left, then $I \in \{C_j^3, C_j^4\}$ and by Lemma 5 we get that this amounts to p'. Finally, if I weakly intersects \mathcal{E}_j to the left, then this amounts to q', if e_j is in the cut-set, or to 0, otherwise. As for the number of edges between intervals of type C, by Lemma 5 one can see that this is equal to $4|E_G(X, Y)|$. Summing up, we get:

$$2nm(p'+q') + 2p'm + (2q'+4)|E_G(X,Y)|$$

Denote the value 2nm(p'+q') + 2p'm by β_3 , and note that this is independent of [A, B].

Let us now count the number of edges of the cut-set among link intervals. For this, denote by \mathcal{L}^{j} the set of link intervals in the *j*-th region, i.e., $\mathcal{L}^{j} = \{L_{i}^{2j}, L_{i}^{2j-1} \mid i \in \{1, \ldots, n\}\}$. Also, denote by V_{A}^{j} the set of indices $i \in \{1, \ldots, n\}$ such that $\{L_{i}^{2j-1}, L_{i}^{2j}\} \subseteq A$; define V_{B}^{j} analogously and let $a = |V_A^j|$ and $b = |V_B^j|$. We count the number of edges of the cut between intervals of \mathcal{L}^j , for every $j \in \{1, \ldots, m\}$, and between intervals of \mathcal{L}^j and intervals of \mathcal{L}^{j+1} , for every $j \in \{1, \ldots, m-1\}$, and then we sum up. So consider a region $j \in \{1, \ldots, m\}$, and observe that, because [A, B] is alternating partitioned, we get that either j is odd and V_{A}^{j} contains exactly the indices of the vertices within Y, while V_B^j contains the indices of the vertices within X, or j is even and the reverse occurs. More formally: if j is odd, then $V_A^j = \{i \in \{1, ..., n\} \mid v_i \in Y\}$ and $V_B^j = \{i \in \{1, ..., n\} \mid v_i \in X\}$; and if j is even, then $V_A^j = \{i \in \{1, ..., n\} \mid v_i \in X\}$ and $V_B^j = \{i \in \{1, ..., n\} \mid v_i \in Y\}$. In either case, since for each index in V_A^j (resp. V_B^j), there is a pair of intervals in $\mathcal{L}^j \cap A$ (resp. $\mathcal{L}^j \cap B$), we get that the number of edges of the cut between intervals of \mathcal{L}^{j} is equal to 4|X||Y| = 4ab. Now, suppose $j \in \{1, \ldots, m-1\}$; we count the edges of the cut between \mathcal{L}^j and \mathcal{L}^{j+1} . Again because [A, B] is alternating partitioned, we know that if $V_A^j = \{i_1, \ldots, i_a\}$, then $V_B^{j+1} = V_A^j$, while $V_A^{j+1} = V_B^j = \{1, \ldots, n\} \setminus V_A^j$. Supposing $i_1 < \cdots < i_a$, this implies that there are exactly 4 edges between $\{L_{i_{a'}}^{2j+1}, L_{i_{a'}}^{2j+2}\}$ and $\{L_{i_{a''}}^{2j-1}, L_{i_{a''}}^{2j}\}$ for each $a', a'' \in \{1, \ldots, a\}$ with a' < a''. Summing up we get that there are $4\sum_{a'=1}^{a'}(a-a') = 4\frac{a(a-1)}{2} = 2a(a-1)$ edges between $\mathcal{L}^j \cap A$ and $\mathcal{L}^{j+1} \cap B$. Analogously we can conclude that there are 2b(b-1) edges between $\mathcal{L}^j \cap B$ and $\mathcal{L}^{j+1} \cap A$. Summing up with the previous 4ab, for every $j \in \{1, ..., m-1\}$, we get $2a^2 - 2a + 2b^2 - 2b + 4ab = 2[(a+b)^2 - (a+b)]$ edges of the cut incident to \mathcal{L}^{j} minus the number of edges of the cut between \mathcal{L}^{j} and \mathcal{L}^{j-1} , as these get counted in \mathcal{L}^{j-1} . Recall that a+b=|X|+|Y|=n to see that this gives us 2n(n-1) edges. Finally, summing up for all $j \in \{1, \ldots, m-1\}$ and summing also the edges between link intervals in \mathcal{L}^m , we get that the number of edges of the cut incident to link intervals is equal to:

$$\sum_{j=1}^{m-1} 2n(n-1) + 4|X||Y| = n(n-1)(3n-2) + 4|X||Y|$$

Observe that $4(n-1) \leq 4|X||Y| \leq n^2$, and denote the value n(n-1)(3n-2) by β_4 .

Now, observe that it remains to count the number of edges of the cut-set between link intervals and intervals of type C, and between intervals of type C and vertex grained gadgets. We start with the latter. Given an edge $e_j = v_i v_{i'}$, with i < i', there are exactly n - i vertex grained gadgets covered by C_j^1, C_j^2 , and n - i' vertex grained gadgets covered by C_j^3, C_j^4 . Together with the q edges between each of these intervals of type C and the corresponding vertex gadgets (namely, \mathcal{H}_i^j and $\mathcal{H}_{i'}^j$), we get a total of 2(n - i)(p + q) + 2(n - i')(p + q) + 4q. Even though we cannot give a precise value below, observe that this value can be exactly computed during the construction. The upper bound is given just to make it explicit that this is a polynomial function. Also, below, for $e_j = v_i v_{i'}$, the value ℓ_j denotes i and r_j denotes i'.

$$\beta_{5} = \sum_{j=1}^{m} [2(n-r_{j})(p+q) + 2(n-\ell_{j})(p+q) + 4q] \\ = \sum_{j=1}^{m} [4n(p+q) - 2(r_{j}+\ell_{j})(p+q) + 4q] \\ \leq 4m[n(p+q)+q].$$

Finally, we count the number of edges of the cut between link intervals and intervals of type C. This is the only part of the counting that will not be exact. Again, consider an edge $e_j = v_i v_{i'}$, and first consider the interval C_j^1 ; we will see that the arguments hold for C_j^2 , and that analogous arguments hold for C_j^3 , C_j^4 . Observe that C_j^1 intersects exactly the following link intervals: $L_{i''}^{2j-1}$ and $L_{i''}^{2j}$ for every $i'' \in \{1, \ldots, n\}$; and $L_{i''}^{2j-2}$ and $L_{i''}^{2j-3}$ for every $i'' \in \{i+1, \ldots, n\}$. This is a total of less than 4n link intervals. Because an analogous argument can be applied to C_j^2 , C_j^3 , C_j^4 , we get a total of 16n possible edges in the cut-set, for each value of j, totalling $16nm = 24n^2$.

Let $\beta = \sum_{i=1}^{5} \beta_i$, and $\gamma = \beta + 4(n-1)$. We now prove that $|E_G(X,Y)| \ge k$ if and only if $|E_{\mathbb{G}}(A,B)| \ge \gamma + (2q'+4)k$. We have proved that:

$$\overbrace{\beta+4(n-1)}^{\gamma} + (2q'+4)|E_G(X,Y)| \leq |E_{\mathbb{G}}(A,B)| \leq \beta + 25n^2 + (2q'+4)|E_G(X,Y)|.$$

If $|E_G(X,Y)| \ge k$, then by the first inequality we have that $|E_{\mathbb{G}}(A,B)| \ge \beta + 4(n-1) + (2q'+4)k$. On the other hand, if $|E_{\mathbb{G}}(A,B)| \ge \beta + 4(n-1) + (2q'+4)k$, then by the second inequality we have that $|E_G(X,Y)| \ge k - \frac{25n^2 - 4(n-1)}{2q'+4} \ge k - \frac{26n^2}{2q'+4}$. Since $q' \ge 13n^2$, we get that $|E_G(X,Y)| > k-1$.

To finish the proof that the reduction works, we simply need to choose appropriate values for p, q, p', q'. Recall all necessary conditions:

- For each (x, y)-grained gadget \mathcal{H} in \mathcal{M} , let t be the number of intervals in $\mathcal{M} \setminus \mathcal{H}$ intersecting \mathcal{H} , ℓ be the number of intervals in $\mathcal{M} \setminus \mathcal{LS}$ intersecting the left short intervals, and r be the number of intervals in $\mathcal{M} \setminus \mathcal{RS}$ intersecting the right short intervals. Then we want that ℓ and r are both odd, and that y > t(x/y - 1) and x > t + 2y (from Lemma 3);
- = q > 4n + p' + q' + 3 (from Lemma 5);
- $q > 3(2n^2 + 2n + q' + 2)$ (from Lemma 6); and

By Lemma 4, we know that in order for the values r, ℓ in the first item to be odd, it suffices to choose q, q' to be odd. Observe that $n \ge 4$ since G is a cubic graph. For a given edge gadget \mathcal{E}_j , we know that there are exactly 2n + 4 intervals in $\mathcal{M} \setminus \mathcal{E}_j$ intersecting it, namely the link intervals and intervals of type C in the *j*-th region. We could just choose $q' \in \{13n^2, 13n^2 + 1\}$ such that q' is odd and $p' = 26n^2 + 2n + 7$. In this case, we have p' > t + 2q' and $q' \ge 13n^2 >$ $(\frac{26n^2 + 2n + 7}{13n^2} - 1)(2n + 4) \ge (\frac{p'}{q'} - 1)t$, since for edge gadgets t = 2n + 4. Similarly, we choose $q \in \{42n^2 + 3n + 10, 42n^2 + 3n + 11\}$ such that q is odd and $p = 84n^2 + 8n + 29$. We now have p > t + 2q and $q \ge 42n^2 + 3n + 10 > (\frac{84n^2 + 8n + 29}{42n^2 + 3n + 10} - 1)(2n + 6) \ge (\frac{p}{q} - 1)t$, since for vertex gadgets t < 2n + 6.

To finish the proof of Theorem 1, it remains to prove that the interval count of our reduction graph is exactly four, which is done in the next subsection.

2.4 Proof of Theorem 1: Bounding the interval count

Consider a cubic graph G on n vertices and m = 3n/2 edges, and orderings π_V, π_E of the vertex set and edge set of G. Denote the triple (G, π_V, π_E) by \mathfrak{G} . First, we want to prove that the interval count of our constructed interval model $\mathcal{M}(\mathfrak{G})$ is at most 4. But observe that the construction of $\mathcal{M}(\mathfrak{G})$ is actually not unique, since the intervals are not uniquely defined; e.g., given such a model, one can obtain a model satisfying the same properties simply by adding $\epsilon > 0$ to all points defining the intervals. In what follows, we provide a construction of a uniquely defined interval model related to \mathfrak{G} that satisfies the desired conditions and has interval count 4.

Consider our constructed interval model $\mathcal{M}(\mathfrak{G})$, and for each $j \in \{1, \ldots, m\}$, denote by \mathcal{S}_j the set of intervals related to the *j*-th region, i.e., $\mathcal{S}_j = \mathcal{E}_j \cup \bigcup_{\ell=1}^4 C_j^\ell \cup \bigcup_{i=1}^n (\mathcal{H}_i^j \cup \{L_i^{2j} \cup L_i^{2j-1}\})$. We show how to accommodate \mathcal{S}_1 within the closed interval [0, 6n-2] in such a way that the same pattern can be adopted in the subsequent regions of $\mathcal{M}(\mathfrak{G})$ too, each time starting at multiples of 4n. More specifically, letting t = 4n, we will accommodate \mathcal{S}_j within $[t \cdot (j-1), 6n-2+t \cdot (j-1)]$. Assume $e_1 = v_i v_{i'}$, with i < i'. Below, we describe exactly which closed interval of the line corresponds to each interval $I \in \mathcal{S}_1$.

- For each $i \in \{1, \ldots, n\}$, the left long intervals of \mathcal{H}_i^1 are equal to $[2i-2, 2i-\frac{3}{2}]$ and the left short intervals are any choice of p distinct points within the open interval $(2i-2, 2i-\frac{3}{2})$, whereas the right long intervals of \mathcal{H}_i^1 are equal to $[2i-\frac{3}{2}, 2i-1]$ and the right short intervals are any choice of p distinct points within the open interval $(2i-\frac{3}{2}, 2i-1]$. Note that open intervals are used to locate the closed intervals of length zero, but that the short intervals themselves are not open.
- C_1^1 and C_1^2 are equal to [2i 1, 2i + 2n 2].
- C_1^3 and C_1^4 are equal to [2i' 1, 2i' + 2n 2].
- The left long intervals of \mathcal{E}_1 are equal to [2n, 4n-1].
- The left short intervals of \mathcal{E}_1 are any choice of p' distinct points in the open interval (2i + 2n 2, 2i' + 2n 2). Again, the open interval is used just to locate the closed intervals of length zero.
- The right long intervals of \mathcal{E}_1 are equal to $[4n 1, 4n \frac{1}{2}]$ and the right short intervals are any choice of p' distinct points within the corresponding open interval.
- For each $i \in \{1, \ldots, n\}$, intervals L_i^1, L_i^2 are equal to [2i 1, 4n + 2(i 1)].



Figure 5 The closed intervals in $S_1 \cup \bigcup_{i=1}^4 \mathcal{H}_i^2$ of a graph on 4 vertices. We consider e_1 to be equal to v_3v_4 . Each colour represents a different interval size. The short intervals are represented by the dots located inside the open (red) intervals. Vertical lines mark the endpoints of the intervals in $S_1 \cup \bigcup_{i=1}^4 \mathcal{H}_i^2$, while the green vertical line marks the beginning of the intervals in S_2 .

The suitable chosen lengths of the above defined closed intervals are (see Figure 5, where we denote by Λ the set of link intervals):

- 1. 0: short intervals of all grained gadgets (dots in Figure 5);
- 2. 1/2: left long and right long intervals of each \mathcal{H}_i^1 , and right long intervals of \mathcal{E}_1 (red intervals in Figure 5);
- 3. 2n-1: intervals C_1^1, \ldots, C_1^4 , and left long intervals of \mathcal{E}_1 (blue intervals in Figure 5);
- 4. 4n 1: intervals L_i^1 and L_i^2 , for every $i \in [n]$ (orange intervals in Figure 5).

Now, let $\mathcal{M}'(\mathfrak{G})$ be the interval model where each S_j is defined exactly as S_1 , except that we shift all the intervals to the right in a way that point 0 now coincides with point $t \cdot (j-1)$. More formally, an interval I in S_j corresponding to the copy of an interval $[\ell, r]$ in S_1 is defined as $[\ell + t \cdot (j-1), r + t \cdot (j-1)]$. Also, we assign the intervals in the (m+1)-th grained gadgets to be at the end of this model, using the same sizes of intervals as above; i.e., \mathcal{H}_i^{m+1} is within the interval $[2i-2+t \cdot m, 2i-1+t \cdot m]$.

We have shown above that $\mathcal{M}'(\mathfrak{G})$ has interval count 4. The following lemma shows that the above chosen intervals satisfy the properties imposed in Subsections 2.1 and 2.2 on our constructed interval model $\mathcal{M}(\mathfrak{G})$.

▶ Lemma 8. Let G be a cubic graph. Then, there exists an interval model $\mathcal{M}(\mathfrak{G})$ with interval count 4 for $\mathfrak{G} = (G, \pi_V, \pi_E)$, for every ordering π_V and π_E of the vertex set and edge set of G, respectively.

Proof. Denote $\mathcal{M}(\mathfrak{G})$ by \mathcal{M} . We need to prove that \mathcal{M} satisfies the conditions of our construction, namely:

- 1. For every $j \in \{1, ..., m\}$ and $i \in \{1, ..., n\}$, link intervals L_i^{2j}, L_i^{2j-1} weakly intersect \mathcal{H}_i^j to the right and weakly intersects \mathcal{H}_i^{j+1} to the left;
- 2. For every $j \in \{1, \ldots, m\}$ and $i, i' \in \{1, \ldots, n\}$, i < i', the grained gadget \mathcal{H}_i^j occurs strictly to the left of $\mathcal{H}_{i'}^j$;
- 3. For every $j \in \{1, \ldots, m\}$, grained gadget \mathcal{E}_j occurs strictly between the right endpoint of \mathcal{H}_n^j and the left endpoint of \mathcal{H}_1^{j+1} ; and
- 4. For every $e_j = v_i v_{i'} \in E(G)$, i < i', intervals C_j^1, C_j^2 weakly intersect \mathcal{H}_i^j to the right and \mathcal{E}_j to the left, while C_j^3, C_j^4 weakly intersect $\mathcal{H}_{i'}^j$ to the right and strongly intersect \mathcal{E}_j to the left.

By construction, we know that the right endpoint of \mathcal{H}_i^j is equal to 2i - 1 + t(j - 1), which is also equal to the left endpoints of L_i^{2j-1}, L_i^{2j} . Also, the left endpoint of \mathcal{H}_i^{j+1} is equal to 2i-2+tj, which is also equal to the right endpoints of L_i^{2j-1}, L_i^{2j} since t = 4n; hence Item 1 follows. As for Item 2, just note that the right endpoint of \mathcal{H}_i^j , which is equal to 2i - 1 + t(j-1), is strictly smaller than the left endpoint of $\mathcal{H}_{i'}^{j}$, which is equal to 2i' - 2 + t(j-1). Indeed, since $i' \geq i+1$, we get $2i'-2 \ge 2(i+1)-2 = 2i > 2i-1$. Now, observe that \mathcal{E}_j is contained in the closed interval $[2n+t(j-1), 4n-\frac{1}{2}+t(j-1)]$, that the right endpoint of \mathcal{H}_n^j is equal to 2n-1+t(j-1), and the left endpoint of \mathcal{H}_1^{j+1} is equal to tj = 4n + t(j-1). Item 3 thus follows. Finally, as we have seen, the right endpoint of \mathcal{H}_i^j is equal to 2i - 1 + t(j - 1), which is equal to the left endpoints of C_j^1, C_j^2 ; hence these weakly intersect \mathcal{H}_i^j to the right. Also, the left endpoint of \mathcal{E}_j is equal to 2n + t(j-1), while the right endpoint of C_j^1, C_j^2 is equal to 2(i-1) + 2n + t(j-1), and all the left short intervals of \mathcal{E}_j are contained in the open interval [2(i-1)+2n+t(j-1),2(i'-1)+2n+t(j-1)]. Therefore we get that C_j^1, C_j^2 weakly intersect \mathcal{E}_j to the left. Analogously, the right endpoint of $\mathcal{H}_{i'}^j$ is equal to 2i'-1+t(j-1), which is equal to the left endpoints of C_j^3, C_j^4 ; hence they weakly intersect $\mathcal{H}_{i'}^j$ to the right. Finally, the right endpoint of C_i^3, C_i^4 is equal to 2(i'-1) + 2n + t(j-1), and all the left short intervals of \mathcal{E}_j are contained in the open interval [2(i-1)+2n+t(j-1), 2(i'-1)+2n+t(j-1)]. Also, the left endpoint of the right long intervals of \mathcal{E}_j is equal to 4n - 1 + t(j-1), which is strictly bigger than 2(i'-1) + 2n + t(j-1) since $i' \leq n$. Therefore, C_j^3, C_j^4 strongly intersect \mathcal{E}_j to the left, finishing the proof of Item 4.

We have just shown that, for any orderings π_V and π_E , there exists a model $\mathcal{M}(\mathfrak{G})$ of interval count 4, where $\mathfrak{G} = (G, \pi_V, \pi_E)$. On the other hand, we prove in the remainder of this section that any graph isomorphic to $\mathbb{G}_{\mathcal{M}(\mathfrak{G})}$ has interval count at least 4. For this, we show that all such graphs contain as an induced subgraph a certain graph of interval count 4, which we denote by H_4 . Next, we define the family $\{H_k\}_{k\geq 2}$ and prove in a more general way that $\mathrm{ic}(H_k) = k$ for every $k \geq 2$.

Let $P_5 = (u_1, \ldots, u_5)$ be a path on 5 vertices. For every graph H', we let $P_5 \circ H'$ be the graph obtained from the disjoint union of P_5 with H' by making u_3 , the central vertex of P_5 , adjacent to every vertex of H'. In other words, $P_5 \circ H'$ is the graph with vertex set $V(P_5) \cup V(H')$ and edge

XX:14 Maximum cut on interval graphs of interval count four is NP-complete

set $E(P_5) \cup E(H') \cup \{u_3v \mid v \in V(H')\}$. Then, for every $k \ge 2$, we let H_k be the graph defined recursively as follows (see Figure 6): $H_2 = K_{1,3}$;

 $H_k = P_5 \circ H_{k-1}$ for k > 2.



Figure 6 Graph H_k for $k \ge 2$.

▶ Lemma 9. For every $k \ge 2$, $ic(H_k) = k$.

Proof. The proof is by induction on k. Since $H_2 = K_{1,3}$ and $ic(K_{1,3}) = 2$ c.f. [25], we obtain that the lemma holds for k = 2. As inductive hypothesis, suppose that $ic(H_{k'}) = k'$ for some $k' \ge 2$. We prove that $ic(H_{k'+1}) = k' + 1$.

First, note that, if $\mathcal{M}_{P_5} = \{I_1, \ldots, I_5\}$ is an interval model of a P_5 , with interval I_i representing vertex u_i , then the precedence relation among the intervals of I_1, \ldots, I_5 is either that of Figure 7 (i.e., I_1 precedes I_3 , which precedes I_5 , and I_2 precedes I_4), or the reverse of the order presented in the figure c.f. [25]. Let \mathcal{M} be an interval model of $H_{k'+1}$. Since $H_{k'+1}$ contains a P_5 as an induced subgraph, assume without loss of generality that $\mathcal{M} \supset \mathcal{M}_{P_5}$ and that, with respect to \mathcal{M} , I_1 precedes I_3 , I_3 precedes I_5 , and I_2 precedes I_4 . This implies that

$$\ell(I_3) \le r(I_2) < \ell(I_4) \le r(I_3). \tag{2}$$

By construction, the only vertex of P_5 which is adjacent to the vertices of $H_{k'}$ is its central vertex u_3 . Consequently, if $\mathcal{M}_{H_{k'}} \subset \mathcal{M}$ is the interval model of $H_{k'}$, then there cannot be any intersection between $\mathcal{M}_{H_{k'}}$ and $\mathcal{M}_{P_5} \setminus \{I_3\}$, i.e., $I' \cap I_i = \emptyset$ for each $I' \in \mathcal{M}_{H_{k'}}$ and each $i \in \{1, \ldots, 5\}$, with $i \neq 3$. Hence, it follows from (2) that

 $\min\{\ell(I') \mid I' \in \mathcal{M}_{H_{k'}}\} > r(I_2) \text{ and } \max\{r(I') \mid I' \in \mathcal{M}_{H_{k'}}\} < \ell(I_4).$

Figure 7 illustrates this fact. As a result, $I_3 \supset I'$ for every $I' \in \mathcal{M}_{H_{L'}}$. This, along with the

Figure 7 Interval model $\mathcal{M}_{H_{k'+1}}$ of $H_{k'+1}$.

inductive hypothesis that $ic(H_{k'}) = k'$, implies that $ic(H_{k'+1}) \ge k' + 1$. On the other hand, it is straightforward that $ic(H_{k'+1}) \le k' + 1$ (for instance, consider the model illustrated in Figure 7). Therefore, $ic(H_{k'+1}) = k' + 1$.

Now, we finally show that $\mathbb{G}_{\mathcal{M}(\mathfrak{G})}$ contains an H_4 as an induced subgraph. Since G is cubic, there exists an edge $e_j = (v_i, v_{i'}) \in E(G)$ such that 1 < i < i'. Let (see Figure 4):

- \blacksquare I_1 (resp. I_2) be a right short (resp. long) interval of \mathcal{H}_1^j ;
- I₃ be the link interval L_1^{2j-1} ;
- If I_4 (resp. I_5) be a left long (resp. short) interval of \mathcal{H}_1^{j+1} ;
- \blacksquare I'_1 (resp. I'_2) be a right short (resp. long) interval of \mathcal{H}^j_i ;
- \blacksquare I'_3 be the interval C^1_j ;
- I'_4 (resp. I'_5) be a left long (resp. short) interval of \mathcal{E}_j ;

- I_1, J_2 and J_3 be three left short intervals of \mathcal{H}_{i+1}^j ; and
- \blacksquare J be a left long interval of \mathcal{H}_{i+1}^j .

The interval graph related to the model comprised by such intervals is isomorphic to H_4 . More specifically, observe first that $\mathcal{J} = \{J, J_1, J_2, J_3\}$ models $K_{1,3}$. Then, notice that $\mathcal{P} = \{I_1, \ldots, I_5\}$ and $\mathcal{P}' = \{I'_1, \ldots, I'_5\}$ model paths on 5 vertices, in this order. Finally observe that I'_3 is adjacent to every $I \in \mathcal{J}$, while there are no edges between \mathcal{J} and $\mathcal{P}' \setminus \{I'_3\}$; hence, $\mathcal{J} \cup \mathcal{P}'$ is a model for H_3 . Similarly, I_3 is adjacent to every $I \in \mathcal{J} \cup \mathcal{P}'$, while there are no edges between $\mathcal{J} \cup \mathcal{P}'$ and $\mathcal{P} \setminus \{I_3\}$; hence $\mathcal{J} \cup \mathcal{P}' \cup \mathcal{P}$ is a model for H_4 . Therefore, $\mathbb{G}_{\mathcal{M}(\mathfrak{G})}$ has an H_4 as an induced subgraph, as we wanted to prove.

3 The interval count of Adhikary et al.'s construction

We provided in Section 2 a reduction from the MAXCUT problem having as input a cubic graph Ginto that of MAXCUT in an interval graph G' having $ic(G') \leq 4$. Although our reduction requires the choice of orderings π_V and π_E of respectively V(G) and E(G) in order to produce the resulting interval model, we have established that we are able to construct an interval model with interval count 4 regardless of the particular choices for π_V and π_E (Lemma 8). Our reduction was based on that of [1], strengthened in order to control the interval count of the resulting model.

This section is dedicated to discuss the interval count of the original reduction [1]. Although the interval count was not of concern in [1], in order to contrast the reduction found there with the one presented in this work, we investigate how interval count varies in the original reduction considering different vertex/edge orderings. First, we establish that the original reduction yields an interval model corresponding to a graph G' such that $ic(G') = O(\sqrt[4]{|V(G')|})$. Second, we exhibit an example of a cubic graph G for which a choice of π_V and π_E yields a model \mathcal{M}' with interval count $\Omega(\sqrt[4]{V(G')})$, proving that this bound is tight for some choices of π_V and π_E . For bridgeless cubic graphs, we are able in Lemma 10 to decrease the upper bound by a constant factor, but to the best of our knowledge $O(\sqrt[4]{|V(G')|})$ is the tightest upper bound. Before we go further analysing the interval count of the original reduction, it is worthy to note that a tight bound on the interval count of a general interval graph G as a function of its number of vertices n is still open. It is known that $ic(G) \leq \lfloor (n+1)/2 \rfloor$ and that there is a family of graphs G for which ic(G) = (n-1)/3 [7,17]. That is, the interval count of a graph can reach $\Theta(n)$.

In the original reduction, given a cubic graph G, an interval graph G' is defined through the construction of one of its models \mathcal{M} , described as follows:

- 1. let $\pi_V = (v_1, v_2, \ldots, v_n)$ and $\pi_E = (e_1, e_2, \ldots, e_m)$ be arbitrary orderings of V(G) and E(G), respectively;
- **2.** for each $v_i \in V(G)$, $e_j \in E(G)$, let $\mathcal{G}(v_i)$ and $\mathcal{G}(e_j)$ denote respectively a (p,q)-grained gadget and a (p', q')-grained gadget, where:
 - $= q = 200n^3 + 1, p = 2q + 7n, \text{ and}$ $= q' = 10n^2 + 1, p' = 2q' + 7n;$
- **3.** for each $v_k \in V(G)$, insert $\mathcal{G}(v_k)$ in \mathcal{M} such that $\mathcal{G}(v_i)$ is entirely to the left of $\mathcal{G}(v_j)$ if and only if i < j. For each $e_k \in E(G)$, insert $\mathcal{G}(e_k)$ in \mathcal{M} entirely to the right of $\mathcal{G}(v_n)$ and such that $\mathcal{G}(e_i)$ is entirely to the left of $\mathcal{G}(e_j)$ if and only if i < j;
- 4. for each $e_j = (v_i, v_{i'}) \in E(G)$, with i < i', four intervals $I_{i,j}^1, I_{i,j}^2, I_{i',j}^1, I_{i',j}^2$ are defined in \mathcal{M} , called *link* intervals, such that:
 - $I_{i,j}^1$ and $I_{i,j}^2$ (resp. $I_{i',j}^1$ and $I_{i',j}^2$) are true twin intervals that weakly intersect $\mathcal{G}(v_i)$ (resp. $\mathcal{G}(v_{i'})$) to the right;

= $I_{i,j}^1$ and $I_{i,j}^2$ (resp. $I_{i',j}^1$ and $I_{i',j}^2$) weakly intersect (resp. strongly intersect) $\mathcal{G}(e_j)$ to the left. By construction, therefore, $I_{i,j}^1$ and $I_{i,j}^2$ (resp. $I_{i',j}^1$ and $I_{i',j}^2$) cover all intervals in grained gadgets associated to a vertex v_{ℓ} with $\ell > i$ (resp. $\ell > i'$) or an edge e_{ℓ} with $\ell < j$.

Note that the number of intervals in \mathcal{M} is independent of what orderings we choose for the vertices and edges of G and, therefore, so is the number of vertices of G'. Let n' = |V(G')|. Since G is cubic,

XX:16 Maximum cut on interval graphs of interval count four is NP-complete

 $m = \frac{3n}{2}$. By construction,

$$n' = n(2p + 2q) + m(2p' + 2q') + 4m = 1200n^{4} + 90n^{3} + 25n^{2} + 21m^{4}$$

and thus $n = \Theta(\sqrt[4]{n'})$. Since the set of intervals covered by any link interval depends on π_V and π_E , distinct sequences yield distinct resulting graphs G' having distinct interval counts.

We show next that $ic(G') = O(\sqrt[4]{n'})$. Note that

- the intervals of all gadgets $\mathcal{G}(v_i)$ and $\mathcal{G}(e_j)$ can use only two interval lengths (one for all short intervals, another for all the long intervals);
- for each $e_j = v_i v_{i'} \in E(G)$, with i < i', both intervals $I_{i,j}^1$ and $I_{i,j}^2$ may be coincident in any model, and therefore may have the same length. The same holds for both intervals $I_{i',j}^1$ and $I_{i',j}^2$.

Therefore, $ic(G') \leq 2m + 2 = 3n + 2 = \Theta(\sqrt[4]{n'})$. Therefore, the NP-completeness result derived from the original reduction in [1] can be strengthened to state that MAXCUT is NP-complete for interval graphs G having interval count $O(\sqrt[4]{|V(G)|})$.

Second, we show that there is a resulting model \mathcal{M}' produced in the reduction, defined in terms of particular orderings π_V, π_E for which $ic(\mathcal{M}') = \Omega(\sqrt[4]{n'})$. Consider the cubic graph G depicted in Figure 8(a) which consists of an even cycle (v_1, v_2, \ldots, v_n) with the addition of the edges $(v_i, v_{i+\frac{n}{2}})$ for all $1 \leq i \leq n/2$. For the ordering $\pi_V = (v_n, v_{n-1}, \ldots, v_1)$ and any ordering π_E of the edges starting with the suborder $(v_1v_2, v_2v_3, \ldots, v_nv_1)$ (i.e. starting with the edges of the cycle), the reduction yields a model \mathcal{M}' for which there is a chain $I_{1,1}^1 \subset I_{2,2}^1 \subset \cdots \subset I_{n,n}^1$ of nested intervals (see Figure 8(b)), which shows that $ic(\mathcal{M}') \geq n$, and thus $ic(\mathcal{M}') = \Omega(\sqrt[4]{n'})$.



Figure 8 (a) A cubic graph G, and (b) a chain of nested intervals in the model \mathcal{M}' .

It can be argued from the proof of NP-completeness for MAXCUT when restricted to cubic graphs [2] that the constructed cubic graph may be assumed to have no bridges. This fact was not used in the original reduction of [1]. In an attempt to obtain a model \mathcal{M} having fewer lengths for bridgeless cubic graphs, we have derived Lemma 10. Although the number of lengths in this new upper bound has decreased by the constant factor of 4/9, it is still $\Theta(n) = \Theta(\sqrt[4]{n'})$.

▶ Lemma 10. Let G be a cubic bridgeless graph with n = |V(G)|. There exist particular orderings π_V of V(G) and π_E of E(G) such that:

- 1. there is a resulting model \mathcal{M} produced in the original reduction of MAXCUT such that $ic(\mathcal{M}) \leq \frac{4n}{3} + 3$.
- 2. for all such resulting models \mathcal{M} , we have that $ic(\mathcal{M}) \geq 5$ if G is not a Hamiltonian graph.

Proof. Let G be a cubic bridgeless graph with $V(G) = \{v_1, v_2, \ldots, v_n\}$. By Petersen's theorem, every cubic bridgeless graph contains a perfect matching, so G admits a perfect matching M. Let $H = G \setminus M$. Therefore, H is 2-regular and, therefore, H consists of a disjoint union of cycles C_1, C_2, \ldots, C_k , for some $k \ge 1$. For all $1 \le i \le k$, let $\pi_V^i = v_1^i, v_2^i, \ldots, v_{k_i}^i$ be an ordering of the vertices of C_i , with $k_i = |C_i|$, such that $(v_j^i, v_{j+1}^i) \in E(C_i)$ for all $1 \le j \le k_i$, where $v_{k_i+1}^i = v_1^i$. Let π_E^i be the ordering $(v_1^i, v_2^i), (v_2^i, v_3^i), \ldots, (v_{k_i-1}^i, v_{k_i}^i), (v_1^i, v_{k_i}^i)$ for all $1 \le i \le k$. Let π_M be any ordering of the edges of M such that $(v_i, v_r) < (v_j, v_s)$ in π_M only if $v_i < v_j$ in π_V . Finally, let π_V be the ordering of V(G) obtained from the concatenation of the orderings $\pi_V^1, \pi_V^2, \ldots, \pi_K^k, \pi_M$.

In order to prove (2.), assume G is not a Hamiltonian graph. Therefore k > 1. Observe that there is the following chain of nested intervals $I_1 \subset I_2 \subset I_3 \subset I_4 \subset I_5$, where

- I is the leftmost interval in $\mathcal{RS}(\mathcal{G}(v_3^2))$,
- $\blacksquare I_2 \text{ is an interval in } \mathcal{RL}(\mathcal{G}(v_3^2)),$
- I₃ is a link interval corresponding to both $\mathcal{G}(v_2^2)$ and $\mathcal{G}(v_1^2v_2^2)$,
- If I_4 is a link interval corresponding to both $\mathcal{G}(v_1^2)$ and $\mathcal{G}(v_1^2 v_{k_2}^2)$, and
- Is is a link interval corresponding to both $\mathcal{G}(v_1^1)$ and $\mathcal{G}(e)$, where e is the edge of M incident to v_1^1 ,

since $\ell(I_5) < \ell(I_4) < \ell(I_3) < \ell(I_2) < \ell(I_1) < r(I_1) < r(I_2) < r(I_3) < r(I_4) < r(I_5)$. Thus, for all such resulting models \mathcal{M} , we have that $ic(\mathcal{M}) \ge 5$.

In order to prove (1.), we show that there exists an interval model \mathcal{M} , produced by the original reduction of MAXCUT considering orderings π_V and π_E , such that $ic(\mathcal{M}) \leq \frac{4n}{3} + 3$, where n = |V(G)|. Let L_1 be the set of all link intervals of the grained gadgets corresponding to edges of \mathcal{M} , that is, $L_1 = \{I_{i,k}^1, I_{i,k}^2, I_{j,k}^1, I_{j,k}^2 : e_k = (i, j) \in \mathcal{M}\}$. Moreover, let L_2 be the set of all link intervals of the grained gadgets corresponding to edges of \mathcal{M} , that is, that is, $L_1 = \{I_{i,k}^1, I_{i,k}^2, I_{j,k}^1, I_{j,k}^2 : e_k = (i, j) \in \mathcal{M}\}$. Moreover, let L_2 be the set of all link intervals of the grained gadgets corresponding to the edges $(v_1^i, v_{k_i}^i)$ of C_i and the vertex v_1^i for all $1 \leq i \leq k$, that is, Note that $|L_2| = k \leq n/3$ and $|L_1| = 4 \cdot |\mathcal{M}| = 2n$. Let $L = L_1 \cup L_2$. Let $\mathcal{M}' = \mathcal{M} \setminus L$. We claim that $ic(\mathcal{M}') \leq 3$. Since each pair of true twins $I_{j,k}^1, I_{j,k}^2$ and $I_{i,k}^1, I_{i,k}^2$ in L_1 can have the same length in \mathcal{M} , it follows from this claim that $ic(\mathcal{M}) \leq |L_1| + \frac{|L_2|}{2} + ic(\mathcal{M}') \leq \frac{n}{3} + n + 3 = \frac{4n}{3} + 3$, holding the result. It remains to show that the claim indeed holds.

To prove the claim, let \mathcal{M}'' be the interval model obtained from \mathcal{M}' by removing all intervals corresponding to the grained gadgets (or, in other words, by keeping only the intervals corresponding to link intervals). It is easily seen that \mathcal{M}'' is a proper interval model, that is, no interval is properly contained in another. Therefore, the interval graph corresponding to \mathcal{M}'' is a proper interval graph and \mathcal{M}'' can be modified so that their intervals have all a single length. Since it is possible to bring all grained gadgets back to \mathcal{M}'' using two more lengths, we have that $ic(\mathcal{M}') \leq 3$, as claimed.

As a concluding remark, we note that the interval count of the interval model \mathcal{M} produced in the original reduction is highly dependent on the assumed orderings of V(G) and E(G), and may achieve $ic(\mathcal{M}) = \Omega(\sqrt[4]{n'})$. The model \mathcal{M}' produced in our reduction enforces that $ic(\mathcal{M}') = 4$ which is invariant for any such orderings. On the perspective of the recognition problem for interval graphs with interval count k, with fixed $k \geq 2$, for which very little is known, our NP-completeness result on a class of bounded interval count graphs is also of interest.

Acknowledgements

We thank Vinicius F. Santos who shared Reference [1], and anonymous referees for many valuable suggestions, including improving the interval count from 5 to 4.

Data availability statement

Data sharing not applicable to this article as no datasets were generated or analysed during the current study.

— References

- Ranendu Adhikary, Kaustav Bose, Satwik Mukherjee, and Bodhayan Roy. Complexity of maximum cut on interval graphs. In Kevin Buchin and Éric Colin de Verdière, editors, 37th International Symposium on Computational Geometry, SoCG 2021, June 7-11, 2021, Buffalo, NY, USA (Virtual Conference), volume 189 of LIPIcs, pages 7:1–7:11, Dagstuhl, Germany, 2021. Schloss Dagstuhl - Leibniz-Zentrum für Informatik. doi:10.4230/LIPIcs.SoCG.2021.7.
- 2 Piotr Berman and Marek Karpinski. On some tighter inapproximability results (extended abstract). In Jirí Wiedermann, Peter van Emde Boas, and Mogens Nielsen, editors, Automata, Languages and Programming, 26th International Colloquium, ICALP'99, Prague, Czech Republic, July 11-15, 1999, Proceedings, volume 1644 of Lecture Notes in Computer Science, pages 200–209, Berlin, Heidelberg, 1999. Springer. doi:10.1007/3-540-48523-6_17.

XX:18 Maximum cut on interval graphs of interval count four is NP-complete

- 3 Hans L. Bodlaender, Celina M. H. de Figueiredo, Marisa Gutierrez, Ton Kloks, and Rolf Niedermeier. Simple max-cut for split-indifference graphs and graphs with few p₄'s. In Celso C. Ribeiro and Simone L. Martins, editors, *Experimental and Efficient Algorithms, Third International Workshop, WEA 2004, Angra dos Reis, Brazil, May 25-28, 2004, Proceedings, volume 3059 of Lecture Notes in Computer Science*, pages 87–99, Berlin, Heidelberg, 2004. Springer. doi:10.1007/978-3-540-24838-5_7.
- 4 Hans L. Bodlaender, Ton Kloks, and Rolf Niedermeier. SIMPLE MAX-CUT for unit interval graphs and graphs with few p₄s. *Electron. Notes Discret. Math.*, 3:19–26, 1999. doi:10.1016/ S1571-0653(05)80014-9.
- 5 J. Adrian Bondy and Uppaluri S. R. Murty. *Graph Theory*. Graduate Texts in Mathematics. Springer, New York, 2008. doi:10.1007/978-1-84628-970-5.
- 6 Arman Boyaci, Tínaz Ekim, and Mordechai Shalom. A polynomial-time algorithm for the maximum cardinality cut problem in proper interval graphs. *Inf. Process. Lett.*, 121:29–33, 2017. doi:10.1016/j.ipl.2017.01.007.
- 7 Márcia R. Cerioli, Fabiano de S. Oliveira, and Jayme Luiz Szwarcfiter. The interval count of interval graphs and orders: a short survey. J. Braz. Comput. Soc., 18(2):103–112, 2012. doi:10.1007/s13173-011-0047-1.
- 8 Márcia R. Cerioli, Fabiano S. Oliveira, and Jayme L. Szwarcfiter. On counting interval lengths of interval graphs. *Discret. Appl. Math.*, 159(7):532-543, 2011. URL: https://www. sciencedirect.com/science/article/pii/S0166218X10002581, doi:https://doi.org/10. 1016/j.dam.2010.07.006.
- Dibyayan Chakraborty, Sandip Das, Florent Foucaud, Harmender Gahlawat, Dimitri Lajou, and Bodhayan Roy. Algorithms and complexity for geodetic sets on planar and chordal graphs. In Yixin Cao, Siu-Wing Cheng, and Minming Li, editors, 31st International Symposium on Algorithms and Computation, ISAAC 2020, December 14-18, 2020, Hong Kong, China (Virtual Conference), volume 181 of LIPIcs, pages 7:1-7:15, Dagstuhl, Germany, 2020. Schloss Dagstuhl Leibniz-Zentrum für Informatik. doi:10.4230/LIPIcs.ISAAC.2020.7.
- 10 Johanne Cohen, Fedor V. Fomin, Pinar Heggernes, Dieter Kratsch, and Gregory Kucherov. Optimal linear arrangement of interval graphs. In Rastislav Kralovic and Pawel Urzyczyn, editors, Mathematical Foundations of Computer Science 2006, 31st International Symposium, MFCS 2006, Stará Lesná, Slovakia, August 28-September 1, 2006, Proceedings, volume 4162 of Lecture Notes in Computer Science, pages 267–279, Berlin, Heidelberg, 2006. Springer. doi:10.1007/11821069_24.
- 11 Derek G. Corneil, Hiryoung Kim, Sridhar Natarajan, Stephan Olariu, and Alan P. Sprague. Simple linear time recognition of unit interval graphs. *Inf. Process. Lett.*, 55(2):99–104, 1995. doi:10.1016/0020-0190(95)00046-F.
- 12 Celina M. H. de Figueiredo, Alexsander Andrade de Melo, Fabiano de S. Oliveira, and Ana Silva. Maximum cut on interval graphs of interval count five is NP-complete. CoRR, abs/2012.09804, 2020. arXiv:2012.09804.
- 13 Celina M. H. de Figueiredo, Alexsander Andrade de Melo, Fabiano de S. Oliveira, and Ana Silva. Maximum cut on interval graphs of interval count four is NP-complete. In Filippo Bonchi and Simon J. Puglisi, editors, 46th International Symposium on Mathematical Foundations of Computer Science, MFCS 2021, August 23-27, 2021, Tallinn, Estonia, volume 202 of LIPIcs, pages 38:1–38:15, Dagstuhl, Germany, 2021. Schloss Dagstuhl Leibniz-Zentrum für Informatik. doi:10.4230/LIPIcs.MFCS.2021.38.
- 14 Celina M. H. de Figueiredo, João Meidanis, and Célia Picinin de Mello. A linear-time algorithm for proper interval graph recognition. Inf. Process. Lett., 56(3):179–184, 1995. doi:10.1016/0020-0190(95)00133-W.
- 15 Celina M.H. de Figueiredo, Alexsander A. de Melo, Diana Sasaki, and Ana Silva. Revising Johnson's table for the 21st century. *Discret. Appl. Math.*, 2021. URL: https://www.sciencedirect.com/science/article/pii/S0166218X21002109, doi:https://doi.org/10.1016/j.dam.2021.05.021.

- 16 Tínaz Ekim, Aysel Erey, Pinar Heggernes, Pim van 't Hof, and Daniel Meister. Computing minimum geodetic sets of proper interval graphs. In David Fernández-Baca, editor, LATIN 2012: Theoretical Informatics - 10th Latin American Symposium, Arequipa, Peru, April 16-20, 2012. Proceedings, volume 7256 of Lecture Notes in Computer Science, pages 279–290, Berlin, Heidelberg, 2012. Springer. doi:10.1007/978-3-642-29344-3_24.
- Peter C. Fishburn. Interval graphs and interval orders. *Discret. Math.*, 55(2):135–149, 1985.
 doi:10.1016/0012-365X(85)90042-1.
- 18 M. R. Garey, David S. Johnson, and Larry J. Stockmeyer. Some simplified NP-complete graph problems. *Theor. Comput. Sci.*, 1(3):237–267, 1976. doi:10.1016/0304-3975(76)90059-1.
- Yuan Jinjiang and Zhou Sanming. Optimal labelling of unit interval graphs. Applied Math., 10(3):337–344, sep 1995. doi:10.1007/bf02662875.
- 20 David S. Johnson. The NP-completeness column: An ongoing guide. J. Algorithms, 6(3):434–451, 1985. doi:10.1016/0196-6774(85)90012-4.
- 21 Pavel Klavík, Yota Otachi, and Jirí Sejnoha. On the classes of interval graphs of limited nesting and count of lengths. *Algorithmica*, 81(4):1490–1511, 2019. doi:10.1007/s00453-018-0481-y.
- 22 Jan Kratochvíl, Tomás Masarík, and Jana Novotná. U-bubble model for mixed unit interval graphs and its applications: The maxcut problem revisited. In Javier Esparza and Daniel Král', editors, 45th International Symposium on Mathematical Foundations of Computer Science, MFCS 2020, August 24-28, 2020, Prague, Czech Republic, volume 170 of LIPIcs, pages 57:1–57:14, Dagstuhl, Germany, 2020. Schloss Dagstuhl - Leibniz-Zentrum für Informatik. doi:10.4230/LIPIcs.MFCS.2020.57.
- 23 Dániel Marx. A short proof of the NP-completeness of minimum sum interval coloring. Oper. Res. Lett., 33(4):382–384, 2005. doi:10.1016/j.orl.2004.07.006.
- 24 Sara Nicoloso, Majid Sarrafzadeh, and X. Song. On the sum coloring problem on interval graphs. Algorithmica, 23(2):109–126, 1999. doi:10.1007/PL00009252.
- 25 FS Roberts. Indifference graphs, f. harary (ed.), proof techniques in graph theory, 1969.