

# A COMPLEX ANALOGUE OF TODA'S THEOREM

SAUGATA BASU

ABSTRACT. Toda [24] proved in 1989 that the (discrete) polynomial time hierarchy,  $\mathbf{PH}$ , is contained in the class  $\mathbf{P}^{\#\mathbf{P}}$ , namely the class of languages that can be decided by a Turing machine in polynomial time given access to an oracle with the power to compute a function in the counting complexity class  $\#\mathbf{P}$ . This result which illustrates the power of counting is considered to be a seminal result in computational complexity theory. An analogous result (with a compactness hypothesis) in the complexity theory over the reals (in the sense of Blum-Shub-Smale real machines [4]) was proved in [2]. Unlike Toda's proof in the discrete case, which relied on sophisticated combinatorial arguments, the proof in [2] is topological in nature in which the properties of the topological join is used in a fundamental way. However, the constructions used in [2] were semi-algebraic – they used real inequalities in an essential way and as such do not extend to the complex case. In this paper, we extend the techniques developed in [2] to the complex projective case. A key role is played by the complex join of quasi-projective complex varieties. As a consequence we obtain a complex analogue of Toda's theorem. The results contained in this paper, taken together with those contained in [2], illustrate the central role of the Poincaré polynomial in algorithmic algebraic geometry, as well as, in computational complexity theory over the complex and real numbers – namely, the ability to compute it efficiently enables one to decide in polynomial time all languages in the (compact) polynomial hierarchy over the appropriate field.

## 1. INTRODUCTION AND MAIN RESULTS

**1.1. History and Background.** The primary motivation for this paper comes from classical (i.e. discrete) computational complexity theory. In classical complexity theory, there is a seminal result due to Toda [24] linking the complexity of counting with that of deciding sentences with a fixed number of quantifier alternations.

More precisely, Toda's theorem gives the following inclusion (see Section 1.3.1 below or refer to [17] for precise definitions of the complexity classes appearing in the theorem).

**Theorem 1.1** (Toda [24]).

$$\mathbf{PH} \subset \mathbf{P}^{\#\mathbf{P}}.$$

In other words, any language in the (discrete) polynomial hierarchy can be decided by a Turing machine in polynomial time, given access to an oracle with the power to compute a function in  $\#\mathbf{P}$ .

---

*Date:* March 1, 2010.

*1991 Mathematics Subject Classification.* Primary 14F25, 14Q15; Secondary 68Q15.

*Key words and phrases.* Polynomial hierarchy, Betti numbers, constructible sets, Toda's theorem.

The author was supported in part by an NSF grant CCF-0915954.

*Remark 1.2.* The proof of Theorem 1.1 in [24] is quite non-trivial. While it is obvious that the classes  $\mathbf{P}$ ,  $\mathbf{NP}$ ,  $\mathbf{coNP}$  are contained in  $\mathbf{P}^{\#\mathbf{P}}$ , the proof for the higher levels of the polynomial hierarchy is quite intricate and proceeds in two steps: first proving that the  $\mathbf{PH} \subset \mathbf{BP} \cdot \oplus \cdot \mathbf{P}$  (using previous results of Schönning [19], and Valiant and Vazirani [25]), and then showing that  $\mathbf{BP} \cdot \oplus \cdot \mathbf{P} \subset \mathbf{P}^{\#\mathbf{P}}$ . Aside from the obvious question about what should be a proper analogue of the complexity class  $\#\mathbf{P}$  over the reals or complex numbers, because of the presence of the intermediate complexity class in the proof, there seems to be no direct way of extending such a proof to real or complex complexity classes in the sense of Blum-Shub-Smale model of computation [4, 21]. The proof of the main theorem (Theorem 2.1) of this paper, which can be seen as a complex analogue of Theorem 1.1, proceeds along completely different lines and is mainly topological in nature.

In the late eighties Blum, Shub and Smale [4, 21] introduced the notion of Turing machines over more general fields, thereby generalizing the classical problems of computational complexity theory such as  $\mathbf{P}$  vs.  $\mathbf{NP}$  to corresponding problems over arbitrary fields (such as the real, complex,  $p$ -adic numbers etc.) If one considers languages accepted by a Blum-Shub-Smale machine over a finite field one recovers the classical notions of discrete complexity theory. Over the last two decades there has been a lot of research activity towards proving real as well as complex analogues of well known theorems in discrete complexity theory. The first steps in this direction were taken by the authors Blum, Shub, and Smale (henceforth B-S-S) themselves, when they proved the  $\mathbf{NP}_C$ -completeness of the problem of deciding whether a systems of  $n + 1$  polynomial equations in  $n$  variables of has a solution (in affine space) (this is the complex analogue of Cook-Levin's theorem that the satisfiability problem is  $\mathbf{NP}$ -complete in the discrete case), and subsequently through the work of several researchers (Koiran, Bürgisser, Cucker, Meer to name a few) a well-established complexity theory over the reals as well as complex numbers have been built up, which mirrors closely the discrete case.

Indeed, one of the main attractions of the Blum-Shub-Smale computational model is that it provides a framework to prove complexity results over more general structures, than just finite fields, with the hope that such results will help to unravel the algebro-geometric underpinnings of the basic separation questions amongst complexity classes. It is also often interesting to investigate complex (as well as real) analogues of results in discrete complexity theory, because doing so reveals underlying geometric and topological phenomenon not visible in the discrete case. From this viewpoint it is quite natural to seek complex (as well as real) analogues of Toda's theorem; and as we will see in this paper (see also [2]), Toda's theorem properly interpreted over the real and complex numbers gives an unexpected connection between two important but distinct strands of algorithmic algebraic geometry – namely, *decision problems* involving quantifier elimination on one hand, and the problems of *computing topological invariants* of constructible sets on the other. Indeed, the original result of Toda, together with its real and complex counter-parts seem to suggest a deeper connection of a model-theoretic nature, between the problems of efficient quantifier-elimination and efficient computation of certain discrete invariants of definable sets in a structure, which might be an interesting problem on its own to explore further in the future.

**1.2. Recent Work.** There has been a large body of recent research on obtaining appropriate real (as well as complex) analogues of results in discrete complexity

theory, especially those related to counting complexity classes (see [16, 6, 8, 7]). In [2] a real analogue of Toda's theorem was proved (with a compactness hypothesis). In this paper we prove a similar result in the complex case. Even though the basic approach is similar in both cases, the topological tools in the complex case are different enough to merit a separate treatment. This is elaborated further in the next section (the main difficulty in extending the real arguments in [2] to the complex case is that we can no longer use inequalities in our constructions).

**1.3. Definitions of complexity classes.** In order to formulate our result it is first necessary to define precisely complex counter-parts of the discrete polynomial time hierarchy  $\mathbf{PH}$  and the discrete complexity class  $\#\mathbf{P}$ , and this is what we do next.

**1.3.1. Complex counter-parts of  $\mathbf{PH}$  and  $\#\mathbf{P}$ .** For the rest of the paper  $\mathbb{C}$  will denote an algebraically closed field of characteristic zero (there is no essential loss in assuming that  $\mathbb{C} = \mathbb{C}$ ) (indeed by a transfer argument it suffices to prove all our results in this case). By a *complex machine* we will mean a machine in the sense of Blum-Shub-Smale [4] over the ground field  $\mathbb{C}$ .

*Notational convention.* Since in what follows we will be forced to deal with multiple blocks of variables in our formulas, we follow a notational convention by which we denote blocks of variables by bold letters with superscripts (e.g.  $\mathbf{X}^i$  denotes the  $i$ -th block), and we use non-bold letters with subscripts to denote single variables (e.g.  $X_j^i$  denotes the  $j$ -th variable in the  $i$ -th block). We use  $\mathbf{x}^i$  to denote a specific value of the block of variables  $\mathbf{X}^i$ . We will call a quantifier-free first-order formula (in the language of fields),  $\phi(\mathbf{X}^1; \dots; \mathbf{X}^\omega)$ , having several blocks of variables  $(\mathbf{X}^1, \dots, \mathbf{X}^\omega)$  to be *multi-homogeneous* if each polynomial appearing in it is multi-homogeneous in the blocks of variables  $(\mathbf{X}^1, \dots, \mathbf{X}^\omega)$  and such that  $\phi$  is satisfied whenever any one of the blocks  $\mathbf{X}^i = 0$ . Clearly such a formula defines a *constructible* subset of  $\mathbb{P}_{\mathbb{C}}^{k_1} \times \dots \times \mathbb{P}_{\mathbb{C}}^{k_\omega}$  where the block  $\mathbf{X}^i$  is assumed to have  $k_i + 1$  variables. If  $\omega = 1$ , that is there is only one block of variables, then we call  $\phi$  a *homogeneous* formula.

**Notation 1.3** (Realization). More generally, let

$$\Phi(\mathbf{X}^1; \dots; \mathbf{X}^\sigma) \stackrel{\text{def}}{=} (\mathbf{Q}_1 \mathbf{Y}^1) \dots (\mathbf{Q}_\omega \mathbf{Y}^\omega) \phi(\mathbf{X}^1; \dots; \mathbf{X}^\sigma; \mathbf{Y}^1; \dots; \mathbf{Y}^\omega)$$

be a (quantified) multi-homogeneous formula, with  $\mathbf{Q}_i \in \{\exists, \forall\}$ ,  $1 \leq i \leq \omega$ ,  $\phi$  a quantifier-free multi-homogeneous formula, and  $\mathbf{X}^i$  (resp.  $\mathbf{Y}^j$ ) is a block of  $k_i + 1$  (resp.  $\ell_j + 1$ ) variables. We denote by  $\mathcal{R}(\Phi) \subset \mathbb{P}_{\mathbb{C}}^{k_1} \times \dots \times \mathbb{P}_{\mathbb{C}}^{k_\sigma}$  the constructible set which is the *realization* of the formula  $\Phi$ ; i.e.,

$$\begin{aligned} \mathcal{R}(\Phi(\mathbf{X})) &= \{(\mathbf{x}^1, \dots, \mathbf{x}^\sigma) \in \mathbb{P}_{\mathbb{C}}^{k_1} \times \dots \times \mathbb{P}_{\mathbb{C}}^{k_\sigma} \mid \\ &(\mathbf{Q}_1 \mathbf{y}^1 \in \mathbb{P}_{\mathbb{C}}^{\ell_1}) \dots (\mathbf{Q}_\omega \mathbf{y}^\omega \in \mathbb{P}_{\mathbb{C}}^{\ell_\omega}) \phi(\mathbf{x}^1; \dots; \mathbf{x}^\sigma; \mathbf{y}^1; \dots; \mathbf{y}^\omega)\}. \end{aligned}$$

Sometimes, in order to emphasize the block structure in a multi-homogeneous formula, we will write the quantifications as  $(\exists \mathbf{Y} \in \mathbb{P}_{\mathbb{C}}^\ell)$  (resp.  $(\forall \mathbf{Y} \in \mathbb{P}_{\mathbb{C}}^\ell)$ ) instead of just  $(\exists \mathbf{Y})$  (resp.  $(\forall \mathbf{Y})$ ). This is purely notational and does not affect the syntax of the formula.

We say that two multi-homogeneous formulas,  $\Phi$  and  $\Psi$ , are *equivalent* if  $\mathcal{R}(\Phi) = \mathcal{R}(\Psi)$ . Clearly, equivalent multi-homogeneous formulas must have identical number of blocks of free variables, and the corresponding block sizes must also be equal.

Since the notion of multi-homogeneous formulas might look a bit unusual at first glance from the point of view of logic, we illustrate below how to homogenize non-homogeneous formulas by considering the following simple example (which is a building block for the “repeated squaring” technique used to prove doubly exponential lower bounds for (real) quantifier elimination [9]).

**Example 1.4.** Let  $\Phi(X)$  be the following (existentially) quantified non-homogeneous formula expressing the fact, that  $X^4 = 1$ .

$$\Phi(X) \stackrel{\text{def}}{=} \exists Y (Y^2 - 1 = 0) \wedge (Y - X^2 = 0).$$

A multi-homogeneous version of the same formula is given by:

$$\Phi^h(X_0 : X_1) \stackrel{\text{def}}{=} \exists ((Y_0 : Y_1) \in \mathbb{P}_C^1) (Y_1^2 - Y_0^2 = 0) \wedge (X_0^2 Y_1 - X_1^2 Y_0 = 0).$$

Notice that the quantifier-free bi-homogeneous formula

$$\Psi^h(X_0 : X_1; Y_0 : Y_1) \stackrel{\text{def}}{=} (Y_1^2 - Y_0^2 = 0) \wedge (X_0^2 Y_1 - X_1^2 Y_0 = 0)$$

defines a constructible subset of  $\mathbb{P}_C^1 \times \mathbb{P}_C^1$ , and that the affine part of the constructible subset of  $\mathbb{P}_C^1$  defined by  $\Phi^h$  coincides with the constructible subset of  $C^1$  defined by  $\Phi(X)$ .

**1.4. Complex analogue of PH.** We recall the definition of the polynomial hierarchy over  $C$ . It mirrors the discrete case very closely (see [23]).

**Definition 1.5** (The class  $\mathbf{P}_C$ ). Let  $k(n)$  be any polynomial in  $n$ . A sequence

$$\left( T_n \subset C^{k(n)} \right)_{n>0}$$

of constructible subsets is said to belong to the class  $\mathbf{P}_C$  if there exists a B-S-S machine  $M$  over  $C$  (see [4, 3]), such that for all  $\mathbf{x} \in C^{k(n)}$ , the machine  $M$  tests membership of  $\mathbf{x}$  in  $T_n$  in time bounded by a polynomial in  $n$ .

**Definition 1.6** (The classes  $\Sigma_{C,\omega}$  and  $\Pi_{C,\omega}$ ). Let  $k(n), k_1(n), \dots, k_\omega(n)$  be polynomials in  $n$ . A sequence

$$\left( S_n \subset C^{k(n)} \right)_{n>0}$$

of constructible subsets is said to be in the complexity class  $\Sigma_{C,\omega}$ , if for each  $n > 0$ , the constructible set  $S_n$  is described by a first order formula

$$(1.1) \quad (Q_1 \mathbf{Y}^1) \cdots (Q_\omega \mathbf{Y}^\omega) \phi_n(X_1, \dots, X_{k(n)}, \mathbf{Y}^1, \dots, \mathbf{Y}^\omega),$$

with  $\phi_n$  a quantifier free formula in the first order theory of  $C$ , and for each  $i, 1 \leq i \leq \omega$ ,  $\mathbf{Y}^i = (Y_1^i, \dots, Y_{k_i(n)}^i)$  is a block of  $k_i(n)$  variables,  $Q_i \in \{\exists, \forall\}$ , with  $Q_j \neq Q_{j+1}$ ,  $1 \leq j < \omega$ ,  $Q_1 = \exists$ , and the sequence

$$\left( T_n \subset C^{k(n)} \times C^{k_1(n)} \times \dots \times C^{k_\omega(n)} \right)_{n>0}$$

of constructible subsets defined by the quantifier-free formulas  $(\phi_n)_{n>0}$  belongs to the class  $\mathbf{P}_C$ .

Similarly, the complexity class  $\Pi_{C,\omega}$  is defined as in Definition 1.6, with the exception that the alternating quantifiers in (1.1) start with  $Q_1 = \forall$ .

Since, adding an additional block of quantifiers on the outside (with new variables) does not change the set defined by a quantified formula we have the following inclusions:

$$\Sigma_{C,\omega} \subset \Pi_{C,\omega+1}, \text{ and } \Pi_{C,\omega} \subset \Sigma_{C,\omega+1}.$$

Note that by the above definition the class  $\Sigma_{C,0} = \Pi_{C,0}$  is the familiar class  $\mathbf{P}_C$ , the class  $\Sigma_{C,1} = \mathbf{NP}_C$  and the class  $\Pi_{C,1} = \mathbf{co-NP}_C$ .

**Definition 1.7** (Complex polynomial hierarchy). The complex polynomial time hierarchy is defined to be the union

$$\mathbf{PH}_C \stackrel{\text{def}}{=} \bigcup_{\omega \geq 0} (\Sigma_{C,\omega} \cup \Pi_{C,\omega}) = \bigcup_{\omega \geq 0} \Sigma_{C,\omega} = \bigcup_{\omega \geq 0} \Pi_{C,\omega}.$$

As in the real case studied in [2] for technical reasons we need to restrict to compact constructible sets. However, unlike in [2] where the compact languages consisted of closed semi-algebraic subsets of spheres, in this paper we consider closed subsets of projective spaces instead. This is a much more natural choice for defining compact complex complexity classes (indeed, the sphere is not a constructible set over the complex numbers).

We now define the compact analogue of  $\mathbf{PH}_C$  that we will denote  $\mathbf{PH}_C^c$ . Unlike in the non-compact case, we will assume all variables vary over certain compact sets (namely complex projective spaces of varying dimensions).

We first need to be precise about what we mean by a complexity class of sequences of constructible subsets of complex projective spaces.

**Notation 1.8** (Affine cones). For any constructible subset  $S \subset \mathbb{P}_C^k$  we denote by  $C(S) \subset C^{k+1}$  the affine cone over  $S$ . More generally, if  $S \subset \mathbb{P}_C^{k_1} \times \dots \times \mathbb{P}_C^{k_\omega}$  is a constructible subset, then  $C(S) \subset C^{k_1+1} \times \dots \times C^{k_\omega+1}$  will denote the union of  $L^1 \times \dots \times L^\omega$  such that each  $L^i \subset C^{k_i+1}$  is a line through the origin, such that the point in  $\mathbb{P}_C^{k_1} \times \dots \times \mathbb{P}_C^{k_\omega}$  represented by  $L^1 \times \dots \times L^\omega$  is in  $S$ .

**Definition 1.9.** Let  $k(n)$  be a polynomial in  $n$ . We say that a sequence

$$\left( S_n \subset \mathbb{P}_C^{k(n)} \right)_{n>0}$$

of constructible subsets is in the complexity class  $\mathbf{P}_C$ , if the sequence of affine cones  $(C(S_n) \subset C^{k(n)+1})_{n>0} \in \mathbf{P}_C$ . More generally, suppose that  $m(n)$  is a polynomial in  $n$  and,  $(k_i(n))_{i>0}$  is a polynomially computable sequence of polynomials in  $n$ . We say that a sequence

$$\left( S_n \subset \mathbb{P}_C^{k_1(n)} \times \dots \times \mathbb{P}_C^{k_{m(n)}(n)} \right)_{n>0}$$

of constructible subsets is in the complexity class  $\mathbf{P}_C$ , if the sequence of affine cones  $(C(S_n) \subset C^{k_1(n)+1} \times \dots \times C^{k_{m(n)}(n)+1})_{n>0} \in \mathbf{P}_C$ .

*Remark 1.10.* Note that in the above definition if  $m(n)$  is a constant, then we can use the classical Segre embedding (see [20, Chap. 1, Sec. 5]) and obtain a polynomial time equivalent sequence of constructible subsets of projective spaces, and there is no need for the more general definition involving products of projective spaces. However, if  $m(n)$  grows polynomially in  $n$ , then the Segre embedding

requires exponentially many variables. On the other hand it seems perfectly reasonable that, for example, the sequence  $(\underbrace{\mathbb{P}_{\mathbb{C}}^n \times \cdots \times \mathbb{P}_{\mathbb{C}}^n}_{n \text{ times}})_{n>0}$  should be in  $\mathbf{P}_{\mathbb{C}}$  (see also Example 1.12 below). This is indeed so by the definition given above.

**Definition 1.11** (Compact projective version of  $\Sigma_{\mathbb{C},\omega}$ ). Suppose that  $m(n)$  is a polynomial in  $n$  and,  $(k_i(n))_{i>0}$  a polynomially computable sequence of polynomials in  $n$ . We say that a sequence

$$\left( S_n \subset \mathbb{P}_{\mathbb{C}}^{k_1(n)} \times \cdots \times \mathbb{P}_{\mathbb{C}}^{k_{m(n)}(n)} \right)_{n>0}$$

of constructible subsets is in the complexity class  $\Sigma_{\mathbb{C},\omega}^c$ , if there exists polynomials

$$\ell_1(n), \dots, \ell_{\omega}(n),$$

such that for each  $n > 0$ ,  $S_n$  is described by a first order formula

$$(\mathbf{Q}_1 \mathbf{Y}^1 \in \mathbb{P}_{\mathbb{C}}^{\ell_1(n)}) \cdots (\mathbf{Q}_{\omega} \mathbf{Y}^{\omega} \in \mathbb{P}_{\mathbb{C}}^{\ell_{\omega}(n)}) \phi_n(\mathbf{X}^1; \dots; \mathbf{X}^{m(n)}; \mathbf{Y}^1; \dots; \mathbf{Y}^{\omega}),$$

with  $\phi_n$  a quantifier-free first order multi-homogeneous formula defining a *closed* (in the Zariski topology) subset of  $\mathbb{P}_{\mathbb{C}}^{k_1(n)} \times \cdots \times \mathbb{P}_{\mathbb{C}}^{k_{m(n)}(n)} \times \mathbb{P}_{\mathbb{C}}^{\ell_1(n)} \times \cdots \times \mathbb{P}_{\mathbb{C}}^{\ell_{\omega}(n)}$ , and for each  $i, 1 \leq i \leq \omega$ ,  $\mathbf{X}^i = (X_0^i, \dots, X_{k_i(n)}^i)$  is a block of  $k_i(n) + 1$  variables,  $\mathbf{Y}^i = (Y_0^i, \dots, Y_{\ell_i(n)}^i)$  is a block of  $\ell_i(n) + 1$  variables,  $\mathbf{Q}_i \in \{\exists, \forall\}$ ,  $\mathbf{Q}_1 = \exists$ , and the sequence of constructible sets  $(T_n)_{n>0}$  defined by the formulas  $(\phi_n)_{n>0}$  belongs to the class  $\mathbf{P}_{\mathbb{C}}$ .

**Example 1.12.** We give a very natural example of a language in  $\Sigma_{\mathbb{C},1}^c$  (i.e. the compact version of  $\mathbf{NP}_{\mathbb{C}}$ ). Let  $k(n,d) = \binom{n+d}{d}$  and identify

$$\underbrace{\mathbb{P}_{\mathbb{C}}^{k(n,d)-1} \times \cdots \times \mathbb{P}_{\mathbb{C}}^{k(n,d)-1}}_{n+1 \text{ times}}$$

with systems of  $n + 1$  homogeneous polynomials in  $n$  variables of degree  $d$ . Let

$$S_{n,d} \subset \underbrace{\mathbb{P}_{\mathbb{C}}^{k(n,d)-1} \times \cdots \times \mathbb{P}_{\mathbb{C}}^{k(n,d)-1}}_{n+1 \text{ times}}$$

be defined by

$$S_{n,d} = \{(P_1; \cdots; P_{n+1}) \in P_i \in \mathbb{P}_{\mathbb{C}}^{k(n,d)-1} \mid \exists \mathbf{x} = (x_0 : \cdots : x_n) \in \mathbb{P}_{\mathbb{C}}^n \text{ with } P_1(\mathbf{x}) = \cdots = P_{n+1}(\mathbf{x}) = 0\}.$$

In other words,  $S_{n,d}$  is the set of systems of  $(n + 1)$  homogeneous polynomial equations of degree  $d$ , which have a zero in  $\mathbb{P}_{\mathbb{C}}^n$ . Then it is clear from the definition of the class  $\Sigma_{\mathbb{C},1}^c$  that for any fixed  $d > 0$ ,

$$\left( S_{n,d} \subset \underbrace{\mathbb{P}_{\mathbb{C}}^{k(n,d)-1} \times \cdots \times \mathbb{P}_{\mathbb{C}}^{k(n,d)-1}}_{n+1 \text{ times}} \right)_{n>0} \in \Sigma_{\mathbb{C},1}^c.$$

Note that it is *not known* if for any fixed  $d$

$$\left( S_{n,d} \subset \underbrace{\mathbb{P}_{\mathbb{C}}^{k(n,d)-1} \times \cdots \times \mathbb{P}_{\mathbb{C}}^{k(n,d)-1}}_{n+1 \text{ times}} \right)_{n>0}$$

is  $\mathbf{NP}_{\mathbb{C}}$ -complete, while the non-compact version of this language i.e. the language consisting of systems of polynomials having a zero in  $C^n$  (instead of  $\mathbb{P}_{\mathbb{C}}^n$ ), has been shown to be  $\mathbf{NP}_{\mathbb{C}}$ -complete for  $d \geq 2$  [3].

We define analogously the class  $\mathbf{\Pi}_{\mathbb{C},\omega}^c$ , and finally define:

**Definition 1.13.** The *compact projective polynomial hierarchy* over  $\mathbb{C}$  is defined to be the union

$$\mathbf{PH}_{\mathbb{C}}^c \stackrel{\text{def}}{=} \bigcup_{\omega \geq 0} (\Sigma_{\mathbb{C},\omega}^c \cup \mathbf{\Pi}_{\mathbb{C},\omega}^c) = \bigcup_{\omega \geq 0} \Sigma_{\mathbb{C},\omega}^c = \bigcup_{\omega \geq 0} \mathbf{\Pi}_{\mathbb{C},\omega}^c.$$

Notice that the constructible subsets belonging to any language in  $\mathbf{PH}_{\mathbb{C}}^c$  are all compact (in fact Zariski closed subsets of complex projective spaces).

*Remark 1.14.* The compact classes introduced above might be of interest in their own right. As remarked earlier it is not known whether the compact language

$$\left( S_{n,d} \subset \underbrace{\mathbb{P}_{\mathbb{C}}^{k(n,d)-1} \times \cdots \times \mathbb{P}_{\mathbb{C}}^{k(n,d)-1}}_{n+1 \text{ times}} \right)_{n>0}$$

in Example 1.12 is  $\mathbf{NP}_{\mathbb{C}}$ -complete. It is important to resolve this question in order to understand whether the hardness of solving polynomial systems over  $\mathbb{C}$  is due to the non-compactness of the (affine) solution space, or due to some intrinsic algebraic reasons.

1.4.1. *Complex projective analogue of  $\#\mathbf{P}$ .* We now define the complex analogue of  $\#\mathbf{P}$  (cf. the class  $\#\mathbf{P}_{\mathbb{R}}^{\dagger}$  defined in [2] in the real case).

We first need a notation.

**Notation 1.15** (Poincaré polynomial). In case  $\mathbb{C} = \mathbb{C}$ , for any constructible subset  $S \subset \mathbb{P}_{\mathbb{C}}^k$  we denote by  $b_i(S)$  the  $i$ -th Betti number (that is the rank of the singular homology group  $H_i(S) = H_i(S, \mathbb{Z})$ ) of  $S$ .

We also let  $P_S \in \mathbb{Z}[T]$  denote the *Poincaré polynomial* of  $S$ , namely

$$(1.2) \quad P_S(T) \stackrel{\text{def}}{=} \sum_{i \geq 0} b_i(S) T^i.$$

*Remark 1.16.* Over an arbitrary algebraically closed field  $\mathbb{C}$  of characteristic 0, ordinary singular homology is not well defined. We use a modified homology theory (which agrees with singular homology in case  $\mathbb{C} = \mathbb{C}$  and which is homotopy invariant) as done in [1] in case of semi-algebraic sets over arbitrary real closed fields (see [1], page 279). (Note that by taking real and imaginary parts, every constructible set is a semi-algebraic set over an appropriate real closed field.)

For the rest of the paper we will assume  $\mathbb{C} = \mathbb{C}$ , noting that all the results generalize to arbitrary algebraically closed fields of characteristic 0 using the transfer principle.

**Definition 1.17** (The class  $\#\mathbf{P}_C^\dagger$ ). We say a sequence of constructible functions

$$(f_n : \mathbb{P}_C^n \rightarrow \mathbb{Z}[T])_{n>0}$$

is in the class  $\#\mathbf{P}_C^\dagger$ , if there exists a polynomial  $m(n)$ , and a polynomially computable sequence of polynomials  $(k_i(n))_{i>0}$ , and a language

$$\left( S_n \subset \mathbb{P}_C^n \times \mathbb{P}_C^{k_1(n)} \times \cdots \times \mathbb{P}_C^{k_m(n)(n)} \right)_{n>0} \in \mathbf{P}_C,$$

such that

$$f_n(\mathbf{x}) = P_{S_n, \mathbf{x}}$$

for each  $\mathbf{x} \in \mathbb{P}_C^n$ , where  $S_{n, \mathbf{x}} = S_n \cap \pi^{-1}(\mathbf{x})$  and  $\pi : \mathbb{P}_C^n \times \mathbb{P}_C^{k_1(n)} \times \cdots \times \mathbb{P}_C^{k_m(n)(n)} \rightarrow \mathbb{P}_C^n$  is the projection along the last co-ordinates.

*Remark 1.18.* We make a few remarks about the class  $\#\mathbf{P}_C^\dagger$  defined above. First of all notice that the class  $\#\mathbf{P}_C^\dagger$  is quite robust. For instance, given two sequences  $(f_n)_{n>0}, (g_n)_{n>0} \in \#\mathbf{P}_C^\dagger$  it follows (by taking disjoint union of the corresponding constructible sets) that  $(f_n + g_n)_{n>0} \in \#\mathbf{P}_C^\dagger$ , and also  $(f_n g_n)_{n>0} \in \#\mathbf{P}_C^\dagger$  (by taking Cartesian product of the corresponding constructible sets and using the multiplicative property of the Poincaré polynomials, which itself is a consequence of the Kunneth formula in homology theory.)

*Remark 1.19.* The connection between counting points of varieties and their Betti numbers is more direct over fields of positive characteristic via the zeta function. The zeta function of a variety defined over  $\mathbb{F}_p$  is the exponential generating function of the sequence whose  $n$ -th term is the number of points in the variety over  $\mathbb{F}_{p^n}$ . The zeta function of such a variety turns out to be a rational function in one variable (a deep theorem of algebraic geometry first conjectured by Andre Weil [26] and proved by Dwork [12] and Deligne [10, 11]), and its numerator and denominator are products of polynomials whose degrees are the Betti numbers of the variety with respect to a certain ( $\ell$ -adic) co-homology theory. The point of this remark is that the problems of “counting” varieties and computing their Betti numbers, are connected at a deeper level, and thus our choice of definition for a complex analogue of  $\#\mathbf{P}$  is not altogether ad hoc.

*Remark 1.20.* A different definition of the class  $\#\mathbf{P}_C^\dagger$  (more in line with previous work of Burgisser et al. [8]) would be obtained by replacing in Definition 1.17 the Poincaré polynomial,  $P_S(T)$ , by the Euler-Poincaré characteristic i.e. the value of  $P_S$  at  $T = -1$ . The Euler-Poincaré characteristic is additive (at least in the category of compact varieties), and thus has some attributes of being a discrete analogue of volume. But at the same time it should be noted that the Euler-Poincaré characteristic is a rather weak invariant – for instance, it does not determine the number of connected components of a given variety. Also notice that in the case of finite fields referred to in Remark 1.19, all the Betti numbers, not just their alternating sum, enter (as degrees of factors) in the rational expression for the zeta function of a variety. While it would certainly be a much stronger reduction result if one could obtain a Toda-type theorem using only the Euler-Poincaré characteristic instead of the whole Poincaré polynomial, it is at present unclear if such a theorem can be proven.

## 2. STATEMENTS OF THE MAIN THEOREMS

We can now state the main result of this paper.

**Theorem 2.1** (Complex analogue of Toda's theorem).

$$\mathbf{PH}_C^c \subset \mathbf{P}_C^{\#\mathbf{P}_C^\dagger}.$$

*Remark 2.2.* We leave it as an open problem to prove Theorem 2.1 with  $\mathbf{PH}_C$  instead of  $\mathbf{PH}_C^c$  on the left hand side. However, we also note that many theorems of complex algebraic geometry take their most satisfactory form in the case of complete varieties, which is the setting considered in this paper.

As a consequence of our method, we obtain a reduction (Theorem 2.5) that might be of independent interest. We first define the following two problems:

**Definition 2.3** (Compact general decision problem with at most  $\omega$  quantifier alternations ( $\mathbf{GDP}_{C,\omega}^c$ )). The input and output for this problem are as follows.

- **Input.** A sentence  $\Phi$

$$(\mathbf{Q}_1 \mathbf{X}^1 \in \mathbb{P}^{k_1}) \cdots (\mathbf{Q}_\omega \mathbf{X}^\omega \in \mathbb{P}^{k_\omega}) \phi(\mathbf{X}^1; \dots; \mathbf{X}^\omega),$$

where for each  $i, 1 \leq i \leq \omega$ ,  $\mathbf{Q}_i \in \{\exists, \forall\}$ , with  $\mathbf{Q}_j \neq \mathbf{Q}_{j+1}, 1 \leq j < \omega$ , and  $\phi$  is a quantifier-free multi-homogeneous formula defining a *closed* subset  $S$  of  $\mathbb{P}^{k_1} \times \cdots \times \mathbb{P}^{k_\omega}$ .

- **Output.** True or False depending on whether  $\Phi$  is true or false.

**Definition 2.4** (Computing the Poincaré polynomial of constructible sets (*Poincaré*)). The input and output for this problem are as follows.

- **Input.** A quantifier-free homogeneous formula defining a constructible subset  $S \subset \mathbb{P}_C^k$ .
- **Output.** The Poincaré polynomial  $P_S(T)$ .

**Theorem 2.5.** *For every  $\omega > 0$ , there is a deterministic polynomial time reduction in the Blum-Shub-Smale model of  $\mathbf{GDP}_{C,\omega}^c$  to *Poincaré*.*

*Remark 2.6.* We remark that (in contrast to the real case) in the complex case, we are able to prove a slightly stronger result than stated above in Theorem 2.5. Our proof of Theorem 2.5 gives a polynomial time reduction of  $\mathbf{GDP}_{C,\omega}^c$  to the problem of computing the *pseudo-Poincaré* polynomial (defined below, see Eqn. 3.2) of constructible sets. The pseudo-Poincaré polynomial is easily computable from the Poincaré polynomial.

**2.1. Outline of the main ideas and contributions.** The basic idea behind the proof of a real analogue of Toda's theorem in [2] is a topological construction, which given a semi-algebraic set  $S \subset \mathbb{R}^{m+n}$ ,  $p \geq 0$ , and  $\pi : \mathbb{R}^{m+n} \subset \mathbb{R}^n$  the projection along (say) the first  $m$  co-ordinates, constructs *efficiently* a semi-algebraic set,  $D^p(S)$ , such that

$$(2.1) \quad b_i(\pi(S)) = b_i(D^p(S)), \quad 0 \leq i < p.$$

Moreover, membership in  $D^p(S)$  can be tested efficiently if the same is true for  $S$ . Note that this last property will not hold in general for the set  $\pi(S)$  itself (unless of course  $\mathbf{P}_R = \mathbf{NP}_R$ ).

The topological construction used in the definition of  $D^p(X)$  in [2] is the iterated fibered join,  $J_\pi^p(X)$ , of a semi-algebraic set  $X$  with itself over a projection map  $\pi$ .

The fibers of the induced map  $J_\pi(X) : J_\pi^p(X) \rightarrow \pi(X)$ , over a point  $y \in \pi(X)$ , are then ordinary  $(p+1)$ -fold joins of the fiber  $\pi^{-1}(y)$ , and by connectivity properties of the join are  $p$ -connected. It is now possible using a version of the Vietoris-Beagle theorem that the map  $J_\pi(X)$  is a  $p$ -equivalence (see [2] for the precise definition of  $p$ -equivalence). The main construction in [2] was to realize efficiently the fibered join  $J_\pi^p(X)$  up to homotopy by a semi-algebraic set. This construction however is semi-algebraic in nature – i.e. it uses real inequalities in an essential way and thus does not generalize in a straightforward way to the complex case. Thus, a different construction is needed in the complex case.

In the complex case, the role of the fibered join is played by the *complex join fibered over a map* defined below (see Definition 3.19). The fibers of the  $(p+1)$ -fold complex join fibered over a projection  $\pi$ ,  $J_{\mathbb{C},\pi}^p(X)$ , of a compact constructible set  $X$  are not quite  $p$ -connected as in the real case, but are reasonably nice – namely they are homologically equivalent to a projective space of dimension  $p$  (see Proposition 3.15). This allows us to relate the Poincaré polynomial of  $X$  with that of its image  $\pi(X)$ , even though the relation is not as straightforward as in the real case (see Theorem 3.21 below).

We remark that Theorem 3.21 can be used to express directly the Betti numbers of the image under projection of a projective variety in terms of those another projective variety obtained directly without having to perform effective quantifier elimination (which has exponential complexity). The description of this second variety is *much simpler and algebraic* in nature compared to the one used in [2] in the real semi-algebraic case, and thus might be of independent interest. Theorem 3.21 can also be viewed as an improvement over the descent spectral sequence argument used in [13] to bound the Betti numbers of projections (of semi-algebraic sets) in the complex projective case. A similar construction using the projective join is also available in the real case (using  $\mathbb{Z}/2\mathbb{Z}$  coefficients) but we omit its description in the current paper.

Finally, we believe that the compact projective versions of the complex complexity classes introduced in this paper deserve further investigations on their own (see Remark 1.14 below), since many known efficient algorithms in complex algebraic geometry assume some form of compactness (see, for instance, [5]).

The rest of the paper is organized as follows. In Section 3 we state and prove the necessary ingredients from algebraic topology needed to prove the main theorems. In Section 4 we prove the main results of the paper.

### 3. TOPOLOGICAL INGREDIENTS

In this section we state and prove the main topological ingredients necessary for the proof of the main theorems.

**3.1. Alexander-Lefschetz duality.** We will need the classical Alexander-Lefschetz duality theorem in order to relate the Betti numbers of a constructible subset  $S \subset \mathbb{P}_{\mathbb{C}}^{k_1} \times \cdots \times \mathbb{P}_{\mathbb{C}}^{k_\ell}$  with those of its complement  $\mathbb{P}_{\mathbb{C}}^{k_1} \times \cdots \times \mathbb{P}_{\mathbb{C}}^{k_\ell} \setminus S$ .

**Theorem 3.1** (Alexander-Lefschetz duality). *Let  $S \subset \mathbb{P}_{\mathbb{C}}^{k_1} \times \cdots \times \mathbb{P}_{\mathbb{C}}^{k_\ell}$  be a constructible subset. Then for each odd  $i$ ,  $1 \leq i \leq 2k+1$  with  $k = k_1 + \cdots + k_\ell$ , we have that*

$$(3.1) \quad b_{i-1}(S) - b_{i-2}(S) = b_{2k-i}(\mathbb{P}_{\mathbb{C}}^{k_1} \times \cdots \times \mathbb{P}_{\mathbb{C}}^{k_\ell} - S) - b_{2k-i+1}(\mathbb{P}_{\mathbb{C}}^{k_1} \times \cdots \times \mathbb{P}_{\mathbb{C}}^{k_\ell} - S) + b_i(\mathbb{P}_{\mathbb{C}}^{k_1} \times \cdots \times \mathbb{P}_{\mathbb{C}}^{k_\ell}).$$

*Proof.* Lefschetz duality theorem [22] gives for each  $i, 0 \leq i \leq 2k$ ,

$$b_i(\mathbb{P}_{\mathbb{C}}^{k_1} \times \cdots \times \mathbb{P}_{\mathbb{C}}^{k_\ell} - A) = b_{2k-i}(\mathbb{P}_{\mathbb{C}}^{k_1} \times \cdots \times \mathbb{P}_{\mathbb{C}}^{k_\ell}, S).$$

The theorem now follows from the long exact sequence of homology,

$$\cdots \rightarrow H_i(S) \rightarrow H_i(\mathbb{P}_{\mathbb{C}}^{k_1} \times \cdots \times \mathbb{P}_{\mathbb{C}}^{k_\ell}) \rightarrow H_i(\mathbb{P}_{\mathbb{C}}^{k_1} \times \cdots \times \mathbb{P}_{\mathbb{C}}^{k_\ell}, S) \rightarrow H_{i-1}(S) \rightarrow \cdots$$

after noting that  $H_i(\mathbb{P}_{\mathbb{C}}^{k_1} \times \cdots \times \mathbb{P}_{\mathbb{C}}^{k_\ell}) = 0$ , for all  $i \neq 0, 2, 4, \dots, 2k$ .  $\square$

For technical reasons (see Corollary 3.4 below) we need to consider the even and odd parts of the Poincaré polynomial of constructible sets.

Given  $P = \sum_{i \geq 0} a_i T^i \in \mathbb{Z}[T]$ , we write

$$P \stackrel{\text{def}}{=} P^{\text{even}}(T^2) + TP^{\text{odd}}(T^2),$$

where

$$P^{\text{even}}(T) = \sum_{i \geq 0} a_{2i} T^i,$$

and

$$P^{\text{odd}}(T) = \sum_{i \geq 0} a_{2i+1} T^i.$$

We introduce for any  $S \subset \mathbb{P}_{\mathbb{C}}^n$ , a related polynomial,  $Q_S(T)$ , which we call the *pseudo-Poincaré polynomial* of  $S$  defined as follows.

$$(3.2) \quad Q_S(T) \stackrel{\text{def}}{=} \sum_{j \geq 0} (b_{2j}(S) - b_{2j-1}(S)) T^j.$$

In other words,

$$(3.3) \quad Q_S = P_S^{\text{even}} - TP_S^{\text{odd}}.$$

We introduce below notation for several operators on polynomials that we will use later.

**Notation 3.2** (Operators on polynomials). For any polynomial  $Q = \sum_{i \geq 0} a_i T^i \in \mathbb{Z}[T]$  with  $\deg(Q) \leq n$ , we will denote by:

- (A)  $\text{Rec}_n(Q)$  the polynomial  $T^n Q(\frac{1}{T})$ ;
- (B)  $\text{Trunc}_m(Q)$  the polynomial  $\sum_{0 \leq i \leq m} a_i T^i \in \mathbb{Z}[T]$ ; and,
- (C)  $M_P(Q)$  the polynomial  $PQ$ , for any polynomial  $P \in \mathbb{Z}[T]$ .

*Remark 3.3.* Notice that all the operators introduced above are computable in polynomial time.

Using the notation introduced above we have the following easy corollary of Theorem 3.1.

**Corollary 3.4.** *Let  $S \subset \mathbb{P}_{\mathbb{C}}^{k_1} \times \cdots \times \mathbb{P}_{\mathbb{C}}^{k_\ell}$  be any constructible subset, and  $k = k_1 + \cdots + k_\ell$ . Then,*

$$Q_S(T) = -\text{Rec}_k(Q_{\mathbb{P}_{\mathbb{C}}^{k_1} \times \cdots \times \mathbb{P}_{\mathbb{C}}^{k_\ell} - S}) + P_{\mathbb{P}_{\mathbb{C}}^{k_1} \times \cdots \times \mathbb{P}_{\mathbb{C}}^{k_\ell}}^{\text{even}}.$$

**3.2. The complex join of constructible sets.** Let  $X \subset \mathbb{P}_{\mathbb{C}}^k$  and  $Y \subset \mathbb{P}_{\mathbb{C}}^{\ell}$  be two constructible sets defined by homogeneous formulas  $\phi(X_0, \dots, X_k)$  and  $\psi(Y_0, \dots, Y_{\ell})$  respectively, where  $(X_0 : \dots : X_k)$  (respectively  $(Y_0 : \dots : Y_{\ell})$ ) are homogeneous co-ordinates in  $\mathbb{P}_{\mathbb{C}}^k$  (respectively  $\mathbb{P}_{\mathbb{C}}^{\ell}$ ).

**Definition 3.5** (Complex join). The *complex join*,  $J_{\mathbb{C}}(X, Y)$ , of  $X$  and  $Y$  is defined to be the constructible subset of  $\mathbb{P}_{\mathbb{C}}^{k+\ell+1}$  defined by the formula

$$\phi(Z_0, \dots, Z_k) \wedge \psi(Z_{k+1}, \dots, Z_{k+\ell+1}),$$

where  $(Z_0 : \dots : Z_{k+\ell+1})$  are homogeneous coordinates in  $\mathbb{P}_{\mathbb{C}}^{k+\ell+1}$ .

*Remark 3.6.* Firstly, notice that  $J_{\mathbb{C}}(X, Y)$  does not depend on the formulas  $\phi$  and  $\psi$  used to define  $X$  and  $Y$  respectively. Also, notice that if  $X$  and  $Y$  are both empty then so is  $J_{\mathbb{C}}(X, Y)$ . Indeed, if  $X = \emptyset$  (respectively,  $Y = \emptyset$ ) then  $J_{\mathbb{C}}(X, Y)$  is isomorphic to  $Y$  (respectively,  $X$ ). If  $X$  and  $Y$  are both non-empty then  $J_{\mathbb{C}}(X, Y)$  is obtained topologically by joining each point of  $X$  with each point of  $Y$  by a complex projective line,  $\mathbb{P}_{\mathbb{C}}^1$ .

**Example 3.7.** It is easy to check from the above definition that the join,  $J_{\mathbb{C}}(\mathbb{P}_{\mathbb{C}}^k, \mathbb{P}_{\mathbb{C}}^{\ell})$ , of two projective spaces is again a projective space, namely  $\mathbb{P}_{\mathbb{C}}^{k+\ell+1}$ .

*Remark 3.8.* The projective join as defined above is a classical object in algebraic geometry. Amongst many other applications, the complex suspension of a projective variety  $X$  (i.e. the complex join  $J_{\mathbb{C}}(X, \mathbb{P}_{\mathbb{C}}^1)$ ) plays an important role in defining Lawson homology of projective varieties [14]. Within the area of computational complexity theory, the projective join of a variety with a point was used in [18] for proving hardness of the problem of computing Betti numbers of complex varieties.

**Definition 3.9.** For  $p > 0$ , we denote by  $J_{\mathbb{C}}^p(X)$  the  $(p+1)$ -fold iterated complex join of  $X$  with itself.

In other words,

$$J_{\mathbb{C}}^p X = \underbrace{J_{\mathbb{C}}(J_{\mathbb{C}}(\dots(J_{\mathbb{C}}(X, X))\dots))}_{(p+1) \text{ times}}.$$

If  $X \subset \mathbb{P}_{\mathbb{C}}^k$  is defined by a first-order homogeneous formula  $\phi(X_0, \dots, X_k)$ , then  $J_{\mathbb{C}}^p(X) \subset \mathbb{P}_{\mathbb{C}}^{(p+1)(k+1)-1}$  is defined by the homogeneous formula

$$J_{\mathbb{C}}^p(\phi)(X_0^0, \dots, X_k^0, \dots, X_0^p, \dots, X_k^p) \stackrel{\text{def}}{=} \bigwedge_{i=0}^p \phi(X_0^i, \dots, X_k^i).$$

where  $(X_0^0 : \dots : X_k^p)$  are homogeneous co-ordinates in  $\mathbb{P}_{\mathbb{C}}^{(p+1)(k+1)-1}$ .

Note that by Remark 3.6, if  $X$  is empty then  $J_{\mathbb{C}}^p(X)$  is empty for every  $p > 0$ .

**3.3. Properties of the topological join.** We also need to introduce the *topological join* of two spaces. The following is mostly taken from [2].

**Definition 3.10.** The *join*  $X * Y$  of two topological spaces  $X$  and  $Y$  is defined by

$$(3.4) \quad X * Y \stackrel{\text{def}}{=} X \times Y \times \Delta^1 / \sim,$$

where

$$(x, y, t_0, t_1) \sim (x', y', t_0, t_1)$$

if  $t_0 = 1, x = x'$  or  $t_1 = 1, y = y'$ .

Intuitively,  $X * Y$  is obtained by joining each point of  $X$  with each point of  $Y$  by an interval.

We will need the well-known fact that the iterated join of a topological space is highly connected. In order to make this statement precise we first define

**Definition 3.11** ( $p$ -equivalence). A map  $f : A \rightarrow B$  between two topological spaces is called a  **$p$ -equivalence** if the induced homomorphism

$$f_* : H_i(A) \rightarrow H_i(B)$$

is an isomorphism for all  $0 \leq i < p$ , and an epimorphism for  $i = p$ , and we say that  $A$  is  **$p$ -equivalent** to  $B$ .

The following is well known (see, for instance, [15, Proposition 4.4.3]).

**Theorem 3.12.** *Let  $X$  be a compact semi-algebraic set. Then, the  $(p+1)$ -fold join  $\underbrace{X * \dots * X}_{(p+1) \text{ times}}$  is  $p$ -equivalent to a point.*

We will need a particular property of projection maps that we are going to consider later in the paper.

**Notation 3.13.** For any constructible set  $A$ , we denote by  $K(A)$  the collection of all compact (in the Euclidean topology) subsets of  $A$ .

**Definition 3.14.** Let  $f : A \rightarrow B$  be a map between two constructible sets  $A$  and  $B$ . We say that  $f$  **compact covering** if for any  $L \in K(f(A))$ , there exists  $K \in K(A)$  such that  $f(K) = L$ .

### 3.4. Topological properties of the complex join.

**Proposition 3.15.** *Let  $X \subset \mathbb{P}_{\mathbb{C}}^k$  be a non-empty constructible subset and  $p > 0$ . Let*

$$i : J_{\mathbb{C}}^p(X) \hookrightarrow \mathbb{P}_{\mathbb{C}}^{(p+1)(k+1)-1}$$

denote the inclusion map. Then the induced homomorphism

$$i_* : H_j(J_{\mathbb{C}}^p X) \rightarrow H_j(\mathbb{P}_{\mathbb{C}}^{(p+1)(k+1)-1})$$

is an isomorphism for  $0 \leq j < p$ .

Before proving Proposition 3.15 we first fix some notation.

**Notation 3.16** (Hopf map). For any  $k \geq 0$ , we will denote by  $\pi : \mathbb{C}^{k+1} \setminus \{0\} \rightarrow \mathbb{P}_{\mathbb{C}}^k$  the tautological line bundle over  $\mathbb{P}_{\mathbb{C}}^k$ , and by

$$\tilde{\pi} : \mathbf{S}^{2k+1} \rightarrow \mathbb{P}_{\mathbb{C}}^k,$$

the **Hopf fibration**, namely the restriction of  $\pi$  to the unit sphere in  $\mathbb{C}^{k+1}$  defined by the equation  $|z_1|^2 + \dots + |z^{k+1}|^2 = 1$ . Finally for any subset  $S \subset \mathbb{P}_{\mathbb{C}}^k$ , we will denote by  $\tilde{S}$  the subset  $\tilde{\pi}^{-1}(S) \subset \mathbf{S}^{2k+1}$ . Restricting the map  $\tilde{\pi}$  to  $\tilde{S}$  we obtain the restriction of the Hopf fibration to the base  $S$  i.e. we have the following commutative diagram.

$$\begin{array}{ccc} \tilde{S} & \xrightarrow{i} & \mathbf{S}^{2k+1} \\ \downarrow \tilde{\pi} & & \downarrow \tilde{\pi} \\ S & \xrightarrow{i} & \mathbb{P}_{\mathbb{C}}^k \end{array}$$

We need the following lemma.

**Lemma 3.17.** *Let  $X \subset \mathbb{P}_{\mathbb{C}}^k, Y \subset \mathbb{P}_{\mathbb{C}}^{\ell}$  be constructible subsets. Then  $J_{\mathbb{C}}(\widetilde{X}, Y) \subset \mathbf{S}^{2(k+\ell)+3}$  is homeomorphic to the (topological) join  $\widetilde{X} * \widetilde{Y}$ .*

*Proof.* Consider  $\mathbf{x} \in X$  and  $\mathbf{y} \in Y$  and the projective line  $L \subset J_{\mathbb{C}}(X, Y)$  joining  $\mathbf{x}$  and  $\mathbf{y}$ . It is easy to see that the preimage  $\tilde{L} = \tilde{\pi}^{-1}(L) \cong \mathbf{S}^1$  is a topological join of  $\tilde{\pi}^{-1}(\mathbf{x})$  and  $\tilde{\pi}^{-1}(\mathbf{y})$  (each homeomorphic to  $\mathbf{S}^1$ ). Now since  $\widetilde{X}$  (resp.  $\widetilde{Y}$ ) is fibered by the various  $\tilde{\pi}^{-1}(\mathbf{x})$  (resp.  $\tilde{\pi}^{-1}(\mathbf{y})$ ), it follows that  $J_{\mathbb{C}}(\widetilde{X}, Y)$  is homeomorphic to  $\widetilde{X} * \widetilde{Y}$ .  $\square$

*Proof of Proposition 3.15.* It follows from repeated applications of Lemma 3.17 that  $\widetilde{J_{\mathbb{C}}^p(X)}$  is homeomorphic to

$$\underbrace{\widetilde{X} * \cdots * \widetilde{X}}_{(p+1) \text{ times}}.$$

We also have the commutative square

$$\begin{array}{ccc} \widetilde{J_{\mathbb{C}}^p X} & \xrightarrow{i} & \mathbf{S}^{2(p+1)(k+1)-1} \\ \downarrow \tilde{\pi} & & \downarrow \tilde{\pi} \\ J_{\mathbb{C}}^p X & \xrightarrow{i} & \mathbb{P}_{\mathbb{C}}^{(p+1)(k+1)-1} \end{array}$$

and a corresponding square

$$\begin{array}{ccc} H_*(\widetilde{J_{\mathbb{C}}^p X}) & \xrightarrow{i_*} & H_*(\mathbf{S}^{2(p+1)(k+1)-1}) \\ \downarrow \tilde{\pi}_* & & \downarrow \tilde{\pi}_* \\ H_*(J_{\mathbb{C}}^p X) & \xrightarrow{i_*} & H_*(\mathbb{P}_{\mathbb{C}}^{(p+1)(k+1)-1}) \end{array}$$

of induced homomorphisms in the homology groups.

It follows from Theorem 3.12 that if  $X \neq \emptyset$ , then

$$\begin{aligned} H_0(\widetilde{J_{\mathbb{C}}^p(X)}) &\cong \mathbb{Z}, \\ H_i(\widetilde{J_{\mathbb{C}}^p X}) &\cong 0, \quad 0 < i < p. \end{aligned}$$

It is easy to see that for  $p > 0$ ,  $J_{\mathbb{C}}^p(X)$  is simply connected and hence  $\widetilde{J_{\mathbb{C}}^p(X)}$  is a simple  $\mathbf{S}^1$ -bundle (i.e. a  $\mathbf{S}^1$ -bundle with a simply connected base) over  $J_{\mathbb{C}}^p(X)$ .

It now follows by a standard argument (which we expand below) involving the spectral sequence of the bundle  $\tilde{\pi} : \widetilde{J_{\mathbb{C}}^p(X)} \rightarrow J_{\mathbb{C}}^p(X)$ , that for  $0 \leq i < p$ ,

$$(3.5) \quad \begin{aligned} H_i(J_{\mathbb{C}}^p(X)) &\cong \mathbb{Z}, \text{ for } i \text{ even,} \\ H_i(J_{\mathbb{C}}^p(X)) &\cong 0 \text{ for } i \text{ odd.} \end{aligned}$$

(The above claim also follows from the Gysin sequence of the  $\mathbf{S}^1$ -bundle  $\tilde{\pi} : \widetilde{J_{\mathbb{C}}^p(X)} \rightarrow J_{\mathbb{C}}^p(X)$  but we give an independent proof below).

Consider the  $E_2$ -term of the (homological) spectral sequence of the bundle

$$\tilde{\pi} : \widetilde{J_{\mathbb{C}}^p(X)} \rightarrow J^p(X).$$

For  $i, j \geq 0$ , we have that

$$E_2^{i,j} = H_i(J_{\mathbb{C}}^p(X)) \otimes H_j(\mathbf{S}^1).$$

From this we deduce that

$$E_2^{i,0} = E_2^{i,1} = H_i(J_{\mathbb{C}}^p(X)).$$

Also, from the fact that

$$H_0(\widetilde{J_{\mathbb{C}}^p(X)}) = \mathbb{Z},$$

we get that

$$E_2^{0,0} = \mathbb{Z},$$

and hence,

$$E_2^{0,1} = \mathbb{Z}$$

as well. Moreover, we have that

$$E_3^{i,j} = E_4^{i,j} = \dots = E_{\infty}^{i,j}$$

for all  $i \geq 0$  and  $j = 0, 1$ . Now from the fact that the spectral sequence  $E_r$  converges to the homology of  $\widetilde{J_{\mathbb{C}}^p(X)}$  we deduce that

$$\begin{aligned} E_3^{i,j} &= 0 \text{ for } 0 \leq i \leq p-1 \text{ and all } j, \\ E_3^{0,0} &= \mathbb{Z}. \end{aligned}$$

This implies that the differential

$$d_2 : E_2^{i,0} \rightarrow E_2^{i-2,1}$$

is an isomorphism for  $1 \leq i \leq p-1$ . Together with the fact that

$$E_2^{i,0} = E_2^{i,1} = H_i(J_{\mathbb{C}}^p(X)),$$

this immediately implies (3.5). The proposition follows directly from this.  $\square$

In our application we will need the following (rather technical) generalization of Proposition 3.15.

**Proposition 3.18.** *Let  $p, \alpha_0, \dots, \alpha_{\omega} \geq 0$ ,  $N = \prod_{0 \leq j \leq \omega} (\alpha_j + 1)$ , and let  $\phi$  be a homogeneous formula defining a constructible subset of  $\mathbb{P}_{\mathbb{C}}^k$ . Also, let for each  $i, 0 \leq i \leq \omega$ ,  $\Lambda^i \in \{\wedge, \vee\}$ , and let  $\Phi$  denote the multi-homogeneous formula defined by*

$$\Phi \stackrel{\text{def}}{=} \Lambda_{0 \leq i_0 \leq \alpha_0}^0 \cdots \Lambda_{0 \leq i_{\omega} \leq \alpha_{\omega}}^{\omega} J_{\mathbb{C}}^p \phi(\mathbf{X}_{i_0, \dots, i_{\omega}}).$$

Let

$$S = \mathcal{R}(\Phi) \subset \underbrace{\mathbb{P}_{\mathbb{C}}^{(p+1)(k+1)-1} \times \dots \times \mathbb{P}_{\mathbb{C}}^{(p+1)(k+1)-1}}_{N \text{ times}},$$

and let

$$i : S \hookrightarrow \mathbb{P}_{\mathbb{C}}^{(p+1)(k+1)-1} \times \dots \times \mathbb{P}_{\mathbb{C}}^{(p+1)(k+1)-1}$$

denote the inclusion map. Then, the induced homomorphisms

$$i_* : H_j(S) \rightarrow H_j(\mathbb{P}_{\mathbb{C}}^{(p+1)(k+1)-1} \times \dots \times \mathbb{P}_{\mathbb{C}}^{(p+1)(k+1)-1})$$

are isomorphisms for  $0 \leq j < p$ .

*Proof.* Notice that, if  $\omega = 0$  and  $\Lambda^0 = \bigwedge$ , then

$$\bigwedge_{0 \leq i_0 \leq \alpha_0} J_{\mathbb{C}}^p \phi(\mathbf{X}_{i_0})$$

defines the product of the constructible sets  $S_{i_0} = \mathcal{R}(J_{\mathbb{C}}^p \phi(X_{i_0}))$ ,  $0 \leq i_0 \leq \alpha_0$ . The claim follows in this case from Proposition 3.15 and the Kunnetth formula for homology.

If  $\omega = 0$  and  $\Lambda^0 = \bigvee$ , the claim follows from the previous case and a standard argument using the Mayer-Vietoris double complex.

The general case is easily proved using induction on  $\omega$ .  $\square$

**3.5. Complex join fibered over a map and its properties.** In our application we need the complex join fibered over certain maps.

**Definition 3.19** (Complex join fibered over a map). Let  $A \subset \mathbb{P}_{\mathbb{C}}^k \times \mathbb{P}_{\mathbb{C}}^{\ell}$  be a constructible set defined by a first-order multi-homogeneous formula,

$$\phi(X_0, \dots, X_k; Y_0, \dots, Y_{\ell})$$

and let  $\pi_{\mathbf{Y}} : \mathbb{P}_{\mathbb{C}}^k \times \mathbb{P}_{\mathbb{C}}^{\ell} \rightarrow \mathbb{P}_{\mathbb{C}}^k$  be the projection along the  $\mathbf{Y}$ -co-ordinates.

For  $p > 0$ , the  *$p$ -fold complex join of  $A$  fibered over the map  $\pi_{\mathbf{Y}}$* ,  $J_{\mathbb{C}, \mathbf{Y}}^p(A) \subset \mathbb{P}_{\mathbb{C}}^k \times \mathbb{P}_{\mathbb{C}}^{(\ell+1)(p+1)-1}$ , is defined by the formula

$$(3.6) \quad J_{\mathbb{C}, \mathbf{Y}}^p(\phi)(X_0, \dots, X_k; Y_0^0, \dots, Y_{\ell}^0, \dots, Y_0^p, \dots, Y_{\ell}^p) \stackrel{\text{def}}{=} \bigwedge_{i=0}^p \phi(X_0, \dots, X_k; Y_0^i, \dots, Y_{\ell}^i).$$

*Remark 3.20.* There is a natural induced map

$$J_{\mathbb{C}, \mathbf{Y}}^p : J_{\mathbb{C}, \mathbf{Y}}^p(A) \rightarrow \pi_{\mathbf{Y}}(A)$$

sending  $(x_0 : \dots : x_k; y_0^0 : \dots : y_{\ell}^0) \in J_{\mathbb{C}, \mathbf{Y}}^p(A)$  to  $(x_0 : \dots : x_k) \in \pi_{\mathbf{Y}}(A)$ . It is easy to verify from Definition 3.19 that the map  $J_{\mathbb{C}, \mathbf{Y}}^p$  is well defined and is a surjection.

Now, let  $A \subset \mathbb{P}_{\mathbb{C}}^k \times \mathbb{P}_{\mathbb{C}}^{\ell}$  be a constructible subset  $\pi_{\mathbf{Y}} : \mathbb{P}_{\mathbb{C}}^k \times \mathbb{P}_{\mathbb{C}}^{\ell} \rightarrow \mathbb{P}_{\mathbb{C}}^k$  be the projection along the last co-ordinates. Suppose that  $\pi_{\mathbf{Y}}$  is a compact covering. The following theorem relates the Poincaré polynomial of  $J_{\mathbb{C}, \mathbf{Y}}^p(A)$  to that of the image  $\pi_{\mathbf{Y}}(A)$ .

**Theorem 3.21.** *For every  $p \geq 0$ , we have that*

$$(3.7) \quad P_{\pi_{\mathbf{Y}}(A)} = (1 - T^2)P_{J_{\mathbb{C}, \mathbf{Y}}^p(A)} \pmod{T^p}.$$

*Proof.* We first assume that  $A$  is compact. We have the following commutative square.

$$\begin{array}{ccc} J_{\mathbb{C}, \mathbf{Y}}^p(A) & \xrightarrow{i} & \pi_{\mathbf{Y}}(A) \times \mathbb{P}_{\mathbb{C}}^{(p+1)(\ell+1)-1} \\ \downarrow J_{\mathbb{C}, \mathbf{Y}}^p & & \downarrow \pi_{\mathbf{Y}} \\ \pi_{\mathbf{Y}}(A) & \xrightarrow{\text{Id}} & \pi_{\mathbf{Y}}(A) \end{array}$$

The diagram above induces a morphism,  $\phi_r^{i,j} : E_r^{i,j} \rightarrow 'E_r^{i,j}$  between the (homological) Leray spectral sequences of the two vertical maps in the above diagram. Here,  $E_r$  (resp.  $'E_r$ ) denotes the Leray spectral sequence of the map

$J_{\mathbb{C}, \mathbf{Y}}^p : J_{\mathbb{C}, \mathbf{Y}}^p(A) \rightarrow \pi_{\mathbf{Y}}(A)$  (resp.  $\pi_{\mathbf{Y}} : \pi_{\mathbf{Y}}(A) \times \mathbb{P}_{\mathbb{C}}^{(p+1)(\ell+1)-1} \rightarrow \pi_{\mathbf{Y}}(A)$ ). The spectral sequence,  $'E_r$ , of the map  $\pi_{\mathbf{Y}} : \pi_{\mathbf{Y}}(A) \times \mathbb{P}_{\mathbb{C}}^{(p+1)(\ell+1)-1} \rightarrow \pi_{\mathbf{Y}}(A)$  degenerates at the  $'E_2$ -term where

$$'E_2^{i,j} = H_i(\pi_{\mathbf{Y}}(A), \mathcal{H}_j(\mathbb{P}_{\mathbb{C}}^{(p+1)(\ell+1)-1})),$$

where  $H_i(\pi_{\mathbf{Y}}(A), \mathcal{H}_j(\mathbb{P}_{\mathbb{C}}^{(p+1)(\ell+1)-1}))$  denotes the  $i$ -th homology group of  $\pi_{\mathbf{Y}}(A)$  with local coefficients taking values in the fibers  $H_j(\pi_{\mathbf{Y}}^{-1}(\mathbf{x}))$ ,  $\mathbf{x} \in \pi_{\mathbf{Y}}(A)$ . Moreover, it follows from Proposition 3.15 that

$$\phi_2^{i,j} : E_2^{i,j} \rightarrow 'E_2^{i,j}$$

are isomorphisms for  $i + j < p$ . Thus,  $E_{\infty}^{i,j} = 'E_{\infty}^{i,j}$  for  $0 \leq i + j < p$ . This implies that  $H_q(J_{\mathbb{C}, \mathbf{Y}}^p(A)) \cong H_q(\pi_{\mathbf{Y}}(A) \times \mathbb{P}_{\mathbb{C}}^{(p+1)(\ell+1)-1})$  for  $0 \leq q < p$ , and thus

$$(3.8) \quad P_{J_{\mathbb{C}, \mathbf{Y}}^p(A)} = P_{\pi_{\mathbf{Y}}(A) \times \mathbb{P}_{\mathbb{C}}^{(p+1)(\ell+1)-1}} \pmod{T^p}.$$

We also have that

$$(3.9) \quad \begin{aligned} P_{\pi_{\mathbf{Y}}(A) \times \mathbb{P}_{\mathbb{C}}^{(p+1)(\ell+1)-1}} &= P_{\pi_{\mathbf{Y}}(A)} \times P_{\mathbb{P}_{\mathbb{C}}^{(p+1)(\ell+1)-1}} \\ &= P_{\pi_{\mathbf{Y}}(A)} \times (1 + T^2 + \dots + T^{2((p+1)(\ell+1)-1)}) \\ &= P_{\pi_{\mathbf{Y}}(A)} \times (1 - T^2)^{-1} \pmod{T^p}. \end{aligned}$$

The theorem now follows from Eqns. (3.8) and (3.9). The general case follows by taking direct limit over all compact subsets of  $A$ . More precisely, for  $K_1 \subset K_2$  compact subsets of  $A$ , we have for  $0 \leq q < p$  the following commutative square.

$$\begin{array}{ccc} H_q(J_{\mathbb{C}, \mathbf{Y}}^p(K_1)) & \xrightarrow{i_*} & H_q(J_{\mathbb{C}, \mathbf{Y}}^p(K_2)) \\ \downarrow \cong & & \downarrow \cong \\ H_q(\pi_{\mathbf{Y}}(K_1) \times \mathbb{P}_{\mathbb{C}}^{(p+1)(\ell+1)-1}) & \xrightarrow{i_*} & H_q(\pi_{\mathbf{Y}}(K_2) \times \mathbb{P}_{\mathbb{C}}^{(p+1)(\ell+1)-1}) \end{array}$$

where the vertical maps are isomorphisms by the previous case. If we take the direct limit as  $K$  ranges in  $\mathbf{K}(A)$ , we obtain the following:

$$\begin{array}{ccc} \varinjlim H_q(J_{\mathbb{C}, \mathbf{Y}}^p(K)) & \xrightarrow{\cong} & H_q(J_f^p(A)) \\ \downarrow \cong & & \downarrow \\ \varinjlim H_q(\pi_{\mathbf{Y}}(K) \times \mathbb{P}_{\mathbb{C}}^{(p+1)(\ell+1)-1}) & \xrightarrow{\cong} & H_q(\pi_{\mathbf{Y}}(A) \times \mathbb{P}_{\mathbb{C}}^{(p+1)(\ell+1)-1}) \end{array}$$

The isomorphism on the top level comes from the fact that homology and direct limit commute [22]. For the bottom isomorphism, we need the additional fact that since we assume that  $\pi_{\mathbf{Y}}$  is a compact covering we have

$$\varinjlim \{H_q(\pi_{\mathbf{Y}}(K) \times \mathbb{P}_{\mathbb{C}}^{(p+1)(\ell+1)-1}) \mid K \in \mathbf{K}(A)\} = \varinjlim \{H_q(L \times \mathbb{P}_{\mathbb{C}}^{(p+1)(\ell+1)-1}) \mid L \in \mathbf{K}(\pi_{\mathbf{Y}}(A))\}.$$

This proves that the right vertical arrow is also an isomorphism.  $\square$

Using the same notation as in Theorem 3.21 and Eqn (3.2) we have the following easy corollary of Theorem 3.21.

**Corollary 3.22.** *Let  $p = 2m + 1$  with  $m \geq 0$ . Then*

$$(3.10) \quad Q_{\pi_{\mathbf{Y}}(A)} = (1 - T) Q_{J_{\mathbf{C}, \mathbf{Y}}^p(A)} \pmod{T^{m+1}}.$$

*Proof.* The corollary follows directly from Theorem 3.21 and the fact that for any polynomial  $P \in \mathbb{Z}[T]$  we have

$$\begin{aligned} ((1 - T^2)P)^{\text{even}} &= (1 - T)(P)^{\text{even}}, \\ ((1 - T^2)P)^{\text{odd}} &= (1 - T)(P)^{\text{odd}}. \end{aligned}$$

□

As before we need a slightly more general version of Theorem 3.21 as well as Corollary 3.22.

Let  $\alpha_0, \dots, \alpha_\omega \geq 0$ , and  $N = \prod_{0 \leq j \leq m} (\alpha_j + 1)$ . Let  $\phi$  be a homogeneous formula defining a constructible subset of  $\mathbb{P}^{k_1} \times \dots \times \mathbb{P}_{\mathbf{C}}^{k_\sigma} \times \mathbb{P}_{\mathbf{C}}^\ell$ . Also, let for each  $i, 0 \leq i \leq \omega$ ,  $\Lambda^i \in \{\vee, \wedge\}$ , and let  $\Phi$  denote the multi-homogeneous formula defined by

$$\Phi \stackrel{\text{def}}{=} \bigwedge_{0 \leq i_0 \leq \alpha_0} \Lambda^0 \cdots \bigwedge_{0 \leq i_\omega \leq \alpha_\omega} \Lambda^\omega \phi(\mathbf{X}^1; \dots; \mathbf{X}^\sigma; \mathbf{Y}_{i_0, \dots, i_\omega}).$$

Let

$$S = \mathcal{R}(\Phi) \subset \mathbb{P}^{k_1} \times \dots \times \mathbb{P}_{\mathbf{C}}^{k_\sigma} \times \underbrace{\mathbb{P}_{\mathbf{C}}^\ell \times \dots \times \mathbb{P}_{\mathbf{C}}^\ell}_{N \text{ times}},$$

and let  $\pi_{\mathbf{Y}} : \mathbb{P}_{\mathbf{C}}^{k_1} \times \dots \times \mathbb{P}_{\mathbf{C}}^{k_\sigma} \times \underbrace{\mathbb{P}_{\mathbf{C}}^\ell \times \dots \times \mathbb{P}_{\mathbf{C}}^\ell}_{N \text{ times}} \rightarrow \mathbb{P}_{\mathbf{C}}^{k_1} \times \dots \times \mathbb{P}_{\mathbf{C}}^{k_\sigma}$  be the projection

along the  $\mathbf{Y}$ -co-ordinates.

For  $p > 0$ , let

$$J_{\mathbf{C}, \mathbf{Y}}^p(S) \subset \mathbb{P}^{k_1} \times \dots \times \mathbb{P}_{\mathbf{C}}^{k_\sigma} \times \underbrace{\mathbb{P}_{\mathbf{C}}^{(\ell+1)(p+1)-1} \times \dots \times \mathbb{P}_{\mathbf{C}}^{(\ell+1)(p+1)}}_{N \text{ times}}$$

be defined by the formula

$$J_{\mathbf{C}, \mathbf{Y}}^p(\Phi) \stackrel{\text{def}}{=} \bigwedge_{0 \leq i_0 \leq \alpha_0} \Lambda^0 \cdots \bigwedge_{0 \leq i_\omega \leq \alpha_\omega} \Lambda^\omega J_{\mathbf{C}, \mathbf{Y}_{i_0, \dots, i_\omega}}^p \phi(\mathbf{X}^1; \dots; \mathbf{X}^\sigma; \mathbf{Y}_{i_0, \dots, i_\omega}).$$

**Theorem 3.23.** *For every  $p \geq 0$ , we have that*

$$(3.11) \quad P_{\pi_{\mathbf{Y}}(S)} = (1 - T^2)^N P_{J_{\mathbf{C}, \mathbf{Y}}^p(S)} \pmod{T^p}.$$

*Proof.* The proof is identical to that of Theorem 3.21 above using Proposition 3.18 instead of Proposition 3.15 and noticing that by the Kunneth formula for homology, the Poincaré polynomial of

$$\underbrace{\mathbb{P}_{\mathbf{C}}^{(\ell+1)(p+1)-1} \times \dots \times \mathbb{P}_{\mathbf{C}}^{(\ell+1)(p+1)}}_{N \text{ times}}$$

equals  $(1 - T^2)^{-N} \pmod{T^p}$ . □

As before we have the following corollary.

**Corollary 3.24.** *Let  $p = 2m + 1$  with  $m \geq 0$ . Then*

$$(3.12) \quad Q_{\pi_{\mathbf{Y}}(S)} = (1 - T)^N Q_{J_{\mathbf{C}, \mathbf{Y}}^p(S)} \pmod{T^m}.$$

It is clear from the definition that the complex joins of sets in  $\mathbf{P}_{\mathbf{C}}$  also belong to  $\mathbf{P}_{\mathbf{C}}$ . We record this observation in the following formal proposition.

**Proposition 3.25** (Polynomial time membership testing). *Suppose that the sequence of constructible sets  $(S_n \subset \mathbb{P}_C^{k(n)} \times \mathbb{P}_C^{\ell(n)})_{n>0} \in \mathbf{P}_C$ , and  $\mathbf{X}_n = (X_0 : \dots : X_{k(n)})$   $\mathbf{Y}_n = (Y_0 : \dots : Y_{\ell(n)})$  are homogeneous co-ordinates of  $\mathbb{P}_C^{k(n)}$  and  $\mathbb{P}_C^{\ell(n)}$  respectively. Let  $p(n)$  be a polynomial. Then,*

$$\left( J_{C, \mathbf{Y}_n}^{p(n)}(S_n) \subset \mathbb{P}_C^{k(n)} \times \mathbb{P}_C^{(p(n)+1)(\ell(n)+1)-1} \right)_{n>0} \in \mathbf{P}_C.$$

*Proof.* Obvious from the definition of  $(J_{C, \mathbf{Y}_n}^{p(n)}(S_n))_{n>0}$ . □

#### 4. PROOF OF THE MAIN THEOREM

We are now in a position to prove Theorem 2.1. The proof relies on the following key proposition.

**Proposition 4.1.** *Let  $m(n), k_1(n), \dots, k_\omega(n)$  be polynomials, and let*

$$(\Phi_n(\mathbf{X}, \mathbf{Y}))_{n>0}$$

*be a sequence of multi-homogeneous formulas*

$$\Phi_n(\mathbf{X}, \mathbf{Y}) \stackrel{\text{def}}{=} (Q_1 \mathbf{Z}^1 \in \mathbb{P}_C^{k_1}) \cdots (Q_\omega \mathbf{Z}^\omega \in \mathbb{P}_C^{k_\omega}) \phi_n(\mathbf{X}; \mathbf{Y}; \mathbf{Z}^1; \dots; \mathbf{Z}^\omega),$$

*having free variables  $(\mathbf{X}; \mathbf{Y}) = (X_0, \dots, X_{k(n)}; Y_0, \dots, Y_{m(n)})$ , with*

$$Q_1, \dots, Q_\omega \in \{\exists, \forall\},$$

*and  $\phi_n$  a multi-homogeneous quantifier-free formula defining a closed (resp. open) constructible subset*

$$S_n \subset \mathbb{P}_C^k \times \mathbb{P}_C^m \times \mathbb{P}_C^{k_1} \times \cdots \times \mathbb{P}_C^{k_\omega}.$$

*Suppose also that*

$$(\mathcal{R}(\phi_n(\mathbf{X}; \mathbf{Y}; \mathbf{Z}^1; \dots; \mathbf{Z}^\omega)))_{n>0} \in \mathbf{P}_C.$$

*Then there exists:*

(A) *a sequence of quantifier-free multi-homogeneous formulas*

$$(\Theta_n(\mathbf{X}; \mathbf{V}^1; \dots; \mathbf{V}^N))_{n>0},$$

*with  $\mathbf{V}^i = (V_0, \dots, V_{p_i})$ , and  $N, p_1, \dots, p_N$  polynomials in  $n$ , such that  $\Theta_n(\mathbf{x}; \mathbf{V}^1; \dots; \mathbf{V}^N)$  defines a constructible subset  $T_n \subset \mathbb{P}_C^{p_1} \times \cdots \times \mathbb{P}_C^{p_N}$ , with*

$$(T_n)_{n>0} \in \mathbf{P}_C;$$

(B) *polynomial time computable maps*

$$F_n : \mathbb{Z}[T] \rightarrow \mathbb{Z}[T],$$

*such that*

$$Q_{\mathcal{R}(\Phi_n(\mathbf{x}; \cdot))} = F_n(Q_{\mathcal{R}(\Theta_n(\mathbf{x}; \mathbf{V}^1; \dots; \mathbf{V}^N))}).$$

The idea behind the proof of Proposition 4.1 is to use induction on the number,  $\omega$ , of quantifier blocks. When  $\omega = 0$ , the proposition is obvious. When  $\omega > 0$ , then using Corollary 3.22, we can construct a new formula (say  $\Phi'_n$ ) such that  $\Phi'$  has one less block of quantifiers, but such that  $Q_{\mathcal{R}(\Phi'_n)}$  is easily computable from  $Q_{\mathcal{R}(\Phi_n)}$ . One can then use induction to finish the proof. However, a technical complication arises due to the fact that in the projective situation (unlike in the affine situation) we cannot immediately replace two adjacent blocks of the same quantifier by a single block. This is the logical manifestation of the elementary

geometric fact that (unlike affine spaces) the product of two projective spaces is not itself a projective space. In order to overcome this difficulty and carry through the inductive step properly, we need to prove a slightly stronger, but technically more involved proposition, which we state next. Proposition 4.1 will be an immediate corollary of this more general proposition.

**Proposition 4.2.** *Let  $\sigma, \omega \geq 0$  be constants, and*

$$a_0(n), \alpha_0(n), a_1(n), \alpha_1(n), \dots, a_\sigma(n), \alpha_\sigma(n), \\ k(n), k_1(n), \dots, k_\omega(n)$$

*fixed polynomials in  $n$  taking non-negative values for  $n \in \mathbb{N}$ .*

*For  $0 \leq j \leq \sigma$ , let  $\mathbf{W}^j$  denote the tuple of variables*

$$(\dots, \mathbf{W}_{i_0, \dots, i_j}^j, \dots), 0 \leq i_0 \leq \alpha_0, \dots, 0 \leq i_j \leq \alpha_j,$$

*where each  $\mathbf{W}_{i_0, \dots, i_j}^j$  is a block of  $a_j(n) + 1$  variables.*

*Let*

$$(\Phi_n(\mathbf{X}; \mathbf{W}^0; \mathbf{W}^1; \dots; \mathbf{W}^\sigma))_{n>0},$$

*be a sequence of multi-homogeneous formulas defined by*

$$\Phi_n(\mathbf{X}; \mathbf{W}^0; \mathbf{W}^1; \dots; \mathbf{W}^\sigma) \stackrel{def}{=} \\ \Lambda_{0 \leq i_0 \leq \alpha_0}^0 \cdots \Lambda_{0 \leq i_n \leq \alpha_\sigma}^\sigma (Q_1 \mathbf{Z}^1 \in \mathbb{P}_C^{k_1}) \cdots (Q_\omega \mathbf{Z}^\omega \in \mathbb{P}_C^{k_\omega}) \\ \phi_n(\mathbf{X}; \mathbf{W}^0; \mathbf{W}^1; \dots; \mathbf{W}^\sigma; \mathbf{Z}^1; \dots; \mathbf{Z}^\omega),$$

*with*

$$\Lambda^0, \dots, \Lambda^\sigma \in \{\bigvee, \bigwedge\},$$

$$Q_1, \dots, Q_\omega \in \{\exists, \forall\},$$

*and each  $\phi_n$  a multi-homogeneous quantifier-free formula, multi-homogeneous in the blocks of variables,  $\mathbf{X}, \mathbf{Z}^1, \dots, \mathbf{Z}^\omega$  and  $(\mathbf{W}_{i_0, \dots, i_j, 0}^j, \dots, \mathbf{W}_{i_0, \dots, i_j, \alpha_j}^j)$  for  $0 \leq j \leq \sigma$ . Suppose also that each  $\phi_n$  defines a closed (resp. open) constructible set, and that*

$$(\mathcal{R}(\phi_n))_{n>0} \in \mathbf{P}_C.$$

*Then there exists:*

(A) *a sequence of quantifier-free multi-homogeneous formulas*

$$(\Theta_n(\mathbf{X}; \mathbf{V}^1; \dots; \mathbf{V}^N))_{n>0},$$

*with  $\mathbf{V}^i = (V_0, \dots, V_{p_i})$ , and  $N, p_1, \dots, p_N$  polynomials in  $n$ , such that  $\Theta_n(\mathbf{x}; \mathbf{V}^1; \dots; \mathbf{V}^N)$  defines a constructible subset  $T_n \subset \mathbb{P}_C^{p_1} \times \dots \times \mathbb{P}_C^{p_N}$ , with*

$$(T_n)_{n>0} \in \mathbf{P}_C;$$

(B) *polynomial time computable maps*

$$F_n : \mathbb{Z}[T] \rightarrow \mathbb{Z}[T],$$

*such that*

$$Q_{\mathcal{R}(\Phi_n(\mathbf{x}; \cdot))} = F_n(Q_{\mathcal{R}(\Theta_n(\mathbf{x}; \mathbf{V}^1; \dots; \mathbf{V}^N))}).$$

*Proof of Proposition 4.2.* The proof is by induction on  $\omega$ .

If  $\omega = 0$ , we let  $\Theta_n = \Phi_n$ , and  $F_n$  to be the identity map. Since there are no quantifiers, for each  $n \geq 0$  the constructible set defined by  $\Theta_n$  and  $\Phi_n$  are the same, and thus the Betti numbers of the sets defined by  $\Theta_n$  and  $\Phi_n$  are equal.

If  $\omega > 0$ , we have the following two cases.

(A) Case 1,  $Q_1 = \exists$ : First note that  $\Phi_n$  defines a constructible subset of  $\mathbb{P}_{\mathbb{C}}^{k(n)} \times U_n$ , where

$$U_n = (\mathbb{P}_{\mathbb{C}}^{(a_0+1)(\alpha_0+1)-1})^{m_0} \cdots \times (\mathbb{P}_{\mathbb{C}}^{(a_j+1)(\alpha_j+1)-1})^{m_j} \times \cdots \times (\mathbb{P}_{\mathbb{C}}^{(a_\sigma+1)(\alpha_\sigma+1)-1})^{m_\sigma},$$

where for  $0 \leq j \leq \sigma$ ,

$$m_j(n) = \prod_{0 \leq i \leq j-1} (\alpha_i(n) + 1).$$

The formula  $\Phi_n$  is equivalent to the following formula.

$$\bar{\Phi}_n \stackrel{\text{def}}{=} \cdots (\exists \mathbf{Z}^{1, i_0, \dots, i_\sigma} \in \mathbb{P}_{\mathbb{C}}^{k_1}) \cdots \Lambda_{0 \leq i_0 \leq \alpha_0}^0 \cdots \Lambda_{0 \leq i_\sigma \leq \alpha_\sigma}^\sigma \\ (\mathbb{Q}_2 \mathbf{Z}^2 \in \mathbb{P}_{\mathbb{C}}^{k_2}) \cdots (\mathbb{Q}_\omega \mathbf{Z}^\omega \in \mathbb{P}_{\mathbb{C}}^{k_\omega}) \phi_n(\mathbf{X}; \mathbf{W}^0; \dots, \mathbf{W}^\sigma; \mathbf{Z}^{1, i_0, \dots, i_\sigma}; \mathbf{Z}^2 \dots; \mathbf{Z}^\omega),$$

where the blocks of existential quantifiers in the beginning are indexed by the tuples

$$(i_0, \dots, i_\sigma), 0 \leq i_0 \leq \alpha_0(n), \dots, 0 \leq i_\sigma \leq \alpha_\sigma(n).$$

and the number of such blocks is

$$\alpha(n) = \prod_{i=0}^{\sigma} (\alpha_i(n) + 1).$$

Let

$$m(n) = \sum_{j=0}^{\sigma} m_j(n) ((a_j(n) + 1)(\alpha_j(n) + 1) - 1).$$

(Note that  $m(n)$  is the (complex) dimension of  $U_n$  defined previously.)

Consider the sequence

$$\left( J_{\mathbb{C}, \mathbf{Z}^1}^{2m(n)+1}(\bar{\Phi}_n) \right)_{n>0},$$

where

$$J_{\mathbb{C}, \mathbf{Z}^1}^{2m(n)+1}(\bar{\Phi}_n) \stackrel{\text{def}}{=} \Lambda_{0 \leq i_0 \leq \alpha_0}^0 \cdots \Lambda_{0 \leq i_\sigma \leq \alpha_\sigma}^\sigma \\ J_{\mathbb{C}, \mathbf{Z}^{1, i_0, \dots, i_\sigma}}^{2m+1}((\mathbb{Q}_2 \mathbf{Z}^2 \in \mathbb{P}_{\mathbb{C}}^{k_2}) \cdots (\mathbb{Q}_\omega \mathbf{Z}^\omega \in \mathbb{P}_{\mathbb{C}}^{k_\omega}) \phi_n(\mathbf{X}; \mathbf{W}^0; \dots; \mathbf{W}^\sigma; \mathbf{Z}^{1, i_0, \dots, i_\sigma}; \mathbf{Z}^2 \dots; \mathbf{Z}^\omega)).$$

Note that the formula

$$J_{\mathbb{C}, \mathbf{Z}^1}^{2m(n)+1}(\bar{\Phi}_n) = \Lambda_{0 \leq i_0 \leq \alpha_0}^0 \cdots \Lambda_{0 \leq i_\sigma \leq \alpha_\sigma}^\sigma \Lambda_{0 \leq i_{\sigma+1} \leq \alpha_{\sigma+1}}^{\sigma+1} \\ (\mathbb{Q}_2 \mathbf{Z}^2 \in \mathbb{P}_{\mathbb{C}}^{k_2}) \cdots (\mathbb{Q}_\omega \mathbf{Z}^\omega \in \mathbb{P}_{\mathbb{C}}^{k_\omega}) \\ \phi_n(\mathbf{X}; \mathbf{W}^0; \dots; \mathbf{W}^\sigma; \mathbf{W}^{\sigma+1}; \mathbf{Z}^2; \dots; \mathbf{Z}^\omega).$$

with  $\Lambda^{\sigma+1} = \bigwedge$ ,  $\alpha_{\sigma+1} = 2m + 1$ , and  $\mathbf{W}^{\sigma+1} = \mathbf{Z}^{1, i_0, \dots, i_\sigma, i_{\sigma+1}}$ .

Observe that the number of quantifiers in the formulas  $J_{\mathbb{C}, \mathbf{Z}^1}^{2m(n)+1}(\bar{\Phi}_n)$ , is  $\omega - 1$ .

Moreover,  $J_{\mathbb{C}, \mathbf{Z}^1}^{2m(n)+1}(\bar{\Phi}_n)$  satisfy by Proposition 3.25 the required polynomial time hypothesis, and has the same shape as the formulas  $\Phi_n$ . We can thus apply the induction hypothesis to this sequence to obtain a sequence  $(\Theta_n)_{n>0}$ , as well as a sequence of polynomial time computable maps  $(G_n)_{n>0}$ . By inductive hypothesis we can suppose that for each  $\mathbf{x} \in \mathbb{P}_{\mathbb{C}}^{k(n)}$

$$Q_{\mathcal{R}(J_{\mathbb{C}, \mathbf{Z}^1}^{2m(n)+1}(\bar{\Phi}_n)(\mathbf{x}; \cdot))} = G_n(Q_{\mathcal{R}(\Theta_n(\mathbf{x}; \cdot))}).$$

Using Corollary 3.22 we have

$$\begin{aligned} Q_{\mathcal{R}(\Phi_n(\mathbf{x}; \cdot))} &= (1 - T)^{\alpha(n)} Q_{\mathcal{R}(J_{\mathbb{C}, \mathbf{Z}^1}^{2m(n)+1}(\bar{\Phi}_n)(\mathbf{x}; \cdot))} \pmod{T^{m(n)+1}} \\ &= (1 - T)^{\alpha(n)} G_n(Q_{\mathcal{R}(\Theta_n(\mathbf{x}; \cdot))}) \pmod{T^{m(n)+1}}. \end{aligned}$$

We set

$$F_n = \text{Trunc}_{m(n)} \circ M_{(1-T)^{\alpha(n)}} \circ G_n$$

(see Notation 3.2). This completes the induction in this case.

(B) Case 2,  $Q_1 = \forall$ :

The formula  $\Phi_n$  is equivalent to the following formula.

$$\begin{aligned} \bar{\Phi}_n &\stackrel{\text{def}}{=} \dots (\forall \mathbf{Z}^{1, i_0, \dots, i_\sigma} \in \mathbb{P}_{\mathbb{C}}^{k_1}) \dots \\ &\quad \bigwedge_{0 \leq i_0 \leq \alpha_0}^0 \dots \bigwedge_{0 \leq i_\sigma \leq \alpha_\sigma}^\sigma \\ &(\mathbb{Q}_2 \mathbf{Z}^2 \in \mathbb{P}_{\mathbb{C}}^{k_2}) \dots (\mathbb{Q}_\omega \mathbf{Z}^\omega \in \mathbb{P}_{\mathbb{C}}^{k_\omega}) \phi_n(\mathbf{X}; \mathbf{W}^0; \dots, \mathbf{W}^\sigma; \mathbf{Z}^{1, i_0, \dots, i_\sigma}; \mathbf{Z}^2 \dots; \mathbf{Z}^\omega). \end{aligned}$$

where the blocks of universal quantifiers in the beginning are indexed by the tuples

$$(i_0, \dots, i_\sigma), 0 \leq i_0 \leq \alpha_0(n), \dots, 0 \leq i_\sigma \leq \alpha_\sigma(n).$$

Consider the sequence

$$\left( J_{\mathbb{C}, \mathbf{Z}^1}^{2m+1}(-\bar{\Phi}_n) \right)_{n>0},$$

where

$$\begin{aligned} &J_{\mathbb{C}, \mathbf{Z}^1}^{2m+1}(-\bar{\Phi}_n) \stackrel{\text{def}}{=} \\ &\neg \bigwedge_{0 \leq i_0 \leq \alpha_0}^0 \dots \bigwedge_{0 \leq i_\sigma \leq \alpha_\sigma}^\sigma \\ &J_{\mathbb{C}, \mathbf{Z}^{1, i_0, \dots, i_\sigma}}^{2m+1}((\mathbb{Q}_2 \mathbf{Z}^2 \in \mathbb{P}_{\mathbb{C}}^{k_2}) \dots (\mathbb{Q}_\omega \mathbf{Z}^\omega \in \mathbb{P}_{\mathbb{C}}^{k_\omega}) \phi_n(\mathbf{X}; \mathbf{W}^0; \dots; \mathbf{W}^\sigma; \mathbf{Z}^{1, i_0, \dots, i_\sigma}; \mathbf{Z}^2 \dots; \mathbf{Z}^\omega)). \end{aligned}$$

Note that the formula

$$\begin{aligned} &J_{\mathbb{C}, \mathbf{Z}^1}^{2m+1}(\bar{\Phi}_n) = \\ &\quad \bar{\bigwedge}_{0 \leq i_0 \leq \alpha_0}^0 \dots \bar{\bigwedge}_{0 \leq i_\sigma \leq \alpha_\sigma}^\sigma \bigwedge_{0 \leq i_{\sigma+1} \leq \alpha_{\sigma+1}}^{\sigma+1} \\ &\quad (\bar{\mathbb{Q}}_2 \mathbf{Z}^2 \in \mathbb{P}_{\mathbb{C}}^{k_2}) \dots (\bar{\mathbb{Q}}_\omega \mathbf{Z}^\omega \in \mathbb{P}_{\mathbb{C}}^{k_\omega}) \\ &\quad \neg \phi_n(\mathbf{X}; \mathbf{W}^0; \dots; \mathbf{W}^\sigma; \mathbf{W}^{\sigma+1}; \mathbf{Z}^2; \dots; \mathbf{Z}^\omega). \end{aligned}$$

with  $\Lambda^{\sigma+1} = \bigwedge$ ,  $\alpha_{\sigma+1} = 2m + 1$ , and  $\mathbf{W}^{\sigma+1} = \mathbf{Z}^{1, i_0, \dots, i_{\sigma}, i_{\sigma+1}}$ , and

$$\begin{aligned}\bar{\Lambda}^i &= \bigvee \text{ if } \Lambda^i = \bigwedge, \bar{\Lambda}^i = \bigwedge \text{ if } \Lambda^i = \bigvee, \\ \bar{Q}_i &= \exists \text{ if } Q_i = \forall, \bar{Q}_i = \forall \text{ if } Q_i = \exists.\end{aligned}$$

Observe that the number of quantifiers in the formulas  $J_{\mathbf{C}, \mathbf{Z}^1}^{2m+1}(\bar{\Phi}_n)$ , is  $\omega - 1$ .

Moreover,  $J_{\mathbf{C}, \mathbf{Z}^1}^{2m+1}(\bar{\Phi}_n)$  satisfy by Proposition 3.25 the required polynomial time hypothesis, and have the same shape as the formulas  $\Phi_n$ . We can thus apply the induction hypothesis to this sequence to obtain a sequence  $(\Theta_n)_{n>0}$ , as well as a sequence of polynomial time computable maps  $(G_n)_{n>0}$ . By inductive hypothesis we can suppose that for each  $\mathbf{x} \in \mathbb{P}_{\mathbf{C}}^{k(n)}$

$$Q_{\mathcal{R}(J_{\mathbf{C}, \mathbf{Z}^1}^{2m+1}(\bar{\Phi}_n)(\mathbf{x}; \cdot))} = G_n(Q_{\mathcal{R}(\Theta_n(\mathbf{x}; \cdot))}).$$

Using Corollary 3.22 we have

$$\begin{aligned}Q_{\mathcal{R}(\bar{\Phi}_n(\mathbf{x}; \cdot))} &= (1 - T)^{\alpha(n)} Q_{\mathcal{R}(J_{\mathbf{C}, \mathbf{Z}^1}^{2m+1}(\bar{\Phi}_n)(\mathbf{x}; \cdot))} \pmod{T^{m(n)+1}} \\ &= (1 - T)^{\alpha(n)} G_n(Q_{\mathcal{R}(\Theta_n(\mathbf{x}; \cdot))}) \pmod{T^{m(n)+1}}.\end{aligned}$$

The sets  $K_n = \mathcal{R}(\Phi_n(\mathbf{x}; \cdot))$  are constructible; so by Corollary 3.4 (corollary to Theorem 3.1), we have

$$Q_{K_n}(T) = -\text{Rec}_m(\text{Trunc}_m(Q_{U_n - K_n})) + P_{U_n}.$$

We set  $F_n$  to be the operator defined by

$$F_n(Q) = -\text{Rec}_m(\text{Trunc}_m(M_{(1-T)^{\alpha(n)}}(G_n(Q)))) + P_{U_n}.$$

This completes the induction in this case as well.  $\square$

*Proof of Proposition 4.1.* Proposition 4.1 is a special case of Proposition 4.2 with  $\sigma = 0$ ,  $\alpha_0 = 0$ , and  $\mathbf{Y} = \mathbf{W}^0$ .  $\square$

*Proof of Theorem 2.1.* Follows immediately from Proposition 4.1 in the special case  $m = 0$ . In this case the sequence of formulas  $(\Phi_n)_{n>0}$  correspond to a language in the polynomial hierarchy and for each  $n$ ,  $\mathbf{x} = (x_0 : \dots : x_{k(n)}) \in S_n \subset \mathbb{P}_{\mathbf{C}}^{k(n)}$  if and only if

$$F_n(Q_{\mathcal{R}(\Theta_n(\mathbf{x}; \cdot))})(0) > 0$$

and this last condition can be checked in polynomial time with advice from the class  $\#\mathbf{P}_{\mathbf{C}}^{\dagger}$ .  $\square$

*Remark 4.3.* It is interesting to observe that in complete analogy with the proof of the classical Toda's theorem the proof of Theorem 2.1 also requires just one call to the oracle at the end.

*Proof of Theorem 2.5.* Follows from the proof of Proposition 4.1 since the formula  $\Theta_n$  is clearly computable in polynomial time from the given formula  $\Phi_n$  as long as the number of quantifier alternations  $\omega$  is bounded by a constant.  $\square$

## 5. ACKNOWLEDGEMENTS

The author benefited from discussions with Donu Arapura, Andrei Gabrielov, and James McClure at various stages of this work.

## REFERENCES

1. S. Basu, R. Pollack, and M.-F. Roy, *Algorithms in real algebraic geometry*, Algorithms and Computation in Mathematics, vol. 10, Springer-Verlag, Berlin, 2006 (second edition). MR 1998147 (2004g:14064)
2. S. Basu and T. Zell, *Polynomial hierarchy, Betti numbers, and a real analogue of Toda's Theorem*, Found. Comput. Math. **to appear** (available at [arXiv.org:0812.1200](https://arxiv.org/abs/0812.1200)) (2009).
3. L. Blum, F. Cucker, M. Shub, and S. Smale, *Complexity and real computation*, Springer-Verlag, New York, 1998, With a foreword by Richard M. Karp. MR 99a:68070
4. L. Blum, M. Shub, and S. Smale, *On a theory of computation and complexity over the real numbers: NP-completeness, recursive functions and universal machines*, Bull. Amer. Math. Soc. (N.S.) **21** (1989), no. 1, 1–46. MR 90a:68022
5. Peter Bürgisser and Felipe Cucker, *On a Problem Posed by Steve Smale*, 2009.
6. P. Bürgisser and F. Cucker, *Variations by complexity theorists on three themes of Euler, Bézout, Betti, and Poincaré*, Complexity of computations and proofs (Jan Krajíček, ed.), Quad. Mat., vol. 13, Dept. Math., Seconda Univ. Napoli, Caserta, 2004, pp. 73–151. MR 2131406 (2006c:68053)
7. ———, *Counting complexity classes for numeric computations. II. Algebraic and semialgebraic sets*, J. Complexity **22** (2006), no. 2, 147–191. MR 2200367 (2007b:68059)
8. P. Bürgisser, F. Cucker, and M. Lotz, *Counting complexity classes for numeric computations. III. Complex projective sets*, Found. Comput. Math. **5** (2005), no. 4, 351–387. MR 2189543 (2006h:68039)
9. J. H. Davenport and J. Heintz, *Real quantifier elimination is doubly exponential*, Journal of Symbolic Computation **5** (1988), no. 1/2, 29–35.
10. P. Deligne, *La conjecture de Weil. I*, Inst. Hautes Études Sci. Publ. Math. (1974), no. 43, 273–307. MR 0340258 (49 #5013)
11. ———, *La conjecture de Weil. II*, Inst. Hautes Études Sci. Publ. Math. (1980), no. 52, 137–252. MR 601520 (83c:14017)
12. B. Dwork, *On the rationality of the zeta function of an algebraic variety*, American Journal of Mathematics **82** (1960), no. 3, 631–648.
13. A. Gabrielov, N. Vorobjov, and T. Zell, *Betti numbers of semialgebraic and sub-Pfaffian sets*, J. London Math. Soc. (2) **69** (2004), no. 1, 27–43. MR 2025325 (2004k:14105)
14. H. Blaine Lawson, Jr., *Algebraic cycles and homotopy theory*, Ann. of Math. (2) **129** (1989), no. 2, 253–291. MR 986794 (90h:14008)
15. J. Matoušek, *Using the Borsuk-Ulam theorem*, Universitext, Springer-Verlag, Berlin, 2003, Lectures on topological methods in combinatorics and geometry, Written in cooperation with Anders Björner and Günter M. Ziegler. MR 1988723 (2004i:55001)
16. K. Meer, *Counting problems over the reals*, Theoret. Comput. Sci. **242** (2000), no. 1-2, 41–58. MR 1769145 (2002g:68041)
17. C. Papadimitriou, *Computational complexity*, Addison-Wesley, 1994.
18. Peter Scheiblechner, *On the complexity of deciding connectedness and computing Betti numbers of a complex algebraic variety*, J. Complexity **23** (2007), no. 3, 359–379. MR 2330991 (2009d:14020)
19. U. Schöning, *Probabilistic complexity classes and lowness*, J. Comput. System Sci. **39** (1989), no. 1, 84–100. MR 1013721 (91b:68041a)
20. Igor R. Shafarevich, *Basic algebraic geometry. 1*, second ed., Springer-Verlag, Berlin, 1994, Varieties in projective space, Translated from the 1988 Russian edition and with notes by Miles Reid. MR 1328833 (95m:14001)
21. M. Shub and S. Smale, *On the intractability of Hilbert's Nullstellensatz and an algebraic version of “NP ≠ P?”*, Duke Math. J. **81** (1995), no. 1, 47–54 (1996), A celebration of John F. Nash, Jr. MR 1381969 (97h:03067)
22. Edwin H. Spanier, *Algebraic topology*, McGraw-Hill Book Co., New York, 1966. MR 0210112 (35 #1007)

23. L. Stockmeyer, *The polynomial-time hierarchy*, Theoret. Comput. Sci. **3** (1976), no. 1, 1–22 (1977). MR 0438810 (55 #11716)
24. S. Toda, *PP is as hard as the polynomial-time hierarchy*, SIAM J. Comput. **20** (1991), no. 5, 865–877. MR 1115655 (93a:68047)
25. L. G. Valiant and V. V. Vazirani, *NP is as easy as detecting unique solutions*, Theoret. Comput. Sci. **47** (1986), no. 1, 85–93. MR 871466 (88i:68021)
26. A. Weil, *Number of solutions of equations over finite fields*, Bulletin of the American Mathematical Society **55** (1949), 497–508.

DEPARTMENT OF MATHEMATICS, PURDUE UNIVERSITY, WEST LAFAYETTE, IN 47906, U.S.A.  
*E-mail address:* `sbasu@math.purdue.edu`