
Preprint No. M 14/05

**Copositivity tests based on the
linear complementarity problem**

Carmo Bras, Gabriele Eichfelder and Joaquim
Judice

28. August 2014

Impressum:

Hrsg.: Leiter des Instituts für Mathematik
Weimarer Straße 25
98693 Ilmenau
Tel.: +49 3677 69-3621
Fax: +49 3677 69-3270
<http://www.tu-ilmenau.de/math/>

Copositivity tests based on the Linear Complementarity Problem

Carmo Brás¹, Gabriele Eichfelder² and Joaquim Júdice³

August 28, 2014

Abstract

Copositivity tests are presented based on new necessary and sufficient conditions requiring the solution of linear complementarity problems (LCP). Methodologies involving Lemke’s method, an enumerative algorithm and a linear mixed-integer programming formulation are proposed to solve the required LCPs. A new necessary condition for (strict) copositivity based on solving a Linear Program (LP) is also discussed, which can be used as a preprocessing step. The algorithms with these three different variants are thoroughly applied to test matrices from the literature and to max-clique instances with matrices up to dimension 496×496 . We compare our procedures with three other copositivity tests from the literature as well as with a general global optimization solver. The numerical results are very promising and equally good and in many cases better than the results reported elsewhere.

Key Words: conic optimization, copositivity, complementarity problems, linear programming

Mathematics subject classifications (MSC 2000): 15A48, 90C33, 65F30, 65K99, 90C26,

1 Introduction

The notion of (strict) copositivity of a matrix [11, 25] is well known in the area of linear complementarity problems (LCP) in the context of existence results and results on the successful termination of Lemke’s algorithm — a well known simplex-like vertex following algorithm for LCPs [11, 17, 25]. An $n \times n$ matrix is called \mathbb{R}_+^n -copositive (or \mathbb{R}_+^n -semidefinite or shortly copositive), if it generates a quadratic form

¹Faculdade de Ciências e Tecnologia, Universidade Nova de Lisboa, Portugal, mb@fct.unl.pt

²Institute of Mathematics, Technische Universität Ilmenau, Po 10 05 65, D-98684 Ilmenau, Germany, Gabriele.Eichfelder@tu-ilmenau.de

³Instituto de Telecomunicações, Portugal, joaquim.judice@co.it.pt

which takes no negative values on the nonnegative orthant \mathbb{R}_+^n . The strict copositive matrices are those for which the quadratic form is even positive on $\mathbb{R}_+^n \setminus \{0\}$.

In the last decade there has been an increasing interest in this property of a matrix, and in linear optimization problems over the cone of copositive matrices. For recent surveys on copositive programming see [2, 5, 10, 15, 19]. This interest is primarily based on the fact that some hard problems as the Maximum Clique problem (see [4, 13]) were shown to have a reformulation as a copositive program. Burer showed in [9] that, under weak assumptions, every quadratic program with linear constraints can be formulated as a copositive program, even if some of the variables are binary. Hence efficient numerical tests for a matrix on copositivity are essential.

The problem of determining whether a matrix is not copositive is NP-complete [26]. As discussed in [3] various authors have proposed such test, but there are only a few implemented numerical algorithms which apply to general symmetric matrices without any structural assumptions or dimensional restrictions and which are not just recursive, i.e., do not rely on information taken from all principal submatrices. There are some quite recent implementations which satisfy both criteria to full extend: in [7, 8] the first algorithm was proposed by Bundfuss and Dür, see also the modification and improvements by Žilinskas and Dür [32], by Sponsel, Bundfuss and Dür [28] and by Tanaka and Yoshise [29]. Later, another algorithm was presented by Bomze and Eichfelder [3].

Both approaches combine necessary and sufficient criteria for copositivity with a branch-and-bound algorithm. The branching is done in a data driven way and consists in a partitioning of the standard simplex in subsimplices. For each one of the subsimplices it is tested whether a sufficient criteria for copositivity is satisfied or whether a necessary condition is violated. The approaches use thereby different necessary and sufficient criteria but both approaches obtain better results in verifying that a matrix is not copositive than proving that a matrix is copositive. Moreover, the algorithms are also in most cases more successful to show copositivity for matrices which are even strict copositive. The first approach by Bundfuss and Dür is thereby based on the evaluation of a set of inequalities on each subsimplex, while the second one by Bomze and Eichfelder requires to solve convex quadratic problems and linear problems as sufficient criteria.

We present in this paper a new numerical approach for testing a matrix on copositivity also without any assumptions on the matrix and without using information from submatrices. We give new necessary and sufficient conditions. These are based on the relation between the (global) quadratic optimization problem which consists in minimizing the quadratic form of the matrix over the standard simplex, which is equivalent to testing the matrix on copositivity, and a mathematical program with linear complementarity constraints (MPLCC). We derive conditions by studying some linear complementarity problems (LCP) which deliver feasible solutions for the MPLCC. These conditions require the determination of solutions of LCPs. We propose an enumerative algorithm [22], Lemke's algorithm [11, Chapter 4.4] and a mixed integer formulation (MIP) [21] for this purpose.

Some of the new necessary conditions are easy and fast to verify as they require for instance the application of Lemke's algorithm only. Thus these conditions can also

be evaluated as an additional criteria in each iteration of other copositivity tests as the ones mentioned above. Moreover, we use known preprocessing results and combine them with a new preprocessing step based on solving linear problems (LP). We test the derived procedures on some famous matrices from the literature as well as on maximum-clique instances from the DIMACS challenge and generated smaller instances from the maximum clique problem. These matrices are also used as test instances for the above mentioned approaches. The considered matrices have up to dimension 496×496 . The numerical results show that the procedures are quite efficient for showing that a matrix is not strictly copositive or even not copositive. A hybrid algorithm combining some of the procedures discussed in this paper has proven to be successful for all the instances. In particular, the algorithm has been able to establish that the maximum clique is a lower-bound for each one of the instances. More numerical effort is needed to verify that matrices are strictly copositive or copositive but not strictly copositive (which is a well known drawback also for the above mentioned approaches).

Instead of applying algorithms which were especially designed for testing a matrix on copositivity one could also directly apply a global optimization solver, as for instance BARON, to the quadratic optimization problem mentioned above. In this paper, for the first time, a numerical copositivity test is compared to a general global optimization solver. For the predefined allowed time BARON fails for five of the large instances while our proposed hybrid algorithm can solve all the instances.

The remaining of this article is structured as follows: in Section 2 we give necessary and sufficient conditions for copositivity based on a reformulation as a mathematical program with linear complementarity constraints (MPLCC). From that we derive conditions based on the solutions of LCPs which we again characterize by solutions of MIPs. We also give necessary conditions for copositivity based on LPs. In Section 3 we present the algorithms which consist of two basic steps 0 and 1 (preprocessing and applying Lemke's method) and a step 2 for which we present three different possible procedures. Numerical experiments with these techniques are reported in Section 4. In the last section we give some conclusions and an outlook on possible extensions of the proposed methods to be done in the future.

2 Sufficient and necessary conditions for copositivity

In this section we first recall some basic notations before we study the relation between a matrix to be copositive and the solutions of a mathematical program with linear complementarity constraints (MPLCC). We derive necessary and sufficient conditions based on linear complementarity problems (LCPs) and a mixed integer formulation of one of these LCPs. Finally, we also give some easy to verify necessary conditions for (strict) copositivity based on linear problems (LPs).

2.1 Copositivity and Global Optimization

We recall the definition of (strict) copositivity:

Definition 2.1. A real $n \times n$ matrix M is denoted copositive if $x^\top Mx \geq 0$ for all $x \in \mathbb{R}_+^n$, and strictly copositive if $x^\top Mx > 0$ for all $x \in \mathbb{R}_+^n \setminus \{0\}$.

Any strictly copositive matrix is copositive. As a real $n \times n$ matrix M is (strictly) copositive if and only if the symmetric matrix $\frac{1}{2}(M + M^\top)$ is (strictly) copositive, we restrict our examinations to symmetric matrices. Let \mathcal{S}^n denote the real linear space of real symmetric $n \times n$ matrices. The set of copositive symmetric $n \times n$ matrices is a convex cone and the interior of the cone of copositive matrices is the set of strictly copositive matrices (cf. [7, 16]). We denote the cone of copositive matrices by C and its interior, the set of strictly copositive matrices, by SC .

Definition 2.2. A real $n \times n$ matrix M is said to be a S_0 matrix ($M \in S_0$) if there exists a point $0 \neq x \geq 0$ such that $Mx \geq 0$.

Testing a matrix on copositivity is related to quadratic programming:

Lemma 2.3. Let $M \in \mathcal{S}^n$ and let \bar{x} be a (global) minimal solution of the quadratic optimization problem

$$\begin{aligned} QP: \quad \min \quad & f(x) := \frac{1}{2}x^\top Mx \\ \text{s.t.} \quad & e^\top x = 1 \\ & x \geq 0 \end{aligned} \tag{1}$$

where $e \in \mathbb{R}^n$ denotes the vector with all components equal to one. Then

- (a) $M \in C$ if and only if $f(\bar{x}) \geq 0$,
- (b) $M \in SC$ if and only if $f(\bar{x}) > 0$.
- (c) $M \notin C$ if and only if there exists a feasible x with $f(x) < 0$.
- (d) $M \notin SC$ if there exists a feasible x with $f(x) = 0$.

Note that QP (1) is solvable as the feasible set is nonempty, compact and the objective function is continuous. As the feasible set is given by linear constraints, any global minimal solution of QP (1) satisfies the KKT-conditions, i.e. there exists $\lambda_0 \in \mathbb{R}$ and $w \in \mathbb{R}^n$ such that

$$\begin{aligned} Mx &= \lambda_0 e + w \\ x &\geq 0, \quad w \geq 0 \\ x^\top w &= 0, \quad e^\top x = 1 \end{aligned}$$

We write $x \in K$ if there exists some $w \in \mathbb{R}^n$ and some $\lambda_0 \in \mathbb{R}$ such that (x, λ_0, w) satisfies the above conditions. For any $x \in K$ it holds

$$f(x) = \frac{1}{2}x^\top Mx = \frac{1}{2}(\lambda_0 e^\top x + w^\top x) = \frac{\lambda_0}{2}.$$

Therefore $\lambda_0 = x^\top Mx$. Hence we can consider the following Mathematical Program with Linear Complementarity Constraints:

$$\begin{aligned} \text{MPLCC:} \quad & \min \quad \frac{1}{2}\lambda_0 \\ & \text{s.t.} \quad w = Mx - \lambda_0 e \\ & \quad x \geq 0, w \geq 0 \\ & \quad x^\top w = 0, e^\top x = 1 \\ & \quad \lambda_0 \in \mathbb{R} \end{aligned} \tag{2}$$

If \bar{x} is a minimal solution of QP (1) then there exists some $\bar{\lambda}_0 \in \mathbb{R}$ and some $\bar{w} \in \mathbb{R}^n$ such that $(\bar{x}, \bar{\lambda}_0, \bar{w})$ is a feasible solution of MPLCC (2) with the same objective function value. On the other hand, any feasible point $(\bar{x}, \bar{\lambda}_0, \bar{w})$ of MPLCC (2) gives a feasible solution of QP (1) with the same objective function value. Hence both problems are equivalent in the sense that they have the same objective function value and a minimal solution of one problem directly gives a minimal solution of the other problem. We conclude from Lemma 2.3:

Corollary 2.4. *(a) $M \in C$ if and only if MPLCC (2) has a (global) minimal solution $(\bar{x}, \bar{\lambda}_0, \bar{w})$ with $\bar{\lambda}_0 \geq 0$.*

(b) $M \in SC$ if and only if MPLCC (2) has a (global) minimal solution $(\bar{x}, \bar{\lambda}_0, \bar{w})$ with $\bar{\lambda}_0 > 0$.

2.2 Conditions for copositivity based on LCPs

For solving the MPLCC (2) we study its feasible set which contains a linear complementarity problem (LCP). The general form of an LCP is given as follows:

$$\begin{aligned} \text{LCP:} \quad & \text{Find } x \in \mathbb{R}^n \text{ and } w \in \mathbb{R}^n \text{ such that} \\ & \quad w = q + Mx \\ & \quad x \geq 0, w \geq 0 \\ & \quad x^\top w = 0 \end{aligned} \tag{3}$$

where $M \in \mathcal{S}^n$ and $q \in \mathbb{R}^n$ are given. We also use the notation $\text{LCP}(q, M)$ for representing a LCP with a given vector q and matrix M . We denote a pair $(x, w) \in \mathbb{R}^n \times \mathbb{R}^n$ feasible for $\text{LCP}(q, M)$ if

$$w = q + Mx, \quad x \geq 0 \quad \text{and} \quad w \geq 0.$$

A point $(x, \lambda_0, w) \in \mathbb{R}^n \times \mathbb{R} \times \mathbb{R}^n$ is feasible for the MPLCC (2) if and only if $e^\top x = 1$ and (x, w) satisfies $\text{LCP}(-\lambda_0 e, M)$. Hence, a feasible solution of MPLCC (2) can be found by first determining a solution $\bar{x} \neq 0$ of $\text{LCP}(-\lambda_0 e, M)$ and then by setting

$$\tilde{x} := \frac{1}{e^\top \bar{x}} \bar{x}.$$

Thereby, it is enough to study the problem $\text{LCP}(-\lambda_0 e, M)$ for $\lambda_0 = 0$, a positive and a negative λ_0 to cover all cases of $\lambda_0 \in \mathbb{R}$ due to the following result:

Lemma 2.5. *Let $\bar{\lambda}_0 \in \mathbb{R}$ and $\tilde{\lambda}_0 \in \mathbb{R}$ be given. If $\bar{\lambda}_0 \cdot \tilde{\lambda}_0 > 0$ then the following holds:*

LCP($-\bar{\lambda}_0 e, M$) has a solution $\bar{x} \neq 0$ if and only if LCP($-\tilde{\lambda}_0 e, M$) has a solution

$$\frac{\bar{\lambda}_0}{\tilde{\lambda}_0} \bar{x} \neq 0 .$$

Proof. If LCP($-\bar{\lambda}_0 e, M$) has a solution $\bar{x} \neq 0$, then let $J := \{i \in \{1, \dots, n\} \mid \bar{x}_i > 0\}$ and $L := \{1, \dots, n\} \setminus J$. Then due to complementarity

$$\begin{aligned} 0 &= -\bar{\lambda}_0 e_J + M_{JJ} \bar{x}_J, & 0 &\leq -\bar{\lambda}_0 e_L + M_{LJ} \bar{x}_J, \\ \bar{x}_J &> 0, & \bar{x}_L &= 0. \end{aligned}$$

Hence \bar{x} also satisfies

$$\begin{aligned} 0 &= -\left(\frac{\tilde{\lambda}_0}{\bar{\lambda}_0} \bar{\lambda}_0\right) e_J + M_{JJ} \bar{x}_J, & 0 &\leq -\left(\frac{\tilde{\lambda}_0}{\bar{\lambda}_0} \bar{\lambda}_0\right) e_L + M_{LJ} \bar{x}_J, \\ \bar{x}_J &> 0, & \bar{x}_L &= 0. \end{aligned}$$

Multiplying with $\tilde{\lambda}_0/\bar{\lambda}_0$ yields

$$\begin{aligned} 0 &= -\tilde{\lambda}_0 e_J + M_{JJ} \left(\frac{\tilde{\lambda}_0}{\bar{\lambda}_0} \bar{x}_J\right), & 0 &\leq -\tilde{\lambda}_0 e_L + M_{LJ} \left(\frac{\tilde{\lambda}_0}{\bar{\lambda}_0}\right) \bar{x}_J, \\ \frac{\tilde{\lambda}_0}{\bar{\lambda}_0} \bar{x}_J &> 0, & \frac{\tilde{\lambda}_0}{\bar{\lambda}_0} \bar{x}_L &= 0. \end{aligned}$$

Hence $\frac{\tilde{\lambda}_0}{\bar{\lambda}_0} \bar{x}$ is a solution of LCP($-\tilde{\lambda}_0 e, M$). □

We conclude with Corollary 2.4 sufficient conditions for the (strict) copositivity and for M being not (strictly) copositive based on LCP($-\lambda_0 e, M$) for $\lambda_0 = 0$ and $\lambda_0 = -1$. Knowledge about nonzero solutions of LCP($-\lambda_0 e, M$) for $\lambda_0 > 0$ does not imply the strict copositivity of the matrix. It has to be guaranteed that there are no nonzero solutions for $\lambda_0 \leq 0$.

Corollary 2.6. (a) *If LCP($0, M$) has a solution $\bar{x} \neq 0$ then $M \notin SC$.*

(b) *If LCP(e, M) has a solution $\bar{x} \neq 0$ then $M \notin C$ and thus also $M \notin SC$.*

(c) *If no nonzero solution of LCP(e, M) and of LCP($0, M$) exists then $M \in SC$ and thus $M \in C$.*

(d) *If no nonzero solution of LCP(e, M) exists then $M \in C$.*

Example 2.7. *For the matrix*

$$M = \begin{pmatrix} 1 & -2 \\ -2 & 1 \end{pmatrix}$$

the point $\bar{x} = (1 \ 1)^\top$ with $\bar{w} = (0 \ 0)$ is a nonzero solution of LCP(e, M) and hence $M \notin C$ (and $M \notin SC$) by Corollary 2.6.

The solvability of the LCP for any q gives another necessary condition for strict copositivity.

Lemma 2.8. *[11, Theorem 3.8.5] If $M \in SC$, then for each $q \in \mathbb{R}^n$ the problem $LCP(q, M)$ has a solution.*

A solution can be found by Lemke's method, see for instance [11, Chapter 4.4], which is a simplex-like vertex following algorithm, that uses basic feasible solutions of a system of the form

$$w = q + x_0 d + Mx, \quad x \geq 0, \quad x_0 \geq 0, \quad w \geq 0 \quad (4)$$

where d is a positive vector (note that $x^\top w = 0$ in each iteration of the method). This method guarantees to terminate in a finite number of iterations if all the basic feasible solutions of the system (4) are nondegenerate. For the steps of Lemke's method see page 14 and for a discussion on the complexity of this algorithm see for instance [1]. The procedure either finds a solution of the LCP or it terminates in an unbounded ray. For some classes of matrices, including SC , this later termination can not occur and Lemke's method always terminates with a solution of LCP, cf. [11, Theorem 4.4.9]. We derive the following necessary condition for SC which we use in Step 1 of our algorithm:

Corollary 2.9. *If Lemke's method applied to $LCP(q, M)$ for some $q \in \mathbb{R}^n$ terminates in an unbounded ray then $M \notin SC$.*

If M is only a copositive matrix, then the $LCP(q, M)$ does not need to have a solution. For instance, this is the case for M the zero matrix and q negative.

Example 2.10. *For the matrix M of Example 2.7 Lemke's method for solving $LCP(-e, M)$ (i.e. $\lambda_0 = 1$ in $LCP(-\lambda_0 e, M)$) terminates in an unbounded ray. Hence, $M \notin SC$.*

The drawback of the sufficient conditions for (strict) copositivity of Corollary 2.6 is that one needs to guarantee that there is no nonzero solution of these LCPs, which is much more difficult than computing a solution for these problems. This indicates why in practice establishing strict copositivity (copositivity) is more difficult than showing that a matrix is not SC (C).

Finally, we study another LCP which gives a certificate for $M \notin SC$ and $M \notin C$. For that, let

$$p := \begin{pmatrix} 0 \\ -1 \end{pmatrix} \in \mathbb{R}^{n+1} \quad \text{and} \quad Q := \begin{pmatrix} M & e \\ e^\top & 0 \end{pmatrix} \in \mathbb{R}^{(n+1) \times (n+1)}. \quad (5)$$

Then $LCP(p, Q)$ is equivalent to find $(x, \mu_0) \in \mathbb{R}^{n+1}$ such there exists some $(w, r_0) \in \mathbb{R}^{n+1}$ with

$$\begin{aligned} \begin{pmatrix} w \\ r_0 \end{pmatrix} &= \begin{pmatrix} 0 \\ -1 \end{pmatrix} + \begin{pmatrix} M & e \\ e^\top & 0 \end{pmatrix} \begin{pmatrix} x \\ \mu_0 \end{pmatrix} \\ \begin{pmatrix} x \\ \mu_0 \end{pmatrix} &\geq 0, \quad \begin{pmatrix} w \\ r_0 \end{pmatrix} \geq 0 \\ \begin{pmatrix} x \\ \mu_0 \end{pmatrix}^\top \begin{pmatrix} w \\ r_0 \end{pmatrix} &= 0. \end{aligned} \quad (6)$$

The following result holds:

Lemma 2.11. (a) *LCP(p, Q) has a solution (x, μ₀) if and only if M ∉ SC.*

(b) *LCP(p, Q) has a solution (x, μ₀) with μ₀ > 0 if and only if M ∉ C.*

Proof. (a) If LCP(p, Q) has a solution then either $r_0 = 0$ and $\mu_0 \geq 0$ or $r_0 > 0$ and $\mu_0 = 0$. In the first case we immediately obtain that there is a feasible point $(x, \lambda_0, w) \in \mathbb{R}^n \times \mathbb{R} \times \mathbb{R}^n$ for MPLCC (2) with $\lambda_0 = -\mu_0 \leq 0$ and hence, by Corollary 2.4, $M \notin SC$. In the second case, there exists some $x \geq 0$ with $x \neq 0$ and $0 = x^\top w = x^\top Mx$ which implies $M \notin SC$. On the other hand, if $M \notin SC$, then there exists some feasible $(x, \lambda_0, w) \in \mathbb{R}^n \times \mathbb{R} \times \mathbb{R}^n$ with $\lambda_0 \leq 0$ for MPLCC (2) which delivers with $r_0 = 0$ and $\mu_0 = -\lambda_0 \geq 0$ a feasible solution for LCP(p, Q).

(b) First let LCP(p, Q) have a solution with $\mu_0 > 0$ and thus $r_0 = 0$. Then there is a feasible point $(x, \lambda_0, w) \in \mathbb{R}^n \times \mathbb{R} \times \mathbb{R}^n$ for MPLCC (2) with $\lambda_0 = -\mu_0 < 0$ and hence, by Corollary 2.4, $M \notin C$. On the other hand, if $M \notin C$, then there exists some feasible $(x, \lambda_0, w) \in \mathbb{R}^n \times \mathbb{R} \times \mathbb{R}^n$ with $\lambda_0 < 0$ for MPLCC (2) which delivers with $r_0 = 0$ and $\mu_0 = -\lambda_0 > 0$ such a feasible solution for LCP(p, Q).

□

Then we can state the following copositivity tests:

Corollary 2.12. (a) *If LCP(p, Q) has a solution (x, μ₀) with μ₀ = 0 then M ∉ SC.*

(b) *If LCP(p, Q) has a solution (x, μ₀) with μ₀ > 0 then M ∉ C and thus also M ∉ SC.*

(c) *If LCP(p, Q) has no solution, then M ∈ SC and thus M ∈ C.*

(d) *If LCP(p, Q) has no solution (x, μ₀) with μ₀ > 0, then M ∈ C.*

Example 2.13. *For the matrix M of Example 2.7 LCP(p, Q) has a solution (x, μ₀) with $x = (1/2, 1/2)^\top$ and $\mu_0 = 1/2 > 0$ and thus, by Corollary 2.12, $M \notin C$.*

In order to use Corollary 2.12(d) one needs to know all solutions of LCP(p, Q). For obtaining the complete solution set we will apply the enumerative algorithm described in [22]. We also exploit the following reformulation of the LCP(p, Q) as a mixed integer linear program (MIP) [21]:

Lemma 2.14. *The mixed integer linear program*

$$\begin{aligned} \text{MIP}_1 : \quad & \max \quad \alpha \\ & \text{s.t.} \quad 0 \leq Qy + \alpha p \leq z \\ & \quad \quad 0 \leq y \leq e - z \\ & \quad \quad z \in \{0, 1\}^{n+1} \\ & \quad \quad 0 \leq \alpha \leq 1. \end{aligned}$$

with p and Q as in (5) has a solution, and $(x^*, \mu_0^*) = \frac{y^*}{\alpha^*}$ is a solution, if and only if MIP_1 has a feasible solution (α^*, y^*, z^*) with $\alpha^* > 0$.

Proof. This result follows from Proposition 2.5 in [21] by noting that $p \neq 0$ and that it is enough that (α^*, y^*, z^*) is feasible (and not additionally optimal) in the proof given there. \square

Note that this MIP is feasible and y_{n+1} corresponds to the μ_0 variable of the LCP(p, Q) (with $\mu_0 = \frac{1}{\alpha^*} y_{n+1}^*$). Therefore by Corollary 2.12, the following result holds.

Corollary 2.15. (a) If MIP_1 has a feasible solution (α, y, z) with $\alpha > 0$ then $M \notin SC$.

(b) If MIP_1 has a feasible solution (α, y, z) with $\alpha > 0$ and $y_{n+1} > 0$ then $M \notin C$.

(c) MIP_1 has a global optimal value equal to zero if and only if $M \in SC$.

(d) If MIP_1 has no feasible solution (α, y, z) with $\alpha > 0$ and $y_{n+1} = 0$ then $M \in C$.

The case of $\alpha > 0$ and $y_{n+1} = 0$ in the computed solution of MIP_1 is the most difficult, as we cannot conclude that M is C or not from the solution of MIP_1 . In this case, we may consider a MIP of the form:

$$\begin{aligned} MIP_2 : \quad & \max \quad y_{n+1} \\ & \text{s.t.} \quad 0 \leq Qy + \alpha p \leq z \\ & \quad 0 \leq y \leq e - z \\ & \quad z \in \{0, 1\}^{n+1} \\ & \quad 0 \leq \alpha \leq 1 \\ & \quad \alpha \geq \varepsilon, \end{aligned}$$

where ε is a positive tolerance (usually $\varepsilon = 10^{-4}$). Then by Corollary 2.15, the following result holds:

Corollary 2.16. Let $\varepsilon > 0$. If MIP_2 has a feasible solution (α, y, z) with $y_{n+1} > 0$ then $M \notin C$.

As one can see from Corollary 2.15 it is easier to show for a matrix $M \notin SC$ than $M \notin C$. There is a further possibility for establishing that $M \notin C$ by making use of the test on a related matrix that is not SC:

Lemma 2.17. Let $M, H \in \mathcal{S}^n$ and $H \in SC$. Then $M \in C$ if and only if $M + \beta H \in SC$ for all $\beta > 0$.

Proof. The implication \Rightarrow is obvious from the definitions of matrices SC and C . Now, if $M \notin C$, there exists a $0 \neq \bar{x} \geq 0$ such that $\bar{x}^\top M \bar{x} < 0$. Since $H \in SC$ then

$$\beta = -\frac{\bar{x}^\top M \bar{x}}{\bar{x}^\top H \bar{x}}$$

is positive and satisfies

$$\bar{x}^\top (M + \beta H) \bar{x} = 0.$$

Hence $M + \beta H \notin SC$. \square

2.3 Conditions for copositivity based on LPs

Recall that for $M \in \mathcal{S}^n$ it holds

$$-M \in S_0 \Leftrightarrow \exists x \geq 0, x \neq 0 : Mx \leq 0$$

and thus $-M \in S_0$ implies $M \notin SC$. Thereby, $-M \in S_0$ if and only if the system

$$Mx \leq 0, x \geq 0, e^\top x = 1$$

has a solution, which can be found by solving a linear program (LP):

Lemma 2.18. *Let $c \in \mathbb{R}_+^n$. If*

$$LP_1 : \inf\{c^\top x \mid Mx \leq 0, x \geq 0, e^\top x = 1\}$$

has a minimal solution, then $M \notin SC$.

Note that, as LP_1 is bounded from below, it has a minimal solution if and only if it is feasible. The dual problem to LP_1 is $\sup_{y \in \mathbb{R}_+^n} \min_{i=1, \dots, n} \{c_i + (My)_i\}$ which is unbounded if and only if there exists $y \in \mathbb{R}_+^n$ such that for all $i \in \{1, \dots, n\}$ it holds $(My)_i > 0$.

Moreover, $M \notin S_0$, i.e.

$$0 \neq x \geq 0 \Rightarrow (Mx)_i < 0 \text{ for some } i$$

implies $M \notin C$. Note that $M \notin S_0$ if and only if the system

$$Mx \geq 0, x \geq 0, e^\top x = 1$$

has no solution, which can be verified by considering again a linear optimization problem:

Lemma 2.19. *Let $c \in \mathbb{R}_+^n$. If*

$$LP_2 : \inf\{c^\top x \mid Mx \geq 0, x \geq 0, e^\top x = 1\},$$

has no feasible solution, then $M \notin C$ and thus also $M \notin SC$.

Lemmas 2.18 and 2.19 give easy to verify necessary conditions for a matrix not to be (strict) copositive.

Example 2.20. *For the matrix M of Example 2.7 the system*

$$Mx \leq 0, \quad x \geq 0, \quad e^\top x = 1$$

has the solution $(1/2, 1/2)$. Thus, by Lemma 2.18, $M \notin SC$.

The system

$$\begin{array}{l} Mx \geq 0 \\ x \geq 0 \\ e^\top x = 1 \end{array} \Leftrightarrow \begin{cases} x_1 \geq 2x_2 \\ x_2 \geq 2x_1 \\ x_1 + x_2 = 1 \\ x_i \geq 0, i = 1, 2 \end{cases}$$

has no feasible solution. By Lemma 2.19 $M \notin C$.

3 Algorithm

Below we collect the results of the previous sections and give an algorithm for testing a matrix on (strict) copositivity. In the algorithm we also combine some preprocessing steps based on the conditions given in Section 2.3 with well known results from the literature.

3.1 Preprocessing

In addition to the results of Lemma 2.18 and 2.19 we will make use of the following preprocessing steps which are based on the collection in [3], see also [30].

Lemma 3.1. *Let $M = [m_{ij}] \in \mathcal{S}^n$ and choose an arbitrary $i \in \{1, \dots, n\}$.*

- (a) *If $m_{ii} < 0$, then $M \notin C$.*
- (b) *if $m_{ii} = 0$, then $M \notin SC$.*
- (c) *if $m_{ii} = 0 > m_{ij}$ for some $j \in \{1, \dots, n\}$, then $M \notin C$.*

The preprocessing steps used later in our main algorithm are summarized in Algorithm 1. We used $c = e$ in the LPs (e is a vector of ones).

Algorithm 1 Preprocessing for C and for SC

Input: matrix $M \in \mathcal{S}^n$

(Part (1):)

if $m_{ii} < 0$ for any $i \in \{1, \dots, n\}$ **then**
 $M \notin C$ and stop.

end if

if $m_{ii} = 0$ for any $i \in \{1, \dots, n\}$ **then**
 $M \notin SC$.

end if

if $m_{ii} = 0 > m_{ij}$ for any $i \neq j, i, j \in \{1, \dots, n\}$ **then**
 $M \notin C$ and stop.

end if

Let $c \in \mathbb{R}_+^n$.

(Part (2):)

if LP₁ has a feasible solution **then**
 $M \notin SC$.

end if

(Part (3):)

if LP₂ has no feasible solution **then**
 $M \notin C$ and stop.

end if

3.2 Outline of the algorithm for (strict) copositivity

Algorithm 2 gives the structure of our main algorithm. In Step 1 we make use of Corollary 2.9. For Step 2 we propose three different procedures in this section.

Algorithm 2 Test for C and SC

Input: Matrix $M \in \mathcal{S}^n$

STEP 0: Apply Algorithm 1 (Preprocessing).

if STEP 0 was not conclusive **then**

STEP 1: Use Lemke's method with different initial complementary basic solutions to solve $\text{LCP}(-e, M)$.

if Lemke's method terminates in an unbounded ray **then**

$M \notin SC$.

end if

STEP 2: Apply Procedures 1-3 to be discussed in Section 3.3.

end if

Note that it depends on the choice of the procedure in Step 2 whether the Algorithm 2 guarantees to find $M \notin C$, $M \in C$ or $M \in SC$.

3.3 Procedures for Step 2

As discussed in Section 2.2, we can solve $\text{LCP}(p, Q)$ to derive proofs for the matrix to be in C, in SC or not copositive. We may apply an enumerative algorithm, Lemke's method and the mixed linear integer formulation for solving this LCP. Each of the following three procedures are discussed next and can be used for Step 2 of Algorithm 2.

Procedure 1: Solution of $\text{LCP}(p, Q)$ by an enumerative algorithm and apply Corollary 2.12.

An efficient enumerative method for the Linear Complementarity Problem (LCP) has been proposed by Júdice, Faustino and Ribeiro [22]. This method finds a solution of the LCP by exploring a binary tree generated by the dichotomy $x_i = 0$ or $w_i = 0$ associated with the complementary condition, see Fig. 1.




Figure 1: Branching procedure of the enumerative method.

In each node of the tree, the algorithm finds a stationary point of the nonconvex quadratic program of the form

$$\begin{aligned}
\min \quad & f(x, w, \mu_0, r_0) = x^\top w + \mu_0 r_0 \\
\text{s.t.} \quad & w = Mx + \mu_0 e \\
& r_0 = e^\top x - 1 \\
& x_i = 0, \quad i \in I \\
& w_j = 0, \quad j \in J \\
& x, w, \mu_0, r_0 \geq 0,
\end{aligned} \tag{7}$$

where I and J are the index sets defined by the fixed variables, i.e. $I := \{i \in \{1, \dots, n\} : x_i = 0 \text{ fixed}\}$ (and also additionally $\mu_0 = 0$ may be fixed) and $J := \{i \in \{1, \dots, n\} : w_i = 0 \text{ fixed}\}$ (and eventually additionally $r_0 = 0$ fixed), respectively, in the path of the tree from this node to the root. Furthermore the algorithm contains some heuristic rules for choosing the node and the pair of complementary variables for branching. For details on the algorithm we refer to [22].

It is important to note that it is much easier to compute a solution for an LCP (when it exists) than showing that an LCP has no solution. So, as expected it is much easier to establish that a matrix is not copositive than proving that it has this property. However, the algorithm can at least in theory show that any matrix is C or not.

Procedure 2: Solution of LCP(p, Q) with Lemke's algorithm and $n + 1$ initial complementary basic solutions and apply Corollary 2.12.

Lemke's method is a pivotal algorithm that uses basic feasible solutions of the following general LCP (GLCP)

$$\begin{aligned}
v &= p + x_0 d + Qu \\
u &\geq 0, \quad v \geq 0, \quad x_0 \geq 0 \\
u^\top v &= 0
\end{aligned} \tag{8}$$

where in our setting $v = [w \ r_0]^\top$, $u = [x \ \mu_0]^\top$ and $d \in \mathbb{R}^{n+1}$ is a positive vector (usually $d = e$). To guarantee such a solution of the GLCP (8) the algorithm starts with a basic solution given by

$$\bar{u} = 0, \quad \bar{x}_0 = \frac{1}{d_r}, \quad v = p + \bar{x}_0 d, \tag{9}$$

where $d_r := \min\{d_i : i = 1, \dots, n + 1\} > 0$. All the variables u_i and the variable v_r are nonbasic and the remaining variables are basic. Hence there exists exactly a complementary pair (v_r, u_r) of nonbasic variables. In the next iteration the algorithm chooses u_r (complementary of the previous leaving variable) as the entering variable which interchanges with a leaving basic variable (found by the common minimum quotient rule) [11]. If such a leaving variable does not exist the algorithm stops in an unbounded ray. Otherwise, a new basic solution of the GLCP (8) is obtained and either $x_0 = 0$ and a solution of the LCP is at hand or the procedure is repeated.

We rewrite the linear constraints of GLCP (8) in the form

$$Ay = p, y \geq 0 \quad (10)$$

where $A = [I \ -Q \ -d] \in \mathbb{R}^{(n+1) \times (2n+3)}$, $y = [v \ u \ x_0]^\top \in \mathbb{R}^{2n+3}$ and $p \in \mathbb{R}^{n+1}$. A basic feasible solution for the system (10) is defined by two sets $J = \{k_1, \dots, k_{n+1}\}$ and $L = \{1, \dots, 2n+3\} \setminus J$ of basic and nonbasic variables respectively, such that

$$y_{k_i} = \bar{p}_i - \sum_{j \in L} \bar{a}_{ij} y_j, \quad i = 1, \dots, n+1. \quad (11)$$

This solution is given by $y_{k_i} = \bar{p}_i$, $i = 1, \dots, n+1$, $y_j = 0$, $j \in L$ and is called a basic feasible solution of the GLCP (8). A nonsingular Basis matrix $B \in \mathbb{R}^{(n+1) \times (n+1)}$ is associated to this basic solution and consists of the columns of A corresponding to the basic variables [25]. The steps of Lemke's method for solving LCP(p, Q) can now be presented as follows.

Lemke's method

Initial Step: Let $d_r := \min\{d_i, i = 1, \dots, n+1\}$ and start with a basic feasible solution of GLCP (8), where the u_i variables and v_r are nonbasic and the remaining variables are basic.

General Step: Let $y_s, s \in L$ be the nonbasic variable that is complementary of the variable that has become nonbasic in the previous iteration.

- (i) If $\bar{a}_{is} \leq 0$ for all $i = 1, \dots, n+1$, stop the algorithm with the termination in an unbounded ray.
- (ii) Compute $t \in \{1, \dots, n+1\}$ such that

$$t = \min \left\{ r \in \{1, \dots, n+1\} : \frac{\bar{p}_r}{\bar{a}_{rs}} = \min \left\{ \frac{\bar{p}_i}{\bar{a}_{is}} : \bar{a}_{is} > 0 \right\} \right\}. \quad (12)$$

Perform a pivotal operation which interchanges the nonbasic variable y_s with the basic variable y_{k_t} associated with line t in (12). If $x_0 = 0$ after such an operation, stop the algorithm with a solution of the LCP. Otherwise, repeat the general step.

An important feature of this algorithm is the reduced computational effort per iteration. In fact, this work essentially consists of solving a linear system of equations with the Basis matrix.

Lemke's method can start with any basic feasible solution of GLCP (8). A possible choice is to consider the vector x equal to one of the canonical basis vectors e^i instead of the null vector in the initial basic solution of GLCP (8). Therefore the basic variables of this initial basic solution are the variable $y_i = x_i$, the variable $y_{n+1} = \mu_0$ and the variables $y_j = w_j$, $j \in \{1, \dots, n\} \setminus \{i\}$. In this way we can construct $(n+1)$ initial basic solutions (including the trivial with $x = 0$) for initializing Lemke's method. Note that for each one of these basic solutions $d = Be$, where B is the associated Basis matrix and $e \in \mathbb{R}^{n+1}$ is a vector of ones.

An obvious drawback of this approach is that the process is not conclusive when it does not guarantee that a matrix is not C (or at least not SC).

Procedure 3: Solution of MIP_1 and MIP_2 and apply Corollaries 2.15 and 2.16.

In procedure 3 we aim again at solving the problem $LCP(p, Q)$ but this time by using its reformulation as an MIP. We start by finding a feasible solution with $\alpha > 0$ of MIP_1 and apply Corollary 2.15. If MIP_1 has a feasible solution with $\alpha > 0$ and $y_{n+1} = 0$, then we find a feasible solution of MIP_2 and apply Corollary 2.16. If the global optimal value of MIP_1 is equal to zero, then $M \in C$ (and even $M \in SC$). This discussion confirms that it is much easier to show that a matrix is not C (or not SC) than establishing that it is copositive.

4 Numerical results

We test the basic steps (Step 0 and 1) of our algorithm and all 3 procedures on several test instances from the literature — like the famous Horn matrix. We also consider generated Max-clique instances and instances from the DIMACS collection [14] with matrices up to size 496×496 . We compare the results with those from the literature on copositivity tests by Bundfuss and Dür [8], Žilinskas and Dür [32], and Bomze and Eichfelder [3]. Moreover, as testing copositivity is equivalent to determining a global solution of the quadratic optimization problem (1), see Lemma 2.3, we apply the global optimization solver BARON (with default parameters settings) and compare with our results.

All experiments have been performed on a Pentium IV (Intel) with 3.0 GHz and 2 GBytes of RAM memory, using the operating system Linux. The algorithm was implemented in the General Algebraic Modeling System (GAMS) language (Rev 118 Linux/Intel) [6] and the solvers CPLEX [12] (Version 9.1), MINOS [24] (Version 5.51) and BARON [27] (version 22.7.2) were used to solve the MIP and nonlinear optimization problems, and LP was solved with CPLEX. Running times presented in these sections are always given in CPU seconds. The maximum CPU time allowed for all procedures is 7200 seconds.

4.1 Test matrices

The non-copositive matrices $M_1 \notin C$ and $M_2 \notin C$ are taken from Bomze and Eichfelder [3] and Kaplan [23], respectively:

$$M_1 = \begin{pmatrix} 1 & -0.72 & -0.59 & 1 \\ -0.72 & 1 & -0.6 & -0.46 \\ -0.59 & -0.6 & 1 & -0.6 \\ 1 & -0.46 & -0.6 & 1 \end{pmatrix}, \quad M_2 = \begin{pmatrix} 1 & -0.72 & -0.59 & -0.6 \\ -0.72 & 1 & 0.21 & -0.46 \\ -0.59 & 0.21 & 1 & -0.6 \\ -0.6 & -0.46 & -0.6 & 1 \end{pmatrix}.$$

The matrix $M_3 \in SC$ is from Kaplan [23] and $M_4 \in SC$ is a principal submatrix of a matrix from Kaplan [23, Ex. 1]:

$$M_3 = \begin{pmatrix} 1 & 0.9 & -0.54 & 0.21 \\ 0.9 & 1 & -0.03 & 0.78 \\ -0.54 & -0.03 & 1 & 0.52 \\ 0.21 & 0.78 & 0.52 & 1 \end{pmatrix}, \quad M_4 = \begin{pmatrix} 1 & 0.9 & -0.54 \\ 0.9 & 1 & -0.03 \\ -0.54 & -0.03 & 1 \end{pmatrix}.$$

The matrix $M_5 \in C \setminus SC$ is from Väliäho [31] and the famous Horn matrix M_6 is also an example with $M_6 \in C \setminus SC$ [18]:

$$M_5 = \begin{pmatrix} 1 & -1 & 1 & 2 & -3 \\ -1 & 2 & -3 & -3 & 4 \\ 1 & -3 & 5 & 6 & -4 \\ 2 & -3 & 6 & 5 & -8 \\ -3 & 4 & -4 & -8 & 16 \end{pmatrix}, \quad M_6 = \begin{pmatrix} 1 & -1 & 1 & 1 & -1 \\ -1 & 1 & -1 & 1 & 1 \\ 1 & -1 & 1 & -1 & 1 \\ 1 & 1 & -1 & 1 & -1 \\ -1 & 1 & 1 & -1 & 1 \end{pmatrix}$$

The Hoffman-Pereira matrix $M_7 \in C \setminus SC$ [20] is another example of this type:

$$M_7 = \begin{pmatrix} 1 & -1 & 1 & 0 & 0 & 1 & -1 \\ -1 & 1 & -1 & 1 & 0 & 0 & 1 \\ 1 & -1 & 1 & -1 & 1 & 0 & 0 \\ 0 & 1 & -1 & 1 & -1 & 1 & 0 \\ 0 & 0 & 1 & -1 & 1 & -1 & 1 \\ 1 & 0 & 0 & 1 & -1 & 1 & -1 \\ -1 & 1 & 0 & 0 & 1 & -1 & 1 \end{pmatrix}.$$

Thus $M_1, M_2 \notin C$, $M_3, M_4 \in SC$ and $M_5, M_6, M_7 \in C \setminus SC$.

4.2 Maximum clique instances

Next to the above test matrices we also used test instances for the maximum clique problem (based on a reformulation as a linear optimization problem over the cone of copositive matrices) from the DIMACS collection [14], cf. [3, 8, 32] and generated instances (cf. [32]). For a simple, i.e. loopless and undirected, graph $G = (V, E)$ with node set $V = \{1, \dots, n\}$ and edge set E , a clique C is a subset of V such that every pair of nodes in C is connected by an edge in E . A clique C is said to be a maximum clique if it contains the most elements among all cliques, and its size $\omega(G)$ is called the (maximum) clique number. The maximum clique problem can be reformulated as a copositive optimization problem

$$\omega(G) = \min \{ \lambda \in \mathbb{N} \mid \lambda(E_n - A_G) - E_n \text{ is copositive} \} \quad (13)$$

with E_n the $n \times n$ all-ones matrix and $A_G = [a_{ij}]_{i,j}$ the adjacency matrix of the graph G , i.e. $a_{ij} = 1$ if $\{i, j\} \in E$, and $a_{ij} = 0$ else, $i, j \in \{1, \dots, n\}$. According to [28, Prop. 3.2] it holds

$$\lambda(E_n - A_G) - E_n \begin{cases} \in SC & \text{if } \lambda > \omega(G) \\ \in C \setminus SC & \text{if } \lambda = \omega(G) \\ \notin C & \text{if } \lambda < \omega(G). \end{cases}$$

Thus if $\lambda(E_n - A_G) - E_n \notin C$ we can conclude that $\omega(G) \geq \lambda + 1$. In Table 1 we include the characteristics of the graphs from DIMACS [14] collection and the generated graphs (cf. [32]). The number n of nodes gives the dimension $n \times n$ of the examined matrices. The small instances (with $n \in [14, 22]$) as well as eight large instances of Table 1 were also tested in [3].

Table 1: Generated small instances [32] and large instances from DIMACS collection [14].

MATRIX	N	E	$\omega(G)$
c-fat14-1	14	52	6
Brock14	14	51	5
Brock16	16	59	5
Brock18	18	78	5
Brock20	20	95	5
Morgen14	14	50	5
Morgen16	16	59	5
Morgen18	18	60	5
Morgen20	20	67	5
Morgen22	22	68	5
Johnson6-2-4	15	45	3
Johnson6-4-4	15	45	3
Johnson7-2-4	21	105	3
Jagota14	14	31	6
Jagota16	16	57	8
Jagota18	18	84	10
sanchis14	14	50	5
sanchis16	16	50	5
sanchis18	18	50	5
sanchis20	20	50	5
sanchis22	22	50	5

MATRIX	N	E	$\omega(G)$
Brock200-1	200	14834	21
Brock200-2	200	9876	12
Brock200-3	200	12048	15
Brock200-4	200	13089	17
c-fat200-1	200	1534	12
c-fat200-2	200	3235	24
c-fat200-5	200	8473	58
Hamming6-2	64	1824	32
Hamming6-4	64	704	4
Hamming8-2	256	31616	128
Hamming8-4	256	20864	16
Johnson8-2-4	28	210	4
Johnson8-4-4	70	1855	14
Johnson16-2-4	120	5460	8
Johnson32-2-4	496	107880	16
Keller4	171	9435	11
Mann-a9	45	918	16
Mann-a27	378	70551	126

4.3 Numerical Experiments: Steps 0 and 1

In Tables 2, 3 and 4 we report the performance of Steps 0 and 1 of the algorithm where IT and T denote respectively, the number of LP iterations and time of execution required by Lemke's algorithm.

We have run all the steps of the algorithm even in the cases where one of the previous steps is conclusive. The numbers (1), (2) and (3) in Step 0 refer to the Parts as marked in Algorithm 2.

Furthermore in Step 1, for instances where Lemke's method was inconclusive for the initial trivial basis ($x_i = 0$, for all $i = 1, \dots, n$), i.e., Lemke's method found a solution, then we applied the method with different initial basic solutions as described in Procedure 2 for Step 2. The process is repeated until the method terminates in an unbounded ray (Case: UNB. RAY) or finds a solution of the LCP in each one of the $(n + 1)$ applications of the method (Case: SOL.).

The numerical results indicate that the preprocessing phase does not help much for these instances. Actually there are only two cases where the preprocessing was effective. On the contrary, the use of the Lemke's method in Step 1 seems quite promising for establishing that a matrix is not *SC*. Furthermore in many cases it was enough to apply Lemke's algorithm only once with the trivial basic feasible

Table 2: Performance of the algorithm for the 7 matrices of Section 4.1 (Steps 0 and 1).

MATRIX	STEP 0			STEP 1					
	(1)	(2)	(3)	TRIVIAL BASIS		OTHER BASIS		UNB. RAY	SOL.
	$M \notin C$	$M \notin SC$	$M \notin C$	IT	T	IT	T	$M \notin SC$	
M_1	-	✓	✓	3	0.00E+00			✓	
M_2	-	✓	✓	4	0.00E+00			✓	
M_3	-	-	-	2	0.00E+00	8	0.00E+00		✓
M_4	-	-	-	2	0.00E+00	5	3.13E-02		✓
M_5	-	-	-	3	0.00E+00			✓	
M_6	-	-	-	2	0.00E+00			✓	
M_7	-	-	-	2	0.00E+00			✓	

Table 3: Performance of the algorithm for small matrices $\lambda(E_n - A_G) - E_n$, with $\lambda = \omega(G) - 1$ (Steps 0 and 1).

MATRIX	STEP 0			STEP 1					
	(1)	(2)	(3)	TRIVIAL BASIS		OTHER BASIS		UNB. RAY	SOL.
	$M \notin C$	$M \notin SC$	$M \notin C$	IT	T	IT	T	$M \notin SC$	
c-fat14-1	-	-	-	6	3.13E-02			✓	
Brock14	-	-	-	4	0.00E+00			✓	
Brock16	-	-	-	3	0.00E+00	9	0.00E+00	✓	
Brock18	-	-	-	4	0.00E+00			✓	
Brock20	-	-	-	4	0.00E+00			✓	
Morgen14	-	-	-	5	0.00E+00			✓	
Morgen16	-	-	-	4	0.00E+00			✓	
Morgen18	-	-	-	2	0.00E+00	15	3.13E-02	✓	
Morgen20	-	-	-	4	0.00E+00			✓	
Morgen22	-	-	-	4	0.00E+00			✓	
Johnson6-2-4	-	-	-	3	0.00E+00			✓	
Johnson6-4-4	-	-	-	3	0.00E+00			✓	
Johnson7-2-4	-	-	-	3	0.00E+00			✓	
Jagota14	-	-	-	4	0.00E+00	11	0.00E+00	✓	
Jagota16	-	-	-	5	0.00E+00	15	0.00E+00	✓	
Jagota18	-	-	-	6	0.00E+00	19	3.13E-02	✓	
sanchis14	-	-	-	5	0.00E+00			✓	
sanchis16	-	-	-	2	0.00E+00	26	0.00E+00	✓	
sanchis18	-	-	-	5	0.00E+00			✓	
sanchis20	-	-	-	5	0.00E+00			✓	
sanchis22	-	-	-	3	0.00E+00	4	0.00E+00	✓	

solution of GLCP (8). The computational effort of Lemke's algorithm in this last case is quite small. So, as a final conclusion of this first study, Steps 0 and 1 should be included in a more elaborated algorithm to verify whether a given matrix is SC or C or not.

4.4 Numerical Experiments: Step 2

4.5 Procedure 1

In Tables 5, 6 and 7 we report the performance of the enumerative algorithm to solve $LCP(p, Q)$ in order to detect copositivity or non-copositivity of the matrices. If the algorithm finds a solution then it gives an indication of just $M \notin SC$ or $M \notin C$ depending on the value of μ_0 to be equal to zero or positive respectively. So, the algorithm stops as soon as it finds a complementary solution with $\mu_0 > 0$. If this is not the case, the search process continues until showing that the $LCP(p, Q)$ has

Table 4: Performance of the algorithm for matrices $\lambda(E_n - A_G) - E_n$, with $\lambda = \omega(G) - 1$ (Steps 0 and 1).

MATRIX	STEP 0			STEP 1					
	(1)	(2)	(3)	TRIVIAL BASIS		OTHER BASIS		UNB. RAY	SOL.
	$M \notin C$	$M \notin SC$	$M \notin C$	IT	T	IT	T	$M \notin SC$	
Brock200-1	-	-	-	131	9.38E-02	2589	5.72E+00		✓
Brock200-2	-	-	-	7	2.50E-02	1246	3.05E+00		✓
Brock200-3	-	-	-	11	1.09E-02	1881	4.23E+00		✓
Brock200-4	-	-	-	12	7.81E-02	2062	5.14E+00		✓
c-fat200-1	-	-	-	12	3.13E-02			✓	
c-fat200-2	-	-	-	24	6.25E-02			✓	
c-fat200-5	-	-	-	58	1.25E-01			✓	
Hamming6-2	-	-	-	32	3.13E-02			✓	
Hamming6-4	-	-	-	4	0.00E+00			✓	
Hamming8-2	-	-	-	128	5.78E-01			✓	
Hamming8-4	-	-	-	16	1.25E-01			✓	
Johnson8-2-4	-	-	-	3	0.00E+00			✓	
Johnson8-4-4	-	-	-	14	0.00E+00			✓	
Johnson16-2-4	-	-	-	8	3.13E-02			✓	
Johnson32-2-4	-	-	-	16	4.53E-01			✓	
Keller4	-	-	-	7	3.13E-02	1131	2.39E+00		✓
Mann-a9	-	-	-	9	0.00E+00	468	9.38E-02		✓
Mann-a27	-	-	-	27	3.59E-01	14040	1.27E+02		✓

no solution ($M \in SC$ and $M \in C$) or the maximum CPU time of 7200 seconds is attained. In the following tables the notation NODES, IT and T stands respectively, for the total number of nodes, iterations and time used by the enumerative algorithm. We marked with (-) and (*) respectively, the instances for which a complementary solution does not exist ($M \in SC$) and the algorithm was not able to find such a solution within the CPU time allowed.

Table 5: (Procedure 1) Performance of the enumerative algorithm to solve LCP(p, Q) for matrices of Sections 4.1.

MATRIX	NODES	IT	T	μ_0	$M \in SC$	$M \notin SC$	$M \notin C$
M_1	1	4	5.80E-02	9.19E-02			✓
M_2	1	5	5.80E-02	1.16E-01			✓
M_3	14	19	1.74E+00	-	✓		
M_4	12	16	1.42E+00	-	✓		
M_5	1	8	5.90E-02	0.00E+00		✓	
M_6	1	2	5.90E-02	0.00E+00		✓	
M_7	1	2	5.80E-02	0.00E+00		✓	

The numerical results indicate that the enumerative method was efficient to establish that the matrix is at least not SC in general. However, there were three instances where the method was unable to terminate under the maximum time allowed. Furthermore, the algorithm required a small amount of effort for the smallest instances (Tables 5 and 6) but this effort increases very much when the dimension increases.

4.6 Procedure 2

Tables 8, 9 and 10 include the performance of Lemke's algorithm for solving LCP(p, Q) by changing the initial basic solution for each application k of the method ($k = 1, \dots, n + 1$). The process is repeated until the method finds a solution with $\mu_0 > 0$

Table 6: (Procedure 1) Performance of the enumerative algorithm to solve $LCP(p, Q)$ for small matrices $\lambda(E_n - A_G) - E_n$, with $\lambda = \omega(G) - 1$.

MATRIX	NODES	IT	T	μ_0	$M \notin SC$	$M \notin C$
c-fat14-1	3	15	4.71E-01	1.67E-01		✓
Brock14	1	18	5.90E-02	0.00E+00	✓	
Brock16	10	52	1.02E+00	0.00E+00	✓	
Brock18	1	8	6.00E-02	0.00E+00	✓	
Brock20	1	8	6.10E-02	0.00E+00	✓	
Morgen14	1	12	6.00E-02	2.00E-01		✓
Morgen16	1	13	6.00E-02	0.00E+00	✓	
Morgen18	32	98	7.96E+00	0.00E+00	✓	
Morgen20	1	16	6.10E-02	2.00E-01		✓
Morgen22	1	15	6.10E-02	0.00E+00	✓	
Johnson6-2-4	1	19	6.00E-02	0.00E+00	✓	
Johnson6-4-4	1	19	5.90E-02	0.00E+00	✓	
Johnson7-2-4	1	19	6.10E-02	2.86E-02		✓
Jagota14	3	33	1.39E-01	1.67E-01		✓
Jagota16	13	103	3.05E+00	1.25E-01		✓
Jagota18	23	236	8.36E+00	1.00E-01		✓
sanchis14	12	69	1.40E+00	0.00E+00	✓	
sanchis16	10	28	1.02E+00	0.00E+00	✓	
sanchis18	1	8	6.00E-02	2.00E-01		✓
sanchis20	1	9	6.10E-02	2.00E-01		✓
sanchis22	80	226	7.35E+01	2.00E-01		✓

Table 7: (Procedure 1) Performance of the enumerative algorithm to solve $LCP(p, Q)$ for matrices $\lambda(E_n - A_G) - E_n$, with $\lambda = \omega(G) - 1$.

MATRIX	NODES	IT	T	μ_0	$M \notin SC$	$M \notin C$
Brock200-1	730	5.35E+04	7.20E+03	0.00E+00	✓	
Brock200-2	760	5.43E+04	7.20E+03	*		
Brock200-3	734	6.19E+04	7.20E+03	*		
Brock200-4	718	4.14E+04	7.20E+03	*		
c-fat200-1	1	1.60E+01	2.81E-01	8.33E-02		✓
c-fat200-2	1	2.80E+01	2.83E-01	4.17E-02		✓
c-fat200-5	1	6.10E+01	2.88E-01	1.72E-02		✓
Hamming6-2	1	4.53E+02	9.20E-02	3.13E-02		✓
Hamming6-4	1	4.20E+01	8.00E-02	0.00E+00	✓	
Hamming8-2	1	7.11E+03	2.25E+00	7.81E-03		✓
Hamming8-4	7	3.99E+03	2.61E+00	6.25E-02		✓
Johnson8-2-4	1	1.50E+01	6.20E-02	0.00E+00	✓	
Johnson8-4-4	1	5.71E+02	1.02E-01	7.14E-02		✓
Johnson16-2-4	1	1.43E+02	1.40E-01	0.00E+00	✓	
Johnson32-2-4	1	1.26E+03	2.71E+00	0.00E+00	✓	
Keller4	641	7.85E+04	2.93E+03	9.09E-02		✓
Mann-a9	4	6.50E+01	1.84E-01	0.00E+00	✓	
Mann-a27	840	3.80E+04	7.20E+03	0.00E+00	✓	

($M \notin C$) or only computes one solution with $\mu_0 = 0$ ($M \notin SC$) or terminates in an unbounded ray for all the $n + 1$ initial basic solutions (no conclusion is given). The notation IT stands for the total number of pivotal iterations required by Lemke's algorithm for the visited basis.

The numerical results indicate that Lemke's method was able to declare the matrices not SC or not C for all the instances but four. Actually three of these instances were the ones for which the enumerative method was not conclusive. The computational effort also increases much with the increasing of the dimension of the problems.

Table 8: (Procedure 2) Performance of the Lemke's algorithm with different initial basis for the matrices of Section 4.1.

MATRIX	TRIVIAL BASIS		OTHER BASIS		$M \notin SC$	$M \notin C$
	IT	T	IT	T		
M_1	1	0.00E+00	3	0.00E+00		✓
M_2	1	0.00E+00	4	0.00E+00		✓
M_3	1	0.00E+00	7	4.69E-02		
M_4	1	0.00E+00	6	0.00E+00		
M_5	1	0.00E+00	19	0.00E+00	✓	
M_6	1	0.00E+00	15	0.00E+00	✓	
M_7	1	0.00E+00	22	3.13E-02	✓	

Table 9: (Procedure 2) Performance of the Lemke's algorithm for small matrices $\lambda(E_n - A_G) - E_n$, with $\lambda = \omega(G) - 1$.

MATRIX	TRIVIAL BASIS		OTHER BASIS		$M \notin SC$	$M \notin C$
	IT	T	IT	T		
c-fat14-1	1	0.00E+00	6	0.00E+00		✓
Brock14	1	0.00E+00	15	0.00E+00		✓
Brock16	1	0.00E+00	59	3.13E-02	✓	
Brock18	1	0.00E+00	37	0.00E+00		✓
Brock20	1	0.00E+00	24	0.00E+00		✓
Morgen14	1	0.00E+00	16	0.00E+00		✓
Morgen16	1	0.00E+00	19	0.00E+00		✓
Morgen18	1	0.00E+00	7	3.13E-02		✓
Morgen20	1	0.00E+00	13	0.00E+00		✓
Morgen22	1	0.00E+00	28	0.00E+00		✓
Johnson6-2-4	1	0.00E+00	3	0.00E+00		✓
Johnson6-4-4	1	0.00E+00	3	0.00E+00		✓
Johnson7-2-4	1	0.00E+00	89	3.13E-02		✓
Jagota14	1	0.00E+00	14	0.00E+00		✓
Jagota16	1	0.00E+00	18	0.00E+00		✓
Jagota18	1	0.00E+00	22	0.00E+00		✓
sanchis14	1	0.00E+00	5	3.13E-02		✓
sanchis16	1	0.00E+00	45	0.00E+00		✓
sanchis18	1	0.00E+00	5	0.00E+00		✓
sanchis20	1	0.00E+00	5	0.00E+00		✓
sanchis22	1	0.00E+00	22	0.00E+00		✓

4.7 Procedure 3

In the next tables 11-13 we report the performance of the solver CPLEX to find a feasible solution for the problems MIP_1 and MIP_2 . The symbol (*) stays for problems where the solver CPLEX was unable to find a solution with $\alpha > 0$ within 3600 seconds of CPU time allowed. In this case, CPLEX gives the feasible solution $\alpha = 0$ for MIP_1 . We used the notations NODES, IT and T respectively, for the number of nodes, iterations and CPU seconds required by the solver.

The numerical results indicate that using the MIP formulation of the $LCP(p, Q)$ seems to be an interesting approach for showing that a matrix is not SC or not C . Like the remaining procedures, the computational effort is reduced for the smallest problems but tends to increase with an increase of the dimension of the problems. The procedure could not give an indication of noncopositivity for four instances. However, Lemke's method has shown noncopositivity for all these four matrices.

Table 10: (Procedure 2) Performance of the Lemke's algorithm with different initial basis for matrices $\lambda(E_n - A_G) - E_n$, with $\lambda = \omega(G) - 1$.

MATRIX	TRIVIAL BASIS		OTHER BASIS		$M \notin SC$	$M \notin C$
	IT	T	IT	T		
Brock200-1	1	9.38E-02	2.86E+03	6.41E+00		
Brock200-2	1	9.38E-02	1.43E+03	3.27E+00		
Brock200-3	1	9.38E-02	1.89E+03	5.82E+00		
Brock200-4	1	9.38E-02	2.12E+03	6.77E+00		
c-fat200-1	1	3.13E-02	1.20E+01	6.25E-02		✓
c-fat200-2	1	6.25E-02	2.40E+01	9.38E-02		✓
c-fat200-5	1	6.25E-02	5.80E+01	1.25E-01		✓
Hamming6-2	1	3.13E-02	3.20E+01	3.13E-02		✓
Hamming6-4	1	0.00E+00	4.00E+00	0.00E+00		✓
Hamming8-2	1	2.97E-01	1.28E+02	7.03E-01		✓
Hamming8-4	1	2.19E-01	1.60E+01	1.88E-01		✓
Johnson8-2-4	1	3.13E-02	2.20E+01	0.00E+00		✓
Johnson8-4-4	1	0.00E+00	2.31E+02	1.25E-01		✓
Johnson16-2-4	1	4.69E-02	8.00E+00	3.13E-02		✓
Johnson32-2-4	1	6.88E-01	1.60E+01	1.00E+00		✓
Keller4	1	6.25E-02	1.33E+03	2.47E+00	✓	
Mann-a9	1	0.00E+00	1.80E+01	0.00E+00		✓
Mann-a27	1	4.06E-01	4.73E+04	4.19E+02	✓	

Table 11: (Procedure 3) Performance of applying CPLEX to MIP_1 and MIP_2 for the matrices of Section 4.1.

MATRIX	MIP ₁					MIP ₂				
	$(\alpha \neq 0, \alpha \neq 0, \alpha = 0 \ y_{n+1} = 0)$		IT	NODES	T	$(\alpha \neq 0, \alpha \neq 0, \alpha = 0 \ y_{n+1} > 0)$		IT	NODES	T
	$M \in SC$	$M \notin SC$				$M \in C$	$M \notin C$			
M_1		✓	10	0	0.00E+00					
M_2		✓	9	0	0.00E+00					
M_3	✓		20	5	0.00E+00					
M_4	✓		20	5	0.00E+00					
M_5		✓	22	3	0.00E+00	✓		68	26	0.00E+00
M_6		✓	21	5	0.00E+00	✓		32	10	0.00E+00
M_7		✓	56	12	0.00E+00	✓		59	18	0.00E+00

4.8 Summary of numerical results

As a conclusion of this numerical study, we suggest to use in Step 2 Procedure 2 (Lemke's method) first and then Procedure 3 (mixed integer formulation) when the Procedure 2 is not conclusive. It is important to add that such a hybrid method was able to establish that all the matrices but one (Mann-a27) of the maximum clique collection are not C .

The numerical experiments also show that it is easier in general to establish that a matrix M is not SC than showing that M is not C . For that reason we make use of Lemma 2.17 to establish that the last matrix Mann-a27 of the clique collection is not C . Let $M := (\omega(G) - 1)(E_n - A_G) - E_n$ be this matrix. The matrix $H := E_n - A_G$ is a nonnegative matrix with positive diagonal elements and $H \in SC$ [11, Chapter 3]. Applying the hybrid method to the matrix $P := M + 0.1H$ (i.e., $\beta = 0.1$) then the algorithm terminates in Step 1 with the indication that $P \notin SC$. The algorithm used more than one initial basic solution, and required 2.56E+04 iterations and 2.67E+02 CPU time. So, matrix Mann-a27 for $\lambda = \omega(G) - 1$ is not C .

As a final conclusion of this numerical study, the hybrid algorithm including Steps 0,

Table 12: (Procedure 3) Performance of applying CPLEX to MIP_1 and MIP_2 for small matrices $\lambda(E_n - A_G) - E_n$, with $\lambda = \omega(G) - 1$.

MATRIX	MIP ₁						MIP ₂					
	$(\alpha \neq 0, \quad (\alpha \neq 0, \quad \alpha = 0 \quad y_{n+1} = 0) \quad y_{n+1} > 0)$			IT	NODES	T	$y_{n+1} = 0 \quad y_{n+1} > 0$			IT	NODES	T
	$M \in SC$	$M \notin SC$	$M \notin C$				$M \in C$	$M \notin C$				
c-fat14-1	✓			546	68	1.00E-02		✓	561	9.00E+01	2.00E-02	
Brock14	✓			281	34	1.00E-02		✓	932	2.12E+02	2.00E-02	
Brock16			✓	205	25	1.00E-02						
Brock18	✓			14	0	0.00E+00		✓	1474	3.71E+02	3.00E-02	
Brock20	✓			374	54	2.00E-02		✓	2957	4.19E+02	7.00E-02	
Morgen14			✓	69	1	1.00E-02						
Morgen16	✓			40	4	1.00E-02		✓	1038	1.99E+02	3.00E-02	
Morgen18	✓			304	29	2.00E-02		✓	1406	2.40E+02	6.00E-02	
Morgen20	✓			478	66	2.00E-02		✓	1786	2.89E+02	6.00E-02	
Morgen22	✓			199	23	1.00E-02		✓	1414	2.78E+02	5.00E-02	
Johnson6-2-4	✓			179	19	1.00E-02		✓	239	4.90E+01	1.00E-02	
Johnson6-4-4	✓			122	8	0.00E+00		✓	160	2.80E+01	1.00E-02	
Johnson7-2-4	✓			340	38	2.00E-02		✓	1026	1.21E+02	2.00E-02	
Jagota14			✓	266	53	1.00E-02						
Jagota16			✓	719	137	2.00E-02						
Jagota18			✓	1934	297	4.00E-02						
sanchis14	✓			169	27	1.00E-02		✓	696	1.75E+02	2.00E-02	
sanchis16			✓	182	20	1.00E-02						
sanchis18			✓	37	1	0.00E+00						
sanchis20			✓	42	2	0.00E+00						
sanchis22			✓	62	1	1.00E-02						

1 and 2 as discussed above was able to establish noncopositivity for all the maximum clique matrices. This means that such a procedure was able to give a lower bound of $\omega(G)$ for all these problems. Table 14 demonstrates this behavior of the hybrid algorithm and shows the superiority of this algorithm over the approaches discussed by Bomze and Eichfelder [3], Bundfuss and Dür [8] and Žilinskas and Dür [32].

4.9 Global solution with Baron

Tables 15, 16 and 17 report the performance of the solver BARON (with default parameters settings) for finding the global minimum of problem (1). We marked with (*) the problems for which the solver was not able to prove the optimality of the solution within the 7200 CPU seconds allowed and we report the best upper bound obtained by the solver for the limited time of execution. For all the small instances $\lambda(E_n - A_G) - E_n$ this upper bound allows to conclude that $M \notin C$ but it is inconclusive for five of the bigger matrices. These results clearly indicate that is better to employ our new hybrid method to establish noncopositivity than an efficient global optimizer for studying this property by exploiting the definition of a copositive matrix.

5 Conclusions

In this paper we introduce a number of procedures based on the linear complementarity problem and linear programming that proved to be useful for studying the copositivity or noncopositivity of a matrix. A hybrid algorithm has been constructed

Table 13: (Procedure 3) Performance of applying CPLEX to MIP_1 and MIP_2 for matrices $\lambda(E_n - A_G) - E_n$, with $\lambda = \omega(G) - 1$.

MATRIX	MIP ₁					
	$\alpha = 0$	$(\alpha \neq 0, y_{n+1} = 0)$	$(\alpha \neq 0, y_{n+1} > 0)$	IT	NODES	T
	$M \in SC$	$M \notin SC$	$M \notin C$			
Brock200-1			✓	3.46E+06	8.59E+04	9.13E+02
Brock200-2			✓	2.47E+06	5.48E+04	5.29E+02
Brock200-3			✓	7.21E+06	1.79E+05	1.55E+03
Brock200-4			✓	8.83E+04	1.96E+03	4.86E+01
c-fat200-1		✓		1.31E+05	2.34E+03	5.84E+01
c-fat200-2		✓		5.31E+05	5.95E+03	1.54E+02
c-fat200-5	*			1.61E+07	3.57E+05	3.60E+03
Hamming6-2			✓	4.52E+07	3.20E+06	3.13E+03
Hamming6-4		✓		2.90E+05	1.62E+04	1.13E+01
Hamming8-2	*			5.88E+06	1.48E+05	3.60E+03
Hamming8-4	*			1.13E+07	1.33E+05	3.60E+03
Johnson8-2-4		✓		4.60E+01	0.00E+00	1.00E-02
Johnson8-4-4			✓	1.40E+05	5.07E+03	1.06E+01
Johnson16-2-4			✓	3.47E+07	5.68E+05	3.60E+03
Johnson32-2-4			✓	1.59E+06	7.75E+03	3.60E+03
Keller4		✓		2.42E+07	5.67E+05	3.60E+03
Mann-a9		✓		4.43E+06	3.78E+05	4.00E+02
Mann-a27	*			1.53E+06	6.31E+04	3.60E+03

MATRIX	MIP ₂				
	$y_{n+1} = 0$	$y_{n+1} > 0$	IT	NODES	T
	$M \in C$	$M \notin C$			
Brock200-1					
Brock200-2					
Brock200-3					
Brock200-4					
c-fat200-1		✓	9.92E+04	4.63E+03	3.72E+01
c-fat200-2		✓	2.67E+05	2.40E+04	8.11E+01
c-fat200-5					
Hamming6-2					
Hamming6-4		✓	4.23E+04	2.55E+03	1.94E+00
Hamming8-2					
Hamming8-4					
Johnson8-2-4		✓	2.22E+03	3.49E+02	7.00E-02
Johnson8-4-4					
Johnson16-2-4					
Johnson32-2-4					
Keller4		✓	2.22E+07	1.84E+06	3.60E+03
Mann-a9		✓	3.28E+07	1.80E+07	3.60E+03
Mann-a27					

based on these procedures and has shown to perform well to establish noncopositivity of matrices of the so-called maximum clique collection that are usually used as test instances for similar procedures.

Numerical results with these instances indicate that the hybrid algorithm is more efficient to detect that a matrix is not strictly copositive than showing that it is not copositive. This conclusion has been exploited to establish the noncopositivity of one of the matrices of the maximum clique set by showing that a related matrix is not strictly copositive. In our opinion, such type of approach should deserve more attention in the future. Recently, a similar strategy was suggested by Sponsel et al. in [28, Theorem 3.3].

It is also interesting to investigate the performance of the algorithm discussed in

Table 14: Comparison of lower bounds for DIMACS collection with the results in [3], [8] and [32].

MATRIX	$\omega(G)$	LOWER BOUNDS				
		HYBRID	ALGORITHM	in [3]	in [8]	in [32]
Brock200-1	21		21			13
Brock200-2	12		12		9	10
Brock200-3	15		15		11	11
Brock200-4	17		17	7		13
c-fat200-1	12		12			
c-fat200-2	24		24			
c-fat200-5	58		58			
Hamming6-2	32		32	32	28	32
Hamming6-4	4		4		4	4
Hamming8-2	128		128	128		128
Hamming8-4	16		16	16	12	16
Johnson8-2-4	4		4	4	4	4
Johnson8-4-4	14		14	14	14	14
Johnson16-2-4	8		8	8	8	8
Johnson32-2-4	16		16			16
Keller4	11		11	6	9	8
Mann-a9	16		16		16	16
Mann-a27	126		126			121

Table 15: Performance of the solver Baron for the matrices of Section 4.1.

MATRIX	UPPER BOUND	NODES	T	$M \in SC$	$M \in C$	$M \notin C$
M_1	-9.19E-02	129	8.40E-02			✓
M_2	-1.16E-01	365	2.00E-01			✓
M_3	2.30E-01	21	3.90E-02	✓		
M_4	2.30E-01	7	3.00E-02	✓		
M_5	-7.40E-17	1.42E+03	9.25E-01		✓	
M_6	0.00E+00	6.98E+04	4.68E+01		✓	
M_7	0.00E+00	1.73E+05	1.53E+02		✓	

Table 16: Performance of the solver Baron for small matrices $\lambda(E_n - A_G) - E_n$, with $\lambda = \omega(G) - 1$.

MATRIX	UPPER BOUND	NODES	T	$M \in C$	$M \notin C$
c-fat14-1	-1.67E-01	3.57E+05	5.76E+02		✓
Brock14	-2.00E-01*	4.60E+05	7.20E+03		✓
Brock16	-2.00E-01*	8.96E+05	1.61E+03		✓
Brock18	-2.00E-01*	4.60E+05	7.20E+03		✓
Brock20	-2.00E-01*	2.51E+06	7.20E+03		✓
Morgen14	-2.00E-01*	1.51E+05	7.20E+03		✓
Morgen16	-2.00E-01*	4.73E+05	7.20E+03		✓
Morgen18	-2.00E-01*	2.86E+06	7.20E+03		✓
Morgen20	-2.00E-01*	2.34E+06	7.20E+03		✓
Morgen22	-2.00E-01*	1.85E+06	5.83E+03		✓
Johnson6-2-4	-3.33E-01*	1.80E+06	4.30E+03		✓
Johnson6-4-4	-3.33E-01*	1.72E+06	4.09E+03		✓
Johnson7-2-4	-3.33E-01*	2.92E+04	7.20E+03		✓
Jagota14	-1.67E-01	3.06E+04	5.27E+01		✓
Jagota16	-1.25E-01	6.62E+05	1.42E+03		✓
Jagota18	-1.00E-01*	2.11E+06	7.20E+03		✓
sanchis14	-2.00E-01*	3.83E+05	5.15E+02		✓
sanchis16	-2.00E-01*	3.93E+05	7.13E+02		✓
sanchis18	-2.00E-01*	3.68E+05	6.37E+02		✓
sanchis20	-2.00E-01*	4.02E+06	7.20E+03		✓
sanchis22	-2.00E-01*	3.29E+06	7.20E+03		✓

Table 17: Performance of the solver Baron for matrices $\lambda(E_n - A_G) - E_n$, with $\lambda = \omega(G) - 1$.

MATRIX	UPPER BOUND	NODES	T	$M \in C$	$M \notin C$
Brock200-1	5.26E-02*	7.49E+02	7.20E+03		
Brock200-2	1.00E-01*	7.75E+02	7.20E+03		
Brock200-3	7.69E-02*	8.92E+02	7.20E+03		
Brock200-4	6.67E-02*	8.62E+02	7.20E+03		
c-fat200-1	-8.33E-02*	4.20E+03	7.20E+03		✓
c-fat200-2	-4.17E-02*	3.68E+03	7.20E+03		✓
c-fat200-5	-1.72E-02*	3.06E+03	7.20E+03		✓
Hamming6-2	-3.13E-02*	6.68E+04	7.20E+03		✓
Hamming6-4	-2.50E-01*	3.80E+04	7.20E+03		✓
Hamming8-2	-7.81E-03*	1.19E+03	7.20E+03		✓
Hamming8-4	-6.25E-02*	9.15E+02	7.20E+03		✓
Johnson8-2-4	-2.50E-01*	1.35E+06	7.20E+03		✓
Johnson8-4-4	-7.14E-02*	4.04E+04	7.20E+03		✓
Johnson16-2-4	-1.25E-01*	5.31E+03	7.20E+03		✓
Johnson32-2-4	-6.25E-02*	3.31E+02	7.20E+03		✓
Keller4	-9.09E-02*	1.21E+03	7.20E+03		✓
Mann-a9	-6.25E-02*	3.91E+05	7.20E+03		✓
Mann-a27	6.84E-02*	1.02E+03	7.20E+03		

this paper to instances with copositive matrices. Finally the use of these techniques to provide lower and upper bounds of copositive programming formulations of some structured global optimization problems (such as the maximum clique problem) should deserve attention in the near future.

Acknowledgements

The work of the first author was partially supported by the Portuguese Foundation for Science and Technology through the project PEst-OE/MAT/UI0297/2014 (CMA).

References

- [1] Adler, I. and Verma, S.: The Linear Complementarity Problem, Lemke Algorithm, Perturbation, and the Complexity Class PPAD. Manuscript, Department of IEOR, University of California, Berkeley, CA 94720, February (2011).
- [2] Bomze, I.M.: Copositive optimization – recent developments and applications. *Eur. J. Oper. Res.* **216**, 509–520 (2012).
- [3] Bomze, I.M. and Eichfelder, G.: Copositivity detection by difference-of-convex decomposition and ω -subdivision. *Mathematical Programming Ser. A* **138**, 365–400 (2013).
- [4] Bomze, I.M., Dür, M., de Klerk, E., Roos, C., Quist, A.J., Terlaky, T.: On copositive programming and standard quadratic optimization problems. *J. Glob. Optim.* **13**, 369–387 (1998).
- [5] Bomze, I.M., Schachinger, W. and Uchida, G.: Think co(mpletely)positive ! Matrix properties, examples and a clustered bibliography on copositive optimization. *J. Global Optim.* **52**, 425–445 (2012).

- [6] Brooke, A., Kendrick, D., Meeraus, A. and Raman, R.: *GAMS a User's Guide*. GAMS Development Corporation, Washington (1998).
- [7] Bundfuss, S.: *Copositive Matrices, Copositive Programming, and Applications*. Dissertation, Technischen Universität Darmstadt (2009).
- [8] Bundfuss, S., Dür, M.: Algorithmic copositivity detection by simplicial partition. *Linear Algebra Appl.* **428**, 1511–1523 (2008).
- [9] Burer, S.: On the copositive representation of binary and continuous nonconvex quadratic programs. *Math. Program.* **120**, 479–495 (2009).
- [10] Burer, S.: Copositive programming. In: M.F. Anjos and J.-B. Lasserre (eds.), *Handbook of Semidefinite, Cone and Polynomial Optimization: Theory, Algorithms, Software and Applications*, Springer Series in Operations Research and Management Science (2011).
- [11] Cottle, R.W., Pang, J.-S. and Stone, R.E. *The Linear Complementarity Problem*. SIAM, New York (2009).
- [12] CPLEX, I.: 11.0 Users Manual. ILOG SA, Gentilly, France (2008).
- [13] de Klerk, E., Pasechnik, D.V.: Approximation of the stability number of a graph via copositive programming. *SIAM J. Optim.* **12**, 875–892 (2002).
- [14] DIMACS: Second DIMACS Challenge. Test instances available at <http://dimacs.rutgers.edu/challenges>, last accessed 13 Jan. 2010.
- [15] Dür, M.: Copositive Programming – a survey. In: M. Diehl, F. Glineur, E. Jarlebring, W. Michiels (eds.), *Recent Advances in Optimization and its Applications in Engineering*, pp. 3-20. Springer, New York (2010).
- [16] Eichfelder, G. and Jahn, J.: Set-Semidefinite Optimization. *Journal of Convex Analysis* **15**, 767–801 (2008).
- [17] Facchinei, F. and Pang, J.-S., *Finite-Dimensional Variational Inequalities and Complementarity Problems*. Springer-Verlag, New York (2003).
- [18] Hall, M. Jr., Newman, M.: Copositive and completely positive quadratic forms. *Proc. Camb. Phil. Soc.* **59**, 329–339 (1963).
- [19] Hiriart-Urruty, J.-B., Seeger, A.: A Variational Approach to Copositive Matrices. *SIAM Rev.* **52** 593–629 (2010).
- [20] Hoffman, A.J. and Pereira, F.: On copositive matrices with -1,0,1 entries. *Journal of Combinatorial Theory (A)* **14**, 302–309 (1973).
- [21] Horst, R., Pardalos, P.M. and Thoai, N. *Introduction to Global Optimization*. Kluwer Academic Publishers, Dordrecht (2000).

- [22] Júdice, J., Faustino, A. and Ribeiro, I.: On the solution of NP-hard linear complementarity problems. *Top* **10**, 125–145 (2002).
- [23] Kaplan, W.: A copositivity probe. *Linear Algebra Appl.* **337**, 237–251 (2001).
- [24] Murtagh, B., Saunders, M., Murray, W., Gill, P., Raman, R. and Kalvelagen, E.: MINOS-NLP. Systems Optimization Laboratory, Stanford University, Palo Alto, CA.
- [25] Murty, K.G., *Linear Complementarity, Linear and Nonlinear Programming*. Heldermann Verlag, Berlin, (1988).
- [26] Murty, K.G. and Kabadi, S.N.: Some NP-complete problems in quadratic and linear programming. *Math. Programming* **39**, 117–129 (1987).
- [27] Sahinidis, N. and Tawarmalani, M.: BARON 7.2.5: Global Optimization of Mixed-Integer Nonlinear Programs. GAMS Development Corporation, Washington (2005).
- [28] Sponsel, J., Bundfuss, S., Dür, M.: An improved algorithm to test copositivity *J. Glob. Optim.* **52**, 537–551 (2012).
- [29] Tanaka, A., Yoshise, A.: An LP-based Algorithm to Test Copositivity. Discussion Paper Series No. 1320, Department of Social Systems and Management, University of Tsukuba (2014).
- [30] Väliäho, H.: Criteria for copositive matrices. *Linear Algebra Appl.* **81**, 19–34 (1986).
- [31] Väliäho, H.: Quadratic-programming criteria for copositive matrices. *Linear Algebra Appl.* **119**, 163–182 (1989).
- [32] Žilinskas, J., Dür, M.: Depth-first simplicial partition for copositivity detection, with an application to Maxclique. *Optim. Methods. Softw.* **26**, 499–510 (2011).