# Constructing Permutation Arrays from Groups

Sergey Bereg<sup>\*</sup> Avi Levy<sup>†</sup> I. Hal Sudborough<sup>\*</sup>

#### Abstract

Let M(n, d) be the maximum size of a permutation array on n symbols with pairwise Hamming distance at least d. We use various combinatorial, algebraic, and computational methods to improve lower bounds for M(n, d). We compute the Hamming distances of affine semilinear groups and projective semilinear groups, and unions of cosets of AGL(1, q) and PGL(2, q) with Frobenius maps to obtain new, improved lower bounds for M(n, d). We give new randomized algorithms. We give better lower bounds for M(n, d) also using new theorems concerning the contraction operation. For example, we prove a quadratic lower bound for M(n, n-2) for all  $n \equiv 2 \pmod{3}$  such that n+1 is a prime power.

## 1 Introduction

Two permutations  $\pi$  and  $\sigma$  on n symbols have Hamming distance  $hd(\pi, \sigma) = d$  if, for exactly d distinct elements  $x, \pi(x) \neq \sigma(x)$ . The Hamming distance of a permutation array A, denoted by hd(A), is the minimum  $hd(\pi, \sigma)$  for all permutations  $\pi \neq \sigma$ in A. Arrays of permutations on n symbols and with Hamming distance at least d between any two permutations in the array have been used for error correcting codes in communication over very noisy power line channels [13, 19]. For positive integers n and d, with  $d \leq n$ , denote by M(n, d) the maximum size of such an array. Constructing maximum size permutation arrays is difficult and, except for special cases of n and d, work has generally been limited to finding good upper and lower bounds. It is known that sharply k-transitive groups G of permutations on n symbols form a maximum size permutation array for pairwise Hamming distance d = n - k + 1[10]. Except for the exceptional case, i.e. 4- and 5-transitive Mathieu groups [4, 7, 8],

<sup>\*</sup>Department of Computer Science, Erik Jonsson School of Engineering and Computer Science, University of Texas at Dallas.

<sup>&</sup>lt;sup>†</sup>Department of Mathematics, University of Washington.

sharply k-transitive permutation groups on n symbols are only known for k = 2and k = 3 and for n a power of a prime and one more than a power of a prime, respectively, [18]. Furthermore, it is known that sharply k-transitive groups do not exist otherwise. Combinatorial arguments [5, 11] are known, which give upper and lower bounds for M(n, d), for all n and d. There are also computational approaches that have been used to construct good permutation arrays for small values of n. Due to the growth rate of n!, these computational approaches have generally been limited to small values of n and the use of automorphism groups to factor the set of all permutations into a smaller space and hence extend the range of computational processes [5, 14, 21]. It is also known that  $M(n, n - 1) \ge kn$ , where k is the number of mutually orthogonal latin squares (MOLS) of order n [6], which means computing large collections of MOLS is related to searching for large permutation arrays. Other techniques that have been used include permutation polynomials [5], and special groups, such as the Mathieu groups  $M_{22}$ ,  $M_{23}$  and  $M_{24}$  [4].

Let *n* be a positive integer and  $Z_n = \{0, 1, 2, ..., n-1\}$ . Let  $S_n$  be the symmetric group on  $Z_n$  with composition defined by  $(\pi\sigma)(i) = \sigma(\pi(i))$ . If *G* is a group, then  $G\sigma = \{g\sigma \mid g \in G\}$  and  $\sigma G = \{\sigma g \mid g \in G\}$  are called a *right coset* of *G* and a *left coset* of *G*, respectively, with the *representative*  $\sigma$  [12].

Our results, giving infinitely many improved lower bounds are obtained by describing collections of cosets of groups. We show that groups  $A\Gamma L(1, n = p^k)$  and  $P\Gamma L(2, n = p^k)$ , where p is prime, obtained from the affine general linear group AGL(1, n) and the projective general group PGL(2, n) by adding k Frobenius endomorphisms, both have pairwise Hamming distance  $n - p^{k^*}$ , where  $k^*$  denotes the largest proper<sup>1</sup> factor of k. For example,  $A\Gamma L(1, n = 2^k)$ , when k is prime, consists of kn(n-1) permutations on n symbols and has pairwise Hamming distance  $2^k - 2$ , so  $M(n, n - 2) \ge kn(n - 1)$ ,  $P\Gamma L(2, n = 2^k)$ , when k is prime, consists of k(n + 1)n(n - 1) permutations on n + 1 symbols and has pairwise Hamming distance  $2^k - 2$ , so  $M(n + 1, n - 2) \ge k(n + 1)n(n - 1)$ . We also show that the groups AGL(1, n) and PGL(2, n), where  $n = p^k$ , together with the coset defined by the single Frobenius endomorphism  $f(x) = x^p$ , has pairwise Hamming distance n - p.

Using a coset technique and computations involving random choices, we give several improved lower bounds for M(n, d), For example, we show that  $M(13, 5) \ge 6,639,048$  and  $M(14,5) \ge 58,227,624$ , which improve previous lower bounds.

<sup>&</sup>lt;sup>1</sup>A factor of k is *proper* if it is smaller than k.

## 2 New Theoretical Bounds

In this section we use group-theoretic techniques to obtain lower bounds on the sizes of permutation arrays with relatively large Hamming distance. First we consider permutation arrays obtained from semilinear groups over finite fields. Then we investigate a general procedure of modifying permutation arrays called contraction, and prove sharp Hamming distance bounds which involve the cycle structure of permutations in the permutation array.

Let  $\mathbb{F}_n$  be a field of order n. Then  $n = p^k$ , where p is a prime number throughout the remainder of the paper. A polynomial  $f(x) \in \mathbb{F}_n[x]$  is called a *permutation polynomial* if it is injective. In this case, f permutes the elements of  $\mathbb{F}_n$ .

We consider the affine general semilinear group  $A\Gamma L(1,n)$  and the projective general semilinear group  $P\Gamma L(2,n)$ . These groups are described in the appendix. It is well known that, for any permutations  $\tau, \sigma, \rho \in S_n$ ,  $hd(\sigma\tau, \sigma\rho) = hd(\tau, \rho)$ .

The following lemma shows that computing the Hamming distance of a group G does not require computing the Hamming distance between all the pairs of element in G. It shows that the computation time is O(|G|) rather than  $O(|G|^2)$ .

**Lemma 1.** Let G be a group of permutations  $G \subseteq S_n$  with |G| > 1. The Hamming distance of G, i.e. hd(G), is equal  $hd(e, G \setminus \{e\})$ , where e is the identity permutation.

*Proof.* Pick any two distinct permutations  $a, b \in G$ . Then,  $hd(a, b) = hd(e, a^{-1}b)$ . Since  $a^{-1}b$  is in G and is not the identity, the result follows.

**Lemma 2.** Let G and H be subgroups of  $S_n$ , such that  $G = \bigcup_{0 \le i \le r} a_i H$ , for some r > 0, where  $a_0 = e$ . Then,

$$hd(G) = \min(\{hd(a_i, H) \mid 0 < i \le r\}, hd(e, H \setminus \{e\})).$$

*Proof.* Observe that, since  $G = \bigcup_{0 \le i \le r} a_i H$ , then  $G = \bigcup_{0 \le i \le r} H a_i^{-1}$ . Indeed, if  $g \in G$ , then  $g = a_i h$  for some  $0 \le i \le r$  and  $h \in H$ . Then  $g^{-1} \in G$  and  $g^{-1} = h^{-1} a_i^{-1} \in H a_i^{-1}$ . So,

$$hd(G) = \min(\{hd(e, a_iH) \mid 0 < i \le r\}, hd(e, H \setminus \{e\})) \\= \min(\{hd(e, Ha_i^{-1}) \mid 0 < i \le r\}, hd(e, H \setminus \{e\})) \\= \min(\{hd(a_i, H) \mid 0 < i \le r\}, hd(e, H \setminus \{e\})).$$

**Theorem 1.** The Hamming distance of  $A\Gamma L(1,n)$  is  $n - p^{k^*}$ , where  $n = p^k$  and  $k^*$  denotes the largest proper factor of k.

Since  $A\Gamma L(1, n)$  contains kn(n-1) elements, we conclude the following.

**Corollary 1.** Let  $n = p^k$  and let  $k^*$  be the largest proper factor of  $k \ge 1$ . Then  $M(n, n - p^{k^*}) \ge kn(n-1)$ . In particular,

$$M(n, n-2) \ge kn(n-1), \qquad n = 2^k,$$

where k is prime. For example,  $M(2048, 2046) \ge 11 \cdot 2048 \cdot 2047 = 46114816$ .

**Corollary 2.** Let  $n = p^k$  for any  $k \ge 1$ . Let s be the smallest prime factor of k. Then  $M(n, n-p) \ge s \cdot q(q-1)$ . For example,

$$M(16, 14) \ge 2 \cdot 16 \cdot 15 = 480$$
  

$$M(64, 62) \ge 2 \cdot 64 \cdot 63 = 8064$$
  

$$M(81, 78) \ge 2 \cdot 81 \cdot 80 = 12960$$
  

$$M(256, 254) \ge 2 \cdot 256 \cdot 255 = 130560$$
  

$$M(512, 510) \ge 3 \cdot 512 \cdot 511 = 784896.$$

**Theorem 2.** The Hamming distance of  $P\Gamma L(2, n)$  is  $n - p^{k^*}$ , where  $k^*$  denotes the largest proper factor of k.

Since  $P\Gamma L(2, n)$  contains k(n+1)n(n-1) elements, we conclude the following.

**Corollary 3.** Let  $n = p^k$  for any  $k \ge 1$ . Let  $k^*$  be the largest proper factor of k. Then  $M(n + 1, n - p^{k^*}) \ge k(n + 1)n(n - 1)$ . In particular when  $n = 2^p$ , we have  $M(n + 1, n - 2) \ge p(n + 1)n(n - 1)$  and therefore

$$M(9,6) \ge 3 \cdot 9 \cdot 8 \cdot 7 = 1512$$
  

$$M(17,14) \ge 2 \cdot 17 \cdot 16 \cdot 15 = 8160$$
  

$$M(28,24) \ge 3 \cdot 28 \cdot 27 \cdot 26 = 58968$$
  

$$M(33,30) \ge 5 \cdot 33 \cdot 32 \cdot 31 = 163680.$$

Using the following lemma, one can compute the Hamming distance simply by counting roots of polynomials.

For any polynomial f with coefficients over a finite field  $\mathbb{F}_n$ , let r(f) denote the number of roots of f in  $\mathbb{F}_n$ . If f and g are polynomials that define permutations on  $\mathbb{F}_n$ , then the Hamming distance between the permutations f and g is equal to n minus the number of roots in the polynomial f - g, i.e. hd(f,g) = n - r(f - g). This is easily seen by observing that f(x) = g(x) is equivalent to f(x) - g(x) = 0; in other words, x is a root of the polynomial f - g.

**Lemma 3.** For any distinct polynomials  $f, g \in \mathbb{F}_n[x]$  we have

$$hd(f,g) = n - r(f - g).$$

The number of roots of polynomials of AGL(1, n) and Frobenius mappings is well known but we give a proof for completeness.

Proof of Theorem 1. Let H = AGL(1,n). Since  $A\Gamma L(1,n) = \bigcup_{0 \le i < k} x^{p^i} H$  and hd(H) = n-1, by Lemma 2 it suffices to prove  $hd(x^{p^i}, H) \ge n-p^{k^*}$  for all  $1 \le i < k$ . By Lemma 3, it suffices to prove  $r(x^{p^i} + ax + b) \le p^{k^*}$  for any  $1 \le i < k$ , and  $a \ne 0, b \in \mathbb{F}_n$ . Fix  $a \ne 0, b \in \mathbb{F}_n$ . We show that

$$r(t_{a,b}(x)) \le r(t_a(x)) \le r(t(x)), \tag{1}$$

where  $t_{a,b}(x) = x^{p^i} + ax + b$ ,  $t_a(x) = x^{p^i} + ax$ , and  $t(x) = x^{p^i} - x$ .

If  $t_{a,b}(x)$  has no root in  $\mathbb{F}_n$ , then clearly  $r(t_{a,b}(x)) \leq r(t_a(x))$ . Suppose that  $t_{a,b}(x)$  has a root, say  $y_0$ . For any root y of  $t_{a,b}(x)$ , we have  $t_a(y-y_0) = (y-y_0)^{p^i} + a(y-y_0) = y^{p^i} - y_0^{p^i} + ay - ay_0 = t_a(y) - t_a(y_0) = 0$ , where the second equation follows from the property  $(a+b)^p = a^p + b^p$  of Frobenius endomorphisms [4]. Thus,  $y - y_0$  is a root of  $t_a$ . Since the mapping  $y \to (y-y_0)$  is an injection, the first inequality of (1) follows.

We prove the second inequality of (1), by showing  $r(s_a(x)) \leq r(s(x))$ , where  $s_a(x) = t_a(x)/x = x^{p^{i-1}} + a$ , and  $s(x) = t(x)/x = x^{p^{i-1}} - 1$ . Suppose that a = 0. Then 0 is the only root of  $s_a(x)$ . Since s(x) also has the root 1, the inequality is trivially true. So, assume that  $a \neq 0$ . Then 0 is not a root of  $s_a(x)$ . We may also assume that  $s_a(x)$  has at least one root; otherwise, the inequality is trivially true. Let  $z_0$  be a root of  $s_a(x)$ . As z ranges over all roots of  $s_a(x)$ , map z to  $z/z_0$ . Observe that:

$$s\left(\frac{z}{z_0}\right) = \left(\frac{z}{z_0}\right)^{p^i - 1} - 1 = \frac{z^{p^i - 1}}{z_0^{p^i - 1}} - 1 = \frac{-a}{-a} - 1 = 0.$$

So,  $z/z_0$  is a root of s(x). Since the map is injective, it follows that  $r(t_a(x)) \leq r(t(x))$ .

Let S be the set of all roots of  $t(x) = x^{p^i} - x$ . Observe that S forms a finite field, since the set of roots are closed under the operations of addition, multiplication, and division. Thus, S is a subfield of  $\mathbb{F}_n$ , and hence the cardinality of S divides the cardinality of  $\mathbb{F}_n$ , which is  $p^k$ . So,  $|S| = p^j$ , for some j, where  $j \mid k$ . Now consider the extension of  $t(x) = x^{p^i} - x$  into its splitting field [9, 16]. In this field, the expanded root set forms  $\mathbb{F}_{p^i}$ . So, S is a subfield of  $\mathbb{F}_{p^i}$ , and so  $j \mid i$ . Thus, j divides both i and k, i.e.  $j = r(t(x)) \leq p^{gcd(i,k)} \leq p^{k^*}$ . *Proof of Theorem 2.* From the definition of  $P\Gamma L(2, n)$  we have

$$f(x) = \begin{cases} \frac{ax^{p^i} + b}{cx^{p^i} + d} & \text{if } x \in \mathbb{F}_n \text{ and } cx^{p^i} + d \neq 0, \\ \infty & \text{if } x \in \mathbb{F}_n, cx^{p^i} + d = 0 \text{ and } ax^{p^i} + b \neq 0, \\ a/c & \text{if } x = \infty \text{ and } c \neq 0, \\ \infty & \text{if } x = \infty, c = 0 \text{ and } a \neq 0. \end{cases}$$

It follows that  $S = \text{STABILIZER}_{P\Gamma L(2,n)}(\infty)$  is isomorphic to  $A\Gamma L(1,n)$ .<sup>2</sup> So,  $hd(S) = hd(A\Gamma L(1,n)) = n - p^{k^*}$  by Theorem 1. Observe that  $P\Gamma L(2,n) = \bigcup_{k=0}^{n-1} \pi_k S$ , where  $\pi_k$   $(1 \le k < n)$  is a permutation in  $P\Gamma L(2,n)$  that maps k to  $\infty$ , and  $\pi_0$  is the identity permutation. (Such permutations are in  $P\Gamma L(2,n)$ , because it is sharply 3-transitive and hence 2-transitive.) By Lemma 2,

$$hd(P\Gamma L(2,n)) = \min(\{hd(\pi_i, S) \mid 1 \le i < n\}, \{hd(e, S - \{e\})\}).$$

As  $hd(\pi_i, S) = hd(S) = hd(A\Gamma L(1, n)) \ge n - p^{k^*}$ , the result follows.

### 2.1 Contraction

In this section we describe a general method of modifying a permutation array called *contraction*. Contraction allows one to transfer a permutation array from  $S_n$  to  $S_{n-m}$  without affecting the Hamming distance by too much. First, we explain the idea of contraction for m = 1.

The contraction of a permutation  $\sigma$  on  $S_n$ , denoted by  $\sigma^{CT}$ , is the permutation on  $S_{n-1}$  defined by the following, where  $0 \le x \le n-2$ :

$$\sigma^{CT}(x) = \begin{cases} \sigma(x) & \text{if } \sigma(x) \neq n-1, \text{ and} \\ \sigma(n-1) & \text{if } \sigma(x) = n-1. \end{cases}$$
(2)

That is, the contraction  $\sigma^{CT}$  of  $\sigma$  is formed by substituting  $\sigma(n-1)$  for n-1 and deleting the symbol n-1 altogether. For instance, if  $\sigma(n-1) = n-1$ , then  $\sigma^{CT}$  is formed by simply deleting the symbol n-1. For a PA A on  $S_n$ , let  $A^{CT} = \{\sigma^{CT} \mid \sigma \in A\}$ . In general, as the contraction operation leaves most of any permutations values untouched, the Hamming distance of  $A^{CT}$  (as we shall see) is at least as large as three less than the Hamming distance of A itself (see also [22]). Specifically, a decrease of three in the Hamming distance between two permutations  $\rho$  and  $\sigma$  occurs if and only if, for some integers i, j < n-1, and symbols r, s < n-1: (a)  $\rho(i) = n-1$  and

<sup>&</sup>lt;sup>2</sup>Recall that, for a permutation group G on a set X, the *stabilizer* of an element  $x \in X$  is the set of permutations  $\{g \in G : g(x) = x\}$ .

 $\sigma(i) = r$ , (b)  $\rho(j) = s$  and  $\sigma(j) = n - 1$ , and (c)  $\rho(n - 1) = r$  and  $\sigma(n - 1) = s$ . It follows that there is a decrease of three in the Hamming distance between  $\rho$  and  $\sigma$  if and only if the permutation  $\rho^{-1}\sigma$  contains the cycle (n - 1 r s) of length 3. If A is a group, since the order of a 3-cycle is three, the order of the group must be divisible by 3 (by Lagrange's theorem [4]). So, if A is a group whose order is not divisible by 3, there can be no 3-cycle and, therefore, the contraction operation decreases the Hamming distance by at most 2.

positions ... 
$$i$$
 ...  $j$  ...  $n-1$   
 $\rho = \dots n-1$  ...  $s$  ...  $r$   
 $\sigma = \dots r$  ...  $n-1$  ...  $s$ 

Of course the contraction operation can be applied iteratively. One can first contract to a permutation on  $S_{n-1}$ , then  $S_{n-2}$ , then  $S_{n-3}$ , ... We investigate conditions on the group A that is contracted in order to understand the number of times successive contractions decrease the Hamming distance by 2.

**Lemma 4.** Let  $\sigma$  and  $\tau$  be two permutations in  $S_n$ . If  $hd(\sigma^{CT}, \tau^{CT}) = hd(\sigma, \tau) - 2$ then (i) n-1 is in a cycle C of length at least two in the cycle decomposition of  $\sigma^{-1}\tau$ , and (ii) the cycle decomposition of  $(\sigma^{CT})^{-1}\tau^{CT}$  is the same except that the length of C is decreased by two by removing n-1 and either the next or the previous element in C.

*Proof.* (i) follows from the fact that, if  $\sigma$  and  $\tau$  have n-1 in the same position, then  $hd(\sigma^{CT}, \tau^{CT}) = hd(\sigma, \tau)$ .

(ii) If one of the permutations, say  $\sigma$ , has n-1 in position n-1, then  $\sigma(i) = \tau(n-1)$  where *i* is the position of n-1 in  $\tau$ . Then  $C = (\sigma(i), n-1)$  and it is deleted in the cycle decomposition of  $(\sigma^{CT})^{-1}\tau^{CT}$ .

$\tau =$	 n-1	 s
$\sigma =$	 s	 n-1
positions	 i	 n-1

Suppose that  $\sigma(n-1) \neq n-1 \neq \tau(n-1)$  as shown below.

positions	 i	 j	 n-1
$\sigma =$	 n-1	 a	 b
$\tau =$	 c	 n-1	 d

Since  $hd(\sigma^{CT}, \tau^{CT}) = hd(\sigma, \tau) - 2$ , either b = c or a = d. Then C is  $(a \ n - 1 \ b \ d \dots)$ and  $(b \ a \ n - 1 \ c \dots)$  in these cases. The corresponding cycles in  $(\sigma^{CT})^{-1}\tau^{CT}$  are  $C - \{n - 1, b\}$  and  $C - \{a, n - 1\}$ . The lemma follows. **Theorem 3.** Suppose a permutation array  $P \subset S_n$  has Hamming distance d. Let  $Q \subseteq S_{n-2}$  denote the permutation array obtained from P by applying the contraction operation two times.

- (a) The Hamming distance of Q is at least d 6.
- (b) Suppose, for any  $\sigma, \tau \in P$ , the cycle decomposition of  $\sigma^{-1}\tau$  contains no 3-cycle and no 5-cycle. Then the Hamming distance of Q is at least d 4.

*Proof.* The part (a) follows form the fact that the Hamming distance decreases by at most three for each for each contraction operation.

The part (b) for k = 1 follows from the above discussion. We prove part (b) for k = 2. Suppose to the contrary that  $hd(Q) \leq d - 5$ . Let  $\sigma$  and  $\tau$  be two permutations of P such that  $hd(\sigma'', \tau'') \leq d - 5$  where  $\sigma'' = (\sigma')^{CT}, \tau'' = (\tau')^{CT}$  and  $\sigma' = \sigma^{CT}, \tau' = \tau^{CT}$ . Since  $\sigma^{-1}\tau$  contains no 3-cycle,  $hd(\sigma', \tau') \geq d - 2$ . Then the contraction operation on  $\sigma'$  and  $\tau'$  decreases the Hamming distance by exactly three. Therefore  $(\sigma')^{-1}\tau'$  contains a 3-cycle, say (a, b, n - 1).

positions	 i	 j	 n-1
$\sigma' =$	 b	 n-1	 a
$\tau' =$	 n-1	 a	 b

Since  $\sigma^{-1}\tau$  contains no 3-cycle and  $hd(\sigma', \tau') = d-2$ , by Lemma 4,  $\sigma^{-1}\tau$  contains a 5-cycle. Contradiction.

**Corollary 4.** (i) For each prime power n such that  $n = 2 \pmod{3}$ ,

$$M(n-1, n-3) \ge n(n-1).$$

(ii) For each prime power n such that  $n = 2 \pmod{3}$  and  $n \neq 0, 1 \pmod{5}$ ,

$$M(n-2, n-5) \ge n(n-1).$$

*Proof.* The bound can be obtained by applying the contraction operation on the affine general linear group AGL(1, n). Let  $\sigma$  and  $\tau$  be any two distinct permutations from AGL(1, n). Since  $\sigma^{-1}\tau \in AGL(1, n)$  and the order of AGL(1, n) is not multiple of 3, the bound of (i) follows. The bound of (ii) follows from Theorem 3 (b) and the fact that the order of AGL(1, n) is not multiple of 3 and 5.

Infinitely many new bounds can be obtained from Corollary 4 (c). We show some examples.

$$M(31, 29) \ge 32 \cdot 31 = 992 \quad \text{for } n = 32,$$
  

$$M(40, 38) \ge 41 \cdot 40 = 1640 \quad \text{for } n = 41,$$
  

$$M(46, 44) \ge 47 \cdot 46 = 2162 \quad \text{for } n = 47.$$

**Theorem 4.** For each prime power q such that  $q = 2 \pmod{3}$ ,

$$M(q, q-3) \ge (q+1)q(q-1).$$

Proof. Let  $A_q = (PGL(2, q+1))^{CT}$ . It suffices to show that  $hd(A_q) \ge q-3$ . Consider two permutations  $\sigma$  and  $\tau$  from PGL(2, q+1). If  $hd(\sigma, \tau) = q+1$  then  $hd(\sigma^{CT}, \tau^{CT}) \ge q-2$ . Suppose that  $hd(\sigma, \tau) \le q$ . Then  $\sigma^{-1}\tau$  has a fixed point k. A point stabilizer of PGL(2, n) is isomorphic to the affine general linear group AGL(1, q). Since its order is not multiple of 3,  $\sigma^{-1}\tau$  has no 3-cycle. Therefore  $hd(\sigma^{CT}, \tau^{CT}) \ge q-3$  and the claim follows.

An example:  $M(32, 29) \ge 33 \cdot 32 \cdot 31 = 32736$ .

## **3** Experimental Results

Computer searches for arrays of permutations on  $Z_n$  with pairwise Hamming distance d are difficult. The number of all permutations, namely n!, is large, even for small values of n. Many previous search methods have used automorphism groups to factor the space of all permutations into collections of sets of permutations with considerably smaller cardinality [5, 14, 21]. With this smaller cardinality, one constructs a graph G(n, d), whose nodes correspond to sets of permutations, with an edge between two sets  $S_1$  and  $S_2$  if the Hamming distance between permutations in the sets is at least d [5, 14, 21]. One then uses a program to find a large clique in G(n, d).

We give a different type of algorithm. We call it the *coset method*. We start with a group G that forms a good PA for M(n, d'), where d' > d. As exact values for M(q, q - 1) arise from sharply 2-transitive groups AGL(1, q), and exact values of M(q + 1, q - 1) arise from sharply 3-transitive groups PGL(2, q), where q is a power of a prime, one easily finds a group with which to start. Also, the exact value of M(n, n), for all n, arises from a cyclic group of order n. So, in fact, there is always a group with which to start. We search for a permutation, say  $\pi$ , at Hamming distance at least d from G. It follows that the entire coset  $\pi G$  is at distance at least d from G. That is,

$$hd(\pi G, G) = \max\{hd(\pi g_1, g_2) \mid g_1, g_2 \in G\}$$
  
= max{hd(\pi, g\_2 g\_1^{-1} \mid g\_1, g\_2 \in G}  
= max{hd(\pi, g) \mid g \in G},

since  $g_2g_1^{-1} \in G$  by properties of a group. So, one has a PA of cardinality  $2 \cdot |G|$ , by finding a single permutation  $\pi$ . Moreover, recording the coset representative  $\pi$  is sufficient, as all other permutations in the coset are obtained from the group G and  $\pi$ .

The process can be iterated. Suppose we have found k coset representatives  $\pi_0, \pi_1, \ldots, \pi_{k-1}$ , where  $\pi_0$  is the identity permutation, so  $\pi_0 G$  is the group G and the collection of all such cosets is a PA of cardinality  $k \cdot |G|$ . One can continue by finding a permutation  $\pi_k$  such that  $hd(\pi_k, \pi_i G) \ge d$ , for all  $0 \le i < k$ . Such a permutation gives us a new coset, namely  $\pi_k G$ , and a PA of cardinality  $(k+1) \cdot |G|$ .

One implementation of this method guesses a new permutation  $\pi_k$  randomly and then checks (1) that  $hd(\pi_k, \pi_i G) \ge d$ , for all  $0 \le i < k$ , and (2)  $hd(\pi_k, \pi_i G) = d$ , for some i ( $0 \le i < k$ ). The second condition may need explanation. Recall the combinatorial Gilbert-Varshimov (GV) lower bound [10, 17, 20] for M(n, d), namely  $N_{GV}(n, d)$ , where  $N_{GV}(n, d)$  is given by:

$$N_{GV}(n,d) = \frac{n!}{V(n,d-1)},$$
(3)

where V(n, d-1), the number of permutations that are at distance less than d from a given permutation, e.g. the identity, is  $V(n, d-1) = \sum_{k=0}^{d-1} {n \choose k} D_k$ , where  $D_k$  is the number of *derangements* on k symbols. Note that the ratio given in (3) is a lower bound for the number of times one can select another permutation without choosing two with Hamming distance less than d. It is calculated with the assumption that all sets of permutations too close to different permutations are disjoint. (It should be noted that one can often get far more than  $N_{GV}(n, d)$  permutations in a PA, because the assumption that the spheres are disjoint is not required. In fact, this combinatorial lower bound has recently been improved [11].) Of course, such sets need not be disjoint and one can choose a larger set of permutations by eliminating this condition. We do this by requiring each new permutation selected is at distance *exactly* d to one chosen before.

This simple technique makes it feasible to compute and verify large PA's. Previously, the implicit algorithm justifying the GV lower bound was not considered practical, due to large space requirements for keeping track of permutations already chosen and the large time needed for computing Hamming distances. As we have seen in Lemma 1, when the PA is a group and some of its cosets, one need not check the distance between every pair of permutations and one can store the set of coset representatives instead of the set of all permutations. It is worth noting that in [5], the authors computed a PA of size 58,322 for M(16, 9) and stated that this lower bound is not as good as what is given by the GV lower bound<sup>3</sup>. They stated, "... However, the GV lower bound is not constructive." On the other hand, the coset method starting with the group G = AGL(1, 16) found 5,739 cosets of G for Hamming distance 9 and, hence, obtained a lower bound of 1,377,360 for M(16,9).

Also, in [21] the authors computed a PA of size 20,908,800 for M(12,4) and then stated "... it is too large to check fully, but has been extensively checked." In contrast, the coset method starting with the Mathieu group G of order 12 [4, 7, 8] found 638 cosets of G for Hamming distance 4 and, hence, obtained a lower bound of 60,635,520 for M(12,4). Furthermore, since testing of correctness of G and its cosets takes far less time using group-theoretic properties, verification was done in a few minutes using a computer.

Verifying the Hamming distance of a PA A on  $Z_n$  of size N generally involves computing  $\binom{N}{2}$  pairs of permutations of n symbols, which is  $O(N^2n)$  time. When Ais a group, using Lemma 1, one only need compute the distance between the identity and the other N-1 elements, so O(Nn) time. In fact, if A is a group consisting of the identity permutation and its cyclic shifts or A consists of this cyclic group and, say, k of its cosets, the computation time is reduced to O(N) and O(kN), respectively. We discuss testing algorithms in Section 3.1.

The coset method has been used to obtain several new lower bounds. Many of the new lower bounds are given in Tables 1 and 2. The PA's that justify the lower bounds are available at our web site [1]. Sometimes finding a suitable coset representative takes considerable computation time and, hence, when found it should be recorded. For example, when started with the group G = PGL(2, 19) the coset method with difficulty found one coset of G for Hamming distance 16. It can be described by one of its representatives, for example:

#### 1, 2, 3, 4, 6, 8, 15, 5, 19, 18, 10, 7, 17, 16, 12, 20, 9, 11, 13, 14.

Thus,  $M(20, 16) \ge 13,680$ . Some other lower bounds obtained are:  $M(13,5) \ge 10,454,400$   $M(13,6) \ge 1,805,760$   $M(13,7) \ge 380,160$   $M(14,5) \ge 60,445,440$   $M(14,6) \ge 10,834,560$   $M(14,7) \ge 1,900,800$   $M(14,8) \ge 380,160$  $M(14,9) \ge 21,840$ 

<sup>3</sup>Note that  $N_{GV}(16,9) = 97,579.$ 

$$\begin{split} &M(15,6) \geq 58,734,720 \\ &M(15,7) \geq 15,491,520 \\ &M(15,8) \geq 1,900,800 \\ &M(15,9) \geq 181,272 \\ &M(15,10) \geq 32,760 \\ &M(16,7) \geq 70,709,760 \\ &M(16,8) \geq 16,061,760 \\ &M(16,9) \geq 1,377,360 \\ &M(16,10) \geq 164,880. \end{split}$$

### 3.1 Efficient Testing of a New Permutation

Suppose that we have a permutation array P consisting of k left cosets of a group G, i.e.  $P = \bigcup_{i=1}^{k} P_i$ . The critical step of our randomized construction is the computation of distance  $hd(\pi, P)$  where  $\pi$  is a random permutation. The definition of Hamming distance suggests the computation  $hd(\pi, P) = \min_{\sigma \in P} hd(\pi, \sigma)$ . Then the running time is O(n|P|) = O(nk|G|). We show that, if G is cyclic or has the cyclic subgroup, the algorithm for testing  $\pi$  can be improved. Let  $C_n$  denote the cyclic group, i.e.

$$C_n = \{g_j \mid g_j = (j, j+1, \dots, n-1, 0, 1, \dots, j-1), j \in Z_n\}.$$

**Lemma 5.** The Hamming distance from a permutation  $\pi \in S_n$  to  $C_n$  can be computed in O(n) time.

*Proof.* The Hamming distance  $hd(\pi, C_n)$  can be computed as

$$hd(\pi, C_n) = \min_{0 \le j \le n-1} \{ hd(\pi, g_j) \}.$$

The straightforward computation of  $hd(\pi, g_j)$  takes O(n) time. We show that it can be computed in O(1) amortized time.

Let D[0..n-1] be an array and we store a tentative distance of  $hd(\pi, g_j)$  in D[j]. The algorithm has 2 steps.

- 1. We initialize D[j] = n for all  $j \in Z_n$ .
- 2. For each  $m \in Z_n$ , subtract one from D[j] where  $j \equiv (\pi(m) m) \mod n$ .

Clearly, the running time is O(n). We show that the algorithm is correct. If D[j] decreases for some m, then the m-th element of  $\pi$  is  $j + m \pmod{n}$ . Thus,  $\pi$  and

 $g_j$  have matching elements in *m*-th position. In the end, D[j] is equal to n - n' where n' is the number of matching elements of  $\pi$  and  $g_j$ . The claim follows since  $hd(\pi, g_j) = n - n'$ .

**Lemma 6.** Suppose that  $G = C_n$  and let  $\pi_i$ , (i = 1, 2, ..., k) be a representative of the coset  $P_i$ , i.e.  $P_i = \pi_i C_n$ . Then, for any permutation  $\pi \in S_n$ , the Hamming distance  $hd(\pi, P)$  can be computed in O(kn) time.

*Proof.* The Hamming distance  $hd(\pi, P)$  can be computed as

$$hd(\pi, P) = hd(\pi, \bigcup_{1 \le i \le k} P_i) = \min_{1 \le i \le k} \{hd(\pi, P_i)\}$$

It suffices to show that  $d_i = hd(\pi, P_i)$  can be computed in O(n) time, for any *i*. Since  $P_i = \pi_i C_n$  and  $d_i = hd(\pi_i^{-1}\pi, C_n)$ , we first compute  $\sigma = \pi_i^{-1}\pi$  and then  $hd(\sigma, C_n)$  using the algorithm from Lemma 5.

We generalize Lemma 6 as follows.

**Theorem 5.** If  $C_n$  is a subgroup of G then the Hamming distance  $hd(\pi, P)$  can be computed in O(k|G|) time for any permutation  $\pi \in S_n$ .

*Proof.* Let  $G/C_n = \{\sigma C_n \mid \sigma \in G\}$  be the set of left cosets of  $C_n$  in G. Let  $\sigma_1, \sigma_2, \ldots, \sigma_m$  be representatives of these cosets. The Hamming distance  $hd(\pi, P)$  can be computed as

$$hd(\pi, P) = hd(\pi, \bigcup_{1 \le i \le k} \pi_i G) = hd(\pi, \bigcup_{\substack{1 \le i \le k \\ 1 \le t \le m}} \pi_i \sigma_t C_n) = \min_{\substack{1 \le i \le k \\ 1 \le t \le m}} \{hd(\pi, \pi_i \sigma_t C_n)\}.$$

Since  $hd(\pi, \pi_i \sigma_t C_n) = hd(\alpha, C_n)$  where  $\alpha = (\pi_i \sigma_t)^{-1} \pi$ . The Hamming distance  $hd(\alpha, C_n)$  can be computed in O(n) time using the algorithm from Lemma 5. The total time for computing  $hd(\pi, P)$  is O(kmn). It can be written as O(k|G|) since |G| = mn.

### 4 New tables and conclusions

We give in Tables 1 and 2 an updated partial list of lower bounds for M(n, d), with  $n \ge 9$  and  $d \ge 4$ . We also created a webpage [1] that allows one to obtain the PA's, or coded versions of them, for verification. Not all of our new results appear in the table, many are described by theorems in Section 2.

We use the following notation in the table to describe how the results are obtained:

- **a** a bound derived from  $M(n, d-1) \ge M(n, d)$ .
- **b** a bound derived from  $M(n+1, d) \ge M(n, d)$ .
- **d** a bound derived from  $M(n-1, d) \ge M(n, d)/n$ . For example, we have  $M(15, 12) \ge 2520$ , because  $M(16, 12) \ge 40320$ .
- $\mathbf{t}$  the Gilbert-Varshamov lower bound [10, 17, 20].
- **g** a lower bound based on known permutation groups. This includes the Mathieu groups  $M_{11}, M_{12}, M_{22}, M_{23}, M_{24}$  and groups AGL(1, n) or AGL(2, n), which are sharply 2-transitive and sharply 3-transitive, respectively.
- $\mathbf{m}$  a bound derived from mutually orthogonal Latin squares [6].
- $\mathbf{u}$  a result obtained by *partitioning and extending*, which is contained in [2].
- ${f r}$  a result obtained by the coset method using random search. Our improved lower bounds are in bold. The previous lower bounds are given at the bottom of the cell. When a lower bound is obtained by the coset technique, the number of cosets is given at the cell.
- **c** a result obtained by *contraction*, which is an improvement of the  $M(n-1, d-3) \ge M(n, d)$  result, described in [22]. For example,  $M(23, 20) \ge 12144$  comes from  $M(24, 22) \ge 12144$ .

We use capital letters for the following references: A - [5], B - [11], C - [14], D - [15], E - [21], F - [22], G - [3].

We note that  $M(9,6) \ge 1512$  and  $M(17,14) \ge 8160$  follow from Corollaries 1 and 2.

	d=4	d=5	d=6	d=7	d=8	d=9	d=10	d=11	d=12	d=13	d=14	d=15	d=16	wh
9	-	-	-	-	-	-	-	-	-	-	-	_	-	ich
	-	-	-	-	-	-	-	-	-	-		-	-	th
	$18576_C$	3024 <sub>E</sub>	$1512_E$	504 <sub>g</sub>	$/2_g$	9	-	-	-	-	-	-	-	e p
10	-	-	-	-	-	-	-	-	-	-		-	-	eri
	150400	-	-	$1504_u$	-	-	- 10	-	-	-	-	-	-	nut
1 1	$150480_E$	19440 <sub>C</sub>	8640 <sub>E</sub>	1484 <sub>C</sub>	720 <sub>g</sub>	49 <sub>D</sub>	10	-	-	-	-	-	-	tat
11	-	-	-	-	-	-	-	-	-	-	-	-	-	ion
	-	- 205020	- 05040	- 7020	- 7020	- 207 -	- 110	- 11	-	-	-	-	-	(n)s a
12	1742400 <u>E</u>	$203920_E$	95040 <sub>E</sub>	1920 <sub>a</sub>	1920 <sub>g</sub>	2910	110g	11	-	-	-	-		re (
12	_	-	-	-	-	-	-	-	_	_	_		_	, v def
	$20908800_{F}$	$2376000_{F}$	$190080_{F}$	95040	95040	1320	1320	112 <i>c</i>	12					ine
13	638	110	20	4			g		-	_	_	_	-	d e
	$60635520_{r}$	$10454400_r$	$1900800_r$	$380160_{r}$	-	-	-	-	-	-	_	_	-	ne n,
	41712480 <sub>A</sub>	$2376000_{b}^{\prime}$	$271908_t$	$95040_{a}^{'}$	95040 <sub>b</sub>	$6474_{C}$	1320 <sub>b</sub>	$276_{C}$	$156_{q}$	13	-	_	-	and
14	-	636	115	21	4	10	-	-	-	-	-	-	-	$\frac{1}{1}$ th
	-	$60445440_r$	$10929600_r$	$1995840_r$	$380160_r$	$21840_r$	-	-	-	-	-	-	-	ne o
	$550368000_A$	$22767826_B$	890338 <sub>t</sub>	$97547_{t}$	$95040_{b}$	6552 <sub>a</sub>	8736 <sub>C</sub>	2184 <sub>a</sub>	$2184_{g}$	59 <sub>C</sub>	14	-	-	colu
15	-	-	618	163	20	83	15	-	-	-	-	-	-	nm
	-	-	$58734720_r$	$15491520_r$	$1900800_r$	$181272_r$	$32760_r$	-	-	-		-	-	ns
	4.01E9 <sub>B</sub>	$263832788_B$	8991655 <sub>t</sub>	888533 <sub>t</sub>	97572 <sub>t</sub>	12014 <sub>t</sub>	6076 <sub>a</sub>	7540 <sub>C</sub>	2520 <sub>d</sub>	315 <sub>C</sub>	90 <sub>C</sub>	15	-	d a
16	-	-	-	744	169	5739	687	-	-	-	-	-	-	n , re
	-	-	-	$70709760_r$	$16061760_r$	$1377360_r$	$164880_r$	-	-	-	-	-	-	the
	$1.268E11_B$	3.317E9 <sub>B</sub>	-	8972298 <sub>t</sub>	888755 <sub>t</sub>	97579 <sub>t</sub>	40320 <sub>a</sub>	40320 <sub>a</sub>	$40320_{E}$	1376 <sub>C</sub>	$1376_{C}$	240 <sub>g</sub>	16	H e
17	-	-	-	-	791	2298	303	46	-	3		-	-	am
	-	-	-	-	$75176640_r$	$9375840_r$	$1236240_r$	$187680_r$	-	$12240_{r}$	8160	-	-	lmi
	7.93 $E11_B$	-	-	-	8974885 <sub>t</sub>	888727 <sub>t</sub>	97569 <sub>t</sub>	$12014_t$	83504 <sub>A</sub>	4080 <sub>a</sub>	4080 <sub>a</sub>	$4080_{g}$	$272_{g}$	

the set  $S_n$ , g distance.

	d=11	d=12	d=13	d=14	d=15	d=16	d=17	d=18	d=19	d=20	d=21	d=22	d=23	d=24	wh
18	251	40	5	-	-	-	-	-	-	-	-	-	-	-	ich
	$1228896_r$	$195840_r$	$24480_{r}$	$12240_{u}$	$8160_{u}$	-	-	-	-	-	-	-	-	-	<u>4</u> !:
	97569 <sub>t</sub>	83504 <sub>t</sub>	4896 <sub>a</sub>	4896 <sub>a</sub>	4896 <sub>a</sub>	4896 <sub>g</sub>	90 <sub>C</sub>	18	-	-	-	-	-	_	<u>Т</u> ы Я
19	-	3572	486	-	-	-	-	-	-	-	-	-	-	-	oer
	-	$1221624_r$	$166212_r$	-	$12240_G$	-	-	-	-	-	-	-	-	-	nu
	888729 <sub>t</sub>	97569 <sub>t</sub>	65322 <sub>a</sub>	$65322_{F}$	-	4896 <sub>b</sub>	342 <sub>a</sub>	342 <sub>g</sub>	19	-	-	-	-	_	Ita
20	-	1299	215	28	3	2	-	-	-	-	-	-	-	-	$r_{10}$
	-	$8885160_{r}$	$1470600_r$	$191520_r$	$20520_r$	$13680_r$	-	-	-	-	-	-	-	-	ns
	8974608 <sub>t</sub>	888729 <sub>t</sub>	$97569_t$	12014	65322	6840 <sub>a</sub>	6840 <sub>a</sub>	6840 <sub>g</sub>	120 <sub>C</sub>	20	-	-	-	-	n, a
21	-	-	-	174	23	5	2	-	-	-	-	-	-	-	de S
	-	-	-	$1190160_r$	$157320_r$	$34200_{r}$	$13680_u$	-	$333_G$	-	-	-	-	-	efir The
	-	-	888729 <sub>t</sub>	97569 <sub>t</sub>	65322 <sub>b</sub>	6840 <sub>a</sub>	6840 <sub>a</sub>	6840 <sub>b</sub>	147 <sub>a</sub>	147 <sub>C</sub>	21	-	-	_	ere
22	-	-	-	-	47233	-	1250	-	-	-	-	-	-	-	or
	-	-	-	-	$1039126_r$	-	$27500_r$	$13680_u$	$1100_{G}$	$528_d$	$104_u$	-	-	- 1	l, a
	-	-	-	-	443520 <sub>a</sub>	443520 <sub>g</sub>	6840 <sub>a</sub>	6840 <sub>b</sub>	-	220 <sub>A</sub>	121 <sub>C</sub>	22	-	-	bu' Mo
23	-	-	-	-	-	-	-	-	-	-	-	-	-	-	$\frac{th}{b}$
	-	-	-	-	-	-	-	-	$12144_a$	$12144_c$	-	-	-	-	re
	-	-	-	-	$10200960_a$	$10200960_{g}$	291456 <sub>a</sub>	291456 $_{A}$	-	-	506 <sub>a</sub>	506 <sub>g</sub>	23	-	iat iolu
24	-	-	-	-	-	-	89	-	2	-	-	-	-	-	um:
	-	-	-	-	-	-	$1080816_r$	-	$24288_r$	-	-	-	-	-	ns d
	-	-	-	-	-	$2.44E8_g$	97569 <sub>t</sub>	291456 <sub>b</sub>	23782 <sub>a</sub>	$23782_{F}$	12144 <sub>a</sub>	$12144_{g}$	$168_m$	24	$\frac{d}{d}$
25	-	-	-	-	-	-	-	-	-	-	-	-	-	-	$\frac{h}{re}$
	-	-	-	-	-	-	-	-	$61200_d$	$15600_a$	$15600_{c}$	-	-	-	$\frac{1}{10}$
	-	-	-	-	-	-	-	-	-	-	-	$12144_{b}$	600 <sub>a</sub>	600 <sub>g</sub>	H e
26	-	-	-	-	-	-	-	-	102	10	2	-	-	-	[an
	-	-	-	-	-	-	-	-	$1591200_r$	$156000_r$	$31200_{r}$	-	-	-	lg 1
	-	-	-	-	-	-	-	-	_	-	12144 <sub>t</sub>	-	-	$15600_{g}$	ing

) the set  $S_n$ , ig distance.

# References

- [1] Table of cosests. http://www.utdallas.edu/~sxb027100/cosets.
- [2] S. Bereg, L. Morales, and I. H. Sudborough. Extending permutation arrays: improving MOLS bounds. *Designs, Codes and Cryptography*, 2016. to appear.
- [3] S. Bereg, L. Morales, and I. H. Sudborough. Parallel and sequential partition and extension techniques. Manuscript, The University of Texas Dallas, 2016.
- [4] P. Cameron. *Permutation Groups*. London Mathematical Society Student Texts. Cambridge University Press, 1999.
- [5] W. Chu, C. J. Colbourn, and P. Dukes. Constructions for permutation codes in powerline communications. Des. Codes Cryptogr., 32(1-3):51-64, 2004.
- [6] C. J. Colbourn, T. Kløve, and A. C. H. Ling. Permutation arrays for powerline communication and mutually orthogonal latin squares. *IEEE Transactions on Information Theory*, 50(6):1289–1291, 2004.
- [7] J. H. Conway, R. T. Curtis, S. P. Norton, R. A. Parker, and R. A. Wilson. Atlas of Finite Groups: Maximal Subgroups and Ordinary Characters for Simple Groups. Oxford, England: Clarendon Press, 1985.
- [8] J. Dixon and B. Mortimer. *Permutation Groups*. New York: Springer-Verlag, 1996.
- [9] D. S. Dummit and R. M. Foote. Abstract Algebra (2nd ed.). John Wiley & Sons, 1999.
- [10] P. Frankl and M. Deza. On the maximum number of permutations with given maximal or minimal distance. *Journal of Combinatorial Theory, Series A*, 22(3):352 – 360, 1977.
- [11] F. Gao, Y. Yang, and G. Ge. An improvement on the Gilbert-Varshamov bound for permutation codes. *IEEE Transactions on Information Theory*, 59(5):3059– 3063, 2013.
- [12] I. N. Herstein. Topics in Algebra. John Wiley & Sons, 2nd edition edition, 1975.
- [13] S. Huczynska. Powerline communication and the 36 officers problem. Philosophical Transactions of the Royal Society of London A: Mathematical, Physical and Engineering Sciences, 364(1849):3199–3214, 2006.

- [14] I. Janiszczak, W. Lempken, P. R. J. Östergård, and R. Staszewski. Permutation codes invariant under isometries. *Des. Codes Cryptography*, 75(3):497–507, 2015.
- [15] I. Janiszczak and R. Staszewski. An improved bound for permutation arrays of length 10. Technical Report 4, Institute for Experimental Mathematics, University Duisburg-Essen, 2008.
- [16] R. Lidl and H. Niederreiter. *Finite Fields*. Cambridge University Press, 2nd edition, 1997.
- [17] F. J. MacWilliams and N. J. A. Sloane. The Theory of Error-Correcting Codes. Amsterdam: North-Holland/Elsevier, 1977.
- [18] D. Passman. Sharp transitivity. In *Permutation Groups*. New York: Benjamin, Inc., 1968.
- [19] N. Pavlidou, A. H. Vinck, J. Yazdani, and B. Honary. Power lines communications: State of the art and future trends. *IEEE Communications Magazine*, pages 34–40, 2003.
- [20] V. S. Pless and W. C. H. (Editors). Handbook of Coding Theory. Amsterdam: North-Holland/Elsevier, 1998.
- [21] D. H. Smith and R. Montemanni. A new table of permutation codes. Des. Codes Cryptogr., 63(2):241–253, 2012.
- [22] L. Yang, K. Chen, and L. Yuan. New constructions of permutation arrays. CoRR, abs/0801.3987, 2008.

### Appendix

*Finite Fields.* Let  $n = p^k$  be a prime power. There is a field  $\mathbb{F}_n$  with n elements, unique up to isomorphism. We consider groups over the field  $\mathbb{F}_n$ .

Groups. The group AGL(1, n) consists of the affine linear transformations

$$AGL(1,n) = \{ax + b | a, b \in \mathbb{F}_n, a \neq 0\},\$$

where the group operation is function composition. This group is sharply 2-transitive and has n(n-1) elements.

Denote the symbols of  $Z_{n+1}$  by  $0, 1, 2, \ldots, n-1, \infty$ . The permutations of PGL(2, n) are  $g: x \to \frac{ax+b}{cx+d}$  on  $Z_{n+1}$  such that  $a, b, c, d \in GF(n), ad \neq bc, g(\infty) = a/c, g(-d/c) = \infty$  if  $c \neq 0$  and  $g(\infty) = \infty$  if c = 0. Then |PGL(2, n)| = (n+1)n(n-1).

Recall that  $n = p^k$ . The group of affine semilinear polynomials  $A\Gamma L(1, n)$  arises as a semidirect product of AGL(1, n) with a cyclic group of order k. It is generated by iteratively composing the Frobenius automorphism  $x^p$  with the elements of AGL(1, n). Equivalently,

$$A\Gamma L(1, n) = \{ax^{p^{i}} + b \mid a, b \in \mathbb{F}_{n}, a \neq 0, 0 \le i < k\}$$

This group has kn(n-1) elements.

The group of projective semilinear polynomials  $P\Gamma L(2, n)$  arises as a semidirect product of PGL(2, n) with a cyclic group of order k generated by the Frobenius automorphism. This group has k(n + 1)n(n - 1) elements.