Restricted power domination and zero forcing problems

Chassidy Bozeman* Boris Brimkov[†] Craig Erickson[‡]

Daniela Ferrero[§] Mary Flagg[¶] Leslie Hogben[∥]

November 15, 2017

Abstract

Power domination in graphs arises from the problem of monitoring an electric power system by placing as few measurement devices in the system as possible. A power dominating set of a graph is a set of vertices that observes every vertex in the graph, following a set of rules for power system monitoring. A practical problem of interest is to determine the minimum number of additional measurement devices needed to monitor a power network when the network is expanded and the existing devices remain in place. In this paper, we study the problem of finding the smallest power dominating set that contains a given set of vertices X. We also study the related problem of finding the smallest zero forcing set that contains a given set of vertices X. The sizes of such sets in a graph G are respectively called the restricted power domination number and restricted zero forcing number of G subject to X. We derive several tight bounds on the restricted power domination and zero forcing numbers of graphs, and relate them to other graph parameters. We also present exact and algorithmic results for computing the restricted power domination number, including integer programs for general graphs and a linear time algorithm for graphs with bounded treewidth. We also use restricted power domination to obtain a parallel algorithm for finding minimum power dominating sets in trees.

Keywords: Power domination, restricted power domination, zero forcing, restricted zero forcing

AMS subject classification 05C69, 05C50, 05C57, 94C15

^{*}Department of Mathematics, Iowa State University, Ames, IA 50011, USA (cbozeman@iastate.edu).

[†]Department of Computational and Applied Mathematics, Rice University, Houston, TX, 77005, USA (boris.brimkov@rice.edu)

[‡]Department of Mathematics and Computer Science, Grand View University, Des Moines, IA, USA (cerickson@grandview.edu)

[§]Department of Mathematics, Texas State University, San Marcos, TX 78666, USA (dfer-rero@txstate.edu).

[¶]Department of Mathematics, Computer Science and Cooperative Engineering, University of St. Thomas, 3800 Montrose, Houston, TX 77006, USA (flaggm@stthom.edu).

Department of Mathematics, Iowa State University, Ames, IA 50011, USA and American Institute of Mathematics, 600 E. Brokaw Road, San Jose, CA 95112, USA (hogben@aimath.org).

1 Introduction

The study of the power domination number of a graph arose from the question of how to monitor electric power networks at minimum cost, and was recast in graph theoretical terms by Haynes et al. in [17]. In this model, vertices represent electric nodes and edges represent connections via transmission lines. Electric power companies need to continuously monitor their networks, and one method to do this is to place Phase Measurement Units (PMUs) at selected locations in the system. Because of the cost of a PMU, it is important to minimize the number of PMUs used while maintaining the ability to observe the entire system. The physical laws by which PMUs can observe the network give rise to propagation rules governing power domination; see Section 2 for formal definitions.

Electric networks are frequently modified, sometimes by building an extension with new nodes and lines. The labor to install a PMU and the (non-portable) infrastructure that must be put at the node to support the PMU are significant parts of the total cost of installing a PMU at a given node. Thus it seems useful to determine the minimum number of additional PMUs needed (and the locations where they should be placed) when an existing network is expanded and the existing PMUs remain in place. In this paper we consider the problem of finding a power dominating set that contains a given set of vertices X (locations of the existing PMUs) and minimizes the number of additional PMUs needed, or equivalently, that minimizes the total number of PMUs used subject to the constraint that the vertices in X are included in the solution.

The process of zero forcing was introduced independently in combinatorial matrix theory (where it refers to forcing zero entries in a null vector of a matrix) [1], and in mathematical physics in the control of quantum systems [6]. Zero forcing is closely related to power domination, because power domination can be described as a domination step followed by the zero forcing process (or, since power domination came first, zero forcing can be described as power domination without the domination step). The problem of finding a zero forcing set that is minimum subject to containing a set X of vertices is also considered here. Zero forcing sets containing specific vertices have been used in [3] to show the correctness of the Wavefront algorithm for zero forcing (which is implemented in [7]), and explicitly or implicitly in certain other zero forcing proofs [4, 22]. This concept may also have application to control of quantum systems where certain controls are automatically available.

This paper is organized as follows. In the next section, we recall some graph theoretic notions, specifically those related to power domination and zero forcing, and define restricted power domination and zero forcing. Additionally, we show that our study of restricted power domination falls under the general study of restricted domination parameters initiated by Goddard and Henning [15] as a generalization of the work by Sanchis [23]. In Section 3, we present structural results and bounds for restricted power domination and restricted zero forcing. In Section 4, we give exact results and algorithms for the restricted power domination and restricted zero forcing numbers of certain families of graphs. We conclude with some final remarks and open questions in Section 5.

2 Preliminaries

A graph G = (V, E) is an ordered pair consisting of a finite nonempty set of vertices V = V(G) and a set of edges E = E(G) containing unordered pairs of distinct vertices (all graphs discussed in this paper are simple, undirected, and finite). The order of G is denoted by n = |V(G)|. Two vertices u and v are adjacent, or neighbors, if $\{u, v\} \in E$. For $v \in V$, the neighborhood of v is $N(v) = \{u \in V : \{u, v\} \in E\}$ (or $N_G(v)$ if G is needed for clarity), and the closed neighborhood of v is the set $N[v] = N(v) \cup \{v\}$. Similarly, for a set of vertices S, $N[S] = \bigcup_{v \in S} N[v]$. The degree of a vertex v, denoted deg v, is the cardinality of the set N(v).

In a graph G, a vertex v dominates all vertices in $N_G[v]$, a set of vertices S dominates all vertices in $N_G[S]$, and in particular, when $N_G[S] = V(G)$ we say that S is a dominating set of G. A minimum dominating set is a dominating set of minimum cardinality, and its cardinality is the domination number of G, denoted G.

In [17] the authors introduced the related concept of power domination by presenting propagation rules in terms of vertices and edges in a graph. Here we use a simplified version of the propagation rules that is equivalent to the original (see [5]). For a set $S \subseteq V$ in a graph G = (V, E), define $PD(S) \subseteq V$ recursively:

- 1. PD(S) = N[S].
- 2. While there exists $v \in PD(S)$ such that $|N(v) \setminus PD(S)| = 1$: $PD(S) = PD(S) \cup N(v)$.

A set $S \subseteq V(G)$ is a power dominating set of a graph G if PD(S) = V(G) at the conclusion of the process above. A minimum power dominating set is a power dominating set of minimum cardinality, and the power domination number of G, denoted by $\gamma_P(G)$, is the cardinality of a minimum power dominating set.

We recall the following well-known upper bounds on the domination and power domination numbers of a graph that will be used throughout.

Theorem 2.1 (Ore's Theorem). [18, Theorem 2.1] If G is a graph on $n \ge 2$ vertices and G has no isolated vertices, then $\gamma(G) \le \frac{n}{2}$.

Theorem 2.2. [5, 27] If G is a connected graph on $n \geq 3$ vertices, then $\gamma_P(G) \leq \frac{n}{3}$.

Zero forcing can be described as a coloring game on the vertices of G. The color change rule is: If u is a blue vertex and exactly one neighbor w of u is white, then change the color of w to blue. We say u forces w and denote this by $u \to w$. Given a set B of blue vertices (all other vertices being white), the closure of B, denoted cl(B), is the set of blue vertices obtained after the color change rule is applied until no new vertex can be forced; it can be shown that cl(B) is uniquely determined by B (see [1]). A zero forcing set is a set for which cl(B) = V(G); a minimum zero forcing set is a zero forcing set of minimum cardinality, and the zero forcing number of G, denoted Z(G), is the cardinality of a minimum zero forcing set.

A chronological list of forces \mathcal{F} associated with a zero forcing set B is a sequence of forces applied to obtain cl(B) in the order they are applied. A forcing chain for a chronological list of forces is a maximal sequence of vertices (v_1, \ldots, v_k) such that the force $v_i \to v_{i+1}$ is

in \mathcal{F} for $1 \leq i \leq k-1$; we say the forcing chains are associated with B. Each forcing chain produces a distinct path in G, one of whose endpoints is in B; the other is called a terminal (and the two can coincide).

Definition 2.3. Let G = (V, E) be a graph and let $X \subseteq V$. A set $S \subseteq V(G)$ is a power dominating set of G subject to X if S is a power dominating set of G and $X \subseteq S$. The restricted power domination number of G subject to X is the minimum number of vertices in a power dominating set that contains X, and is denoted by $\gamma_P(G; X)$.

A set $S \subseteq V(G)$ is a zero forcing set of G subject to X if S is a zero forcing set of G and $X \subseteq S$. The restricted zero forcing number of G subject to X is the minimum number of vertices in a zero forcing set that contains X, and is denoted by Z(G; X).

Observation 2.4. For any graph G, $\gamma_P(G; \emptyset) = \gamma_P(G)$ and $Z(G; \emptyset) = Z(G)$.

Example 2.5. For a path P_n and any nonempty set X, $\gamma_P(P_n; X) = |X|$, because any nonempty set of vertices is a power dominating set of a path. If $X = \{x\}$ and x is an endpoint of the path, then Z(G; X) = 1 = Z(G); if x is not an endpoint of the path, then Z(G; X) = 2.

The study of restricted power domination and restricted zero forcing fit into the general study of restricted domination parameters initiated by Goddard and Henning in [15] as a generalization of the work of Sanchis [23] and Henning [19] that investigated the restricted version of standard graph domination. One motivation for Sanchis' study of restricted domination was a problem in resource allocation in distributed computer systems. We follow the notation introduced by Henning (to avoid a conflict with an existing notation) and generalized by Goddard and Henning: Let G be a graph, and $X \subseteq V(G)$. Define $r(G, X, \gamma)$ to be the cardinality of a smallest dominating set of G that contains G. The G-restricted domination number of G is G-restricted domination number of G-restricted domination G-restricted domination number of G-rest

Returning to the more general setting and using the terminology of [15], let π be a property of sets, such as being a dominating, power dominating, or zero forcing set and denote the minimum cardinality of a π -set by f_{π} . Given a set of vertices X, define $r(G, X, f_{\pi})$ to be the minimum cardinality of a π -set containing X. Thus, $\gamma_P(G; X) = r(G, X, \gamma_P)$ and Z(G; X) = r(G, X, Z). Given a non-negative integer k, the k-restricted π -number of G is $r_k(G, f_{\pi}) = \max\{r(G, X, f_{\pi}) : X \subseteq V(G), |X| = k\}$.

Remark 2.6. Since for any graph G, $\gamma_P(G) \leq \gamma(G)$, it follows that $r(G, X, \gamma_P) \leq r(G, X, \gamma)$ for any $X \subseteq V(G)$. As a consequence, $r_k(G, \gamma_P) \leq r_k(G, \gamma)$ and $r(G, X, \gamma_P) \leq r_k(G, \gamma)$ for k = |X|.

Other generalizations and extensions of power domination and zero forcing have also been explored; for example, variants of the problems obtained by modifying the color change rules have received significant attention [8, 9, 14, 26], as has the problem of studying the number of timesteps involved in the process of zero forcing or power dominating a graph [13, 20, 21].

Algorithmically, given a graph G = (V, E) and sets $X \subseteq V$ and $S \subseteq V$, deciding if S is a power dominating set of G subject to X is simply a matter of determining whether S contains X and whether S is a power dominating set. As a consequence, given an arbitrary graph

G = (V, E), an arbitrary set of vertices $X \subseteq V$, and an integer $k \ge |X|$, the computational complexity of deciding if a graph G has a power dominating set of cardinality k is equivalent to that of determining if G has a power dominating set of cardinality k that also contains X. The former problem has been proven to be NP-complete for bipartite graphs [17, Theorem 5], chordal graphs [17, Theorem 5], circle and planar graphs [16, Theorem 1].

3 Structural results and bounds

In this section, we present a number of structural results and bounds on the restricted power domination and restricted zero forcing numbers of a graph. We begin with the following simple bounds.

Proposition 3.1. For any graph G = (V, E) and any set $X \subseteq V$,

$$\gamma_P(G) \leq \gamma_P(G; X) \leq \gamma_P(G) + |X|,$$

$$Z(G) \leq Z(G; X) \leq Z(G) + |X|,$$

$$|X| \leq \gamma_P(G; X) \leq Z(G; X),$$

and all bounds are tight.

Proof. The lower bounds are all immediate from the definitions. The first two upper bounds follow by adding the vertices of X into a power dominating or zero forcing set, and the last follows from the fact that any zero forcing set is a power dominating set. For tightness: Let X consist of s leaves in $K_{1,p}$ where $1 \le s \le p-2$, and note that $\gamma_P(K_{1,p};X) = 1+s = \gamma_P(K_{1,p}) + |X|$. Various choices of X in the path can be used to show the remaining bounds are tight; see Example 2.5.

By a similar reasoning, the following more general bounds hold.

Proposition 3.2. For any graph G = (V, E) and any sets $Y \subseteq X \subseteq V$,

$$\gamma_P(G;Y) \le \gamma_P(G;X) \le \gamma_P(G;Y) + |X\backslash Y|,$$

 $Z(G;Y) \le Z(G;X) \le Z(G;Y) + |X\backslash Y|,$

and all bounds are tight.

In the next section, we give structural results related to the restricted power dominating sets of a graph. These lead to improved bounds on the restricted power domination and zero forcing numbers in Sections 3.2 and 3.3. In Section 3.4 we establish a relationship between the restricted power domination number and the restricted zero forcing number.

3.1 Structural results for power domination

The next lemma states well-known results about mandatory vertices in power dominating sets; we include the brief proof for completeness (and because it has sometimes been misstated in the literature without proof). A *leaf* is a vertex of degree one.

Lemma 3.3. Let G = (V, E) be a graph.

- (1) If $v \in V$ is incident to three or more leaves, then v is contained in every minimum power dominating set of G.
- (2) If $v \in V$ is incident to exactly two leaves, then every minimum power dominating set contains either v or one of its two leaf neighbors.
- (3) There is a minimum power dominating set of G that contains every vertex v that is incident to exactly two leaves.

Proof. First let $v \in V$ be incident to at least three leaves and suppose there is a minimum power dominating set S of G that does not contain v. If S excludes two or more of the leaves of G incident to v, then those leaves cannot be dominated or forced at any step. Thus, S excludes at most one leaf incident to v, which means S contains at least two leaves ℓ_1 and ℓ_2 incident to v. Then, $(S \setminus \{\ell_1, \ell_2\}) \cup \{v\}$ is a smaller power dominating set than S, which is a contradiction.

Now consider the case in which $v \in V$ is incident to exactly two leaves, ℓ_1 and ℓ_2 , and suppose there is a minimum power dominating set S of G such that $\{v, \ell_1, \ell_2\} \cap S = \emptyset$. Then neither ℓ_1 nor ℓ_2 can be dominated or forced at any step, contradicting the assumption that S is a power dominating set. If S is a power dominating set that contains ℓ_1 or ℓ_2 , say ℓ_1 , then $(S \setminus \{\ell_1\}) \cup \{v\}$ is also a power dominating set and has the same cardinality. Applying this to every vertex incident to exactly two leaves produces the minimum power dominating set required by (3).

Definition 3.4. Given a graph G = (V, E) and a set $X \subseteq V$, define $\ell_r(G, X)$ as the graph obtained by attaching r leaves to each vertex in X. If $X = \{v_1, \ldots, v_k\}$, we denote the r leaves attached to vertex v_i as $\ell_i^1, \ldots, \ell_i^r$ for each $i = 1, \ldots, k$, so that $V(\ell_r(G, X)) = V \cup \{\ell_i^j : 1 \le i \le k, 1 \le j \le r\}$ and $E(\ell_r(G, X)) = E \cup \{v_i \ell_i^j : 1 \le i \le k, 1 \le j \le r\}$.

We will now establish a relationship between $\gamma_P(G;X)$ and $\ell_2(G,X)$ using Lemma 3.3.

Proposition 3.5. For any graph G = (V, E) and any set $X \subseteq V$, $\gamma_P(G; X) = \gamma_P(\ell_2(G, X))$.

Proof. First note that any power dominating set of G that contains X is also a power dominating set of $\ell_2(G,X)$. Thus, $\gamma_P(\ell_2(G,X)) \leq \gamma_P(G;X)$.

By Lemma 3.3, there exists a minimum power dominating set S of $\ell_2(G, X)$ that contains all vertices in X. Clearly S is a power dominating set of G containing X, and this implies $\gamma_P(G; X) \leq \gamma_P(\ell_2(G, X))$.

Since the motivation for studying the power domination number subject to a given set stems from the problem of optimizing PMU placement when the electric grid is expanded, we sometimes assume that all graphs are connected and have at least three vertices (however, subgraphs of a given graph are allowed to be disconnected). This can make the statements of some of the following results cleaner, and they are easy to adapt to disconnected graphs that may have small components: Since K_1 and K_2 each have a power domination number of one,

$$\gamma_P(G; X) = \sum_{\{i: |C_i| \ge 3\}} \gamma_P(C_i; X \cap C_i) + \sum_{\{i: |C_i| \le 2\}} \max\{|X \cap C_i|, 1\}$$

where C_1, \ldots, C_r are the connected components of G.

Corollary 3.6. Let G be a connected graph of order $n \geq 3$ and $X \subseteq V(G)$. Then $\gamma_P(G; X) \leq \left| \frac{n+2|X|}{3} \right|$ and this bound is tight.

Proof. By Proposition 3.5, $\gamma_P(G;X) = \gamma_P(\ell_2(G,X))$. Since $|V(\ell_2(G,X))| = n + 2|X| \ge 3$, it follows from Theorem 2.2 that $\gamma_P(\ell_2(G,X)) \le \left\lfloor \frac{n+2|X|}{3} \right\rfloor$.

To see that this bound is tight, let G = (V, E) and $X \subseteq V$ and construct G' = (V', E') by attaching two leaves to each vertex in $V \setminus X$, that is, let $G' = \ell_2(G, V \setminus X)$. Then $\gamma_P(G'; X) = |V| = \left\lfloor \frac{|V'| + 2|X|}{3} \right\rfloor$.

As noted in Remark 2.6, $\gamma_P(G;X) \leq r_k(G,\gamma)$ for $X \subseteq V(G)$ and k=|X|. Let G be a connected graph of order n and minimum degree at least 2. It was shown in [19] that $r_k(G,\gamma) \leq \frac{2n+3k}{5}$ for every integer k such that $1 \leq k \leq n$. As a consequence, for every nonempty set $X \subseteq V(G)$, $\gamma_P(G;X) \leq \frac{2n+3|X|}{5}$. However, Corollary 3.6 gives a better bound. Lemma 3.3 ensures that X is contained in *some* but not *every* minimum power dominat-

Lemma 3.3 ensures that X is contained in *some* but not *every* minimum power dominating set of $\ell_2(G, X)$. For example, in Figure 3.1 let G be the path with vertex set $\{3, 4, 5\}$, let $X = \{3, 5\}$, and note that $\{1, 5\}$ is a minimum power dominating set of $\ell_2(G, X)$ that does not contain X.




Figure 3.1: A graph used in several examples.

The next result shows that X is contained in every minimum power dominating set of $\ell_3(G, X)$. The proof is analogous to the proof of Proposition 3.5 and is omitted.

Proposition 3.7. For any graph G = (V, E) and any set $X \subseteq V$, a set S is a minimum power dominating set of G subject to X if and only if S is a minimum power dominating set of $\ell_3(G, X)$.

Lemma 3.8. Let G be any graph and let u be a leaf vertex of G. Then the following are equivalent:

- 1. $\gamma_P(G; \{u\}) = \gamma_P(G u) + 1$.
- 2. There is some minimum power dominating set of G subject to $\{u\}$ in which u does not need to perform a dominating step or a force.
- 3. There is some minimum power dominating set of G subject to $\{u\}$ of the form $S \cup \{u\}$ where S is a minimum power dominating set of G u.

Furthermore, $\gamma_P(G; \{u\}) = \gamma_P(G - u)$ if and only if u needs to perform a dominating step or a force for any minimum power dominating set of G subject to $\{u\}$.

Proof. Since u is a leaf, a dominating step is the same as a force for u. Furthermore, $\gamma_P(G-u) \leq \gamma_P(G) \leq \gamma_P(G-u) + 1$ and $\gamma_P(G) \leq \gamma_P(G; \{u\}) \leq \gamma_P(G-u) + 1$. We show that $(2) \Rightarrow (3) \Rightarrow (1) \Rightarrow (2)$. Then the last statement follows from $(1) \Leftrightarrow (2)$.

If there is some minimum power dominating set S' of G subject to $\{u\}$ in which u does not need to perform a dominating step or a force, then $S = S' \setminus \{u\}$ is a power dominating set of G - u, and S is minimum since $\gamma_P(G) - 1 \leq \gamma_P(G - u)$.

If there is some minimum power dominating set of G subject to $\{u\}$ of the form $S \cup \{u\}$ where S is a minimum power dominating set of G - u, then $\gamma_P(G; \{u\}) = \gamma_P(G - u) + 1$.

Now suppose that $\gamma_P(G; \{u\}) = \gamma_P(G - u) + 1$. Let w be the neighbor of u in G. Then $\gamma_P(G - u) + 1 = \gamma_P(G - uw) = \gamma_P(G - uw; \{u\})$. Since $\gamma_P(G - uw; \{u\}) = \gamma_P(G; \{u\})$, the presence of the edge incident to u does not affect the restricted power domination number, so u does not need to perform a force in some minimum power dominating set S' of G subject to $\{u\}$.

3.2 Bounds for restricted power domination

If W is a subset of the vertices of a graph H, the subgraph of H induced by W is denoted by H[W].

Proposition 3.9. Let G' = (V', E') be a graph, $V \subset V'$, G = G'[V], and S be a power dominating set of G. Let t be the number of isolated vertices of $G'[V' \setminus V]$. Then,

$$\gamma_P(G';S) \le |S| + \frac{|V' \setminus V|}{2} + \frac{t}{2}$$

and this bound is tight.

Proof. Let X_1 be the set of isolated vertices of $G'[V' \setminus V]$. By Theorem 2.1, the $|V' \setminus V| - t$ non-isolated vertices of $G'[V' \setminus V]$ can be dominated by a set X_2 of size at most $\frac{|V' \setminus V| - t}{2}$. Thus, the set $X_1 \cup X_2$ can dominate all vertices in $G'[V' \setminus V]$ in the first time step, and then the rest of G' can be power dominated by S. Thus, $\gamma_P(G'; S) \leq |S| + \frac{|V' \setminus V| - t}{2} + t = |S| + \frac{|V' \setminus V|}{2} + \frac{t}{2}$.

To see that the bound is tight, consider $G = P_n \square P_2$ with n > 3, denote the vertices of G by (i,j) for $1 \le i \le n$ and j = 1, 2, and let $S = \{(1,1)\}$. Construct G' from G by adding one K_2 with both vertices adjacent to (1,2) and another K_2 with both vertices adjacent to (n,2) (see Figure 3.2 for an example). Since for both $\ell = 1$ and $\ell = n$, one of $(\ell,2)$ or a vertex from $N_{G'}[(\ell,2)] \setminus V(G)$ must be in any power dominating set of G', $\gamma_P(G';S) = 3 = 1 + \frac{4}{2} + \frac{0}{2}$. \square




Figure 3.2: Graph G' for Proposition 3.9.

The next proposition is a basic extension result.

Proposition 3.10. Let G' = (V', E') be a graph, $V \subset V'$, G = G'[V], H_1, \ldots, H_k be the components of $G'[V' \setminus V]$, and S be a power dominating set of G. For $1 \leq i \leq k$, let $N_i = V(H_i) \cap N_{G'}[V \setminus N_G[S]]$. Then

$$\gamma_P(G'; S) \le \sum_{i=1}^k \gamma_P(H_i; N_i) + |S|$$

and this bound is tight.

Proof. For i = 1, ..., k, let S_i be any power dominating set of H_i that contains N_i . We show $\bigcup_{i=1}^k S_i \cup S$ is a power dominating set of G', so choosing minimum power dominating sets S_i subject to the restrictions establishes the inequality. Suppose $y \in N_{G'}[H_i] \cap V$, which implies y has a neighbor $x \in V(H_i)$. If $y \notin N_G[S]$, then $x \in N_i$, so S_i dominates y. If $y \in N_G[S]$, then y is dominated by S. Since all neighbors of H_i that are not in H_i are dominated, all the vertices of H_i can be power dominated by S_i . Once all vertices not in G are power dominated, S can power dominate G.

To see that the bound is tight, let G' be the graph in Figure 3.1, $G = G'[\{1, 2, 3, 4\}]$, and $S = \{3\}$. Then $\gamma_P(G'; S) = 2 = 1 + 1 = \gamma_P(G'[\{5, 6, 7\}]; \{5\}) + |S|$. Larger examples may be constructed by subdividing the edge between 3 and 4 repeatedly.

Remark 3.11. The statement of Proposition 3.10 remains valid when N_i is re-defined as a subset of $V(H_i)$ which is a minimum dominating set of $V(H_i) \cap (N_{G'}[V] \setminus N_{G'}[S])$. The proof for this modified statement is similar to the proof of Proposition 3.10, except that the order is different: Since all the vertices in $N_{G'}[V] \setminus V$ are dominated, the vertices in G can be power dominated, followed by the vertices of H_i , i = 1, ..., k.

Proposition 3.12. Let G' = (V', E') be a graph, $V \subset V'$, G = G'[V], and S be a power dominating set of G. Suppose each component of $G'[V' \setminus V]$ has at least 3 vertices. Then,

$$\gamma_P(G';S) \le |S| + \frac{|V' \setminus V|}{3} + |N_{G'}[V \setminus N_G[S]] \cap (V' \setminus V)|$$

and this bound is tight.

Proof. Let H_1, \ldots, H_k be the components of $G'[V' \setminus V]$. For $1 \leq i \leq k$, let $N_i = V(H_i) \cap N_{G'}[V \setminus N_G[S]]$. From Proposition 3.10, $\gamma_P(G'; S) \leq \sum_{i=1}^k \gamma_P(H_i; N_i) + |S|$. From Proposition 3.1, $\gamma_P(H_i; N_i) \leq \gamma_P(H_i) + |N_i|$ for $i = 1, \ldots, k$, and $\gamma_P(H_i) \leq \frac{|V(H_i)|}{3}$ by Theorem 2.2. Thus, we conclude $\gamma_P(G'; S) \leq \sum_{i=1}^k \frac{|V(H_i)|}{3} + \sum_{i=1}^k |N_i| + |S|$. Since $\sum_{i=1}^k |V(H_i)| = |V' \setminus V|$ and $\sum_{i=1}^k |N_i| = N_{G'}[V \setminus N_G[S]] \cap (V' \setminus V)$, we obtain $\gamma_P(G'; S) \leq \frac{|V' \setminus V|}{3} + |N_{G'}[V \setminus N_G[S]] \cap (V' \setminus V) + |S|$.

To see that the bound is tight, consider $G = K_n$ with vertices $V = \{v_1, \ldots, v_n\}$. For each $i = 1, \ldots, n$, add three vertices a_i, b_i and c_i , and also add the edges $v_i a_i, a_i b_i$ and $a_i c_i$. Let G' be the resulting graph so |V'| = 4n (see Figure 3.3 for an example). Then, G = G'[V], $S = \{v_1\}$ is a power dominating set of G, and the connected components of $G[V' \setminus V]$ are the paths b_i, a_i, c_i for $i = 1, \ldots, n$. It is easily verified that $S \cup \{b_i : i = 1, \ldots, n\}$ is a power dominating set and every S-restricted power dominating set must contain S and at least one vertex in each component of $V' \setminus V$, so $\gamma_P(G'; S) = 1 + n$. Since $N_G[S] = V$, clearly $V \setminus N_G[S] = \emptyset$, so $|N_{G'}[V \setminus N_G[S]] \cap (V' \setminus V)| = 0$. Since $|V' \setminus V| = 3n$ and |S| = 1, the upper bound is $|S| + \frac{|V' \setminus V|}{3} + |N_{G'}[V \setminus N_G[S]] \cap (V' \setminus V)| = 1 + n + 0 = \gamma_P(G'; S)$. \square




Figure 3.3: Graphs $G = K_4$ and G' for Proposition 3.12.

Proposition 3.13. Let G = (V, E) be a graph, and $V = V_1 \cup V_2$ be a partition of V. Let $N_1 = N[V_2] \cap V_1$ and $N_2 = N[V_1] \cap V_2$, and let $G_1 = G[V_1]$ and $G_2 = G[V_2]$. Let $W_1 \subseteq V_1$ and $W_2 \subseteq V_2$ be sets of vertices such that $W_1 \cup W_2$ dominates $N_1 \cup N_2$ in G. Then,

$$\gamma_P(G) \le \gamma_P(G; W_1 \cup W_2) \le \gamma_P(G_1; W_1) + \gamma_P(G_2; W_2)$$

and these bounds are tight.

Proof. Let S_1 and S_2 be sets that realize $\gamma_P(G_1; W_1)$ and $\gamma_P(G_2; W_2)$, respectively. By construction, $S_1 \cup S_2$ contains $W_1 \cup W_2$. Moreover, the vertices in $S_1 \cup S_2$ can dominate $N_1 \cup N_2$ in G in the first time step, and then S_1 and S_2 , respectively, can force the vertices of V_1 and V_2 in G independently, since the vertices in V_1 (respectively, V_2) are not adjacent to any uncolored vertices outside V_1 (respectively, V_2). Thus, $S_1 \cup S_2$ is a power dominating set of G, so $\gamma_P(G; W_1 \cup W_2) \leq \gamma_P(G_1; W_1) + \gamma_P(G_2; W_2)$.

To see that the bound is tight, let $G_1 = K_{1,p}$ and $G_2 = K_{1,q}$ with $p, q \ge 4$. Label the centers of the stars c_1 and c_2 , respectively. Let u_1 and v_1 be two leaves of G_1 and u_2 and v_2 be two leaves of G_2 . Let G be the graph formed from the disjoint union of G_1 and G_2 by adding all possible edges between a vertex in $\{u_1, v_1\}$ and a vertex in $\{u_2, v_2\}$ (see Figure 3.4 for an example). For the obvious partition of the vertices into $V(G_1), V(G_2)$, let $W_1 = \{c_1\}$ and $W_2 = \{c_2\}$. Then $\gamma_P(G) = 2 = \gamma_P(G; W_1 \cup W_2) = \gamma_P(G[V_1]; W_1) + \gamma_P(G[V_2]; W_2)$. \square




Figure 3.4: Graph G for Proposition 3.13.

3.3 Bounds for restricted zero forcing

The next two results are the zero forcing versions of Remark 3.11 and Proposition 3.13.

Proposition 3.14. Let G' = (V', E') be a graph, $V \subset V'$, G = G'[V], H_1, \ldots, H_k be the components of $G'[V' \setminus V]$, and B be a zero forcing set of G. For $1 \leq i \leq k$, let $N_i = N_{G'}[V] \cap V(H_i)$. Then,

$$Z(G'; B) \le \sum_{i=1}^{k} Z(H_i; N_i) + |B|$$

and this bound is tight.

Proof. For i = 1, ..., k, let B_i be any zero forcing set of H_i that contains N_i . We show $\bigcup_{i=1}^k B_i \cup B$ is a zero forcing set of G', so choosing minimum zero forcing sets B_i subject to the restrictions establishes the inequality. Since all vertices in $N_{G'}[V] \setminus V$ are initially colored blue, B can force all vertices of G. After this, since all vertices in $N_{G'}[V(H_i)] \setminus V(H_i)$ are colored blue for each H_i , B_i will be able to force $V(H_i)$ for $1 \le i \le k$.

To see that the bound is tight, let G' be the graph in Figure 3.1, $G = G'[\{1, 2, 3\}]$, and $B = \{1\}$. Then $H_1 = \{4, 5, 6, 7\}$, $N_1 = \{4\}$, and $Z(G'; B) = 3 = Z(H_1; N_1) + |B|$.

Proposition 3.15. Let G = (V, E) be a graph, and $V = V_1 \cup V_2$ be a partition of V. Let $N_1 = N[V_2] \cap V_1$ and $N_2 = N[V_1] \cap V_2$, and let $G_1 = G[V_1]$ and $G_2 = G[V_2]$. Then,

$$Z(G) \le \min(Z(G_1) + Z(G_2; N_2), Z(G_1; N_1) + Z(G_2)),$$

and this bound is tight.

Proof. Without loss of generality, assume $Z(G_1) + Z(G_2; N_2) \leq Z(G_1; N_1) + Z(G_2)$. Start with an initial set $B = B_1 \cup B_2$ where B_1 and B_2 are minimum zero forcing sets of G_1 and G_2 , respectively, and $N_2 \subseteq B_2$. Given a vertex v in V_1 , its only white neighbors belong to G_1 , hence G_1 can force all of G_1 . Once G_1 is colored blue, G_2 is a zero forcing set of G_2 and will force the rest of the graph. To see that this bound is tight, consider the graph G_1 shown in Figure 3.5. Partition the vertices as $G_1 = \{1, 2, 3\}$ and $G_2 = \{4, 5, 6, 7, 8\}$, so $G_1 = \{2\}$ and $G_2 = \{6, 7, 8\}$. Then $G_1 = \{2\}$ and $G_2 = \{6, 7, 8\}$. Then $G_2 = \{2\}$ and $G_3 =$




Figure 3.5: Graph G for Proposition 3.15.

We conclude this section by considering Z(G;X) when X consists of a single vertex v. To this end, we recall the concept of zero forcing spread of a vertex, as defined in [11]. Let G be a graph and v be a vertex in G. The zero forcing spread of v is defined as $z_v(G) = Z(G) - Z(G - v)$.

It is shown in [11] that for any vertex v in a graph G, $-1 \le z_v(G) \le 1$, and that $z_v(G) = 1$ if and only if there exists a minimum zero forcing set of G containing v such that v does not perform a force. Furthermore, if $z_v(G) = -1$, then v is not contained in any minimum zero forcing set of G. Since $Z(G) \le Z(G; \{v\}) \le Z(G) + 1$, we have the following result.

Proposition 3.16. For a graph G and $v \in V(G)$

$$Z(G; \{v\}) = \begin{cases} Z(G) & \text{if } z_v(G) = 1\\ Z(G) + 1 & \text{if } z_v(G) = -1. \end{cases}$$

For a graph G and vertex $v \in V(G)$ with $z_v(G) = 0$, it may be that $Z(G; \{v\}) = Z(G)$ or $Z(G; \{v\}) = Z(G) + 1$. In Figure 3.6, $z_v(G) = 0$ and $Z(G; \{v\}) = Z(G) = 2$, while $z_u(G) = 0$ and $Z(G; \{u\}) = 3 = Z(G) + 1$.

Note that by numbering the desired vertex v as the lowest number 0, the Bruteforce algorithm implemented in the software [7] will return the desired vertex in a minimum zero forcing set if this is possible. Thus the Bruteforce algorithm can be used to determine whether $Z(G; \{v\}) = Z(G)$ or $Z(G; \{v\}) = Z(G) + 1$.




Figure 3.6: Graph G in which $z_v(G) = 0$ and $Z(G; \{v\}) = Z(G)$, while $z_u(G) = 0$ and $Z(G; \{u\}) = Z(G) + 1$.

3.4 Connections between restricted power domination and restricted zero forcing

Let the maximum degree of G be denoted by $\Delta(G)$.

Remark 3.17. If S is a power dominating set of G that contains X, then N[S] is a zero forcing set of G that contains N[X]. Since $|N[S]| \leq (\Delta(G) + 1)|S|$, we have

$$Z(G; N[X]) \le (\Delta(G) + 1)\gamma_P(G; X). \tag{1}$$

The next results improve the bound in (1); their proofs use ideas from the proofs of [10, Lemma 2] and [2, Theorem 3.2], but differ in ensuring that the set X is not inadvertently discarded.

Lemma 3.18. Let G be a graph with no isolated vertices and let $S = \{u_1, \ldots, u_t\}$ be a power dominating set for G that contains X. Then $Z(G; X) \leq \sum_{i=1}^t \deg u_i$ and this bound is tight.

Proof. For $i = 1, \ldots, t$, define

$$U_i = \begin{cases} N[u_i] \setminus \{v_i\} & \text{where } v_i \in N(u_i) \setminus S \text{ if } N(u_i) \setminus S \neq \emptyset \\ \{u_i\} & \text{if } N(u_i) \setminus S = \emptyset, \end{cases}$$

and $B = \bigcup_{i=1}^t U_i$. Then $|B| \leq \sum_{i=1}^t \deg u_i$, because $|N[u_i] \setminus \{v_i\}| \leq \deg u_i + 1 - 1 = \deg u_i$ when $N(u_i) \setminus S \neq \emptyset$, and when $N(u_i) \setminus S = \emptyset$, $|\{u_i\}| = 1$ and $\deg u_i \geq 1$.

We show that B is a zero forcing set of G. Since $B \supseteq S \supseteq X$, this implies $Z(G;X) \le |B| \le \sum_{i=1}^t \deg u_i$. Color all the vertices in B blue and the vertices in $V(G) \setminus B$ white. Then every vertex of S is colored blue, and for $i = 1, \ldots, t$, at most one vertex in $N[u_i]$ is white (v_i) is the only possibility). So u_i can immediately force v_i if necessary, and N[S] is colored blue. Since S is a power dominating set of G, N[S] is a zero forcing set of G, and therefore G is a zero forcing set of G.

Whenever $Z(G) \leq \sum_{i=1}^t \deg u_i$ is tight and $X \subseteq S$, then the bound $Z(G;X) \leq \sum_{i=1}^t \deg u_i$ is tight. A path with S = X consisting of an endpoint is an example.

Theorem 3.19. Let G be a graph with $\Delta(G) \geq 1$. Then $\left\lceil \frac{Z(G;X)}{\Delta(G)} \right\rceil \leq \gamma_P(G;X)$ and this bound is tight.

Proof. Suppose G has connected components G_1, \ldots, G_h . Let $X_j = X \cap V(G_j)$ and suppose G_i is a component that has an edge, so G_i does not have isolated vertices. Choose a minimum power dominating set $S_i = \{u_1^{(i)}, \ldots, u_{t_i}^{(i)}\} \supseteq X_i$ for G_i , so $t_i = \gamma_P(G_i; X_i)$. Then by Lemma 3.18, $Z(G_i; X_i) \le \sum_{i=1}^{t_i} \deg u_i^{(i)} \le t_i \Delta(G_i) = \gamma_P(G_i; X_i) \Delta(G_i)$. Thus $\gamma_P(G_i; X_i) \ge \left\lceil \frac{Z(G_i; X_i)}{\Delta(G_i)} \right\rceil \ge \left\lceil \frac{Z(G_i; X_i)}{\Delta(G_i)} \right\rceil$.

Since $\Delta(G) \geq 1$, we have $\gamma_P(G_j; X_j) \geq \left\lceil \frac{Z(G_j; X_j)}{\Delta(G)} \right\rceil$ for every component G_j of G (including isolated vertices). Thus,

$$\gamma_P(G;X) = \sum_{j=1}^h \gamma_P(G_j;X_j) \ge \sum_{j=1}^h \left\lceil \frac{Z(G_j;X_j)}{\Delta(G)} \right\rceil \ge \left\lceil \frac{\sum_{j=1}^h Z(G_j;X_j)}{\Delta(G)} \right\rceil = \left\lceil \frac{Z(G;X)}{\Delta(G)} \right\rceil.$$

Whenever $\left\lceil \frac{Z(G)}{\Delta(G)} \right\rceil \leq \gamma_P(G)$ is tight and $X \subseteq S$, then $\left\lceil \frac{Z(G;X)}{\Delta(G)} \right\rceil \leq \gamma_P(G;X)$ is tight. An example is K_n because $Z(K_n) = \Delta(K_n) = n-1$ and $\gamma_P(K_n) = 1$.

4 Exact and algorithmic results

In this section, we present exact results and algorithms for $\gamma_P(G; X)$ and Z(G; X) for certain families of graphs.

4.1 Graphs with bounded treewidth

Our first result uses Proposition 3.7 to show that the restricted power domination number of any graph with bounded treewidth can be computed efficiently.

Theorem 4.1. For any graph G = (V, E) with bounded treewidth and any set $X \subseteq V$, a minimum power dominating set of G subject to X can be computed in O(n) time.

Proof. The treewidth of G does not change when an arbitrary number of leaves are appended to some or all of the vertices of G, so if G has treewidth at most k then $\ell_3(G,X)$ also has treewidth at most k. By Proposition 3.7, a set S is a minimum power dominating set of G subject to X if and only if S is a minimum power dominating set of $\ell_3(G,X)$. Since $|V(\ell_3(G,X))| \leq 4|V(G)|$, the linear time algorithm presented by Guo et al. in [16] applied to $\ell_3(G,X)$ yields a minimum power dominating set of $\ell_3(G,X)$ in O(n) time. Thus, $\gamma_P(G;X)$ can also be obtained in O(n) time.

Next we use some results of Section 3 to design a parallel algorithm for computing the power domination numbers of trees.

Theorem 4.2. Let T be a tree, v be a vertex of T with $\deg v \geq 2, V_1, \ldots, V_k$ be the vertex sets of the components of T - v, and $T_i = T[V_i \cup \{v\}]$ for $1 \leq i \leq k$. Partition the indices $1, \ldots, k$ as follows:

$$I = \{i : \gamma_P(T_i; \{v\}) = \gamma_P(T_i) \text{ and } 1 + \gamma_P(T_i - v) \neq \gamma_P(T_i; \{v\})\},$$

$$I' = \{i : \gamma_P(T_i; \{v\}) = \gamma_P(T_i) \text{ and } 1 + \gamma_P(T_i - v) = \gamma_P(T_i; \{v\})\},$$

$$J = \{i : \gamma_P(T_i; \{v\}) \neq \gamma_P(T_i)\}.$$

Let $g = -k + \sum_{i=1}^{k} \gamma_{P}(T_{i}; \{v\})$. Then,

$$\gamma_P(T) = \begin{cases} g+1 & \text{if } |I| \ge 2 \text{ or } |J| = 0\\ g & \text{if } |I| \le 1 \text{ and } |J| \ge 1. \end{cases}$$

Proof. Note that v is a leaf in each T_i . If v is in a power dominating set of T, forcing can occur independently in each T_i after v and its neighbors have been dominated, so $\gamma_P(T; \{v\}) = g+1$. Since $\gamma_P(T) \leq \gamma_P(T; \{v\}) \leq \gamma_P(T) + 1$, it follows that $g \leq \gamma_P(T) \leq g + 1$.

By Lemma 3.8, I is the index set of the subtrees of T for which it is beneficial to include v in a power dominating set, and where v performs a dominating step or force; I' is the index set of the subtrees of T for which it is beneficial to include v in a power dominating set, but where v does not need to perform a dominating step or force; J is the index set of the subtrees of T for which it is not beneficial to include v in a power dominating set.

For $1 \leq i \leq k$, let S_i be a set realizing $\gamma_P(T_i; \{v_i\})$ in T_i . Note that by definition of J, for $i \in J$, S_i can be chosen such that $S_i \setminus \{v\}$ is a set realizing $\gamma_P(T_i)$. By Lemma 3.8, for $i \in I'$, S_i can be chosen such that v does not need to perform a force. Suppose first that $|I| \leq 1$ and $|J| \geq 1$; we claim that $\bigcup_{i=1}^k (S_i \setminus \{v\})$ is a power dominating set of T. To see why, note that for each $i \in J$, the set $S_i \setminus \{v\}$ will force all of $V(T_i)$ in T, including v. Then for $i \in I'$, all components T_i can be forced by the sets $S_i \setminus \{v\}$, $i \in I'$, since v is colored but does not need to perform a force in those components. Finally, if there is a component T_{i^*} , $i^* \in I$, v will have a single uncolored neighbor at this step of the forcing process (which is in T_{i^*}), and it can force this neighbor; since v is a leaf in T_{i^*} , this is the same as dominating its neighbor. Thus, $S_{i^*} \setminus \{v\}$ can power dominate T_{i^*} after all other components are colored. Since $S_i \setminus \{v\}$, $1 \leq i \leq k$ are pairwise disjoint, it follows that

$$\left| \bigcup_{i=1}^{k} (S_i \setminus \{v\}) \right| = \sum_{i=1}^{k} |S_i \setminus \{v\}| = \sum_{i=1}^{k} (\gamma_P(T_i; \{v\}) - 1) = g \ge \gamma_P(T) \ge g.$$

Now suppose $|I| \geq 2$; then, v is beneficial and forcing in at least 2 components, thus it must be contained in some minimum power dominating set of T. Similarly, if |J| = 0 but v is not in a minimum power dominating set of T, then none of the sets $S_i \setminus \{v\}$ can force v. In both cases, it follows that $\gamma_P(T) = g + 1$.

Note that by Theorem 4.1, the parameter g and the index sets I and J in Theorem 4.2 can be determined by computing the ordinary power domination numbers of several smaller trees. If the vertex v is picked in a way that separates T into roughly equally-sized components, Theorem 4.2 gives an efficient way to compute $\gamma_P(T)$ in parallel, since several processors can be used to compute the power domination numbers of the smaller trees independently.

4.2 Integer programming formulations

In this section, we present integer programming models for finding a minimum restricted power dominating set of a graph. Some of the following definitions and models are adapted from [3]. A fort of a graph G = (V, E) is a nonempty set $F \subset V$ such that no vertex outside F is adjacent to exactly one vertex in F.

Proposition 4.3. Let G = (V, E) be a graph and F be any fort of G.

- 1. If S is a power dominating set of G, then $S \cap N[F] \neq \emptyset$.
- 2. If B is a zero forcing set of G, then $S \cap F \neq \emptyset$.

Proof. We prove the first statement; the proof of the second is a simplified version of that proof. Suppose to the contrary that $S \cap N[F] = \emptyset$, and let v be the first vertex in F to be observed by S. Since no neighbor of v is in S, v is not observed in the domination step. Thus, v must be forced by some vertex w not in F. However, by definition of a fort, w is adjacent to at least two vertices in F, so w cannot force v; this is a contradiction. \square

We now present an integer program which can be used to find a restricted power dominating set subject to X. The binary variable p_v indicates whether vertex v is in the restricted power dominating set; S is the set of all forts of the given graph G = (V, E).

Model 4.4. IP model for restricted power domination based on forts

min
$$\sum_{v \in V} p_v$$
s.t.:
$$\sum_{v \in N[F]} p_v \ge 1 \qquad \forall F \in \mathcal{S}$$

$$p_v = 1 \qquad \forall v \in X$$

$$p_v \in \{0, 1\} \qquad \forall v \in V$$

$$(3)$$

Theorem 4.5. The optimum of Model 4.4 is equal to $\gamma_P(G;X)$.

Proof. Let p be an optimal solution of Model 4.4, and let S be the set of all vertices v for which $p_v = 1$. By constraint (3), S contains X. Suppose to the contrary that S is not a power dominating set of G, which means $cl(N[S]) \neq V$. If any vertex $u \in cl(N[S])$ is adjacent to exactly one vertex $v \in V \setminus cl(N[S])$, then u could force v, contradicting the definition of cl(N[S]). Thus, $V \setminus cl(N[S])$ is a fort. Moreover, since $V \setminus cl(N[S])$ does not contain any vertices of N[S], $N[V \setminus cl(N[S])]$ does not contain any vertex of S. This means constraint (2) is violated by the fort $V \setminus cl(N[S])$, a contradiction. It follows that S is a restricted power dominating set of S subject to S, and is minimum due to the objective function.

Next, let S be a minimum restricted power dominating set of G subject to X, and p be the vector whose nonzero entries are indexed by the vertices in S. By Proposition 4.3, for every $F \in \mathcal{S}$, N[F] contains an element of S; thus, p satisfies constraint (2). Moreover, S contains X, so p satisfies constraint (3). Thus, p is a feasible solution of Model 4.4.

By an argument similar to that given in the proof of Theorem 4.5, it follows that if constraint (2) of Model 4.4 is replaced by $\sum_{v \in F} p_v \geq 1 \ \forall F \in \mathcal{S}$, the optimum of the modified integer program will be equal to Z(G;X).

Since a graph (e.g. K_n) could have an exponential number of forts, Model 4.4 must in general be solved using constraint generation. This approach requires a practical method for finding violated constraints; to this end, we present an auxiliary integer program for generating violated fort constraints in Model 4.6. In this model, S is the set of all vertices for which $p_v = 1$ in the current optimal solution of Model 4.4; thus, S is constant in Model 4.6. The binary variable f_v indicates whether vertex v is in the fort.

Model 4.6. IP model for finding violated fort constraints

min
$$\sum_{v \in V} f_v$$
s.t.:
$$\sum_{v \in V} f_v \ge 1$$

$$f_w + \sum_{u \in N(w) \setminus \{v\}} f_u \ge f_v \qquad \forall (v, w) \text{ with } v \in V, w \in N(v)$$

$$f_v = 0 \qquad \forall v \in cl(N[S])$$

$$f_v \in \{0, 1\} \qquad \forall v \in V$$

$$(4)$$

$$f_v = 0 \qquad \forall v \in cl(N[S])$$

$$f_v \in \{0, 1\} \qquad \forall v \in V$$

$$(6)$$

Theorem 4.7. Model 4.6 finds a minimum-size violated fort.

Proof. Let f be a solution of Model 4.6, and let F be the set of vertices of G for which $f_v = 1$. By constraint (4), F is not empty. By constraint (5), every neighbor of a vertex in F must either be in F, or have at least two neighbors in F. Thus, F is a fort of G. Furthermore, by constraint (6), no vertex in F is in cl(N[S]). Therefore F is a violated fort with respect to the current solution S of Model 4.4, and is minimum due to the objective function.

Next, let F be a violated fort of G; let $f_v = 1$ for $v \in F$, and $f_v = 0$ for $v \notin F$. Since a fort is nonempty by definition, F must contain at least one vertex; therefore, constraint (4) is satisfied. Also by definition, any neighbor w of a vertex v in F must either be in F or have at least one other neighbor in F; therefore, constraint (5) is satisfied. Since F is a violated fort, no vertex of F can be in cl(N[S]); therefore, constraint (6) is satisfied. Thus, f is a feasible solution to Model 4.6.

4.3 Graphs with polynomially-many terminals

In this section, we identify certain conditions that allow the restricted zero forcing number of a graph to be computed efficiently, given a zero forcing set of a proper subgraph.

Proposition 4.8. Let X be a minimum zero forcing set of a graph G = (V, E) and let G_T be the graph obtained by adding a vertex v* to G and connecting it to all vertices in a set $T \subset V$.

- 1. If T is a set of terminals of forcing chains associated with X, then X is a zero forcing set of G_T , i.e., $Z(G_T; X) = Z(G)$.
- 2. If X is a zero forcing set of G_T , then at least one vertex in T is a terminal of a forcing chain associated with X in G.

Proof. If T is a set of terminals of forcing chains associated with X for some chronological list of forces, then any force performed in G can also be performed in G_T at the same step, since v^* would not interfere with any forces performed between vertices of G. Then, v^* could be forced by any of its neighbors in G_T in the last step. Thus, X is also a zero forcing set of G_T , so $|X| \geq Z(G_T; X) \geq |X| = Z(G)$.

Now suppose X is a zero forcing set of G_T . Let \mathcal{F} be a chronological list of forces for X in G_T , and let $u \in T$ be the vertex which forces v^* . If v^* does not perform a force, then $\mathcal{F}\setminus (u\to v^*)$ is a valid chronological list of forces for X in G, and u is a terminal of a forcing chain since it does not force any vertex in G. If v^* does perform a force in G_T , say $v^*\to w$, let $\mathcal{F}'\subset \mathcal{F}$ be the subsequence of \mathcal{F} beginning with the first force and ending with the force directly before the force $v^*\to w$. Then the set of vertices in V(G) colored by \mathcal{F}' is also a zero forcing set of G. Let \mathcal{F}'' be a chronological list of forces for this set in G, and note that u does not perform a force in \mathcal{F}'' , since if it did, it could not have forced v^* in G_T ; moreover, some other vertex in V(G) forces w. Then, combining \mathcal{F}' and \mathcal{F}'' , we obtain a chronological list of forces for G associated with X in which u is the terminal of a forcing chain. \square

Note that the necessary conditions in Proposition 4.8 are sometimes sufficient and the sufficient conditions are sometimes necessary. Note also that while the conditions assure that X is a zero forcing set of G_T , they do not assure that X is a minimum zero forcing set of G_T . For example, let G be the star $K_{1,4}$ and T be a set containing two leaves of G; then Z(G) = 3, and any zero forcing set of G is a zero forcing set of G_T , but G_T also has a zero forcing set of size 2.

Proposition 4.9. Let X be a minimum zero forcing set of G, $\{u_1, \ldots, u_k\}$ be a set of terminals of forcing chains associated with X, and H_1, \ldots, H_k be disjoint connected graphs with $V(H_i) \cap V(G) = \{u_i\}$ for $1 \le i \le k$. Let $G' = G \cup H_1 \cup \cdots \cup H_k$. Then

$$Z(G';X) = |X| - k + \sum_{i=1}^{k} Z(H_i; \{u_i\}).$$

Proof. For $1 \leq i \leq k$, let X_i be a minimum zero forcing of H_i containing u_i . Then, the set $X \cup ((X_1 \cup \cdots \cup X_k) \setminus \{u_1, \ldots, u_k\})$ is a zero forcing set of G', since X can force all vertices in V(G) without any of the vertices in $V(G') \setminus V(G)$ interfering in this forcing process, and then each H_i can be forced independently by X_i . Thus, since X and each $X_i \setminus \{u_i\}$ are pairwise disjoint, and since $|X_i \setminus \{u_i\}| = Z(H_i; \{u_i\}) - 1$, it follows that $Z(G'; X) \leq |X| - k + \sum_{i=1}^k Z(H_i; \{u_i\})$.

Now suppose that B' is a minimum zero forcing set of G' that contains X. If B' contains some $u_i \notin X$, then the set $B' \setminus \{u_i\}$ is also a zero forcing set of G', since it contains X and can therefore force u_i in some step of the forcing process. Thus, without loss of generality,

B' excludes all $u_i \notin X$. For i = 1, ..., k, let $X_i = B' \cap V(H_i)$. If $u_i \notin B'$, then $X_i' = X_i \cup \{u_i\}$ is a zero forcing set of H_i , since no vertex other than u_i can be forced by a vertex outside $V(H_i)$. Similarly, for each $u_i \in B'$, $X_i' = X_i$ is a zero forcing set of H_i . Notice that $u_i \in X_i'$ for i = 1, ..., k. Hence, $Z(G'; X) = |B'| \ge |X \cup \bigcup_{i=1}^k (X_i' \setminus \{u_i\})| = |X| + \sum_{i=1}^k (|X_i'| - 1) \ge |X| - k + \sum_{i=1}^k Z(H_i; \{u_i\})$.

In view of Proposition 4.9, given a graph with the structure of G', with subgraph G and a minimum zero forcing set X of G, one could determine whether Z(G';X) is as in the statement of the proposition in polynomial time, if there are polynomially-many sets of vertices which are terminals of forcing chains associated with X (since then it could be checked whether u_1, \ldots, u_k belong to one of these sets). Graphs with this property include paths, cycles, and complete graphs. However, the next result shows that in general, one could not efficiently enumerate all sets which are terminals of forcing chains associated with X.

Proposition 4.10. A graph with a fixed minimum zero forcing set or minimum power dominating set X can have exponentially-many sets of vertices that are terminals of forcing chains associated with X.

Proof. Let G' = (V, E) be the disjoint union of k copies of C_5 , where the ith copy of C_5 has vertex set $\{a_i, b_i, c_i, v_i, u_i\}$ and edge set $\{a_ib_i, b_ic_i, c_iv_i, v_iu_i, u_ia_i\}$. Let $G = (V \cup \{x\}, E \cup \{xv_i : 1 \le i \le k\} \cup \{xu_i : 1 \le i \le k\})$, and $X = \{x\}$. It is easy to see that X is a minimum power dominating set of G. Initially x dominates $\{u_i : 1 \le i \le k\} \cup \{v_i : 1 \le i \le k\}$. In each copy of C_5 , possible forcing chains associated with u_i and v_i include $\{v_i \to c_i \to b_i \to a_i\}$, $\{u_i \to a_i, v_i \to c_i \to b_i\}$, $\{u_i \to a_i \to b_i, v_i \to c_i\}$, and $\{u_i \to a_i \to b_i \to c_i\}$, with sets of terminals $\{u_i, a_i\}$, $\{a_i, b_i\}$, $\{b_i, c_i\}$, and $\{c_i, v_i\}$, respectively. Thus, there are $\Omega(4^k)$ distinct sets of vertices that are terminals of forcing chains associated with X.

Similarly, $X = \{a_1\} \cup \{u_i : 1 \le i \le k\} \cup \{v_i : 3 \le i \le k\}$ is a minimum zero forcing set of G and there are $\Omega(4^{k-1})$ distinct sets of vertices which are terminals of forcing chains associated with X.

Finally, we briefly discuss when the quantity characterized in Proposition 4.10 is polynomial in the order of the graph. Trivially, if Z(G) = O(1) or Z(G) = n - O(1), then there are polynomially-many sets of vertices which are terminals of forcing chains associated with some fixed zero forcing set X. It could also happen that these cardinalities are bounded by a polynomial even when Z(G) does not have an extremal value.

5 Concluding remarks

In this paper, we have studied the restricted power domination and restricted zero forcing problems, which have practical applications in power network monitoring and control of quantum systems. In particular, we have derived tight bounds on the restricted power domination and restricted zero forcing numbers of graphs, and related them to their non-restricted analogues as well as to other graph parameters. We also presented exact and algorithmic results for computing the restricted power domination and restricted zero forcing

numbers of several families of graphs, including graphs with bounded treewidth, and graphs whose subgraphs have zero forcing sets with polynomially many sets of terminals. We used the notion of restricted power domination to develop a parallel algorithm for the power domination number of trees, whereby a tree is partitioned into smaller trees whose power dominating sets can be computed independently.

We also presented integer programming formulations for computing the restricted power domination numbers of arbitrary graphs; these models use a set cover framework with constraint generation, rather than a step-dependent formulation that is sometimes used in the literature (see, e.g., [12] and the bibliography therein).

Future work can focus on deriving other bounds on the restricted power domination and zero forcing numbers, and relating them to other restricted domination parameters in the framework of the work of Sanchis, Goddard, and Henning [15, 23, 19]. The proposed exact algorithms can also be extended to more general graphs. For instance, since the power domination number of block graphs can be computed efficiently [24, 25], it may be possible to extend Theorems 4.1 and 4.2 to block graphs, i.e., to compute the restricted power domination number of a block graph in O(n) time, and to compute the power domination number of a block graph in parallel using restricted power domination. Computational results for general graphs and refinements of the integer programming Models 4.4 and 4.6 can also be pursued.

Acknowledgements

This research began at the American Institute of Mathematics workshop *Zero forcing and its applications* with support from National Science Foundation, DMS-1128242. The work of BB is supported in part by the National Science Foundation, Grant No. 1450681. The authors thank AIM and NSF.

References

- [1] AIM Minimum Rank Special Graphs Work Group (F. Barioli, W. Barrett, S. Butler, S. M. Cioabă, D. Cvetković, S. M. Fallat, C. Godsil, W. Haemers, L. Hogben, R. Mikkelson, S. Narayan, O. Pryporova, I. Sciriha, W. So, D. Stevanović, H. van der Holst, K. Vander Meulen, A. Wangsness). Zero forcing sets and the minimum rank of graphs. Linear Algebra Appl., 428, 1628–1648, 2008.
- [2] K.F. Benson, D. Ferrero, M. Flagg, V. Furst, L. Hogben, V. Vasilevska, B. Wissman. Zero forcing and power domination for graph products. To appear in *Australas. J. Combin.* http://arxiv.org/abs/1510.02421.
- [3] B. Brimkov, C.C. Fast, I.V. Hicks. Computational approaches for zero forcing and related problems. https://arxiv.org/abs/1704.02065.
- [4] B. Brimkov, C.C. Fast, I.V. Hicks. Graphs with extremal connected forcing numbers. http://arxiv.org/abs/1701.08500.

- [5] D.J. Brueni, L.S. Heath. The PMU placement problem. SIAM J. Discrete Math., 19, 744–761, 2005.
- [6] D. Burgarth, V. Giovannetti. Full control by locally induced relaxation. *Phys. Rev. Lett.* PRL 99, 100501, 2007.
- [7] S. Butler, L. DeLoss, J. Grout, H.T. Hall, J. LaGrange, J.C.-H.Lin, T. McKay, J. Smith, G. Tims. Minimum Rank Library (*Sage* programs for calculating bounds on the minimum rank of a graph, and for computing zero forcing parameters). Available at https://sage.math.iastate.edu/home/pub/84/. For more information contact Jephian C.-H. Lin at jephianlin@gmail.com.
- [8] G.J. Chang, P. Dorbec, M. Montassier, A. Raspaud. Generalized power domination of graphs. *Discrete Appl. Math.*, 160: 1691–1698, 2012.
- [9] G.J. Chang, N. Roussel. On the k-power domination of hypergraphs. J. Combin. Optim., 30(4), 1095-1106, 2015.
- [10] N. Dean, A. Ilic, I. Ramirez, J. Shen, K. Tian. On the power dominating sets of hypercubes. IEEE 14th International Conference on Computational Science and Engineering (CSE), 488 - 491, 2011.
- [11] C.J. Edholm, L. Hogben, M. Huynh, J. LaGrange, D.D. Row. Vertex and edge spread of zero forcing number, maximum nullity, and minimum rank of a graph. *Linear Algebra Appl.*, 436, 4352-4372, 2012.
- [12] N. Fan, J.P. Watson. Solving the connected dominating set problem and power dominating set problem by integer programming. *Combin. Optim. Appl.*, 371–383, 2012.
- [13] D. Ferrero, L. Hogben, F.H.J. Kenter, M. Young. Note on power propagation time and lower bounds for the power domination number. *J. Combin. Optim.*, 34(3), 736–741, 2017.
- [14] D. Ferrero, L. Hogben, F.H.J. Kenter, M. Young. The relationship between k-forcing and k-power domination. To appear in *Discrete Math.* http://arxiv.org/abs/1701.08386.
- [15] W. Goddard, M. A. Henning. Restricted domination parameters in graphs. *J. Combin. Optim.*, 13(4), 353-363, 2007.
- [16] J. Guo, R. Niedermeier, D. Raible. Improved algorithms and complexity results for power domination in graphs. *Algorithmica*, 52, 177–202, 2008.
- [17] T.W. Haynes, S.M. Hedetniemi, S.T. Hedetniemi, M.A. Henning. Domination in graphs applied to electric power networks. *SIAM J. Discrete Math.*, 15, 519–529, 2002.
- [18] T.W. Haynes, S.M. Hedetniemi, P.J. Slater. Fundamentals of domination in graphs applied to electric power networks. Marcel Dekker, New York, 1998.

- [19] M. A. Henning. Restricted domination in graphs. Discrete Math., 254, 175–189, 2002.
- [20] L. Hogben, N. Kingsley, S. Meyer, S. Walker, M. Young. Propagation time for zero forcing on a graph. Discrete Appl. Math., 160(13), 1994–2005, 2012.
- [21] C.S. Liao. Power domination with bounded time constraints. *J. Combin. Optim.*, 31(2), 725–742, 2016.
- [22] D. Row. A technique for computing the zero forcing number of a graph with a cut-vertex. Linear Algebra Appl. 436, 4423–4432, 2012.
- [23] L.A. Sanchis. Bounds related to domination in graphs with minimum degree two. *J. Graph Theory*, 25, 139–152, 1997.
- [24] C. Wang, L. Chen, C. Lu. k-Power domination in block graphs. J. Combin. Optim. 31(2), 865-873, 2016.
- [25] G. Xu, L. Kang, E. Shan, M. Zhao. Power domination in block graphs. Theoret. Comput. Sci. 359, 299–305, 2006.
- [26] B. Yang. Lower bounds for positive semidefinite zero forcing and their applications. *J. Combin. Optim.*, 33(1), 81–105, 2017.
- [27] M. Zhao, L. Kang, G.J. Chang. Power domination in graphs. Discrete Math. 306, 1812– 1816, 2006.