

Tight bounds on the maximal perimeter and the maximal width of convex small polygons

Christian Bingane*

June 3, 2022

Abstract

A small polygon is a polygon of unit diameter. The maximal perimeter and the maximal width of a convex small polygon with $n = 2^s$ vertices are not known when $s \geq 4$. In this paper, we construct a family of convex small n -gons, $n = 2^s$ and $s \geq 3$, and show that the perimeters and the widths obtained cannot be improved for large n by more than a/n^6 and b/n^4 respectively, for certain positive constants a and b . In addition, assuming that a conjecture of Mossinghoff is true, we formulate the maximal perimeter problem as a nonlinear optimization problem involving trigonometric functions and, for $n = 2^s$ with $3 \leq s \leq 7$, we provide global optimal solutions.

Keywords Planar geometry, polygons, isodiametric problems, maximal perimeter, maximal width, global optimization

1 Introduction

The *diameter* of a polygon is the largest Euclidean distance between pairs of its vertices. A polygon is said to be *small* if its diameter equals one. For a given integer $n \geq 3$, the maximal perimeter problem consists in finding a convex small n -gon with the longest perimeter. The problem was first investigated by Reinhardt [1] in 1922, and later by Datta [2] in 1997. They proved that

- for all $n \geq 3$, the value $2n \sin \frac{\pi}{2n}$ is an upper bound on the perimeter of a convex small n -gon;
- when n is odd, the regular small n -gon is an optimal solution, but it is unique only if n is prime;
- when n is even, the regular small n -gon is not optimal;
- when n has an odd factor, there are finitely many optimal solutions [3–5] and they are all equilateral.

When n is a power of 2, the maximal perimeter problem is solved for $n \leq 8$. In 1987, Tamvakis [6] found the unique convex small 4-gon with the longest perimeter, shown in Figure 1b. In 2007, Audet, Hansen, and Messine [7] used both geometrical arguments and methods of global optimization to determine the unique convex small 8-gon with the longest perimeter, illustrated in Figure 3c.

The diameter graph of a small polygon is the graph with the vertices of the polygon, and an edge between two vertices exists only if the distance between these vertices equals one. Figure 1, Figure 2, and Figure 3 represent diameter graphs of some convex small polygons. The solid lines illustrate pairs

*Département de mathématiques et de génie industriel, Polytechnique Montréal, Montreal, QC, Canada. Email: christian.bingane@polymtl.ca




Figure 1: Two convex small 4-gons $(P_4, L(P_4)), W(P_4))$




Figure 2: Three convex small 6-gons $(P_6, L(P_6)), W(P_6))$

of vertices which are unit distance apart. Mossinghoff [8] conjectured that, for $n \geq 4$ power of 2, the diameter graph of a convex small n -gon with maximal perimeter has a cycle of length $n/2 + 1$, plus $n/2 - 1$ additional pendant edges, arranged so that all but two particular vertices of the cycle have a pendant edge. For example, Figure 1b and Figure 3c exhibit the diameter graphs of optimal n -gons when $n = 4$ and when $n = 8$ respectively. We point out that numerical values in figures and tables in this paper are rounded at the last reported digit.

The *width* of a polygon in some direction is the distance between two parallel lines perpendicular to this direction and supporting the polygon from below and above. The width of a polygon is the minimum width for all directions. For a given integer $n \geq 3$, the maximal width problem consists in finding a convex small n -gon with the largest width. This problem was partially solved by Bezdek and Fodor [9] in 2000. They proved that

- for all $n \geq 3$, the value $\cos \frac{\pi}{2n}$ is an upper bound on the width of a convex small n -gon;




Figure 3: Four convex small 8-gons $(P_8, L(P_8)), W(P_8))$

- when n has an odd factor, a convex small n -gon is optimal for the maximal width problem if and only if it is optimal for the maximal perimeter problem;
- when $n = 4$, there are infinitely many optimal convex small 4-gons, including the 4-gon illustrated in Figure 1b.

When $n \geq 8$ is a power of 2, the maximal width is only known for the first open case $n = 8$. In 2013, Audet, Hansen, Messine, and Ninin [10] combined geometrical and analytical reasoning as well as methods of global optimization to prove that there are infinitely many optimal convex small 8-gons, including the 8-gon illustrated in Figure 3d.

For $n = 2^s$ with integer $s \geq 4$, exact solutions in both problems appear to be presently out of reach. However, tight lower bounds on the maximal perimeter and the maximal width may be obtained analytically. For instance, Mossinghoff [8] constructed convex small n -gons, for $n = 2^s$ with $s \geq 3$, and proved that the perimeters obtained cannot be improved for large n by more than $\pi^5/(16n^5)$. We can also show that, when $n = 2^s$ with $s \geq 2$, the value $\cos \frac{\pi}{2n-2}$ is a lower bound on the maximal width and this bound cannot be improved for large n by more than $\pi^2/(4n^3)$. In this paper, we propose tighter lower bounds on both the maximal perimeter and the maximal width of convex small n -gons when $n = 2^s$ and integer $s \geq 3$. Thus, the main result of this paper is the following:

Theorem 1. *For a given integer $n \geq 3$, let $\bar{L}_n := 2n \sin \frac{\pi}{2n}$ denote an upper bound on the perimeter $L(P_n)$ of a convex small n -gon P_n , and $\bar{W}_n := \cos \frac{\pi}{2n}$ denote an upper bound on its width $W(P_n)$. If $n = 2^s$ with $s \geq 3$, then there exists a convex small n -gon B_n such that*

$$\begin{aligned} L(B_n) &= 2n \sin \frac{\pi}{2n} \cos \left(\frac{\pi}{2n} - \frac{1}{2} \arcsin \left(\frac{1}{2} \sin \frac{2\pi}{n} \right) \right), \\ W(B_n) &= \cos \left(\frac{\pi}{n} - \frac{1}{2} \arcsin \left(\frac{1}{2} \sin \frac{2\pi}{n} \right) \right), \end{aligned}$$

and

$$\begin{aligned} \bar{L}_n - L(B_n) &= \frac{\pi^7}{32n^6} + O\left(\frac{1}{n^8}\right), \\ \bar{W}_n - W(B_n) &= \frac{\pi^4}{8n^4} + O\left(\frac{1}{n^6}\right). \end{aligned}$$

The remainder of this paper is organized as follows. Section 2 recalls principal results on the maximal perimeter and the maximal width of convex small polygons. We prove Theorem 1 in Section 3. Tight bounds on the maximal width of unit-perimeter n -gons, $n = 2^s$ and $s \geq 3$, are deduced from Theorem 1 in Section 4. Under the assumption that Mossinghoff's conjecture is true, a nonlinear optimization problem involving trigonometric functions is proposed for the maximal perimeter problem in Section 5. Global optimal solutions obtained by using AMPL with the solver Couenne [11] are given for $n = 2^s$ with $3 \leq s \leq 7$. Section 6 concludes the paper.

2 Perimeters and widths of convex small polygons

2.1 Maximal perimeter and maximal width

Let $L(P)$ denote the perimeter of a polygon P and $W(P)$ its width. For a given integer $n \geq 3$, let R_n denote the regular small n -gon. We have

$$L(R_n) = \begin{cases} 2n \sin \frac{\pi}{2n} & \text{if } n \text{ is odd,} \\ n \sin \frac{\pi}{n} & \text{if } n \text{ is even,} \end{cases}$$

and

$$W(R_n) = \begin{cases} \cos \frac{\pi}{2n} & \text{if } n \text{ is odd,} \\ \cos \frac{\pi}{n} & \text{if } n \text{ is even.} \end{cases}$$

We remark that $L(R_n) < L(R_{n-1})$ [12] and $W(R_n) < W(R_{n-1})$ for all even $n \geq 4$. The polygon R_n does not have maximum perimeter nor maximum width for any even $n \geq 4$. Indeed, when n is even, one can construct a convex small n -gon with a longer perimeter and a larger width than R_n by adding a vertex at distance 1 along the mediatix of an angle in R_{n-1} . We denote this n -gon by R_{n-1}^+ and we have

$$\begin{aligned} L(R_{n-1}^+) &= (2n-2) \sin \frac{\pi}{2n-2} + 4 \sin \frac{\pi}{4n-4} - 2 \sin \frac{\pi}{2n-2}, \\ W(R_{n-1}^+) &= \cos \frac{\pi}{2n-2}. \end{aligned}$$

When n has an odd factor m , we construct another family of convex equilateral small n -gons as follows:

1. Consider a regular small m -gon R_m ;
2. Transform R_m into a Reuleaux m -gon by replacing each edge by a circle's arc passing through its end vertices and centered at the opposite vertex;
3. Add at regular intervals $n/m - 1$ vertices within each arc;
4. Take the convex hull of all vertices.

We denote these n -gons by $R_{m,n}$ and we have

$$\begin{aligned} L(R_{m,n}) &= 2n \sin \frac{\pi}{2n}, \\ W(R_{m,n}) &= \cos \frac{\pi}{2n}. \end{aligned}$$

The 6-gon $R_{3,6}$ is illustrated in Figure 2c.

Theorem 2 (Reinhardt [1], Datta [2]). *For all $n \geq 3$, let L_n^* denote the maximal perimeter among all convex small n -gons and let $\bar{L}_n := 2n \sin \frac{\pi}{2n}$.*

- When n has an odd factor m , $L_n^* = \bar{L}_n$ is achieved by finitely many equilateral n -gons [3–5], including $R_{m,n}$. The optimal n -gon $R_{m,n}$ is unique if m is prime and $n/m \leq 2$.
- When $n = 2^s$ with integer $s \geq 2$, $L(R_n) < L_n^* < \bar{L}_n$.

When $n = 2^s$, the maximal perimeter L_n^* is only known for $s \leq 3$. Tamvakis [6] found that $L_4^* = 2 + \sqrt{6} - \sqrt{2}$, and this value is achieved only by R_3^+ , shown in Figure 1b. Audet, Hansen, and Messine [7] found that $L_8^* \approx 3.121147$, and this value is only achieved by B_8^* , shown in Figure 3c.

Theorem 3 (Bezdek and Fodor [9]). *For all $n \geq 3$, let W_n^* denote the maximal width among all convex small n -gons and let $\bar{W}_n := \cos \frac{\pi}{2n}$.*

- When n has an odd factor, $W_n^* = \bar{W}_n$ is achieved by a convex small n -gon with maximal perimeter $L_n^* = \bar{L}_n$.
- When $n = 2^s$ with integer $s \geq 2$, $W(R_n) < W_n^* < \bar{W}_n$.

When $n = 2^s$, the maximal width W_n^* is only known for $s \leq 3$. Bezdek and Fodor [9] showed that $W_4^* = \frac{1}{2}\sqrt{3}$, and this value is achieved by infinitely many convex small 4-gons, including R_3^+ shown in Figure 1b. Audet, Hansen, Messine, and Ninin found that $W_8^* = \frac{1}{4}\sqrt{10 + 2\sqrt{7}}$, and this value is also achieved by infinitely many convex small 8-gons, including B_8 shown in Figure 3d. It is interesting to note that while the optimal 4-gon for the maximal perimeter problem is also optimal for the maximal width problem, the optimal 8-gon for the maximal perimeter problem is not optimal for the maximal width problem.




Figure 4: Tamvakis polygons $(T_n, L(T_n), W(T_n))$

2.2 Lower bounds on the maximal perimeter and the maximal width

For $n = 2^s$ with integer $s \geq 2$, let T_n denote the convex n -gon obtained by subdividing each bounding arc of a such Reuleaux triangle into either $\lceil n/3 \rceil$ or $\lfloor n/3 \rfloor$ subarcs of equal length, then taking the convex hull of the endpoints of these arcs. For a real number a , $\lceil a \rceil$ is the least integer greater than or equal to a , and $\lfloor a \rfloor$ is the greatest integer less than or equal to a . We illustrate T_n for some n in Figure 4. For each n , the perimeter of T_n is given by

$$L(T_n) = \begin{cases} \frac{4n-4}{3} \sin \frac{\pi}{2n-2} + \frac{2n+4}{3} \sin \frac{\pi}{2n+4} & \text{if } n = 3k + 1, \\ \frac{4n+4}{3} \sin \frac{\pi}{2n+2} + \frac{2n-4}{3} \sin \frac{\pi}{2n-4} & \text{if } n = 3k + 2. \end{cases}$$

We note that T_4 is optimal for the maximal perimeter problem and we can show that

$$\overline{L}_n - L(T_n) = \frac{\pi^3}{4n^4} + O\left(\frac{1}{n^5}\right)$$

for all $n = 2^s$ and $s \geq 2$. By contrast,

$$\begin{aligned} \overline{L}_n - L(R_n) &= \frac{\pi^3}{8n^2} + O\left(\frac{1}{n^4}\right), \\ \overline{L}_n - L(R_{n-1}^+) &= \frac{5\pi^3}{96n^3} + O\left(\frac{1}{n^4}\right) \end{aligned}$$

for all even $n \geq 4$. Tamvakis asked if T_n is also optimal when $s \geq 3$. Obviously, T_8 is not optimal, i.e., $L(T_8) < L_8^*$.

For all $n = 2^s$ with integer $s \geq 2$, let P_n^* denote a convex small n -gon with the longest perimeter.

Conjecture 1 (Mossinghoff [8]). For all $n = 2^s$ with integer $s \geq 2$, the diameter graph of P_n^* has a cycle of length $n/2 + 1$, plus $n/2 - 1$ additional pendant edges, arranged so that all but two particular vertices of the cycle have a pendant edge.

Conjecture 2 (Mossinghoff [8]). For all $n = 2^s$ with integer $s \geq 2$, P_n^* has an axis of symmetry corresponding to one particular pendant edge in its diameter graph.

Conjecture 1 is proven for $n = 4$ [6] and $n = 8$ [7]. Conjecture 2 is only proven for $n = 4$ [6], but it is shown numerically for $n = 8$ in [7]. Mossinghoff [8] constructed a family of convex small n -gons M_n having the diameter graph described in Conjectures 1 and 2. These polygons have the property that

$$\overline{L}_n - L(M_n) = \frac{\pi^5}{16n^5} + O\left(\frac{1}{n^6}\right)$$




Figure 5: Mossinghoff polygons $(M_n, L(M_n), W(M_n))$

when $n = 2^s$ and $s \geq 3$. We show M_n for some n in Figure 5.

On the other hand, for all $n = 2^s$ and integer $s \geq 3$,

$$W(T_n) = \begin{cases} \cos \frac{\pi}{2n-2} & \text{if } n = 3k+1, \\ \cos \frac{\pi}{2n-4} & \text{if } n = 3k+2, \end{cases}$$

$$W(M_n) = \cos \left(\frac{\pi}{2n} + \frac{\pi^2}{4n^2} - \frac{\pi^2}{2n^3} \right),$$

and we can show that $W(R_{n-1}^+) \geq \max\{W(T_n), W(M_n)\}$. Note that

$$\overline{W}_n - W(R_n) = \frac{3\pi^2}{8n^2} + O\left(\frac{1}{n^4}\right),$$

$$\overline{W}_n - W(R_{n-1}^+) = \frac{\pi^2}{4n^3} + O\left(\frac{1}{n^4}\right)$$

for all even $n \geq 4$.

3 Proof of Theorem 1

We use cartesian coordinates to describe an n -gon P_n , assuming that a vertex v_i , $i = 0, 1, \dots, n-1$, is positioned at abscissa x_i and ordinate y_i . Placing the vertex v_0 at the origin, we set $x_0 = y_0 = 0$. We also assume that the n -gon P_n is in the half-plane $y \geq 0$.

For all $n = 2^s$ with integer $s \geq 3$, consider the n -gon P_n having an $(n/2 + 1)$ -length cycle: $v_0 - v_1 - \dots - v_k - \dots - v_{\frac{n}{4}} - v_{\frac{n}{4}+1} - \dots - v_{\frac{n}{2}-k+1} - \dots - v_{\frac{n}{2}} - v_0$ plus $n/2 - 1$ pendant edges: $v_0 - v_{\frac{n}{2}+1}$, $v_k - v_{k+\frac{n}{2}+1}$, $v_{\frac{n}{2}-k+1} - v_{n-k}$, $k = 1, \dots, n/4 - 1$, as illustrated in Figure 6. We assume that P_n has the edge $v_0 - v_{\frac{n}{2}+1}$ as axis of symmetry and for all $k = 1, \dots, n/4 - 1$, the pendant edge $v_k - v_{k+\frac{n}{2}+1}$ bisects the angle $\angle v_{k-1}v_kv_{k+1}$.

Let $\alpha_0 := \angle v_{\frac{n}{2}+1}v_0v_1$, $2\alpha_k := \angle v_{k-1}v_kv_{k+1}$ for all $k = 1, \dots, n/4 - 1$, and $\alpha_{\frac{n}{4}} := \angle v_{\frac{n}{4}-1}v_{\frac{n}{4}}v_{\frac{n}{4}+1}$. Since P_n is symmetric, we have

$$\alpha_0 + 2 \sum_{k=1}^{n/4-1} \alpha_k + \alpha_{n/4} = \frac{\pi}{2}, \quad (1)$$

and

$$L(P_n) = 4 \sin \frac{\alpha_0}{2} + 8 \sum_{k=1}^{n/4-1} \sin \frac{\alpha_k}{2} + 4 \sin \frac{\alpha_{n/4}}{2}, \quad (2a)$$

$$W(P_n) = \min_{k=0,1,\dots,n/4} \cos \frac{\alpha_k}{2}. \quad (2b)$$




Figure 6: Definition of variables $\alpha_0, \alpha_1, \dots, \alpha_{\frac{n}{4}}$ for B_n : Case of $n = 8$ vertices

By placing the vertex v_0 at $(0, 0)$ in the plane, and the vertex $v_{\frac{n}{2}+1}$ at $(0, 1)$, we have

$$x_1 = \sin \alpha_0 = -x_{\frac{n}{2}}, \quad (3a)$$

$$y_1 = \cos \alpha_0 = y_{\frac{n}{2}}, \quad (3b)$$

$$x_k = x_{k-1} - (-1)^k \sin \left(\alpha_0 + 2 \sum_{j=1}^{k-1} \alpha_j \right) = -x_{\frac{n}{2}-k+1} \quad \forall k = 2, 3, \dots, n/4 \quad (3c)$$

$$y_k = y_{k-1} - (-1)^k \cos \left(\alpha_0 + 2 \sum_{j=1}^{k-1} \alpha_j \right) = y_{\frac{n}{2}-k+1} \quad \forall k = 2, 3, \dots, n/4, \quad (3d)$$

$$x_{k+\frac{n}{2}+1} = x_k + (-1)^k \sin \left(\alpha_0 + 2 \sum_{j=1}^{k-1} \alpha_j + \alpha_k \right) = -x_{n-k} \quad \forall k = 1, 2, \dots, n/4 - 1, \quad (3e)$$

$$y_{k+\frac{n}{2}+1} = y_k + (-1)^k \cos \left(\alpha_0 + 2 \sum_{j=1}^{k-1} \alpha_j + \alpha_k \right) = y_{n-k} \quad \forall k = 1, 2, \dots, n/4 - 1. \quad (3f)$$

We also have

$$x_{\frac{n}{4}} = -1/2 = -x_{\frac{n}{4}+1}. \quad (4)$$

since the edge $v_{\frac{n}{4}} - v_{\frac{n}{4}+1}$ is horizontal and $\|v_{\frac{n}{4}} - v_{\frac{n}{4}+1}\| = 1$.

For all $k = 0, 1, \dots, n/4$, suppose $\alpha_k = \frac{\pi}{n} + (-1)^k \beta$ with $\beta = \beta(n)$ satisfying $|\beta| < \frac{\pi}{n}$. Then (1) is verified and (2) becomes

$$L(P_n) = n \sin \left(\frac{\pi}{2n} + \frac{\beta}{2} \right) + n \sin \left(\frac{\pi}{2n} - \frac{\beta}{2} \right) = 2n \sin \frac{\pi}{2n} \cos \frac{\beta}{2}, \quad (5a)$$

$$W(P_n) = \cos \left(\frac{\pi}{2n} + \frac{|\beta|}{2} \right). \quad (5b)$$




Figure 7: Polygons $(B_n, L(B_n), W(B_n))$ defined in Theorem 1

Coordinates (x_i, y_i) in (3) are given by

$$\begin{aligned} x_k &= \sum_{j=1}^k (-1)^{j-1} \sin \left((2j-1) \frac{\pi}{n} + (-1)^{j-1} \beta \right) \\ &= \frac{\sin \frac{2k\pi}{n} \sin \left(\beta - (-1)^k \frac{\pi}{n} \right)}{\sin \frac{2\pi}{n}} = -x_{\frac{n}{2}-k+1} \quad \forall k = 1, 2, \dots, n/4, \quad (6a) \end{aligned}$$

$$\begin{aligned} y_k &= \sum_{j=1}^k (-1)^{j-1} \cos \left((2j-1) \frac{\pi}{n} + (-1)^{j-1} \beta \right) \\ &= \frac{\sin \left(\frac{\pi}{n} - \beta \right) + \cos \frac{2k\pi}{n} \sin \left(\beta - (-1)^k \frac{\pi}{n} \right)}{\sin \frac{2\pi}{n}} = y_{\frac{n}{2}-k+1} \quad \forall k = 1, 2, \dots, n/4, \quad (6b) \end{aligned}$$

$$x_{k+\frac{n}{2}+1} = x_k + (-1)^k \sin \frac{2k\pi}{n} = -x_{n-k} \quad \forall k = 1, 2, \dots, n/4 - 1, \quad (6c)$$

$$y_{k+\frac{n}{2}+1} = y_k + (-1)^k \cos \frac{2k\pi}{n} = y_{n-k} \quad \forall k = 1, 2, \dots, n/4 - 1. \quad (6d)$$

Finally, β is chosen so that (4) is satisfied. It follows, from (6a),

$$\frac{\sin \left(\beta - \frac{\pi}{n} \right)}{\sin \frac{2\pi}{n}} = -\frac{1}{2} \Rightarrow \beta = \beta_0(n) = \frac{\pi}{n} - \arcsin \left(\frac{1}{2} \sin \frac{2\pi}{n} \right) = \frac{\pi^3}{2n^3} + \frac{\pi^5}{8n^5} + O \left(\frac{1}{n^7} \right).$$

Let B_n denote the n -gon obtained by setting $\beta = \beta_0(n)$. From (5), we have

$$\begin{aligned} L(B_n) &= 2n \sin \frac{\pi}{2n} \cos \left(\frac{\pi}{2n} - \frac{1}{2} \arcsin \left(\frac{1}{2} \sin \frac{2\pi}{n} \right) \right), \\ W(B_n) &= \cos \left(\frac{\pi}{n} - \frac{1}{2} \arcsin \left(\frac{1}{2} \sin \frac{2\pi}{n} \right) \right), \end{aligned}$$

and

$$\begin{aligned} \overline{L}_n - L(B_n) &= \frac{\pi^7}{32n^6} + \frac{11\pi^9}{768n^8} + O \left(\frac{1}{n^{10}} \right), \\ \overline{W}_n - W(B_n) &= \frac{\pi^4}{8n^4} + \frac{11\pi^6}{192n^6} + O \left(\frac{1}{n^8} \right). \end{aligned}$$

By construction, B_n is small and convex for all $n = 2^s$ and $s \geq 3$. We illustrate B_n for some n in Figure 7. This completes the proof of Theorem 1. \square

We implemented all polygons presented in this work as a MATLAB package: OPTIGON, freely available on GitHub [13]. In OPTIGON, MATLAB functions that give the coordinates of the vertices

Table 1: Perimeters of B_n

n	$L(R_n)$	$L(R_{n-1}^+)$	$L(T_n)$	$L(M_n)$	$L(B_n)$	\bar{L}_n	$\frac{L(B_n) - L(M_n)}{\bar{L}_n - L(M_n)}$
8	3.0614674589	3.1181091119	3.1190543124	3.1209757852	3.1210621230	3.1214451523	0.1839
16	3.1214451523	3.1361407965	3.1364381783	3.1365320240	3.1365427675	3.1365484905	0.6524
32	3.1365484905	3.1402809876	3.1403234211	3.1403306141	3.1403310687	3.1403311570	0.8374
64	3.1403311570	3.1412710339	3.1412767980	3.1412772335	3.1412772496	3.1412772509	0.9211
128	3.1412772509	3.1415130275	3.1415137720	3.1415138006	3.141513801123	3.141513801144	0.9606

Table 2: Widths of B_n

n	$W(R_n)$	$W(R_{n-1}^+)$	$W(B_n)$	\bar{W}_n	$\frac{W(B_n) - W(R_{n-1}^+)}{\bar{W}_n - W(R_{n-1}^+)}$
8	0.9238795325	0.9749279122	0.9776087734	0.9807852804	0.4577
16	0.9807852804	0.9945218954	0.9949956687	0.9951847267	0.7148
32	0.9951847267	0.9987165072	0.9987837929	0.9987954562	0.8523
64	0.9987954562	0.9996891820	0.9996980921	0.9996988187	0.9246
128	0.9996988187	0.9999235114	0.9999246565	0.9999247018	0.9619

are provided. An algorithm developed in [14] to estimate the maximal area of a small n -gon [1] when $n \geq 6$ is even can be also found.

Table 1 shows the perimeters of B_n , along with the upper bounds \bar{L}_n , the perimeters of R_n , R_{n-1}^+ , T_n , and M_n for $n = 2^s$ and $3 \leq s \leq 7$. As suggested by Theorem 1, when n is a power of 2, B_n provides a tighter lower bound on the maximal perimeter L_n^* compared to the best prior convex small n -gon M_n . For instance, we can note that $L_{128}^* - L(B_{128}) < \bar{L}_{128} - L(B_{128}) < 2.15 \times 10^{-11}$. By analysing the fraction $\frac{L(B_n) - L(M_n)}{\bar{L}_n - L(M_n)}$ of the length of the interval $[L(M_n), \bar{L}_n]$ where $L(B_n)$ lies, it is not surprising that $L(B_n)$ approaches \bar{L}_n much faster than $L(M_n)$ does as n increases. After all, $L(B_n) - L(M_n) \sim \frac{\pi^5}{16n^5}$ for large n .

Table 2 displays the widths of B_n , along with the upper bounds \bar{W}_n , the widths of R_n and R_{n-1}^+ . Again, when $n = 2^s$, B_n provides a tighter lower bound for the maximal width W_n^* compared to the best prior convex small n -gon R_{n-1}^+ . We also remark that $W(B_n)$ approaches \bar{W}_n much faster than $W(R_{n-1}^+)$ does as n increases. It is interesting to note that $W(B_8) = W_8^*$, i.e., B_8 is an optimal solution for the maximal width problem when $n = 8$.

Proposition 1 and Proposition 2 highlight some interesting properties of B_n .

Proposition 1. *Let $n = 2^s$ with integer $s \geq 3$.*

1. *The coordinates of the vertex $v_{\frac{n}{4}}$ in B_n are $(-1/2, 1/2)$.*
2. *For all $k = 1, \dots, n/4 - 1$, the pendant edge $v_k - v_{k+\frac{n}{2}+1}$ of B_n passes through the point $u = (0, 1/2)$.*

Proof. Let $n = 2^s$ with integer $s \geq 3$ and $\beta = \frac{\pi}{n} - \arcsin\left(\frac{1}{2} \sin \frac{2\pi}{n}\right)$.

1. We have, from (6a),

$$x_{\frac{n}{4}} = \frac{\sin\left(\beta - \frac{\pi}{n}\right)}{\sin \frac{2\pi}{n}} = -\frac{1}{2},$$

$$y_{\frac{n}{4}} = \frac{\sin\left(\frac{\pi}{n} - \beta\right)}{\sin \frac{2\pi}{n}} = \frac{1}{2}.$$

2. For all $k = 1, \dots, n/4 - 1$, coordinates (x_i, y_i) in (6) are

$$\begin{aligned} x_k &= \frac{\sin \frac{2k\pi}{n} \sin \left(\beta - (-1)^k \frac{k\pi}{n} \right)}{\sin \frac{2\pi}{n}}, & x_{k+\frac{n}{2}+1} &= x_k + (-1)^k \sin \frac{2k\pi}{n}, \\ y_k &= \frac{1}{2} + \frac{\cos \frac{2k\pi}{n} \sin \left(\beta - (-1)^k \frac{k\pi}{n} \right)}{\sin \frac{2\pi}{n}}, & y_{k+\frac{n}{2}+1} &= y_k + (-1)^k \cos \frac{2k\pi}{n}. \end{aligned}$$

It follows that, for all $k = 1, \dots, n/4 - 1$,

$$\frac{x_{k+\frac{n}{2}+1} - x_k}{y_{k+\frac{n}{2}+1} - y_k} = \tan \frac{2k\pi}{n} = \frac{x_k}{y_k - \frac{1}{2}},$$

i.e., the pendant edge $v_k - v_{k+\frac{n}{2}+1}$ passes through the point $u = (0, 1/2)$.

□

Proposition 2. Let $n = 2^s$ with integer $s \geq 3$. The area of B_n is $\frac{n}{8} \sin \frac{2\pi}{n}$, which is the area of the regular small n -gon R_n .

Proof. Let $n = 2^s$ with integer $s \geq 3$ and $\beta = \frac{\pi}{n} - \arcsin \left(\frac{1}{2} \sin \frac{2\pi}{n} \right)$. Let A_0 be the area of the quadrilateral $uv_1v_{\frac{n}{2}+1}v_{\frac{n}{2}}$, A_k be the area of the quadrilateral $uv_{k+1}v_{k+\frac{n}{2}+1}v_{k-1}$ for all $k = 1, \dots, n/4 - 1$, and $A_{\frac{n}{4}}$ be the area of the triangle $uv_{\frac{n}{4}+1}v_{\frac{n}{4}-1}$, where $u = (0, 1/2)$. The area of B_n is given by

$$A(B_n) = A_0 + 2 \sum_{k=1}^{n/4-1} A_k + 2A_{\frac{n}{4}}.$$

We have

$$\begin{aligned} A_0 &= \frac{1}{2} \|v_{\frac{n}{2}+1} - u\| \|v_{\frac{n}{2}} - v_1\| = \frac{1}{2} \sin \left(\frac{\pi}{n} + \beta \right), \\ A_k &= \frac{1}{2} \|v_{k+\frac{n}{2}+1} - u\| \|v_{k-1} - v_{k+1}\| \\ &= \begin{cases} \frac{1}{2} \sin \left(\frac{\pi}{n} + \beta \right) & \text{if } k \text{ is even,} \\ \frac{1}{2} \sin \frac{2\pi}{n} - \frac{1}{2} \sin \left(\frac{\pi}{n} + \beta \right) & \text{if } k \text{ is odd,} \end{cases} \end{aligned}$$

for all $k = 1, \dots, n/4 - 1$, and

$$A_{\frac{n}{4}} = \frac{1}{2} (x_{\frac{n}{4}+1}(y_{\frac{n}{4}-1} - 1/2) - (y_{\frac{n}{4}+1} - 1/2)x_{\frac{n}{4}-1}) = \frac{1}{4} \sin \left(\frac{\pi}{n} + \beta \right).$$

Thus,

$$A(B_n) = \frac{n}{8} \sin \left(\frac{\pi}{n} + \beta \right) + \frac{n}{8} \left(\sin \frac{2\pi}{n} - \sin \left(\frac{\pi}{n} + \beta \right) \right) = \frac{n}{8} \sin \frac{2\pi}{n}.$$

□

4 Tight bounds on the maximal width of unit-perimeter polygons

Let \hat{P} denote the polygon obtained by contracting a small polygon P so that $L(\hat{P}) = 1$. Thus, the width of the unit-perimeter polygon \hat{P} is given by $W(\hat{P}) = W(P)/L(P)$. For a given integer $n \geq 3$,

$$W(\hat{R}_n) = \begin{cases} \frac{1}{2n} \cot \frac{\pi}{2n} & \text{if } n \text{ is odd,} \\ \frac{1}{n} \cot \frac{\pi}{n} & \text{if } n \text{ is even.} \end{cases}$$

Table 3: Widths of \hat{B}_n

n	$W(\hat{R}_n)$	\bar{w}_{n-1}	$W(\hat{B}_n)$	\bar{w}_n	$\frac{W(\hat{B}_n) - \bar{w}_{n-1}}{\bar{w}_n - \bar{w}_{n-1}}$
8	0.3017766953	0.3129490191	0.3132295145	0.3142087183	0.2227
16	0.3142087183	0.3171454818	0.3172268776	0.3172865746	0.5769
32	0.3172865746	0.3180374156	0.3180504765	0.3180541816	0.7790
64	0.3180541816	0.3182439224	0.3182457366	0.3182459678	0.8870
128	0.3182459678	0.3182936544	0.3182938926	0.3182939071	0.9428

We remark that $W(\hat{R}_n) < W(\hat{R}_{n-1})$ for all even $n \geq 4$. The polygon \hat{R}_n does not have maximum width for any even $n \geq 4$. When n is even, one can construct a unit-perimeter n -gon with the same width as \hat{R}_{n-1} by adding a vertex in the middle of a side of \hat{R}_{n-1} .

When n has an odd factor m , one can note that

$$W(\hat{R}_{m,n}) = \frac{1}{2n} \cot \frac{\pi}{2n}.$$

Theorem 4 (Audet, Hansen, and Messine [15]). *For all $n \geq 3$, let w_n^* denote the maximal width among all unit-perimeter n -gons and let $\bar{w}_n := \frac{1}{2n} \cot \frac{\pi}{2n}$.*

- When n has an odd factor m , $w_n^* = \bar{w}_n$ is achieved by finitely many equilateral n -gons [3–5], including $\hat{R}_{m,n}$. The optimal n -gon $\hat{R}_{m,n}$ is unique if m is prime and $n/m \leq 2$.
- When $n = 2^s$ with integer $s \geq 2$, $W(\hat{R}_n) < \bar{w}_{n-1} \leq w_n^* < \bar{w}_n$.

When $n = 2^s$, the maximal width w_n^* of unit-perimeter n -gons is only known for $s = 2$. Audet, Hansen, and Messine [15] showed that $w_4^* = \frac{1}{4}\sqrt{6\sqrt{3}-9} > \bar{w}_3 = \frac{1}{6}\sqrt{3}$. For $s \geq 3$, exact solutions appear to be presently out of reach. However, it is interesting to note that

$$W(\hat{B}_n) = \frac{1}{2n} \left(\cot \frac{\pi}{2n} - \tan \left(\frac{\pi}{2n} - \frac{1}{2} \arcsin \left(\frac{1}{2} \sin \frac{2\pi}{n} \right) \right) \right)$$

is a tighter lower bound compared to \bar{w}_{n-1} on w_n^* when $n = 2^s$ and $s \geq 3$. Indeed, we can show that, for all $n = 2^s$ and integer $s \geq 3$,

$$\bar{w}_n - W(\hat{B}_n) = \frac{1}{2n} \tan \left(\frac{\pi}{2n} - \frac{1}{2} \arcsin \left(\frac{1}{2} \sin \frac{2\pi}{n} \right) \right) = \frac{\pi^3}{8n^4} + O\left(\frac{1}{n^6}\right),$$

while

$$\begin{aligned} \bar{w}_n - W(\hat{R}_n) &= \frac{\pi}{4n^2} + O\left(\frac{1}{n^4}\right), \\ \bar{w}_n - \bar{w}_{n-1} &= \frac{\pi}{6n^3} + O\left(\frac{1}{n^4}\right) \end{aligned}$$

for all even $n \geq 4$.

Table 3 lists the widths of \hat{B}_n , along with the upper bounds \bar{w}_n , the lower bounds \bar{w}_{n-1} , and the widths of \hat{R}_n for $n = 2^s$ and $3 \leq s \leq 7$. As n increases, it is not surprising that $W(\hat{B}_n)$ approaches \bar{W}_n much faster than \bar{w}_{n-1} does.

5 Solving the maximal perimeter problem

For any $n = 2^s$ with integer $s \geq 3$, we can construct a convex small n -gon B_n^* with a longer perimeter than B_n by adjusting the angles $\alpha_0, \alpha_1, \dots, \alpha_{\frac{n}{4}}$ from the parametrization of Section 3 to maximize the




Figure 8: Variables $\alpha_0, \alpha_1, \dots, \alpha_{\frac{n}{2}-1}$ for $L(Q_n^*)$: Case of $n = 8$ vertices

perimeter $L(P_n)$ in (2a) [8]. Hence, $L(B_n^*)$ is the optimal value of the problem:

$$L(B_n^*) = \max_{\alpha} \quad L(P_n) = 4 \sin \frac{\alpha_0}{2} + \sum_{k=1}^{n/4-1} 8 \sin \frac{\alpha_k}{2} + 4 \sin \frac{\alpha_{n/4}}{2} \quad (7a)$$

$$\text{s. t.} \quad \alpha_0 + \sum_{k=1}^{n/4-1} 2\alpha_k + \alpha_{n/4} = \pi/2, \quad (7b)$$

$$\sin \alpha_0 - \sum_{k=2}^{n/4} (-1)^k \sin \left(\alpha_0 + \sum_{i=1}^{k-1} 2\alpha_i \right) = -1/2, \quad (7c)$$

$$0 \leq \alpha_k \leq \pi/6 \quad \forall k = 0, 1, \dots, n/4 - 1, \quad (7d)$$

$$0 \leq \alpha_{n/4} \leq \pi/3, \quad (7e)$$

$$L(P_n) \geq L(B_n). \quad (7f)$$

This formulation was used in [7] for $n = 8$ to find the convex small 8-gon of maximal perimeter.

For each $n = 2^s$ with integer $s \geq 2$, one can also construct a convex small n -gon Q_n^* with the same diameter graph as R_{n-1}^+ but larger perimeter. Using a similar parametrization as in Section 3, we can show that

$$L(Q_n^*) = \max_{\alpha} \quad L(P_n) = \sum_{k=0}^{n/2-1} 4 \sin \frac{\alpha_k}{2} \quad (8a)$$

$$\text{s. t.} \quad \sum_{k=0}^{n/2-1} \alpha_k = \pi/2, \quad (8b)$$

$$\sum_{k=0}^{n/2-2} (-1)^k \sin \left(\sum_{i=0}^k \alpha_i \right) = 1/2, \quad (8c)$$

$$0 \leq \alpha_0 \leq \pi/6, \quad (8d)$$

$$0 \leq \alpha_k \leq \pi/3 \quad \forall k = 1, 2, \dots, n/2 - 1, \quad (8e)$$

$$L(P_n) \geq L(R_{n-1}^+). \quad (8f)$$

The variables $\alpha_0, \alpha_1, \dots, \alpha_{\frac{n}{2}-1}$ are defined in Figure 8. Clearly, $Q_4^* \equiv R_3^+$ and $L(Q_4^*) = L_4^*$.

We solved both Problems (7) and (8) on the NEOS Server 6.0 using AMPL with the solver Couenne 0.5.8, which is a branch-and-bound algorithm that aims at finding global optima of noncon-

Table 4: Perimeters of B_n^* and Q_n^*

n	$L(Q_n^*)$	$L(B_n)$	$L(B_n^*)$	\bar{L}_n	$\frac{L(B_n^*) - L(B_n)}{\bar{L}_n - L(B_n)}$
8	3.1195976652 [8]	3.1210621230	3.1211471341 [7, 8]	3.1214451523	0.2219
16	3.1364309268	3.1365427675	3.1365439563 [8]	3.1365484905	0.2077
32	3.1403237758	3.1403310687	3.1403310858 [8]	3.1403311570	0.1945
64	3.1412767891	3.1412772496	3.1412772498 [16]	3.1412772509	0.1907
128	3.1415137723	3.141513801123	3.141513801127	3.141513801144	0.1899

Table 5: Angles $\alpha_0^*, \alpha_1^*, \dots, \alpha_{\frac{n}{4}}^*$ of B_n^*

n	π/n	i	α_{8i}^*	α_{8i+1}^*	α_{8i+2}^*	α_{8i+3}^*	α_{8i+4}^*	α_{8i+5}^*	α_{8i+6}^*	α_{8i+7}^*
8	0.392699	0	0.435281	0.368535	0.398447					
16	0.196350	0	0.201226	0.191978	0.199873	0.194672	0.196525			
32	0.0981748	0	0.0987786	0.0975863	0.0987333	0.0976772	0.0986041	0.0978448	0.0984101	0.0980628
		1	0.0981803							
64	0.0490874	0	0.0491627	0.0490125	0.0491613	0.0490154	0.049157	0.0490211	0.0491501	0.0490293
		1	0.0491407	0.0490398	0.0491293	0.049052	0.0491164	0.0490657	0.0491022	0.0490802
		2	0.0490876							
128	0.0245437	0	0.0245531	0.0245343	0.0245531	0.0245344	0.0245529	0.0245346	0.0245527	0.0245348
		1	0.0245524	0.0245352	0.024552	0.0245356	0.0245515	0.0245361	0.024551	0.0245367
		2	0.0245504	0.0245374	0.0245497	0.0245381	0.0245489	0.0245389	0.0245481	0.0245397
		3	0.0245473	0.0245405	0.0245464	0.0245414	0.0245455	0.0245423	0.0245446	0.0245432
		4	0.0245437							

vex mixed-integer nonlinear optimization problems [11]. We have made AMPL codes available in OPTIGON [13].

Table 4 gives the optimal values $L(B_n^*)$ and $L(Q_n^*)$ for $n = 2^s$ and $3 \leq s \leq 7$, along with the perimeters of B_n and the upper bounds \bar{L}_n . Couenne took less than 1 second to compute each $L(B_n^*)$ or $L(Q_n^*)$ except for $L(Q_{16}^*)$, which was computed in 36 minutes. The results in Table 4 support the following key points:

1. The optimal perimeter $L(B_n^*)$ for each $n \leq 64$ computed agrees with the best value found in the literature.
2. For all $n = 2^s$ and $s \geq 3$, $L(Q_n^*) < L(B_n) < L(B_n^*)$, i.e., Q_n^* is a suboptimal solution.
3. As n increases, the fraction $\frac{L(B_n^*) - L(B_n)}{\bar{L}_n - L(B_n)}$ appears to approach a scalar $b^* \in (0, 1)$, i.e., $\bar{L}_n - L(B_n^*) = O(1/n^6)$.
4. For $n = 8$, $L(B_8^*) = L_8^*$.

The optimal angles α_k^* that produce B_n^* and Q_n^* are given in Tables 5 and 6, respectively. We observe a pattern of damped oscillation, converging in an alternating manner to a mean value around π/n . For Q_n^* , this observation leads to the following theorem:


Theorem 5. Suppose $n = 2^s$ with integer $s \geq 2$. Then there exists a convex small n -gon Q_n such that

$$L(Q_n) = 2n \sin \frac{\pi}{2n} \cos \left(\frac{\pi}{8} - \frac{1}{2} \arcsin \left(\frac{1}{\sqrt{2}} \cos \frac{\pi}{n} \right) \right),$$

$$W(Q_n) = \cos \left(\frac{\pi}{2n} + \frac{\pi}{8} - \frac{1}{2} \arcsin \left(\frac{1}{\sqrt{2}} \cos \frac{\pi}{n} \right) \right),$$

Table 6: Angles $\alpha_0^*, \alpha_1^*, \dots, \alpha_{\frac{n}{2}-1}^*$ of \mathbb{Q}_n^*

n	π/n	i	α_{8i}^*	α_{8i+1}^*	α_{8i+2}^*	α_{8i+3}^*	α_{8i+4}^*	α_{8i+5}^*	α_{8i+6}^*	α_{8i+7}^*
8	0.392699	0	0.301375	0.480058	0.355776	0.433588				
16	0.196350	0	0.172189	0.219956	0.175546	0.216429	0.182713	0.210185	0.192054	0.201725
32	0.0981748	0	0.0920622	0.104242	0.0922572	0.103986	0.0927078	0.103531	0.0933908	0.102886
		1	0.0942718	0.10207	0.0953079	0.101105	0.0964509	0.100022	0.0976502	0.988561
64	0.0490874	0	0.0475548	0.0506167	0.0475665	0.0505995	0.0475937	0.0505686	0.0476362	0.0505242
		1	0.0476935	0.0504667	0.0477649	0.0503966	0.0478497	0.0503143	0.0479468	0.0502207
		2	0.0480553	0.0501163	0.0481739	0.0500023	0.0483013	0.0498794	0.0484363	0.0497487
		3	0.0485773	0.0496115	0.048723	0.0494689	0.0488718	0.0493222	0.0490222	0.0491729
128	0.0245437	0	0.0241605	0.0249269	0.024161	0.0249258	0.0241627	0.0249237	0.0241653	0.0249209
		1	0.0241688	0.0249171	0.0241733	0.0249124	0.0241787	0.0249069	0.024185	0.0249005
		2	0.0241922	0.0248933	0.0242002	0.0248853	0.0242091	0.0248765	0.0242189	0.0248669
		3	0.0242294	0.0248565	0.0242407	0.0248454	0.0242528	0.0248336	0.0242655	0.0248211
		4	0.024279	0.0248079	0.024293	0.0247941	0.0243077	0.0247797	0.024323	0.0247647
		5	0.0243388	0.0247492	0.0243551	0.0247332	0.0243718	0.0247167	0.0243889	0.0246999
		6	0.0244064	0.0246826	0.0244243	0.024665	0.0244424	0.0246471	0.0244607	0.0246289
		7	0.0244793	0.0246105	0.0244979	0.0245919	0.0245167	0.0245732	0.0245355	0.0245544


 Figure 9: Polygons $(\mathbb{Q}_n, L(\mathbb{Q}_n), W(\mathbb{Q}_n))$ defined in Theorem 5

and

$$\begin{aligned}\bar{L}_n - L(\mathbb{Q}_n) &= \frac{\pi^5}{32n^4} + O\left(\frac{1}{n^6}\right), \\ \bar{W}_n - W(\mathbb{Q}_n) &= \frac{\pi^3}{8n^3} + O\left(\frac{1}{n^4}\right).\end{aligned}$$

In particular, for $n = 4$, $L(\mathbb{Q}_4) = L_4^*$ and $W(\mathbb{Q}_4) = W_4^*$.

Proof. The proof is similar to that of Theorem 1. □

For all $n = 2^s$ with integer $s \geq 2$, the diameter graph of \mathbb{Q}_n has a cycle of length $n - 1$ plus one pendant edge. We represent some polygons \mathbb{Q}_n in Figure 9. They are all symmetrical with respect to the vertical pendant edge. In \mathbb{Q}_n , the angles $\alpha_0, \alpha_1, \dots, \alpha_{\frac{n}{2}-1}$ defined in Figure 8 are given by $\alpha_k = \pi/n - (-1)^k \gamma$ for all $k = 0, 1, \dots, n/2 - 1$, with

$$\gamma = \frac{\pi}{4} - \arcsin\left(\frac{1}{\sqrt{2}} \cos \frac{\pi}{n}\right) = \frac{\pi^2}{2n^2} - \frac{\pi^4}{6n^4} + O\left(\frac{1}{n^6}\right).$$

We can show that $L(\mathbb{Q}_n) > L(\mathbb{R}_{n-1}^+)$ and $W(\mathbb{Q}_n) < W(\mathbb{R}_{n-1}^+)$ when $n = 2^s$ with integer $s \geq 3$.

6 Conclusion

Tighter lower bounds on the maximal perimeter and the maximal width of convex small n -gons were provided when n is a power of 2. For all $n = 2^s$ with integer $s \geq 3$, we constructed a convex small n -gon B_n whose perimeter and width cannot be improved for large n by more than $\frac{\pi^7}{32n^6}$ and $\frac{\pi^4}{8n^4}$ respectively.

In addition, under the assumption that Mosshinghoff's conjecture is true, we formulated the maximal perimeter problem as a nonlinear optimization problem involving trigonometric functions and provided global optimal n -gons for $n = 2^s$ and $3 \leq s \leq 7$.

Acknowledgements

The author thanks Charles Audet, Professor at Polytechnique Montréal, for helpful discussions on extremal small polygons and helpful comments on early drafts of this paper.

References

- [1] K. Reinhardt, "Extremale polygone gegebenen durchmessers," *Jahresbericht der Deutschen Mathematiker-Vereinigung*, vol. 31, pp. 251–270, 1922.
- [2] B. Datta, "A discrete isoperimetric problem," *Geometriae Dedicata*, vol. 64, no. 1, pp. 55–68, 1997.
- [3] M. J. Mossinghoff, "Enumerating isodiametric and isoperimetric polygons," *Journal of Combinatorial Theory, Series A*, vol. 118, no. 6, pp. 1801–1815, 2011.
- [4] K. G. Hare and M. J. Mossinghoff, "Sporadic Reinhardt polygons," *Discrete & Computational Geometry*, vol. 49, no. 3, pp. 540–557, 2013.
- [5] K. G. Hare and M. J. Mossinghoff, "Most Reinhardt polygons are sporadic," *Geometriae Dedicata*, vol. 198, no. 1, pp. 1–18, 2019.
- [6] N. K. Tamvakis, "On the perimeter and the area of the convex polygon of a given diameter," *Bulletin of the Greek Mathematical Society*, vol. 28, pp. 115–132, 1987.
- [7] C. Audet, P. Hansen, and F. Messine, "The small octagon with longest perimeter," *Journal of Combinatorial Theory, Series A*, vol. 114, no. 1, pp. 135–150, 2007.
- [8] M. J. Mossinghoff, "Isodiametric problems for polygons," *Discrete & Computational Geometry*, vol. 36, no. 2, pp. 363–379, 2006.
- [9] A. Bezdek and F. Fodor, "On convex polygons of maximal width," *Archiv der Mathematik*, vol. 74, no. 1, pp. 75–80, 2000.
- [10] C. Audet, P. Hansen, F. Messine, and J. Ninin, "The small octagons of maximal width," *Discrete & Computational Geometry*, vol. 49, no. 3, pp. 589–600, 2013.
- [11] P. Belotti, J. Lee, L. Liberti, F. Margot, and A. Wächter, "Branching and bounds tightening techniques for non-convex MINLP," *Optimization Methods & Software*, vol. 24, no. 4-5, pp. 597–634, 2009.
- [12] C. Audet, P. Hansen, and F. Messine, "Ranking small regular polygons by area and by perimeter," *Journal of Applied and Industrial Mathematics*, vol. 3, no. 1, pp. 21–27, 2009.

- [13] C. Bingane, “OPTIGON: Extremal small polygons.” <https://github.com/cbingane/optigon>, September 2020.
- [14] C. Bingane, “Largest small polygons: A sequential convex optimization approach,” *Optimization Letters*, 2022.
- [15] C. Audet, P. Hansen, and F. Messine, “Isoperimetric polygons of maximum width,” *Discrete & Computational Geometry*, vol. 41, no. 1, pp. 45–60, 2009.
- [16] M. J. Mossinghoff, “An isodiametric problem for equilateral polygons,” in *Tapas in Experimental Mathematics* (T. Amdeberhan and V. H. Moll, eds.), vol. 457 of *Contemporary Mathematics*, pp. 237–252, American Mathematical Society, 2008.