# Semicoercive Variational Inequalities - From Existence to Numerical Solution of Nonmonotone Contact Problems

Nina Ovcharova · Joachim Gwinner

May 17, 2022

Dedicated to the memory of Professor V.F. Demyanov and Professor S. Schaible

**Abstract** In this paper we present a novel numerical solution procedure for semicoercive hemivariational inequalities. As a model example we consider a unilateral semicoercive contact problem with nonmonotone friction and provide numerical results for benchmark tests

**Keywords** Semicoercivity  $\cdot$  Pseudomonotone bifunction  $\cdot$  Hemivariational inequality  $\cdot$  Plus function  $\cdot$  Smoothing approximation  $\cdot$  Finite element discretization  $\cdot$  Unilateral contact **AMS** 74G15, 74M15, 74S05

# 1 Introduction

This paper presents a contribution to constructive nonsmooth analysis [1] in view of application to nonsmooth continuum mechanics [2]. It is devoted to the numerical solution of nonmonotone semicoercive contact problems in solid mechanics modelled by hemivariational inequalities, a class of variational inequalities (VIs) introduced and studied by Panagiotopoulos [3], see also [4,5,6]. Here we treat unilateral contact problems with nonmonotone friction [7], which occur with adhesion and delamination in the delicate situation, where the body is not fixed along some boundary part, but is only subjected to surface tractions and body forces. Thus there is a loss of coercivity leading to so-called semicoercive or noncoercive variational problems [8,9].

The existence theory of hemivariational inequalities (HVIs) and of more abstract topologically (in the sense of Brézis [10]) pseudomonotone VIs, also in the semicoercive case

N. Ovcharova

Institute of Mathematics, Department of Aerospace Engineering, Universität der Bundeswehr München, Germany, E-mail: nina.ovcharova@unibw.de (🗷)

# J. Gwinner

Institute of Mathematics, Department of Aerospace Engineering, Universität der Bundeswehr München, Germany, E-mail: joachim.gwinner@unibw.de

is well documented in the literature. Without claim of completeness we can cite [11-28] in chronological order

While there are studies of numerical solution methods for *coercive* HVIs, see the book [29] of Haslinger, Miettinen, and Panagiotopoulos and also the more recent paper [30], and there are works on the numerical treatment of semicoercive *monotone* VIs, see the book of Kaplan and Tichatschke [31] and the papers [32,33,34,35,36,37,38], to the best of the authors' knowledge, efficient methods for the numerical solution of semicoercive HVIs supported by rigorous mathematical analysis are missing. It is the purpose of the present paper to initiate work in this direction. To this end we extend the convergence result in [39] for a general approximation scheme for the solution of pseudomonotone VIs to the semicoercive case. Then based on this result, we combine finite element discretization with regularization techniques of nondifferentiable optimization [40,41] to arrive at a novel solution procedure for semicoercive HVIs.

The paper is organized as follows. In the next section we revisit the existence theory for pseudomonotone VIs that is is necessary for the subsequent development. Here we exhibit also an interesting relation between a sign condition for the data of the pseudomonotone VI, that is crucial for existence, and the more recent notion of well-positioned convex sets in the stability theory for semicoercive pseudomonotone VIs due to Adly, Théra, and Ernst [20, 42]. Then we adapt the general approximation scheme of Glowinski [43] to pseudomonotone VIs and provide convergence results thus extending [39, Theorem 3.1] to the semicoercive situation. Then we apply this convergence theorem to HVIs in linear elasticity, combine finite element discretization with regularization techniques of nondifferentiable optimization from [40,41] to obtain a novel solution procedure for semicoercive HVIs. To demonstrate its efficiency we finally provide numerical results for benchmark tests.

#### 2 Semicoercive pseudomonotone variational inequality

In this section we revisit the existence theory for pseudomonotone VIs that is necessary for the subsequent treatment of HVIs. Here we follow [44,45,33,46] for the setting of pseudomonotone bifunctions and semicoercivity. Moreover, we also exhibit a relation to the more recent notion of well-positioned convex sets introduced and studied in [20,42].

Let V be a reflexive Banach space and  $V^*$  its dual. We denote by  $\langle \cdot, \cdot \rangle$  the duality pairing between V and  $V^*$ , and by  $\| \cdot \|$  and  $\| \cdot \|_*$  the norm and the dual norm on V and  $V^*$ , respectively. We are concerned with a linear semicoercive operator A from V to  $V^*$ , i.e. there exists some positive constant  $c_0$  such that

$$\langle Av, v \rangle \ge c_0 |v|^2 \quad \forall v \in V,$$

where  $|\cdot|$  denotes a lower semicontinuous seminorm on V. Thus, the kernel Y of  $|\cdot|$ , defined by

$$Y = \{ y \in V : |y| = 0 \}$$

is a closed nontrivial subspace.

Further, let  $g \in V^*$  be a continuous linear form,  $K \subseteq V$  a nonvoid closed convex set such that  $0 \in K$ , and consider  $\varphi : K \times K \to \mathbb{R}$  such that  $\varphi(\cdot, \cdot)$  is pseudomonotone,  $\varphi(\cdot, v)$  is upper semicontinuous on the intersection of K with any finite dimensional subspace of V and, moreover, there exists some positive constant c such that

$$\varphi(v,0) \le c||v|| \quad \forall v \in V. \tag{1}$$

We recall that the bifunction  $\varphi(\cdot,\cdot)$  is pseudomonotone if  $u_n \rightharpoonup u$  (weakly) in V and  $\liminf_{n \to \infty} \varphi(u_n,u) \ge 0$  implies that  $\limsup_{n \to \infty} \varphi(u_n,v) \le \varphi(u,v)$  for all  $v \in V$ .

In this paper we deal with the semicoercive variational problem  $VI(A, \varphi, g, K)$ : Find  $u \in K$  such that

$$\langle Au, v - u \rangle + \varphi(u, v) \ge \langle g, v - u \rangle \quad \forall v \in K.$$

From [44,33] we adopt the following assumption:

(A1<sup>s</sup>) for any sequence  $\{v_n\}$  with  $|v_n| \to 0$ ,  $v_n \to v$  and  $||v_n|| \ge \eta$  for some  $\eta > 0$  there exists a subsequence  $\{v_{n_k}\}_{k \in \mathbb{N}}$  such that  $v_{n_k} \to v$  in V.

According to [44], [45, Section 5.3] this condition is fulfilled if the norm  $\|\cdot\|$  on V is equivalent to  $\||\cdot\||+|\cdot|$ , where  $\||\cdot\||$  is another norm on V, the dimension of Y is finite and there exists  $\alpha>0$  such that

$$\inf_{v \in V} |||v - y||| \le \alpha |v| \quad \forall v \in V,$$

from which the seminorm  $|\cdot|$  is continuous.

In addition, we resume from [44,33] that

(A2<sup>s</sup>) either (i)  $Y \cap K$  is bounded or (ii) g satisfies

$$\langle g, y \rangle < 0 \quad \forall y \in \{Y \cap K\} \setminus \{0\}.$$

Condition  $(A2^s)$  (ii) implies that the linear functional  $g \in V^*$  is bounded from above on  $K \cap Y$ . In other words, g belongs to the barrier cone of the set  $Y \cap K$  defined by

$$B(Y \cap K) = \{ g \in V^* : \sup_{v \in Y \cap K} \langle g, v \rangle < \infty \}.$$

Let the interior of the barrier cone be nonempty. Then we can show that  $g \in \text{int} B(Y \cap K)$ . According to the terminology used in [20,42], this is equivalent to the property that the set  $Y \cap K$  is well-positioned.

**Proposition 2.1** *Let*  $int B(Y \cap K) \neq \emptyset$ . Suppose  $(A1^s)$ . Then condition  $(A2^s)$  (ii) implies that  $g \in int B(Y \cap K)$ .

*Proof* Suppose by contradiction that g does not belong to  $\operatorname{int} B(Y \cap K)$ . Then, for any sequence  $\{\varepsilon_n\}$  with  $\varepsilon_n \to 0^+$ , there exist  $\|h_n\|_* \le 1$  and  $y_n \in Y \cap K$  such that

$$\langle g + \varepsilon_n h_n, y_n \rangle > n.$$
 (2)

If  $y_n = 0$ , we trivially get a contradiction. Therefore, let  $y_n \neq 0$ . Using (A2<sup>s</sup>) (ii), it follows that

$$\varepsilon_n ||y_n|| \ge \varepsilon_n ||h_n||_* ||y_n|| > \langle g + \varepsilon_n h_n, y_n \rangle > n,$$

and therefore,  $\lim_{n\to\infty} \|y_n\| = +\infty$ . Define  $t_n := \frac{1}{\|y_n\|}$  and consider the sequence  $\{t_n y_n\}$ . Since  $0 \in K$  and K is convex it follows that  $t_n y_n \in K$  for almost all n. Moreover,  $t_n y_n \in Y$ . Since  $\|t_n y_n\| = 1$ , we can extract a subsequence that converges weakly to some  $\tilde{y} \in V$ . In virtue of  $(A1^s)$  we can pass to a strongly convergent subsequence. Hence  $\tilde{y} \neq 0$ . Since  $Y \cap K$  is closed, we can conclude that  $\tilde{y} \in Y \cap K \setminus \{0\}$ .

Multiplying (2) by  $t_n$ , passing to the limit as  $n \to \infty$  and taking into account (A2<sup>s</sup>) (ii) again, we obtain

$$0 > \langle g, \tilde{y} \rangle \ge \liminf_{n \to \infty} n t_n \ge 0,$$

which yields a contradiction.  $\Box$ 

If *K* is a nonvoid convex cone - note that this is the relevant case in view of our subsequent application to HVIs -, the converse is also true.

**Proposition 2.2** Let K be a nonvoid convex cone. Then

$$g \in int B(Y \cap K) \Rightarrow \langle g, v \rangle < 0 \quad \forall y \in \{Y \cap K\} \setminus \{0\}.$$

*Proof* First, since *K* is cone, the barrier cone coincides with the polar cone, i.e.

$$B(Y \cap K) = (Y \cap K)^{-}. \tag{3}$$

Because  $g \in \operatorname{int}(Y \cap K)^-$ , there exists  $\varepsilon > 0$  such that for all  $h \in V^*$  with  $||h||_* \le 1$  we have  $g + \varepsilon h \in (Y \cap K)^-$ .

Now assume that there exists  $\tilde{y} \in (Y \cap K) \setminus \{0\}$  such that  $\langle g, \tilde{y} \rangle \geq 0$ . Then by an easy consequence of the Hahn-Banach extension theorem there exists  $\tilde{h} \in V^*$  such that  $\|\tilde{h}\|_* = 1$  and  $\langle \tilde{h}, \tilde{y} \rangle = \|\tilde{y}\|$ . Then

$$\langle g + \varepsilon \tilde{h}, \tilde{y} \rangle \geq \varepsilon \langle \tilde{h}, \tilde{y} \rangle = \varepsilon ||\tilde{y}|| > 0$$

contradicting that  $g + \varepsilon \tilde{h} \in (Y \cap K)^-$ .  $\square$ 

Remark 2.1 The barrier cone in finite-dimensional setting has a nonempty interior. While this is trivial for a compact  $Y \cap K$ , let us show this in the relevant case of a convex closed cone K. Then indeed, we have  $Y \cap K = \sum_{k=1}^{n} \mathbb{R}_{-}w_k$  with  $||w_k|| = 1$ . Construct  $h_k \in V^*$  such

that

$$||h_k||_* = 1$$
 and  $\langle h_k, w_l \rangle = \delta_{kl}$ .

Let  $h \in V^*$  be such that  $||h||_* \le \frac{1}{2}$ . Hence,  $h = \sum_{k=1}^n \gamma_k h_k$  and  $\langle h, w_k \rangle = \gamma_k$ . Moreover, we have

$$\frac{1}{2} \ge ||h||_* = \sup_{\|\nu\|=1} |\langle h, \nu \rangle| \ge |\langle h, w_k \rangle| = |\gamma_k|.$$

With  $e := \sum_{k=1}^{n} h_k$  we have  $e + h = \sum_{k=1}^{n} (1 + \gamma_k) h_k \in (Y \cap K)^-$ , which due to (3) implies the nonemptiness of the interior of the barrier cone.

Let us now turn back to the problem  $VI(A, \varphi, g, K)$ . We provide the subsequent existence theorem in view of application to HVIs. Since in HVIs, the bifunction  $\varphi$  (see the later definition (8)), cannot be controlled by the seminorm, but only by the norm (see (1) that is proved in [40, Lemma 15] for HVIs), we have to strengthen the condition  $(A2^s)$  (ii) to

(iii) 
$$\exists c > 0 : \langle g, y \rangle < -c$$
 for all  $y \in Y \cap K$  with  $||y|| = 1$ .

Then we can show the following existence result without any compact imbedding assumption.

**Theorem 2.1** Let  $A: V \to V^*$  be a semicoercive linear operator and  $\varphi: V \times V \to \mathbb{R}$  be a pseudomonotone bifunction such that (1) holds and such that  $\varphi(\cdot, v)$  is upper semicontinuous on the intersection of K with any finite dimensional subspace of V. Under assumptions ( $AI^s$ ) and ( $A2^s$ ) (i) or (iii), the problem  $VI(A, \varphi, g, K)$  has a solution.

*Proof* For the proof we use a recession argument that goes back to Stampacchia [47], Hess [48] and Schatzman [49]. Similar reasoning based on a recession analysis can be found in [44,32,33,12,13,16,17,18,21,24]. Another approach to existence results for semicoercive pseudomonotone variational inequalities is a regularization procedure based on adding of a coercive term, see [16,19,22,28].

In view of [44, Theorem 3.9], [48, Proposition] it is sufficient to show the existence of a constant R > 0 such that

$$\langle Av, -v \rangle + \varphi(v, 0) + \langle g, v \rangle < 0 \quad \forall v \in K \quad \text{with} \quad ||v|| = R.$$

Assume the contrary, i.e., there exists a sequence  $v_n \in K$  such that  $||v_n|| \to \infty$  and

$$\langle Av_n, -v_n \rangle + \varphi(v_n, 0) + \langle g, v_n \rangle \ge 0.$$

Hence, using the semicoercivity of the operator A and (1), it follows that

$$|c_0|v_n|^2 \le \langle Av_n, v_n \rangle \le \langle g, v_n \rangle + \varphi(v_n, 0) \le ||g||_* ||v_n|| + c||v_n||. \tag{4}$$

Set  $y_n = \frac{v_n}{\|v_n\|}$ . Since  $0 \in K$  and K is convex, it follows that  $y_n \in K$  for large enough n as well. Moreover,  $\|y_n\| = 1$  and consequently, we can extract a subsequence that converges weakly to some y in the weakly closed set K.

Now, we show that  $|y_n|_V \to 0$ . Assume not. Consequently, there exists a subsequence, again denoted by  $\{y_n\}$ , such that  $|y_n| \ge c_3 > 0$ . Dividing (4) by  $||v_n||$  yields

$$c_0 ||v_n|| ||y_n||^2 \le ||g||_* + c$$

and hence.

$$|y_n| \le \frac{\|g\|_* + c}{c_0|y_n|\|v_n\|} \le \frac{\|g\|_* + c}{c_0 c_3 \|v_n\|}.$$

Since  $||v_n|| \to \infty$ , we arrive at a contradiction, and therefore  $|y_n| \to 0$ . Further, by (A1<sup>s</sup>), we can extract a subsequence again denoted by  $\{y_n\}$  that converges strongly to y in V. Since  $||\cdot||$  is continuous, we have ||y|| = 1 and, in particular,  $y \neq 0$ . Moreover, by the continuity of  $|\cdot|$  it follows that |y| = 0. In conclusion,  $y \in Y \cap K$  and  $y \neq 0$ .

Next we claim that  $\lambda y$  belongs to K for any  $\lambda > 0$ . Indeed, because of  $||v_n|| \to \infty$ , there exists  $n_0$  such that  $||v_n|| > \lambda$  for all  $n \ge n_0$ . By convexity,  $\lambda y_n \in K$  for all  $n \ge n_0$ , and by the closedness of K,  $\lambda y \in K$ . Hence, if  $Y \cap K$  is bounded, the existence of  $y \in Y \cap K$ ,  $y \ne 0$ , leads immediately to a contradiction. Otherwise, we obtain from (4) that

$$0 \le \langle g, v_n \rangle + c \|v_n\|. \tag{5}$$

Dividing (5) by  $||v_n||$  we arrive in the limit at

$$0 \le \langle g, y \rangle + c$$
,

which is a contradiction to (iii). □

Some comments on the conditions (A2<sup>s</sup>) are in order now. For bilateral contact with given  $\underline{h} \leq \overline{h}$  in  $L^{\infty}(\Gamma_c)$  on the boundary part  $\Gamma_c$  of a bounded domain  $\Omega$  and the constraint set  $K = \{u \in V : \underline{h} \leq u | \Gamma_c \leq \overline{h}\}$  in a Sobolev space V on  $\Omega$ , simply set  $K = K - \underline{h}$ , where  $\underline{h} \in V$  extends  $\underline{h}$  to  $\Omega$ . Then the existence theorem applies to the bounded set  $Y \cap K$ . - To simplify the interpretation of the conditions involved in the more delicate unbounded case, assume V is a Hilbert space, as in the subsequent section on HVIs. Condition (A2<sup>s</sup>) (ii)

postulates that the applied force g forms an obtuse angle with the directions y of escape. In contrast, condition (iii) demands together with the Cauchy-Schwarz inequality that for all  $y \in Y \cap K$  with ||y|| = 1.

$$-1 \le \frac{1}{\|g\|_*} \langle g, y \rangle < -\frac{c}{\|g\|_*}.$$

This means that the directions y of escape should stay in a given angle range with the applied force g and moreover g should be large enough, ||g|| > c.

# 3 General approximation acheme for semicoercive pseudomonotone variational inequalities

Let T be a directed set and  $\{V_t\}_{t\in T}$  be a family of finite-dimensional subspaces of V. While K is contained in V,  $K_t$  is a nonempty, closed convex subset of  $V_t$ , not necessarily contained in K. Therefore, for the approximation of K by  $K_t$ , we employ Mosco convergence, see the hypotheses (H1) - (H2) below.

- (H1) If  $\{v_{t'}\}_{t'\in T'}$  weakly converges to v in V,  $v_{t'}\in K_{t'}$   $(t'\in T')$  for a subnet  $\{K_{t'}\}_{t'\in T'}$  of the net  $\{K_t\}_{t\in T}$ , then  $v\in K$ .
- (H2) For any  $v \in K$  and any  $t \in T$  there exists  $v_t \in K_t$  such that  $v_t \to v$  in V.

Since  $0 \in K$ , by a translation argument, we can simply assume that  $0 \in K_t$  for all  $t \in T$ . Further, we replace  $\varphi$  by some some approximation  $\varphi_t$  satisfying

(H3) For any  $t \in T$ ,  $\varphi_t$  is pseudomonotone,  $\varphi_t(\cdot, v)$  is upper semicontinuous on the intersection of  $K_t$  with any finite dimensional subspace of V, and

$$\varphi_t(u_t,0) \leq c \|u_t\| \quad \forall u_t \in K_t.$$

(H4) For any nets  $\{u_t\}$  and  $\{v_t\}$  such that  $u_t \in K_t$ ,  $v_t \in K_t$ ,  $u_t \rightharpoonup u$ , and  $v_t \to v$  in V it follows that

$$\limsup_{t\in T} \varphi_t(u_t,v_t) \leq \varphi(u,v).$$

Altogether the problem  $VI(A, \varphi, g, K)$  is approximated by the problem  $VI(A, \varphi_t, g, K_t)$ : Find  $u_t \in K_t$  such that

$$\langle Au_t, v_t - u_t \rangle + \varphi_t(u_t, v_t) \ge \langle g, v_t - u_t \rangle \quad \forall v_t \in K_t.$$

In some computations it will be necessary to replace also A and g by some approximations  $A_t$  and  $g_t$ , defined for example by a numerical integration procedure which is used in the finite element discretization. But this is rather standard; therefore we do not elaborate on this aspect. In view of our applications to hemivariational inequalities, we assume also that  $\varphi(u,u)=0$  for all  $u\in V$ .

**Theorem 3.1** Under assumptions  $(A1^s)$ - $(A2^s)$ , and hypotheses (H1) - (H4), the family  $\{u_t\}$  of solutions to  $VI(A, \varphi_t, g, K_t)$  is uniformly bounded in V. Moreover, there exists a subnet  $\{u_{t'}\}_{t' \in T'}$  of  $\{u_t\}$  that converges weakly in V to a solution u of the problem  $VI(A, \varphi, g, K)$  and satisfies  $\lim_{t' \in T'} |u_{t'} - u| = 0$ .

*Proof* The existence and uniform boundedness of the family  $\{u_t\}$  can be shown by using the same arguments as those used to prove Theorem 2.1. Further, we can extract a subnet, again denoted by  $\{u_t\}$ , such that  $u_t \rightharpoonup u$ . In view of (H1),  $u_t$  belongs to  $u_t$ . Since  $u_t \mapsto \langle Av, v \rangle$  is convex and continuous, hence weakly lower semicontinuous,

$$\langle Au, u \rangle \le \liminf_{t \in T} \langle Au_t, u_t \rangle.$$
 (6)

Now, take an arbitrary  $v \in K$ . By (H2), there exists a net  $\{v_t\}$  such that  $v_t \in K_t$  and  $v_t \to v$  in V. By (H4) and (6), we get from  $VI(A, \varphi_t, g, K_t)$  that for any  $v \in K$ 

$$\begin{split} \langle Au, v - u \rangle + \varphi(u, v) &\geq \limsup_{t \in T} \langle Au_t, v_t - u_t \rangle + \limsup_{t \in T} \varphi_t(u_t, v_t) \\ &\geq \limsup_{t \in T} \left\{ \langle Au_t, v_t - u_t \rangle + \varphi_t(u_t, v_t) \right\} \\ &\geq \lim_{t \in T} \langle g, v_t - u_t \rangle = \langle g, v - u \rangle \end{split}$$

and consequently, u is a solution to  $VI(A, \varphi, g, K)$ .

Finally, we show the convergence with respect to the seminorm  $|\dot{}|$ . For this purpose, by (H2), we find a subnet  $\{\bar{u}_t\}$ ,  $\bar{u}_t \in K_t$  such that  $\bar{u}_t \to u$ . We start with the relation

$$c|\bar{u}_t - u_t|^2 \le \langle A\bar{u}_t, \bar{u}_t - u_t \rangle + \langle Au_t, u_t - \bar{u}_t \rangle. \tag{7}$$

The first term goes to zero, since  $A\bar{u}_t \to Au$  and  $\bar{u}_t - u_t \rightharpoonup 0$  in V.

From the definition of  $VI(A, \varphi_t, g, K_t)$  it follows that

$$\langle Au_t, u_t - \bar{u}_t \rangle \leq \langle g, u_t - \bar{u}_t \rangle + \varphi_t(u_t, \bar{u}_t).$$

Hence by (H4),

$$\limsup_{t\in T}\langle Au_t, u_t - \bar{u}_t\rangle \leq \limsup_{t\in T} \varphi_t(u_t, \bar{u}_t) \leq \varphi(u, u) = 0,$$

and therefore, (7) entails in the limit that  $\limsup_{t \in T} c|\bar{u}_t - u_t|^2 \le 0$ , hence,  $|\bar{u}_t - u_t| \to 0$ . Further, using the triangle inequality, we get in the limit

$$0 \le \lim_{t \in T} |u_t - u| \le \lim_{t \in T} |\bar{u}_t - u_t|_V + \lim_{t \in T} |\bar{u}_t - u|_V = 0$$

and the proof is complete.  $\Box$ 

# 4 Approximation of a semicoercive hemivariational inequality

Let  $V = H^1(\Omega; \mathbb{R}^d)$  (d = 2,3) and K be a nonempty closed convex subset of V which will be specified later. Let  $\Omega \subset \mathbb{R}^d$  be a bounded domain with a Lipschitz boundary  $\partial \Omega$ . Decompose  $\partial \Omega$  into a Neumann part  $\Gamma_N$  and a contact part  $\Gamma_C$  with positive measure. Note that here the Dirichlet part is assumed to be empty.

We prescribe surface tractions  $\mathbf{t} \in L^2(\Gamma_N; \mathbb{R}^d)$  on  $\Gamma_N$  and nonmonotone, generally multivalued boundary conditions on  $\Gamma_c$ . Moreover, we suppose also that the body is subject to volume force  $\mathbf{f} \in L^2(\Omega; \mathbb{R}^d)$ .

Let  $a(\cdot, \cdot)$  be the bilinear form of the linear elasticity, i.e.

$$a(u,v) = \int_{\Omega} C_{ijhk} \varepsilon_{ij}(u) \varepsilon_{hk}(v) dx =: \langle Au, v \rangle,$$

where  $C_{ijhk} \in L^{\infty}$  and  $\mathscr{C} = \{C_{ijhk}\}$  is assumed to be uniformly positive definite, and  $\varepsilon(u)$  is the symmetric strain tensor defined by

$$\varepsilon_{ij} = \frac{1}{2}(u_{i,j} + u_{j,i}).$$

Then, the bilinear form is semicoercive and the space of rigid body motions is

$$\ker a(\cdot,\cdot) = \{u \in H^1(\Omega; \mathbb{R}^2) : a(u,u) = 0\} \neq \{0\}.$$

In particular,

(i) if 
$$\Omega \subset \mathbb{R}^2$$
 then  $\ker a(\cdot, \cdot) = \{u(x) = (a_1 - bx_2, a_2 + bx_1), a_1, a_2, b \in \mathbb{R}, \forall x \in \Omega\};$ 

(ii) if 
$$\Omega \subset \mathbb{R}^3$$
 then  $\ker a(\cdot, \cdot) = \{u(x) = a + b \land x, a, b \in \mathbb{R}^3, \forall x \in \Omega\}.$ 

We define

$$\varphi(u,v) := \int_{\Gamma_c} f^0(\gamma u(s); \gamma v(s) - \gamma u(s)) \, ds \quad \forall u,v \in V,$$
 (8)

the linear form

$$\langle g, v \rangle := \int_{\Omega} \mathbf{f} \cdot v \, dx + \int_{\Gamma_N} \mathbf{t} \cdot \gamma v \, ds,$$

and consider the semicoercive hemivariational inequality: Find  $u \in K$  such that

$$\langle Au, v - u \rangle + \varphi(u, v) > \langle g, v - u \rangle \quad \forall v \in K.$$
 (9)

Here,  $f^0(\xi;\eta)$  is the generalized Clarke derivative [50] of a locally Lipschitz function  $f:\mathbb{R}^d\to\mathbb{R}$  at  $\xi\in\mathbb{R}^d$  in the direction  $\eta\in\mathbb{R}^d$ , and  $\gamma$  stands for the trace operator from  $H^1(\Omega;\mathbb{R}^d)$  into  $L^2(\Gamma_c;\mathbb{R}^d)$  with the norm  $\gamma_0$ . It is worthwhile to note that the generalized Clarke derivative coinsides with the classical directional derivative only for Clarke regular functions, like convex and maximum-type functions.

Further, we require the following growth condition on the locally Lipschitz superpotential  $f: \mathbb{R} \to \mathbb{R}$ :

(i) 
$$|\eta| \le c_3(1+|\xi|)$$
 for all  $\eta \in \partial f(\xi)$  with  $c_3 > 0$ ;

(ii) 
$$\eta(-\xi) < c_4|\xi|$$
 for all  $\eta \in \partial f(\xi)$  with  $c_4 > 0$ ;

According to [40, Lemma 15] the bifunction  $\varphi: V \times V \to \mathbb{R}$  is well-defined, pseudomonotone, upper-semicontinuous and the condition (1) is satisfied with

$$c := c_4 \operatorname{meas}(\Gamma_c)^{1/2} \gamma_0. \tag{10}$$

According to Theorem 2.1, the HVI (9) has at least one solution provided that  $\langle g, v \rangle < -c$  for all  $v \in K \cap \ker a(\cdot, \cdot)$  with ||v|| = 1.

The approximation of the HVI is based first on the smoothing of the nonsmooth functional  $\varphi$  defined on the contact boundary and then, on discretizing of the regularized problem by finite elements. For more details we refer to [41,40].

# 4.1 Regularization

Let  $S: \mathbb{R}^d \times \mathbb{R}_{++} \to \mathbb{R}$  be a continuously differentiable approximation of f in the sense that

$$\lim_{z \to x, \varepsilon \to 0^+} S(z, \varepsilon) = f(x) \quad \forall x \in \mathbb{R}^d.$$

Define  $DJ_{\varepsilon}:V\to V^*$  by

$$\langle DJ_{\varepsilon}(u), v \rangle = \int_{\Gamma_{\varepsilon}} \nabla_{x} S(\gamma u(s), \varepsilon) \cdot \gamma v(s) ds.$$

The regularized problem reads now as follows: find  $u_{\varepsilon} \in K$  such that

$$\langle Au_{\varepsilon} - g, v - u_{\varepsilon} \rangle + \langle DJ_{\varepsilon}(u_{\varepsilon}), v - u_{\varepsilon} \rangle \ge 0 \quad \forall v \in K.$$
 (11)

#### 4.2 Finite Element Discretization

We consider a regular triangulation  $\mathcal{T}_h$  of  $\Omega$  and define

$$V_h = \{ v_h \in C(\bar{\Omega}; \mathbb{R}^d) : v_h|_T \in (\mathbb{P}_1)^d, \, \forall T \in \mathscr{T}_h \}$$

as a space of all continuous piecewise linear functions. Here,  $\mathbb{P}_1$  consists of all polynomials of degree at most one. In addition, we have a family  $\{K_h\}$  of nonempty closed convex subsets of  $V_h$  that will be specified later on, not necessarily contained in K, such that (H1) and (H2) are satisfied. We use trapezoidal quadrature rule to approximate  $\langle DJ_{\mathcal{E}}(\cdot), \cdot \rangle$  as follows

$$\langle DJ_{\varepsilon}(u_h), v_h \rangle \approx \frac{1}{2} \sum_{i} |P_i P_{i+1}| \left[ \nabla_x S(\gamma u_h(P_i), \varepsilon) \cdot \gamma v_h(P_i) + \nabla_x S(\gamma u_h(P_{i+1}), \varepsilon) \cdot \gamma v_h(P_{i+1}) \right]$$
  
=:  $\varphi_{\varepsilon,h}(u_h, v_h)$ ,

where we have summed over all sides  $(P_i, P_{i+1})$  of the triangles of  $\mathcal{T}^h$  whose union gives the contact boundary  $\Gamma_c$ .

The discretization of the regularized problem (11) reads now: Find  $u_{\varepsilon h} \in K_h$  such that

$$a(u_{\varepsilon h}, v_h - u_{\varepsilon h}) + \varphi_{\varepsilon, h}(u_h, v_h) \ge \langle g, v_h - u_{\varepsilon h} \rangle \quad \forall v_h \in K_h. \tag{12}$$

According to [40],  $\varphi_{\varepsilon,h}$  is pseudomonotone and satisfies the hypotheses (H3) and (H4). Therefore, due to Theorem 3.1 with  $t = (\varepsilon, h) \in T = \mathbb{R}_{++} \times \mathbb{R}_{++}$ , there exists a solution  $u_{\varepsilon h}$  to the discrete regularized problem (12), the family  $\{u_{\varepsilon h}\}$  is uniformly bounded and any weak accumulation point of  $\{u_{\varepsilon h}\}$  is a solution of the problem (9).

#### **5 Numerical Results**

#### 5.1 Statement of the problem

As a model example we consider an unilateral contact of an elastic body with a rigid foundation under given forces and a nonmonotone friction law on the contact boundary. Let  $\Omega$  be the linear elastic body represented by the unit square  $1m \times 1m$  with modulus of elasticity  $E = 2.15 \times 10^{11} N/m^2$  and Poisson's ration v = 0.29 (steel). The boundary  $\Gamma := \partial \Omega$  is decomposed into a Neumann part and a contact part. We emphasize that in all our benchmark




Fig. 1 2D benchmark examples with force distribution and boundary decomposition. In both cases  $u_1 = 0$  on  $\Gamma_3$ ,  $u_2$ - arbitrary on  $\Gamma_3$ 

examples no Direchlet boundary part is assumed. In particular, on the part  $\Gamma_3$  of the boundary we assume that the horizontal displacement  $u_1$  is zero, but the vertical displacement  $u_2$  is not fixed, see Fig. 1.

The linear Hooke's law is given by

$$\sigma_{ij}(\mathbf{u}) = \frac{Ev}{1 - v^2} \delta_{ij} tr(\varepsilon(\mathbf{u})) + \frac{E}{1 + v} \varepsilon_{ij}(\mathbf{u}), \quad i, j = 1, 2,$$
(13)

where  $\delta_{ij}$  is the Kronecker symbol and

$$tr(\boldsymbol{\varepsilon}(\mathbf{u})) := \boldsymbol{\varepsilon}_{11}(\mathbf{u}) + \boldsymbol{\varepsilon}_{22}(\mathbf{u}).$$

The body is loaded with horizontal forces  $F_1$  on  $\Gamma_1$  and vertical forces  $F_2$  on  $\Gamma_2$ . The volume forces are neglected. In our experiments we have used the data:

$$\mathbf{F}_1 = (\pm P, 0)$$
 with  $P = 1 \times 10^6 N/m^2$   
 $\mathbf{F}_2 = (0, -Q)$  with  $Q = 1 \times 10^6 N/m^2$ .

Further, let **n** be the unit outer normal vector on the boundary  $\partial \Omega$ . The stress vector on the surface is decomposed into the normal, respectively, the tangential stress:

$$\sigma_n := \sigma(u)\mathbf{n} \cdot \mathbf{n}, \quad \sigma_t := \sigma(u)\mathbf{n} - \sigma_n\mathbf{n}.$$

In addition, we assume that

$$\begin{cases} u_2(s) \geq 0 & s \in \Gamma_c \\ -\sigma_t(s) \in \partial j(u_1(s)) & \text{for a.a. } s \in \Gamma_c. \end{cases}$$

The assumed nonmonotone multivalued law  $\partial j$  holding in the tangential (horizontal) direction is depicted in Fig. 2 with parameters  $\delta = 9.0 \times 10^{-6} m$ ,  $\gamma_1 = 1.0 \times 10^3 N/m^2$  and  $\gamma_2 = 0.5 \times 10^3 N/m^2$ . Notice that here j is a minimum of one convex quadratic and one linear function, i.e.

$$j(x) = \min\{\frac{\gamma_1}{2\delta}x^2, \gamma_2 x\}.$$

Let

$$V = \{ \mathbf{v} \in H^1(\Omega; \mathbb{R}^2) : v_1 = 0 \text{ on } \Gamma_3 \}$$

and

$$K = \{ \mathbf{v} \in V : v_2 \ge 0 \text{ on } \Gamma_c \}$$




Fig. 2 A nonmonotone friction law

be the convex set of all admissible displacements. The weak formulation of this contact problem leads to the following hemivariational inequality: Find  $\mathbf{u} \in V$  such that for all  $\mathbf{v} \in V$ 

$$a(\mathbf{u}, \mathbf{v} - \mathbf{u}) + \int_{\Gamma_r} j^0(u_1(s); v_1(s) - u_1(s)) \, ds \ge \langle \mathbf{g}, \mathbf{v} - \mathbf{u} \rangle. \tag{14}$$

Here,  $a(\mathbf{u}, \mathbf{v})$  is the energy bilinear form of linear elasticity

$$a(\mathbf{u}, \mathbf{v}) = \int_{\Omega} \sigma_{ij}(\mathbf{u}) \varepsilon_{ij}(\mathbf{u}) dx \quad \mathbf{u}, \mathbf{v} \in V$$

with  $\sigma$ ,  $\varepsilon$  related by means of (13) and the linear form  $\langle \mathbf{g}, \cdot \rangle$  defined by

$$\langle \mathbf{g}, \mathbf{v} \rangle = \pm P \int_{\Gamma_1} v_1 \, ds - Q \int_{\Gamma_2} v_2 \, ds.$$

From Theorem 2.1, we obtain the existence of at least one solution provided that

$$\pm P \int_{\Gamma_1} a_1 - bx_2(s) \, ds - Q \int_{\Gamma_2} a_2 + bx_1(s) \, ds < -\gamma_2 \operatorname{meas} (\Gamma_c)^{1/2}$$

for all  $a_1, a_2, b \in \mathbb{R}$  satisfying  $\int_{\Gamma_3} a_1 - bx_2(s) ds = 0$  and  $\int_{\Gamma_s} a_2 + bx_1(s) ds \ge 0$ .

Note that now we use the decomposition of  $\mathbf{u}$  into normal and tangental parts and therefore,  $\gamma_0 = 1$  in (10).

# 5.2 Regularization and discretization

We solve this problem numerically following the method presented in Section 4 by first regularizing the hemivariational inequality (14) and then discretizing the regularized problem by the finite element method. This procedure leads to a smooth optimization problem that can be finally solved by using global minimization algorithms like trust region methods.

For this purpose we fix  $\varepsilon$  and use  $S : \mathbb{R} \times \mathbb{R}_{++} \to \mathbb{R}$  defined by

$$S(x,\varepsilon) := \begin{cases} g_1(x) & \text{if } (i) \text{ holds} \\ \frac{1}{2\varepsilon} [g(x) - g_1(x)]^2 + \frac{1}{2} (g(x) + g_1(x)) + \frac{\varepsilon}{8} & \text{if } (ii) \text{ holds} \\ g_2(x) & \text{if } (iii) \text{ holds}, \end{cases}$$

to approximate the maximum function  $-j(x) = \max\{-\frac{\gamma_1}{2\delta}x^2, -\gamma_2x\}$ . The cases (i), (ii), (iii) are defined below, respectively, by

(i) 
$$g_2(x) - g_1(x) \le -\frac{\varepsilon}{2}$$

(ii) 
$$-\frac{\varepsilon}{2} \le g_2(x) - g_1(x) \le \frac{\varepsilon}{2}$$

(iii) 
$$g_2(x) - g_1(x) \ge \frac{\varepsilon}{2}$$
.

Let  $\{\mathscr{T}_h\}$  be a regular triangulation of  $\Omega$  and  $\{x_i\}$  be the set of all vertices of the triangles of  $\{\mathscr{T}_h\}$ . We use continuous piecewise linear functions to approximate the displacements. Thus, V and K are approximated, respectively, by

$$V_h = \{ v_h \in C(\overline{\Omega}; \mathbb{R}^2) : v_{h|_T} \in (\mathbb{P}_1)^2, \ \forall T \in \mathcal{T}_h, \ v_{h1}(x_i) = 0 \ \forall x_i \in \overline{\Gamma}_3 \}$$
$$K_h = \{ v_h \in V_h : v_{h2}(x_i) \ge 0 \ \forall x_i \in \overline{\Gamma}_c \}.$$

The approximation of (14) now reads as follows: Find  $u_h \in V_h$  such that

$$a(u_h, v_h - u_h) + \langle -DJ_h(u_h), v_h - u_h \rangle \ge P \int_{\Gamma_E^1} (v_{h1} - u_{h1}) dx_2 \quad \forall v_h \in K_h,$$

where

$$\langle DJ_h(u_h), v_h \rangle = \frac{1}{2} \sum |P_i P_{i+1}| \left[ \frac{\partial S}{\partial x} (u_{h1}(P_i), \varepsilon) v_{h1}(P_i) + \frac{\partial S}{\partial x} (u_{h1}(P_{i+1}), \varepsilon) v_{h1}(P_{i+1}) \right]. \quad (15)$$

The discretized regularized problem (15) is put to work using the following steps. First, we use a condensation technique based on a Schur complement to reduce the total number of unknowns in (15) and pass to a finite-dimensional variational inequality problem formulated only in terms of the contact displacements. The obtained problem is re-written as a mixed complementarity problem, which is further reformulated as a system of nonlinear equations by means of the Fischer-Burmeister function  $f(a,b) = \sqrt{a^2 + b^2} - (a+b)$ . Finally, we apply an appropriate merit function and obtain an equivalent smooth, unconstrained minimization problem, which is numerically solved by using *lsqnonlin* - MATLAB function based on trust region method. The maximal number of iteration in *lsqnonlin* has been fixed to 100. The regularization parameter  $\varepsilon$  is set to 0.1. The tangential component  $u_1$  along  $\Gamma_c$  for the two models (see Fig. 1) and four different mesh sizes h = 1/4, 1/8, 1/16 and 1/32 in [m] is captured in Fig. 4 (a), respectively, Fig. 5 (b). Fig. 3 illustrates the computed complete displacement field on the whole boundary  $\Gamma$  for same mesh sizes.


#### References

- 1. Demyanov, V.F.; Rubinov, A.M.: Constructive nonsmooth analysis. Approximation & Optimization, 7. Peter Lang, Frankfurt am Main, 1995.
- Demyanov, V.F.; Stavroulakis, G.E.; Polyakova, L.N.; Panagiotopoulos, P.D.: Quasidifferentiability
  and nonsmooth modelling in mechanics, engineering and economics. Nonconvex Optimization and its
  Applications, 10. Kluwer Academic Publishers, Dordrecht, 1996.
- Panagiotopoulos, P.D.: Hemivariational inequalities. Applications in mechanics and engineering, Berlin, Springer, 1993.
- Naniewicz, Z.; Panagiotopoulos, P. D.: Mathematical Theory of Hemivariational Inequalities and Applications, Marcel Dekker, New York, 1995.
- Panagiotopoulos, P.D.: Inequality problems in mechanics and application. Convex and nonconvex energy functions, Basel, Birkhäuser, 1998.


 Goeleven, D., Motreanu, D., Dumont, Y., Rochdi, M.: Variational and Hemivariational Inequalities: Theory, Methods and Applications, Vol. II: Unilateral problems, Kluwer, Dordrecht, 2003.

- Baniotopoulos, C.C.; Haslinger, J.; Morávková, Z.: Contact problems with nonmonotone friction: discretization and numerical realization, Comput. Mech. Vol. 40, 157-165 (2007)
- 8. Attouch, H.; Buttazzo, G.; Michaille, G.: Variational Analysis in Sobolev and BV Spaces, MPS-SIAM Series on Optimization. SIAM, Philadelphia, 2006.
- 9. Penot, J.-P.: Npncoercive problems and asymptotic conditions, Asymptot. Anal. 49 (2006), 205 215.
- Brézis, H.: Equations et inequations non linéaires dans les espaces vectoriels en dualité, Ann. Inst. Fourier 18 (1968), 115 – 175.
- 11. Panagiotopoulos, P. D.: Coercive and semicoercive hemivariational inequalities. Nonlinear Anal. 16 (1991), no. 3, 209 231.
- Goeleven, D.; Théra, M.: Semicoercive variational hemivariational inequalities. J. Global Optim. 6 (1995), no. 4, 367 – 381.
- Adly, S.; Goeleven, D.; Théra, M.: Recession mappings and noncoercive variational inequalities, Nonlinear Anal. Series A, 26 (1996), no. 9, 1573–1603.
- Goeleven, D.: On Noncoercive Variational Problems and Related Results, Pitman Research Notes in Mathematics Series 357, Longman, Harlow, 1996.
- 15. Dinca, G.; Panagiotopoulos, P.D.; Pop, G.: An existence result on noncoercive hemivariational inequalities. Ann. Fac. Sci. Toulouse Math. (6) 6 (1997), no. 4, 609 632.
- Gwinner, J.: A note on pseudomonotone functions, regularization, and relaxed coerciveness, Nonlinear Anal. 30 (1997), 4217 – 4227.
- 17. Chadli, O.; Chbani, Z.; Riahi, H.: Some existence results for coercive and noncoercive hemivariational inequalities, Appl. Anal. 69 (1998), no. 1-2, 125 131.
- Chadli, O.; Chbani, Z.; Riahi, H.: Recession methods for equilibrium problems and applications to variational and hemivariational inequalities, Discrete Contin. Dynam. Systems 5 (1999), no. 1, 185 – 196
- Naniewicz, Z.: Semicoercive variational-hemivariational inequalities with unilateral growth conditions,
   J. Global Optim. 17 (2000), no. 1-4, 317 337.
- Adly, S.; Théra, M.; Ernst, E.: Stability of the solution set of non-coercive variational inequalities, Commun. Contemp. Math. 4 (2002), no. 1, 145–160.
- Liu, Zhen-Hai: Elliptic variational hemivariational inequalities, Appl. Math. Lett. 16 (2003), no. 6, 871– 876
- 22. Chadli, O.; Schaible, S. ; Yao, J. C.: Regularized equilibrium problems with application to noncoercive hemivariational inequalities, J. Optim. Theory Appl. 121 (2004), no. 3, 571 596.
- 23. Adly, S.: Stability of linear semi-coercive variational inequalities in Hilbert spaces: application to the Signorini-Fichera problem, J. Nonlinear Convex Anal. 7 (2006), no. 3, 325 334.
- 24. Chadli, O.; Liu, Z.; Yao, J. C.: Applications of equilibrium problems to a class of noncoercive variational inequalities, J. Optim. Theory Appl. 132 (2007), no. 1, 89 110.
- Costea, N.; Radulescu, V. D.: Hartman-Stampacchia results for stably pseudomonotone operators and non-linear hemivariational inequalities. Appl. Anal. 89 (2010), no. 2, 175 – 188.
- Costea, N.; Matei, A.: Contact models leading to variational-hemivariational inequalities. J. Math. Anal. Appl. 386 (2012), no. 2, 647 – 660.
- 27. Tang, Guo-ji; Huang, Nan-jing: Existence theorems of the variational-hemivariational inequalities, J. Global Optim. 56 (2013), no. 2, 605 622.
- 28. Lahmdan, A.; Chadli, O.; Yao, J. C.: Existence of solutions for noncoercive hemivariational inequalities by an equilibrium approach under pseudomonotone perturbation. J. Optim. Theory Appl. 160 (2014), no. 1, 49 66.
- Haslinger, J., Miettinen, M., Panagiotopoulos, P.D.: Finite Element Methods for Hemivariational Inequalities. Kluwer Academic Publishers. Dordrecht. 1999.
- Hintermüller, M.; Kovtunenko, V.A.; Kunisch K.: Obstacle Problems with Cohesion: A Hemivariational Inequality Approach and Its Efficient Numerical Solution, SIAM J. Optim. 21 (2011), no. 2, 491 - 516.
- 31. Kaplan, A.; Tichatschke, R.: Stable Methods for Ill-posed Variational Problems: Prox-Regularization of Elliptic Variational Inequalities and Semi-infinite Problems, Akademie-Verlag, Berlin, 1994.
- 32. Gwinner, J.: Discretization of semicoercive variational inequalities, Aequationes Math. 42 (1991), no. 1, 72 79
- Gwinner, J.: A Discretization Theory for Monotone Semicoercive Problems and Finite Element Convergence for p-Harmonic Signorini Problems, ZAMM J. of Applied Mathematics and Mechanics 74 (1984), no. 9, 417 427.
- 34. Spann, W:: Error estimates for the approximation of semicoercive variational inequalities. Numer. Math. 69 (1994), no. 1, 103 116.

- 35. Dostál, Z.; Friedlander, A.; Santos, S.A.: Solution of coercive and semicoercive contact problems by FETI domain decomposition, Domain decomposition methods, 10 (Boulder, CO, 1997), Contemp. Math. 218, 82–93, Amer. Math. Soc., Providence, 1998.
- 36. Adly, S.; Goeleven, D.: A discretization theory for a class of semi-coercive unilateral problems, Numer. Math. 87 (2000), no. 1, 1–34.
- Dostál, Z.; Horák, D.: Scalable FETI with optimal dual penalty for semicoercive variational inequalities, Current trends in scientific computing (Xi'an, 2002), Contemp. Math. 329, 79–88, Amer. Math. Soc., Providence, 2003.
- 38. Namm, R.V.; Woo, Gyungsoo; Xie, Shu-Sen; Yi, Sucheol: Solution of semicoercive Signorini problem based on a duality scheme with modified Lagrangian functional, J. Korean Math. Soc. 49 (2012), no. 4, 843–854.
- Gwinner, J.; Ovcharova, N.: From solvability and approximation of variational inequalities to solution of nondifferentiable optimization problems in contact mechanics, Optimization 64 (2015), no. 8, 1683 – 1702
- Ovcharova, N.: Regularization Methods and Finite Element Approximation of Hemivariational Inequalities with Applications to Nonmonotone Contact Problems, PhD Thesis, Universität der Bundeswehr München, Cuvillier Verlag, Göttingen, 2012.
- 41. Ovcharova, N., Gwinner J.: A study of regularization techniques of nondifferentiable optimization in view of application to hemivariational inequalities, J. Optim. Theory Appl. 162 (2014), no. 3, 754-778.
- 42. Adly, Samir; Théra, Michel; Ernst, Emil: Well-positioned closed convex sets and well-positioned closed convex functions, J. Global Optim. 29 (2004), no. 4, 337–351.
- Glowinski, R: Numerical Methods for Nonlinear Variational Problems, Reprint of the 1984 original, Springer-Verlag, Berlin, 2008.
- 44. Gwinner, J.: Nichtlineare Variationsungleichungen mit Anwendungen, PhD Thesis, Universität Mannheim, Haag + Herchen Verlag, Frankfurt, 1978.
- 45. Jeggle, H.: Nichtlineare Funktionalanalysis, Teubner Stuttgart, 1979.
- 46. Goeleven, D.; Gwinner, J.: On semicoerciveness, a class of variational inequalities, and an application to von Kármán plates, Math. Nachr. 244 (2002), 89–109.
- 47. Stampacchia, G.: Variational inequalities. In: Theory and Applications of Monotone Operators, Edizione Oderisi, Gubbio, 101-192 (1969)
- 48. Hess, P.: On semicoercive nonlinear problems, Indiana Univ. Math. J. 23 (1974), 645-654.
- Schatzman, M.: Problèmes aux limites non linéaires, non coercifs, Ann. Scuola Norm. Sup. Pisa (3) 27 (1974), 641–686.
- 50. Clarke, F: Optimization and Nonsmooth Analysis, John Wiley, New York, 1983.
- 51. Gwinner, J.: *hp*-FEM convergence for unilateral contact problems with Tresca friction in plane linear elastostatics, J. Comput. Appl. Math. 254 (2013), 175 184.
- Facchinei, F., Pang, J-S.: Finite-Dimensional Variational Inequalities and Complementarity Problems, Vol. I, Vol. II, Springer, New York, 2003.



(a) Reference and deformed configurations in cm corresponding to Fig. 1 (a)



(b) Reference and deformed configurations in cm corresponding to Fig. 1 (b)

Fig. 3 The complete displacement field on the whole boundary  $\Gamma$ 




Fig. 4 Wall left: Left image shows the tangential component  $u_1$  on  $\Gamma_c$  for 4 discretization parameters h=1/4 (red), h=1/8 (green), h=1/16 (dark blue), h=1/32 (light blue) in [m]. The right image shows the distribution of the tangential stress  $-\sigma_t$  along  $\Gamma_c$  for the same 4 scenarios




Fig. 5 Wall right: Left image shows the tangential component  $u_1$  on  $\Gamma_c$  for 4 discretization parameters h=1/4 (red), h=1/8 (green), h=1/16 (dark blue), h=1/32 (light blue) in [m]. The right image shows the distribution of the tangential stress  $-\sigma_t$  along  $\Gamma_c$  for the same 4 scenarios