

# A NOTE ON MAXIMAL SUBGROUPS OF FREE IDEMPOTENT GENERATED SEMIGROUPS OVER BANDS

IGOR DOLINKA

Department of Mathematics and Informatics, University of Novi Sad,  
Trg Dositeja Obradovića 4, 21101 Novi Sad, Serbia  
E-mail: dockie@dmf.uns.ac.rs

## Abstract

We prove that all maximal subgroups of the free idempotent generated semigroup over a band  $B$  are free for all  $B$  belonging to a band variety  $\mathbf{V}$  if and only if  $\mathbf{V}$  consists either of left seminormal bands, or of right seminormal bands.

Let  $S$  be a semigroup, and let  $E = E(S)$  be the set of its idempotents; in fact,  $E$ , along with the multiplication inherited from  $S$ , is a partial algebra. It turns out to be fruitful to restrict further the domain of the partial multiplication defined on  $E$  by considering only the pairs  $e, f \in E$  for which either  $ef \in \{e, f\}$  or  $fe \in \{e, f\}$  (i.e.  $\{ef, fe\} \cap \{e, f\} \neq \emptyset$ ). Note that if  $ef \in \{e, f\}$  then  $fe$  is an idempotent, and the same is true if we interchange the roles of  $e$  and  $f$ . Such unordered pairs  $\{e, f\}$  are called *basic pairs* and their products  $ef$  and  $fe$  are *basic products*.

The *free idempotent generated semigroup over  $E$*  is defined by the following presentation:

$$\text{IG}(E) = \langle E \mid e \cdot f = ef \text{ such that } \{e, f\} \text{ is a basic pair} \rangle.$$

Here  $ef$  denotes the product of  $e$  and  $f$  in  $S$  (which is again an idempotent of  $S$ ), while  $\cdot$  stands for the concatenation operation in the free semigroup  $E^+$  (also to be interpreted as the multiplication in its quotient  $\text{IG}(E)$ ). An important feature

---

*Mathematics subject classification numbers:* 20M05, 20M10, 20F05

*Key words and phrases:* free idempotent generated semigroup, band, maximal subgroup

The support of the Ministry of Education and Science of the Republic of Serbia, through Grant No. 174019, is gratefully acknowledged.

of  $\text{IG}(E)$  is that there is a natural homomorphism from  $\text{IG}(E)$  onto the subsemigroup of  $S$  generated by  $E$ , and the restriction of  $\phi$  to the set of idempotents of  $\text{IG}(E)$  is a basic-product-preserving bijection onto  $E$ , see e.g. [5, 9, 13].

An important background to these definitions is the notion of the *biordered set* [7] of idempotents of a semigroup and its abstract counterpart. The biordered set of idempotents of  $S$  is just a partial algebra on  $E(S)$  obtained by restricting the multiplication from  $S$  to basic pairs of idempotents. In this way we have that if  $B$  is a band (an idempotent semigroup), then, even though there is an everywhere defined multiplication on  $E(B) = B$ , its biordered set [3] is in general still a partial algebra. Another way of treating biordered sets is to consider them as relational structures  $(E(S), \leq^{(l)}, \leq^{(r)})$ , where the set of idempotents  $E(S)$  is equipped by two quasi-order relations defined by

$$\begin{aligned} e &\leq^{(l)} f \text{ if and only if } ef = e, \\ e &\leq^{(r)} f \text{ if and only if } fe = e. \end{aligned}$$

One of the main achievements of [4, 5, 9] is the result that the class of biordered sets considered as relational structures is *axiomatisable*: there is in fact a finite system of formulæ satisfied by biordered sets such that any set endowed with two quasi-orders satisfying the axioms in question is a biordered set of idempotents of some semigroup. In this sense we can speak about the free idempotent generated semigroup over a biordered set  $E$ . A fundamental fact which justifies the term ‘free’ is that  $\text{IG}(E)$  is the free object in the category of all semigroups  $S$  whose biordered set of idempotents is isomorphic to  $E$ : if  $\psi : E \rightarrow E(S)$  is any isomorphism of biordered sets, then it uniquely extends (via the canonical injection of  $E$  into  $\text{IG}(E)$ ) to a homomorphism  $\psi' : \text{IG}(E) \rightarrow S$  whose image is the subsemigroup of  $S$  generated by  $E(S)$ . This is also true if  $\psi$  is a (surjective) homomorphism of biordered sets (taken as relational structures), so that the freeness property of  $\text{IG}(E)$  carries over to even wider categories of semigroups.

In this short note we consider  $\text{IG}(B)$ , the free idempotent generated semigroup over (the biordered set of) a band  $B$ ; more precisely, we are interested in the question whether the maximal subgroups of these semigroups are free. It was conjectured in [8] that each maximal subgroup of any semigroup of the form  $\text{IG}(E)$  is a free group. Recently, this was disproved [1] (see also [2]), where a certain 72-element semigroup was found whose biordered set  $E$  of idempotents yields a maximal subgroup in  $\text{IG}(E)$  isomorphic to  $\mathbb{Z} \oplus \mathbb{Z}$ , the rank 2 free abelian group. Here we will see that a particular 20-element regular band suffices for the same purpose. In fact, as proved by Gray and Ruškuc in [6], *every* group can be isomorphic to a maximal subgroup of some  $\text{IG}(E)$ , while the assumption that the semigroup  $S$  with  $E = E(S)$  is finite yields a sole restriction that the groups

in question are finitely presented. This puts forward many new questions, one of which is the characterisation of bands  $B$  for which all subgroups of  $\text{IG}(B)$  are free.

More specifically, as a first approximation to the latter question, we may ask for a description of all varieties  $\mathbf{V}$  of bands with the property that for each  $B \in \mathbf{V}$  the maximal subgroups of  $\text{IG}(B)$  are free. To facilitate the discussion, we depict in Fig. 1 the bottom part of the lattice  $\mathcal{L}(\mathbf{B})$  of all band varieties, along with their standard labels (see also [15, Diagram II.3.1]).




Figure 1: The bottom part of the lattice of all varieties of bands

The main result of this note is the following.

**Theorem 1.** *Let  $\mathbf{V}$  be a variety of bands. Then  $\text{IG}(B)$  has all its maximal subgroups free for all  $B \in \mathbf{V}$  if and only if  $\mathbf{V}$  is contained either in **LSNB** or in **RSNB**.*

This theorem is a direct consequence of the following two propositions.

**Proposition 2.** *For any left (right) seminormal band  $B$ , all maximal subgroups of  $\text{IG}(B)$  are free.*

**Proposition 3.** *There exists a regular band  $B$  such that  $\text{IG}(B)$  has a maximal subgroup isomorphic to  $\mathbb{Z} \oplus \mathbb{Z}$ .*

The first of these propositions is a generalisation of the well known result of Pastijn [13, Theorem 6.5] (cf. also [10, 12]) that all maximal subgroups of

$\text{IG}(B)$  are free for any normal band  $B$ . The other one supplies a simpler example with the same non-free maximal subgroup than the one considered in [1, Section 5]. The method used is the one from [6], which is based on the Reidemeister-Schreier type rewriting process for obtaining presentations of maximal subgroups of semigroups developed in [11]. So, before turning to the proofs of the above two propositions, we briefly present this general method yielding presentations for maximal subgroups of  $\text{IG}(E)$ ,  $E = E(S)$ , for an arbitrary semigroup  $S$ , and then we explain its particular case when  $S$  is a band. Along the way, we assume some familiarity with the most basic notions of semigroup theory, such as Green's relations and the structure of bands, see, for example, [7, 15].

Let  $S$  be a semigroup and let  $D$  be a  $\mathcal{D}$ -class of  $S$  containing an idempotent  $e_0 \in E(S)$ . We are going to label the  $\mathcal{R}$ -classes contained in  $D$  by  $R_i$ ,  $i \in I$ , while  $L_j$ ,  $j \in J$ , is the list of all  $\mathcal{L}$ -classes of  $D$ . The  $\mathcal{H}$ -class  $R_i \cap L_j$  will be denoted by  $H_{ij}$ . Define  $\mathcal{K} = \{(i, j) : H_{ij} \text{ is a group}\}$ ; as is well known,  $(i, j) \in \mathcal{K}$  if and only if  $H_{ij}$  contains an idempotent, which we denote by  $e_{ij}$ . There is no loss of generality if we assume that both  $I$  and  $J$  contain an index 1, so that  $e_0 = e_{11}$ .

For a word  $\mathbf{w} \in E^*$ , let  $\overline{\mathbf{w}}$  denote the image of  $\mathbf{w}$  under the canonical monoid homomorphism of  $E^*$  into  $S^1$ : in other words, when  $\mathbf{w}$  is non-empty,  $\overline{\mathbf{w}}$  is just the element of  $S$  obtained by multiplying in  $S$  the idempotents the concatenation of which is  $\mathbf{w}$ . We say that a system of words  $\mathbf{r}_j, \mathbf{r}'_j \in E^*$ ,  $j \in J$ , is a *Schreier system of representatives* for  $D$  if for each  $j \in J$ :

- the right multiplications by  $\overline{\mathbf{r}_j}$  and  $\overline{\mathbf{r}'_j}$  are mutually inverse  $\mathcal{R}$ -class preserving bijections  $L_1 \rightarrow L_j$  and  $L_j \rightarrow L_1$ , respectively (so, in particular, right multiplication by  $\mathbf{r}_1$  is the identity mapping on  $L_1$ );
- each prefix of  $\mathbf{r}_j$  coincides with  $\mathbf{r}_{j'}$  for some  $j' \in J$  (in particular, the empty word is just  $\mathbf{r}_1$ ).

It is well-known that such a Schreier system always exists. In the following, we assume that one particular Schreier system has been fixed.

In addition, we will assume that a mapping  $i \mapsto j(i)$  has been specified such that  $(i, j(i)) \in \mathcal{K}$ : such  $j(i)$  must exist for each  $i \in I$ , since  $D$  is a regular  $\mathcal{D}$ -class (as it contains an idempotent), and so each  $\mathcal{R}$ -class  $R_i$  must contain an idempotent. The index  $j(i) \in J$  is called the *anchor* of  $R_i$ .

Finally, call a *square* a quadruple of idempotents  $(e, f, g, h)$  in  $D$  such that

$$\begin{array}{ccccc} e & \mathcal{R} & & f & \\ & & \mathcal{L} & & \mathcal{L} \\ g & \mathcal{R} & & h & . \end{array}$$

Then there are  $i, k \in I$  and  $j, \ell \in J$  such that  $e \in H_{ij}$ ,  $f \in H_{i\ell}$ ,  $g \in H_{kj}$  and  $h \in H_{k\ell}$ . For an idempotent  $\varepsilon \in S$  we say that it *singularises* the square  $(e, f, g, h)$  if any of the following two cases takes place:

- (a)  $\varepsilon e = e$  and  $\varepsilon g = g$ , while  $e = f\varepsilon$ ; or
- (b)  $e = \varepsilon g$ , along with  $e\varepsilon = e$  and  $f\varepsilon = f$ .

Note that case (a) implies  $\varepsilon f = f$ ,  $\varepsilon h = h$ ,  $e\varepsilon = e$  and  $g = g\varepsilon = h\varepsilon$ , while conditions  $\varepsilon e = e$ ,  $f = \varepsilon f = \varepsilon h$ ,  $g\varepsilon = g$  and  $h\varepsilon = h$  follow from (b). The square  $(e, f, g, h)$  is *singular* if it is singularised by some idempotent of  $S$ . Let  $\Sigma$  be the set of all quadruples  $(i, k; j, \ell) \in I \times I \times J \times J$  (to be called *singular rectangles*) such that  $(e_{ij}, e_{i\ell}, e_{kj}, e_{k\ell})$  is a singular square in  $D$ .

The required general result of [6] can be now paraphrased as follows.

**Theorem 4** (Theorem 5 of [6]). *Let  $S$  be a semigroup with a non-empty set of idempotents  $E = E(S)$ . With the notation as above, the maximal subgroup of the free idempotent generated semigroup  $\text{IG}(E)$  containing  $e_{11} \in E$  is presented by  $\langle \Gamma \mid \mathfrak{R} \rangle$ , where  $\Gamma = \{f_{ij} : (i, j) \in \mathcal{K}\}$ , while  $\mathfrak{R}$  consists of three types of relations:*

- (i)  $f_{i,j(i)} = 1$  for all  $i \in I$ ;
- (ii)  $f_{ij} = f_{i\ell}$  for all  $i \in I$  and  $j, \ell \in J$  such that  $\mathbf{r}_j \cdot e_{i\ell} = \mathbf{r}_\ell$ ;
- (iii)  $f_{ij}^{-1} f_{i\ell} = f_{kj}^{-1} f_{k\ell}$  for all  $(i, k; j, \ell) \in \Sigma$ .

For our purpose, we would like to focus on the particular case when  $S$  is a band. Then, clearly,  $\mathcal{K} = I \times J$  and  $D = \{e_{ij} : i \in I, j \in J\}$ . Since  $\mathcal{D} = \mathcal{J}$  in any band, the set of all  $\mathcal{D}$ -classes of  $B$  is partially ordered; it instantly turns out that, by definition, if  $\varepsilon$  singularises a square  $(e, f, g, h)$  in  $D$ , then  $D_\varepsilon \geq D$ . Now any such  $\varepsilon \in B$  induces a pair of transformations on  $I$  and  $J$ , respectively, in the following sense. For each  $i \in I$  and  $j \in J$  there are  $i', k \in I$  and  $j', \ell \in J$  such that  $\varepsilon e_{ij} = e_{i'\ell}$  and  $e_{ij}\varepsilon = e_{kj'}$ . One immediately sees that it must be  $\ell = j$  and  $k = i$ , so that  $B$  acts on the left on  $I$  and on the right on  $J$ . Thus it is convenient to write the transformation  $\sigma = \sigma_\varepsilon^{(l)}$  induced by  $\varepsilon$  on  $I$  to the left of its argument (so that  $ee_{ij} = e_{\sigma(i)j}$ ), while the analogous transformation  $\sigma' = \sigma_\varepsilon^{(r)}$  on  $J$  is written to the right (resulting in the rule  $e_{ij}e = e_{i(j)\sigma'}$ ).

**Corollary 5.** *Let  $B$  be a band, let  $D$  be a  $\mathcal{D}$ -class of  $B$ , and let  $e_{11} \in D$ . Then the maximal subgroup  $G_{e_{11}}$  of  $\text{IG}(B)$  containing  $e_{11}$  is presented by  $\langle \Gamma \mid \mathfrak{R} \rangle$ , where  $\Gamma = \{f_{ij} : i \in I, j \in J\}$  and  $\mathfrak{R}$  consists of relations*

$$f_{i1} = f_{1j} = f_{11} = 1 \tag{1}$$

for all  $i \in I$  and  $j \in J$ , and

$$f_{ij}^{-1} f_{i\ell} = f_{kj}^{-1} f_{k\ell}, \quad (2)$$

where for some  $\varepsilon \in B$  such that  $D_\varepsilon \geq D$  the indices  $i, k \in I$ ,  $j, \ell \in J$  satisfy one of the following two conditions:

- (a)  $\sigma_\varepsilon^{(l)}(i) = i$ ,  $\sigma_\varepsilon^{(l)}(k) = k$ , and  $(j)\sigma_\varepsilon^{(r)} = (\ell)\sigma_\varepsilon^{(r)} = \ell$ ,
- (b)  $\sigma_\varepsilon^{(l)}(i) = \sigma_\varepsilon^{(l)}(k) = k$ ,  $(j)\sigma_\varepsilon^{(r)} = j$  and  $(\ell)\sigma_\varepsilon^{(r)} = \ell$ .

*Proof.* Since  $\mathcal{K} = I \times J$ , we have a generator  $f_{ij}$  for each  $i \in I$  and  $j \in J$ . Furthermore, the same reason allows us to choose  $j(i) = 1$  as the anchor for each  $i \in I$ . Such a choice will imply that the relations of type (i) from Theorem 4 take the form  $f_{i1} = 1$ ,  $i \in I$ . In particular, we have  $f_{11} = 1$ . As for the Schreier system, we can choose  $\mathbf{r}_1$  to be the empty word,  $\mathbf{r}_j = e_{1j}$  for all  $j \in J \setminus \{1\}$  and  $\mathbf{r}'_j = e_{11}$  for all  $j \in J$ . The system  $\mathbf{r}_j$ ,  $j \in J$ , of words over  $E$  is obviously prefix-closed. Since  $e_{i1}e_{ij} = e_{ij}$  and  $e_{ij}e_{11} = e_{i1}$  holds for all  $i \in I$ ,  $j \in J$ , the right multiplications by  $e_{ij}$  and  $e_{11}$  are indeed mutually inverse bijections between  $L_1$  and  $L_j$  and between  $L_j$  and  $L_1$ , respectively. Hence, the relations of type (ii) reduce to  $f_{11} = f_{1j}$ , that is,  $f_{1j} = 1$ , for all  $j \in J$ . Thus we have all the relations (1). Finally, the conditions (a) and (b) express precisely the singularisation of a square  $(e_{ij}, e_{i\ell}, e_{kj}, e_{k\ell})$  in  $D$  by an element  $\varepsilon \in B$ ; therefore, the relations (2) correspond to relations of type (iii).  $\square$

Rectangles  $(i, k; j, \ell) \in I \times J$  of type (a) will be said to be *left-right* singular, while those of type (b) are *up-down* singular (with respect to  $\varepsilon$ ). Another, more compact way of expressing condition (a) is  $i, k \in \text{Im } \sigma_\varepsilon^{(l)}$ ,  $\ell \in \text{Im } \sigma_\varepsilon^{(r)}$  and  $(j, \ell) \in \text{Ker } \sigma_\varepsilon^{(r)}$ , while (b) is equivalent to  $k \in \text{Im } \sigma_\varepsilon^{(l)}$ ,  $(i, k) \in \text{Ker } \sigma_\varepsilon^{(l)}$  and  $j, \ell \in \text{Im } \sigma_\varepsilon^{(r)}$ .

We can now turn to proving our aforementioned result.

*Proof of Proposition 2.* Without any loss of generality, assume that  $B \in \mathbf{RSNB}$  (the case when  $B$  belongs to  $\mathbf{LSNB}$  is dual). Recall (e.g. from [15, Proposition II.3.8]) that the variety  $\mathbf{RSNB}$  satisfies (and is indeed defined by) the identity  $tuv = tvtuv$ . Therefore, if  $B = \bigcup_{\alpha \in Y} B_\alpha$  is the greatest semilattice decomposition of  $B$ ,  $a \in B$  and  $x, y \in D = B_\alpha$  for some  $\alpha \in Y$ , then  $x = xyx$  and  $y = yxy$ . Hence, we have  $ax = ax(yx) = ayxaxyx$  and  $ay = ay(xy) = axyayxy$ , implying  $ax\mathcal{R}ay$ . In particular, for any  $\varepsilon \in B$  such that  $D_\varepsilon \geq D$ ,  $\varepsilon e_{ij}\mathcal{R}\varepsilon e_{k\ell}$  holds in  $D$  for all  $i, k \in I$ ,  $j, \ell \in J$ , so the transformation  $\sigma_\varepsilon^{(l)}$  is a constant function on  $I$ .

We conclude that there are no proper (non-degenerate) rectangles  $(i, k; j, \ell)$  that are left-right singular with respect to some  $\varepsilon \in B$ . In other words, all proper

singular rectangles in  $I \times J$ —and thus all nontrivial relations of  $G_{e_{11}}$ —are of the up-down kind:

$$f_{ij}^{-1}f_{i\ell} = f_{k_0j}^{-1}f_{k_0\ell},$$

where  $j, \ell$  are two fixed points of  $\sigma_\varepsilon^{(r)}$ ,  $i \in I$  is arbitrary, and (since in this context  $\sigma_\varepsilon^{(l)}$  is constant)  $\text{Im } \sigma_\varepsilon^{(l)} = \{k_0\}$ , for some  $\varepsilon \in B$ . However, now it is straightforward to deduce the relation (2) for *all*  $i, k \in I$  and fixed points  $j, \ell$  of  $\sigma_\varepsilon^{(r)}$ . Thus we are led to define an equivalence  $\theta_B$  of  $\bigcup_{\varepsilon \in B, D_\varepsilon \geq D} \text{Im } \sigma_\varepsilon^{(r)} = J$  which is the transitive closure of the relation  $\rho_B$  defined by  $(j_1, j_2) \in \rho_B$  if and only if  $j_1, j_2 \in \text{Im } \sigma_\varepsilon^{(r)}$  for some  $\varepsilon \in B$ . Now it is almost immediate to see that for all  $i, k \in I$  and  $j, \ell \in J$  such that  $(j, \ell) \in \theta_B$  we have that

$$f_{ij}^{-1}f_{i\ell} = f_{kj}^{-1}f_{k\ell}$$

holds in  $G_{e_{11}}$ . This immediately implies  $f_{k\ell} = 1$  for all  $k \in I$  and  $\ell \in 1/\theta_B$ , as well as

$$f_{kj} = f_{k\ell}$$

for all  $k \in I$ , whenever  $(j, \ell) \in \theta_B$ . So, let  $j_1 = 1, j_2, \dots, j_m \in J$  be a cross-section of  $J/\theta_B$ . Then it is straightforward to eliminate all the relations from the presentation of  $G_{e_{11}}$  while reducing its generating set to

$$\{f_{ij_r} : i \in I \setminus \{1\}, 2 \leq r \leq m\}.$$

In other words,  $G_{e_{11}}$  is a free group of rank  $(|I| - 1)(m - 1)$ .  $\square$

*Proof of Proposition 3.* Let  $B$  be the subband of the free regular band on four generators  $a, b, c, d$  consisting of two  $\mathcal{D}$ -classes: a  $2 \times 2$  class  $D_1$  consisting of elements  $ab, aba, ba, bab$  and a  $4 \times 4$  class  $D_0$  consisting of elements of the form  $\mathbf{u}_1\mathbf{v}\mathbf{u}_2$ , where  $\mathbf{u}_1, \mathbf{u}_2 \in \{ab, ba\}$  and  $\mathbf{v} \in \{cd, cdc, dc, dcd\}$ . So, we can take  $I = \{abcd, abdc, bacd, badc\}$ , the set of all initial parts of words from  $D_0$ , and  $J = \{cdba, dcba, cdab, dcab\}$ , the set of all final parts of those words. A direct computation shows that

$$\begin{aligned} \sigma_{ab}^{(l)} &= \sigma_{aba}^{(l)} = \begin{pmatrix} abcd & abdc & badc & bacd \\ abcd & abdc & abdc & abcd \end{pmatrix}, \\ \sigma_{ba}^{(l)} &= \sigma_{bab}^{(l)} = \begin{pmatrix} abcd & abdc & badc & bacd \\ bacd & badc & badc & bacd \end{pmatrix}, \\ \sigma_{ab}^{(r)} &= \sigma_{bab}^{(r)} = \begin{pmatrix} cdba & cdab & dcab & dcba \\ cdab & cdab & dcab & dcab \end{pmatrix}, \\ \sigma_{ba}^{(r)} &= \sigma_{aba}^{(r)} = \begin{pmatrix} cdba & cdab & dcab & dcba \\ cdba & cdba & dcba & dcba \end{pmatrix}. \end{aligned}$$

If we enumerate (for brevity of further calculations)  $abcd \rightarrow 1, abdc \rightarrow 2, badc \rightarrow 3, bacd \rightarrow 4$  and  $cdab \rightarrow 1, cdab \rightarrow 2, dcab \rightarrow 3, dcba \rightarrow 4$ , we get

$$\begin{aligned}\sigma_{ab}^{(l)} = \sigma_{aba}^{(l)} &= \begin{pmatrix} 1 & 2 & 3 & 4 \\ 1 & 2 & 2 & 1 \end{pmatrix}, & \sigma_{ba}^{(l)} = \sigma_{bab}^{(l)} &= \begin{pmatrix} 1 & 2 & 3 & 4 \\ 4 & 3 & 3 & 4 \end{pmatrix}, \\ \sigma_{ab}^{(r)} = \sigma_{bab}^{(r)} &= \begin{pmatrix} 1 & 2 & 3 & 4 \\ 2 & 2 & 3 & 3 \end{pmatrix}, & \sigma_{ba}^{(r)} = \sigma_{aba}^{(r)} &= \begin{pmatrix} 1 & 2 & 3 & 4 \\ 1 & 1 & 4 & 4 \end{pmatrix}.\end{aligned}$$

Hence, the list of singular rectangles is exhausted by:

$$\begin{aligned}(1, 2; 1, 2), (1, 2; 3, 4), (3, 4; 1, 2), (3, 4; 3, 4), \\ (1, 4; 2, 3), (1, 4; 1, 4), (2, 3; 2, 3), (2, 3; 1, 4).\end{aligned}$$

This results in  $f_{11} = f_{12} = f_{13} = f_{14} = f_{21} = f_{31} = f_{41} = f_{22} = f_{44} = 1$  and

$$\begin{aligned}f_{23} = f_{24}, \quad f_{24} = f_{34}, \quad f_{43}^{-1} = f_{33}^{-1} f_{34} \\ f_{32} = f_{42}, \quad f_{42} = f_{43}, \quad f_{23} = f_{32}^{-1} f_{33}.\end{aligned}$$




Figure 2: The rectangles in  $D_0$  singularised by elements of  $D_1$

If we denote  $x = f_{23}$  and  $y = f_{32}$  we obviously remain with these two generators for  $G_{abcdba}$  and a single relation

$$yx = f_{33} = xy,$$

so  $G_{abcdba} \cong \mathbb{Z} \oplus \mathbb{Z}$ . □

This completes the proof of Theorem 1.

**Remark 6.** The band  $B$  from the previous proof can be also realised as a regular subband of the free band  $FB_3$  on three generators  $a, b, c$  whose elements are from  $D'_1 = \{ab, aba, ba, bab\}$  and  $D'_0 = \{\mathbf{u}\mathbf{c}\mathbf{v} : \mathbf{u}, \mathbf{v} \in D'_1\}$ .



We finish the note by several problems that might be subjects of future research in this direction.

**Problem 1.** Characterise all bands  $B$  with the property that  $\text{IG}(B)$  has a non-free maximal subgroup.

**Problem 2.** Characterise all groups that arise as maximal subgroups of  $\text{IG}(B)$  for some band  $B$ . The same problem stands for regular bands  $B$ , and in fact for  $B \in \mathbf{V}$  for any particular band variety  $\mathbf{V} \geq \mathbf{RB}$ .

**Problem 3.** Given a band variety  $\mathbf{V}$  and an integer  $n \geq 1$ , describe the maximal subgroups of  $\text{IG}(\mathfrak{F}_n \mathbf{V})$ , where  $\mathfrak{F}_n \mathbf{V}$  denotes the  $\mathbf{V}$ -free band on a set of  $n$  free generators [16].

**Acknowledgement.** The author is grateful to the anonymous referee, whose careful reading, comments and suggestions significantly improved the presentation of the results.

## References

- [1] M. BRITTENHAM, S. W. MARGOLIS and J. MEAKIN, Subgroups of free idempotent generated semigroups need not be free, *J. Algebra*, **321** (2009), 3026–3042.
- [2] M. BRITTENHAM, S. W. MARGOLIS and J. MEAKIN, Subgroups of free idempotent generated semigroups: full linear monoids, manuscript, 17 pp. [arXiv:1009.5683](#)
- [3] D. EASDOWN, Biordered sets of bands, *Semigroup Forum*, **29** (1984), 241–246.
- [4] D. EASDOWN, Biordered sets are biordered subsets of idempotents of semigroups, *J. Austral. Math. Soc. Ser. A*, **37** (1984), 258–268.
- [5] D. EASDOWN, Biordered sets come from semigroups, *J. Algebra*, **96** (1985), 581–591.
- [6] R. GRAY and N. RUŠKUC, On maximal subgroups of free idempotent generated semigroups, *Israel J. Math.*, to appear.
- [7] P. M. HIGGINS, *Techniques of Semigroup Theory*, Oxford University Press, Oxford, 1992.
- [8] B. MCELWEE, Subgroups of the free semigroup on a biordered set in which principal ideals are singletons, *Comm. Algebra*, **30** (2002), 5513–5519.
- [9] K. S. S. NAMBOORIPAD, Structure of regular semigroups. I, *Mem. Amer. Math. Soc.*, **22** (1979), no. 224, vii+119 pp.
- [10] K. S. S. NAMBOORIPAD and F. PASTIJN, Subgroups of free idempotent generated regular semigroups, *Semigroup Forum*, **21** (1980), 1–7.
- [11] N. RUŠKUC, Presentations for subgroups of monoids, *J. Algebra*, **220** (1999), 365–380.

- [12] F. PASTIJN, Idempotent generated completely 0-simple semigroups, *Semigroup Forum*, **15** (1977), 41–50.
- [13] F. PASTIJN, The biorder on the partial groupoid of idempotents of a semigroup, *J. Algebra* **65** (1980), 147–187.
- [14] M. PETRICH, The translational hull in semigroups and rings, *Semigroup Forum*, **1** (1970), 283–360.
- [15] M. PETRICH, *Lectures in Semigroups*, Wiley, New York, 1977.
- [16] M. PETRICH and P. V. SILVA, Structure of relatively free bands, *Comm. Algebra*, **30** (2002), 4165–4187.