Gorenstein Projective Objects in Comma Categories*†

Yeyang Peng, Rongmin Zhu, Zhaoyong Huang[‡]

Department of Mathematics, Nanjing University, Nanjing 210093, Jiangsu Province, P. R. China

Abstract

Let \mathcal{A} and \mathcal{B} be abelian categories and $\mathbf{F}: \mathcal{A} \to \mathcal{B}$ an additive and right exact functor which is perfect, and let $(\mathbf{F}, \mathcal{B})$ be the left comma category. We give an equivalent characterization of Gorenstein projective objects in $(\mathbf{F}, \mathcal{B})$ in terms of Gorenstein projective objects in \mathcal{B} and \mathcal{A} . We prove that there exists a left recollement of the stable category of the subcategory of $(\mathbf{F}, \mathcal{B})$ consisting of Gorenstein projective objects modulo projectives relative to the same kind of stable categories in \mathcal{B} and \mathcal{A} . Moreover, this left recollement can be filled into a recollement when \mathcal{B} is Gorenstein and \mathbf{F} preserves projectives.

1 Introduction

As a generalization of finitely generated projective modules, Auslander and Bridger [2] introduced finitely generated modules of Gorenstein dimension zero over a commutative noetherian local ring. Then Enochs and Jenda [6] generalized it to Gorenstein projective modules (not necessarily finitely generated) over an arbitrary ring. The properties of Gorenstein projective modules and related modules have been studied widely, see [1, 2, 5–7, 14–16] and references therein.

Let Λ and Γ be arbitrary rings and M a (finitely generated) (Λ, Γ) -bimodule, and let $T := \begin{pmatrix} \Lambda & M \\ 0 & \Gamma \end{pmatrix}$ be the upper triangular ring. Recall from [16] that the (Λ, Γ) -bimodule M is called *compatible* if the following two conditions are satisfied: (1) if Q^{\bullet} is an exact sequence of finitely generated projective Γ -modules, then $M \otimes_{\Gamma} Q^{\bullet}$ is exact; and (2) if P^{\bullet} is a complete finitely generated Λ -projective resolution, then $\operatorname{Hom}_{\Lambda}(P^{\bullet}, M)$ is exact. Let Λ and Γ be artin algebras and the bimodule ${}_{\Lambda}M_{\Gamma}$ compatible. Then finitely generated Gorenstein projective T-modules can be constructed from finitely generated Gorenstein projective T-modules and finitely generated Gorenstein projective T-modules ([16, Theorem 1.4]). Moreover, there exists a left recollement of the stable category $\underline{GP(T)}$ of the category of finitely generated Gorenstein projective T-modules modulo projectives relative to $\underline{GP(\Lambda)}$ and $\underline{GP(\Gamma)}$ ([16, Theorem 3.3]), and this left recollement can be filled into a recollement when T is Gorenstein and ${}_{\Lambda}M$ is projective ([16, Theorem 3.5]). Under some conditions, Enochs, Cortés-Izurdiaga and Torrecillas proved that T is (strongly) CM-free if and only if so are Λ and Γ ([5, Theorem 4.1]).

Let \mathcal{A} and \mathcal{B} be abelian categories and $\mathbf{F}: \mathcal{A} \to \mathcal{B}$ an additive functor. The left comma category $(\mathbf{F}, \mathcal{B})$ was introduced in [8]. Note that module categories of upper triangular matrix rings are comma

^{*2010} Mathematics Subject Classification: 18G25, 18E10, 18E30.

[†]Keywords: Gorenstein objects, Comma categories, Perfect functors, Recollements.

[‡]E-mail address: pengyy@smail.nju.edu.cn, rongminzhu@hotmail.com, huangzy@nju.edu.cn.

categories and that the left comma category $(\mathbf{F}, \mathcal{B})$ is abelian if \mathbf{F} is right exact ([8, 13]). The aim of this paper is to generalize the results mentioned above from module categories of upper triangular matrix rings to comma categories. The paper is organized as follows.

In Section 2, we give some terminology and some preliminary results.

For an abelian category \mathcal{A} , we use $\mathcal{GP}(\mathcal{A})$ to denote the subcategory of \mathcal{A} consisting of Gorenstein projective objects, and use $\underline{\mathcal{GP}(\mathcal{A})}$ to denote the stable category of $\mathcal{GP}(\mathcal{A})$ modulo projectives. Motivated by the definition of compatible bimodules [16], we introduce the so-called *perfect* functors between abelian categories (Definition 3.3). Let \mathcal{A} and \mathcal{B} be abelian categories and $\mathbf{F}: \mathcal{A} \to \mathcal{B}$ an additive and right exact functor such that \mathbf{F} is perfect, and let $(\mathbf{F}, \mathcal{B})$ be the left comma category. Then we give an equivalent characterization of Gorenstein projective objects in $(\mathbf{F}, \mathcal{B})$ in terms of Gorenstein projective objects in \mathcal{B} and \mathcal{A} .

Theorem 1.1. (Theorem 3.5) The following statements are equivalent for an object $\binom{Y}{X}_{\phi}$ in $(\mathbf{F}, \mathcal{B})$.

(1)
$$\binom{Y}{X}_{\phi} \in \mathcal{GP}((\mathbf{F}, \mathcal{B})).$$

(2) $\phi : \mathbf{F}Y \to X$ is injective in \mathcal{B} , $\operatorname{Coker} \phi \in \mathcal{GP}(\mathcal{B})$ and $Y \in \mathcal{GP}(\mathcal{A})$.

As an application, we get that the Gorenstein projective objects coincide with projective objects in $(\mathbf{F}, \mathcal{B})$ if and only if both \mathcal{A} and \mathcal{B} also possess the same property (Corollary 3.9).

In Section 4, we prove the following

Theorem 1.2. (Theorem 4.6) There exists a left recollement

$$\underbrace{\mathcal{GP}(\mathcal{B})} \xrightarrow{i^*} \underbrace{\mathcal{GP}((\mathbf{F},\mathcal{B}))} \xrightarrow{j^*} \underbrace{\mathcal{GP}(\mathcal{A})}.$$

Moreover, this left recollement can be filled into a recollement when \mathcal{B} is Gorenstein and \mathbf{F} preserves projectives (Theorem 4.8).

2 Preliminaries

In this section, we give some notions and some preliminary results.

Let \mathcal{A} be an abelian category and all subcategories of \mathcal{A} are full and closed under isomorphisms. We use $\mathcal{P}(\mathcal{A})$ and $\mathcal{I}(\mathcal{A})$ to denote the subcategories of \mathcal{A} consisting of projective and injective objects respectively. For an object A in \mathcal{A} , $\operatorname{pd}_{\mathcal{A}} A$ and $\operatorname{id}_{\mathcal{A}} A$ are the projective and injective dimensions of A respectively. For a subcategory \mathcal{X} of \mathcal{A} , set

$$\operatorname{pd}_A \mathcal{X} := \sup \{ \operatorname{pd}_A A \mid A \in \mathcal{X} \} \text{ and } \operatorname{id}_A \mathcal{X} := \sup \{ \operatorname{id}_A A \mid A \in \mathcal{X} \}.$$

By using a standard argument, we have the following generalized horseshoe lemma.

Lemma 2.1. Let A be an abelian category and

$$0 \longrightarrow Y \stackrel{f}{\longrightarrow} X \stackrel{g}{\longrightarrow} Z \longrightarrow 0$$

an exact sequence in A.

(1) Let

$$Y \xrightarrow{c^{-1}} C^0 \xrightarrow{c^0} C^1 \xrightarrow{c^1} \cdots$$

be a complex and

$$0 \longrightarrow Z \xrightarrow{d^{-1}} D^0 \xrightarrow{d^0} D^1 \xrightarrow{d^1} \cdots$$


an exact sequence in A. If $\operatorname{Ext}_{A}^{1}(\operatorname{Ker} d^{i}, C^{i})=0$ for any $i\geq 0$, then there exist morphisms

$$\partial^{-1} = \begin{pmatrix} d^{-1}g \\ \sigma^{-1} \end{pmatrix} : X \longrightarrow D^0 \oplus C^0 \text{ and } \partial^i = \begin{pmatrix} d^i & 0 \\ \sigma^i & c^i \end{pmatrix} : D^i \oplus C^i \longrightarrow D^{i+1} \oplus C^{i+1}$$

with $\sigma^i:D^i\to C^{i+1}$ for any $i\geq 0$, such that

$$0 \longrightarrow X \xrightarrow{\partial^{-1}} D^0 \oplus C^0 \xrightarrow{\partial^0} D^1 \oplus C^1 \xrightarrow{\partial^1} \cdots \xrightarrow{\partial^{i-1}} D^i \oplus C^i \xrightarrow{\partial^i} \cdots$$

is a complex in A and the following diagram with exact rows



commutes. Moreover, the middle column is exact if and only if the left column is exact.

(2) Let

$$\cdots \xrightarrow{e_2} E_1 \xrightarrow{e_1} E_0 \xrightarrow{e_0} Y \longrightarrow 0$$

be an exact sequence and

$$\cdots \xrightarrow{f_2} F_1 \xrightarrow{f_1} F_0 \xrightarrow{f_0} Z \longrightarrow 0$$


a complex in A. If $\operatorname{Ext}_{A}^{1}(F_{i}, \operatorname{Im} e_{i}) = 0$ for any $i \geq 0$, then there exist morphisms

$$\partial_0 = (\pi_0, fe_0) : F^0 \oplus E^0 \longrightarrow X \text{ and } \partial_i = \begin{pmatrix} f_i & 0 \\ \pi_i & e_i \end{pmatrix} : F_i \oplus E_i \longrightarrow F_{i-1} \oplus E_{i-1}$$

with $\pi_i: F_i \to E_{i-1}$ for any $i \geq 1$, such that

$$\cdots \xrightarrow{\partial_{i+1}} F_i \oplus E_i \longrightarrow \cdots \xrightarrow{\partial_2} F_1 \oplus E_1 \xrightarrow{\partial_1} F_0 \oplus E_0 \xrightarrow{\partial_0} X \longrightarrow 0$$

is a complex in A and the following diagram with exact rows



commutes. Moreover, the middle column is exact if and only if the right column is exact.

Definition 2.2. ([8]) Let \mathcal{A} be an abelian category and $\mathbf{F}: \mathcal{A} \longrightarrow \mathcal{A}$ an additive endofunctor. The *right* trivial extension of \mathcal{A} by \mathbf{F} , denoted by $\mathcal{A} \ltimes \mathbf{F}$, is defined as follows. An object in $\mathcal{A} \ltimes \mathbf{F}$ is a morphism $\alpha: \mathbf{F} A \longrightarrow A$ for an object A in \mathcal{A} such that $\alpha \cdot \mathbf{F}(\alpha) = 0$; and a morphism in $\mathcal{A} \ltimes \mathbf{F}$ is a pair $(\mathbf{F} \gamma, \gamma)$ of morphisms in \mathcal{A} such that the following diagram

$$FA \xrightarrow{F\gamma} FA'$$

$$\downarrow^{\alpha} \qquad \downarrow^{\alpha'}$$

$$A \xrightarrow{\gamma} A'$$

is commutative.

Definition 2.3. ([8]) Let \mathcal{A} and \mathcal{B} be abelian categories and $\mathbf{F}: \mathcal{A} \longrightarrow \mathcal{B}$ an additive functor. We define the *left comma category* (\mathbf{F}, \mathcal{B}) as follows. The *objects* of the category are $\binom{A}{B}_{\phi}$ with $A \in \mathcal{A}, \in \mathcal{B}$ and $\phi \in \operatorname{Hom}_{\mathcal{B}}(\mathbf{F}A, B)$; and the *morphisms* of the category are pairs $\binom{\alpha}{\beta}$ of morphisms in $\mathcal{A} \times \mathcal{B}$ such that the following diagram

$$FA \xrightarrow{F\alpha} FA'$$

$$\downarrow^{\phi} \qquad \qquad \downarrow^{\phi'}$$

$$B \xrightarrow{\beta} B'$$

is commutative.

Remark 2.4. ([8, Section 1])

(1) The above functor \mathbf{F} induces a functor $\widetilde{\mathbf{F}}: \mathcal{A} \times \mathcal{B} \to \mathcal{A} \times \mathcal{B}$ by $\widetilde{\mathbf{F}}(A, B) = (0, \mathbf{F}A)$ and $\widetilde{\mathbf{F}}(\alpha, \beta) = (0, \mathbf{F}\alpha)$. It is not difficult to show that $(\mathbf{F}, \mathcal{B})$ and $(\mathcal{A} \times \mathcal{B}) \ltimes \widetilde{\mathbf{F}}$ are isomorphic.

(2) If **F** is right exact, then $\mathcal{A} \ltimes \mathbf{F}$ is abelian. Hence by the isomorphism of $(\mathcal{A} \times \mathcal{B}) \ltimes \widetilde{\mathbf{F}}$ and $(\mathbf{F}, \mathcal{B})$, it is clear that if **F** is right exact, then $(\mathbf{F}, \mathcal{B})$ is abelian.

Recall that a sequence in \mathcal{A} is called $\operatorname{Hom}(-,\mathcal{P}(\mathcal{A}))$ -exact if it is exact after applying the functor $\operatorname{Hom}(-,P)$ for any $P\in\mathcal{P}(\mathcal{A})$.

Definition 2.5. ([7]) An object $G \in \mathcal{A}$ is called *Gorenstein projective* if there exists a $\text{Hom}(-, \mathcal{P}(\mathcal{A}))$ -exact exact sequence

$$\cdots \to Q_1 \to Q_0 \to Q^0 \to Q^1 \to \cdots$$

in \mathcal{A} will all Q_i, Q^i projective, such that $G \cong \operatorname{Im}(Q_0 \to Q^0)$; in this case, this exact sequence is called a complete \mathcal{A} -projective resolution of G.

We write $\mathcal{GP}(\mathcal{A}) := \{G \in \mathcal{A} \mid G \text{ is Gorenstein projective}\}$. It is well known that $\mathcal{GP}(\mathcal{A})$ is a Frobenius category such that each object in $\mathcal{P}(\mathcal{A})$ is projective-injective in $\mathcal{GP}(\mathcal{A})$ and its stable category $\mathcal{GP}(\mathcal{A})$ modulo $\mathcal{P}(\mathcal{A})$ is a triangulated category.

3 Gorenstein projective objects

From now on, assume that \mathcal{A} and \mathcal{B} are abelian categories and $\mathbf{F}: \mathcal{A} \longrightarrow \mathcal{B}$ is an additive and right exact functor, and $(\mathbf{F}, \mathcal{B})$ is the left comma category. Then $(\mathbf{F}, \mathcal{B})$ is abelian by Remark 2.4(2). It is known from [8] that the projective object in $(\mathcal{A} \times \mathcal{B}) \ltimes \widetilde{\mathbf{F}}$ is of the form $(\widetilde{\mathbf{F}}(P, Q) \oplus \widetilde{\mathbf{F}}^2(P, Q) \longrightarrow (P, Q) \oplus \widetilde{\mathbf{F}}(P, Q))$ with P projective in \mathcal{A} and Q projective in \mathcal{B} . So by Remark 2.4(1), we have the following

Lemma 3.1. The projective object in $(\mathbf{F}, \mathcal{B})$ is of the form $\begin{pmatrix} 0 \\ Q \end{pmatrix} \oplus \begin{pmatrix} P \\ \mathbf{F}P \end{pmatrix}$ with P projective in \mathcal{A} and Q projective in \mathcal{B} .

The following result generalizes [5, Proposition 2.8(1)].

Proposition 3.2. Let $\binom{M_1}{M_2}$ be an object in $(\mathbf{F}, \mathcal{B})$. If $\operatorname{pd}_{\mathcal{B}} \mathbf{F} \mathcal{P}(\mathcal{A}) < \infty$, then $\operatorname{pd}_{(\mathbf{F}, \mathcal{B})} \binom{M_1}{M_2} < \infty$ if and only if $\operatorname{pd}_{\mathcal{A}} M_1 < \infty$ and $\operatorname{pd}_{\mathcal{B}} M_2 < \infty$.

Proof. Let $\mathrm{pd}_{(\mathbf{F},\mathcal{B})}\binom{M_1}{M_2}<\infty$. Then by Lemma 3.1, we have the following exact sequence of finite length

$$0 \longrightarrow \begin{pmatrix} 0 \\ P_n \end{pmatrix} \oplus \begin{pmatrix} Q_n \\ \mathbf{F}Q_n \end{pmatrix} \longrightarrow \cdots \longrightarrow \begin{pmatrix} 0 \\ P_2 \end{pmatrix} \oplus \begin{pmatrix} Q_2 \\ \mathbf{F}Q_2 \end{pmatrix} \longrightarrow \begin{pmatrix} 0 \\ P_1 \end{pmatrix} \oplus \begin{pmatrix} Q_1 \\ \mathbf{F}Q_1 \end{pmatrix} \longrightarrow \begin{pmatrix} 0 \\ P_0 \end{pmatrix} \oplus \begin{pmatrix} Q_0 \\ \mathbf{F}Q_0 \end{pmatrix} \longrightarrow \begin{pmatrix} M_1 \\ M_2 \end{pmatrix} \longrightarrow 0$$

in $(\mathbf{F}, \mathcal{B})$ with all Q_i projective in \mathcal{A} and all P_i projective in \mathcal{B} . Hence we have exact sequences

$$0 \longrightarrow Q_n \longrightarrow \cdots \longrightarrow Q_2 \longrightarrow Q_1 \longrightarrow Q_0 \longrightarrow M_1 \longrightarrow 0, \tag{3.1}$$

$$0 \longrightarrow P_n \oplus \mathbf{F}Q_n \longrightarrow \cdots \longrightarrow P_2 \oplus \mathbf{F}Q_2 \longrightarrow P_1 \oplus \mathbf{F}Q_1 \longrightarrow P_0 \oplus \mathbf{F}Q_0 \longrightarrow M_2 \longrightarrow 0$$
 (3.2)

in \mathcal{A} and \mathcal{B} respectively. By (3.1), we have $\operatorname{pd}_{\mathcal{A}} M_1 < \infty$. Since $\operatorname{pd}_{\mathcal{B}} \mathbf{F} Q_i < \infty$ for any $0 \leq i \leq n$ by assumption, we have $\operatorname{pd}_{\mathcal{B}} M_2 < \infty$ by (3.2).

Conversely, assume $\operatorname{pd}_{\mathcal{A}} M_1 < \infty$ and $\operatorname{pd}_{\mathcal{B}} M_2 < \infty$. Let

$$0 \longrightarrow Q_n \xrightarrow{\delta_n^1} \cdots \xrightarrow{\delta_3^1} Q_2 \xrightarrow{\delta_2^1} Q_1 \xrightarrow{\delta_1^1} Q_0 \xrightarrow{\delta_0^1} M_1 \longrightarrow 0$$

be a projective resolution of M_1 in \mathcal{A} . Then $\operatorname{pd}_{\mathcal{A}} K_i^1 < \infty$, where $K_i^1 := \operatorname{Ker} \delta_{i-1}^1$ for any $1 \le i \le n+1$. Fix a projective presentation $P_0 \to M_2$ of M_2 in \mathcal{B} . Then we can construct a projective presentation $\binom{Q_0}{P_0 \oplus \mathbf{F} Q_0} \to \binom{M_1}{M_2}$ of $\binom{M_1}{M_2}$ in $(\mathbf{F}, \mathcal{B})$. If $\binom{K_1^1}{K_2^2}$ is its kernel, then there exists an exact sequence

$$0 \longrightarrow K_1^2 \longrightarrow P_0 \oplus \mathbf{F} Q_0 \longrightarrow M_2 \longrightarrow 0$$

in \mathcal{B} . Because $\operatorname{pd}_{\mathcal{B}} \mathbf{F} Q_0 < \infty$ by assumption, we have $\operatorname{pd}_{\mathcal{B}} K_1^2 < \infty$. Repeating this procedure, we get a projective resolution

$$\cdots \xrightarrow{\delta_3} \begin{pmatrix} 0 \\ P_2 \end{pmatrix} \oplus \begin{pmatrix} Q_2 \\ \mathbf{F}Q_2 \end{pmatrix} \xrightarrow{\delta_2} \begin{pmatrix} 0 \\ P_1 \end{pmatrix} \oplus \begin{pmatrix} Q_1 \\ \mathbf{F}Q_1 \end{pmatrix} \xrightarrow{\delta_1} \begin{pmatrix} 0 \\ P_0 \end{pmatrix} \oplus \begin{pmatrix} Q_0 \\ \mathbf{F}Q_0 \end{pmatrix} \xrightarrow{\delta_0} \begin{pmatrix} M_1 \\ M_2 \end{pmatrix} \xrightarrow{\delta_0} 0$$

of $\binom{M_1}{M_2}$ in $(\mathbf{F}, \mathcal{B})$ such that if $\binom{K_i^1}{K_i^2}$ is the kernel of δ_{i-1} , then $\mathrm{pd}_{\mathcal{B}} K_i^2 < \infty$. Since $Q_{n+1} = 0$, we have $\mathrm{Ker} \, \delta_n = \binom{0}{K_{n+1}^2}$. As $\mathrm{pd}_{\mathcal{B}} \, K_{n+1}^2 < \infty$, we have a projective resolution

$$0 \longrightarrow P_{n+m} \longrightarrow \cdots \longrightarrow P_{n+3} \longrightarrow P_{n+2} \longrightarrow P_{n+1} \longrightarrow K_{n+1}^2 \longrightarrow 0$$

of K_{i+1}^2 in \mathcal{B} , which induces the finite projective resolution

$$0 \longrightarrow \begin{pmatrix} 0 \\ P_{n+m} \end{pmatrix} \cdots \longrightarrow \begin{pmatrix} 0 \\ P_{n+3} \end{pmatrix} \longrightarrow \begin{pmatrix} 0 \\ P_{n+2} \end{pmatrix} \longrightarrow \begin{pmatrix} 0 \\ P_{n+1} \end{pmatrix} \longrightarrow \begin{pmatrix} 0 \\ K_{n+1}^2 \end{pmatrix} \longrightarrow 0$$

$$\begin{pmatrix} 0 \\ K_{n+1}^2 \end{pmatrix} \text{ in } (\mathbf{F}, \mathcal{B}). \text{ This means } \mathrm{pd}_{(\mathbf{F}, \mathcal{B})} \operatorname{Ker} \delta_n = \mathrm{pd}_{(\mathbf{F}, \mathcal{B})} \begin{pmatrix} 0 \\ K_{n+1}^2 \end{pmatrix} < \infty, \text{ and hence } \mathrm{pd}_{(\mathbf{F}, \mathcal{B})} \begin{pmatrix} M_1 \\ M_2 \end{pmatrix} < \infty. \qquad \square$$

Motivated by the definition of compatible bimodules in [16, Definition 1.1], we introduce the following

Definition 3.3. The functor **F** is called *perfect* if the following two conditions are satisfied.

- (P1) If \mathcal{Q}^{\bullet} is an exact sequence of projective objects in \mathcal{A} , then $\mathbf{F}\mathcal{Q}^{\bullet}$ is exact.
- (P2) If \mathcal{P}^{\bullet} is a complete \mathcal{B} -projective resolution, then $\operatorname{Hom}(\mathcal{P}^{\bullet}, \mathbf{F}Q)$ is exact for any $Q \in \mathcal{P}(\mathcal{A})$.

For a ring R, $\operatorname{Mod} R$ is the category of left R-modules and $\operatorname{mod} R$ is the category of finitely generated left R-modules. Let Λ and Γ be artin algebras. If M is a compatible (Λ, Γ) -bimodule, then the tensor functor $M \otimes_{\Gamma} -$ is perfect. Let $T := \begin{pmatrix} \Lambda & M \\ 0 & \Gamma \end{pmatrix}$ be the upper triangular matrix algebra. Then $\operatorname{mod} T$ is the left comma category $(M \otimes_{\Gamma} -, \operatorname{mod} \Lambda)$.

Lemma 3.4. The following statements are equivalent.

- (1) **F** satisfies (P2);
- (2) $\operatorname{Ext}_{\mathcal{B}}^{1}(G, \mathbf{F}Q) = 0$ for any $G \in \mathcal{GP}(\mathcal{B})$ and $Q \in \mathcal{P}(\mathcal{A})$;
- (3) $\operatorname{Ext}_{\mathcal{B}}^{\geq 1}(G, \mathbf{F}Q) = 0$ for any $G \in \mathcal{GP}(\mathcal{B})$ and $Q \in \mathcal{P}(\mathcal{A})$.

Proof. The implications $(1) \Rightarrow (3) \Rightarrow (2)$ are trivial. Applying the functor $\text{Hom}_{\mathcal{B}}(-, \mathbf{F}Q)$ to a complete \mathcal{B} -projective resolution of G, we get $(2) \Rightarrow (1)$.

We now give an equivalent characterization of Gorenstein projective objects in the left comma category $(\mathbf{F}, \mathcal{B})$. It is a generalization of [16, Theorem 1.4].

Theorem 3.5. If **F** is perfect, then the following statements are equivalent for an object $\binom{Y}{X}_{\phi}$ in $(\mathbf{F}, \mathcal{B})$.

- (1) $\binom{Y}{X}_{\phi} \in \mathcal{GP}((\mathbf{F}, \mathcal{B})).$
- (2) $\phi : \mathbf{F}Y \to X$ is injective in \mathcal{B} , $\operatorname{Coker} \phi \in \mathcal{GP}(\mathcal{B})$ and $Y \in \mathcal{GP}(\mathcal{A})$.

In this case, $X \in \mathcal{GP}(\mathcal{B})$ if and only if $\mathbf{F}Y \in \mathcal{GP}(\mathcal{B})$.

Proof. (2) \Rightarrow (1) Assume that $\phi : \mathbf{F}Y \to X$ is injective in \mathcal{B} , Coker $\phi \in \mathcal{GP}(\mathcal{B})$ and $Y \in \mathcal{GP}(\mathcal{A})$. Then we have a complete \mathcal{A} -projection resolution

$$(\mathcal{Q}^{\bullet}, q^{\cdot}) := \cdots \longrightarrow Q^{-1} \longrightarrow Q^{0} \xrightarrow{q^{0}} Q^{1} \longrightarrow \cdots$$

with $Y = \operatorname{Ker} q^0$. Since $\mathbf{F} \mathcal{Q}^{\bullet}$ is exact by (P1), we have the following exact sequence

$$0 \longrightarrow \mathbf{F}Y \longrightarrow \mathbf{F}Q^0 \xrightarrow{Fq^0} \mathbf{F}Q^1 \xrightarrow{Fq^1} \cdots$$

Since $\operatorname{Coker} \phi \in \mathcal{GP}(\mathcal{B})$, we have a complete \mathcal{B} -projective resolution

$$\cdots \longrightarrow P^{-1} \longrightarrow P^0 \xrightarrow{d^0} P^1 \longrightarrow \cdots$$

with $\operatorname{Coker} \phi = \operatorname{Ker} d^0$, so $\operatorname{Ker} d^i \in \mathcal{GP}(\mathcal{B})$, and hence $\operatorname{Ext}^1_{\mathcal{B}}(\operatorname{Ker} d^i, \mathbf{F}Q^i) = 0$ for any $i \geq 0$. Applying Lemma 2.1(1) to the exact sequence

$$0 \longrightarrow \mathbf{F} Y \xrightarrow{\phi} X \longrightarrow \operatorname{Coker} \phi \longrightarrow 0$$
,

we obtain an exact sequence

$$0 \longrightarrow X \stackrel{\partial^{-1}}{\longrightarrow} P^0 \oplus \mathbf{F} Q^0 \stackrel{\partial^0}{\longrightarrow} P^1 \oplus \mathbf{F} Q^1 \stackrel{\partial^1}{\longrightarrow} \cdots$$

with $\partial^i = \begin{pmatrix} d^i & 0 \\ \sigma^i & Fq^i \end{pmatrix}$ and $\sigma^i : P^i \to FQ^{i+1}$ for any $i \ge 0$, such that the following diagram with exact rows

$$0 \longrightarrow \mathbf{F}Y \longrightarrow \mathbf{F}Q^{0} \longrightarrow \mathbf{F}Q^{1} \longrightarrow \cdots$$

$$\downarrow \qquad \qquad \downarrow \qquad \qquad \downarrow$$

$$0 \longrightarrow X \longrightarrow P^{0} \oplus \mathbf{F}Q^{0} \longrightarrow P^{1} \oplus \mathbf{F}Q^{1} \longrightarrow \cdots$$

commutes. By a dual argument we get the following diagram with exact rows

$$\cdots \longrightarrow \mathbf{F}Q^{-2} \longrightarrow \mathbf{F}Q^{-1} \longrightarrow \mathbf{F}Y \longrightarrow 0$$

$$\downarrow \qquad \qquad \downarrow \qquad \qquad \downarrow$$

$$\cdots \longrightarrow P^{-2} \oplus \mathbf{F}Q^{-2} \longrightarrow P^{-1} \oplus \mathbf{F}Q^{-1} \longrightarrow X \longrightarrow 0.$$

Combining these two diagrams to get the following diagram with exact rows

Actually, we have the following exact sequence of projective objects

$$L^{\bullet} = \cdots \longrightarrow \begin{pmatrix} Q^{-1} \\ P^{-1} \oplus \mathbf{F} Q^{-1} \end{pmatrix} \longrightarrow \begin{pmatrix} Q^{0} \\ P^{0} \oplus \mathbf{F} Q^{0} \end{pmatrix} \longrightarrow \begin{pmatrix} Q^{1} \\ P^{1} \oplus \mathbf{F} Q^{1} \end{pmatrix} \longrightarrow \cdots$$

in $(\mathbf{F}, \mathcal{B})$. Since each L^i is a projective object in $(\mathbf{F}, \mathcal{B})$, applying $\operatorname{Hom}(L^i, -)$ to the exact sequence:

$$0 \longrightarrow \begin{pmatrix} 0 \\ \widetilde{P} \oplus \widetilde{FQ} \end{pmatrix} \longrightarrow \begin{pmatrix} \widetilde{Q} \\ \widetilde{P} \oplus \widetilde{FQ} \end{pmatrix} \longrightarrow \begin{pmatrix} \widetilde{Q} \\ 0 \end{pmatrix} \longrightarrow 0$$

we get the following exact sequence of complexes

$$0 \longrightarrow \operatorname{Hom}(L^{\bullet}, \left(\begin{smallmatrix} 0 \\ \widehat{P} \oplus \widetilde{FQ} \end{smallmatrix}\right)) \longrightarrow \operatorname{Hom}(L^{\bullet}, \left(\begin{smallmatrix} \widetilde{Q} \\ \widehat{P} \oplus \widetilde{FQ} \end{smallmatrix}\right)) \longrightarrow \operatorname{Hom}(L^{\bullet}, \left(\begin{smallmatrix} \widetilde{Q} \\ 0 \end{smallmatrix}\right)) \longrightarrow 0,$$

that is,

$$0 \longrightarrow \operatorname{Hom}(P^{\bullet}, \widetilde{P} \oplus \widetilde{FQ}) \longrightarrow \operatorname{Hom}(L^{\bullet}, \left(\begin{smallmatrix} \widetilde{Q} \\ \widetilde{P} \oplus \widetilde{FQ} \end{smallmatrix} \right)) \longrightarrow \operatorname{Hom}(Q^{\bullet}, \widetilde{Q}) \longrightarrow 0.$$

Since P^{\bullet} is a complete \mathcal{B} -projective resolution, it follows that $\operatorname{Hom}(P^{\bullet}, \widetilde{P})$ is exact. By (P2), $\operatorname{Hom}(P^{\bullet}, \widetilde{FQ})$ is exact. Since Q^{\bullet} is a complete \mathcal{A} -projective resolution, it follows that $\operatorname{Hom}(Q^{\bullet}, \widetilde{Q})$ is exact. Thus $\operatorname{Hom}(L^{\bullet}, \binom{\widetilde{Q}}{\widetilde{P} \oplus \widetilde{FQ}})$ is also exact. Therefore we conclude that L^{\bullet} is a complete $(\mathbf{F}, \mathcal{B})$ -projective resolution and $\binom{Y}{X}_{\phi} \in \mathcal{GP}((\mathbf{F}, \mathcal{B}))$.

 $(1) \Rightarrow (2)$ Let $\binom{Y}{X}_{\phi} \in \mathcal{GP}((\mathbf{F}, \mathcal{B}))$. Then we have a complete $(\mathbf{F}, \mathcal{B})$ -projective resolution

$$L^{\bullet} := \cdots \longrightarrow \begin{pmatrix} Q^{-1} \\ P^{-1} \oplus \mathbf{F} Q^{-1} \end{pmatrix} \longrightarrow \begin{pmatrix} Q^{0} \\ P^{0} \oplus \mathbf{F} Q^{0} \end{pmatrix} \xrightarrow{\begin{pmatrix} d^{\prime 0} \\ \partial^{0} \end{pmatrix}} \begin{pmatrix} Q^{1} \\ P^{1} \oplus \mathbf{F} Q^{1} \end{pmatrix} \longrightarrow \cdots$$


such that $\operatorname{Ker} \binom{d'^0}{\partial^0} = \binom{Y}{X}_{\phi}$. Then we get an exact sequence $(Q^{\bullet}, d'^{\bullet})$ of projective objects in \mathcal{A} with $\operatorname{Ker} d'^0 = Y$ and the following exact sequence

$$V^{\bullet} := \cdots \longrightarrow P^{-1} \oplus \mathbb{F}Q^{-1} \longrightarrow P^{0} \oplus \mathbb{F}Q^{0} \xrightarrow{\partial^{0}} P^{1} \oplus \mathbb{F}Q^{1} \longrightarrow \cdots$$

with Ker $\partial^0 = X$. By (P1), $\mathbf{F}Q^{\bullet}$ is exact. Since $\begin{pmatrix} d'^i \\ \partial^i \end{pmatrix} : \begin{pmatrix} Q^i \\ P^i \oplus \mathbf{F}Q^i \end{pmatrix} \to \begin{pmatrix} Q^{i+1} \\ P^{i+1} \oplus \mathbf{F}Q^{i+1} \end{pmatrix}$ is a morphism in $(\mathbf{F}, \mathcal{B})$, we get that ∂^i is of the form $\partial^i = \begin{pmatrix} d^i & 0 \\ \sigma^i & \mathbf{F}d'^i \end{pmatrix}$ where $\sigma^i : P^i \to \mathbf{F}Q^{i+1}$ for any i. We have the exact sequence of complexes

$$0 \longrightarrow \mathbf{F} Q^{\bullet} \longrightarrow V^{\bullet} \longrightarrow P^{\bullet} \longrightarrow 0$$

with P^{\bullet} exact. So we get the following diagram with exact columns and rows



such that $\operatorname{Ker} \mathbf{F} d^{0} = \mathbf{F} Y$. Applying the snake lemma we get the following exact sequence

$$0 \longrightarrow \operatorname{Ker} \mathbf{F} d'^0 \longrightarrow \operatorname{Ker} \partial^0 \longrightarrow \operatorname{Ker} d^0 \longrightarrow \operatorname{Im} \mathbf{F} d'^1 \longrightarrow \operatorname{Im} \partial^1 \longrightarrow \operatorname{Im} d^1 \longrightarrow 0$$

that is,

$$0 \longrightarrow \mathbf{F}Y \stackrel{\phi}{\longrightarrow} X \stackrel{\pi'}{\longrightarrow} \operatorname{Ker} d^0 \longrightarrow \operatorname{Im} \mathbf{F} d'^1 \longrightarrow \operatorname{Im} \partial^1 \longrightarrow \operatorname{Im} d^1 \longrightarrow 0.$$

Because the morphism $\operatorname{Im} \mathbf{F} d'^1 \to \operatorname{Im} \partial^1$ is injective, it follows that π' is surjective. Hence $\operatorname{Ker} d^0 \cong \operatorname{Coker} \phi$. Since $\operatorname{Hom}(L^{\bullet}, \binom{0}{\mathcal{P}}) \cong \operatorname{Hom}(P^{\bullet}, \mathcal{P})$ and L^{\bullet} is a complete projection resolution, it follows that $\operatorname{Hom}(P^{\bullet}, \mathcal{P})$ is exact. Hence P^{\bullet} is a complete \mathcal{B} -projective resolution and $\operatorname{Coker} \phi \in \mathcal{GP}(\mathcal{B})$. By (P2), $\operatorname{Hom}(P^{\bullet}, \mathbf{F}Q)$ is exact. Similarly, since each L^i is a projective object in $(\mathbf{F}, \mathcal{B})$, applying $\operatorname{Hom}(L^i, -)$ to the exact sequence

$$0 \longrightarrow \begin{pmatrix} 0 \\ \widetilde{P} \oplus \widetilde{FQ} \end{pmatrix} \longrightarrow \begin{pmatrix} \widetilde{Q} \\ \widetilde{P} \oplus \widetilde{FQ} \end{pmatrix} \longrightarrow \begin{pmatrix} \widetilde{Q} \\ 0 \end{pmatrix} \longrightarrow 0,$$

we get the following exact sequence of complexes

$$0 \longrightarrow \operatorname{Hom}(P^{\bullet}, \widetilde{P} \oplus \widetilde{FQ}) \longrightarrow \operatorname{Hom}(L^{\bullet}, (\widetilde{\widetilde{P}}_{\oplus \widetilde{FQ}})) \longrightarrow \operatorname{Hom}(Q^{\bullet}, \widetilde{Q}) \longrightarrow 0.$$

Since L^{\bullet} is a complete projective resolution, $\operatorname{Hom}(L^{\bullet}, \binom{\widetilde{Q}}{\widetilde{P} \oplus \widetilde{FQ}})$ is exact, and then $\operatorname{Hom}(Q^{\bullet}, \widetilde{Q})$ is also exact. It follows that $Y \in \mathcal{GP}(\mathcal{A})$.

As an application of Theorem 3.5, we have the following

Corollary 3.6. Let F be perfect. Then

- (1) If $(\mathbf{F}, \mathcal{B})$ has finitely many isomorphism classes of indecomposable Gorenstein projective objects, then so have \mathcal{A} and \mathcal{B} .
- (2) If $\mathcal{GP}(\mathcal{B}) = \mathcal{P}(\mathcal{B})$, then $\binom{0}{P}$ and $\binom{Y}{\mathbf{F}Y}$ are exactly all indecomposable Gorenstein projective objects in $(\mathbf{F}, \mathcal{B})$, where Y runs over all indecomposable objects in $\mathcal{GP}(\mathcal{A})$ and P runs over all indecomposable objects in $\mathcal{P}(\mathcal{B})$.

(3) If $\mathcal{GP}(A) = \mathcal{P}(A)$, then $\binom{0}{X}$ and $\binom{Q}{\mathbf{F}Q}$ are exactly all indecomposable Gorenstein projective objects in $(\mathbf{F}, \mathcal{B})$, where Q runs over all indecomposable objects in $\mathcal{P}(A)$ and X runs over all the indecomposable objects in $\mathcal{GP}(\mathcal{B})$.

Proof. (1) Let $X \in \mathcal{GP}(\mathcal{B})$ and $Y \in \mathcal{GP}(\mathcal{A})$. Then by Theorem 3.5, both $\binom{0}{X}$ and $\binom{Y}{\mathbf{F}Y}$ are Gorenstein projective objects in $(\mathbf{F}, \mathcal{B})$. The assertion follows.

(2) + (3) Let $\binom{Y}{X}_{\phi}$ be Gorenstein projective in $(\mathbf{F}, \mathcal{B})$. Then by Theorem 3.5, there exists an exact sequence

$$0 \longrightarrow \mathbf{F} Y \xrightarrow{\phi} X \longrightarrow \operatorname{Coker} \phi \longrightarrow 0$$

in \mathcal{B} with Coker $\phi \in \mathcal{GP}(\mathcal{B})$ and $Y \in \mathcal{GP}(\mathcal{A})$.

If $\mathcal{GP}(\mathcal{B}) = \mathcal{P}(\mathcal{B})$, then $\operatorname{Coker} \phi \in \mathcal{P}(\mathcal{B})$ and the above exact sequence splits. If $\mathcal{GP}(\mathcal{A}) = \mathcal{P}(\mathcal{A})$, then $Y \in \mathcal{P}(\mathcal{A})$. By Lemma 3.4, we have $\operatorname{Ext}_{\mathcal{B}}^{\geq 1}(\operatorname{Coker} \phi, \mathbf{F}Y) = 0$. So the above exact sequence also splits. So, in both cases, we have $X = \mathbf{F}Y \oplus \operatorname{Coker} \phi$ and $\binom{Y}{X}_{\phi} = \binom{Y}{\mathbf{F}Y} \oplus \binom{0}{\operatorname{Coker} \phi}$. The assertions (2) and (3) follow.

Example 3.7. Let k be a field and T a finite-dimensional k-algebra given by the quiver

$$\begin{pmatrix}
\uparrow \\
1 \\
\downarrow \\
3 \longrightarrow 2 \longleftarrow 4$$

with relation $\gamma^3 = 0$. Then

$$T = \begin{pmatrix} e_1 T e_1 & e_1 T (1 - e_1) \\ 0 & (1 - e_1) T (1 - e_1) \end{pmatrix},$$

where e_1 is the idempotent corresponding to the vertex 1. We have that $\Gamma := (1 - e_1)T(1 - e_1)$ is a finite-dimensional k-algebra given by the quiver


$$3 \longrightarrow 2 \longleftarrow 4$$
,

and $\Lambda := e_1 T e_1$ is a finite-dimensional k-algebra given by the quiver



with relation $\gamma^3 = 0$. Take $\mathcal{A} := \text{mod } \Gamma$, $\mathcal{B} := \text{mod } \Lambda$ and $\mathbf{F} := M \otimes_{\Gamma} - \text{with } M = e_1 T (1 - e_1)$. Then $(\mathbf{F}, \mathcal{B}) = \text{mod } T$. We have $\Lambda M \cong \Lambda \Lambda \oplus_{\Lambda} \Lambda \oplus_{\Lambda} \Lambda \oplus_{\Lambda} \Lambda$ and $M_{\Gamma} \cong I(2)_{\Gamma} \oplus I(2)_{\Gamma} \oplus I(2)_{\Gamma}$. Since Γ is hereditary, pd $M_{\Gamma} \leq 1$ and \mathbf{F} is perfect. Since Λ is self-injective, each module in mod Λ is Gorenstein projective.

Then by Corollary 3.6, all indecomposable Gorenstein projective modules in $\operatorname{mod} T$ are as follows.



Example 3.8. Let k be a field and T a finite-dimensional k-algebra given by the quiver

$$5 \xrightarrow{\beta} 4 \xrightarrow{\gamma} 2 \xrightarrow{\alpha_1} \xrightarrow{\alpha_2} 3$$

with the relation $\alpha_2\alpha_1=\alpha_3\alpha_2=\alpha_1\alpha_3=0$. Then

$$T = \begin{pmatrix} (e_1 + e_2 + e_3)T(e_1 + e_2 + e_3) & (e_1 + e_2 + e_3)T(e_4 + e_4) \\ 0 & (e_4 + e_5)T(e_4 + e_5) \end{pmatrix},$$

where e_i is the idempotent corresponding to the vertex i for any $1 \le i \le 5$. We have that $\Gamma := (e_4 + e_5)T(e_4 + e_5)$ is a finite-dimensional k-algebra given by the quiver


$$5 \longrightarrow 4$$

and $\Lambda := (e_1 + e_2 + e_3)T(e_1 + e_2 + e_3)$ is a finite-dimensional k-algebra given by the quiver

$$\begin{array}{c}
1 \\
\alpha_1 \\
2 \\
\end{array}$$

$$\begin{array}{c}
\alpha_3 \\
3
\end{array}$$

with relation $\alpha_2\alpha_1 = \alpha_3\alpha_2 = \alpha_1\alpha_3 = 0$. Take $\mathcal{A} := \operatorname{mod}\Gamma$, $\mathcal{B} := \operatorname{mod}\Lambda$ and $\mathbf{F} := M \otimes_{\Gamma} - \operatorname{with}M = (e_1 + e_2 + e_3)T(e_4 + e_5)$. Then $(\mathbf{F},\mathcal{B}) = \operatorname{mod}T$. We have ${}_{\Lambda}M \cong {}_{\Lambda}P(2) \oplus_{\Lambda}P(2)$ and $M_{\Gamma} \cong P(5)_{\Gamma} \oplus P(5)_{\Gamma}$, and so \mathbf{F} is perfect. Notice that Λ is self-injective and Γ is hereditary, so by Corollary 3.6, all indecomposable Gorenstein projective modules in $\operatorname{mod} T$ are as follows.



By Theorem 3.5, we also have the following

Corollary 3.9. If **F** is perfect, then $\mathcal{GP}((\mathbf{F},\mathcal{B})) = \mathcal{P}((\mathbf{F},\mathcal{B}))$ if and only if $\mathcal{GP}(\mathcal{A}) = \mathcal{P}(\mathcal{A})$ and $\mathcal{GP}(\mathcal{B}) = \mathcal{P}(\mathcal{B})$.

Proof. We first prove the necessity. Let Y be Gorenstein projective in \mathcal{A} . Then by Theorem 3.5, $\binom{Y}{\mathbf{F}Y}$ is Gorenstein projective in $(\mathbf{F}, \mathcal{B})$. So $\binom{Y}{\mathbf{F}Y}$ is projective in $(\mathbf{F}, \mathcal{B})$ by assumption, and hence Y is projective in \mathcal{A} . Now let X be Gorenstein projective in \mathcal{B} . Then by Theorem 3.5, $\binom{0}{X}$ is Gorenstein projective in $(\mathbf{F}, \mathcal{B})$. So $\binom{0}{X}$ is projective in $(\mathbf{F}, \mathcal{B})$ by assumption, and hence X is projective in \mathcal{B} .

We next prove the sufficiency. Let $\binom{Y}{X}_{\phi}$ be Gorenstein projective in $(\mathbf{F}, \mathcal{B})$. Then we have the following exact sequence

$$0 \longrightarrow \mathbf{F}Y \longrightarrow X \longrightarrow \operatorname{Coker} \phi \longrightarrow 0$$

in \mathcal{B} with $\operatorname{Coker} \phi \in \mathcal{GP}(\mathcal{B})$ and $Y \in \mathcal{GP}(\mathcal{A})$ by Theorem 3.5. So $\operatorname{Coker} \phi$ is projective in \mathcal{B} and Y is projective in \mathcal{A} by assumption, and hence $X = \mathbf{F}Y \oplus \operatorname{Coker} \phi$ and $\binom{Y}{X}_{\phi} = \binom{Y}{Y} \oplus \binom{0}{\operatorname{Coker} \phi}$. Thus $\binom{Y}{X}_{\phi}$ is projective in $(\mathbf{F}, \mathcal{B})$ by Lemma 3.1.

Recall from [5] that a ring R is called *strongly left CM-free* if each Gorenstein projective module in Mod R is projective. Let Λ and Γ be arbitrary rings and M a (Λ, Γ) -bimodule, and let $T := \begin{pmatrix} \Lambda & M \\ 0 & \Gamma \end{pmatrix}$ be the upper triangular matrix ring. Then Mod T is the left comma category $(M \otimes_{\Gamma} -, \operatorname{Mod} \Lambda)$. If M_{Γ} has finite flat dimension and ΛM has finite projective dimension, then the functor $M \otimes_{\Gamma} -$ is perfect. So, as an immediate consequence of Corollary 3.9, we have the following

Corollary 3.10. Let Λ and Γ be arbitrary rings and M a (Λ, Γ) -bimodule, and let T be the upper triangular matrix ring as above. If M_{Γ} has finite flat dimension and ΛM has finite projective dimension, then T is strongly left CM-free if and only if so are Λ and Γ .

The above corollary generalized [5, Theorem 4.1], where the assumption that Λ is left Gorenstein regular is needed.

4 Recollements

Definition 4.1. ([9, 13]) A recollement, denoted by $(\mathcal{A}, \mathcal{B}, \mathcal{C})$, of abelian categories is a diagram

$$\mathcal{A} \xrightarrow{i^*} \mathcal{B} \xrightarrow{j_!} \mathcal{C}$$

$$\stackrel{i^*}{\longleftarrow} \mathcal{B} \xrightarrow{j_*} \mathcal{C}$$

of abelian categories and additive functors such that

- (1) $(i^*, i_*), (i_*, i^!), (j_!, j^*)$ and (j^*, j_*) are adjoint pairs.
- (2) $i_*, j_!$ and j_* are fully faithful.
- (3) Im $i_* = \text{Ker } j^*$.

The following lemma is fundamental in this section.

Lemma 4.2. ([13, Example 2.12]) There exists the following recollement of abelian categories:

$$\mathcal{B} \xrightarrow{i^*} i_* \longrightarrow (\mathbf{F}, \mathcal{B}) \xrightarrow{j_!} \mathcal{A},$$

where

$$i^*: \begin{pmatrix} Y \\ X \end{pmatrix}_{\phi} \mapsto \operatorname{Coker} \phi, \ i_*: X \mapsto \begin{pmatrix} 0 \\ X \end{pmatrix}, \ i^!: \begin{pmatrix} Y \\ X \end{pmatrix} \mapsto X,$$

$$j_!: Y \mapsto \begin{pmatrix} Y \\ \mathbf{F}Y \end{pmatrix}, \ j^*: \begin{pmatrix} Y \\ X \end{pmatrix} \mapsto Y, \ j_*: Y \mapsto \begin{pmatrix} Y \\ 0 \end{pmatrix}.$$

Definition 4.3. ([4]) Let \mathcal{C}' , \mathcal{C} and \mathcal{C}'' be triangulated categories. The diagram of exact functors

$$C' \xrightarrow{i^*} \stackrel{i^*}{\longrightarrow} C \xrightarrow{j_!} \stackrel{j_!}{\longrightarrow} C''$$

$$(4.1)$$

is a recollement of C relative to C' and C'', if the following four conditions are satisfied.

- (R1) $(i^*, i_*), (i_*, i^!), (j_!, j^*)$ and (j^*, j_*) are adjoint pairs.
- (R2) i_* , $j_!$ and j_* are fully faithful.
- (R3) $j^*i_* = 0$.
- (R4) For each object $X \in \mathcal{C}$, the counits and units give rise to the following distinguished triangles

$$j_!j^*(X) \xrightarrow{\epsilon_X} X \xrightarrow{\eta_X} i_*i^*(X) \longrightarrow j_!j^*(X)[1],$$

$$i_*i^!(X) \xrightarrow{\omega_X} X \xrightarrow{\zeta_X} j_*j^*(X) \longrightarrow i_*i^!(X)[1],$$

where [1] is the shift functor.

A left recollement of C relative to C' and C'' is a diagram of exact functors consisting of the upper two rows in the diagram (4.1) satisfying all the conditions which involve only the functors $i^*, i_*, j_!, j^*$.

The following result is useful in the sequel.

Lemma 4.4. ([11, Section 1]) Let (4.1) be a diagram of triangulated categories. Then the following statements are equivalent.

- (1) The diagram (4.1) is a recollement.
- (2) The conditions (R1), (R2) and $\operatorname{Im} i_* = \operatorname{Ker} j^*$ are satisfied.
- (3) The conditions (R1), (R2) and $\operatorname{Im} j_! = \operatorname{Ker} i^*$ are satisfied.
- (4) The conditions (R1), (R2) and $\text{Im } j_* = \text{Ker } i^!$ are satisfied.


Remark 4.5. Each assertion in Lemma 4.4(2)–(4) involving only the functors $i^*, i_*, j_!, j^*$ is equivalent to that the upper two rows in the diagram (4.1) is a left recollement.

The following result is a generalization of [16, Theorem 3.3].

Theorem 4.6. If **F** is perfect, then there exists a left recollement

$$\underbrace{\mathcal{GP}(\mathcal{B})} \xrightarrow{i^*} \underbrace{\mathcal{GP}((\mathbf{F},\mathcal{B}))} \xrightarrow{j^*} \underbrace{\mathcal{GP}(\mathcal{A})}.$$

Proof. We first construct the functors involved. By Theorem 3.5, we know the form of Gorenstein projective objects in $(\mathbf{F}, \mathcal{B})$. If a morphism $\binom{X}{Y}_{\phi} \to \binom{X'}{Y'}_{\phi'}$ factors through a projective object $\binom{0}{P} \oplus \binom{Q}{\mathbf{F}Q}$, then we have the following diagram with exact rows



Hence the functor i^* in Lemma 4.2 induces a functor which we still denote by $i^*: \underline{\mathcal{GP}((\mathbf{F},\mathcal{B}))} \to \underline{\mathcal{GP}(\mathcal{B})}$. By Lemma 4.2, we have the functor i_* given by $Y \to \binom{0}{Y}$. It is obvious a functor $\underline{\mathcal{GP}(\mathcal{B})} \to \underline{\mathcal{GP}((\mathbf{F},\mathcal{B}))}$. If a morphism $Y \to Y'$ in \mathcal{B} factors through a projective object P, then $\binom{0}{Y} \to \binom{0}{Y'}$ factors through a projective object $\binom{0}{P}$ in (\mathbf{F},\mathcal{B}) . Hence i_* induces a functor $i_*:\underline{\mathcal{GP}(\mathcal{B})} \to \underline{\mathcal{GP}((\mathbf{F},\mathcal{B}))}$, which is fully faithful.

By Lemma 4.2, we have the functor $j_!$ given by $A \to \binom{A}{\mathbf{F}A}$. It is a functor $\mathcal{GP}(A) \to \mathcal{GP}((\mathbf{F}, \mathcal{B}))$ by Theorem 3.5. If a morphism $X \to X'$ in \mathcal{A} factors through a projective object Q, then $\binom{X}{\mathbf{F}X} \to \binom{X'}{\mathbf{F}X'}$ factors through a projective object $\binom{Q}{\mathbf{F}Q}$ in $(\mathbf{F}, \mathcal{B})$. Hence $j_!$ induces a functor $j_! : \underline{\mathcal{GP}(A)} \to \underline{\mathcal{GP}((\mathbf{F}, \mathcal{B}))}$, which is fully faithful.

By Lemma 4.2, we have the functor j^* given by $\binom{X}{Y} \to X$. It is a functor from $\mathcal{GP}((\mathbf{F},\mathcal{B})) \to \mathcal{GP}(\mathcal{A})$ by Theorem 3.5. If a morphism $\binom{X}{Y} \to \binom{X'}{Y'}$ in (\mathbf{F},\mathcal{B}) factors through a projective object $\binom{Q}{P} \oplus \binom{0}{P}$, then $X \to X'$ factors through a projective object Q in \mathcal{A} . Hence j^* induces a functor $j^* : \underline{\mathcal{GP}((\mathbf{F},\mathcal{B}))} \to \underline{\mathcal{GP}(\mathcal{A})}$. It follows easily from [10, Chapter I, Section 2] that i_*, j^* constructed above are exact functors. By Lemma 4.2, we have that both (i^*, i_*) and $(j_!, j^*)$ are adjoint pairs. Thus i^* and $j_!$ are exact functors by [12, Lemma 8.3].

By construction, we have $\operatorname{Im} i_* \subseteq \operatorname{Ker} j^*$ and $\operatorname{Ker} j^* = \{\binom{X}{Y} \in \underline{\mathcal{GP}((\mathbf{F},\mathcal{B}))} \mid X \in \mathcal{P}(\mathcal{A})\}$. Let $\binom{X}{Y} \in \operatorname{Ker} j^*$. By Theorem 3.5, we have the following exact sequence

$$0 \longrightarrow \mathbf{F} X \xrightarrow{\phi} Y \longrightarrow \operatorname{Coker} \phi \longrightarrow 0$$

in \mathcal{B} with $\operatorname{Coker} \phi \in \mathcal{GP}(\mathcal{B})$. Then $\operatorname{Ext}^1_{\mathcal{B}}(\operatorname{Coker} \phi, \mathbf{F}X) = 0$ by Lemma 3.4. So the above exact splits and $Y \cong \mathbf{F}X \oplus \operatorname{Coker} \phi$. Thus we have

$$\begin{pmatrix} X \\ Y \end{pmatrix} \cong \begin{pmatrix} X \\ \mathbf{F}X \end{pmatrix} \oplus \begin{pmatrix} 0 \\ \operatorname{Coker} \phi \end{pmatrix} = i_*(\operatorname{Coker} \phi),$$

which implies $\operatorname{Ker} j^* \subseteq \operatorname{Im} i_*$

Finally, applying Lemma 4.4(2) and Remark 4.5, we get the required left recollement.

It is natural to ask when the left recollement in Theorem 4.6 can be filled into a recollement. In the following, we will study this question.

Recall from [3] that an abelian category \mathcal{B} with enough projective and injective objects is called Gorenstein if $\operatorname{pd}_{\mathcal{B}} \mathcal{I}(\mathcal{B}) < \infty$ and $\operatorname{id}_{\mathcal{B}} \mathcal{P}(\mathcal{B}) < \infty$.

Lemma 4.7. Let \mathbf{F} be perfect. If \mathcal{B} is Gorenstein and \mathbf{F} preserves projectives, then \mathbf{F} preserves Gorenstein projectives.

Proof. Let $Y \in \mathcal{A}$ be Gorenstein projective. Then there exists a complete $\mathcal{P}(\mathcal{A})$ -resolution

$$Q^{\bullet} := \cdots \to Q_1 \to Q_0 \xrightarrow{d} Q^0 \to Q^1 \to \cdots$$

in \mathcal{A} such that $Y \cong \operatorname{Im} d$. Since \mathbf{F} is perfect, $\mathbf{F}Q^{\bullet}$ is exact and $\mathbf{F}Y \cong \operatorname{Ker} \mathbf{F}d$. If \mathbf{F} preserves projectives, then all terms in $\mathbf{F}Q^{\bullet}$ are projective in \mathcal{B} . Let $P \in \mathcal{B}$ be projective. Because \mathcal{B} is Gorenstein by assumption, we have $\operatorname{id}_{\mathcal{B}} P < \infty$. So $\operatorname{Hom}(\mathbf{F}Q^{\bullet}, P)$ is exact, and hence $\mathbf{F}Y$ is Gorenstein projective. \square

As a generalization of [16, Theorem 3.5], we have the following

Theorem 4.8. Let \mathbf{F} be perfect. If \mathcal{B} is Gorenstein and \mathbf{F} preserves projectives, then there exists a recollement

$$\underbrace{\mathcal{GP}(\mathcal{B})} \xrightarrow{\overset{i^*}{\longleftarrow} i_*} \underbrace{\mathcal{GP}((\mathbf{F}, \mathcal{B}))} \underbrace{\overset{\longleftarrow j_!}{\longleftarrow} j_*^*} \underbrace{\mathcal{GP}(\mathcal{A})}.$$

Proof. By Theorem 4.6, there exists the following left recollement

$$\mathcal{GP}(\mathcal{B}) \xrightarrow{i^*} \mathcal{GP}((\mathbf{F}, \mathcal{B})) \xrightarrow{j^*} \mathcal{GP}(\mathcal{A}).$$

By Lemma 4.2, we have the functor $i^!$ given by $\binom{X}{Y} \to Y$. It is a functor $\mathcal{GP}((\mathbf{F},\mathcal{B})) \to \mathcal{GP}(\mathcal{B})$. If a morphism $\binom{X}{Y}_{\phi} \to \binom{X'}{Y'}_{\phi'}$ in (\mathbf{F},\mathcal{B}) factors through a projective object $\binom{0}{P} \oplus \binom{Q}{\mathbf{F}Q}$ with Q projective in \mathcal{A} and P projective in \mathcal{B} , then $Y \to Y'$ factors through $P \oplus \mathbf{F}Q$. Since \mathbf{F} preserves projectives, we have that $\mathbf{F}Q$ is projective in \mathcal{B} , and so $P \oplus \mathbf{F}Q$ is also projective in \mathcal{B} . Hence $i^!$ induces a functor $i^! : \mathcal{GP}((\mathbf{F},\mathcal{B})) \to \mathcal{GP}(\mathcal{B})$. By Lemma 4.2, $(i_*,i^!)$ is an adjoint pair.

We claim that there exists a fully faithful functor $j_*: \underline{\mathcal{GP}(A)} \to \underline{\mathcal{GP}((\mathbf{F}, \mathcal{B}))}$ given by $X \to {X \choose P}$ with $P \in \mathcal{P}(\mathcal{B})$, such that there exists an exact sequence

$$0 \longrightarrow \mathbf{F} X \xrightarrow{\phi} P \longrightarrow \operatorname{Coker} \phi \longrightarrow 0$$

in \mathcal{B} with Coker $\phi \in \mathcal{GP}(\mathcal{B})$.

Let $X \in \mathcal{GP}(\mathcal{A})$. By Lemma 4.7, $\mathbf{F}X \in \mathcal{GP}(\mathcal{B})$ and there exists an exact sequence

$$0 \longrightarrow \mathbf{F} X \xrightarrow{\phi} P \longrightarrow \operatorname{Coker} \phi \longrightarrow 0$$

in \mathcal{B} with $P \in \mathcal{P}(\mathcal{B})$ and $\operatorname{Coker} \phi \in \mathcal{GP}(\mathcal{B})$. Let $g: X \to X'$ be a morphism in $\mathcal{GP}(\mathcal{B})$ and $P' \in \mathcal{P}(\mathcal{B})$ such that

$$0 \longrightarrow \mathbf{F} X' \xrightarrow{\phi'} P' \longrightarrow \operatorname{Coker} \phi' \longrightarrow 0$$

is an exact sequence in \mathcal{B} with $\operatorname{Coker} \phi' \in \mathcal{GP}(\mathcal{B})$. Since $\operatorname{Ext}^1_{\mathcal{B}}(\operatorname{Coker} \phi, P') = 0$, we have the following diagram with exact rows

$$0 \longrightarrow \mathbf{F}X \xrightarrow{\phi} P \longrightarrow \operatorname{Coker} \phi \longrightarrow 0$$

$$\downarrow^{\mathbf{F}g} \quad \downarrow^{f} \quad \downarrow$$

$$0 \longrightarrow \mathbf{F}X' \xrightarrow{\phi'} P' \longrightarrow \operatorname{Coker} \phi' \longrightarrow 0.$$

If there exists a morphism $f': P \to P'$ such that $f'\phi = \phi' \mathbf{F} g$, then f' - f factors through Coker ϕ . Since $\operatorname{Coker} \phi \in \mathcal{GP}(\mathcal{B})$, we have a monomorphism $\rho: \operatorname{Coker} \phi \to \widetilde{P}$ with \widetilde{P} projective in \mathcal{B} . Then we easily see that $\binom{g}{f} - \binom{g}{f'}$ factors through the projective object $\binom{0}{\widetilde{P}}$ in $(\mathbf{F}, \mathcal{B})$ and hence $\binom{g}{f} = \binom{g}{f'}$ in $\underline{\mathcal{GP}((\mathbf{F}, \mathcal{B}))}$. Note that if we take $g = \operatorname{id}_X$, this also proves that the object $\binom{X}{P} \in \underline{\mathcal{GP}((\mathbf{F}, \mathcal{B}))}$ is independent of the choice of P. Thus we get a functor $j'_*: \mathcal{GP}(\mathcal{B}) \to \mathcal{GP}((\mathbf{F}, \mathcal{B}))$.

Assume that $g: X \to X'$ in \mathcal{A} factors through a projective object Q with $g = g_2g_1$. Since $\mathbf{F}Q$ is projective in \mathcal{B} by assumption, it is injective in $\mathcal{GP}(\mathcal{B})$, therefore there exists a morphism $\alpha: P \to \mathbf{F}Q$ such that $\mathbf{F}g_1 = \alpha\phi$. Since $(f - \phi'\mathbf{F}g_2\alpha)\phi = 0$, there exists $\widetilde{f}: \operatorname{Coker}\phi \to P'$ such that $(f - \phi'\mathbf{F}g_2\alpha) = \widetilde{f}\pi$. Let $\eta: \operatorname{Coker}\phi \to P_1$ be a monomorphism with $P_1 \in \mathcal{P}(\mathcal{B})$. Then we get $\beta: P_1 \to P'$ such that $\widetilde{f} = \beta\eta$. Thus $\binom{g}{f}$ factors through the projective object $\binom{Q}{FQ} \oplus \binom{0}{P_1}$ in $(\mathbf{F}, \mathcal{B})$ with $\binom{g}{f} = \binom{g_2}{(\phi'\mathbf{F}g_2,\beta)}\binom{g_1}{\binom{\alpha}{\eta\pi}}$. Therefore j'_* induces a functor $j_*: \underline{\mathcal{GP}(\mathcal{B})} \to \underline{\mathcal{GP}((\mathbf{F},\mathcal{B}))}$ which given by $X \to \binom{X}{P}$ and $g \to \binom{g}{f}$. If $\binom{g}{f}$ factors through a projective object $\binom{0}{P} \oplus \binom{Q}{\mathbf{F}Q}$ in $(\mathbf{F}, \mathcal{B})$, then g factors through the projective object Q. Thus Q is fully faithful. The claim is proved.

Let $\binom{g}{f}:\binom{X}{Y}\to\binom{X'}{P}$ be a morphism in $\mathcal{GP}((\mathbf{F},\mathcal{B}))$. By Theorem 3.5, there exists an exact sequence

$$0 \longrightarrow \mathbf{F} X' \xrightarrow{\phi} P \longrightarrow \operatorname{Coker} \phi \longrightarrow 0$$

in \mathcal{B} with P projective and $\operatorname{Coker} \phi \in \mathcal{GP}(\mathcal{B})$. Then $\binom{g}{f}$ factors through the projective object $\binom{0}{P'} \oplus \binom{Q}{\mathbf{F}Q}$ in $(\mathbf{F}, \mathcal{B})$ if and only if $g: X \to X'$ factors through the projective object Q in \mathcal{B} . It follows that the isomorphism

$$\operatorname{Hom}_{\underline{\mathcal{GP}(\mathcal{A})}}(X,X') \cong \operatorname{Hom}_{\underline{\mathcal{GP}((\mathbf{F},\mathcal{B}))}}(\binom{X}{Y},\binom{X'}{P})$$

is natural in both variables and (j^*, j_*) is an adjoint pair.

Finally, applying Lemma 4.4, we get the required recollement.

Acknowledgement. This research was partially supported by NSFC (Grant No. 11571164) and a Project Funded by the Priority Academic Program Development of Jiangsu Higher Education Institutions.

References

- J. Asadollahi and S. Salarian, Gorenstein objects in triangulated categories, J. Algebra 281 (2004), 264–286.
- [2] M. Auslander and M. Bridger, Stable Module Theory, Mem. Amer. Math. Soc. 94, Amer. Math. Soc., Providence, RI, 1969.
- [3] A. Beligiannis and I. Reiten, Homological and Homotopical Aspects of Torsion Theories, Mem. Amer. Math. Soc. 188, Amer. Math. Soc., Providence, RI, 2007.
- [4] A. A. Beĭlinson, J. Bernstein and P. Deligne, Faisceaux Pervers, Astérisque 100, Soc. Math. France, Paris, 1982.
- [5] E. E. Enochs, M. Cortés-Izurdiaga and B. Torrecillas, Gorenstein conditions over triangular matrix rings, J. Pure Appl. Algebra 218 (2014), 1544–1554.
- [6] E. E. Enochs and O. M. G. Jenda, Gorenstein injective and projective modules, Math. Z. 220 (1995), 611–633.
- [7] E. E. Enochs and O. M. G. Jenda, Relative Homological Algebra, Vol. 1, Second revised and extended edition, De Gruyter Expositions in Math. 30, Walter de Gruyter, Berlin, 2011.
- [8] R. M. Fossum, P. A. Griffith and I. Reiten, Trivial Extensions of Abelian Categories (Homological Algebra of Trivial Extensions of Abelian Categories with Applications to Ring Theory), Lect. Notes in Math. 456, Springer-Verlag, Berlin-Heidelberg-New York, 1975.
- [9] V. Franjou and T. Pirashvili, Comparison of abelian categories recollements, Doc. Math. 9 (2004), 41–56.
- [10] D. Happel, Triangulated Categories in the Representation Theory of Finite Dimensional Algebras, Lond. Math. Soc. Lecture Note Ser. 119, Cambridge University Press, Cambridge, 1988.
- [11] O. Iyama, K. Kato and J.-I. Miyachi, Recollement on homotopy categories and Cohen-Macaulay modules, J. K-Theory 8 (2011), 507–541.
- [12] B. Keller, Derived categories and their uses, Handbook of algebra, Vol. 1, Edited by M. Hazewinkel, North-Holland, Amsterdam, 1996, pp.671–701.
- [13] C. Psaroudakis, Homological theory of recollements of abelian categories, J. Algebra 398 (2014), 63–110.

- [14] C. Psaroudakis, Ø. Skartsæterhagen and Ø. Solberg, Gorenstein categories, singular equivalences and finite generation of cohomology rings in recollements, Trans. Amer. Math. Soc. (Ser. B) 1 (2014), 45–95.
- [15] B. L. Xiong and P. Zhang, Gorenstein-projective modules over triangular matrix artin algebras, J. Algebra Appl. 11 (2012), 65–80.
- [16] P. Zhang, Gorenstein-projective modules and symmetric recollements, J. Algebra 388 (2013), 65–80.