Multiple invariants conserving Runge-Kutta type methods for Hamiltonian problems

Luigi Brugnano^{*} Yajuan Sun[†]

Warmly dedicated to celebrate the 80-th birthday of John Butcher

Abstract

In a recent series of papers, the class of energy-conserving Runge-Kutta methods named *Hamiltonian BVMs (HBVMs)* has been defined and studied. Such methods have been further generalized for the efficient solution of general conservative problems, thus providing the class of *Line Integral Methods (LIMs)*. In this paper we derive a further extension, which we name *Enhanced Line Integral Methods (ELIMs)*, more tailored for Hamiltonian problems, allowing for the conservation of multiple invariants of the continuous dynamical system. The analysis of the methods is fully carried out and some numerical tests are reported, in order to confirm the theoretical achievements.

Keywords: Hamiltonian problems, energy-conserving methods, multiple invariants, discrete line-integral methods, HBVMs, LIMs, ELIMs, EHBVMs.

MSC: 65P10, 65L05.

1 Introduction

Hamiltonian problems arise in many fields of application, ranging from the nano-scale of molecular dynamics to the macro-scale of celestial mechanics. Such problems are in the following form:

$$y' = J\nabla H(y), \qquad y(0) = y_0 \in \mathbb{R}^{2m}, \tag{1}$$

where the state vector is often partitioned as

$$y = \begin{pmatrix} q \\ p \end{pmatrix}, \qquad q, p \in \mathbb{R}^m$$

with q the vector of the positions and p the vector of the generalized momenta. Moreover,

$$J = \begin{pmatrix} 0 & I_m \\ -I_m & 0 \end{pmatrix} = -J^T = -J^{-1},$$
(2)

^{*}Dipartimento di Matematica e Informatica "U.Dini", Università di Firenze, Italy (luigi.brugnano@unifi.it)

[†]Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing, China (sunyj@lsec.cc.ac.cn)

and $H(y) \equiv H(q, p)$ is the Hamiltonian function defining the problem. From (1) and (2), it is straightforward to derive that $H(y(t)) \equiv H(y_0)$ for $t \ge 0$, since

$$\frac{\mathrm{d}}{\mathrm{d}t}H(y(t)) = \nabla H(y(t))^T y'(t) = \nabla H(y(t))^T J \nabla H(y(t)) = 0$$

due to the fact that J is skew-symmetric. For isolated mechanical systems, the Hamiltonian has the physical meaning of the total energy of the system and, therefore, it is of interest to derive methods which are able to preserve this property in the discrete solution. For the continuous problem, it can be seen that the *symplecticity* of the map implies the property of energy conservation of the given system, so that a relevant line of investigation, concerning the efficient numerical solution of such problems, has been that of devising *symplectic methods*, namely methods for which the discrete map inherits the property of symplecticity (see, e.g., [19, 32, 33]). In particular, in [32] the existence of infinitely many symplectic Runge-Kutta methods was proved, and an algebraic criterion for symplectic Runge-Kutta methods was provided.

Nevertheless, unless the continuous case, in the discrete setting the symplecticity of the map doesn't imply energy-conservation (see also [11]), so that a different line of investigation has been that of looking for energy-conserving methods. One of the first approaches along this line is represented by *discrete gradient methods* [20, 29], which are based upon the definition of a discrete counterpart of the gradient operator, so that energy conservation for the numerical solution is guaranteed at each step and for any choice of the integration step-size. A different approach is based on the concept of *time finite element methods*, which has led to the definition of energy-conserving Runge-Kutta methods [1, 2, 34, 35], based on a local Galerkin approximations for the equation. A partially related approach is given by discrete line integral methods [26, 27, 28], where the key idea is to exploit the relation between the method itself and the discrete line integral, i.e., the discrete counterpart of the line integral in conservative vector fields. This, in turn, allows exact conservation for polynomial Hamiltonians of arbitrarily high-degree, resulting in the class of methods later named Hamiltonian Boundary Value Methods (HBVMs), which have been developed in a series of papers [8, 9, 10, 7, 11, 12, 13, 4, 5] (we refer to [6] for a systematic presentation of this approach). Another approach, strictly related to the latter one, is given by the *averaged* vector field method [31] and its generalizations [21], which have been also analysed in the framework of B-series [16, 24, 17] (i.e., methods admitting a Taylor expansion with respect to the step-size). In particular, the close connection between the limit formulae of HBVMs and the methods described in [21] has been thoroughly analyzed in [10].

For sake of completeness, we also mention that attempts aiming to obtain methods that, in a weaker sense, have both the property of symplecticity and energy-conservation have been also considered (see, e.g., [25, 14, 36]).

Sometimes, the dynamical system defined by (1) has additional invariants, besides the Hamiltonian. It is therefore interesting to devise methods which are able to preserve all of them in the discrete solution. The approach based on the discrete line integrals, which HBVMs rely on, has been then used to cope with this problem, leading to the class of *Line Integral Methods (LIMs)* which are able to preserve any number of invariants for general conservative problems [3] (see also [6]). In this paper, we consider a different generalization of HBVMs, still based on the concept of discrete line integral, which is able to provide multiple

invariants conserving methods, which are more efficient than LIMs, when the problem is in the form (1). For sake of completeness, we mention that a multiple invariants conserving version of discrete gradients is mentioned in [29] (though without providing any example) and an example of such methods is given in [30], using an antisymmetric tensor taking discrete gradients of all integrals to be preserved as input. Additional multiple invariants conserving methods, obtained by using discrete gradients, are defined in [18].

With this premise, the paper is organized as follows: in Section 2 we recall the basic facts about HBVMs; in Section 3 we define their multiple invariants conserving extension; in Section 4 we provide numerical tests for the new presented methods; finally, in Section 5 we give some conclusions.

2 HBVMs

Let us consider a polynomial approximation to the solution of (1), over the interval [0, h], in the form

$$\sigma'(ch) = \sum_{j=0}^{s-1} P_j(c)\gamma_j(\sigma), \qquad c \in [0,1],$$
(3)

where $\{P_j\}_{j\geq 0}$ is the family of Legendre polynomials, shifted and scaled in order to be orthonormal on the interval [0, 1],

$$\deg P_j = j, \qquad \int_0^1 P_j(x) P_j(x) \mathrm{d}x = \delta_{ij}, \qquad \forall i, j \ge 0.$$
(4)

By imposing the initial condition $\sigma(0) = y_0$, and setting $y_1 \equiv \sigma(h) \approx y(h)$, the coefficients $\gamma_j(\sigma)$ are determined by imposing the conservation of energy at t = h. This implies that

$$0 = H(y_1) - H(y_0) = H(\sigma(h)) - H(\sigma(0)) = \int_0^h \nabla H(\sigma(t))^T \sigma'(t) dt$$

= $h \int_0^1 \nabla H(\sigma(\tau h))^T \sigma'(\tau h) d\tau.$ (5)

By taking into account of (3), one then requires [10]:

$$\sum_{j=0}^{s-1} \left[\int_0^1 P_j(\tau) \nabla H(\sigma(\tau h)) \mathrm{d}\tau \right]^T \gamma_j(\sigma) = 0, \tag{6}$$

which holds true, provided that

$$\gamma_j(\sigma) = \eta_j J \int_0^1 P_j(\tau) \nabla H(\sigma(\tau h)) d\tau, \qquad j = 0, \dots, s-1,$$
(7)

where $\eta_0, \ldots, \eta_{s-1}$ are arbitrary constants. HBVMs are then obtained by setting

$$\eta_j = 1, \qquad j = 0, \dots, s - 1,$$

resulting in an approximation of order 2s to y(h) [10, 12]:

$$\sigma(h) - y(h) = O(h^{2s+1})$$

In particular, by considering the orthonormality of the polynomial basis, one obtains that

$$y_1 \equiv \sigma(h) = y_0 + h\gamma_0(\sigma) = y_0 + \int_0^h J\nabla H(\sigma(t)) dt.$$

This latter expression clearly shows that this polynomial approximation generalizes that defined in [31]. However, the resulting polynomial approximation, given by

$$\sigma(ch) = y_0 + h \sum_{j=0}^{s-1} \int_0^c P_j(x) dx \, \gamma_j(\sigma), \qquad c \in [0,1],$$
(8)

provides an effective numerical method only when the integrals appearing in (7) are conveniently approximated by means of a quadrature formula. If this latter formula is defined at the k Gauss-Legendre points in [0, 1],

$$0 < c_1 < \ldots < c_k < 1, \tag{9}$$

(i.e., $P_k(c_i) = 0, i = 1, ..., k$) and corresponding quadrature weights

$$b_1, \dots, b_k > 0, \tag{10}$$

one then obtains a HBVM(k, s) method which can be cast as a k-stage Runge-Kutta method, with abscissae (9), weights (10), and Butcher matrix given by

$$A = \mathcal{I}_s \mathcal{P}_s^T \Omega, \tag{11}$$

where

$$\mathcal{P}_s = (P_{j-1}(c_i)), \ \mathcal{I}_s = \left(\int_0^{c_i} P_{j-1}(x) \mathrm{d}x\right) \in \mathbb{R}^{k \times s}, \qquad \Omega = \mathrm{diag}(b_1, \dots, b_k).$$

The corresponding polynomial approximation is then given by [10, 12]

$$u(ch) = y_0 + h \sum_{j=0}^{s-1} \int_0^c P_j(x) dx \left[\sum_{\ell=1}^k b_\ell P_j(c_\ell) J \nabla H(u_\ell) \right]$$

$$\equiv y_0 + h \sum_{j=0}^{s-1} \int_0^c P_j(x) dx \, \hat{\gamma}_j, \qquad c \in [0, 1], \qquad (12)$$

where

$$u_{\ell} \equiv u(c_{\ell}h), \qquad \ell = 1, \dots, k, \tag{13}$$

are nothing but the stages of the Runge-Kutta method. It can be proved that [8, 10, 12], for all $k \ge s$, a HBVM(k, s) method:

- has order 2s;
- is symmetric;
- when k = s it reduces to the s-stage Gauss-Legendre method;
- is energy-conserving for all polynomial Hamiltonians of degree not larger than 2k/s. Differently, the error in the Hamiltonian is $O(h^{2k+1})$, when H is suitably regular.

From the last point, a *practical* conservation of the Hamiltonian follows, also considering that the computational complexity of the method is s, independently of k. Indeed, by reformulating the discrete problem generated by the method in terms of the s unknown coefficients $\{\hat{\gamma}_j\}$ appearing in (12), one obtains the discrete problem [11]

$$\boldsymbol{\gamma} = \mathcal{P}_s^T \Omega \otimes J \, \nabla H \left(\mathbf{e} \otimes y_0 + h \mathcal{I}_s \otimes I \, \boldsymbol{\gamma} \right),$$

where

$$\mathbf{e} = \begin{pmatrix} 1\\ \vdots\\ 1 \end{pmatrix} \in \mathbb{R}^k, \qquad \boldsymbol{\gamma} = \begin{pmatrix} \hat{\gamma}_0\\ \vdots\\ \hat{\gamma}_{s-1} \end{pmatrix},$$

which has (block)-size s, independently of k.

It is worth mentioning that, because of the existing relations between the integrals of the polynomials $\{P_j\}$ and the polynomials themselves, matrix (11) can be also written as

$$A = \mathcal{P}_{s+1} \hat{X}_s \mathcal{P}_s^T \Omega, \tag{14}$$

where

$$\hat{X}_{s} = \begin{pmatrix} \frac{1}{2} & -\xi_{1} & & \\ \xi_{1} & 0 & \ddots & \\ & \ddots & \ddots & -\xi_{s-1} \\ & & & \xi_{s-1} & 0 \\ \hline & & & & \xi_{s} \end{pmatrix} \equiv \begin{pmatrix} X_{s} \\ 0 \dots 0 & \xi_{s} \end{pmatrix},$$

with

$$\xi_i = \left(2\sqrt{4i^2 - 1}\right)^{-1}, \qquad i = 1, \dots, s$$

By considering that, for k = s,

$$\mathcal{P}_{s+1} = \left(\begin{array}{cc} \mathcal{P}_s & 0 \end{array} \right), \qquad \mathcal{P}_s^T \Omega = \mathcal{P}_s^{-1},$$

one then sees that (14) can be regarded as a generalization of the W-transformation for collocation methods, as defined by Hairer and Wanner [23, page 79].

3 Multiple invariants conserving HBVMs

We now use again the approach based on line integrals, to define a multiple invariants conserving version of HBVM(k, s) methods. Though the basic idea is similar to that used in [3], nevertheless, the obtained methods are definitely different from those described in that reference: the similarity between the two classes of methods stems from the use of the same, straightforward, methodological tool, given by discrete line integrals [26].

In more details, we now use the fact that energy conservation is gained, with $\gamma_j(\sigma)$ in the form of (7), whichever η_j is. Assume then that

$$L: \mathbb{R}^{2m} \to \mathbb{R}^{\nu} \tag{15}$$

is a set of ν (functionally independent) smooth invariants for the dynamical system (1), besides the Hamiltonian *H*. Consequently, one has

$$\nabla L(y)^T J \nabla H(y) = 0 \in \mathbb{R}^{\nu}, \qquad \forall y, \tag{16}$$

where $\nabla L(y)^T$ is the Jacobian matrix of L. We will now extend the approach described in the previous section, in order to impose their conservation. For sake of simplicity, we shall at first define a polynomial approximation $\sigma \in \Pi_s$ (where $s > \nu$), at a continuous level (i.e., similar to (3)–(8)), then passing to define a fully discrete approximation $u \in \Pi_s$. Clearly, by setting σ in the form (3), we gain energy-conservation by repeating similar steps as done until (8). The difference, in such a case, is obtained by setting

$$\eta_{j} = 1, \qquad j = 0, \dots, s - \nu - 1,$$

$$\eta_{j} = \left[1 - h^{2(s-1-j)}\alpha_{j}\right], \qquad j = s - \nu, \dots, s - 1,$$
(17)

with the coefficients $\{\alpha_j\}$ determined in order to obtain the conservation of the ν additional invariants (15)-(16), even though, in principle, any subset of the ν coefficients $\{\eta_j\}$ could be used for this purpose. By setting, as before, the new approximation

$$y_1 \equiv \sigma(h) \approx y(h),$$

from (3), (7), and (17) one then obtains, by requiring conservation of all invariants,

$$0 = L(y_1) - L(y_0) = L(\sigma(h)) - L(\sigma(0)) = \int_0^h \nabla L(\sigma(t))^T \sigma'(t) dt$$
$$= h \int_0^1 \nabla L(\sigma(\tau h))^T \sigma'(\tau h) d\tau = h \sum_{j=0}^{s-1} \left[\int_0^1 P_j(\tau) \nabla L(\sigma(\tau h)) d\tau \right]^T \gamma_j(\sigma) d\tau$$
$$\equiv h \left[\sum_{j=0}^{s-1} \phi_j(\sigma)^T \bar{\gamma}_j(\sigma) - \sum_{j=s-\nu}^{s-1} h^{2(s-1-j)} \alpha_j \phi_j(\sigma)^T \bar{\gamma}_j(\sigma) \right],$$

where (see (7) and (17)), for all $j \ge 0$:

$$\phi_{j}(\sigma) = \int_{0}^{1} P_{j}(\tau) \nabla L(\sigma(\tau h)) d\tau \in \mathbb{R}^{2m \times \nu},$$

$$\bar{\gamma}_{j}(\sigma) = \int_{0}^{1} P_{j}(\tau) J \nabla H(\sigma(\tau h)) d\tau \in \mathbb{R}^{2m}.$$
(18)

Consequently, energy-conservation is "for free" and, moreover, the conservation of the invariants is gained provided that

$$\sum_{j=s-\nu}^{s-1} h^{2(s-1-j)} \alpha_j \phi_j(\sigma)^T \bar{\gamma}_j(\sigma) = \sum_{j=0}^{s-1} \phi_j(\sigma)^T \bar{\gamma}_j(\sigma).$$
(19)

By defining the matrix

$$\Gamma(\sigma) = \left[h^{2(\nu-1)}\phi_{s-\nu}(\sigma)^T \bar{\gamma}_{s-\nu}(\sigma), \dots, h^0 \phi_{s-1}(\sigma)^T \bar{\gamma}_{s-1}(\sigma) \right] \in \mathbb{R}^{\nu \times \nu}$$
(20)

and the vectors

$$\boldsymbol{\alpha} = \begin{pmatrix} \alpha_{s-\nu} \\ \vdots \\ \alpha_{s-1} \end{pmatrix}, \quad \mathbf{b}(\sigma) = \sum_{j=0}^{s-1} \phi_j(\sigma)^T \bar{\gamma}_j(\sigma) \quad \in \mathbb{R}^{\nu},$$

equation (19) can be recast in vector form as

.

$$\Gamma(\sigma)\boldsymbol{\alpha} = \mathbf{b}(\sigma). \tag{21}$$

The following results then hold true.

Lemma 1 Let $\psi : [0,h] \to V$, with V a vector space, admit a Taylor expansion at 0. Then, for all $j \ge 0$:

$$\int_0^1 P_j(\tau)\psi(\tau h)\mathrm{d}\tau = O(h^j).$$

<u>Proof.</u> By taking into account (4), one obtains:

$$\int_{0}^{1} P_{j}(\tau)\psi(\tau h)d\tau = \int_{0}^{1} P_{j}(\tau)\sum_{n\geq 0} \frac{\psi^{(n)}(0)}{n!}\tau^{n}h^{n}d\tau = \sum_{n\geq 0} \frac{\psi^{(n)}(0)}{n!}h^{n}\int_{0}^{1} P_{j}(\tau)\tau^{n}d\tau$$
$$= \sum_{n\geq j} \frac{\psi^{(n)}(0)}{n!}h^{n}\int_{0}^{1} P_{j}(\tau)\tau^{n}d\tau = O(h^{j}). \Box$$

Lemma 2 If H is suitably regular, then the right-hand side of problem (1) can be expanded as

$$J\nabla H(y(ch)) = \sum_{j\geq 0} P_j(c)\bar{\gamma}_j(y), \qquad c\in[0,1],$$

where $\bar{\gamma}(y)$ is defined according to (18).

<u>Proof.</u> See [13]. \Box

Lemma 3 With reference to (21), one has: $\mathbf{b}(\sigma) = O(h^{2s})$.

<u>Proof.</u> From (16) and (18) one obtains:

$$\sum_{j\geq 0} \phi_j(\sigma)^T \bar{\gamma}_j(\sigma) = 0.$$

Consequently, by virtue of Lemma 1,

$$\mathbf{b}(\sigma) = \sum_{j=0}^{s-1} \phi_j(\sigma)^T \bar{\gamma}_j(\sigma) = -\sum_{j\geq s} \phi_j(\sigma)^T \bar{\gamma}_j(\sigma) = O(h^{2s}). \square$$

Lemma 4 Matrix $\Gamma(\sigma)$ in (21) has $O(h^{2s-2})$ entries.

<u>Proof.</u> The proof follows immediately from (20) and Lemma 1.

In order to simplify the subsequent arguments, we make the following assumption on matrix $\Gamma(\sigma)$:¹

Assumption 1 Matrix $\Gamma(\sigma)$ is nonsingular.

The following result then easily follows from Lemmas 3 and 4.

Theorem 1 Under Assumption 1, the vector $\boldsymbol{\alpha}$ in (21) has $O(h^2)$ entries.

Remark 1 We observe that, in order for Theorem 1 to hold, it is necessary that

$$s > \nu.$$
 (22)

In fact, when $s = \nu$, from (19) one obtains that the products $h^{2(s-1-j)}\alpha_j$ are all equal to 1 and then, from (17) it follows that $\eta_j = 0, j = 0, \ldots, s - 1$. Consequently, in the sequel we shall assume that

$$\eta_0 = 1. \tag{23}$$

We can now state the following result.

Corollary 1 Under Assumption 1, the method conserves all the invariants. Moreover, $\sigma(h) - y(h) = O(h^{2s+1})$.

<u>Proof.</u> The first part of the proof follows from the definition of the method. The second part of the proof strictly follows the technique used in [13]. Let then $y(t; \omega, z)$ be the solution

¹Actually, it would suffice the system (21) to be consistent, but the arguments would become more involved.

of problem (1) satisfying the initial condition $y(\omega) = z$. Moreover, let $\Phi(t, \tau)$ be the corresponding fundamental matrix solution of the associated variational problem. Consequently, from Lemmas 1 and 2, Theorem 1, and from (17), one obtains:

$$\begin{split} \sigma(h) - y(h) &= y(h; h, \sigma(h)) - y(h; 0, \sigma(0)) = \int_0^h \frac{\mathrm{d}}{\mathrm{d}t} y(h; t, \sigma(t)) \mathrm{d}t \\ &= \int_0^h \left[\frac{\partial}{\partial \omega} y(h; \omega, \sigma(t)) |_{\omega=t} + \frac{\partial}{\partial z} y(h; t, z) |_{z=\sigma(t)} \sigma'(t) \right] \mathrm{d}t \\ &= \int_0^h \Phi(h, t) \left[-J \nabla H(\sigma(t)) + \sigma'(t) \right] \mathrm{d}t \\ &= h \int_0^1 \Phi(h, \tau h) \left[-\sum_{j \ge 0} P_j(\tau) \bar{\gamma}_j(\sigma) + \sum_{j=0}^{s-1} P_j(\tau) \gamma_j(\sigma) \right] \mathrm{d}\tau \\ &= -h \int_0^1 \Phi(h, \tau h) \left[\sum_{j \ge s} P_j(\tau) \bar{\gamma}_j(\sigma) + \sum_{j=s-\nu}^{s-1} P_j(\tau) h^{2(s-1-j)} \alpha_j \bar{\gamma}_j(\sigma) \right] \mathrm{d}\tau \\ &= -h \sum_{j \ge s} \left[\underbrace{\int_0^1 P_j(\tau) \Phi(h, \tau h) \mathrm{d}\tau}_{=O(h^j)} \right]_{\bar{\gamma}_j(\sigma)}^{=O(h^j)} - h \sum_{j=s-\nu}^{s-1} \left[\underbrace{\int_0^1 P_j(\tau) \Phi(h, \tau h) \mathrm{d}\tau}_{=O(h^j)} \right] h^{2(s-1-j)} \underbrace{\frac{O(h^{2s+1})}{O(h^{2s+1})} \prod_{j=O(h^{2s+1})}^{O(h^{2s+1})} \right] h^{2(s-1-j)} \underbrace{\frac{O(h^{2s+1})}{O(h^{2s+1})} \prod_{j=O(h^{2s+1})}^{O(h^{2s+1})} \prod_{j=O(h^{2s+1})}^{O(h^{2s+1})} \left[\frac{D(h^{2s+1})}{O(h^{2s+1})} \right] h^{2(s-1-j)} \underbrace{\frac{D(h^{2s+1})}{O(h^{2s+1})}}_{=O(h^{2s+1})} \prod_{j=O(h^{2s+1})}^{O(h^{2s+1})} \prod_{j=O(h^{2s+$$

3.1 Discretization and ELIM(r, k, s) methods

As is clear, the polynomial approximation $\sigma \in \Pi_s$ defined above doesn't yet provide a numerical method: this will be obtained once the integrals in (18) are approximated by means of a suitable quadrature formula. As in the case of LIM(r, k, s) methods in [3], for this purpose we choose the abscissae

$$0 < \hat{c}_1 < \ldots < \hat{c}_r < 1,$$
 (24)

placed at the r Gauss-Legendre points in [0,1], and the corresponding weights

$$\hat{b}_1, \dots, \hat{b}_r > 0, \tag{25}$$

besides (9)–(10) previously considered. In so doing, we obtain a new polynomial approximation, say $u \in \Pi_s$, defined by replacing the integrals with the given quadrature formula, having order 2r or 2k, depending on the chosen abscissae. By setting u_{ℓ} formally defined as in (13), and (see (24))

$$\hat{u}_{\ell} \equiv u(\hat{c}_{\ell}), \qquad \ell = 1, \dots, r,$$

for $0 \le j \le s - 1$ one then obtains:

$$\hat{\phi}_{j} = \sum_{\ell=1}^{r} \hat{b}_{\ell} P_{j}(\hat{c}_{\ell}) \nabla L(\hat{u}_{\ell}) \equiv \phi_{j}(u) - \Psi_{j}(h),$$

$$\hat{\gamma}_{j} = \sum_{\ell=1}^{k} b_{\ell} P_{j}(c_{\ell}) J \nabla H(u_{\ell}) \equiv \bar{\gamma}_{j}(u) - \Delta_{j}(h),$$
(26)

in place of (18) where, by denoting

$$\mu(j) = \lfloor \frac{2j}{s} \rfloor, \qquad j \in \{r, k\},$$
(27)

and assuming L and H suitably regular,

$$\Psi_{j}(h) = \begin{cases} 0, & \text{if } L \in \Pi_{\mu(r)}, \\ O(h^{2r-j}), & \text{otherwise,} \end{cases}$$
(28)

and

$$\Delta_{j}(h) = \begin{cases} 0, & \text{if } H \in \Pi_{\mu(k)}, \\ O(h^{2k-j}), & \text{otherwise.} \end{cases}$$
(29)

Remark 2 Actually, for any invariant in (26) one could use a different quadrature formula, depending on the required accuracy. Nevertheless, for sake of brevity, we shall hereafter consider only the use of two (possibly) different quadratures: (9)–(10) for the $\{\hat{\gamma}_j\}$, and (24)-(25) for the $\{\hat{\phi}_j\}$. However, the generalization is straightforward.

Setting by $\hat{\eta}_j$ and $\hat{\alpha}_j$, respectively, the discrete approximations to (17), the new polynomial approximation is then given by

$$u(ch) = y_0 + h \sum_{j=0}^{s-1} \int_0^c P_j(x) dx \,\hat{\eta}_j \hat{\gamma}_j \qquad (30)$$

$$\equiv y_0 + h \left[\sum_{j=0}^{s-1} \int_0^c P_j(x) dx \,\hat{\gamma}_j - \sum_{j=s-\nu}^{s-1} \int_0^c P_j(x) dx \, h^{2(s-1-j)} \hat{\alpha}_j \hat{\gamma}_j \right],$$

with the scalars $\hat{\alpha}_j$ satisfying the equation (compare with (19)):

$$\sum_{j=s-\nu}^{s-1} h^{2(s-1-j)} \hat{\alpha}_j \hat{\phi}_j^T \hat{\gamma}_j = \sum_{j=0}^{s-1} \hat{\phi}_j^T \hat{\gamma}_j.$$
(31)

Similarly as previously done in (20)–(21), by defining the matrix

$$\hat{\Gamma} = \left[h^{2(\nu-1)} \hat{\phi}_{s-\nu}^T \hat{\gamma}_{s-\nu}, \dots, h^0 \hat{\phi}_{s-1}^T \hat{\gamma}_{s-1} \right] \in \mathbb{R}^{\nu \times \nu}$$
(32)

and the vectors

$$\hat{\boldsymbol{\alpha}} = \begin{pmatrix} \hat{\alpha}_{s-\nu} \\ \vdots \\ \hat{\alpha}_{s-1} \end{pmatrix}, \quad \hat{\mathbf{b}} = \sum_{j=0}^{s-1} \hat{\phi}_j^T \hat{\gamma}_j \quad \in \mathbb{R}^{\nu},$$

equation (31) can be recast in vector form as

$$\hat{\Gamma}\hat{\boldsymbol{\alpha}} = \hat{\mathbf{b}}.\tag{33}$$

Since the number of the additional invariants (16) has to satisfy (22) (and, then, (23) holds true), similarly as in the case of HBVM(k, s), the new approximation is given by

$$y_1 \equiv u(h) = y_0 + h\hat{\gamma}_0 = y_0 + h\sum_{\ell=1}^k b_\ell J \nabla H(u_\ell).$$
(34)

Definition 1 We shall denote by ELIM(r, k, s) (Enhanced LIM(r, k, s)) the methods defined by (30)–(34). In particular, for similarity with the GHBVM(k, s) $\equiv LIM(k, k, s)$ methods in [3], when r = k we shall speak about an EHBVM(k, s) (Enhanced HBVM(k, s)) method.

The following results then easily follow, providing a discrete counterpart of Theorem 1.

Theorem 2 Under Assumption 1, for all $r, k \geq s$ matrix $\hat{\Gamma}$ is nonsingular, for all sufficiently small step-sizes h, and the vector $\hat{\alpha}$ has $O(h^2)$ entries.

We can now state the following results, concerning the order of accuracy of the discrete solution, as well as of the invariants, provided by ELIM(r, k, s) methods.

Theorem 3 Assume that the Hamiltonian function defining problem (1) is a polynomial of degree less than or equal to $\mu(k)$ as defined in (27). Then, ELIM(r, k, s) method is energy-conserving, provided that $r \geq s$. Differently, for all general and suitably regular H, one obtains

$$H(y_1) - H(y_0) = O(h^{2k+1}), \qquad \forall k \ge s,$$

provided that $r \geq s$.

Proof. One has:

$$H(y_{1}) - H(y_{0}) = H(u(h)) - H(u(0)) = \int_{0}^{h} \nabla H(u(t))^{T} u'(t) dt$$

$$= h \int_{0}^{1} \nabla H(u(\tau h))^{T} u'(\tau h) d\tau = h \int_{0}^{1} \nabla H(u(\tau h))^{T} \sum_{j=0}^{s-1} P_{j}(\tau) \hat{\eta}_{j} \hat{\gamma}_{j} d\tau$$

$$= h \sum_{j=0}^{s-1} \left[\int_{0}^{1} \nabla H(u(\tau h)) P_{j}(\tau) d\tau \right]^{T} \hat{\eta}_{j} \hat{\gamma}_{j} = h \sum_{j=0}^{s-1} \hat{\eta}_{j} \bar{\gamma}_{j}(u)^{T} J \hat{\gamma}_{j} = (*).$$

The first part of the proof easily follows from the fact that, if $H \in \Pi_{\mu(k)}$, then

$$\hat{\gamma}_j = \bar{\gamma}_j(u), \qquad j = 0, \dots, s-1,$$

so that (*) = 0, since J is skew-symmetric. In general, assuming that H is suitably regular, one has (see (26) and (29)):

$$(*) = h \sum_{j=0}^{s-1} \hat{\eta}_j \bar{\gamma}_j(u)^T J(\bar{\gamma}_j(u) - \Delta_j(h)) = -h \sum_{j=0}^{s-1} \underbrace{\widehat{\eta}_j}_{=O(h^j)} \underbrace{\bar{\gamma}_j(u)^T}_{=O(h^j)} J \underbrace{\widehat{\Delta}_j(h)}_{=O(h^{2k+1})} = O(h^{2k+1}). \square$$

Using similar arguments, by means of (28) it is possible to prove the following result.

Theorem 4 Assume that the invariants (16) of problem (1) are polynomials of degree less than or equal to $\mu(r)$ as defined in (27). Then, EHBVM(r, k, s) method is invariantsconserving, provided that $k \geq s$. For all general and suitably regular L, one obtains

$$L(y_1) - L(y_0) = O(h^{2r+1}), \qquad \forall r \ge s,$$

provided that $k \geq s$.

Next result concerns the order of accuracy of the numerical solution.

Theorem 5 Assuming that both H and L are suitably regular, for all $r, k \ge s$ the numerical solution generated by a ELIM(r, k, s) method satisfies

$$y_1 - y(h) = O(h^{2s+1}).$$

That is, the method has order 2s.

<u>Proof.</u> The proof proceeds in a similar way as that of Corollary 1. By using the same notation in that corollary, one has:

$$\begin{split} y_1 - y(h) &= y(h; h, u(h)) - y(h, 0, u(0)) = \int_0^h \frac{d}{dt} y(h; t, u(t)) dt \\ &= \int_0^h \left[\frac{\partial}{\partial \omega} y(h; \omega, u(t)) |_{\omega=t} + \frac{\partial}{\partial z} y(h; t, z) |_{z=u(t)} u'(t) \right] dt \\ &= \int_0^h \Phi(h, t) \left[-J \nabla H(u(t)) + u'(t) \right] dt \\ &= h \int_0^1 \Phi(h, \tau h) \left[-\sum_{j \ge 0} P_j(\tau) \bar{\gamma}_j(u) + \sum_{j=0}^{s-1} P_j(\tau) \hat{\eta}_j \hat{\gamma}_j \right] d\tau \\ &= h \int_0^1 \Phi(h, \tau h) \left[-\sum_{j \ge 0} P_j(\tau) \bar{\gamma}_j(u) + \sum_{j=0}^{s-1} P_j(\tau) \hat{\eta}_j(\bar{\gamma}_j(u) - \Delta_j(h)) \right] d\tau \\ &= h \int_0^1 \Phi(h, \tau h) \left[-\sum_{j \ge 0} P_j(\tau) \Delta_j(h) - \sum_{j \ge s} P_j(\tau) \bar{\gamma}_j(u) - \sum_{j=s-\nu} P_j(\tau) h^{2(s-1-j)} \hat{\alpha}_j \bar{\gamma}_j(u) \right] d\tau \\ &= h \int_0^{s-1} \Phi(h, \tau h) \left[-\sum_{j=0}^{s-1} P_j(\tau) \Delta_j(h) - \sum_{j \ge s} P_j(\tau) \bar{\gamma}_j(u) - \sum_{j=s-\nu} P_j(\tau) h^{2(s-1-j)} \hat{\alpha}_j \bar{\gamma}_j(u) \right] d\tau \\ &= h \sum_{j=0}^{s-1} \left[\underbrace{ -\int_0^1 P_j(\tau) \Phi(h, \tau h) d\tau}_{=O(h^{j})} \right]^{=O(h^{2k-j})} \Delta_j(h) - h \sum_{j \ge s} \left[\underbrace{ \int_0^1 P_j(\tau) \Phi(h, \tau h) d\tau}_{=O(h^{j})} \right]^{=O(h^{j})} \hat{\gamma}_j(u) \\ &= O(h^{2k+1}) + O(h^{2s+1}) + O(h^{2s+1}) = O(h^{2s+1}). \Box \end{split}$$

For sake of completeness, we also mention the following result, whose proof is straightforward (see, e.g., [8, 6]). **Theorem 6** Provided that the abscissae (9) and (24) are symmetrically distributed in the interval [0,1], the ELIM(r,k,s) method is symmetric.

3.2 Runge-Kutta type formulation of ELIM(r, k, s) methods

Though the method (30)–(34) is not strictly a Runge-Kutta method, nevertheless, it admits a Runge-Kutta type formulation which is quite useful to represent it. In more details, we already saw that a HBVM(k, s) methods is a k-stage Runge-Kutta method defined by the following Butcher tableau (see (11))

$$\frac{\mathbf{c} \quad \mathcal{I}_s \mathcal{P}_s^T \Omega}{\mathbf{b}^T},$$

where, as usual, **c** is the vector of the abscissae and **b** is the vector of the weights. Moreover, we recall that the only formal difference between a HBVM(k, s) method and an ELIM(r, k, s)method consists in the coefficients $\hat{\eta}_1, \ldots, \hat{\eta}_{s-1}$ which may assume values different from 1 (indeed, $\hat{\eta}_0 = 1$, as stated in (23)). Consequently, by introducing the diagonal matrix

$$\Sigma_s = \operatorname{diag}(1, \hat{\eta}_1, \dots, \hat{\eta}_{s-1}),$$

one obtains the following Runge-Kutta type formulation of an ELIM(r, k, s) method:

$$\frac{\mathbf{c} \quad \mathcal{I}_s \Sigma_s \mathcal{P}_s^T \Omega}{\mathbf{b}^T}.$$

As an example, HBVM(2,2) is the usual 2-stage Gauss method, whereas ELIM(r,2,2) is given by

$$\frac{\frac{1}{2} - \frac{\sqrt{3}}{6}}{\frac{1}{4} + (\hat{\eta}_1 - 1)\frac{\sqrt{3}}{12}} \quad \frac{1}{4} - (\hat{\eta}_1 + 1)\frac{\sqrt{3}}{12}}{\frac{1}{2} + \frac{\sqrt{3}}{6}} \quad \frac{1}{4} + (\hat{\eta}_1 + 1)\frac{\sqrt{3}}{12} \quad \frac{1}{4} - (\hat{\eta}_1 - 1)\frac{\sqrt{3}}{12}}{\frac{1}{2}} \quad \frac{1}{2}$$

As expected, when $\hat{\eta}_1 = 1$ one retrieves the usual 2-stage Gauss method.

4 Numerical tests

We here report a few numerical tests, aimed to assess the theoretical findings, as well as to compare the Enhanced Line Integral Methods (ELIMs), here introduced, with the Line Integral Methods (LIMs) defined in [3]. This will be done on a Hamiltonian problem possessing multiple invariants. The generated discrete problems are solved by means of a fixed-point iteration, even though the efficient implementation of both classes of methods deserves a further investigation.

In order to compare the methods, it will be useful to consider that, for a given problem possessing ν invariants besides the Hamiltonian, one has:

$$\frac{\text{cost of 1 LIM}(r_1, k_1, s) \text{ fixed-point iteration}}{\text{cost of 1 ELIM}(r_2, k_2, s) \text{ fixed-point iteration}} \approx \frac{k_1 + (\nu + 1)r_1}{k_2 + \nu r_2}.$$
(35)

Consequently, when $r_1 = k_1 = r_2 = k_2 \equiv k$, one obtains that²

$$\frac{\text{cost of 1 GHBVM}(k, s) \text{ fixed-point iteration}}{\text{cost of 1 EHBVM}(k, s) \text{ fixed-point iteration}} \approx \frac{\nu + 2}{\nu + 1}.$$
(36)

That said, the problem that we consider is the well known Kepler problem [22, 3], defined by the Hamiltonian

$$H(q,p) = \frac{1}{2} \|p\|_2^2 + \frac{1}{\|q\|_2}, \qquad q, p \in \mathbb{R}^2.$$
(37)

When the initial condition is chosen as

$$(q_0^T, p_0^T) = \left(\begin{array}{ccc} 1 - \varepsilon, & 0, & 0, & \sqrt{\frac{1+\varepsilon}{1-\varepsilon}} \end{array} \right), \qquad \varepsilon \in [0, 1),$$

its solution is periodic, with period 2π , and is given by an ellipse of eccentricity ε in the q-plane. This problem admits two additional (independent) invariants of motion, besides the Hamiltonian (37), given by the *angular momentum*

$$L_1(q,p) = q^T J_2 p, \qquad J_2 = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix},$$
 (38)

and the Laplace-Runge-Lenz (LRL) vector, resulting in the following conserved quantity:

$$L_2(q,p) = (e_1^T p) L_1(q,p) - \frac{e_2^T q}{\|q\|_2},$$
(39)

where, as usual, $e_1, e_2 \in \mathbb{R}^2$ are the two unit vectors.

We solve this problem, considering an eccentricity $\varepsilon = 0.6$, by using the following methods:

- the symplectic 3-stage Gauss method (GAUSS3);
- the (practically) energy-conserving HBVM(12,3) method;
- the EHBVM(12,3) method (i.e., ELIM(12,12,3)) and the GHBVM(12,3) method (i.e., LIM(12,12,3)) in [3], where it is imposed *only* the (practical) conservation of the angular momentum (38) besides the Hamiltonian (37);
- the EHBVM(12,3) and GHBVM(12,3) methods as above, where it is imposed *both* the (practical) conservation of the angular momentum (38) and of the LRL vector (39) besides the Hamiltonian (37).

In Table 1 we list the measured errors after 10 periods, thus confirming that, according to Theorem 5, all methods are sixth-order. Moreover, in Table 3 we list the maximum norm for the vector $\hat{\alpha}$ defined in (33), over the same interval, for the EHBVM(12,3) method:

• by imposing *only* the conservation of the angular momentum besides the Hamiltonian. Here, $\alpha_h^{(1)} = \max_{n=1,\dots,\frac{T}{h}} \|\hat{\boldsymbol{\alpha}}_n\|_{\infty}$;

²We recall that [3] $\text{LIM}(k, k, s) \equiv \text{GHBVM}(k, s)$, and (see Definition 1) $\text{ELIM}(k, k, s) \equiv \text{EHBVM}(k, s)$.

• by imposing *both* the conservation of the angular momentum and of the LRL vector besides the Hamiltonian. As before, $\alpha_h^{(2)} = \max_{n=1,\dots,\frac{T}{h}} \|\hat{\boldsymbol{\alpha}}_n\|_{\infty}$.

The obtained results confirm that the entries of the vector $\hat{\alpha}$ are actually $O(h^2)$, as predicted by Theorem 2. For sake of completeness, in Figure 1 we plot the two components of the vector $\hat{\alpha}$ in the second case, when a step-size $h = \pi/30$ is used: their periodic behavior, in accordance with that of the solution, is clearly evident.

In order to compare the computational costs of EHBVM(12,3) and GHBVM(12,3), in Table 2 we also list the total number of fixed-point iterations needed for solving the discrete problems generated when computing the results listed in Table 1. From Table 2, one sees that GHBVM(12,3) requires approximately the same number of iterations as those needed by GAUSS3 and HBVM(12,3) methods (this fact was already known from [3]), whereas EHBVM(12,3) requires some extra iteration, which increase with the number of conserved invariants. However, according to (36) one fixed-point iteration for GHBVM(12,3), when preserving ν invariants besides the Hamiltonian, costs

$$\frac{2+\nu}{1+\nu}, \qquad \nu = 1, 2,$$

times than that of the corresponding EHBVM(12,3) method. This, in turn, shows that, for the considered problem, EHBVMs are more efficient than GHBVMs.

At last, concerning the conservation of the invariants, by using a constant step-size h = 0.1, we have solved the problem over the interval $[0, 10^3]$, obtaining the following results:

- concerning the conservation of the Hamiltonian (37), all methods are (practically) energy-conserving, except the symplectic 3-stage Gauss method. However, the Hamiltonian error turns out to be bounded, as expected, as confirmed by the plot in Figure 2;
- concerning the conservation of the angular momentum (38), all methods conserve this invariant, except HBVM(12,3). However, the error appears to be bounded, as is shown in Figure 3;
- concerning the conservation of the LRL vector (39), all methods exhibit a drift, except EHBVM(12,3) and GHBVM(12,3), when this invariant is required to be conserved, as is shown in Figure 4. In particular: the drifts of the GAUSS3 and HBVM(12,3) methods are practically the same. Both of them are slightly larger than that shown by the GHBVM(12,3) method which is, in turn, larger than that of EHBVM(12,3) method, when only the invariants (37) and (38) are imposed to be conserved.³

5 Conclusions

In this paper, we have used the technique of *discrete line integrals* introduced by Iavernaro and Pace [26] to define an extension of the energy-conserving methods named HBVMs, in order to cope with the conservation of multiple invariants for Hamiltonian problems. This

³These results agree with the analysis in [15].

Table 1: Error in the numerical solution for the 3-stage Gauss method (E_G) , HBVM(12,3) (E_H) , and EHBVM(12,3) and GHBVM(12,3) with only angular momentum conserved $(E_E^{(1)})$ and $E_G^{(1)}$, respectively), and with both angular momentum and LRL vector conserved $(E_E^{(2)})$ and $E_G^{(2)}$, respectively).

	U ·	-	\$ /									
h	E_G	order	E_H	order	$E_{E}^{(1)}$	order	$E_{G}^{(1)}$	order	$E_{E}^{(2)}$	order	$E_{G}^{(2)}$	order
$\pi/30$	1.942e-03	-	4.587e-05	-	1.017e-05	-	4.049e-05	-	1.928e-05	-	4.367e-05	-
$\pi/60$	2.817e-05	6.1	7.375e-07	6.0	1.644e-07	6.0	6.505e-07	6.0	3.052e-07	6.0	6.868e-07	6.0
$\pi/120$	4.346e-07	6.0	1.161e-08	6.0	2.591e-09	6.0	1.023e-08	6.0	4.785e-09	6.0	1.075e-08	6.0
$\pi/240$	6.771e-09	6.0	1.816e-10	6.0	4.030e-11	6.0	1.599e-10	6.0	7.509e-11	6.0	1.677e-10	6.0
$\pi/480$	1.052e-10	6.0	1.815e-12	6.6	4.718e-13	6.4	2.201e-12	6.2	1.413e-12	5.7	2.346e-12	6.2

Table 2: Total number of fixed-point iterations for solving the discrete problems when using the 3-stage Gauss method (GAUSS3), HBVM(12,3) method, and EHBVM(12,3) and GH-BVM(12,3) with only angular momentum conserved (EHBVM₁(12,3) and GHBVM₁(12,3), respectively), and with both angular momentum and LRL vector conserved (EHBVM₂(12,3) and GHBVM₂(12,3), respectively).

h	GAUSS3	HBVM(12,3)	$EHBVM_1(12,3)$	$GHBVM_1(12,3)$	$EHBVM_2(12,3)$	$GHBVM_2(12,3)$		
$\pi/30$	6705	6775	7256	6779	7474	6781		
$\pi/60$	11147	11244	12691	11247	13407	11249		
$\pi/120$	19085	19343	21664	19339	23012	19348		
$\pi/240$	33876	34752	37511	34743	39117	34753		
$\pi/480$	61501	61959	65125	61967	68217	61970		

has resulted in an "enhanced" version of the *Line Integral Methods (LIMs)* introduced in [3]. Consequently, we have named the new methods *Enhanced Line Integral Methods (ELIMs)*. The analysis of such methods has been carried out, proving that the original order of HBVMs is retained by the new methods. At last, a few numerical tests clearly confirm the theoretical findings.

Acknowledgements

This paper emerged from the visit of the first author at the Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing, China, in December 2012–January 2013. This has been possible because of the support provided by the Academy. The second author was supported by the Foundation for Innovative Research Groups of the NNSFC (11021101).

The authors wish to tank an anonymous referee, for his comments and suggestions, which helped to improve the original manuscript.

References

- [1] P. Betsch, P. Steinmann. Inherently energy conserving time finite elements for classical mechanics. J. Comput. Phys. 160 (2000) 88–116.
- [2] C.L. Bottasso. A new look at finite elements in time: a variational interpretation of Runge–Kutta methods. *Appl. Numer. Math.* **25** (1997) 355–368.

Table 3: Quadratic convergence of the maximum norm of the vector $\hat{\boldsymbol{\alpha}}$, for the Kepler problem, by using the EHBVM(12,3) method, when imposing only the angular momentum conservation $(\alpha_h^{(1)})$, and both angular momentum and LRL vector conservation $(\alpha_h^{(2)})$.

h	$lpha_h^{(1)}$	order	$\alpha_h^{(2)}$	order
$\pi/30$	4.530e-3	—	1.246e-2	—
$\pi/60$	1.155e-3	2.0	3.195e-3	2.0
$\pi/120$	2.902e-4	2.0	8.040e-4	2.0
$\pi/240$	7.265e-5	2.0	2.013e-4	2.0
$\pi/480$	1.837e-5	2.0	5.055e-5	2.0




Figure 1: Components of the vector $\hat{\boldsymbol{\alpha}}$ for the fully conservative EHBVM(12,3) method, $h = \pi/30$.

- [3] L. Brugnano, F. Iavernaro. Line integral methods which preserve all invariants of conservative problems. J. Comput. Appl. Math. 236 (2012) 3905–3919.
- [4] L. Brugnano, F. Iavernaro. Recent advances in the numerical solution of conservative problems. *AIP Conf. Proc.* **1493** (2012) 175–182.
- [5] L. Brugnano, F. Iavernaro. Geometric integration by playing with matrices. AIP Conf. Proc. 1479 (2012) 16–19.
- [6] L. Brugnano, F. Iavernaro. Line integral methods and their application to the numerical solution of conservative problems. arXiv:1301.2367
- [7] L. Brugnano, F. Iavernaro, T. Susca. Numerical comparisons between Gauss-Legendre methods and Hamiltonian BVMs defined over Gauss points. *Monografias de la Real Acedemia de Ciencias de Zaragoza* 33 (2010) 95–112.




Figure 2: Hamiltonian error by using the 3-stage Gauss method, h = 0.1.

- [8] L. Brugnano, F. Iavernaro, D. Trigiante. Analisys of Hamiltonian Boundary Value Methods (HBVMs) for the numerical solution of polynomial Hamiltonian dynamical systems. (2009) arXiv:0909.5659v1
- [9] L. Brugnano, F. Iavernaro, D. Trigiante. Hamiltonian BVMs (HBVMs): A family of "Drift Free" methods for integrating polynomial Hamiltonian problems. *AIP Conf. Proc.* 1168 (2009) 715–718.
- [10] L. Brugnano, F. Iavernaro, D. Trigiante. Hamiltonian Boundary Value Methods (Energy Preserving Discrete Line Methods). JNAIAM J. Numer. Anal. Ind. Appl. Math. 5,1-2 (2010) 17–37.
- [11] L. Brugnano, F. Iavernaro, D. Trigiante. A note on the efficient implementation of Hamiltonian BVMs. J. Comput. Appl. Math. 236 (2011) 375–383.
- [12] L. Brugnano, F. Iavernaro, D. Trigiante. The lack of continuity and the role of infinite and infinitesimal in numerical methods for ODEs: the case of symplecticity. *Appl. Math. Comput.* **218** (2012) 8053–8063.
- [13] L. Brugnano, F. Iavernaro, D. Trigiante. A simple framework for the derivation and analysis of effective one-step methods for ODEs. Appl. Math. Comput. 218 (2012) 8475–8485.
- [14] L. Brugnano, F. Iavernaro, D. Trigiante. Energy and quadratic invariants preserving integrators based upon Gauss collocation formulae. SIAM J. Numer. Anal. 50, 6 (2012) 2897–2916.
- [15] M. Calvo, M.P. Laburta, J.I. Montijano, L. Rández. Error growth in the numerical integration of periodic orbits. *Math. Comput. Simulation* 81 (2011) 2646–2661.
- [16] E. Celledoni, R.I. McLachlan, B. Owren, G.R.W. Quispel. Energy-preserving integrators and the structure of B-series. *Found. Comput. Math.* **10** (2010) 673–693.




Figure 3: Angular momentum error by using the HBVM(12,3) method, h = 0.1.

- [17] E. Celledoni, B. Owren and Y. Sun. The minimal stage, energy preserving Runge-Kutta method for polynomial Hamiltonian systems is the Averaged Vector Field method. *Math. Comp.* (2013) In press.
- [18] M. Dahlby, B. Owren, T. Yaguchi. Preserving multiple first integrals by discrete gradients. J. Phys. A: Math. Theor. 44 (2011) 305205 (14pp).
- [19] Feng Kang. On difference schemes and symplectic geometry. In Proceedings of the 1984 Beijing symposium on differential geometry and differential equations. Science Press, Beijing, 1985, pp. 42–58.
- [20] O. Gonzales. Time integration and discrete Hamiltonian systems. J. Nonlinear Sci. 6 (1996) 449–467.
- [21] E. Hairer. Energy preserving variant of collocation methods. JNAIAM J. Numer. Anal. Ind. Appl. Math. 5,1-2 (2010) 73–84.
- [22] E. Hairer, C. Lubich, G. Wanner. *Geometric Numerical Integration. Structure-Preserving Algorithms for Ordinary Differential Equations*, Second ed., Springer, Berlin, 2006.
- [23] E. Hairer, G.Wanner. Solving Ordinary Differential Equations II. Stiff and Differential-Algebraic Problems, 2nd edition. Springer-Verlag, Berlin, 1996.
- [24] E. Hairer, C.J. Zbinden. On conjugate symplecticity of B-series integrators. IMA Journal of Numerical Analysis 33 (2013) 57–79.
- [25] C. Kane, J.E. Marsden, M. Ortiz, Symplectic-energy-momentum preserving variational integrators, J. Math. Phys. 40 (1999) 3353–3371.
- [26] F. Iavernaro, B. Pace. s-Stage trapezoidal methods for the conservation of Hamiltonian functions of polynomial type. AIP Conf. Proc. 936 (2007) 603–606.




Figure 4: Error in the LRL vector, h = 0.1.

- [27] F. Iavernaro, B. Pace. Conservative Block-Boundary Value Methods for the solution of polynomial Hamiltonian Systems. AIP Conf. Proc. 1048 (2008) 888–891.
- [28] F. Iavernaro, D. Trigiante. High-order symmetric schemes for the energy conservation of polynomial Hamiltonian problems. JNAIAM J. Numer. Anal. Ind. Appl. Math. 4,1-2 (2009) 87–101.
- [29] R.I. McLachlan, G.R.W. Quispel, N. Robidoux. Geometric integration using discrete gradient. *Phil. Trans. R. Soc. Lond. A* 357 (1999) 1021–1045.
- [30] G.R.W. Quispel, H.W. Capel. Solving ODE's numerically while preserving all first integrals. Unpublished report, 1999.
- [31] G.R.W. Quispel, D.I. McLaren. A new class of energy-preserving numerical integration methods. J. Phys. A: Math. Theor. 41 (2008) 045206 (7pp).
- [32] J.M. Sanz-Serna. Runge-Kutta schemes for Hamiltonian systems. BIT 28 (1988) 877– 883.
- [33] Y.B. Suris. On the canonicity of mappings that can be generated by methods of Runge– Kutta type for integrating systems $x'' = \partial U/\partial x$. U.S.S.R. Comput. Math. Math. Phys. **29**, 1 (1989) 138–144.
- [34] Q. Tang, C. Chen. Continuous finite element methods for Hamiltonian systems. Appl. Math. Mech. 28,8 (2007) 1071–1080.
- [35] W. Tang, Y. Sun. Time finite element methods: a unified framework for numerical discretizations of ODEs. Appl. Math. Comput. 219, 4 (2012) 2158–2179.

[36] D. Wang, A. Xiao, X. Li. Parametric symplectic partitioned Runge-Kutta methods with energy-preserving properties for Hamiltonian systems. *Comput. Phys. Comm.* 184 (2013) 303–310.