arXiv:1902.11031v1 [math.RA] 28 Feb 2019

Tensor product of dimension effect algebras

Anna Jenčová and Sylvia Pulmannová*

Abstract

Dimension effect algebras were introduced in (A. Jenčová, S. Pulmannová, Rep. Math. Phys. **62** (2008), 205-218), and it was proved that they are unit intervals in dimension groups. We prove that the effect algebra tensor product of dimension effect algebras is a dimension effect algebra, which is the unit interval in the unital abelian po-groups tensor product of the corresponding dimension groups.

1 Introduction

In [12], the notion of a dimension effect algebra was introduced as a counterpart of the notion of a dimension group. Recall that a dimension group (or a Riesz group) is a directed, unperforated interpolation group. By [5], dimension groups can be also characterized as direct limits of directed systems of simplicial groups. In analogy with the latter characterization, dimension effect algebras were defined as direct limits of directed systems of finite effect algebras with the Riesz decomposition property. It is well known that the latter class of effect algebras corresponds to the class of finite MV-algebras, and in analogy with simplicial groups, we call them simplicial effect algebras. It turns out that dimension effect algebras are exactly the unit intervals in unital dimension groups, and simplicial effect algebras are exactly the unit intervals in unital simplicial groups. In [12], an intrinsic characterization

^{*}Mathematical Institute, Slovak Academy of Sciences, Štefánikova 49, SK-814 73 Bratislava, Slovakia; jenca@mat.savba.sk, pulmann@mat.savba.sk. Supported by grant VEGA No.2/0069/16 and by the grant of the Slovak Research and Development Agency grant No. APVV-16-0073.

of dimension effect algebras was found, and also a categorical equivalence between countable dimension effect algebras and unital AF C*-algebras was shown [12, Theorem 5.2].

In this paper we continue the study of dimension effect algebras. In particular, we study the tensor product of dimension effect algebras in the category of effect algebras. We recall that the tensor product in the category of effect algebras exists, and its construction was described in [3]. We first prove that the tensor product of simplicial effect algebras is again a simplicial effect algebra and is (up to isomorphism) the unit interval in the tensor product of the corresponding unital simplicial groups (Theorem 4.3). Then we extend this result to any dimension effect algebras, using the fact that every dimension effect algebra is a direct limit of a directed system of simplicial effect algebras. Namely, we prove that the tensor product of dimension effect algebras is a dimension effect algebra (Theorem 4.5), and is (up to isomorphism) the unit interval in the tensor product of the corresponding dimension groups (Corollary 4.6). We conjecture that this last statement holds more generally for tensor products of interval effect algebras and their universal groups.

We note that the categorical equivalence between effect algebras with RDP and interpolation groups proved in [12, Theorem 3.8], or the known constructions of tensor products in the category of interval effect algebras [8] cannot be applied here, since the category of effect algebras is much larger than the category of effect algebras with RDP or interval effect algebras.

In the last section, we apply our results to the interval $\mathbb{R}^+[0,1]$ and construct a directed system of simplicial groups that has this interval as its direct limit.

2 Preliminaries

The notion of an effect algebra was introduced by D.J. Foulis and M.K. Bennett in [6]. An alternative definition of so called *D*-poset was introduced in [13]. Effect algebras and D-posets are categorically equivalent structures [4].

2.1 Definition. An *effect algebra* is an algebraic system $(E; 0, 1, \oplus)$, where \oplus is a partial binary operation and 0 and 1 are constants, such that the following axioms are satisfied for $a, b, c \in E$:

- (i) if $a \oplus b$ is defined the $b \oplus a$ is defined and $a \oplus b = b \oplus a$ (commutativity);
- (ii) if $a \oplus b$ and $(a \oplus b) \oplus c$ are defined, then $a \oplus (b \oplus c)$ is defined and $(a \oplus b) \oplus c = a \oplus (b \oplus c)$ (associativity);
- (iii) for every $a \in E$ there is a unique $a^{\perp} \in E$ such that $a \oplus a^{\perp} = 1$;
- (iv) if $a \oplus 1$ is defined then a = 0.

In what follows, if we write $a \oplus b$, $a, b \in E$, we tacitly assume that $a \oplus b$ is defined in E. The operation \oplus can be extended to the \oplus -sum of finitely many elements by recurrence in an obvious way. Owing to commutativity and associativity, the element $a_1 \oplus a_2 \oplus \cdots \oplus a_n$ is ambiguously defined. In any effect algebra a partial order can be defined as follows: $a \leq b$ if there is $c \in E$ with $a \oplus c = b$. In this partial order, 0 is the smallest and 1 is the greatest element in E. Moreover, if $a \oplus c_1 = a \oplus c_2$, then $c_1 = c_2$, and we define $c = b \oplus a$ iff $a \oplus c = b$. In particular, $1 \oplus a = a^{\perp}$ is called the *orthosupplement* of a. We say that $a, b \in E$ are *orthogonal*, written $a \perp b$, iff $a \oplus b$ exists in E. It can be shown that $a \perp b$ iff $a \leq b^{\perp}$. An effect algebra which is a lattice with respect to the above ordering is called a *lattice effect algebra*.

Let *E* and *F* be effect algebras. A mapping $\phi : E \to F$ is an *effect algebra* morphism iff $\phi(1) = 1$ and $\phi(e \oplus f) = \phi(e) \oplus \phi(f)$ whenever $e \oplus f$ is defined in *E*. The category of effect algebras with effect algebra morphisms will be denoted by **EA**.

2.1 Interval effect algebras and RDP

Important examples of effect algebras are obtained in the following way. Let $(G, G^+, 0)$ be a (additively written) partially ordered abelian group with a positive cone G^+ and neutral element 0. For $a \in G^+$ define the interval $G[0, a] := \{x \in G : 0 \le x \le a\}$. Then G[0, a] can be endowed with a structure of an effect algebra by defining $x \perp y$ iff $x + y \le a$, and then putting $a \oplus b := a + b$. Effect algebras obtained in this way are called *interval effect algebras*. We note that a prototype of effect algebras is the interval [0, I] in the group of self-adjoint operators on a Hilbert space, so-called algebra of Hilbert space effects. Hilbert space effects play an important role in quantum measurement theory, and the abstract definition was motivated by this

example. On the other hand, there are effect algebras that are not interval effect algebras, see e.g. [14].

The partially ordered abelian group G is directed if $G = G^+ - G^+$. An element $u \in G^+$ is an order unit if for all $a \in G$, $a \leq nu$ for some $n \in \mathbb{N}$. If G has an order unit u, it is directed, indeed, if $g \leq nu$, then g = nu - (nu - g). An element $u \in G^+$ is called a generating unit if every $a \in G^+$ is a finite sum of (not necessarily different) elements of the interval G[0, u]. Clearly, a generating unit is an order unit, the converse may be false.

If G and H are partially ordered abelian groups, then a group homomorphism $\phi: G \to H$ is *positive* if $\phi(G^+) \subseteq H^+$. An isomorphism $\phi: G \to H$ is an *order isomorphism* if $\phi(G^+) = H^+$. If G and H have order units u and v, respectively, then a positive homomorphism $\phi: G \to H$ is called *unital* if $\phi(u) = v$. The category of partially ordered abelian groups having an order unit, with positive unital homomorphisms will be denoted by **POG**.

Relations between interval effect algebras and partially ordered abelian groups are described in the following theorem, proved in [1]. Recall that a mapping $\phi : E \to K$, where E is an effect algebra and K is any abelian group, is called a *K*-valued measure on E if $\phi(a \oplus b) = \phi(a) + \phi(b)$ whenever $a \oplus b$ is defined in E.

2.2 Theorem. Let E be an interval effect algebra. Then there exists a unique (up to isomorphism) partially ordered directed abelian group (G, G^+) and an element $u \in G^+$ such that the following conditions are satisfied:

- (i) E is isomorphic to the interval effect algebra $G^+[0, u]$.
- (ii) *u* is a generating unit.
- (iii) Every K-valued measure $\phi : E \to K$ can be extended uniquely to a group homomorphism $\phi^* : G \to K$.

The group G in the preceding theorem is called a *universal group* for E, and will be denoted by G_E . In what follows we consider a property that ensures that a partially ordered group with order unit is the universal group for its unit interval. There are examples (see [7, Example 11.3, 11.5]) that show that this is not true in general.

A partially ordered abelian group G is said to have the *Riesz interpolation* property (RIP), or to be an *interpolation group*, if given a_i, b_j $(1 \le i \le m, 1 \le j \le n)$ with $a_i \le b_j$ for all i, j, there exists $c \in G$ such that $a_i \le c \le b_j$ for all *i*, *j*. The Riesz interpolation property is equivalent to the Riesz decomposition property (RDP): given $a_i, b_j \in G^+$, $(1 \le i \le m, 1 \le j \le n)$ with $\sum a_i = \sum b_j$, there exist $c_{ij} \in G^+$ with $a_i = \sum_j c_{ij}, b_j = \sum_i c_{ij}$. An equivalent definition of the RDP is as follows: given a, b_i in G^+ , $i \le n$ with $a \le \sum_{i \le n} b_i$, there exist $a_i \in G^+$ with $a_i \le b_i, i \le n$, and $a = \sum_{i \le n} a_i$. To verify these properties, it is only necessary to consider the case m = n = 2 (cf. [9, 10]).

For interpolation groups we have the following theorem [16], [12, Theorem 3.5].

2.3 Theorem. Let G be an interpolation group with order unit u. Put $E := G^+[0, u]$. Then (G, u) is the universal group for E.

In a similar way as for partially ordered abelian groups, RDP can be defined for effect algebras. We say that an effect algebra E has the *Riesz* decomposition property (RDP) if one of the following equivalent properties is satisfied:

(R1)
$$a \leq b_1 \oplus b_2 \oplus \cdots \oplus b_n$$
 implies $a = a_1 \oplus a_2 \oplus \cdots \oplus a_n$ with $a_i \leq b_i, i \leq n$;

(R2) $\bigoplus_{i \leq m} a_i = \bigoplus_{j \leq m} b_j$, $m, n \in \mathbb{N}$, implies $a_i = \bigoplus_j c_{ij}$, $i \leq m$, and $b_j = \bigoplus_i c_{ij}$, $j \leq n$, where $c_{ij} \in E$.

Similarly as for partially ordered groups, it suffices to consider the case m = n = 2.

Let us remark that RIP can be also defined for effect algebras. In contrast with the case of partially ordered abelian groups, RIP and RDP are not equivalent for effect algebras: RDP implies RIP, but there are examples of effect algebras with RIP which do not have RDP (e.g., the "diamond" is lattice ordered effect algebra that does not satisfy RDP, [4]).

It was proved by Ravindran [18], that every effect algebra with RDP is an interval effect algebra, and its universal group is an interpolation group. Ravindran's result can be extended to a categorical equivalence between the category of effect algebras with RDP with effect algebra morphisms and the category of interpolation groups with order unit with positive unital group homomorphisms, [12, Theorem 3.8].

3 Dimension groups and dimension effect algebras

In this section, we study dimension groups and their effect algebra counterpart, introduced in [12]. These are interpolation groups with some additional properties.

A partially ordered abelian group G is called *unperforated* if given $n \in \mathbb{N}$ and $a \in G$, then $na \in G^+$ implies $a \in G^+$. Every Archimedean, directed abelian group is unperforated [10, Proposition 1.24], and also every lattice ordered abelian group is unperforated [10, Proposition 1.22].

3.1 Definition. [10] A partially ordered group G is a *dimension group* (or a *Riesz group*) if it is directed, unperforated and has the interpolation property.

A simple example of a dimension group is as follows.

3.2 Definition. [10, Definition p. 183], [11] A simplicial group is any partially ordered abelian group that is isomorphic (as partially ordered abelian group) to \mathbb{Z}^n (with the product ordering) for some nonnegative integer n. A simplicial basis for a simplicial group G is any basis (x_1, \ldots, x_n) for G as a free abelian group such that $G^+ = \mathbb{Z}^+ x_1 + \cdots + \mathbb{Z}^+ x_n$.

It was proved by Effros, Handelman and Shen [5] that the dimension groups with order unit are precisely the direct limits of directed systems of simplicial groups with an order unit in the category **POG**.

Note that an element $v \in \mathbb{Z}^r$ is an order unit if and only if all of its coordinates are strictly positive. In this case, the interval $(\mathbb{Z}^+)^r[0, v]$ is the direct product of finite chains $(0, 1, \ldots, v_i), i = 1, 2, \ldots, r$ and therefore is a finite effect algebra with RDP. Conversely, every finite effect algebra with RDP is a unit interval in a simplicial group. Below, such effect algebras will be called *simplicial*.

In analogy with dimension groups, in [12], direct limits of directed systems of simplicial effect algebras have been called *dimension effect algebras*. It was shown that an effect algebra is a dimension effect algebra if and only if its universal group is a dimension group. An intrinsic characterization of dimension effect algebras was found in [12, Theorem 4.2].

For the convenience of the readers, we give a short description of the directed system and direct limit of effect algebras [4, Definition 1.9.36].

A directed system of effect algebras is a family $A_I := (A_i; (f_{ij} : A_j \to A_i, i, j \in I, j \leq i)$ where (I, \leq) is a directed set, A_i is an effect algebra for each $i \in I$, and f_{ij} is a morphism such that

- (i1) $f_{ii} = id_{A_i}$ fir every $i \in I$;
- (i2) if $m \leq j \leq i$ in *I*, then $f_{ij}f_{jm} = f_{im}$.

Let A_I be a directed system of effect algebras, then $\underline{f} := (A; (f_i : A_i \to A; i \in I))$ is called the *direct limit* of A_I iff the following conditions hold:

- (ii1) A is an effect algebra; f_i is a morphism for each $i \in I$;
- (ii2) if $j \leq i$ in I, then $f_i f_{ij} = f_j$ (i.e., f is compatible with A_I);
- (ii3) if $\underline{g} := (B; (g_i : A_i \to B, i \in I))$ is any system compatible with A_I , then there exists exactly one morphism $g : A \to B$ such that $gf_i = g_i$, for every $i \in I$.

It was proved (cf. [4, Theorem 1.9.27]) that the direct limit in the category of effect algebras exists. A sketch of the construction of the direct limit is as follows. Let $A = \bigcup_{i \in I} A_i$ be the disjoint union of $A_i, i \in I$. Define a relation \equiv on A as follows. Put $a \equiv b$ ($a \in A_i, b \in A_j$) if there exists a $k \in I$ with $i, j \leq k$ such that $f_{ki}(a) = f_{kj}(b)$ in A_k . Then \equiv is an equivalence relation, and the quotient $\overline{A} := A / \equiv$ can be organized into an effect algebra with the operation \oplus defined as follows: let \overline{a} denotes the equivalence class corresponding to a. For $a \in A_I, b \in A_j, \ \overline{a} \oplus \overline{b}$ is defined iff there is $k \in I, \ i, j \leq k$ such that $f_{ki}(a) \oplus f_{kj}(b)$ exists in A_k , and then $\overline{a} \oplus \overline{b} = \overline{(f_{ki}(a) \oplus f_{kj}(b))}$ in \overline{A} . For every $i \in I$, define $f_i : A_i \to A / \equiv$ as the natural projection $f_i(a) = \overline{a}$. Then $\lim_{\to} A := (\overline{A}; f_i : A_i \to \overline{A}, i \in I)$ is the desired direct limit.

From this construction, it can be derived that properties involving finite number of elements, such as RDP or being a dimension effect algebra (cf. the characterization in [12, Thm. 4.2]), are preserved under direct limits in **EA**.

4 Tensor product of dimension effect algebras

The tensor product in the category \mathbf{EA} is defined below as an universal bimorphism. We will show that such a tensor product always exists and that it is essentially given by the construction in [3], see also [4, Chap. 4.2].

Let E,F,L be effect algebras. A mapping $\beta:E\times F\to L$ is called a bimorphism if

- (i) $a, b \in E$ with $a \perp b, q \in F$ imply $\beta(a,q) \perp \beta(b,q)$ and $\beta(a \oplus b,q) = \beta(a,q) \oplus \beta(b,q)$;
- (ii) $c, d \in F$ with $c \perp d, p \in E$ imply $\beta(p, c) \perp \beta(p, d)$ and $\beta(p, (c \oplus d)) = \beta(p, c) \oplus \beta(p, d);$
- (iii) $\beta(1,1) = 1$.

4.1 Definition. Let E and F be effect algebras. A pair (T, τ) consisting of an effect algebra T and a bimorphism $\tau : E \times F \to T$ is said to be the *tensor* product of E and F if whenever L is an effect algebra and $\beta : E \times F \to L$ is a bimorphism, there exists a unique morphism $\phi : T \to L$ such that $\beta = \phi \circ \tau$.

It is clear that if the tensor product exists, it is unique up to isomorphism. We will use the notation $E \otimes F$ for the effect algebra T and \otimes for the bimorphism τ : $\tau(e, f) = e \otimes f \in E \otimes F$.

4.2 Theorem. The tensor product always exists in EA.

Proof. The theorem was essentially proved in [3, Theorem 7.2], see also [4, Theorem 4.2.2]. There a somewhat different definition of a tensor product is considered and the bimorphisms are assumed nontrivial, that is, the target algebra is required to satisfy $0 \neq 1$. If at least one such bimorphism exists, it is easy to see that [3] provides a construction of a tensor product in our sense. On the other hand, if there are no nontrivial bimorphisms, then the tensor product is given by the one-element effect algebra $\{0 = 1\}$ and the unique bimorphism $E \times F \to \{0\}$.

The tensor product of dimension groups in the category **POG** was studied by Goodearl and Handelman [11] and it was proved that such a tensor product is a dimension group as well. Recall that the tensor product of G_1 and G_2 in **POG** can be constructed as the abelian group tensor product $G_1 \otimes G_2$, endowed with the positive cone $G_1^+ \otimes G_2^+$, generated by simple tensors of positive elements.

Our aim in this section is to describe the tensor product of dimension effect algebras in the category **EA**. Note that we cannot directly apply the

above result via the categorical equivalence of [12, Theorem 3.8], since the category **EA** is much larger than the category of effect algebras with RDP.

We first consider the case of simplicial effect algebras. Let E and F be simplicial effect algebras, with atoms

$$(e_1,\ldots,e_n),$$
 (f_1,\ldots,f_m)

and unit elements

$$u = \sum_{i} u_i e_i, \qquad v = \sum_{j} v_j f_j,$$

respectively. Then G_E and G_F are simplicial groups and $G_E \otimes G_F$ is a simplicial group with generators

$$g_{ij} = e_i \otimes f_j, i = 1, \dots, n; j = 1, \dots, m.$$

Hence the unit interval $G_E \otimes G_F[0, u \otimes v]$ is a simplicial effect algebra with atoms g_{ij} and and unit element $w = \sum_{i,j} u_i v_j g_{ij}$.

4.3 Theorem. Tensor product of simplicial effect algebras in the category **EA** is a simplicial effect algebra, namely

$$E \otimes F \simeq G_E \otimes G_F[0, u \otimes v].$$

Proof. Let G denote the simplicial effect algebra on the right hand side. Obviously, (bi)morphisms on simplicial effect algebras are uniquely determined by their values on the atoms. Let $\tau : E \times F \to G$ be the bimorphism determined by

$$\tau(e_i, f_j) = g_{ij}, \qquad i = 1, \dots, n, \ j = 1, \dots, m.$$

We need to prove that for any effect algebra H and bimorphism $\beta : E \times F \to H$, there is a morphism $\psi : G \to H$, such that

$$\psi(g_{ij}) = \beta(e_i, f_j), \qquad i = 1, \dots, n, \ j = 1, \dots, m.$$

Since g_{ij} generate G, uniqueness of such a morphism is clear. So let $z \in G$, then $z = \sum_{i,j} z_{ij} g_{ij}$, for $z_{ij} \leq u_i v_j$ for all i and j. There are nonnegative integers q_{ij} , r_{ij} such that

$$z_{ij} = v_j q_{ij} + r_{ij}, \qquad r_{ij} < v_j,$$

then since $v_j q_{ij} \leq z_{ij} \leq u_i v_j$, we have $q_{ij} \leq u_i$, with equality only if $r_{ij} = 0$. Then $a_j := \sum_i q_{ij} e_i \in E$ and $r_{ij} f_j \in F$. We have

$$z = \sum_{j} \left(\sum_{i} q_{ij} v_j g_{ij} + \sum_{i} r_{ij} g_{ij}\right)$$
$$= \sum_{j} \tau(a_j, v_j f_j) + \sum_{i, r_{ij} > 0} \tau(e_i, r_{ij} f_j)$$

Put $a'_j := \sum_{i,r_{ij}>0} e_i$, then $a_j \perp a'_j$. Now we can write

$$\begin{aligned} H \ni 1 &= \beta(u, v) = \sum_{j} \beta(u, v_{j}f_{j}) \\ &= \sum_{j} \left[\beta(a_{j}, v_{j}f_{j}) + \beta(a'_{j}, v_{j}f_{j}) + \beta(u - (a_{j} + a'_{j}), v_{j}f_{j}) \right] \\ &= \sum_{j} \left[\beta(a_{j}, v_{j}f_{j}) + \sum_{i} \beta(e_{i}, r_{ij}f_{j}) + \sum_{i, r_{ij} > 0} \beta(e_{i}, (v_{j} - r_{ij})f_{j}) \right. \\ &+ \beta(u - (a_{j} + a'_{j}), v_{j}f_{j}) \right] \end{aligned}$$

It follows that

$$\sum_{i,j} z_{ij}\beta(e_i, f_j) = \sum_{i,j} [q_{ij}v_j\beta(e_i, f_j) + r_{ij}\beta(e_i, f_j)]$$
$$= \sum_j [\beta(a_j, v_j f_j) + \sum_i \beta(e_i, r_{ij} f_j)]$$

is a well defined element in H and we may put

$$\psi(z) = \sum_{i,j} z_{ij}\beta(e_i, f_j),$$

which clearly defines a morphism $G \to H$.

Let

$$A_I = (A_i; (f_{ij} : A_j \to A_i); i, j \in I, j \leq i),$$

$$B_J = (B_k; (g_{k\ell} : B_\ell \to B_k); k, \ell \in J, \ell \leq k)$$

be directed systems of simplicial effect algebras. Let us define the index set (\mathcal{I}, \leq) as the product $I \times J$ with pointwise ordering. By the previous theorem,

each $A_i \otimes B_k$, $(i,k) \in \mathcal{I}$ is a simplicial effect algebra. Let $(j,\ell) \in \mathcal{I}$ be such that $(j,\ell) \leq (i,k)$, then we have morphisms $f_{ij}: A_j \to A_i$ and $g_{k\ell}: B_\ell \to B_k$. For $a \in A_j$, $b \in B_\ell$, put $\beta(a,b) = f_{ij}(a) \otimes g_{k\ell}(b) \in A_i \otimes B_k$, this defines a bimorphism $A_j \times B_\ell \to A_i \otimes B_k$. By properties of tensor product, this extends to a unique morphism $f_{ij} \otimes g_{k\ell}: A_j \otimes B_\ell \to A_i \otimes B_k$.

4.4 Theorem. Let

$$A_I \otimes B_J := (A_i \otimes B_k; (f_{ij} \otimes g_{k\ell} : A_j \otimes B_\ell \to A_i \otimes B_k), (i, k), (j, \ell) \in \mathcal{I}, (j, \ell) \leq (i, k)).$$

Then $A_I \otimes B_J$ is a directed system of simplicial effect algebras.

Proof. We have to check properties (i1) and (i2). For (i1), note that $f_{ii} = id_{A_i}, g_{kk} = id_{B_k} \text{ imply } f_{ii} \otimes g_{kk} = id_{A_i \otimes B_k}$. For (i2), let $(m, n) \leq (j, \ell) \leq (i, k)$. Then

$$m \le j \le i \implies f_{ij}f_{jm} = f_{im}$$
$$n \le \ell \le k \implies g_{k\ell}g_{\ell n} = g_{kn}$$

and for $a_m \in A_m, b_n \in B_n$,

$$(f_{ij} \otimes g_{k\ell})(f_{jm} \otimes g_{\ell n})(a_m \otimes b_n) = (f_{ij} \otimes g_{k\ell})(f_{jm}(a_m) \otimes g_{\ell n}(b_n))$$

$$= f_{ij}f_{jm}(a_m) \otimes g_{k\ell}g_{\ell n}(b_n)$$

$$= f_{im}(a_m) \otimes g_{kn}(b_n)$$

$$= f_{im} \otimes g_{kn}(a_m \otimes b_n).$$

Since this holds on simple tensors, it extends to whole $A_m \otimes B_n$.

4.5 Theorem. Let A_I, B_J be directed systems of simplicial effect algebras, and let $(\bar{A}; (f_i : A_i \to \bar{A}, i \in I))$ and $(\bar{B}; (g_j : B_j \to \bar{B}, j \in J))$ be their corresponding direct limits. Then $(\bar{A} \otimes \bar{B}; (f_i \otimes g_j : A_i \otimes B_j \to \bar{A} \otimes \bar{B}, i \in I, j \in J))$ is the direct limit of $A_I \otimes B_J$.

Proof. We have to check properties (ii1), (ii2) and (ii3). The first one is clear: since \overline{A} , \overline{B} are effect algebras, $\overline{A} \otimes \overline{B}$ is an effect algebra as well. To prove compatibility, let $(j, \ell) \leq (i, k)$. Then $j \leq i, \ell \leq k$ and we have

$$(f_i \otimes g_k)(f_{ij} \otimes g_{k\ell}) = f_i f_{ij} \otimes g_k g_{k\ell} = f_j \otimes g_\ell.$$

For (ii3), let $(C; (h_{ij} : A_i \otimes B_j \to C, i \in I, j \in J))$ be another system compatible with $A_I \otimes B_J$ (i.e., $h_{ik}(f_{ij} \otimes g_{k\ell}) = h_{j\ell}, j \leq i, \ell \leq k$). Let $a \in \bar{A}$, $b \in \bar{B}$. Since \bar{A} and \bar{B} are direct limits, there are some indices $i \in I, k \in J$ and elements $a_i \in A_i$ and $b_k \in B_k$ such that $a = f_i(a_i)$ and $b = g_k(b_k)$, see the construction on page 7. Define $h(a, b) := h_{ik}(a_i \otimes b_k)$. Then $h : \bar{A} \times \bar{B} \to C$ is a bimorphism, which extends to a morphism $\bar{h} : \bar{A} \otimes \bar{B} \to C$.

4.6 Corollary. Let E and F be dimension effect algebras, and let G_E and G_F be their universal groups with units u_E and u_F . Then the tensor product $E \otimes F$ is isomorphic to the unit interval $[0, u_E \otimes u_F]$ in the tensor product $G_E \otimes G_F$ of their universal groups, that is

$$G_E[0, u_E] \otimes G_F[0, v_F] \simeq G_E \otimes G_F[0, u_E \otimes u_F]$$

Proof. Let $E = \overline{A}$, $F = \overline{B}$ be direct limits of directed systems A_I and B_J . Each A_i , $i \in I$ and B_k , $k \in J$ is a simplicial effect algebra and G_{A_i} , G_{B_k} are simplicial groups. By [12, Theorem 4.1], we obtain that G_E is a direct limit of (G_{A_i}, f_{ij}^*) , where f_{ij}^* are the unique morphisms in **POG**, extending f_{ij} , similarly for G_F .

By Theorem 4.3, $A_i \otimes B_k$ is a simplicial effect algebra and $G_{A_i \otimes B_k} \simeq G_{A_i} \otimes G_{B_k}$. By Theorem 4.5, $E \otimes F$ is the direct limit of the directed system $A_I \otimes B_J$. Since $A_I \otimes B_J$ has RDP, it follows by [12, Theorem 4.1] that the universal group $G_{E \otimes F}$ is a direct limit of the system of universal groups

$$\{G_{A_i\otimes B_k}\simeq G_{A_i}\otimes G_{B_k}, (f_{ij}\otimes g_{k\ell})^*\simeq f_{ij}^*\otimes g_{k\ell}^*\},\$$

Using [11, Lemma 2.2], we obtain

$$G_{E\otimes F}\simeq G_E\otimes G_F, \qquad u_{E\otimes F}=u_E\otimes u_F.$$

5 Conclusions and a conjecture

We have proved that the **EA** tensor product of dimension effect algebras is again a dimension effect algebra. The tensor product $E \otimes F$ is proved to be the direct limit of a directed system of simplicial effect algebras, obtained as a "tensor product" of the directed systems corresponding to dimension effect algebras E and F. It is also proved that $E \otimes F$ is (isomorphic to) the unit interval in the **POG** tensor product of the corresponding universal groups G_E and G_F . We conjecture that this is true for general interval effect algebras. Note that in the category of *interval* effect algebras, the tensor product exists [8, Theorem 9.1] and our conjecture says that it is (isomorphic to) the **EA** tensor product.

A special class of interval effect algebras are the algebras with RDP. It is again an open question whether in this case the **EA** tensor product has RDP. If our conjecture is true, $E \otimes F$ is the unit interval in the **POG** tensor product of groups with RDP. As it was shown in [19, cf. Remark 2.13], the **POG** tensor product of groups with RDP might not have RDP, but in the presence of generating units, RDP holds in an asymptotic form in the sense of [15].

6 An example: $\mathbb{R}[0,1]$

Let us consider the interval [0, 1] in $(\mathbb{R}, \mathbb{R}^+, 0)$. This is clearly a dimension group with order unit 1 and hence the interval [0, 1] is a dimension effect algebra. It was proved in [17] that the **EA** tensor product $[0, 1] \otimes [0, 1]$ is not lattice ordered and thus not isomorphic to [0, 1]. By our results, $[0, 1] \otimes [0, 1]$ is a dimension effect algebra, which is the interval $R \otimes R[0, 1 \otimes 1]$. Note that the fact that the **POG** tensor product $R \otimes R$ is not lattice ordered was shown in [19].

As an example, we will present [0, 1] as a direct limit of a directed system of simplicial effect algebras. The tensor product $[0, 1] \otimes [0, 1]$ is then obtained as a direct limit as in Theorem 4.5.

We first need to introduce some notations. For any n-tuple

$$A = (x_1, \dots, x_n)$$

of elements in \mathbb{R}^+ , let f_A denote the positive group homomorphism

$$f_A: \mathbb{Z}^n \to \mathbb{R}, \quad e_i^n \mapsto x_i, \ i = 1, \dots, n$$

and let

$$L(A) := f_A(\mathbb{Z}^n), \quad L(A)^+ := f_A((\mathbb{Z}^n)^+), \quad L_>(A)^+ := f_A((\mathbb{Z}^n)^+),$$

where $(Z_{>}^{n})^{+} := \{\sum_{i} z_{i} e_{i}^{n} \text{ with } z_{i} > 0 \text{ for all } i = 1, \dots, n\}$. We also use the notations

$$Q(A) := Lin_{\mathbb{Q}}(A), \quad Q(A)^+ := Q(A) \cap \mathbb{R}^+$$

Let us define the index set as

 $\mathcal{I} := \{ A \subset [0, 1], \text{finite}, \mathbb{Q} - \text{linearly independent}, \ 1 \in L_{>}(A)^{+} \}.$

Any $A \subset \mathbb{R}^+$ with cardinality n can be identified with the n-tuple of its elements (x_1, \ldots, x_n) , indexed so that $x_1 < \cdots < x_n$. For $A, B \in \mathcal{I}$, write $B \preceq A$ if $B \subset L(A)^+$. It is easy to see that \preceq is a preorder in \mathcal{I} .

6.1 Proposition. (\mathcal{I}, \preceq) is directed.

For the proof, we need some lemmas.

6.2 Lemma. Let $B = (y_1, \ldots, y_k)$ be a tuple of elements in \mathbb{R}^+ . Assume that for some $1 \leq N < k$,

$$\sum_{i=1}^{N} y_i = \sum_{i=N+1}^{k} y_i$$

Then there is some tuple $A = (x_1, \ldots, x_l)$ of elements in $Q(B)^+$ such that l < k and $y_i \in L(A)^+$, $i = 1, \ldots, k$.

Proof. We proceed by induction on k. By the assumptions, we see that k is at least 2, in which case we have $y_1 = y_2$. Put $A := \{y_2\}$ and we are done.

Now let k > 2 and assume that the assertion is true for tuples of length k - 1. By reindexing and rearranging the sums, we may assume that $y_k = \min\{y_1, \ldots, y_k\}$. Put $y'_1 := y_1 - y_k$, then $y'_1 \in Q(B)^+$ and we have the equality

$$y'_1 + y_2 + \dots + y_N = y_{N+1} + \dots + y_{k-1}$$

containing only k - 1 elements. By the induction hypothesis, there is some tuple $A' = (x_1, \ldots, x_{l'})$ with elements in $Q(B)^+$ and l' < k - 1, and some $(k - 1) \times l'$ matrix Z' with values in nonnegative integers such that

$$y'_1 = f_{A'}(z'_{1\cdot}), \quad y_i = f_{A'}(z'_{i\cdot}), \ i = 2, \dots, k-1,$$

here z'_{i} denotes the *i*-th row of Z'. Let $A = (x_1, \ldots, x_{l'}, y_k)$ and

$$Z = \begin{pmatrix} & 1 \\ & 0 \\ Z' & \vdots \\ & 0 \\ 0 & 1 \end{pmatrix}.$$

Then A is an *l*-tuple of elements in $Q(B)^+$, l = l' + 1 < k and $y_i = f_A(z_i) \in L(A)^+$ for all *i*.

6.3 Lemma. Let $B = (y_1, \ldots, y_k)$ be a tuple of elements in \mathbb{R}^+ . Then there is a \mathbb{Q} -linearly independent tuple $A = (x_1, \ldots, x_n)$ of elements in $Q(B)^+$ such that $y_i \in L(A)^+$, $i = 1, \ldots, k$.

Proof. If B is Q-linearly independent, there is nothing to do. Otherwise, there are some $r_i \in \mathbb{Q}$ such that $\sum_i r_i y_i = 0$ with some $r_i \neq 0$. Clearly, by multiplying by a common denominator, we may assume that $r_i \in \mathbb{Z}$. Assume that the elements are arranged in such a way that

$$r_i \begin{cases} > 0 & \text{for } i = 1, \dots, N \\ < 0 & \text{for } i = N + 1, \dots M \\ = 0 & \text{for } i = M + 1, \dots k. \end{cases}$$

Put $p_i = \prod_{i \neq j \leq M} |r_j|$ and let $y'_i = \frac{y_i}{p_i}$ for $i = 1, \ldots, M$. Clearly, $y'_1, \ldots, y'_M \in Q(B)^+$. Then by multiplying the equality by $\prod_{j=1}^M |r_j|^{-1}$, we obtain

$$\sum_{i=1}^{N} y'_{i} = \sum_{i=N+1}^{M} y'_{i}.$$

Applying Lemma 6.2, there is some *l*-tuple $A' = (x'_1, \ldots, x'_l) \in Q(B)^+$ with l < M such that $y'_i \in L(A')^+$ for $i = 1, \ldots, M$, so that also $y_i = p_i y'_i \in L(A')^+$, $i = 1, \ldots, M$.

We now repeat the same process with $B' = (x'_1, \ldots, x'_l, y_{M+1}, \ldots, y_k)$. Since Q(B') = Q(B) and |B'| < k, after a finite number of steps we obtain a \mathbb{Q} -linearly independent set $A = \{x_1, \ldots, x_n\}$ with the required properties.

Proof of Proposition 6.1. Let $B, C \in \mathcal{I}$, then by Lemma 6.3 there is some \mathbb{Q} -linearly independent tuple $A = (x_1 < \cdots < x_n)$ of elements in $Q(B \cup C)^+$ such that $B \cup C \subset L(A)^+$. By assumptions, $1 \in L_>(B)^+ \subset L(A)^+$, so that $1 = \sum_i z_i x_i$ for unique coefficients $z_1, \ldots, z_n \in \mathbb{Z}^+$. Assume that $z_{i_0} = 0$ for some i_0 . Let $B = (y_1 < \cdots < y_k)$. There are some positive integers v_1, \ldots, v_k such that $1 = \sum_{j=1}^k v_j y_j$ and some nonnegative integers w_1^j, \ldots, w_n^j such that $y_j = \sum_i w_i^j x_i$. It follows that

$$1 = \sum_{j=1}^{\kappa} v_j y_j = \sum_i (\sum_j v_j w_i^j) x_i = \sum_i z_i x_i,$$

so that $\sum_{j} v_{j} w_{i}^{j} = z_{i}$, in particular, $\sum_{j} v_{j} w_{i_{0}}^{j} = 0$. Since all v_{j} are positive, this implies that $w_{i_{0}}^{j} = 0$ for all j and we have

$$y_j = \sum_{i \neq i_0} w_i^j x_i.$$

Hence $B \subset L(A \setminus \{x_{i_0}\})^+$, similarly also $C \subset L(A \setminus \{x_{i_0}\})^+$. It follows that we may assume that $1 \in L_>(A)^+$. This means that $1 = \sum_i z_i x_i$ for positive integers z_i , which implies that we must have $0 < x_i \leq 1$. It follows that $A \in \mathcal{I}$ and \mathcal{I} is directed.

We now construct a directed system of simplicial effect algebras. Let $A \in \mathcal{I}$. Since A is \mathbb{Q} -linearly independent, f_A is a **POG** isomorphism onto its range. Let E_A be the interval $[0, f_A^{-1}(1)]$ in $\mathbb{Z}^{|A|}$ and let $g_A = f_A|_{E_A}$. Then g_A is an effect algebra isomorphism onto the interval [0, 1] in $(L(A), L(A)^+, 0)$. Let $B \in \mathcal{I}, B \leq A$, then since $L(B)^+ \subseteq L(A)^+$, we have $g_B(E_B) \subseteq g_A(E_A)$. Put

$$g_{AB}: E_B \to E_A, \qquad g_{AB} = g_A^{-1} g_B,$$

then it is clear that

$$\mathcal{E} = (E_A, A \in \mathcal{I}; g_{AB}, B \preceq A)$$

is a directed system of simplicial effect algebras.

6.4 Proposition. ([0, 1]; $g_A, A \in \mathcal{I}$) is the direct limit of \mathcal{E} .

Proof. It is clear that $([0,1]; g_A, A \in \mathcal{I})$ is compatible with \mathcal{E} . Note also that any $x \in [0,1]$ is contained in the range of some g_A . Indeed, assume that $x \in \mathbb{Q} \cap [0,1]$, then $x = \frac{m}{n}$ for $n \in \mathbb{N}$, $m \in \mathbb{Z}^+$. Let $A = \{\frac{1}{n}\}$, then $A \in \mathcal{I}$ and we have $E_A = [0,n]_{\mathbb{Z}}$, $x = g_A(m)$. If $x \notin \mathbb{Q}$, then $A = \{x, 1-x\} \in \mathcal{I}$ and $x \in A \subset g_A(E_A)$.

Now let E be an effect algebra and let $k_A : E_A \to E$ be a morphisms for $A \in \mathcal{I}$, such that $(E; k_A, A \in \mathcal{I})$ is compatible with \mathcal{E} . Let $x \in [0, 1]$ be in the range of g_A and put

$$\psi(x) = k_A(g_A^{-1}(x)).$$

Assume that $B \in \mathcal{I}$ is such that x is also in the range of g_B and let $C \in \mathcal{I}$ be such that $A, B \leq C$. Then $g_A(E_A) \subseteq g_C(E_C)$ and by compatibility

$$k_A(g_A^{-1}(x)) = k_C g_{CA}(g_A^{-1}(x)) = k_C(g_C^{-1}(x)).$$

Similarly we obtain that $k_B(g_B^{-1}(x)) = k_C(g_C^{-1}(x))$, hence ψ is a well defined map.

Let $I = \{1\}$, then clearly $I \in \mathcal{I}, E_I = \{0, 1\} \subset \mathbb{Z}$ and we have

$$\psi(0) = k_I(0) = 0, \qquad \psi(1) = k_I(1) = 1$$

since k_I is an effect algebra morphism. Further, let $x_1, x_2, x \in [0, 1]$ be such that $x = x_1 + x_2$. Let $A \in \mathcal{I}$ be such that $x_1, x_2 \in g_A(E_A)$, then clearly also $x \in g_A(E_A)$ and we have $g_A^{-1}(x_1) + g_A^{-1}(x_2) = g_A^{-1}(x)$, since g_A is an isomorphism onto its range. Hence

$$\psi(x) = k_A(g_A^{-1}(x)) = k_A(g_A^{-1}(x_1) + g_A^{-1}(x_2)) = \psi(x_1) + \psi(x_2).$$

This proves that ψ is an effect algebra morphism $[0,1] \to E$. Further, for any $A \in \mathcal{I}$ and $z \in E_A$,

$$k_A(z) = k_A(g_A^{-1}g_A(z)) = \psi g_A(z),$$

so that $k_A = \psi g_A$. Since ψ is obviously the unique map $[0, 1] \to E$ with this property, this proves the statement.

References

- M.K. Bennett and D.J. Foulis: Interval and scale effect algebras, Advances in Appl. Math. 41 (1997), 200-215.
- [2] C.C. Chang: Algebraic analysis of many valued logic, Trans. Amer. Math. Soc 88 (1958) 467-490.
- [3] A. Dvurečenskij: Tensor product of difference posets, *Trans. Amer.Math. Soc.* **347** (1995) 1043-1057.
- [4] A. Dvurečenskij, S. Pulmannová: New Trends in Quantum Structures, Kluwer Acad. Publ., Dordrecht/Boston/London and Ister Science, Bratislava, 2000.
- [5] E.G. Effros, D.E. Handelman, C.L. Shen: Dimension groups and their affine representations, Amer. J. Math. 102 (1980), 384-407.

- [6] D.J. Foulis, M.K. Bennett:Effect algebras and unsharp quantum logics, *Found. Phys.* 24 (1994), 1325-1346.
- [7] D.J. Foulis and R. Greechie: Quantum logics and partially ordered abelian groups, *Handbook of Quantum Logic and Quantum Structures*, K. Engesser, D.M. Gabbay, D. Lehmann, Eds., Elsevier, Amsterdam 2007.
- [8] D.J. Foulis, R.J. Greechie, M.K. Bennett: Sums and products of interval effect algebras, Int. J. Theoret Phys. 33 (1994), 2119-2136.
- [9] L. Fuchs: Riesz groups, Ann. Scuola Norm. Sup. Pisa Cl. Sci. 19 (1965), 1-34.
- [10] K.R. Goodearl: Partially Ordered Abelian Groups with Interpolation, Math. Surveys and Monographs No 20, Amer. Math. Soc., Providence, Rhode Island 1986.
- [11] K.R. Goodearl, D.E. Handelman: Tensor products of dimension groups and K₀ of unit regular rings, *Can. J. Math.* **38** (1968), 633-658.
- [12] A. Jenčová, S. Pulmannová: A note on effect algebras and dimension theory of AF C*-algebras, *Rep. Math. Phys* 62 (2008), 205-218.
- [13] F. Kôpka, F. Chovanec: D-posets, Math. Slovaca 44 (1994), 21-34.
- [14] M. Navara: An orthomodular lattice admitting no group=valued measure, Proc. Amer. Math. soc. 122 (1994), 7-12.
- [15] E. Pardo: Metric completions of ordered groups and K_0 of exchange rings, *Trans. Amer. Math. Soc.* **350** (1998), 913-933.
- [16] S. Pulmannová: Effect algebras with the Riesz decomposition property and AF C*-algebras, Foundations of Physics 29 (1999), 1389-1401.
- [17] S. Pulmannová: Tensor product of divisible effect algebras, Bull. Austral. Math. Soc. 68 (2003), 127-140.
- [18] K. Ravindran: On a structure theory of effect algebras, *PhD Dis*sertation, Kansas State University, Manhattan, Kansas 1996.

[19] F. Wehrung: Tensor product of structures with interpolation, Pacific J. Math. 176 (1996), 267-285.