Quantification of Entanglement of Teleportation in Arbitrary Dimensions

Sk Sazim,¹ Satyabrata Adhikari,² Subhashish Banerjee,² and Tanumoy Pramanik³

¹Institute of Physics, Sainik School Post, Bhubaneswar-751005, Orissa, India

²Indian Institute of Technology Rajasthan, Jodhpur-342011, India

³S. N. Bose National Centre for Basic Sciences, Salt Lake, Kolkata-700098, India

We study bipartite entangled states in arbitrary dimensions and obtain different bounds for the entanglement measures in terms of teleportation fidelity. We find that there is a simple relation between negativity and teleportation fidelity for pure states but for mixed states, an upper bound is obtained for negativity in terms of teleportation fidelity using convex-roof extension negativity (CREN). However, with this it is not clear how to distinguish between states useful for teleportation and positive partial transpose (PPT) entangled states. Further, there exists a strong conjecture in the literature that all PPT entangled states, in 3×3 systems, have Schmidt rank two. This motivates us to develop measures capable of identifying states useful for teleportation fidelity and entanglement measures depending upon Schmidt rank of the states. These relations and bounds help us to determine the amount of entanglement required for teleportation, which we call the "Entanglement of Teleportation". These bounds are used to determine the teleportation fidelity as well as two qubit open quantum systems.

PACS numbers: 03.65.Yz, 03.65.Ud, 03.67.Mn

I. INTRODUCTION

Entanglement [1] lies at the heart of quantum mechanics. For a long time it was considered synonymous with quantum correlations but is now regarded as a subset of quantum correlations, see for example [2]. Entanglement plays a pivotal role in various information processing tasks, including, among others, quantum teleportation [3], super dense coding [4], remote state preparation [5], secret sharing [6], and quantum cryptography [7].

In quantum teleportation, using entangled states as resource, it is possible to transfer quantum information from an unknown qubit to another one placed at a distance. Thus, one of the party, say, Alice makes a two qubit measurement on her qubit and the unknown state in Bell basis, and sends the measurement results through a classical channel to the second party, say, Bob (who is located away from Alice). Accordingly, Bob makes appropriate unitary transformations to obtain the desired state. Thus the ability of teleporting an unknown state depends on the nature of entanglement of the resource state and is called teleportation fidelity.

The situation is very straight forward when we have an unknown qubit to send with the help of a pure entangled state as a resource. However, it is more involved when we have mixed entangled states as a medium of teleportation. For a general two qubit density matrix $\rho = \frac{1}{4}[I \otimes I + \sum_i r_i(\sigma_i \otimes I) + \sum_j s_j(I \otimes \sigma_j) + \sum_{i,j} t_{ij}(\sigma_i \otimes \sigma_j)]$, the teleportation fidelity is a function of the eigenvalues of correlation matrix $T = [t_{ij}]$. Similarly, when we go from qubits to higher dimensional bipartite states the teleportation fidelity is expressed in terms of the singlet fraction of the state. The relation between optimal teleportation fidelity $F(\rho)$ and maximal singlet fraction $f(\rho)$ in a $d \otimes d$ system, if one performs quantum teleportation with the state ρ , is [8]

$$F(\rho) = \frac{df(\rho) + 1}{d+1}.$$
 (1)

Here the singlet fraction is defined as, $f(\rho) = \max_{|\psi\rangle} \langle \psi | \rho | \psi \rangle$, and $|\psi\rangle$ is a maximally entangled state in $d \otimes d$. If $f(\rho) > \frac{1}{d}$ then the parties can perform quantum teleportation with the average fidelity of the teleported qubit exceeding the classical limit $\frac{2}{d+1}$.

In bipartite two qubit states it is known that the total amount of entanglement present in the resource state is useful for teleportation. Then one can ask the following question: How much entanglement is necessary for teleporting an unknown state when we have a bipartite state in arbitrary dimensions? To answer this question one has to quantify the entanglement and find out for what range of entanglement the state can be used as a resource for teleportation. In other words, one needs to establish a relationship between the amount of entanglement and teleportation fidelity. We establish various relations between teleportation fidelity and entanglement measures depending upon Schmidt rank [9, 10] of the states. These relations and bounds help us to answer the above question. Given an arbitrary two-qudit state with Schmidt rank upto three we can predict its utility as a resource for teleportation.

Negativity [11] is a measure of entanglement of a bipartite quantum state described by the density operator ρ and is formally defined in $d \times d$ systems as

$$N(\rho) = \frac{||\rho^{T_A}|| - 1}{d - 1},$$
(2)

where ρ^{T_A} is a partial transpose of ρ with respect to the system A and ||.|| denotes the trace norm. Negativity fails to distinguish separable states from PPT entangled states, that is, bound entangled states. This difficulty can be overcome by the use of convex-roof extension of negativity (CREN) [12].

There is strong evidence in 3×3 systems that bound entangled states exist only for states with Schmidt rank two [13]. Thus the entangled states with Schmidt rank three would, presumably, be useful for teleportation. This provides a strong motivation to study the states with Schmidt rank three from the perspective of teleportation fidelity. The Schmidt number is a very useful entanglement measure [14]. In the literature, there exists different kinds of entanglement measures, expressed in terms of Schmidt numbers, suitable for quantification of the amount of entanglement present in the system.

We quantify the amount of entanglement present in the resource state to find out the bounds within which these states can be useful for teleportation. Thus, we obtain relations connecting entanglement measures with teleportation fidelity using CREN as well as singlet fraction, expressed in terms of Schmidt coefficients. Our results are obtained for arbitrary dimensional bipartite states with at most three non vanishing Schmidt coefficients. We implement our results to detect mixed states useful for teleportation. A monotonous relation between entanglement and teleportation fidelity in mixed two qudit systems could be expected from [15], where a monotonous connection between entanglement and singlet fraction, and hence teleportation fidelity, was established for two qubit mixed states.

The plan of the paper is as follows. In Section 2, we study the relation between negativity and teleportation fidelity for pure as well as mixed systems. Based on our conclusions from Section 2, we establish a relation between singlet fraction and different types of entanglement measures for arbitrary dimensional pure two qudit system with a maximum of three Schmidt coefficients in Section 3. Then we study the bounds of teleportation fidelity and entanglement measures for two special cases, i) arbitrary dimensional pure bipartite state with two Schmidt coefficients, and ii) arbitrary dimensional pure bipartite state with three Schmidt coefficients. These results are used in section 3 (B) to arbitrary dimensional mixed bipartite systems with Schmidt coefficients two and three. In section 4, we apply our results on examples of mixed states, in particular, two qutrit mixed state with Schmidt rank two, and two qubit mixed states generated dynamically by an open system model. Finally, we conclude in section 5.

II. RELATION BETWEEN NEGATIVITY AND TELEPORTATION FIDELITY FOR $d \otimes d$ SYSTEMS

Here we study the relation between negativity and teleportation fidelity for pure as well as mixed systems.

A. Pure Systems

Let H_A and H_B be two Hilbert spaces each with dimension d. In $d \otimes d$ system, any pure state $|\Psi\rangle$ can be expressed as

$$|\Psi\rangle = \sum_{i=1}^{d} \sqrt{\lambda_j} |j\rangle |j\rangle.$$
(3)

The negativity of the state $|\Psi
angle$ is defined as

$$N(|\Psi\rangle) = \frac{2}{d-1} \sum_{i < j} \sqrt{\lambda_i \lambda_j}.$$
 (4)

The singlet fraction for any pure state in $d \otimes d$ system is given by

$$f(|\Psi\rangle) = \frac{1}{d} \left(\sum_{i=1}^{d} \sqrt{\lambda_i}\right)^2.$$
 (5)

The relation between negativity and singlet fraction is given by [16]

$$N(|\Psi\rangle) = \frac{df(|\Psi\rangle) - 1}{d - 1}.$$
(6)

In terms of teleportation fidelity, Eq. (6) reduces to

$$F(|\Psi\rangle) = \frac{2}{d+1} + \frac{(d-1)N(|\Psi\rangle)}{d+1}.$$
 (7)

Therefore, it follows that every entangled pure state in a $d \otimes d$ system is useful for teleportation.

B. Mixed Systems

A bipartite mixed state described can be described by the density operator ρ

$$\rho = \sum_{i} p_{i} |\Psi_{i}\rangle \langle \Psi_{i}|. \tag{8}$$

The negativity of the mixed state ρ can be extended from the pure state by means of convex roof, that is, convex-roof extended negativity (CREN) [12]:

$$N(\rho) = \min_{\{p_i, |\Psi_i\rangle\}} \sum_i p_i N(|\Psi_i\rangle).$$
(9)

The upper bound of the negativity of the mixed state ρ can be expressed in terms of the singlet fraction as

$$N(\rho) = \min_{\{p_i, |\Psi_i\rangle\}} \sum_{i} p_i N(|\Psi_i\rangle)$$

$$\leq \sum_{i} p_i N(|\Psi_i\rangle)$$

$$= \frac{d}{d-1} \sum_{i} p_i f(|\Psi_i\rangle) - \frac{1}{d-1}.$$
(10)

In terms of teleportation fidelity, the bound on negativity is

$$N(\rho) \le \frac{d+1}{d-1} \sum_{i} p_i F(|\Psi_i\rangle) - \frac{2}{d-1}.$$
 (11)

The above inequality (11) measures the upper bound of entanglement contained in the mixed state ρ . From this, it is clear that CREN detects both PPT bound entangled states as well as states useful for teleportation. However, it is not clear how to distinguish between these two classes of states. Further, there exists a strong conjecture in the literature [13] that all PPT entangled states, in 3×3 systems, have Schmidt rank two. This motivates us to develop measures capable of identifying states useful for teleportation and dependent on the Schmidt number.

III. RELATION BETWEEN SINGLET FRACTION AND DIFFERENT ENTANGLEMENT MEASURES FOR TWO QUDIT SYSTEM WITH THREE SCHMIDT COEFFICIENTS

In this section we obtain an explicit relation that will connect entanglement monotones with singlet fraction for a two qudit system of arbitrary dimension. We obtain results in $d \otimes d$ systems with two and three non zero Schmidt coefficients.

A. Bounds on entanglement measures for pure two qudit systems useful for teleportation

Let us consider a bipartite $d \otimes d$ system in which three Schmidt coefficients are non zero. Without any loss of generality we assume that the first three Schmidt coefficients are non zero. Any pure two qudit system with three non zero Schmidt coefficients λ_1 , λ_2 and λ_3 can be written in Schmidt decomposition form as,

$$|\Psi^d\rangle = \sqrt{\lambda_1}|00\rangle + \sqrt{\lambda_2}|11\rangle + \sqrt{\lambda_3}|22\rangle, \qquad (12)$$

with the Schmidt coefficients summing to one, i.e., $\lambda_1 + \lambda_2 + \lambda_3 = 1$. To quantify the amount of entanglement in $|\Psi^d\rangle$ we consider two different entanglement measures $E^{(d,2)}(|\Psi^d\rangle)$ and $E^{(d,3)}(|\Psi^d\rangle)$ which can be defined as [17],

$$E^{(d,2)}(|\Psi^d\rangle) = \sqrt{\frac{2d}{d-1}(\lambda_1\lambda_2 + \lambda_2\lambda_3 + \lambda_1\lambda_3)},$$
 (13)

$$E^{(d,3)}(|\Psi^d\rangle) = \left(\frac{6d^2}{(d-1)(d-2)}\right)^{\frac{1}{3}} (\lambda_1 \lambda_2 \lambda_3)^{\frac{1}{3}}.$$
 (14)

Here $E^{(d,2)}(|\Psi^d\rangle)$ and $E^{(d,3)}(|\Psi^d\rangle)$ denote entanglement measure for a $d \otimes d$ dimensional pure system defined by taking the sum of the product of the Schmidt coefficients taken two or three at a time, respectively. We note that for a Schmidt rank two state, $E^{(d,3)}(|\Psi^d\rangle) = 0$ but $E^{(d,2)}(|\Psi^d\rangle) \neq 0$.

The singlet fraction for the state $|\Psi^d
angle$ is defined as

$$f(|\Psi^d\rangle) = \max_{|\Phi\rangle} |\langle \Phi | \Psi^d \rangle|^2, \tag{15}$$

where the maximum is taken over all maximally entangled states $|\Phi\rangle$ in $d \otimes d$ systems. The singlet fraction $f(|\Psi^d\rangle)$ for pure state $|\Psi^d\rangle$ can also be expressed in terms of Schmidt coefficients [18] as

$$f(|\Psi^d\rangle) = \frac{1}{d} \left(\sqrt{\lambda_1} + \sqrt{\lambda_2} + \sqrt{\lambda_3}\right)^2.$$
(16)

Expanding the the right hand side part of Eq. (16) and using $\lambda_1 + \lambda_2 + \lambda_3 = 1$, we get

$$\sqrt{\lambda_1 \lambda_2} + \sqrt{\lambda_2 \lambda_3} + \sqrt{\lambda_1 \lambda_3} = \frac{df(|\Psi^d\rangle) - 1}{2}.$$
 (17)

Also, we have the following identity

$$\lambda_1 \lambda_2 + \lambda_2 \lambda_3 + \lambda_1 \lambda_3 = (\sqrt{\lambda_1 \lambda_2} + \sqrt{\lambda_2 \lambda_3} + \sqrt{\lambda_1 \lambda_3})^2 -2\sqrt{\lambda_1 \lambda_2 \lambda_3} (\sqrt{\lambda_1} + \sqrt{\lambda_2} + \sqrt{\lambda_3}) (18)$$

Using (13), (14), (16), (17) and (18) we have

$$(E^{(d,2)}(|\Psi^{d}\rangle))^{2} = \frac{d^{3}}{2(d-1)} \left(f(|\Psi^{d}\rangle) - \frac{1}{d}\right)^{2} - \frac{4}{d-1} \sqrt{\frac{d(d-1)(d-2)}{6}} (E^{(d,3)}(|\Psi^{d}\rangle))^{\frac{3}{2}} \times \sqrt{f(|\Psi^{d}\rangle)}.$$
(19)

This establishes the required relationship between the entanglement measures $E^{(d,2)}(|\Psi^d\rangle)$ and $E^{(d,3)}(|\psi^d\rangle)$ with the singlet fraction $f(|\Psi^d\rangle)$ for a pure two qudit system $|\Psi^d\rangle$ with three non vanishing Schmidt coefficients.

Next, we will consider separately the cases of states of Schmidt ranks two and three, respectively. For purpose of clarity, in the discussions to follow, we modify the notation of the entanglement measures discussed above, as $E_j^{d,i}$, where d stands for the $d \otimes d$ dimensional system, j indicates the Schmidt rank of the state under consideration and i is the number of coefficients taken at a time.

1. States with Schmidt Rank Two

When one of the Schmidt coefficients (say, λ_3) is zero, i.e., $E_2^{(d,3)}(|\Psi^d\rangle) = 0$, from Eq. (19), we have

$$E_2^{(d,2)}(|\Psi^d\rangle) = \sqrt{\frac{d^3}{2(d-1)}} \left(f_2(|\Psi^d\rangle) - \frac{1}{d} \right),$$
(20)

where, $f_2(|\Psi^d\rangle)$ denotes the singlet fraction of Schmidt rank two state, and $f_2(|\Psi^d\rangle) > \frac{1}{d}$. If $F_2(|\Psi^d\rangle)$ denotes the teleportation fidelity of Schmidt rank two states, then $E_2^{(d,2)}(|\Psi^d\rangle)$ can be expressed in terms of $F_2(|\Psi^d\rangle)$ as

$$E_2^{(d,2)}(|\Psi^d\rangle) = \sqrt{\frac{d^3}{2(d-1)}} \left[\frac{(d+1)F_2(|\Psi^d\rangle) - 2}{d}\right].$$
 (21)

This establishes the relation between the entanglement monotone and teleportation fidelity of Schmidt rank two states. If the state $|\Psi^d\rangle$ has Schmidt number two and useful for teleportation, then we have [20]

$$\frac{1}{d} < f_2(|\Psi^d\rangle) \le \frac{2}{d}.$$
(22)

Eq. (22) can be recast in terms of teleportation fidelity as

$$\frac{2}{d+1} < F_2(|\Psi^d\rangle) \le \frac{3}{d+1}.$$
(23)

Using Eq. (23), $E_2^{(d,2)}(|\Psi^d\rangle)$ can be seen to be bounded as

$$0 < E_2^{(d,2)}(|\Psi^d\rangle) \le \sqrt{\frac{d}{2(d-1)}}.$$
(24)

When the amount of entanglement lies in the above range we can use the state for teleportation. This quantifies the entanglement required for teleportation for a pure qudit state with two non-vanishing Schmidt coefficients.

2. States with Schmidt Rank Three

Next we take up sates where none of the three Schmidt coefficients are zero, i.e., $E_3^{(d,3)}(|\Psi^d\rangle) \neq 0$.

Using the well known result of arithmetic mean (AM) being greater than or equal to geometric mean (GM) on three real quantities $\sqrt{\lambda_1 \lambda_2}$, $\sqrt{\lambda_1 \lambda_3}$ and $\sqrt{\lambda_2 \lambda_3}$, we have

$$\frac{\sqrt{\lambda_1\lambda_2} + \sqrt{\lambda_1\lambda_3} + \sqrt{\lambda_1\lambda_3}}{3} \ge \left(\sqrt{\lambda_1\lambda_2}\sqrt{\lambda_1\lambda_3}\sqrt{\lambda_2\lambda_3}\right)^{\frac{1}{3}}.$$
(25)

Using Eqs. (14), and (17), we have

$$f_3(|\Psi^d\rangle) \ge \frac{6}{d} \left[\left(\frac{(d-1)(d-2)}{6d^2} \right)^{\frac{1}{3}} E_3^{(d,3)}(|\Psi^d\rangle) \right] + \frac{1}{d}.(26)$$

Since, the singlet fraction $f_3(|\Psi^d\rangle)$ attains its maximum value unity at $\lambda_1 = \lambda_2 = \lambda_3 = \frac{1}{d}$, we have

$$\frac{6}{d} \left[\left(\frac{(d-1)(d-2)}{6d^2} \right)^{\frac{1}{3}} E_3^{(d,3)}(|\Psi^d\rangle) \right] + \frac{1}{d} \le f_3(|\Psi^d\rangle) \le 1.$$
(27)

In terms of teleportation fidelity $F_3(|\Psi^d\rangle)$, the above inequality can be expressed as

$$\frac{2}{d+1} + \frac{6}{d+1} \left(\frac{(d-1)(d-2)}{6d^2} \right)^{\frac{1}{3}} E_3^{(d,3)}(|\Psi^d\rangle) \\ \leq F_3(|\Psi^d\rangle) \leq 1.$$
(28)

Hence, pure entangled states with $E_3^{(d,3)}(|\Psi^d\rangle)$ satisfying Eq. (28) and teleportation fidelity $F_3(|\Psi^d\rangle) > \frac{2}{d+1}$ are Schmidt rank three states useful for teleportation.

B. Bounds on entanglement measures for mixed two qudit systems useful for teleportation

In this section we would like to answer the following questions : (i) What is the minimum amount of entanglement needed to perform teleportation when the mixed state eith Schmidt rank two is used as a resource in a $d \otimes d$ system? (ii) What is the minimum amount of entanglement needed to perform teleportation when the mixed state with Schmidt rank three is used as resource in a $d \otimes d$ system?

Let us consider a mixed qudit state described by the density operator $\rho = \sum_{i=1}^{n} p_i \rho_i$, where $\sum_{i=1}^{n} p_i = 1$ and ρ_i

 $(= |\psi_i\rangle\langle\psi_i|)$ are composite pure states. The singlet fraction $f(\rho)$ of the state ρ is defined as

$$f(\rho) = \max_{U} \langle \psi^{+} | U^{\dagger} \otimes \mathcal{I} \rho U \otimes \mathcal{I} | \psi^{+} \rangle,$$
(29)

where U is the unitary matrix, \mathcal{I} is the identity matrix and $|\psi^+\rangle = \frac{1}{\sqrt{d}} \sum_{k=0}^{d-1} |kk\rangle$ represents a pure maximally entangled state.

The entanglement measure $E^{(d,2)}(|\Psi^d\rangle)$ and $E^{(d,3)}(|\Psi^d\rangle)$ given in Eqs. (13) and (14) for pure states can also be defined for a mixed state ρ as

$$E^{(d,2)}(\rho) = \min \sum_{i=1}^{n} p_i E^{(d,2)}(\rho_i),$$
(30)

and

Ì

$$E^{(d,3)}(\rho) = \min \sum_{i=1}^{n} p_i E^{(d,3)}(\rho_i).$$
(31)

Here the minimum is taken over all pure state decompositions of ρ . Now one may ask a question that, like entanglement measures, does the singlet fraction $f(\rho)$ also have the property [19]

$$f(\rho) = \min \sum p_i f(\rho_i), \qquad (32)$$

where the minimum is taken over all decomposition of ρ . Unfortunately, the answer is no.

1. Two qudit mixed state of Schmidt rank two

From Eq. (19), $E_2^{(d,2)}(\rho_i)$ for any bipartite pure qudit state with Schmidt rank two ρ_i whose $f_2(\rho_i) = \frac{1}{d}$, i.e., for states not useful for teleportation, we have

$$E_2^{(d,2)}(\rho_i) = 0. \tag{33}$$

In general for any bipartite pure qudit state with Schmidt rank two ρ_i useful for teleportation, the entanglement $E_2^{(d,2)}$ is

$$E_2^{(d,2)}(\rho_i) = \sqrt{\frac{d^3}{2(d-1)}} \left(f_2(\rho_i) - \frac{1}{d} \right).$$
(34)

Using Eqs. (30) and (34), we have

$$E_{2}^{(d,2)}(\rho) = \min \sum_{i} p_{i} \sqrt{\frac{d^{3}}{2(d-1)}} \left(f_{2}(\rho_{i}) - \frac{1}{d} \right)$$

$$\leq \sum_{i} p_{i} \sqrt{\frac{d^{3}}{2(d-1)}} \left(f_{2}(\rho_{i}) - \frac{1}{d} \right)$$

$$< \sqrt{\frac{d}{2(d-1)}}, \qquad (35)$$

where the last inequality follows from an application of Eq. (22). Hence, if the mixed state ρ with Schmidt rank two in a $d \otimes d$ system is useful for teleportation then

$$0 < E_2^{(d,2)}(\rho) < \sqrt{\frac{d}{2(d-1)}}.$$
(36)

2. Two qudit mixed state of Schmidt rank three

Using, once again, the result of arithmetic mean (AM) being greater than or equal to geometric mean (GM) on three real quantities $\lambda_1 \lambda_2$, $\lambda_1 \lambda_3$ and $\lambda_2 \lambda_3$ and Eqs. (13), (14) we obtain the following bound on $E_3^{(d,3)}(\rho)$ for two qudit mixed states with Schmidt rank three:

$$0 < E_3^{(d,3)}(\rho) < \left[\frac{d(d-1)}{6}\right]^{\frac{1}{6}} \frac{1}{(d-2)^{1/3}}.$$
 (37)

Comparing Eqs. (37) and (36), we can see that if the entanglement lies in the range $\sqrt{\frac{d}{2(d-1)}}$ to $\left[\frac{d(d-1)}{6}\right]^{\frac{1}{6}} \frac{1}{(d-2)^{1/3}}$ it can be concluded that the state is of Schmidt rank three.

IV. ILLUSTRATIONS AND APPLICATIONS

In this section we provide examples of qubit and qutrit mixed states as applications of our results. This paves the way for detecting states which are useful for teleportation as well as to quantify the amount of entanglement required for teleportation, in realistic settings.

A. Two qutrit mixed states with Schmidt rank two

We consider a two qutrit mixed state with Schmidt rank two [10] given by

$$\rho_f = \frac{5p}{p+2}\rho_c + \frac{2(1-2p)}{p+2}|\phi\rangle\langle\phi|; 0 \le p \le \frac{1}{2}, \qquad (38)$$

where, $\rho_c = \frac{1}{2}(|\chi_0\rangle\langle\chi_0| + |\chi_1\rangle\langle\chi_1|)$. This decomposition for state ρ_f is optimal. Here, $|\chi_0\rangle$ and $|\chi_1\rangle$ are of the form $|\chi_0\rangle = \sqrt{\frac{3}{5}}|\psi\rangle + \sqrt{\frac{2}{5}}|\phi\rangle$ and $|\chi_1\rangle = \sqrt{\frac{3}{5}}|\psi\rangle - \sqrt{\frac{2}{5}}|\phi\rangle$, respectively, and the states $|\psi\rangle, |\phi\rangle$ are given by, $|\psi\rangle = \frac{1}{\sqrt{3}}(|00\rangle + |11\rangle - e^{\frac{i\pi}{3}}|22\rangle)$ and $|\phi\rangle = \frac{1}{\sqrt{2}}(|00\rangle + |11\rangle)$. Also, p is the classical probability of mixing.

We check whether the bounds on $E_2^{(3,2)}(\rho_i)$ works for the above density matrix. For $3 \otimes 3$ dimension, $E_2^{(3,2)}(\rho_f)$ (see Eq. (35)) becomes

$$E_{2}^{(3,2)}(\rho_{f}) = \frac{3\sqrt{3}}{2} \left(\min \sum_{i} p_{i} f_{2}(\rho_{i}) \right) - \frac{\sqrt{3}}{2}$$
$$= \frac{3\sqrt{3}}{2} \left(\min_{\{p\}} \left[\frac{1+p}{2+p} \right] \right) - \frac{\sqrt{3}}{2}$$
$$= \frac{\sqrt{3}}{4}; \quad \text{for } p = 0.$$
(39)

In this calculation we have used the appropriate maximally entangled basis given in [21]. From Eqs. (39) and (24), it can be seen that the state (in Eq. (38)) is useful for teleportation.

B. States generated as a result of Two-Qubit Interaction with a Squeezed Thermal Bath

Open quantum systems is the systematic study of the evolution of the system of interest, such as a qubit, under the influence of its environment, also called the bath or the reservoir. This results in decoherence and dissipation. Consider the Hamiltonian $H = H_S + H_R + H_{SR}$; where S stands for the system of interest, R for reservoir and SR for the system-reservoir interaction. Depending upon the system-reservoir interaction, open systems can be classified into two broad categories, viz., dissipative or QND (quantum non-demolition). In case of QND dephasing occurs without damping the system, i.e., where $[H_S, H_{SR}] = 0$ while decoherence along with dissipation occurs in dissipative systems, i.e., $[H_S, H_{SR}] \neq 0$. [22–24].

1. States generated as a result of Two-Qubit Open System Interacting with a Squeezed Thermal Bath via a Dissipative Interaction

Here we study the dynamics of the bound [Eq. (36)] for a two-qubit open system interacting with a squeezed thermal bath, modeled as a 3-D electromagnetic field (EMF), as well as its specialization to a vacuum bath, where the bath squeezing (r) and temperature (T) are set to zero, and undergoing a dissipative interaction [25]. The model Hamiltonian is

$$H = H_{S} + H_{R} + H_{SR}$$

= $\sum_{n=1}^{2} \hbar \omega_{n} S_{n}^{z} + \sum_{\vec{k}_{s}} \hbar \omega_{k} \left(b_{\vec{k}_{s}}^{\dagger} b_{\vec{k}_{s}} + \frac{1}{2} \right)$
- $i\hbar \sum_{\vec{k}_{s}} \sum_{n=1}^{2} [\vec{\mu}_{n} \cdot \vec{g}_{\vec{k}_{s}}(\vec{r}_{n})(S_{n}^{+} + S_{n}^{-})b_{\vec{k}_{s}} - h.c.].(40)$

Here $\vec{\mu}_n$ are the transition dipole moments, dependent on the different atomic positions \vec{r}_n and S_n^+ (= $\frac{1}{2}|e_n\rangle\langle g_n|$), and $S_n^-(= \frac{1}{2}|g_n\rangle\langle e_n|$) are the dipole raising and lowering operators satisfying the usual commutation relations. $S_n^z(= \frac{1}{2}(|e_n\rangle\langle e_n| - |g_n\rangle\langle g_n|))$ is the energy operator of *n*th atom and $b_{\vec{k}_s}^{\dagger}$, $b_{\vec{k}_s}$ are the creation and anihilation operators of the field mode \vec{k}_s with the wave vector \vec{k} and polarization index s = 1, 2. A key feature of the model is that the system-reservoir (S-R) coupling constant $\vec{g}_{\vec{k}_s}(\vec{r}_n)$ is dependent on the position of the qubit r_n and is

$$\vec{g}_{\vec{k}_s}(\vec{r}_n) = \left(\frac{\omega_k}{2\epsilon\hbar V}\right)^{\frac{1}{2}} \vec{e}_{\vec{k}_s} e^{i\vec{k}\cdot r_n},\tag{41}$$

where V is the normalization volume and $\vec{e}_{\vec{k}s}$ is the unit polarization vector of the field. The position dependence of the coupling leads to interesting dynamical consequences and allows the entire dynamics to be classified into two categories, that is, the independent regime, where the interqubit distance is far enough for each qubit to locally interact with an independent bath or the collective regime, where the qubits are close




FIG. 1: Plot of (i) concurrence C (or $E_2^{(2,2)}$) and (ii) singlet fraction f with respect to the time of evolution t, respectively. Here we consider the case of a vacuum bath (T = r = 0) and the collective decoherence model ($r_{12} = 0.05$).

enough for them to interact with the bath collectively. Asuming an initial system-reservoir separable state, with the system in a separable, and the bath in a squeezed thermal state, with time the qubits develop correlations between themselves via a channel setup by the bath. A master equation for the reduced dynamics of the two qubit system is obtained by tracing out the environment (bath), using the usual Born-Markov and rotating wave approximation (RWA). This can be then solved to obtain the dynamics of the reduced density matrix, whose details are presented in [25], for the general case of a squeezed thermal bath at finite temperature as well as for a vacuum reservoir.

Let the reduced two-qubit density matrix of the system be $\rho_f(t)$. Its spectral decomposition corresponding to its eigenvalues $(\lambda_i(t))$ is,

$$\rho_f(t) = \sum_i \lambda_i(t)\rho_i(t). \tag{42}$$

Here $\rho_i(t) = |\psi_i(t)\rangle \langle \psi_i(t)|, |\psi_i(t)\rangle$ being the eigenvectors corresponding to the eigenvalues $\lambda_i(t)$ ($\sum_i \lambda_i(t) = 1$). For a two qubit state the Eq. (36) becomes

$$0 \le E_2^{(2,2)}(\rho_f(t)) \le 1.$$
(43)

We can easily say that for two-qubit state $E_2^{(2,2)}$ is nothing

2. States generated as a result of Two-Qubit Open System Interacting with a Squeezed Thermal Bath via Quantum Nondemolition Interaction

Now we take up the Hamiltonian, describing a QND interaction of two qubits with the bath as

$$H = H_S + H_R + H_{SR}$$

= $\sum_{n=1}^{2} \hbar \varepsilon_n J_z^n + \sum_k \hbar \omega_k b_k^{\dagger} b_k$
+ $\sum_{n,k} \hbar J_z^n (g_k^n b_k^{\dagger} + g_k^{n*} b_k).$ (44)

but concurrence C. If we look at the Figs. (1), and (2), for the case of a vacuum bath (T = 0, r = 0), concurrence C (or $E_2^{2,2}$) is seen to decrease with time of evolution t, with a predominantly oscillatory behavior in the collective regime (marked by the inter qubit distance $r_{12} < 1$). The singlet fraction f also shows similar behavior. From these two figures, it is clear that when and where C becomes zero, and f is equal to $\frac{1}{2}$.

For the case of a squeezed thermal bath, as the system evolves with time t, concurrence C and f exhibit damped behavior, as seen in Figs. (3) and (4). If we increase the inter-qubit distance r_{12} , then the concurrence C for the system suddenly falls to zero (i.e., sudden death of entanglement in the system). Thus, the system can be used as a resource for teleportation purpose in the range $0 \leq r_{12} < r_d$. Here we define a new term r_d , such that at $r_{12} = r_d$ concurrence C of the system becomes zero. Obviously this r_d will be different for different parameter (T, r) settings. The Figs. (4) depict the abrupt decrease of concurrence C and singlet fraction f as r_{12} increases. In Figs. (5), the behavior of C and f with respect to environmental squeezing parameter r is shown. For rbetween -0.02 to 0.02 both C and f remain almost constant, thereby exhibiting the tendency of squeezing to resist environmental degradation. Beyond this range there is a rapid fall of the depicted quantities.

Here H_S , H_R and H_{SR} stand for the Hamiltonians of the system, reservoir and system-reservoir interaction, respectively. b_k^{\dagger} , b_k denote the creation and annihilation operators for the reservoir oscillator of frequency ω_k , g_k^n stands for the coupling constant (again assumed to be position dependent) for the interaction of the oscillator field with the qubit system and are taken to be $g_k^n = g_k e^{-ik.r_n}$, where r_n is the qubit position. Since $[H_S, H_{SR}] = 0$, the Hamiltonian [Eq. 44)] is of QND type. In the parlance of quantum information theory,




FIG. 2: Plot of (i) concurrence C (or $E_2^{(2,2)}$) and (ii) f with respect to the inter-qubit distance r_{12} , respectively. Here we consider the case of vacuum bath (T = r = 0) and system is at time t = 10.




FIG. 3: Plot of (i) concurrence C (or $E_2^{(2,2)}$) and (ii) singlet fraction f with respect to the time of evolution t, respectively, for a squeezed thermal bath (T = 1, r = 0.1) in the collective regime ($r_{12} = 0.05$).

the noise generated is called the phase damping noise. The position dependence of the coupling constant once more allows for the dynamical classification into the independent and collective regimes. In order to obtain the reduced dynamics of the system , we trace over the reservoir variables, the details of which can be found in [26].

Now we study the behavior of concurrence C (actually $E_2^{2,2}$) and singlet fraction f as the two-qubit system evolves with time t both for collective and localized (independent) decoherence model. It can be noticed from Figs. (6), and (7) that the value of concurrence C is higher and lasts longer in the case of collective decoherence model than in the case of localized decoherence model. As expected, the the singlet fraction f shows similar kind of behavior with time t. When C becomes zero, f becomes equal to $\frac{1}{2}$, i.e., the system at this particular time t cannot be useful for teleportation, otherwise it is useful. Hence the system satisfies the lower bound of Eq. (43), when concurrence C vanishes. The behavior of C and f, under pure dephasing, with respect to environmental squeezing parameter r is depicted in Figs. (8). For r between -0.02 to 0.02 both C and f remain almost constant, thereby exhibiting the tendency of squeezing to resist environmental degradation. Beyond this range there is a fall of the depicted quantities, though the degradation here is smoother than that in Figs. (5).

V. CONCLUSION

We have made a study of entanglement of teleportation for arbitrary $d \otimes d$ dimensional states having Schmidt rank upto three. We found that there is a simple relation between negativity and teleportation fidelity for pure states but for mixed states, an upper bound was obtained for negativity in terms of teleportation fidelity using convex-roof extension negativity (CREN). The existence of a strong conjecture in the literature concerning all PPT entangled states, in 3×3 systems, having Schmidt rank two, motivated us to develop measures capable of identifying states useful for teleportation and dependent on the Schmidt number. This enabled a classification of entanglement as a function of teleportation fidelity, the "Entanglement of Teleportation". These results were then extended to mixed two qudit states, which we illustrated on specific examples of a two qutrit mixed state with Schmidt rank two, and two qubit states dynamically generated by interaction with an appropriate reservoir, for both pure dephasing as well as dissipative




FIG. 4: Plot of (i) concurrence C (or $E_2^{(2,2)}$) and (ii) f with respect to the inter-qubit distance r_{12} , respectively, for a squeezed thermal bath (T = 1, r = 0.1) and time of evolution t = 1.




FIG. 5: Plot of (i) concurrence C (or $E_2^{(2,2)}$) and (ii) f with respect to the squeezing parameter r, respectively, for a thermal bath ($T = 5, r_{12} = 0.05$) and time of evolution t = 2.

interactions. This work thus brings into focus the utility of studying higher dimensional entangled states using measures like "Entanglement of Teleportation" along with negativity. Acknowledgment: T. Pramanik thanks UGC, India for financial support. We thank Prof. A. Mazumdar and Prof. H. S. Sim for useful discussions.

- [1] A. Einstein, B. Podoisky and N. Rosen, Phys. Rev. 47, 777 (1935).
- [2] H. Ollivier and W. H. Zurek, Phys. Rev. Lett. 88, 017901 (2001); L. Henderson and V. Vedral, J. Phys. A 34, 6899 (2001).
- [3] C. H. Bennett, G. Brassard, C. Creapeau, R. Jozsa, A. Pares and W. K. Wooters, Phys. Rev. Lett. 70, 1895 (1993).
- [4] C. H. Bennett and S. Wiesner, Phys. Rev. Lett. 69, 433 (1992).
- [5] A. K. Pati, Phys. Rev. A. 63, 014320-1 (2001); C. H. Bennett,
 D. P. DiVincenzo, P. W. Shor, J. A. Smolin, B. M. Terhal and
 W. K. Wooters, Phys. Rev. Lett. 87, 077902 (2001).
- [6] M. Hillery, V. Buzek and A. Berthiaume, Phys. Rev. A. 59, 1829 (1999).
- [7] N. Gisin, G. Ribordy, W. Tittel and H. Zbinden, Rev. Mod. Phys. 74, 145 (2002).
- [8] M. Horodecki, P. Horodecki, and R. Horodecki, Phys. Rev. A. 60, 1888 (1999).
- [9] E. Schmidt, Math. Ann. 63, 433 (1907).
- [10] Schmidt decomposition [9] is a very good tool to describe com-

posite systems. If $|\Psi\rangle$ is a pure state of composite systems A and B then, $|\Psi\rangle = \sum_{i=1}^{d_A} \sqrt{\lambda_i} |i_A\rangle |i_B\rangle$, represents the Schmidt decomposition of $|\Psi\rangle$, where $|i_A\rangle (i_A = 1, 2, ..., d_A) \in \mathcal{H}_A$ and $|i_B\rangle (i_B = 1, 2, ..., d_B) \in \mathcal{H}_B$ are orthonormal bases for A and B respectively, and $d_A \leq d_B$. Here λ_i are the Schmidt coefficients, non-negative real numbers satisfying the relation $\sum_i \lambda_i = 1$. The number of non-zero Schmidt coefficients of a state is called its Schmidt rank. For bipartite mixed state $(\rho = \sum_i p_i |\Psi_i\rangle \langle \Psi_i|)$ in finite dimension, the Schmidt rank is defined as [20] $SR(\rho) = \inf_{\{\sum_i p_i, |\Psi_i\rangle\}} \sup_i SR(|\Psi_i\rangle)$ where, the infimum is taken over all the ensembles $\{p_i, |\Psi_i\rangle\}$ of pure states with the average state ρ .

- [11] A. Peres, Phys. Rev. Lett. **76**, 1413 (1996); F. Verstraete, K. Audenaert, J. Dehaene and B. D. Moor, J. Phys. A. **34**, 10327 (2001).
- [12] S. Lee, D. P. Chi, S. D. Oh and J. Kim, Phys. Rev. A. 68, 062304 (2003).
- [13] A. Sanpera, D. Bruss and M. Lewenstein, Phys. Rev. A. 63,




FIG. 6: Plot of (i) concurrence C (or $E_2^{(2,2)}$) and (ii) f as a function of the time of evolution t. Here we consider the case of QND interaction (T = 5, r = 0.1), in the collective decoherence regime ($r_{12} = 0.05$).




FIG. 7: Plot of (i) concurrence C (or $E_2^{(2,2)}$) and (ii) f as a function of the time of evolution t, for the case of QND interaction (T = 5, r = 0.1), in the independent decoherence regime $(r_{12} = 1.1)$.

050301 (2001).

- [14] J. Sperling and W. Vogel, Physica Scripta 83, 045002 (2011).
- [15] C. H. Bennett, G. Brassard, S. Popescu, B. Schumacher, J. A. Smolin and W. K. Wooters, Phys. Rev. Lett. 76, 722 (1996).
- [16] M. Horodecki, P. Horodecki and R. Horodecki, Phys. Rev. A 60, 1888 (1999).
- [17] G. Gour, Phys. Rev. A 71, 012318 (2005).
- [18] M. Horodecki and P. Horodecki, Phys. Rev. A 59, 4206 (1999).
- [19] M-J. Zhao, Z-G.Li, S-M. Fei and Z-X. Wang, J. Phys. A:Math. Theor. 43, 275203 (2010).
- [20] B. M. Terhal and P. Horodecki, Phys. Rev. A 61, 040301(R) (2000).

- [21] V. Karimipour and L. Memarzadeh, Phys. Rev. A 73, 012329 (2006).
- [22] S. Banerjee and R. Ghosh, J. Phys. A:Math. Theor. 40, 1273 (2007).
- [23] S. Banerjee and R. Srikanth, Euro. Phys. J. D 46, 335 (2008).
- [24] R. Srikanth and S. Banerjee, Phys. Rev. A 77, 012318 (2008).
- [25] S. Banerjee, V. Ravishankar and R. Srikanth, Ann. of Phys. (N. Y.) 325, 816 (2010).
- [26] S. Banerjee, V. Ravishankar and R. Srikanth, Euro. Phys. J. D 56, 277 (2010).




FIG. 8: Plot of (i) concurrence C (or $E_2^{(2,2)}$) and (ii) f as a function of squeezing parameter r. Here we consider the case of QND interaction (T = 5), in the collective decoherence regime ($r_{12} = 0.05$) and time of evolution t = 2.