On small set of one-way LOCC indistinguishability of maximally entangled states

Yan-Ling Wang¹, Mao-Sheng Li², Zhu-Jun Zheng¹, Shao-Ming Fei^{3,4}

¹Department of Mathematics, South China University of Technology, Guangzhou 510640, P.R.China

²Department of Mathematical of Science, Tsinghua University, Beijing 100084, P.R.China

³School of Mathematical Sciences, Capital Normal University, Beijing 100048, China

⁴Max-Planck-Institute for Mathematics in the Sciences, 04103 Leipzig, Germany

July 10, 2018

Abstract In this paper, we study the one-way local operations and classical communication (LOCC) problem. In $\mathbb{C}^d \otimes \mathbb{C}^d$ with $d \geq 4$, we construct a set of $3\lceil \sqrt{d} \rceil - 1$ one-way LOCC indistinguishable maximally entangled states which are generalized Bell states. Moreover, we show that there are four maximally entangled states which cannot be perfectly distinguished by one-way LOCC measurements for any dimension $d \geq 4$.

1 Introduction

In compound quantum systems, many global operators can not be implemented using only local operations and classical communication (LOCC). This reflects the fundamental feature of quantum mechanics called nonlocality. Meanwhile, the understanding of the limitation of quantum operators that can be implemented by LOCC is also one of the significant subjects in quantum information theory. And local distinguishability of quantum states plays an important role in exploring quantum nonlocality [1, 2]. In the bipartite case, Alice and Bob share a quantum system which is chosen from one of a known set of mutually orthogonal quantum states. Their goal is to identify the given state using only LOCC. The nonlocality of quantum information is therefore revealed when a set of or-

thogonal states can not be distinguished by LOCC. Moreover, the local distinguishability has been found practical applications in quantum cryptography primitives such as secret sharing and data hiding [3, 4].

The question of local discrimination of orthogonal quantum states has received considerable attentions in recent years [5-19]. It is well known that any two orthogonal maximally entangled states can be perfectly distinguished with LOCC [2]. In Refs. [8, 9], the authors proved that a set of d+1 or more maximally entangled states in $d \otimes d$ systems are not perfectly locally distinguishable. Hence it is interesting to ask whether there are locally indistinguishable sets consisting of d or fewer maximally entangled states in $d \otimes d$. For d = 3, Nathanson has shown that any three maximally entangled states can be perfectly distinguished [6]. Recently, the authors in [15, 17] considered one-way LOCC distinguishability and presented sets of d and d-1 indistinguishable maximally entangled states for d = 5, ..., 10. The problem remains open if there exists fewer than d - 1indistinguishable maximally entangled states for arbitrary dimension d. More recently, Nathanson showed that there exist triples of mutually orthogonal maximally entangled states in $\mathbb{C}^d \otimes \mathbb{C}^d$ which cannot be distinguished with one-way LOCC when d is even or $d \equiv 2 \mod 3$ [16]. In addition, the authors in [18] gave a set with $\lceil \frac{d}{2} \rceil + 2$ maximally entangled states in $\mathbb{C}^d \otimes \mathbb{C}^d$ which is one-way LOCC indistinguishable, where $\lceil a \rceil$ means the least integer which is not less than a. And in [19], the authors presented sets with four and five maximally entangled states in $\mathbb{C}^{4m} \otimes \mathbb{C}^{4m}$ which is one-way LOCC indistinguishable but two-way distinguishable. Whether there are four or three one-way LOCC indistinguishable maximally entangled states in arbitrary dimension remains unknown.

In this paper, we give a positive answer to this question when the number of states in the set is four. First for any dimension $d \geq 4$, we give a set of $3\lceil \sqrt{d} \rceil - 1$ one-way LOCC indistinguishable maximally entangled states. Moreover, we can find four maximally entangled states which cannot be perfectly distinguished by one-way LOCC measurements for any dimension $d \geq 4$.

2 Preliminaries

We first introduce some basic results that will be used in proving our theorems. Under the computational base $\{|ij\rangle\}_{i,j=0}^{d-1}$ of Hilbert space $\mathbb{C}^d\otimes\mathbb{C}^d$, the generalized Bell states are defined as follows:

$$|\psi_{nm}\rangle = I \otimes U_{nm}(\frac{1}{\sqrt{d}} \sum_{j=0}^{d-1} |jj\rangle),$$
 (1)

where $U_{mn}=X^mZ^n$ are generalized Pauli matrices constituting a basis of unitary operators, and $X|j\rangle=|j\oplus_d 1\rangle,\ Z|j\rangle=\omega^j|j\rangle,\ \omega=e^{\frac{2\pi\sqrt{-1}}{d}}$. We define $V_{mn}=U_{mn}^T$, where T stands for transpose. It is directly verified that $ZX=\omega XZ$.

Lemma 1. Suppose $U_{mn} = X^m Z^n, U_{m'n'} = X^{m'} Z^{n'}$, we have

$$U_{m'n'}^{\dagger}U_{mn} = \omega^{(m'-m)n'}U_{(m-m' \bmod d)(n-n' \bmod d)}.$$

Proof:

$$U_{m'n'}^{\dagger}U_{mn} = (X^{m'}Z^{n'})^{\dagger}(X^{m}Z^{n})$$

$$= (Z^{\dagger n'}X^{\dagger m'})(X^{m}Z^{n})$$

$$= (Z^{(d-1)n'}X^{(d-1)m'})(X^{m}Z^{n})$$

$$= (Z^{-n'}X^{-m'})(X^{m}Z^{n})$$

$$= Z^{-n'}X^{m-m'}Z^{n}$$

$$= \omega^{(m'-m)n'}X^{m-m'}Z^{n-n'}$$

$$= \omega^{(m'-m)n'}U_{(m-m'\bmod d)(n-n'\bmod d)}.$$

For the convenience of citation, we recall the results given in Refs. [16, 17].

Lemma 2. [17] In $\mathbb{C}^d \otimes \mathbb{C}^d$, $N \leq d$ number of pairwise orthogonal maximally entangled states $|\psi_{n_i m_i}\rangle$, $i=1,2,\ldots,N$, taken from the set given in Eq. (1), can be perfectly distinguished by one-way LOCC $A \to B$, if and only if there exists at least one state $|\alpha\rangle \in \mathcal{H}_B$ for which the states $U_{n_1 m_1} |\alpha\rangle$, $U_{n_2 m_2} |\alpha\rangle$, ..., $U_{n_N m_N} |\alpha\rangle$ are pairwise orthogonal.

On the other hand, the set is perfectly distinguishable by one-way LOCC in the $B \to A$, if and only if there exists at least one state $|\alpha\rangle \in \mathcal{H}_A$ for which the states $V_{n_1m_1}|\alpha\rangle, V_{n_2m_2}|\alpha\rangle, \ldots, V_{n_Nm_N}|\alpha\rangle$ are pairwise orthogonal.

Lemma 3. [16] Given a set of states $S = \{|\psi_i\rangle = (I \otimes U_i)|\phi\rangle\} \subset \mathbb{C}^d \otimes \mathbb{C}^d$, with $|\phi\rangle$ the standard maximally entangled state. The elements of S can be perfectly distinguished with one-way LOCC if and only if there exists a set of states $\{|\phi_k\rangle\} \subset \mathbb{C}^d$ and a set of positive numbers $\{m_k\}$ such that $\sum_k m_k |\phi_k\rangle \langle \phi_k| = I_d$ and $\langle \phi_k | U_j^{\dagger} U_i | \phi_k \rangle = \delta_{ij}$.

In the following, we concentrate ourselves on the set of maximally entangled states. Any maximally entangled state in $\mathbb{C}^d \otimes \mathbb{C}^d$ can be written as $|\psi\rangle = (I \otimes U)|\psi_0\rangle$, where $|\psi_0\rangle = \frac{1}{\sqrt{d}} \sum_{i=1}^d |ii\rangle$, and U is a unitary matrix. Since there is a one to one correspondence between a maximally entangled state $|\psi_i\rangle$ and the unitary matrix U_i , we call the set of unitary matrices $\{U_i\}_{i=1}^d$ the defining unitary matrices of the set of maximally entangled states $\{|\psi_i\rangle\}_{i=1}^d$.

3 Sets of one-way LOCC indistinguishable states

The authors in [18] presented a set with $\lceil \frac{d}{2} \rceil + 2$ generalized Bell states in $\mathbb{C}^d \otimes \mathbb{C}^d$ which is one-way LOCC indistinguishable. In the following, firstly, we also consider the one-way distinguishability of generalized Bell states.

Theorem 1. In $\mathbb{C}^d \otimes \mathbb{C}^d$ $(d \ge 4)$, there exists an orthogonal set with $3\lceil \sqrt{d} \rceil - 1$ maximally entangled states which is one-way LOCC indistinguishable:

$$\{|\psi_{00}\rangle, |\psi_{10}\rangle, \dots, |\psi_{n-1,0}\rangle, |\psi_{2n-1,0}\rangle, |\psi_{3n-1,0}\rangle, |\psi_{4n-1,0}\rangle, \dots, |\psi_{(n-1)n-1,0}\rangle, |\psi_{d-1,0}\rangle, |\psi_{n-1,1}\rangle, |\psi_{2n-1,1}\rangle, |\psi_{3n-1,1}\rangle, |\psi_{4n-1,1}\rangle, \dots, |\psi_{(n-1)n-1,1}\rangle, |\psi_{d-1,1}\rangle\}, \text{ where } n = \lceil \sqrt{d} \rceil.$$

The corresponding unitary matrices are given by

$$\{U_{00}, U_{10}, \dots, U_{n-1,0}, U_{2n-1,0}, U_{3n-1,0}, U_{4n-1,0}, \dots, U_{(n-1)n-1,0}, U_{d-1,0}, U_{n-1,1}, U_{2n-1,1}, U_{3n-1,1}, U_{4n-1,1}, \dots, U_{(n-1)n-1,1}, U_{d-1,1}\}.$$

Proof: If $\{|\psi_{00}\rangle, |\psi_{10}\rangle, \dots, |\psi_{n-1,0}\rangle, |\psi_{2n-1,0}\rangle, |\psi_{3n-1,0}\rangle, \dots, |\psi_{(n-1)n-1,0}\rangle, |\psi_{d-1,0}\rangle, |\psi_{n-1,1}\rangle, |\psi_{2n-1,1}\rangle, \dots, |\psi_{(n-1)n-1,1}\rangle, |\psi_{d-1,1}\rangle\}$ can be one-way LOCC distinguished, then by lemma $2, \exists |\alpha\rangle \neq 0 \in \mathbb{C}^d$, such that the set $\{U_{00}|\alpha\rangle, U_{10}|\alpha\rangle, \dots, U_{n-1,0}|\alpha\rangle, U_{2n-1,0}|\alpha\rangle, U_{3n-1,0}|\alpha\rangle, \dots, U_{(n-1)n-1,0}|\alpha\rangle, U_{d-1,0}|\alpha\rangle, U_{n-1,1}|\alpha\rangle, U_{2n-1,1}|\alpha\rangle, \dots, U_{(n-1)n-1,1}|\alpha\rangle, U_{d-1,1}|\alpha\rangle\}$ are mutually orthogonal.

From the orthogonality of $U_{00}|\alpha\rangle$ and $U_{10}|\alpha\rangle, U_{20}|\alpha\rangle, \dots, U_{n-1,0}|\alpha\rangle$, we obtain

$$\langle \alpha | U_{10} | \alpha \rangle = \sum_{j=0}^{d-1} \omega^j \alpha_j \overline{\alpha}_j = 0,$$

$$\langle \alpha | U_{20} | \alpha \rangle = \sum_{j=0}^{d-1} \omega^{2j} \alpha_j \overline{\alpha}_j = 0,$$

$$\vdots$$

$$\langle \alpha | U_{n-1,0} | \alpha \rangle = \sum_{j=0}^{d-1} \omega^{(n-1)j} \alpha_j \overline{\alpha}_j = 0.$$

Then by the orthogonality of $U_{2n-1,0}|\alpha\rangle$ and $U_{n-1,0}|\alpha\rangle,\ldots,U_{10}|\alpha\rangle,U_{00}|\alpha\rangle$, taking into

account with the lemma 1, we get

$$\langle \alpha | U_{n-1,0}^{\dagger} U_{2n-1,0} | \alpha \rangle = \langle \alpha | U_{n,0} | \alpha \rangle = \sum_{j=0}^{d-1} \omega^{nj} \alpha_j \overline{\alpha}_j = 0,$$

$$\vdots$$

$$\langle \alpha | U_{10}^{\dagger} U_{2n-1,0} | \alpha \rangle = \langle \alpha | U_{2n-2,0} | \alpha \rangle = \sum_{j=0}^{d-1} \omega^{(2n-2)j} \alpha_j \overline{\alpha}_j = 0,$$

$$\langle \alpha | U_{00}^{\dagger} U_{2n-1,0} | \alpha \rangle = \langle \alpha | U_{2n-1,0} | \alpha \rangle = \sum_{j=0}^{d-1} \omega^{(2n-1)j} \alpha_j \overline{\alpha}_j = 0.$$

Similarly, from the orthogonality of $U_{3n-1,0}|\alpha\rangle$, $U_{4n-1,0}|\alpha\rangle$, ..., $U_{(n-1)n-1,0}|\alpha\rangle$, $U_{d-1,0}|\alpha\rangle$ and $U_{n-1,0}|\alpha\rangle$, ..., $U_{10}|\alpha\rangle$, we have:

$$\sum_{j=0}^{d-1} \omega^{(2n)j} \alpha_j \overline{\alpha}_j = \sum_{j=0}^{d-1} \omega^{(2n+1)j} \alpha_j \overline{\alpha}_j = \dots = \sum_{j=0}^{d-1} \omega^{(d-1)j} \alpha_j \overline{\alpha}_j = 0.$$

Putting the above d-1 equations together, we have

$$\sum_{j=0}^{d-1} \omega^j \alpha_j \overline{\alpha}_j = \sum_{j=0}^{d-1} \omega^{2j} \alpha_j \overline{\alpha}_j = \sum_{j=0}^{d-1} \omega^{3j} \alpha_j \overline{\alpha}_j = \dots = \sum_{j=0}^{d-1} \omega^{(d-1)j} \alpha_j \overline{\alpha}_j = 0.$$

Solving these d-1 equations, we have $(\alpha_0 \overline{\alpha}_0, \alpha_1 \overline{\alpha}_1, \cdots, \alpha_{d-1} \overline{\alpha}_{d-1}) = \lambda(1, 1, \cdots, 1)$.

- 1) If $\lambda = 0$, then $(\alpha_0 \overline{\alpha}_0, \alpha_1 \overline{\alpha}_1, \cdots, \alpha_{d-1} \overline{\alpha}_{d-1}) = (0, 0, \cdots, 0)$, that is, $|\alpha\rangle = \mathbf{0}$.
- 2) If $\lambda \neq 0$, then for $\forall i, j$, we have $\alpha_i \overline{\alpha}_j \neq 0$. By the orthogonality of $U_{n-1,1} | \alpha \rangle$ and $U_{n-1,0} | \alpha \rangle, \dots, U_{20} | \alpha \rangle, U_{10} | \alpha \rangle, U_{00} | \alpha \rangle$ and lemma 1, we have

$$\langle \alpha | U_{n-1,0}^{\dagger} U_{n-1,1} | \alpha \rangle = \langle \alpha | U_{01} | \alpha \rangle = \sum_{j=0}^{d-1} \omega^{0j} \alpha_j \overline{\alpha}_{j \oplus_d 1} = 0,$$

$$\vdots$$

$$\langle \alpha | U_{10}^{\dagger} U_{n-1,1} | \alpha \rangle = \langle \alpha | U_{n-2,1} | \alpha \rangle = \sum_{j=0}^{d-1} \omega^{(n-2)j} \alpha_j \overline{\alpha}_{j \oplus_d 1} = 0,$$

$$\langle \alpha | U_{00}^{\dagger} U_{n-1,1} | \alpha \rangle = \langle \alpha | U_{n-1,1} | \alpha \rangle = \sum_{j=0}^{d-1} \omega^{(n-1)j} \alpha_j \overline{\alpha}_{j \oplus_d 1} = 0.$$

By the orthogonality of $U_{2n-1,1}|\alpha\rangle$, $U_{3n-1,1}|\alpha\rangle$, ..., $U_{(n-1)n-1,1}|\alpha\rangle$, $U_{d-1,1}|\alpha\rangle$ and $U_{00}|\alpha\rangle$, $U_{10}|\alpha\rangle$, $U_{20}|\alpha\rangle$, ..., $U_{n-1,0}|\alpha\rangle$, we have

$$\sum_{j=0}^{d-1} \omega^{nj} \alpha_j \overline{\alpha}_{j \oplus_d 1} = \sum_{j=0}^{d-1} \omega^{(n+1)j} \alpha_j \overline{\alpha}_{j \oplus_d 1} = \dots = \sum_{j=0}^{d-1} \omega^{(d-1)j} \alpha_j \overline{\alpha}_{j \oplus_d 1} = 0.$$

From the above equations, $(\alpha_0 \overline{\alpha}_1, \alpha_1 \overline{\alpha}_2, \dots, \alpha_{d-1} \overline{\alpha}_0) = (0, 0, \dots, 0)$ and $\alpha_i \overline{\alpha}_j \neq 0$ are contradictory. Therefore $\{|\psi_{00}\rangle, |\psi_{10}\rangle, \dots, |\psi_{n-1,0}\rangle, |\psi_{2n-1,0}\rangle, |\psi_{3n-1,0}\rangle, \dots, |\psi_{(n-1)n-1,0}\rangle, |\psi_{d-1,0}\rangle, |\psi_{n-1,1}\rangle, |\psi_{2n-1,1}\rangle, \dots, |\psi_{(n-1)n-1,1}\rangle, |\psi_{d-1,0}\rangle\}$ cannot be one-way LOCC distinguished.

Remark: It should be noticed that the above result may be worse than the known $\lceil \frac{d}{2} \rceil + 2$ result [18] in the case of small d. And $3\lceil \sqrt{d} \rceil - 1 \le \lceil \frac{d}{2} \rceil + 2$ when $d \ge 30$, so our theorem gives a smaller one-way LOCC indistinguishable maximal entangled states in this case.

In the above discussions, we restrict ourselves on the one-way LOCC indistinguished generalized Bell states. In the following we consider general orthogonal maximally entangled states that are indistinguishable under one-way LOCC.

Theorem 2. There exist four mutually orthogonal maximally entangled states in $\mathbb{C}^d \otimes \mathbb{C}^d$ which cannot be distinguished under one-way LOCC for odd $d \geq 7$.

Proof: Set d = 2 + r, $r \ge 5$. Let P denote the $r \times r$ permutation matrix,

$$P = \begin{bmatrix} 0 & 0 & \cdots & 0 & 1 \\ 1 & 0 & \cdots & 0 & 0 \\ 0 & 1 & \cdots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & 1 & 0 \end{bmatrix}_{r \times r}.$$

Clearly, $P^r = I$ where I denotes the $r \times r$ identity matrix. We set $U_0 = I_d$,

$$U_1 = \left[egin{array}{cc} \omega X & & \\ & P \end{array}
ight], \quad U_2 = \left[egin{array}{cc} \gamma Z & & \\ & P^2 \end{array}
ight], \quad U_3 = \left[egin{array}{cc} \sigma Y & & \\ & P^{rac{r+1}{2}} \end{array}
ight],$$

where ω, γ and σ are phases satisfying $|\omega| = |\gamma| = |\sigma| = 1$, $\overline{\gamma} \neq \pm i\overline{\omega}^2$, X, Y, Z are the Pauli matrices:

$$X = \left[egin{array}{cc} 0 & 1 \\ 1 & 0 \end{array}
ight], \quad Y = \left[egin{array}{cc} 0 & -i \\ i & 0 \end{array}
ight], \quad Z = \left[egin{array}{cc} 1 & 0 \\ 0 & -1 \end{array}
ight].$$

Let $|\psi_0\rangle$ be the standard maximally entangled state, $|\psi_0\rangle = \sum_{i=0}^{d-1} |ii\rangle$. We construct four maximally entangled states as follows:

$$\{(I\otimes U_0)|\psi_0\rangle, (I\otimes U_1)|\psi_0\rangle, (I\otimes U_2)|\psi_0\rangle, (I\otimes U_3)|\psi_0\rangle\}\subseteq \mathbb{C}^d\otimes \mathbb{C}^d.$$

One can check that these states are mutually orthogonal and maximally entangled.

Suppose that Alice performs an initial measurement $\mathbb{M} = \{M_k\}_{k=1}^n$ on her system and gets the measurement outcome corresponding to some operator M_k $(1 \le k \le n)$ of the following form:

$$M_k = \left[\begin{array}{cc} A_k & C_k^{\dagger} \\ C_k & B_k \end{array} \right] \geqslant 0,$$

where A_k is a 2 × 2 matrix and B_k a $r \times r$ matrix.

By lemma 3, all the measurements of Alice's can be chosen to be rank one. So we suppose all the matrices M_k $(1 \le k \le n)$ are rank one and $M_k = |\phi_k\rangle\langle\phi_k|$ for some $|\phi_k\rangle \in \mathbb{C}^d$. In order to distinguish the above four states by one-way LOCC, we must have

$$0 = \langle \phi_k | U_i^{\dagger} U_i | \phi_k \rangle = Tr(U_i^{\dagger} U_i | \phi_k \rangle \langle \phi_k |) = Tr(U_i | \phi_k \rangle \langle \phi_k | U_i^{\dagger}) = Tr(U_i M_k U_i^{\dagger}), \ i \neq j.$$

That is, $Tr(U_iM_kU_j^{\dagger})=0$, whenever $i\neq j$. By specify choosing i and j, we obtain the following equations:

$$Tr(U_1M_k) = \omega Tr(A_kX) + Tr(B_kP) = 0, (2)$$

$$Tr(U_2M_k) = \gamma Tr(A_kZ) + Tr(B_kP^2) = 0,$$
 (3)

$$Tr(U_3M_k) = \sigma Tr(A_kY) + Tr(B_kP^{\frac{r+1}{2}}) = 0,$$
 (4)

$$Tr(U_2M_kU_1^{\dagger}) = -i\overline{\omega}\gamma Tr(A_kY) + Tr(B_kP) = 0,$$
(5)

$$Tr(U_3 M_k U_1^{\dagger}) = -i\overline{\omega}\sigma Tr(A_k Z) + Tr(B_k P^{\frac{r-1}{2}}) = 0,$$
(6)

$$Tr(U_3M_kU_2^{\dagger}) = -i\overline{\gamma}\sigma Tr(A_kX) + Tr(B_kP^{\frac{r-3}{2}}) = 0.$$
 (7)

From equations (2) and (5), we have

$$\omega \operatorname{Tr}(A_k X) + i \overline{\omega} \gamma \operatorname{Tr}(A_k Y) = 0. \tag{8}$$

After easily calculation, we can obtain $Tr(A_kX) = A_k(1,2) + A_k(2,1)$ and $Tr(A_kY) = iA_k(1,2) - iA_k(2,1)$. Since A_k is a Hermitian matrix, then both $Tr(A_kX)$ and $Tr(A_kY)$ are real numbers. Moving the second term of equation (8) to the right hand side then taking the norm of each side, we have $|Tr(A_kX)| = |Tr(A_kY)|$. If $Tr(A_kX) \neq 0$, then we have $i\overline{\omega}^2\gamma = -\frac{Tr(A_kX)}{Tr(A_kY)} = 1$ or -1. This is contradicted with $\overline{\gamma} \neq \pm i\overline{\omega}^2$. Hence we have $Tr(A_kX) = Tr(A_kY) = 0$. Substituting $Tr(A_kY) = 0$ into equation (4), we

obtain $Tr(B_kP^{\frac{r+1}{2}})=0$. Due to $P^r=I$ and the Hermitian of the matrix B_k , the equality $Tr(B_kP^{\frac{r-1}{2}})=\overline{Tr(B_kP^{\frac{r+1}{2}})}$ holds, which gives rise to $Tr(B_kP^{\frac{r-1}{2}})=0$. Then by equation (6), we obtain $Tr(A_kZ)=0$. Equations $Tr(A_kX)=Tr(A_kY)=Tr(A_kZ)=0$ give that $A_k=tI_2$ for some $t_k\in\mathbb{R}$. Noticing that we have assumed $\mathrm{rank}(M_k)=1$, so $\mathrm{rank}(A_k)\leq 1$. Hence $A_k=\mathbf{0}$ for all $1\leq k\leq n$. But now $\sum_k^n M_{k=1}$ cannot equal to the identity I for the 2×2 matrix of the left upper corner must equal to zero. This makes a contradiction.

Hence, we can conclude that the four states we construct above can not be distinguished by one-way LOCC.

Corollary. There exist four mutually orthogonal maximally entangled states in $\mathbb{C}^d \otimes \mathbb{C}^d$ which cannot be distinguished under one-way LOCC for $d \geq 4$.

Proof: By the above theorem, we only need to check for the cases: d is even and d = 5. For all these cases, it has been showed that there exist three mutually orthogonal maximally entangled states which cannot be distinguished under one-way LOCC in Ref.[16]. And there exists another maximally entangled state orthogonal to all the three states. So after adding such a state, these four states cannot be distinguished by one-way LOCC.

4 Conclusion

We study the one-way LOCC problem and present a set of $3\lceil \sqrt{d} \rceil - 1$ one-way LOCC indistinguishable maximally entangled states which are all generalized Bell states. It should be noticed that if d is large enough, then the number $3\lceil \sqrt{d} \rceil - 1$ is much smaller than the number $\lceil \frac{d}{2} \rceil + 2$ in [18]. But for small d (less than 30), our results are not so good as the known results. In addition to, we have also found four maximally entangled states which cannot be perfectly distinguished by one-way LOCC measurements for any dimension $d \ge 4$. For some particular dimension d, small one-way indistinguishable sets that contain only three states has been given in [16]. The question whether there exist three one-way indistinguishable maximally entangled states for arbitrary $d \ge 4$ remains open.

Acknowledgments We are thankful for the referees' suggestions and careful reading of our paper. This work is supported by NSFC (Grant Nos. 11475178, 11275131).

References

- [1] Bennett C.H., DiVincenzo D.P., Fuchs C.A., Mor T., Rains E., Shor P.W., Smolin J.A. and Wootters W.K.: Quantum nonlocality without entanglement. *Phys. Rev. A*, 59:1070-1091 (1999)
- [2] Walgate J. and Hardy L.: Nonlocality asymmetry and distinguishing bipartite states. Phys. Rev. Lett, 89, 147901 (2002)
- [3] DiVincenzo D.P., Leung D.W. and Terhal B.M.: Quantum data hiding. IEEE Trans. Inf. Theory 48, 580 (2002)
- [4] Markham D. and Sanders B. C.: Graph States for Quantum Secret Sharing. Phys. Rev. A 78, 042309 (2008)
- [5] Walgate J., Short A. J., Hardy L. and Vedral V.: Local Distinguishability of Multipartite Orthogonal Quantum States. *Phys. Rev. Lett* 85, 4972 (2000)
- [6] Nathanson M.: Distinguishing bipartitite orthogonal states using LOCC: Best and worst cases. J. Math. Phys. 46, 062103 (2005)
- [7] Ghosh S., Kar G., Roy A., Sen(De)A. and Sen U.: Distinguishability of Bell states.

 Phys. Rev. Lett 87, 277902 (2001)
- [8] Ghosh S., Kar G., Roy A. and Sarkar D.: Distinguishability of maximally entangled states. *Phys. Rev. A* **70**, 022304 (2004)
- [9] Fan H.: Distinguishability and Indistinguishability by Local Operations and Classical Communication. Phys. Rev. Lett 92, 177905 (2004)
- [10] Yu N., Duan R. and Ying M.: Four Locally Indistinguishable Ququad-Ququad Orthogonal Maximally Entangled States. Phys. Rev. Lett 109, 020506 (2012)
- [11] Cosentino A.: Positive-partial-transpose-indistinguishable states via semidefinite programming. *Phys. Rev. A*, **87**, 012321 (2013)

- [12] Yu N., Duan R. and Ying M.: Distinguishability of Quantum States by Positive Operator-Valued Measures With Positive Partial Transpose. *IEEE Trans. Inf. Theory* Vol. 60, No.4 (2014)
- [13] Li M.-S., Wang Y.-L., Fei S.-M. and Zheng Z.-J.: d locally indistinguishable maximally entangled states in $\mathbb{C}^d \otimes \mathbb{C}^d$. Phys. Rev. A. **91**, 042318 (2015)
- [14] Yu S.-X. and Oh C.H.: Detecting the local indistinguishability of maximally entangled states. arXiv:1502.01274v1(2015)
- [15] Bandyopadhyay S., Ghosh S. and Kar G.: LOCC distinguishability of unilaterally transformable quantum states. *New J. Phys.* **13**, 123013 (2011)
- [16] Nathanson M.: Three maximally entangled states can require two-way local operations and classical communication for local discrimination. *Phys. Rev. A*, 88, 062316 (2013)
- [17] Zhang Z.-C., Wen Q.-Y., Gao F., Tian G.-J. and Cao T.-Q.: One-way LOCC indistinguishability of maximally entangled states. *Quantum Inf. Proc.* **13**,795 (2014)
- [18] Zhang Z.-C., Feng K.-Q., Gao F. and Wen Q.-Y.: Distinguishing maximally entangled states by one-way local operations and classical communication. *Phys. Rev. A.* 91, 012329 (2015)
- [19] Tian G.-J., Yu S.-X., Gao F., Wen Q.-Y. and Oh C.H.: Local discrimination of four or more maximally entangled states. *Phys. Rev. A.* **91**, 052314 (2015)