

Measure of genuine coherence based of quasi-relative entropy

Anna Vershynina

Department of Mathematics, Philip Guthrie Hoffman Hall, University of Houston, 3551 Cullen Blvd., Houston, TX 77204-3008, USA

June 1, 2022

Abstract

We present a genuine coherence measure based on a quasi-relative entropy as a difference between quasi-entropies of the dephased and the original states. The measure satisfies non-negativity and monotonicity under genuine incoherent operations (GIO). It is strongly monotone under GIO in two- and three-dimensions, or for pure states in any dimension, making it a genuine coherence monotone. We provide a bound on the error term in the monotonicity relation under GIO in terms of the trace distance between the original and the dephased states. Moreover, the lower bound on the coherence measure can also be calculated in terms of this trace distance.

1 Introduction

Quantum coherence is a fundamental property of quantum systems, describing the existence of quantum interference. It is widely used in thermodynamics [1, 9, 19], transport theory [28, 38], and quantum optics [13, 30], among few applications. Recently, problems involving coherence included quantification of coherence [2, 22, 26, 27, 31, 40], distribution [25], entanglement [7, 33], operational resource theory [5, 7, 12, 37], correlations [17, 20, 34], with only a few references mentioned in each. See [32] for a more detailed review.

As a golden standard it is taken that any “good” coherence measure should satisfy four criteria presented in [2]: vanishing on incoherent states; monotonicity under incoherent operations; strong monotonicity under incoherent operations, and convexity. Alternatively, the last two properties can be substituted by an additivity for subspace independent states, which was shown in [40].

A number of ways has been proposed as a coherence measure, but only a few satisfy all necessary cri-

teria [2, 41, 42]. A broad class of coherence measure are defined as the minimal distance D to the set of incoherent states \mathcal{I} , as

$$\mathcal{C}_D(\rho) = \min_{\delta \in \mathcal{I}} D(\rho, \delta).$$

In [2], it was shown that coherence vanishes on incoherent states when the distance vanishes only on identical states; the measure is monotone when the distance is contractive under quantum channels; and it is convex when the distance is jointly convex. Strong monotonicity property is more challenging to pinpoint. Measures that satisfy the strong monotonicity that have been introduced up to date, are based on l_1 , relative entropy, Tsallis entropy, and real symmetric concave functions on a probability simplex.

Another approach to generate physically relevant coherence measures is to consider different incoherent operations. The largest class of incoherent operations is called maximally incoherent (MIO), and it consists of all completely-positive trace-preserving (CPTP) maps that preserve the set of incoherent states. The smaller set, called incoherent operations (IO) [2], has Krauss

operators that each preserve the set of incoherent states (see Definition 2.3). A smaller set consists of strictly incoherent operations (SIO) [37, 39], which are the result of action on a primary and ancillary systems that do not generate coherence on a primary system, see Definition 2.6. And the last class of operations that is discussed in this paper, is called genuine incoherent operations (GIO) [10], which act trivially on incoherent states, see Definition 2.4. See [6] for a larger list of incoherent operations, and their comparison. For these types of incoherent operations one may look at similar properties as the ones presented in [2]. Restricted to GIO, one would obtain a measure of genuine coherence when it is non-negative and monotone, or a coherence monotone when it is also strongly monotone under GIO.

In [10], the following genuine coherence measure was proposed:

$$\mathcal{C}_D(\rho) = D(\rho \parallel \Delta(\rho)) ,$$

for a distance D , and $\Delta(\rho)$ being the dephased state in a pre-fixed basis, see Notation 2.2. It was shown that this is a genuine coherence measure if the distance is contractive under unital operations. In fact, the monotonicity holds not only for GIO maps but for dephasing-covariant incoherent operations (DIO) as well (the ones that commute with the dephasing operator).

Here we propose another genuine coherence measure based on a quasi-relative entropy:

$$\mathcal{C}_f(\rho) = S_f(\Delta(\rho)) - S_f(\rho) ,$$

here $S_f(\rho)$ is a quasi entropy, which could be defined in two ways, one of which is $S_f(\rho) = -S_f(\rho \parallel I)$. The motivation for this definition comes from the relative entropy coherence. It was shown [2] that for a relative entropy $S(\cdot \parallel \cdot)$, there is a closed expression of a distance-based coherence measure:

$$\min_{\delta \in \mathcal{I}} S(\rho \parallel \delta) = S(\rho \parallel \Delta(\rho)) = S(\Delta(\rho)) - S(\rho) .$$

In general, for quasi-relative entropies neither of these equalities will hold. This can be seen for Tsallis

relative entropy, which is a particular case of a quasi-relative entropy. The closest incoherent state is given in [27], and it is not a dephased state $\Delta(\rho)$. The second equality does not hold either in general.

We show that quasi-relative entropy coherence, which we call f -coherence, is unique for pure states, non-negative, zero if and only if a state is incoherent, and monotone under GIO maps. Moreover, we give a lower bound on this coherence in terms of a trace distance between a state and its dephased state, we provide an if and only if condition on a GIO map that saturates the monotonicity relation, and bound the error term in the monotonicity relation. Additionally, we investigate when the f -coherence would be monotone under a larger class of SIO maps.

We show that f -coherence saturates strong monotonicity under GIO maps in two- and three-dimensions, and it satisfies the strong monotonicity under GIO maps in any dimensions for pure states.

2 Preliminaries

2.1 Coherence

Let \mathcal{H} be a d -dimensional Hilbert space. Let us fix a basis $\mathcal{E} = \{|j\rangle\}_{j=1}^d$ of vectors in \mathcal{H} .

2.1 Definition. A state δ is called *incoherent* if it can be represented as follows

$$\delta = \sum_j \delta_j |j\rangle \langle j| .$$

2.2 Notation. Denote the set of **incoherent states** for a fixed basis $\mathcal{E} = \{|j\rangle\}_j$ as

$$\mathcal{I} = \{\rho = \sum_j p_j |j\rangle \langle j|\} .$$

A **dephasing** operation in \mathcal{E} basis is the following map:

$$\Delta(\rho) = \sum_j \langle j | \rho | j \rangle |j\rangle \langle j| .$$

2.3 Definition. A CPTP map Φ with the following Kraus operators

$$\Phi(\rho) = \sum_n K_n \rho K_n^* ,$$

is called **the incoherent operation (IO)** or *incoherent CPTP (ICPTP)*, when the Kraus operators satisfy

$$K_n \mathcal{I} K_n^* \subset \mathcal{I}, \text{ for all } n ,$$

besides the regular completeness relation $\sum_n K_n^* K_n = \mathbb{1}$.

Any reasonable measure of coherence $\mathcal{C}(\rho)$ should satisfy the following conditions

- (C1) $\mathcal{C}(\rho) = 0$ if and only if $\rho \in \mathcal{I}$;
- (C2) Non-selective monotonicity under IO maps (monotonicity)

$$\mathcal{C}(\rho) \geq \mathcal{C}(\Phi(\rho)) ;$$

- (C3) Selective monotonicity under IO maps (strong monotonicity)

$$\mathcal{C}(\rho) \geq \sum_n p_n \mathcal{C}(\rho_n) ,$$

where p_n and ρ_n are the outcomes and post-measurement states

$$\rho_n = \frac{K_n \rho K_n^*}{p_n}, \quad p_n = \text{Tr} K_n \rho K_n^* .$$

- (C4) Convexity,

$$\sum_n p_n \mathcal{C}(\rho_n) \geq \mathcal{C} \left(\sum_n p_n \rho_n \right) ,$$

for any sets of states $\{\rho_n\}$ and any probability distribution $\{p_n\}$.

These properties are parallel with the entanglement measure theory, where the average entanglement is not increased under the local operations and classical communication (LOCC). Notice that coherence measures that satisfy conditions (C3) and (C4) also satisfies condition (C2).

In [10] a class of incoherence operations was defined, called genuinely incoherent operations (GIO) as quantum operations that preserve all incoherent states.

2.4 Definition. An IO map Λ is called a **genuinely incoherent operation (GIO)** is for any incoherent state $\delta \in \mathcal{I}$,

$$\Lambda(\delta) = \delta .$$

An operation Λ is GIO if and only if all Kraus representations of Λ has all Kraus operators diagonal in a pre-fixed basis [10].

Conditions (C2), (C3) and (C4) can be restricted to GIO maps to obtain different classes of coherence measures.

2.5 Definition. In this case, a **measure of genuine coherence** satisfies at least (G1) and (G2). And if a coherence measure fulfills conditions (G1), (G2), (G3) it is called **genuine coherence monotone**.

A larger class of IO maps was defined in [37, 39].

2.6 Definition. An IO map Λ is called **strictly incoherent operations (SIO)** if its Kraus representation operator commute with dephasing, i.e. for $\Lambda(\rho) = \sum_j K_j \rho K_j^*$, we have for any j ,

$$K_j \Delta(\rho) K_j^* = \Delta(K_j \rho K_j^*) .$$

Since Kraus operators of GIO maps are diagonal in \mathcal{E} basis, any GIO map is SIO as well, i.e. $\text{GIO} \subset \text{SIO}$, [10].

One may consider an additional property, closely related to the entanglement theory:

- (C5) Uniqueness for pure states: for any pure state $|\psi\rangle$ coherence takes the form:

$$\mathcal{C}(\psi) = S(\Delta(\psi)) ,$$

where S is the von Neumann entropy and Δ is the dephasing operation defined as

$$\Delta(\rho) = \sum_j \langle j | \rho | j \rangle | j \rangle \langle j | .$$

2.2 Quasi-relative entropy

Quantum quasi-relative entropy was introduced by Petz [23, 24] as a quantum generalization of a classical Csiszár's f -divergence [8]. It is defined in the context of von Neumann algebras, but we consider only the Hilbert space setup. Let \mathcal{H} be a finite-dimensional Hilbert space, and ρ and σ be two states (given by density operators).

2.7 Definition. *For strictly positive bounded operators A and B acting on a finite-dimensional Hilbert space \mathcal{H} , and for any continuous function $f : (0, \infty) \rightarrow \mathbb{R}$, the quasi-relative entropy (or sometimes referred to as the f -divergence) is defined as*

$$S_f(A||B) = \text{Tr}(f(L_B R_A^{-1})A) ,$$

where left and right multiplication operators are defined as $L_B(X) = BX$ and $R_A(X) = XA$.

There is a straightforward way to calculate the quasi-relative entropy from the spectral decomposition of operators [16, 36]. Let A and B have the following spectral decomposition

$$A = \sum_j \lambda_j |\phi_j\rangle \langle \phi_j| , \quad B = \sum_k \mu_k |\psi_k\rangle \langle \psi_k| . \quad (2.1)$$

the set $\{|\phi_k\rangle \langle \psi_j|\}_{j,k}$ forms an orthonormal basis of $\mathcal{B}(\mathcal{H})$, the space of bounded linear operators, with respect to the Hilbert-Schmidt inner product defined as $\langle A, B \rangle = \text{Tr}(A^*B)$. By [36], the product of left and right multiplication operators can be written as

$$L_B R_A^{-1} = \sum_{j,k} \frac{\mu_k}{\lambda_j} P_{j,k} , \quad (2.2)$$

where $P_{j,k} : \mathcal{B}(\mathcal{H}) \rightarrow \mathcal{B}(\mathcal{H})$ is defined by

$$P_{j,k}(X) = |\psi_k\rangle \langle \phi_j| \langle \psi_k| X |\phi_j\rangle .$$

The quasi-relative entropy is calculated as follows

$$S_f(A||B) = \sum_{j,k} \lambda_j f\left(\frac{\mu_k}{\lambda_j}\right) |\langle \psi_k| \phi_j \rangle|^2 . \quad (2.3)$$

2.8 Theorem. ([23]) *For states, i.e. trace one positive density matrices ρ and σ , the quasi-relative entropy is bounded below by*

$$S_f(\rho||\sigma) \geq f(1).$$

The equality happens for a non-linear function f if and only if $\rho = \sigma$.

It is natural to require the quasi-relative entropy to be zero for equal state, and therefore we assume throughout the paper that $f(1) = 0$.

For an operator convex function, f , the quasi-relative entropy is jointly convex and monotone under CPTP maps [16]. The equality in monotonicity holds if and only if the map is reversible on these two states, i.e. for two states ρ and σ with $\text{supp } \rho \subset \text{supp } \sigma$, and a CPTP map Λ , the equality

$$S_f(\rho||\sigma) = S_f(\Lambda(\rho)||\Lambda(\sigma))$$

is satisfied if and only if

$$\mathcal{R}_\sigma(\Lambda(\rho)) = \rho ,$$

where \mathcal{R}_σ is the Petz's recovery map defined as

$$\mathcal{R}_\sigma(\omega) = \sigma^{1/2} \Lambda^* \left(\Lambda(\sigma)^{-1/2} \omega \Lambda(\sigma)^{-1/2} \right) \sigma^{1/2} . \quad (2.4)$$

2.9 Assumption. *Throughout the paper we will assume that the function f is operator convex and $f(1) = 0$.*

For any function f , its transpose \tilde{f} is defined as

$$\tilde{f}(x) = x f\left(\frac{1}{x}\right) , \quad x \in (0, \infty) .$$

The transpose \tilde{f} of an operator convex function f on $(0, \infty)$ is operator convex again, [16]. From (2.3) it follows that

$$S_{\tilde{f}}(\rho||\sigma) = S_f(\sigma||\rho) .$$

2.10 Example. *For $f(x) = -\log x$, the quasi-relative entropy becomes the Umegaki relative entropy*

$$S_{-\log}(\rho||\sigma) = S(\rho||\sigma) = \text{Tr}(\rho \log \rho - \rho \log \sigma) .$$

2.11 Example. For $p \in (-1, 1)$ and $p \neq 0$ let us take the function

$$f_p(x) := \frac{1}{p(1-p)}(1-x^p),$$

which is operator convex. The quasi-relative entropy for this function is calculated to be

$$S_{f_p}(\rho|\|\sigma) = \frac{1}{p(1-p)}(1 - \text{Tr}(\sigma^p \rho^{1-p})).$$

2.12 Example. For $p \in (-1, 1)$ take $q = 1-p \in (0, 2)$, the function

$$f_q(x) = \frac{1}{1-q}(1-x^{1-q})$$

is operator convex. The quasi-relative entropy for this function is known as Tsallis q -entropy

$$S_q(\rho|\|\sigma) = \frac{1}{1-q}(1 - \text{Tr}(\rho^q \sigma^{1-q})).$$

3 f -entropy

For a convex, operator monotone decreasing function f , such that $f(1) = 0$, define entropy two ways.

3.1 Definition. The f -entropy is defined as

$$S_f(\rho) := f(1/d) - S_f(\rho|I/d). \quad (3.1)$$

$$\hat{S}_f(\rho) := -S_f(\rho|I). \quad (3.2)$$

Let us use a notation \tilde{S}_f for either S_f or \hat{S}_f .

3.2 Theorem. f -entropy is non-negative, and is zero on pure states.

Proof. Let $\{\lambda_j\}$ be the eigenvalues of ρ . Then from (2.3) we have

$$S_f(\rho) = f(1/d) - \sum_j \lambda_j f\left(\frac{1}{d\lambda_j}\right), \quad (3.3)$$

and

$$\hat{S}_f(\rho) = - \sum_j \lambda_j f\left(\frac{1}{\lambda_j}\right). \quad (3.4)$$

A sequence of eigenvalues $\{\lambda_j\}$ is majorized by a sequence $\{1, 0, \dots, 0\}$. Since a perspective function (or

a transpose function) $xf(1/x)$ is convex for a convex function f [16], this implies that by results on Schur-concavity [14, 21, 29] we have

$$\sum_j \lambda_j f\left(\frac{1}{d\lambda_j}\right) \leq f(1/d).$$

Here, if needed, we adopt a convention $0 \cdot \pm\infty := 0$ [15].

Since f is monotonically decreasing and $f(1) = 0$, for any $0 \leq \lambda_j \leq 1$, $f\left(\frac{1}{\lambda_j}\right) \leq 0$. Thus, $\hat{S}_f \geq 0$.

When $\rho = |\Psi\rangle\langle\Psi|$ is a pure state, there is only one eigenvalue $\lambda = 1$. Then

$$S_f(|\Psi\rangle\langle\Psi|) = f(1/d) - f(1/d) = 0,$$

and

$$\hat{S}_f(|\Psi\rangle\langle\Psi|) = -f(1) = 0. \quad \square$$

3.3 Theorem. The maximum value of f -entropy is reached on the maximally mixed state I/d and it is

$$S_f(\rho) \leq f(1/d),$$

and

$$\hat{S}_f(\rho) \leq -f(d).$$

Proof. From Theorem 2.8, $S_f(\rho|I/d) \geq 0$, or since f is convex, we have

$$\sum_j \lambda_j f\left(\frac{1}{d\lambda_j}\right) \geq f\left(\sum_j \lambda_j \frac{1}{d\lambda_j}\right) = f(1) = 0.$$

Similarly,

$$\sum_j \lambda_j f\left(\frac{1}{\lambda_j}\right) \geq f\left(\sum_j \lambda_j \frac{1}{\lambda_j}\right) = f(d).$$

From (3.3) and (3.4), the result follows. Clearly, when $\rho = I/d$, we have $S_f(I/d) = f(1/d) - 0 = f(1/d)$, and from (3.4) we have $\hat{S}_f(I/d) = -f(d)$. \square

3.4 Theorem. The f -entropies are concave in ρ . Let $\{p_k\}$ be a probability distribution and ρ_k be some states, then for $\rho = \sum_k p_k \rho_k$, we have

$$\tilde{S}_f(\rho) \geq \sum_k p_k \tilde{S}_f(\rho_k).$$

Proof. This immediately follows from the joint convexity of f -divergence [15, 16]. \square

3.5 Theorem. *The f -entropies are invariant under unitaries.*

Proof. Since a unitary operation $U\rho U^*$ does not change the eigenvalues of ρ , and the f -entropies are the functions of eigenvalues of ρ , this implies that f -entropies are invariant under any operations that preserve eigenvalues. \square

3.6 Theorem. *The f -entropies are non-decreasing under unital CPTP maps, i.e. for any linear CPTP map Λ , such that $\Lambda(I) = I$, we have*

$$\tilde{S}_f(\Lambda(\rho)) \geq \tilde{S}_f(\rho) .$$

Proof. Let us denote $\sigma = I$ or $\sigma = I/d$, which corresponds to the appropriate f -entropy. Then

$$\tilde{S}_f(\Lambda(\rho)) - \tilde{S}_f(\rho) = S_f(\rho\|\sigma) - S_f(\Lambda(\rho)\|\sigma) \quad (3.5)$$

$$= S_f(\rho\|\sigma) - S_f(\Lambda(\rho)\|\Lambda(\sigma)) \geq 0 . \quad (3.6)$$

The last equality holds since Λ is unital, and the inequality holds due to the monotonicity of f -divergence under CPTP maps [18, 23, 35]. \square

4 Measure of genuine coherence

In a d -dimensional Hilbert space \mathcal{H} , fix a basis $\mathcal{E} = \{|j\rangle\}_{j=0}^{d-1}$.

4.1 Definition. *For any entropy function S , which is non-decreasing under CPTP maps, define coherence as follows:*

$$\mathcal{C}_S(\rho) := S(\Delta(\rho)) - S(\rho) . \quad (4.1)$$

In particular, for any operator convex and operator monotone decreasing function f , define two f -coherence measures.

4.2 Definition. *For entropy defined in (3.1),*

$$\mathcal{C}_f(\rho) := S_f(\Delta(\rho)) - S_f(\rho) . \quad (4.2)$$

For entropy defined in (3.2),

$$\hat{\mathcal{C}}_f(\rho) := \hat{S}_f(\Delta(\rho)) - \hat{S}_f(\rho) . \quad (4.3)$$

Let us denote $\tilde{\mathcal{C}}_f$ as either one \mathcal{C}_f or $\hat{\mathcal{C}}_f$ for shortness.

If $\{\lambda_j\}$ are the eigenvalues of ρ , and the diagonal elements of ρ in \mathcal{E} basis are $\chi_j = \langle j|\rho|j\rangle$, then from (3.3) and (3.4), we have

$$\mathcal{C}_f(\rho) = \sum_j \lambda_j f\left(\frac{1}{d\lambda_j}\right) - \sum_j \chi_j f\left(\frac{1}{d\chi_j}\right) , \quad (4.4)$$

and

$$\hat{\mathcal{C}}_f(\rho) = \sum_j \lambda_j f\left(\frac{1}{\lambda_j}\right) - \sum_j \chi_j f\left(\frac{1}{\chi_j}\right) . \quad (4.5)$$

4.1 Example

4.1.1 Log

Since $f(x) = -\log(x)$ is operator convex, coherence measure defined above coincides with [2]:

$$\mathcal{C}(\rho) = \hat{\mathcal{C}}_f(\rho) = S_{\log}(\Delta(\rho)) - S_{\log}(\rho) \quad (4.6)$$

$$= \sum_j \lambda_j \log \lambda_j - \sum_j \chi_j \log \chi_j \quad (4.7)$$

$$= S(\Delta(\rho)) - S(\rho) \quad (4.8)$$

$$= S(\rho\|\Delta(\rho)) \quad (4.9)$$

$$= \min_{\delta \in \mathcal{I}} S(\rho\|\delta) . \quad (4.10)$$

4.1.2 Power

The function $f(x) = \frac{1}{1-\alpha}(1-x^{1-\alpha})$ is operator convex for $\alpha \in (0, 2)$. The coherence monotone is then defined as

$$\mathcal{C}_\alpha(\rho) = \frac{d^{\alpha-1}}{1-\alpha} \left[\sum_j \chi_j^\alpha - \sum_j \lambda_j^\alpha \right] = d^{\alpha-1} \hat{\mathcal{C}}_\alpha(\rho) . \quad (4.11)$$

5 Properties

5.1 Uniqueness for pure states.

For any pure state coherence becomes an entropy of a dephased state:

$$\mathcal{C}_S(\psi) = S(\Delta(\psi)) .$$

This holds since entropies are zero on pure states.

5.2 Positivity

5.1 Theorem. \mathcal{C}_S and, in particular, $\tilde{\mathcal{C}}_f$ are non-negative.

Proof. By assumption S is non-decreasing under CPTP maps, it follows that \mathcal{C} is non-negative.

This holds for f -entropies as well due to Theorem 3.6, since the dephasing operation is unital. \square

Clearly, for any incoherent state ρ , coherence $\mathcal{C}_S(\rho) = 0$. Having no information on the saturation condition for a general entropy S , it is impossible to say what happens in the other direction. Consider f -coherences (4.2) and (4.3).

5.2 Theorem. $\tilde{\mathcal{C}}_f(\rho) = 0$ if and only if $\rho \in \mathcal{I}$ is incoherent state.

Proof. In Theorem 3.6, the equality in the only inequality (3.6) holds if and only if there is a recovery map \mathcal{R} such that $\mathcal{R}(\Delta(\rho)) = \rho$ and $\mathcal{R}(I) = I$, [15, 16]. By (2.4), this map admits the following explicit form: denoting $\sigma = I$

$$\mathcal{R}_\sigma(\omega) = \sigma^{1/2} \Delta^* \left(\Delta(\sigma)^{-1/2} \omega \Delta(\sigma)^{-1/2} \right) \sigma^{1/2},$$

where Δ^* is a dual map of Δ . Since Δ is a linear unital GIO map, we have

$$\mathcal{R}_\sigma(\omega) = \Delta^*(\omega). \quad (5.1)$$

Therefore, condition $\mathcal{R}_\sigma(\Delta(\rho)) = \rho$ implies that

$$\rho = \Delta^*(\Delta(\rho)). \quad (5.2)$$

Since $\Delta^* = \Delta$, we have that $\rho = \Delta(\rho)$, which happens if and only if $\rho \in \mathcal{I}$. Thus, $\mathcal{C}_f(\rho) = 0 = \hat{\mathcal{C}}_f(\rho)$ if and only if $\rho \in \mathcal{I}$. \square

A strengthening of the monotonicity inequality for f -divergence was presented in [4]. Using this result, we obtain the following lower bound on f -coherence.

5.3 Theorem. Let f be an operator monotone decreasing function, and $T > 0$. Suppose for some constant $c > 0$, there is a constant $C > 0$ so that $dt \leq CT^{2c} d\mu_f(t)$ for $t \in [T^{-1}, T]$. Then there is an explicitly computable constant $K_f(\rho)$ depending only on

the smallest non-zero eigenvalue of ρ , C and c , such that,

$$\mathcal{C}_f(\rho) \geq K_f(\rho) \|\rho - \Delta(\rho)\|_1^{4(1+c)}. \quad (5.3)$$

Here, $\|A\|_1 = \text{Tr}|A| = \text{Tr}\sqrt{A^*A}$ is the trace-norm of an operator.

From this inequality, the above condition of a zero coherence becomes apparent, i.e. $\mathcal{C}_f(\rho) = 0$ if and only if $\rho \in \mathcal{I}$.

The upper bound given below extends the upper bound for a relative entropy of coherence [2] to any f -coherence.

5.4 Theorem. The coherence is upper bounded by

$$\mathcal{C}_f(\rho) \leq f(1/d),$$

and

$$\hat{\mathcal{C}}_f(\rho) \leq -f(d).$$

The maximum value is reached for a maximally coherent pure state $\rho = |\psi\rangle\langle\psi|$, with $|\psi\rangle = \frac{1}{\sqrt{d}} \sum_j |j\rangle$.

Proof. This follows from the upper bound on the f -entropy Theorem 3.3, and the definition of coherence

$$\tilde{\mathcal{C}}_f(\rho) = \tilde{S}_f(\Delta(\rho)) - \tilde{S}_f(\rho).$$

For a pure state the entropy is zero, $\tilde{S}_f(|\psi\rangle\langle\psi|) = 0$. The dephasing operation applied to the state $|\psi\rangle = \frac{1}{\sqrt{d}} \sum_j |j\rangle$ gives a maximally mixed state I/d . The theorem follows from the fact that the entropy is maximal on maximally mixed state. \square

5.3 Monotonicity

5.5 Theorem. \mathcal{C}_S and, in particular, $\tilde{\mathcal{C}}_f$ is monotone under GIO.

Proof. Any GIO map Λ is also SIO, and, in particular, Λ commutes with the dephasing operation. Therefore, $\Delta(\Lambda(\rho)) = \Lambda(\Delta(\rho)) = \Delta(\rho)$, the last equality is due to the fact that $\Delta(\rho) \in \mathcal{I}$ and Λ as GIO preserves incoherent states. Therefore,

$$\mathcal{C}_S(\rho) - \mathcal{C}_S(\Lambda(\rho)) \quad (5.4)$$

$$= S(\Delta(\rho)) - S(\rho) - S(\Delta(\Lambda(\rho))) + S(\Lambda(\rho)) \quad (5.5)$$

$$= S(\Lambda(\rho)) - S(\rho) \quad (5.6)$$

$$\geq 0, \quad (5.7)$$

since Λ is a CPTP map and S is non-increasing under CPTP maps. For f -coherences, the last inequality holds to the Theorem 3.6 since a GIO map is unital. \square

5.6 Theorem. *For GIO map Λ , the equality*

$$\tilde{\mathcal{C}}_f(\rho) = \tilde{\mathcal{C}}_f(\Lambda(\rho))$$

happens if and only if any Kraus representation of $\Lambda(\rho) = \sum_j K_j \rho K_j^$ must have operators $K_j = \sum_n k_{jn} |n\rangle \langle n|$ that satisfy: for any n, m such that $\langle n | \rho | m \rangle \neq 0$, it must be that*

$$\left| \sum_j \overline{k_{jn}} k_{jm} \right|^2 = 1 .$$

Proof. Similarly, to the positivity section, equality in (5.7) happens if and only if there is a recovery map \mathcal{R} such that $\mathcal{R}(\Lambda(\rho)) = \rho$ and $\mathcal{R}(I) = I$, [15, 16]. By (2.4), this map admits the following explicit form: denoting $\sigma = I$

$$\mathcal{R}_\sigma(\omega) = \sigma^{1/2} \Lambda^* \left(\Lambda(\sigma)^{-1/2} \omega \Lambda(\sigma)^{-1/2} \right) \sigma^{1/2} ,$$

where Λ^* is a dual map of Λ . Since Λ is a linear unital GIO map, we have

$$\mathcal{R}_\sigma(\omega) = \Lambda^*(\omega) . \quad (5.8)$$

Therefore, condition $\mathcal{R}_\sigma(\Lambda(\rho)) = \rho$ implies that

$$\rho = \Lambda^*(\Lambda(\rho)) . \quad (5.9)$$

Denote a Kraus representation of Λ as $\Lambda(\rho) = \sum_j K_j \rho K_j^*$. From [10], since Λ is GIO, any Kraus representation of Λ has diagonal operators, i.e. each $K_j = \sum_n k_{jn} |n\rangle \langle n|$ is diagonal in basis \mathcal{E} . Since $\sum_j K_j^* K_j = I$, we have $\sum_j |k_{jn}|^2 = 1$ for every n . The dual map is $\Lambda^*(\rho) = \sum_j K_j^* \rho K_j$. Therefore, (5.9) becomes

$$\rho = \sum_{ji} K_j^* K_i \rho (K_j^* K_i)^* .$$

Writing both sides in basis \mathcal{E} gives

$$\sum_{nm} \langle n | \rho | m \rangle |n\rangle \langle m| \quad (5.10)$$

$$= \sum_{nm} \sum_{ij} \overline{k_{jn}} k_{in} k_{jm} \overline{k_{im}} \langle n | \rho | m \rangle |n\rangle \langle m| \quad (5.11)$$

$$= \sum_{nm} \left| \sum_j \overline{k_{jn}} k_{jm} \right|^2 \langle n | \rho | m \rangle |n\rangle \langle m| . \quad (5.12)$$

This implies that for every n, m such that $\langle n | \rho | m \rangle \neq 0$ we have

$$\left| \sum_j \overline{k_{jn}} k_{jm} \right|^2 = 1 . \quad (5.13)$$

This clearly confirms that any incoherent state saturates monotonicity for GIO maps.

If ρ is a coherent state, i.e. there exist n, m such that $\langle n | \rho | m \rangle \neq 0$, to saturate monotonicity the map Λ should satisfy (5.13). Note that by Cauchy-Schwarz inequality we have

$$\left| \sum_j \overline{k_{jn}} k_{jm} \right|^2 \leq \sum_j |k_{jn}|^2 \sum_j |k_{jm}|^2 = 1 .$$

The equality above happens if and only if there exists a scalar $\alpha_{nm} \in \mathbb{C}$ such that for any j : $k_{jn} = \alpha_{nm} k_{jm}$. \square

Applying the strengthening of monotonicity inequality for f -divergences [4], we obtain a strengthening on the monotonicity inequality for f -coherence.

5.7 Theorem. *Let Λ be any GIO map. Let f be an operator monotone decreasing function, and $T > 0$. Suppose for some constant $c > 0$, there is a constant $C > 0$ so that $dt \leq CT^{2c} d\mu_f(t)$ for $t \in [T^{-1}, T]$. Then there is an explicitly computable constant $K_f(\rho)$ depending only on the smallest non-zero eigenvalue of ρ , C and c , such that,*

$$\mathcal{C}_f(\rho) - \mathcal{C}_f(\Lambda(\rho)) \geq K_f(\rho) \|\rho - \Lambda^*(\Lambda(\rho))\|_1^{4(1+c)} . \quad (5.14)$$

Proof. Any GIO map Λ commutes with the dephasing operation, therefore, $\Delta(\Lambda(\rho)) = \Lambda(\Delta(\rho)) = \Delta(\rho)$, the

last equality is due to the fact that $\Delta(\rho) \in \mathcal{I}$ and Λ as GIO preserves incoherent states. Using this, we have

$$\mathcal{C}_f(\rho) - \mathcal{C}_f(\Lambda(\rho)) \quad (5.15)$$

$$= S_f(\Delta(\rho)) - S_f(\rho) - S_f(\Delta(\Lambda(\rho))) + S_f(\Lambda(\rho)) \quad (5.16)$$

$$= S_f(\Lambda(\rho)) - S_f(\rho) \quad (5.17)$$

$$= S_f(\rho \| I/d) - S_f(\Lambda(\rho) \| I/d) \quad (5.18)$$

$$= S_f(\rho \| \sigma) - S_f(\Lambda(\rho) \| \Lambda(\sigma)) , \quad (5.19)$$

where $\sigma = I/d$, and since $\sigma \in \mathcal{I}$ and Λ is GIO, $\Lambda(\sigma) = \sigma$.

Applying result in [4], which estimates the error in the monotonicity relation for f -divergence, leads to the desired abound.

□

The next theorem shows that $\tilde{\mathcal{C}}_f$ is not in general monotone under SIO operations.

5.8 Theorem. *If $\tilde{\mathcal{C}}_f$ is monotone under all SIO, then for all states ρ and $|0\rangle \in \mathcal{E}$, we have*

$$\tilde{\mathcal{C}}_f(\rho \otimes I/d) = \tilde{\mathcal{C}}_f(\rho \otimes |0\rangle \langle 0|) .$$

In other words, if \mathcal{C}_f is monotone under SIO, then for all states with eigenvalues $\{\lambda_j\}$ and diagonal elements $\{\chi_j\}$ in the basis \mathcal{E} , the following holds

$$\begin{aligned} & \sum_j \lambda_j f\left(\frac{1}{d\lambda_j}\right) - \sum_j \chi_j f\left(\frac{1}{d\chi_j}\right) \\ &= \sum_j \lambda_j f\left(\frac{1}{d^2\lambda_j}\right) - \sum_j \chi_j f\left(\frac{1}{d^2\chi_j}\right) . \end{aligned} \quad (5.20)$$

And, if $\hat{\mathcal{C}}_f$ is monotone under SIO, then for all states with eigenvalues $\{\lambda_j\}$ and diagonal elements $\{\chi_j\}$ in the basis \mathcal{E} , the following holds

$$\sum_j \lambda_j f\left(\frac{1}{\lambda_j}\right) - \sum_j \chi_j f\left(\frac{1}{\chi_j}\right) \quad (5.21)$$

$$= \sum_j \lambda_j f\left(\frac{d}{\lambda_j}\right) - \sum_j \chi_j f\left(\frac{d}{\chi_j}\right) . \quad (5.22)$$

Proof. First, note that from (4.4) we have: for $|0\rangle \in \mathcal{E}$,

$$\mathcal{C}_f(\rho \otimes |0\rangle \langle 0|) = \sum_j \lambda_j f\left(\frac{1}{d^2\lambda_j}\right) - \sum_j \chi_j f\left(\frac{1}{d^2\chi_j}\right) , \quad (5.23)$$

and

$$\mathcal{C}_f(\rho \otimes I/d) = \mathcal{C}_f(\rho) = \sum_j \lambda_j f\left(\frac{1}{d\lambda_j}\right) - \sum_j \chi_j f\left(\frac{1}{d\chi_j}\right) . \quad (5.24)$$

Moreover,

$$\hat{\mathcal{C}}_f(\rho \otimes |0\rangle \langle 0|) = \hat{\mathcal{C}}_f(\rho) = \sum_j \lambda_j f\left(\frac{1}{\lambda_j}\right) - \sum_j \chi_j f\left(\frac{1}{\chi_j}\right) , \quad (5.25)$$

and

$$\hat{\mathcal{C}}_f(\rho \otimes I/d) = \sum_j \lambda_j f\left(\frac{d}{\lambda_j}\right) - \sum_j \chi_j f\left(\frac{d}{\chi_j}\right) . \quad (5.26)$$

Let us consider two examples of SIO\GIO maps.

1. Let $\Phi(\rho) = I/d$ be the depolarizing quantum channel, which in Kraus form can be written as

$$\Phi(\rho) = I/d = \sum_{ij=0}^{d-1} K_{ij} \rho K_{ij}^* ,$$

where $K_{ij} = \frac{1}{\sqrt{d}} |i\rangle \langle j|$.

Define an operation on a tensor product Hilbert space as follows

$$\Lambda(\omega) = \sum_{ij} (I \otimes K_{ij}) \omega (I \otimes K_{ij})^* . \quad (5.27)$$

Clearly, Λ is not a GIO, since its Kraus operators are not diagonal in $\mathcal{E} \otimes \mathcal{E}$ basis, or since

$$\Lambda(\rho \otimes |0\rangle \langle 0|) = \rho \otimes \Phi(|0\rangle \langle 0|) = \rho \otimes I/d \quad (5.28)$$

$$\neq \rho \otimes |0\rangle \langle 0| \in \mathcal{E} \otimes \mathcal{E} . \quad (5.29)$$

But Λ is SIO, since for any n, m

$$(I \otimes K_{nm})(\Delta(\omega))(I \otimes K_{nm}^*) \quad (5.30)$$

$$= \frac{1}{d} (I \otimes |n\rangle \langle m|) \left(\sum_{ij} \langle ij | \omega | ij \rangle |ij\rangle \langle ij| \right) (I \otimes |m\rangle \langle n|) \quad (5.31)$$

$$= \frac{1}{d} \sum_{ij} \langle ij | \omega | ij \rangle |i\rangle \langle i| \otimes |n\rangle \langle m| |j\rangle \langle j| |m\rangle \langle n| \quad (5.32)$$

$$= \frac{1}{d} \sum_i \langle im | \omega | im \rangle |in\rangle \langle in| , \quad (5.33)$$

and

$$\Delta((I \otimes K_{nm})\omega(I \otimes K_{nm}^*)) \quad (5.34)$$

$$= \frac{1}{d} \sum_{ij} \langle ij | (I \otimes |n\rangle \langle m|) \omega(I \otimes |m\rangle \langle n|) |ij\rangle |ij\rangle \langle ij| \quad (5.35)$$

$$= \frac{1}{d} \sum_i \langle im | \omega |im\rangle |in\rangle \langle in| \quad (5.36)$$

$$(5.37)$$

Therefore, Λ is a SIO map.

For either \mathcal{C}_f or $\hat{\mathcal{C}}_f$, consider

$$\tilde{\mathcal{C}}_f(\Lambda(\rho \otimes |0\rangle \langle 0|)) = \tilde{\mathcal{C}}_f(\rho \otimes \Phi(|0\rangle \langle 0|)) \quad (5.38)$$

$$= \tilde{\mathcal{C}}_f(\rho \otimes I/d) . \quad (5.39)$$

2. Consider another example, let $\Psi(\rho) = |0\rangle \langle 0|$ be the erasure channel, which in Kraus form can be written as

$$\Psi(\rho) = |0\rangle \langle 0| = \sum_{j=0}^{d-1} K_j \rho K_j^*, \quad \text{where } K_j = |0\rangle \langle j| .$$

Define an operation on a tensor product Hilbert space as follows

$$M(\omega) = \sum_j (I \otimes K_j) \omega (I \otimes K_j)^* . \quad (5.40)$$

Clearly, M is not a GIO, since its Kraus operators are not diagonal in $\mathcal{E} \otimes \mathcal{E}$ basis, or since

$$M(\rho \otimes I/d) = \rho \otimes \Psi(I/d) = \rho \otimes |0\rangle \langle 0| \quad (5.41)$$

$$\neq \rho \otimes I/d \in \mathcal{E} \otimes \mathcal{E} . \quad (5.42)$$

But M is SIO, since for any n ,

$$(I \otimes K_n) \Delta(\omega) (I \otimes K_n^*) \quad (5.43)$$

$$= \sum_{ij} \langle ij | \omega |ij\rangle (I \otimes |0\rangle \langle n|) |ij\rangle \langle ij| (I \otimes |n\rangle \langle 0|) \quad (5.44)$$

$$= \sum_i \langle in | \omega |in\rangle |i\rangle \langle i| \otimes |0\rangle \langle 0| . \quad (5.45)$$

and

$$\Delta((I \otimes K_n)\omega(I \otimes K_n^*)) \quad (5.46)$$

$$= \sum_{ij} \langle ij | (I \otimes |0\rangle \langle n|) \omega(I \otimes |n\rangle \langle 0|) |ij\rangle |ij\rangle \langle ij| \quad (5.47)$$

$$= \sum_i \langle in | \omega |in\rangle |i0\rangle \langle i0| . \quad (5.48)$$

Therefore, M is an SIO map.

For either \mathcal{C}_f or $\hat{\mathcal{C}}_f$, consider

$$\tilde{\mathcal{C}}_f(M(\rho \otimes I/d)) = \tilde{\mathcal{C}}_f(\rho \otimes \Psi(I/d)) \quad (5.49)$$

$$= \tilde{\mathcal{C}}_f(\rho \otimes |0\rangle \langle 0|) . \quad (5.50)$$

Now, compare (5.39) and (5.50). In order for monotonicity of f -coherence to hold under all SIO, there must be an equality

$$\tilde{\mathcal{C}}_f(\rho \otimes I/d) = \tilde{\mathcal{C}}_f(\rho \otimes |0\rangle \langle 0|) .$$

Invoking (5.23-5.26) we have the result stated in the theorem. \square

Note that both (5.20) and (5.22) hold for the logarithmic function $f(x) = -\log(x)$, but fail for the power function $f(x) = \frac{1}{1-\alpha}(1-x^{1-\alpha})$. This is in line with the fact that the relative entropy of coherence is monotone under SIO, and it shows that Tsallis coherence fails monotonicity for SIO.

5.4 Strong monotonicity

5.9 Theorem. *f -coherences $\tilde{\mathcal{C}}_f$ saturate strong monotonicity for convex mixtures of diagonal unitaries. Therefore, $\tilde{\mathcal{C}}_f$ saturates strong monotonicity under GIO in two- and three-dimensions.*

Proof. Consider an example of GIO, which is a probabilistic mixture of diagonal unitaries: for some $\alpha_j > 0$, s.t. $\sum_j \alpha_j = 1$, define

$$\Lambda(\rho) = \sum_j \alpha_j U_j \rho U_j^* ,$$

where for some ρ_{jn} the unitaries U_j are diagonal in \mathcal{E} , i.e.

$$U_j = \sum_n e^{i\phi_{jn}} |n\rangle \langle n| .$$

In [10] it has been shown that all GIO are of such form for dimensions two and three, but it is no longer the case for higher dimensions.

Note that for $\sigma = I$ or $\sigma = I/d$ and for all unitaries U , we have

$$S_f(U\rho U^*|\sigma) = S_f(\rho|\sigma) . \quad (5.51)$$

Taking U_j diagonal in \mathcal{E} above, it follows that

$$\Delta(U_j\rho U_j^*) = \Delta(\rho) .$$

Therefore, $\tilde{\mathcal{C}}_f$ saturates the strong monotonicity under convex mixtures of diagonal unitaries:

$$\sum_j \alpha_j \tilde{\mathcal{C}}_f(U_j\rho U_j^*) \quad (5.52)$$

$$= \sum_j \alpha_j [S_f(U_j\rho U_j^*|\sigma) - S_f(\Delta(U\rho U^*)|\sigma)] \quad (5.53)$$

$$= \sum_j \alpha_j [S_f(\rho|\sigma) - S_f(\Delta(\rho)|\sigma)] \quad (5.54)$$

$$= \tilde{\mathcal{C}}_f(\rho) . \quad (5.55)$$

□

5.10 Remark. Expanding the set of operations to include all unitaries (not necessarily diagonal in \mathcal{E}), forces $\tilde{\mathcal{C}}_f$ to be invariant under all unitaries if it is monotone under them. This results from the following observation: if $\tilde{\mathcal{C}}_f$ is monotone under all unitaries U and all states ρ , then, since (5.51) holds, it must be that

$$S_f(\Delta(U\rho U^*)|\sigma) \geq S_f(\Delta(\rho)|\sigma) .$$

But taking a unitary $V = U^*$ and an initial state $\omega = U\rho U^*$ above, results in the opposite inequality:

$$S_f(\Delta(V\omega V^*)|\sigma) = S_f(\Delta(\rho)|\sigma) \quad (5.56)$$

$$\geq S_f(\Delta(U\rho U^*)|\sigma) = S_f(\Delta(\omega)|\sigma) . \quad (5.57)$$

Therefore, the above inequality must be equality, which makes $\tilde{\mathcal{C}}_f$ invariant under unitaries.

5.11 Theorem. *For any pure state ρ , the f -coherences are strongly monotone under GIO maps in any finite dimension.*

Proof. Let us denote $\sigma = I$ or $\sigma = I/d$ depending on the f -coherence we are considering. For a GIO map Λ with Kraus operators K_j , denote

$$p_j = \text{Tr} K_j \rho K_j^*, \quad \rho_j = \frac{1}{p_j} K_j \rho K_j^* .$$

For a pure state ρ , states ρ_j are also pure. Therefore,

$$\tilde{\mathcal{C}}_f(\rho) - \sum_j p_j \tilde{\mathcal{C}}_f(\rho_j) \quad (5.58)$$

$$= \sum_j p_j S_f(\Delta(\rho_j)|\sigma) - S_f(\Delta(\rho)|\sigma) . \quad (5.59)$$

Since any GIO map is an SIO map as well, it follows that

$$\Delta(\rho_j) = \frac{1}{p_j} K_j \Delta(\rho) K_j^* .$$

Dephased state $\Delta(\rho)$ is diagonal in \mathcal{E} basis with eigenvalues χ_j , i.e. $\Delta(\rho) = \sum_j \chi_j |j\rangle \langle j|$. The f -divergence is

$$S_f(\Delta(\rho)|I) = \sum_n \chi_n f\left(\frac{1}{\chi_n}\right) .$$

Kraus operators of GIO map are diagonal in \mathcal{E} basis, $K_j = \sum_n k_{jn} |n\rangle \langle n|$, with $\sum_j |k_{jn}|^2 = 1$ for all j . Then

$$K_j \Delta(\rho) K_j^* = \sum_n \chi_n |k_{jn}|^2 |n\rangle \langle n| .$$

And

$$\sum_j p_j S_f(\Delta(\rho_j)|I) = \sum_{jn} \chi_n |k_{jn}|^2 f\left(\frac{p_j}{\chi_n |k_{jn}|^2}\right) .$$

Since f is convex, we have for every n :

$$\sum_j |k_{jn}|^2 f\left(\frac{p_j}{\chi_n |k_{jn}|^2}\right) \geq f\left(\sum_j \frac{p_j}{\chi_n}\right) \quad (5.60)$$

$$= f\left(\frac{1}{\chi_n}\right) . \quad (5.61)$$

Similarly,

$$S_f(\Delta(\rho)|I/d) = \sum_n \chi_n f\left(\frac{1}{d\chi_n}\right) ,$$

and

$$\sum_j p_j S_f(\Delta(\rho_j)|I/d) = \sum_{jn} \chi_n |k_{jn}|^2 f\left(\frac{p_j}{d\chi_n |k_{jn}|^2}\right) .$$

Because f is convex, for any n :

$$\sum_j |k_{jn}|^2 f\left(\frac{p_j}{d\chi_n |k_{jn}|^2}\right) \geq f\left(\sum_j \frac{p_j}{d\chi_n}\right) \quad (5.62)$$

$$= f\left(\frac{1}{d\chi_n}\right). \quad (5.63)$$

And thus, $\sum_j p_j S_f(\Delta(\rho_j)||\sigma) \geq S_f(\Delta(\rho)||\sigma)$. Which implies that for any pure state ρ , the f -coherence is strongly monotone under GIO:

$$\tilde{\mathcal{C}}_f(\rho) \geq \sum_j p_j \tilde{\mathcal{C}}_f(\rho_j).$$

□

Data sharing not applicable to this article as no datasets were generated or analysed during the current study.

Acknowledgments. A. V. is supported by NSF grants DMS-1812734 and DMS-2105583.

References

- [1] Aberg, J. (2014). Catalytic coherence. *Phys. Rev. Lett.* 113, 150402
- [2] Baumgratz, T., Cramer, M., Plenio, M. B. (2014). Quantifying coherence. *Physical review letters*, 113(14), 140401
- [3] Bromley, T. R., Cianciaruso, M., Adesso, G. (2015). Frozen quantum coherence. *Physical review letters*, 114(21), 210401.
- [4] Carlen, E. A., Vershynina, A. (2018). Recovery and the data processing inequality for quasi-entropies. *IEEE Transactions on Information Theory*, 64(10), 6929-6938.
- [5] Chitambar, E., Gour, G. (2016). Critical examination of incoherent operations and a physically consistent resource theory of quantum coherence. *Physical review letters*, 117(3), 030401
- [6] Chitambar, E., Gour, G. (2016). Comparison of incoherent operations and measures of coherence. *Physical Review A*, 94(5), 052336.
- [7] Chitambar, E., Hsieh, M. H. (2016). Relating the resource theories of entanglement and quantum coherence. *Physical review letters*, 117(2), 020402
- [8] Csiszár, I. (1967). Information-type measures of difference of probability distributions and indirect observation. *studia scientiarum Mathematicarum Hungarica*, 2, 229-318.
- [9] Cwikliski, P., Studziski, M., Horodecki, M., Oppenheim, J. (2015). Limitations on the evolution of quantum coherences: towards fully quantum second laws of thermodynamics. *Physical review letters*, 115(21), 210403
- [10] De Vicente, J. I., Streltsov, A. (2016). Genuine quantum coherence. *Journal of Physics A: Mathematical and Theoretical*, 50(4), 045301.
- [11] Dey, P. K., Chakraborty, D., Char, P., Chattopadhyay, I., Sarkar, D. (2019). Structure of POVM based Resource Theory of Coherence. *arXiv preprint arXiv:1908.01882*.
- [12] Du, S., Bai, Z., Guo, Y. (2015). Conditions for coherence transformations under incoherent operations. *Physical Review A*, 91(5), 052120
- [13] Glauber, R. J. (1963). Coherent and incoherent states of the radiation field. *Physical Review*, 131(6), 2766
- [14] Hardy, G. H., J. E. Littlewood, and G. Pólya (1929). Some simple inequalities satisfied by convex function. *Messenger Math.* 58, 145? 152.
- [15] Hiai, F., Mosonyi, M., Petz, D., Bény, C. (2011). Quantum f -divergences and error correction. *Reviews in Mathematical Physics*, 23(07), 691-747
- [16] Hiai, F., Mosonyi, M. (2017). Different quantum f -divergences and the reversibility of quantum operations. *Reviews in Mathematical Physics*, 29(07), 1750023.

- [17] Hu, M. L., Hu, X., Wang, J., Peng, Y., Zhang, Y. R., Fan, H. (2018). Quantum coherence and geometric quantum discord. *Physics Reports*, 762, 1-100
- [18] Lesniewski, A., Ruskai, M. B. (1999). Monotone Riemannian metrics and relative entropy on non-commutative probability spaces. *Journal of Mathematical Physics*, 40(11), 5702-5724.
- [19] Lostaglio, M., Jennings, D. Rudolph, T. Description of quantum coherence in thermodynamic processes requires constraints beyond free energy. *Nat. Commun.* 6, 6383 (2015)
- [20] Ma, J., Yadin, B., Girolami, D., Vedral, V., Gu, M. (2016). Converting coherence to quantum correlations. *Physical review letters*, 116(16), 160407
- [21] Marshall, A. W., Olkin, I., Arnold, B. C. (2010). Doubly Stochastic Matrices. In *Inequalities: Theory of Majorization and Its Applications* (pp. 29-77). Springer, New York, NY.
- [22] Piani, M., Cianciaruso, M., Bromley, T. R., Napoli, C., Johnston, N., Adesso, G. (2016). Robustness of asymmetry and coherence of quantum states. *Physical Review A*, 93(4), 042107
- [23] Petz, D. (1985). Quasi-entropies for states of a von Neumann algebra. *Publications of the Research Institute for Mathematical Sciences*, 21(4), 787-800.
- [24] Petz, D. (1986). Quasi-entropies for finite quantum systems. *Reports on mathematical physics*, 23(1), 57-65.
- [25] Radhakrishnan, C., Parthasarathy, M., Jambulingam, S., Byrnes, T. (2016). Distribution of quantum coherence in multipartite systems. *Physical review letters*, 116(15), 150504
- [26] Rana, S., Parashar, P., Lewenstein, M. (2016). Trace-distance measure of coherence. *Physical Review A*, 93(1), 012110
- [27] Rastegin, A. E. (2016). Quantum-coherence quantifiers based on the Tsallis relative α entropies. *Physical Review A*, 93(3), 032136
- [28] Rebentrost, P., Mohseni, M., Aspuru-Guzik, A. (2009). Role of quantum coherence and environmental fluctuations in chromophoric energy transport. *J. Phys. Chem. B* 113, 9942
- [29] Schur, I. (1923). *Über eine Klasse von Mittelbildungen mit Anwendungen auf die Determinantentheorie*. *Sitzungsberichte der Berliner Mathematischen Gesellschaft*, 22(9-20), 51.
- [30] Scully, M. O., Zubairy, M. S. (1997). *Quantum Optics* (Cambridge.) Ch, 4, 17
- [31] Shao, L. H., Xi, Z., Fan, H., Li, Y. (2015). Fidelity and trace-norm distances for quantifying coherence. *Physical Review A*, 91(4), 042120
- [32] Streltsov, A., Adesso, G., Plenio, M. B. (2017). Colloquium: Quantum coherence as a resource. *Reviews of Modern Physics*, 89(4), 041003.
- [33] Streltsov, A., Singh, U., Dhar, H. S., Bera, M. N., Adesso, G. (2015). Measuring quantum coherence with entanglement. *Physical review letters*, 115(2), 020403
- [34] Tan, K. C., Kwon, H., Park, C. Y., Jeong, H. (2016). Unified view of quantum correlations and quantum coherence. *Physical Review A*, 94(2), 022329
- [35] Tomamichel, M., Colbeck, R., Renner, R. (2009). A fully quantum asymptotic equipartition property. *IEEE Transactions on information theory*, 55(12), 5840-5847.
- [36] Virosztek, D. (2016). Quantum entropies, relative entropies, and related preserver problems.
- [37] Winter, A., Yang, D. (2016). Operational resource theory of coherence. *Physical review letters*, 116(12), 120404

- [38] Witt, B., Mintert, F. (2013). Stationary quantum coherence and transport in disordered networks. *New J. Phys.* 15, 093020
- [39] Yadin, B., Ma, J., Girolami, D., Gu, M., Vedral, V. (2016). Quantum processes which do not use coherence. *Physical Review X*, 6(4), 041028.
- [40] Yu, X. D., Zhang, D. J., Xu, G. F., Tong, D. M. (2016). Alternative framework for quantifying coherence. *Physical Review A*, 94(6), 060302.
- [41] Zhao, H., Yu, C. S. (2018). Coherence measure in terms of the Tsallis relative α entropy. *Scientific reports*, 8(1), 299
- [42] Zhu, H., Ma, Z., Cao, Z., Fei, S. M., Vedral, V. (2017). Operational one-to-one mapping between coherence and entanglement measures. *Physical Review A*, 96(3), 032316