

# A Primal-Dual Method of Partial Inverses for Composite Inclusions\*

Maryam A. Alghamdi,<sup>‡</sup> Abdullah Alotaibi,<sup>‡</sup> Patrick L. Combettes,<sup>‡,‡</sup> and Naseer Shahzad <sup>‡</sup>

<sup>‡</sup>King Abdulaziz University  
Department of Mathematics  
Sciences Faculty for Girls, P. O. Box 4087  
Jeddah 21491, Saudi Arabia

<sup>‡</sup>King Abdulaziz University  
Department of Mathematics, P. O. Box 80203  
Jeddah 21859, Saudi Arabia

<sup>‡</sup>UPMC Université Paris 06  
Laboratoire Jacques-Louis Lions – UMR CNRS 7598  
75005 Paris, France

## Abstract

Spingarn's method of partial inverses has found many applications in nonlinear analysis and in optimization. We show that it can be employed to solve composite monotone inclusions in duality, thus opening a new range of applications for the partial inverse formalism. The versatility of the resulting primal-dual splitting algorithm is illustrated through applications to structured monotone inclusions and optimization.

**Keywords** convex optimization, duality, method of partial inverses, monotone operator, splitting algorithm

**Mathematics Subject Classifications (2010)** Primary 47H05; Secondary 65K05, 90C25.

---

\*Contact author: P. L. Combettes, [plc@math.jussieu.fr](mailto:plc@math.jussieu.fr), phone: +33 1 4427 6319, fax: +33 1 4427 7200.

# 1 Introduction

Let  $\mathcal{H}$  be a real Hilbert space, let  $A: \mathcal{H} \rightarrow 2^{\mathcal{H}}$  be a set-valued operator, let  $\text{gra } A = \{(x, u) \in \mathcal{H} \times \mathcal{H} \mid u \in Ax\}$  denote the graph of  $A$ , let  $V$  be a closed vector subspace of  $\mathcal{H}$ , and let  $P_V$  and  $P_{V^\perp}$  denote respectively the projectors onto  $V$  and onto its orthogonal complement. The partial inverse of  $A$  with respect to  $V$  is defined through

$$\text{gra } A_V = \{(P_V x + P_{V^\perp} u, P_V u + P_{V^\perp} x) \mid (x, u) \in \text{gra } A\}. \quad (1.1)$$

This operator, which was introduced by Spingarn in [20], can be regarded as an intermediate object between  $A$  and  $A^{-1}$ . A key result of [20] is that, if  $A$  is maximally monotone, problems of the form

$$\text{find } x \in V \text{ and } u \in V^\perp \text{ such that } u \in Ax \quad (1.2)$$

can be solved by applying the proximal point algorithm [19] to the partial inverse  $A_V$ . The resulting algorithm, known as the *method of partial inverses*, has been applied to various problems in nonlinear analysis and optimization; see, e.g., [3, 6, 8, 10, 12, 13, 14, 16, 20, 21, 22]. The goal of the present paper is to propose a new range of applications of the method of partial inverses by showing that it can be applied to solving the following type of monotone inclusions in duality.

**Problem 1.1** Let  $\mathcal{H}$  and  $\mathcal{G}$  be real Hilbert spaces, let  $A: \mathcal{H} \rightarrow 2^{\mathcal{H}}$  and  $B: \mathcal{G} \rightarrow 2^{\mathcal{G}}$  be maximally monotone operators, and let  $L: \mathcal{H} \rightarrow \mathcal{G}$  be a bounded linear operator. Solve the primal inclusion

$$\text{find } x \in \mathcal{H} \text{ such that } 0 \in Ax + L^* B L x \quad (1.3)$$

together with the dual inclusion

$$\text{find } v \in \mathcal{G} \text{ such that } 0 \in -L A^{-1}(-L^* v) + B^{-1} v. \quad (1.4)$$

The operator duality described in (1.3)–(1.4) is an extension of the classical Fenchel-Rockafellar duality setting for functions [18] which has been studied in particular in [1, 5, 11, 15, 17]. Our main result shows that, through a suitable reformulation, Problem 1.1 can be reduced to a problem of the form (1.2) and that the method of partial inverses applied to the latter leads to a splitting algorithm in which the operators  $A$ ,  $B$ , and  $L$  are used separately.

The remainder of the paper is organized as follows. In Section 2, we revisit the method of partial inverses and propose new convergence results. Our method of partial inverses for solving Problem 1.1 is presented in Section 3, together with applications. An alternative implementation of this method is proposed in Section 4, where further applications are provided. Section 5 contains concluding remarks.

**Notation.** The scalar product of a real Hilbert space  $\mathcal{H}$  is denoted by  $\langle \cdot \mid \cdot \rangle$  and the associated norm by  $\|\cdot\|$ ;  $\rightharpoonup$  and  $\rightarrow$  denote, respectively, weak and strong convergence, and  $\text{Id}$  is the identity operator. We denote by  $\mathcal{B}(\mathcal{H}, \mathcal{G})$  the class of bounded linear operators from  $\mathcal{H}$  to a real Hilbert space  $\mathcal{G}$ . Let  $A: \mathcal{H} \rightarrow 2^{\mathcal{H}}$ . The inverse  $A^{-1}$  of  $A$  is defined via  $\text{gra } A^{-1} = \{(u, x) \in \mathcal{H} \times \mathcal{H} \mid (x, u) \in \text{gra } A\}$ ,  $J_A = (\text{Id} + A)^{-1}$  is the resolvent of  $A$ ,  $\text{ran } A = \bigcup_{x \in \mathcal{H}} Ax$  is the range of  $A$ , and  $\text{zer } A = \{x \in \mathcal{H} \mid 0 \in Ax\}$  is the set of zeros of  $A$ . We denote by  $\Gamma_0(\mathcal{H})$  the class of lower semicontinuous convex functions  $f: \mathcal{H} \rightarrow ]-\infty, +\infty]$  such that  $\text{dom } f = \{x \in \mathcal{H} \mid f(x) < +\infty\} \neq \emptyset$ . Let  $f \in \Gamma_0(\mathcal{H})$ . The conjugate of  $f$  is the function  $f^* \in \Gamma_0(\mathcal{H})$  defined by  $f^*: u \mapsto \sup_{x \in \mathcal{H}} (\langle x \mid u \rangle - f(x))$ . For every  $x \in \mathcal{H}$ ,  $f + \|x - \cdot\|^2/2$  possesses a unique minimizer, which is denoted by  $\text{prox}_f x$ . We have

$$\text{prox}_f = J_{\partial f}, \quad \text{where } \partial f: \mathcal{H} \rightarrow 2^{\mathcal{H}}: x \mapsto \{u \in \mathcal{H} \mid (\forall y \in \mathcal{H}) \langle y - x \mid u \rangle + f(x) \leq f(y)\} \quad (1.5)$$

is the subdifferential of  $f$ . See [2] for background on monotone operators and convex analysis.

## 2 A method of partial inverses

Throughout this section,  $\mathcal{K}$  denotes a real Hilbert space. We establish the convergence of a relaxed, error-tolerant method of partial inverses. First, we review some results about partial inverses and the proximal point algorithm.

**Lemma 2.1** [20, Section 2] *Let  $A: \mathcal{K} \rightarrow 2^{\mathcal{K}}$  be maximally monotone, let  $V$  be a closed vector subspace of  $\mathcal{K}$ , and let  $z \in \mathcal{K}$ . Then  $A_V$  is maximally monotone and  $z \in \text{zer } A_V \Leftrightarrow (P_V z, P_{V^\perp} z) \in \text{gra } A$ .*

**Lemma 2.2** *Let  $A: \mathcal{K} \rightarrow 2^{\mathcal{K}}$  be maximally monotone, let  $V$  be a closed vector subspace of  $\mathcal{K}$ , let  $z \in \mathcal{K}$ , and let  $p \in \mathcal{K}$ . Then  $p = J_{A_V} z \Leftrightarrow P_V p + P_{V^\perp}(z - p) = J_A z$ .*

*Proof.* This result, which appears implicitly in [20, Section 4], follows from the equivalences

$$\begin{aligned} P_V p + P_{V^\perp}(z - p) = J_A z &\Leftrightarrow (P_V p + P_{V^\perp}(z - p), z - P_V p - P_{V^\perp}(z - p)) \in \text{gra } A \\ &\Leftrightarrow (P_V p + P_{V^\perp}(z - p), P_V(z - p) + P_{V^\perp} p) \in \text{gra } A \\ &\Leftrightarrow (p, z - p) \in \text{gra } A_V \\ &\Leftrightarrow p = J_{A_V} z, \end{aligned} \tag{2.1}$$

where we have used (1.1).  $\square$

**Lemma 2.3** [8, Remark 2.2(vi)] *Let  $B: \mathcal{K} \rightarrow 2^{\mathcal{K}}$  be a maximally monotone operator such that  $\text{zer } B \neq \emptyset$ , let  $z_0 \in \mathcal{K}$ , let  $(c_n)_{n \in \mathbb{N}}$  be a sequence in  $\mathcal{K}$ , and let  $(\lambda_n)_{n \in \mathbb{N}}$  be a sequence in  $]0, 2[$ . Suppose that  $\sum_{n \in \mathbb{N}} \lambda_n(2 - \lambda_n) = +\infty$  and  $\sum_{n \in \mathbb{N}} \lambda_n \|c_n\| < +\infty$ , and set*

$$(\forall n \in \mathbb{N}) \quad z_{n+1} = z_n + \lambda_n (J_B z_n + c_n - z_n). \tag{2.2}$$

*Then  $J_B z_n - z_n \rightarrow 0$  and  $(z_n)_{n \in \mathbb{N}}$  converges weakly to a point in  $\text{zer } B$ .*

The next theorem analyzes a method of partial inverses and extends the results of [10] and [20].

**Theorem 2.4** *Let  $A: \mathcal{K} \rightarrow 2^{\mathcal{K}}$  be a maximally monotone operator, let  $V$  be a closed vector subspace of  $\mathcal{K}$ , let  $(\lambda_n)_{n \in \mathbb{N}}$  be a sequence in  $]0, 2[$ , and let  $(e_n)_{n \in \mathbb{N}}$  be a sequence in  $\mathcal{K}$  such that  $\sum_{n \in \mathbb{N}} \lambda_n(2 - \lambda_n) = +\infty$  and  $\sum_{n \in \mathbb{N}} \lambda_n \|e_n\| < +\infty$ . Suppose that the problem*

$$\text{find } x \in V \text{ and } u \in V^\perp \text{ such that } u \in Ax \tag{2.3}$$

*has at least one solution, let  $x_0 \in V$ , let  $u_0 \in V^\perp$ , and set*

$$(\forall n \in \mathbb{N}) \quad \begin{cases} p_n = J_A(x_n + u_n) + e_n \\ r_n = x_n + u_n - p_n \\ x_{n+1} = x_n - \lambda_n P_V r_n \\ u_{n+1} = u_n - \lambda_n P_{V^\perp} p_n. \end{cases} \tag{2.4}$$

*Then the following hold:*

- (i)  $P_V(p_n - e_n) - x_n \rightarrow 0$  and  $P_{V^\perp}(r_n + e_n) - u_n \rightarrow 0$ .
- (ii) *There exists a solution  $(\bar{x}, \bar{u})$  to (2.3) such that  $x_n \rightharpoonup \bar{x}$  and  $u_n \rightharpoonup \bar{u}$ .*

*Proof.* Set

$$(\forall n \in \mathbb{N}) \quad \mathbf{z}_n = \mathbf{x}_n + \mathbf{u}_n \quad \text{and} \quad \mathbf{c}_n = P_V \mathbf{e}_n - P_{V^\perp} \mathbf{e}_n. \quad (2.5)$$

Then  $\sum_{n \in \mathbb{N}} \lambda_n \|\mathbf{c}_n\| \leq \sum_{n \in \mathbb{N}} \lambda_n (\|P_V \mathbf{e}_n\| + \|P_{V^\perp} \mathbf{e}_n\|) \leq 2 \sum_{n \in \mathbb{N}} \lambda_n \|\mathbf{e}_n\| < +\infty$ . Furthermore, since  $(\mathbf{x}_n)_{n \in \mathbb{N}}$  lies in  $V$  and  $(\mathbf{u}_n)_{n \in \mathbb{N}}$  lies in  $V^\perp$ , (2.4) can be rewritten as

$$(\forall n \in \mathbb{N}) \quad \begin{cases} \mathbf{p}_n = J_A(\mathbf{x}_n + \mathbf{u}_n) + \mathbf{e}_n \\ \mathbf{r}_n = \mathbf{x}_n + \mathbf{u}_n - \mathbf{p}_n \\ \mathbf{x}_{n+1} = \mathbf{x}_n + \lambda_n (P_V \mathbf{p}_n - \mathbf{x}_n) \\ \mathbf{u}_{n+1} = \mathbf{u}_n + \lambda_n (P_{V^\perp} \mathbf{r}_n - \mathbf{u}_n), \end{cases} \quad (2.6)$$

which yields

$$\begin{aligned} (\forall n \in \mathbb{N}) \quad & P_V \left( \frac{\mathbf{z}_{n+1} - \mathbf{z}_n}{\lambda_n} + \mathbf{z}_n - \mathbf{c}_n \right) + P_{V^\perp} \left( \mathbf{z}_n - \left( \frac{\mathbf{z}_{n+1} - \mathbf{z}_n}{\lambda_n} + \mathbf{z}_n - \mathbf{c}_n \right) \right) \\ &= P_V \left( \frac{\mathbf{z}_{n+1} - \mathbf{z}_n}{\lambda_n} + \mathbf{z}_n - \mathbf{c}_n \right) + P_{V^\perp} \left( \frac{\mathbf{z}_n - \mathbf{z}_{n+1}}{\lambda_n} + \mathbf{c}_n \right) \\ &= P_V \left( \frac{\mathbf{x}_{n+1} - \mathbf{x}_n}{\lambda_n} + \mathbf{x}_n \right) + P_{V^\perp} \left( \frac{\mathbf{u}_n - \mathbf{u}_{n+1}}{\lambda_n} \right) - \mathbf{e}_n \\ &= P_V \mathbf{p}_n + P_{V^\perp} (\mathbf{u}_n - \mathbf{r}_n) - \mathbf{e}_n \\ &= P_V \mathbf{p}_n + P_{V^\perp} (\mathbf{p}_n - \mathbf{x}_n) - \mathbf{e}_n \\ &= \mathbf{p}_n - \mathbf{e}_n \\ &= J_A \mathbf{z}_n. \end{aligned} \quad (2.7)$$

Hence, it follows from (2.5), (2.6), and Lemma 2.2 that

$$(\forall n \in \mathbb{N}) \quad \mathbf{z}_{n+1} = \mathbf{z}_n + \lambda_n (J_{A_V} \mathbf{z}_n + \mathbf{c}_n - \mathbf{z}_n). \quad (2.8)$$

Altogether, we derive from Lemmas 2.1 and 2.3 that

$$J_{A_V} \mathbf{z}_n - \mathbf{z}_n \rightarrow \mathbf{0} \quad (2.9)$$

and that there exists  $\mathbf{z} \in \text{zer } A_V$  such that

$$\mathbf{z}_n \rightharpoonup \mathbf{z}. \quad (2.10)$$

(i): In view of (2.4), (2.5), Lemma 2.2, and (2.9), we have

$$\mathbf{x}_n - P_V (\mathbf{p}_n - \mathbf{e}_n) = \mathbf{x}_n - P_V J_A \mathbf{z}_n = \mathbf{x}_n - P_V J_{A_V} \mathbf{z}_n = P_V (\mathbf{z}_n - J_{A_V} \mathbf{z}_n) \rightarrow \mathbf{0} \quad (2.11)$$

and

$$\mathbf{u}_n - P_{V^\perp} (\mathbf{r}_n + \mathbf{e}_n) = P_{V^\perp} \mathbf{z}_n - P_{V^\perp} (\mathbf{z}_n - J_A \mathbf{z}_n) = P_{V^\perp} J_A \mathbf{z}_n = P_{V^\perp} (\mathbf{z}_n - J_{A_V} \mathbf{z}_n) \rightarrow \mathbf{0}. \quad (2.12)$$

(ii): Since  $\mathbf{z} \in \text{zer } (A_V)$ , it follows from Lemma 2.1 that  $(\bar{\mathbf{x}}, \bar{\mathbf{u}}) = (P_V \mathbf{z}, P_{V^\perp} \mathbf{z})$  solves (2.3). Furthermore, using (2.5), (2.10), and the weak continuity of  $P_V$  and  $P_{V^\perp}$ , we get  $\mathbf{x}_n = P_V \mathbf{z}_n \rightharpoonup P_V \mathbf{z} = \bar{\mathbf{x}}$  and  $\mathbf{u}_n = P_{V^\perp} \mathbf{z}_n \rightharpoonup P_{V^\perp} \mathbf{z} = \bar{\mathbf{u}}$ .  $\square$

**Remark 2.5** Theorem 2.4(ii) was obtained in [10, Section 5] under the additional assumptions that  $\sum_{n \in \mathbb{N}} \|\mathbf{e}_n\| < +\infty$ ,  $\inf_{n \in \mathbb{N}} \lambda_n > 0$ , and  $\sup_{n \in \mathbb{N}} \lambda_n < 2$ ; the original weak convergence result of [20] corresponds to the case when  $(\forall n \in \mathbb{N}) \mathbf{e}_n = \mathbf{0}$  and  $\lambda_n = 1$ . As noted in [8, Remark 2.2(iii)], our assumptions do not require that  $\sum_{n \in \mathbb{N}} \|\mathbf{e}_n\| < +\infty$  and they even allow for situations in which  $(\mathbf{e}_n)_{n \in \mathbb{N}}$  does not converge weakly to 0.

**Remark 2.6** It follows from Lemma 2.1 that any algorithm that constructs a zero of  $A_V$  can be used to solve (2.3). We have chosen to employ the proximal point algorithm of Lemma 2.3 because it features a flexible error model and it allows for under- and over-relaxations which can prove very useful in improving the convergence pattern of the algorithm. In addition, (2.8) leads to the simple implementation (2.4) in terms of the resolvent  $J_A$ . An alternative proximal point method is

$$(\forall n \in \mathbb{N}) \quad z_{n+1} = z_n + \lambda_n (J_{\gamma_n A_V} z_n + c_n - z_n), \quad (2.13)$$

where  $(\gamma_n)_{n \in \mathbb{N}}$  lies  $]0, +\infty[$  (weak convergence conditions under various hypotheses on  $(\lambda_n)_{n \in \mathbb{N}}$ ,  $(\gamma_n)_{n \in \mathbb{N}}$ , and  $(e_n)_{n \in \mathbb{N}}$  exist; see for instance [7, Corollary 4.5] and [10, Theorem 3]). However, due to the presence of the parameters  $(\gamma_n)_{n \in \mathbb{N}}$ , (2.13) results in an algorithm which is much less straightforward to execute than (2.4). This issue is also discussed in [10, 14, 20].

### 3 Primal-dual composite method of partial inverses

The following technical facts will be needed subsequently.

**Lemma 3.1** *Let  $\mathcal{H}$  and  $\mathcal{G}$  be real Hilbert spaces, and let  $L \in \mathcal{B}(\mathcal{H}, \mathcal{G})$ . Set  $\mathcal{K} = \mathcal{H} \oplus \mathcal{G}$  and  $V = \{(x, y) \in \mathcal{K} \mid Lx = y\}$ . Then, for every  $(x, y) \in \mathcal{K}$ , the following hold:*

- (i)  $P_V(x, y) = (x - L^*(\text{Id} + LL^*)^{-1}(Lx - y), y + (\text{Id} + LL^*)^{-1}(Lx - y))$ .
- (ii)  $P_V(x, y) = ((\text{Id} + L^*L)^{-1}(x + L^*y), L(\text{Id} + L^*L)^{-1}(x + L^*y))$ .
- (iii)  $P_{V^\perp}(x, y) = (L^*(\text{Id} + LL^*)^{-1}(Lx - y), -(\text{Id} + LL^*)^{-1}(Lx - y))$ .
- (iv)  $P_{V^\perp}(x, y) = (x - (\text{Id} + L^*L)^{-1}(x + L^*y), y - L(\text{Id} + L^*L)^{-1}(x + L^*y))$ .

*Proof.* Set  $T: \mathcal{K} \rightarrow \mathcal{G}: (x, y) \mapsto Lx - y$ . Then  $T \in \mathcal{B}(\mathcal{K}, \mathcal{G})$ ,  $T^*: \mathcal{G} \rightarrow \mathcal{K}: v \mapsto (L^*v, -v)$ ,  $TT^* = (\text{Id} + LL^*)$  is invertible, and  $V = \ker T$  is a closed vector subspace of  $\mathcal{K}$ . Let us fix  $(x, y) \in \mathcal{K}$ .

(i): Since  $V = \ker T$ , it follows from [2, Example 28.14(iii)] that

$$\begin{aligned} P_V(x, y) &= (x, y) - T^*(TT^*)^{-1}T(x, y) \\ &= (x - L^*(\text{Id} + LL^*)^{-1}(Lx - y), y + (\text{Id} + LL^*)^{-1}(Lx - y)). \end{aligned} \quad (3.1)$$

(i) $\Rightarrow$ (ii): We have

$$\begin{aligned} &(\text{Id} + L^*L)(x - L^*(\text{Id} + LL^*)^{-1}(Lx - y) - (\text{Id} + L^*L)^{-1}(x + L^*y)) \\ &= x + L^*Lx - L^*(\text{Id} + LL^*)^{-1}(Lx - y) - L^*LL^*(\text{Id} + LL^*)^{-1}(Lx - y) - x - L^*y \\ &= L^*(Lx - y) - L^*(\text{Id} + LL^*)^{-1}(Lx - y) - L^*LL^*(\text{Id} + LL^*)^{-1}(Lx - y) \\ &= L^*(\text{Id} + LL^* - \text{Id} - LL^*)(\text{Id} + LL^*)^{-1}(Lx - y) \\ &= 0. \end{aligned} \quad (3.2)$$

Therefore,  $x - L^*(\text{Id} + LL^*)^{-1}(Lx - y) = (\text{Id} + L^*L)^{-1}(x + L^*y)$ . Likewise,

$$\begin{aligned} &(\text{Id} + LL^*)(y + (\text{Id} + LL^*)^{-1}(Lx - y) - L(\text{Id} + L^*L)^{-1}(x + L^*y)) \\ &= y + LL^*y + Lx - y - L(\text{Id} + L^*L)^{-1}(x + L^*y) - LL^*L(\text{Id} + L^*L)^{-1}(x + L^*y) \\ &= L(x + L^*y) - L(\text{Id} + L^*L)^{-1}(x + L^*y) - LL^*L(\text{Id} + L^*L)^{-1}(x + L^*y) \\ &= (L(\text{Id} + L^*L) - L - LL^*L)(\text{Id} + L^*L)^{-1}(x + L^*y) \\ &= 0 \end{aligned} \quad (3.3)$$

and hence  $y + (\text{Id} + LL^*)^{-1}(Lx - y) = L(\text{Id} + L^*L)^{-1}(x + L^*y)$ .

$$(i) \Rightarrow (iii): P_{V^\perp}(x, y) = (x, y) - P_V(x, y) = (L^*(\text{Id} + LL^*)^{-1}(Lx - y), -(\text{Id} + LL^*)^{-1}(Lx - y)).$$

$$(ii) \Rightarrow (iv): P_{V^\perp}(x, y) = (x, y) - P_V(x, y) = (x - (\text{Id} + L^*L)^{-1}(x + L^*y), y - L(\text{Id} + L^*L)^{-1}(x + L^*y)).$$

□

Our main algorithm is introduced and analyzed in the next theorem. It consists of applying (2.4) to the operator  $A: (x, y) \mapsto Ax \times By$  and the subspace  $V = \{(x, y) \in \mathcal{H} \oplus \mathcal{G} \mid Lx = y\}$ .

**Theorem 3.2** *In Problem 1.1, set  $Q = (\text{Id} + L^*L)^{-1}$  and assume that  $\text{zer}(A + L^*BL) \neq \emptyset$ . Let  $(\lambda_n)_{n \in \mathbb{N}}$  be a sequence in  $]0, 2[$ , let  $(a_n)_{n \in \mathbb{N}}$  be a sequence in  $\mathcal{H}$ , and let  $(b_n)_{n \in \mathbb{N}}$  be a sequence in  $\mathcal{G}$  such that  $\sum_{n \in \mathbb{N}} \lambda_n(2 - \lambda_n) = +\infty$  and  $\sum_{n \in \mathbb{N}} \lambda_n \sqrt{\|a_n\|^2 + \|b_n\|^2} < +\infty$ . Let  $x_0 \in \mathcal{H}$  and  $v_0 \in \mathcal{G}$ , and set  $y_0 = Lx_0$ ,  $u_0 = -L^*v_0$ , and*

$$(\forall n \in \mathbb{N}) \quad \begin{cases} p_n = J_A(x_n + u_n) + a_n \\ q_n = J_B(y_n + v_n) + b_n \\ r_n = x_n + u_n - p_n \\ s_n = y_n + v_n - q_n \\ t_n = Q(r_n + L^*s_n) \\ w_n = Q(p_n + L^*q_n) \\ x_{n+1} = x_n - \lambda_n t_n \\ y_{n+1} = y_n - \lambda_n L t_n \\ u_{n+1} = u_n + \lambda_n(w_n - p_n) \\ v_{n+1} = v_n + \lambda_n(Lw_n - q_n). \end{cases} \quad (3.4)$$

Then the following hold:

$$(i) \quad x_n - w_n + Q(a_n + L^*b_n) \rightarrow 0 \quad \text{and} \quad y_n - Lw_n + LQ(a_n + L^*b_n) \rightarrow 0.$$

$$(ii) \quad u_n - r_n + t_n - a_n + Q(a_n + L^*b_n) \rightarrow 0 \quad \text{and} \quad v_n - s_n + Lt_n - b_n + LQ(a_n + L^*b_n) \rightarrow 0.$$

Moreover, there exists a solution  $\bar{x}$  to (1.3) and a solution  $\bar{v}$  to (1.4) such that the following hold:

$$(iii) \quad -L^*\bar{v} \in A\bar{x} \quad \text{and} \quad \bar{v} \in BL\bar{x}.$$

$$(iv) \quad x_n \rightharpoonup \bar{x} \quad \text{and} \quad v_n \rightharpoonup \bar{v}.$$

*Proof.* Set

$$\mathcal{K} = \mathcal{H} \oplus \mathcal{G} \quad \text{and} \quad V = \{(x, y) \in \mathcal{K} \mid Lx = y\} \quad (3.5)$$

and note that

$$V^\perp = \{(u, v) \in \mathcal{K} \mid u = -L^*v\}. \quad (3.6)$$

In addition, set

$$Z = \{(x, v) \in \mathcal{K} \mid -L^*v \in Ax \quad \text{and} \quad v \in BLx\} \quad (3.7)$$

and

$$A: \mathcal{K} \rightarrow 2^{\mathcal{K}}: (x, y) \mapsto Ax \times By. \quad (3.8)$$

We also introduce the set

$$\mathcal{S} = \{(x, u) \in V \times V^\perp \mid u \in Ax\}. \quad (3.9)$$

Observe that

$$\mathcal{S} = \{((x, Lx), (-L^*v, v)) \in \mathcal{K} \times \mathcal{K} \mid (x, v) \in \mathcal{Z}\}. \quad (3.10)$$

Thus (see [5, 15] for the first two equivalences),

$$\text{zer}(A + L^*BL) \neq \emptyset \Leftrightarrow \text{zer}(-LA^{-1}(-L^*) + B^{-1}) \neq \emptyset \Leftrightarrow \mathcal{Z} \neq \emptyset \Leftrightarrow \mathcal{S} \neq \emptyset. \quad (3.11)$$

Now define  $(\forall n \in \mathbb{N})$   $e_n = (a_n, b_n)$ ,  $p_n = (p_n, q_n)$ ,  $r_n = (r_n, s_n)$ ,  $u_n = (u_n, v_n)$ , and  $x_n = (x_n, y_n)$ . Then  $x_0 \in V$  and  $u_0 \in V^\perp$ . Moreover, by [2, Proposition 23.16],  $A$  is maximally monotone and

$$(\forall n \in \mathbb{N}) \quad J_A(x_n + u_n) = (J_A(x_n + u_n), J_B(y_n + v_n)). \quad (3.12)$$

Furthermore, it follows from (3.5) and Lemma 3.1(ii) that

$$(\forall n \in \mathbb{N}) \quad P_V r_n = (Q(r_n + L^*s_n), LQ(r_n + L^*s_n)), \quad (3.13)$$

and from Lemma 3.1(iv) that

$$(\forall n \in \mathbb{N}) \quad P_{V^\perp} p_n = (p_n - Q(p_n + L^*q_n), q_n - LQ(p_n + L^*q_n)). \quad (3.14)$$

Thus, we derive from (3.12), (3.13), and (3.14) that (2.4) yields (3.4). Altogether, since  $\sum_{n \in \mathbb{N}} \lambda_n \|e_n\| = \sum_{n \in \mathbb{N}} \lambda_n \sqrt{\|a_n\|^2 + \|b_n\|^2} < +\infty$ , Theorem 2.4(i) and Lemma 3.1 imply that (i) and (ii) are satisfied, and Theorem 2.4(ii) implies that there exists  $(\bar{x}, \bar{u}) \in \mathcal{S}$  such that  $x_n \rightharpoonup \bar{x}$  and  $u_n \rightharpoonup \bar{u}$ . Therefore, by (3.10), there exists  $(\bar{x}, \bar{v}) \in \mathcal{Z}$  such that  $(x_n, v_n) \rightharpoonup (\bar{x}, \bar{v})$ . Since  $\mathcal{Z} \subset (\text{zer}(A + L^*BL)) \times (\text{zer}(-LA^{-1}(-L^*) + B^{-1}))$  [5, Proposition 2.8(i)], the proof is complete.  $\square$

**Remark 3.3** In the special case when  $A = 0$  and  $L$  has closed range, an algorithm was proposed in [16] to solve the primal problem (1.3), i.e., to find a point in  $\text{zer}(L^*BL)$ . It employs the method of partial inverses in  $\mathcal{G}$  for finding  $y \in V$  and  $v \in V^\perp$  such that  $v \in By$ , where  $V = \text{ran } L$ , and then solves  $Lx = y$ . Each iteration of the resulting algorithm requires the computation of the generalized inverse of  $L$ , which is numerically demanding.

The following application of Theorem 3.2 concerns multi-operator inclusions.

**Problem 3.4** Let  $m$  be a strictly positive integer, and let  $\mathcal{H}$  and  $(\mathcal{G}_i)_{1 \leq i \leq m}$  be real Hilbert spaces. Let  $z \in \mathcal{H}$ , let  $C: \mathcal{H} \rightarrow 2^{\mathcal{H}}$  be maximally monotone, and, for every  $i \in \{1, \dots, m\}$ , let  $B_i: \mathcal{G}_i \rightarrow 2^{\mathcal{G}_i}$  be maximally monotone, let  $o_i \in \mathcal{G}_i$ , and let  $L_i \in \mathcal{B}(\mathcal{H}, \mathcal{G}_i)$ . Solve

$$\text{find } \bar{x} \in \mathcal{H} \text{ such that } z \in C\bar{x} + \sum_{i=1}^m L_i^* B_i(L_i \bar{x} - o_i) \quad (3.15)$$

together with the dual problem

$$\text{find } \bar{v}_1 \in \mathcal{G}_1, \dots, \bar{v}_m \in \mathcal{G}_m \text{ such that} \\ (\forall i \in \{1, \dots, m\}) \quad -o_i \in -L_i C^{-1} \left( z - \sum_{j=1}^m L_j^* \bar{v}_j \right) + B_i^{-1} \bar{v}_i. \quad (3.16)$$

**Corollary 3.5** In Problem 3.4, set  $Q = (\text{Id} + \sum_{i=1}^m L_i^* L_i)^{-1}$  and assume that  $z \in \text{ran}(C + \sum_{i=1}^m L_i^* B_i(L_i \cdot - o_i))$ . Let  $(\lambda_n)_{n \in \mathbb{N}}$  be a sequence in  $]0, 2[$  such that  $\sum_{n \in \mathbb{N}} \lambda_n(2 - \lambda_n) = +\infty$ , let  $(a_n)_{n \in \mathbb{N}}$  be a sequence in  $\mathcal{H}$ , and let  $x_0 \in \mathcal{H}$ . For every  $i \in \{1, \dots, m\}$ , let  $(b_{i,n})_{n \in \mathbb{N}}$  be a sequence in  $\mathcal{G}_i$ , let  $v_{i,0} \in \mathcal{G}_i$ , and set  $y_{i,0} = L_i x_0$ . Suppose that  $\sum_{n \in \mathbb{N}} \lambda_n \sqrt{\|a_n\|^2 + \sum_{i=1}^m \|b_{i,n}\|^2} < +\infty$ , and set  $u_0 = -\sum_{i=1}^m L_i^* v_{i,0}$  and

$$(\forall n \in \mathbb{N}) \quad \begin{cases} p_n = J_C(x_n + u_n + z) + a_n \\ r_n = x_n + u_n - p_n \\ \text{For } i = 1, \dots, m \\ \quad \begin{cases} q_{i,n} = o_i + J_{B_i}(y_{i,n} + v_{i,n} - o_i) + b_{i,n} \\ s_{i,n} = y_{i,n} + v_{i,n} - q_{i,n} \\ t_n = Q(r_n + \sum_{i=1}^m L_i^* s_{i,n}) \\ w_n = Q(p_n + \sum_{i=1}^m L_i^* q_{i,n}) \\ x_{n+1} = x_n - \lambda_n t_n \\ u_{n+1} = u_n + \lambda_n(w_n - p_n) \end{cases} \\ \text{For } i = 1, \dots, m \\ \quad \begin{cases} y_{i,n+1} = y_{i,n} - \lambda_n L_i t_n \\ v_{i,n+1} = v_{i,n} + \lambda_n(L_i w_n - q_{i,n}). \end{cases} \end{cases} \quad (3.17)$$

Then there exists a solution  $\bar{x}$  to (3.15) and a solution  $(\bar{v}_i)_{1 \leq i \leq m}$  to (3.16) such that the following hold:

- (i)  $z - \sum_{i=1}^m L_i^* \bar{v}_i \in C\bar{x}$  and  $(\forall i \in \{1, \dots, m\}) \bar{v}_i \in B_i(L_i \bar{x} - o_i)$ .
- (ii)  $x_n \rightharpoonup \bar{x}$  and  $(\forall i \in \{1, \dots, m\}) v_{i,n} \rightharpoonup \bar{v}_i$ .

*Proof.* Set  $A: \mathcal{H} \rightarrow 2^{\mathcal{H}}: x \mapsto -z + Cx$ ,  $\mathcal{G} = \bigoplus_{i=1}^m \mathcal{G}_i$ ,  $L: \mathcal{H} \rightarrow \mathcal{G}: x \mapsto (L_i x)_{1 \leq i \leq m}$ , and  $B: \mathcal{G} \rightarrow 2^{\mathcal{G}}: (y_i)_{1 \leq i \leq m} \mapsto \times_{i=1}^m B_i(y_i - o_i)$ . Then  $L^*: \mathcal{G} \rightarrow \mathcal{H}: (y_i)_{1 \leq i \leq m} \mapsto \sum_{i=1}^m L_i^* y_i$  and Problem 3.4 is therefore an instantiation of Problem 1.1. Moreover, [2, Propositions 23.15 and 23.16] yield

$$J_A: x \mapsto J_C(x + z) \quad \text{and} \quad J_B: (y_i)_{1 \leq i \leq m} \mapsto (o_i + J_{B_i}(y_i - o_i))_{1 \leq i \leq m}. \quad (3.18)$$

Now set  $(\forall n \in \mathbb{N}) b_n = (b_{i,n})_{1 \leq i \leq m}$ ,  $q_n = (q_{i,n})_{1 \leq i \leq m}$ ,  $s_n = (s_{i,n})_{1 \leq i \leq m}$ ,  $v_n = (v_{i,n})_{1 \leq i \leq m}$ , and  $y_n = (y_{i,n})_{1 \leq i \leq m}$ . In this setting, (3.4) coincides with (3.17) and the claims therefore follow from Theorem 3.2(iii)&(iv).  $\square$

The next application addresses a primal-dual structured minimization problem.

**Problem 3.6** Let  $m$  be a strictly positive integer, and let  $\mathcal{H}$  and  $(\mathcal{G}_i)_{1 \leq i \leq m}$  be real Hilbert spaces. Let  $z \in \mathcal{H}$ , let  $f \in \Gamma_0(\mathcal{H})$ , and, for every  $i \in \{1, \dots, m\}$ , let  $g_i \in \Gamma_0(\mathcal{G}_i)$ ,  $o_i \in \mathcal{G}_i$ , and  $L_i \in \mathcal{B}(\mathcal{H}, \mathcal{G}_i)$ . Solve the primal problem

$$\underset{x \in \mathcal{H}}{\text{minimize}} \quad f(x) + \sum_{i=1}^m g_i(L_i x - o_i) - \langle x \mid z \rangle \quad (3.19)$$

together with the dual problem

$$\underset{v_1 \in \mathcal{G}_1, \dots, v_m \in \mathcal{G}_m}{\text{minimize}} \quad f^*\left(z - \sum_{i=1}^m L_i^* v_i\right) + \sum_{i=1}^m (g_i^*(v_i) + \langle v_i \mid o_i \rangle). \quad (3.20)$$

**Corollary 3.7** In Problem 3.6, set  $Q = (\text{Id} + \sum_{i=1}^m L_i^* L_i)^{-1}$  and assume that

$$z \in \text{ran}\left(\partial f + \sum_{i=1}^m L_i^*(\partial g_i)(L_i \cdot - o_i)\right), \quad (3.21)$$



Let  $(\lambda_n)_{n \in \mathbb{N}}$  be a sequence in  $]0, 2[$  such that  $\sum_{n \in \mathbb{N}} \lambda_n(2 - \lambda_n) = +\infty$ , let  $(a_n)_{n \in \mathbb{N}}$  be a sequence in  $\mathcal{H}$ , and let  $x_0 \in \mathcal{H}$ . For every  $i \in \{1, \dots, m\}$ , let  $(b_{i,n})_{n \in \mathbb{N}}$  be a sequence in  $\mathcal{G}_i$ , let  $v_{i,0} \in \mathcal{G}_i$ , and set  $y_{i,0} = L_i x_0$ . Suppose that  $\sum_{n \in \mathbb{N}} \lambda_n \sqrt{\|a_n\|^2 + \sum_{i=1}^m \|b_{i,n}\|^2} < +\infty$ , and set  $u_0 = -\sum_{i=1}^m L_i^* v_{i,0}$  and

$$(\forall n \in \mathbb{N}) \quad \left\{ \begin{array}{l} p_n = \text{prox}_f(x_n + u_n + z) + a_n \\ r_n = x_n + u_n - p_n \\ \text{For } i = 1, \dots, m \\ \quad \left\{ \begin{array}{l} q_{i,n} = o_i + \text{prox}_{g_i}(y_{i,n} + v_{i,n} - o_i) + b_{i,n} \\ s_{i,n} = y_{i,n} + v_{i,n} - q_{i,n} \\ t_n = Q(r_n + \sum_{i=1}^m L_i^* s_{i,n}) \\ w_n = Q(p_n + \sum_{i=1}^m L_i^* q_{i,n}) \\ x_{n+1} = x_n - \lambda_n t_n \\ u_{n+1} = u_n + \lambda_n(w_n - p_n) \\ \text{For } i = 1, \dots, m \\ \quad \left\{ \begin{array}{l} y_{i,n+1} = y_{i,n} - \lambda_n L_i t_n \\ v_{i,n+1} = v_{i,n} + \lambda_n(L_i w_n - q_{i,n}). \end{array} \right. \end{array} \right. \end{array} \right. \quad (3.22)$$

Then there exists a solution  $\bar{x}$  to (3.19) and a solution  $(\bar{v}_i)_{1 \leq i \leq m}$  to (3.20) such that  $z - \sum_{i=1}^m L_i^* \bar{v}_i \in \partial f(\bar{x})$ ,  $x_n \rightharpoonup \bar{x}$ , and  $(\forall i \in \{1, \dots, m\}) v_{i,n} \rightharpoonup \bar{v}_i \in \partial g_i(L_i \bar{x} - o_i)$ .

*Proof.* Set  $C = \partial f$  and  $(\forall i \in \{1, \dots, m\}) B_i = \partial g_i$ . Then, using the same type of argument as in the proof of [9, Theorem 4.2], we derive from (3.21) that Problem 3.4 reduces to Problem 3.6 and that (3.17) reduces to (3.22). Thus, the assertions follow from Corollary 3.5.  $\square$

## 4 Alternative composite primal-dual method of partial inverses

The partial inverse method (3.4) relies on the implicit assumption that the operator  $(\text{Id} + L^* L)^{-1}$  is relatively easy to implement. In some instances, it may be advantageous to work with  $(\text{Id} + L L^*)^{-1}$  instead. In this section we describe an alternative method tailored to such situations.

**Theorem 4.1** *In Problem 1.1, set  $R = (\text{Id} + L L^*)^{-1}$  and assume that  $\text{zer}(A + L^* B L) \neq \emptyset$ . Let  $(\lambda_n)_{n \in \mathbb{N}}$  be a sequence in  $]0, 2[$ , let  $(a_n)_{n \in \mathbb{N}}$  be a sequence in  $\mathcal{H}$ , and let  $(b_n)_{n \in \mathbb{N}}$  be a sequence in  $\mathcal{G}$  such that  $\sum_{n \in \mathbb{N}} \lambda_n(2 - \lambda_n) = +\infty$  and  $\sum_{n \in \mathbb{N}} \lambda_n \sqrt{\|a_n\|^2 + \|b_n\|^2} < +\infty$ . Let  $x_0 \in \mathcal{H}$  and  $v_0 \in \mathcal{G}$ , and set  $y_0 = L x_0$ ,  $u_0 = -L^* v_0$ , and*

$$(\forall n \in \mathbb{N}) \quad \left\{ \begin{array}{l} p_n = J_A(x_n + u_n) + a_n \\ q_n = J_B(y_n + v_n) + b_n \\ r_n = x_n + u_n - p_n \\ s_n = y_n + v_n - q_n \\ t_n = R(L r_n - s_n) \\ w_n = R(L p_n - q_n) \\ x_{n+1} = x_n + \lambda_n(L^* t_n - r_n) \\ y_{n+1} = y_n - \lambda_n(t_n + s_n) \\ u_{n+1} = u_n - \lambda_n L^* w_n \\ v_{n+1} = v_n + \lambda_n w_n. \end{array} \right. \quad (4.1)$$

Then the conclusions of Theorem 3.2 are true.

*Proof.* The proof is analogous to that of Theorem 3.2 except that we replace (3.13) by

$$(\forall n \in \mathbb{N}) \quad P_{\mathbf{V}} \mathbf{r}_n = (r_n - L^* R(Lr_n - s_n), s_n + R(Lr_n - s_n)), \quad (4.2)$$

and (3.14) by

$$(\forall n \in \mathbb{N}) \quad P_{\mathbf{V}^\perp} \mathbf{p}_n = (L^* R(Lp_n - q_n), -R(Lp_n - q_n)) \quad (4.3)$$

by invoking Lemma 3.1(i)&(iii).  $\square$

Next, we present an application to a coupled inclusions problem. This problem has essentially the same structure as Problem 3.4, except that primal and dual inclusions are interchanged.

**Problem 4.2** Let  $m$  be a strictly positive integer, and let  $(\mathcal{H}_i)_{1 \leq i \leq m}$  and  $\mathcal{G}$  be real Hilbert spaces. Let  $o \in \mathcal{G}$ , let  $D: \mathcal{G} \rightarrow 2^{\mathcal{G}}$  be maximally monotone, and, for every  $i \in \{1, \dots, m\}$ , let  $z_i \in \mathcal{H}_i$ , let  $A_i: \mathcal{H}_i \rightarrow 2^{\mathcal{H}_i}$  be maximally monotone, and let  $L_i \in \mathcal{B}(\mathcal{H}_i, \mathcal{G})$ . Solve the primal problem

$$\text{find } \bar{x}_1 \in \mathcal{H}_1, \dots, \bar{x}_m \in \mathcal{H}_m \text{ such that } (\forall i \in \{1, \dots, m\}) \quad z_i \in A_i \bar{x}_i + L_i^* D\left(\sum_{j=1}^m L_j \bar{x}_j - o\right) \quad (4.4)$$

together with the dual problem

$$\text{find } \bar{v} \in \mathcal{G} \text{ such that } -o \in -\sum_{i=1}^m L_i A_i^{-1}(z_i - L_i^* \bar{v}) + D^{-1} \bar{v}. \quad (4.5)$$

**Corollary 4.3** In Problem 4.2, set  $R = (\text{Id} + \sum_{i=1}^m L_i L_i^*)^{-1}$  and assume that (4.4) has at least one solution. Let  $(\lambda_n)_{n \in \mathbb{N}}$  be a sequence in  $]0, 2[$  such that  $\sum_{n \in \mathbb{N}} \lambda_n(2 - \lambda_n) = +\infty$ , let  $(b_n)_{n \in \mathbb{N}}$  be a sequence in  $\mathcal{G}$ , and let  $v_0 \in \mathcal{G}$ . For every  $i \in \{1, \dots, m\}$ , let  $(a_{i,n})_{n \in \mathbb{N}}$  be a sequence in  $\mathcal{H}_i$ , let  $x_{i,0} \in \mathcal{H}_i$ , and set  $u_{i,0} = -L_i^* v_0$ . Suppose that  $\sum_{n \in \mathbb{N}} \lambda_n \sqrt{\|b_n\|^2 + \sum_{i=1}^m \|a_{i,n}\|^2} < +\infty$ , and set  $y_0 = \sum_{i=1}^m L_i x_{i,0}$  and

$$(\forall n \in \mathbb{N}) \quad \left\{ \begin{array}{l} \text{For } i = 1, \dots, m \\ \quad \left[ \begin{array}{l} p_{i,n} = J_{A_i}(x_{i,n} + u_{i,n} + z_i) + a_{i,n} \\ r_{i,n} = x_{i,n} + u_{i,n} - p_{i,n} \end{array} \right. \\ q_n = o + J_D(y_n + v_n - o) + b_n \\ s_n = y_n + v_n - q_n \\ t_n = R(\sum_{i=1}^m L_i r_{i,n} - s_n) \\ w_n = R(\sum_{i=1}^m L_i p_{i,n} - q_n) \\ \text{For } i = 1, \dots, m \\ \quad \left[ \begin{array}{l} x_{i,n+1} = x_{i,n} + \lambda_n(L_i^* t_n - r_{i,n}) \\ u_{i,n+1} = u_{i,n} - \lambda_n L_i^* w_n \end{array} \right. \\ y_{n+1} = y_n - \lambda_n(t_n + s_n) \\ v_{n+1} = v_n + \lambda_n w_n. \end{array} \right. \quad (4.6)$$

Then there exists a solution  $(\bar{x}_i)_{1 \leq i \leq m}$  to (4.4) and a solution  $\bar{v}$  to (4.5) such that the following hold:

- (i)  $\bar{v} \in D(\sum_{i=1}^m L_i \bar{x}_i - o)$  and  $(\forall i \in \{1, \dots, m\}) \quad z_i - L_i^* \bar{v} \in A_i \bar{x}_i$ .
- (ii)  $v_n \rightharpoonup \bar{v}$  and  $(\forall i \in \{1, \dots, m\}) \quad x_{i,n} \rightharpoonup \bar{x}_i$ .

*Proof.* Set  $B: \mathcal{G} \rightarrow 2^{\mathcal{G}}: y \mapsto D(y - o)$ ,  $\mathcal{H} = \bigoplus_{i=1}^m \mathcal{H}_i$ ,  $L: \mathcal{H} \rightarrow \mathcal{G}: (x_i)_{1 \leq i \leq m} \mapsto \sum_{i=1}^m L_i x_i$ , and  $A: \mathcal{H} \rightarrow 2^{\mathcal{H}}: (x_i)_{1 \leq i \leq m} \mapsto \times_{i=1}^m (-z_i + A_i x_i)$ . Then  $L^*: \mathcal{G} \rightarrow \mathcal{H}: y \mapsto (L_i^* y)_{1 \leq i \leq m}$  and hence Problem 4.2 is a special case of Problem 1.1. On the other hand, [2, Propositions 23.15 and 23.16] yield

$$J_A: (x_i)_{1 \leq i \leq m} \mapsto (J_{A_i}(x_i + z_i))_{1 \leq i \leq m} \quad \text{and} \quad J_B: y \mapsto o + J_D(y - o). \quad (4.7)$$

Now set  $(\forall n \in \mathbb{N}) a_n = (a_{i,n})_{1 \leq i \leq m}$ ,  $p_n = (p_{i,n})_{1 \leq i \leq m}$ ,  $r_n = (r_{i,n})_{1 \leq i \leq m}$ ,  $u_n = (u_{i,n})_{1 \leq i \leq m}$ , and  $x_n = (x_{i,n})_{1 \leq i \leq m}$ . Then (4.1) reduces to (4.6) and we can appeal to Theorem 4.1 to conclude.  $\square$

**Remark 4.4** In Problem 4.2, set  $D = \partial g$ , where  $g \in \Gamma_0(\mathcal{G})$ , and  $(\forall i \in \{1, \dots, m\}) A_i = \partial f_i$ , where  $f_i \in \Gamma_0(\mathcal{H}_i)$  and  $z_i \in \text{ran}(\partial f_i + L_i^*(\partial g)(L_i \cdot -o))$ . Then, arguing as in the proof of Corollary 3.7, we derive from Corollary 4.3 an algorithm for solving the primal problem

$$\underset{x_1 \in \mathcal{H}_1, \dots, x_m \in \mathcal{H}_m}{\text{minimize}} \quad \sum_{i=1}^m (f_i(x_i) - \langle x_i \mid z_i \rangle) + g\left(\sum_{i=1}^m L_i x_i - o_i\right) \quad (4.8)$$

together with the dual problem

$$\underset{v \in \mathcal{G}}{\text{minimize}} \quad \sum_{i=1}^m f_i^*(z_i - L_i^* v) + g^*(v) + \langle v \mid o \rangle \quad (4.9)$$

by replacing  $J_{A_i}$  by  $\text{prox}_{f_i}$  and  $J_D$  by  $\text{prox}_g$  in (4.6).

## 5 Concluding remarks

We have shown that the method of partial inverses can be used to solve composite monotone inclusions in duality and have presented a few applications of this new framework. Despite their apparent complexity, all the algorithms developed in this paper are instances of the method of partial inverses, which is itself an instance of the proximal point algorithm. This underlines the fundamental nature of the proximal point algorithm and its far reaching ramifications. Finally, let us note that in [4] the method of partial inverses was coupled to standard splitting methods for the sum of two monotone operators to solve inclusions of the form  $0 \in Ax + Bx + N_V x$ , where  $N_V$  is the normal cone operator of the closed vector subspace  $V$ . Combining this approach to our results should lead to new splitting methods for more general problems involving composite operators, such as those studied in [9].

**Acknowledgement.** This work was funded by the Deanship of Scientific Research (DSR), King Abdulaziz University, under grant number 4-130/1433 HiCi. The authors, therefore, acknowledge technical and financial support of KAU.

## References

- [1] H. H. Bauschke, R. I. Boţ, W. L. Hare, and W. M. Moursi, Attouch–Théra duality revisited: Paramonotonicity and operator splitting, *J. Approx. Theory*, 164, pp. 1065–1084, 2012.
- [2] H. H. Bauschke and P. L. Combettes, *Convex Analysis and Monotone Operator Theory in Hilbert Spaces*. Springer, New York, 2011.
- [3] H. H. Bauschke, P. L. Combettes, and D. R. Luke, Finding best approximation pairs relative to two closed convex sets in Hilbert spaces, *J. Approx. Theory*, 127, pp. 178–192, 2004.
- [4] L. M. Briceño-Arias, Forward-Douglas-Rachford splitting and forward-partial inverse method for solving monotone inclusions, preprint, <http://arxiv.org/abs/1212.5942>
- [5] L. M. Briceño-Arias and P. L. Combettes, A monotone+skew splitting model for composite monotone inclusions in duality, *SIAM J. Optim.*, 21, pp. 1230–1250, 2011.
- [6] R. S. Burachik, C. Sagastizábal, and S. Scheimberg, An inexact method of partial inverses and a parallel bundle method, *Optim. Methods Softw.*, 21, pp. 385–400, 2006.
- [7] P. L. Combettes, Solving monotone inclusions via compositions of nonexpansive averaged operators, *Optimization*, 53, pp. 475–504, 2004.

- [8] P. L. Combettes, Iterative construction of the resolvent of a sum of maximal monotone operators, *J. Convex Anal.*, 16, pp. 727–748, 2009.
- [9] P. L. Combettes and J.-C. Pesquet, Primal-dual splitting algorithm for solving inclusions with mixtures of composite, Lipschitzian, and parallel-sum type monotone operators, *Set-Valued Var. Anal.*, 20, pp. 307–330, 2012.
- [10] J. Eckstein and D. P. Bertsekas, On the Douglas-Rachford splitting method and the proximal point algorithm for maximal monotone operators, *Math. Programming*, 55, pp. 293–318, 1992.
- [11] J. Eckstein and M. C. Ferris, Smooth methods of multipliers for complementarity problems, *Math. Programming*, 86, pp. 65–90, 1999.
- [12] H. Idrissi, O. Lefebvre, and C. Michelot, Applications and numerical convergence of the partial inverse method, *Lecture Notes in Math.*, 1405, pp. 39–54, 1989.
- [13] B. Lemaire, The proximal algorithm, in: *New Methods in Optimization and Their Industrial Uses*, (J.-P. Penot, Ed.), pp. 73–87. Birkhäuser, Boston, MA, 1989.
- [14] Ph. Mahey, S. Oualibouch, and Pham Dinh Tao, Proximal decomposition on the graph of a maximal monotone operator, *SIAM J. Optim.* 5, pp. 454–466, 1995.
- [15] T. Pennanen, Dualization of generalized equations of maximal monotone type, *SIAM J. Optim.*, 10, pp. 809–835, 2000.
- [16] T. Pennanen, A splitting method for composite mappings, *Numer. Funct. Anal. Optim.*, 23, pp. 875–890, 2002.
- [17] S. M. Robinson, Composition duality and maximal monotonicity, *Math. Programming*, 85, pp. 1–13, 1999.
- [18] R. T. Rockafellar, Duality and stability in extremum problems involving convex functions, *Pacific J. Math.*, 21, pp. 167–187, 1967.
- [19] R. T. Rockafellar, Monotone operators and the proximal point algorithm, *SIAM J. Control Optim.*, 14, pp. 877–898, 1976.
- [20] J. E. Spingarn, Partial inverse of a monotone operator, *Appl. Math. Optim.*, 10, pp. 247–265, 1983.
- [21] J. E. Spingarn, Applications of the method of partial inverses to convex programming: Decomposition, *Math. Programming* 32, pp. 199–223, 1985.
- [22] J. E. Spingarn, A projection method for least-squares solutions to overdetermined systems of linear inequalities, *Linear Algebra and Its Applications*, 86, pp. 211–236, 1987.