

Long time average of first order mean field games and weak KAM theory

P. Cardaliaguet*

October 2, 2018

Abstract

We show that the long time average of solutions of first order mean field game systems in finite horizon is governed by an ergodic system of mean field game type. The well-posedness of this later system and the uniqueness of the ergodic constant rely on weak KAM theory.

Introduction

The aim of this paper is to study the link between the finite horizon first order mean field game system

$$\left\{ \begin{array}{ll} (i) & -\partial_t u^T + H(x, Du^T) = F(x, m^T(t)) \text{ in } (0, T) \times \mathbb{R}^d \\ (ii) & \partial_t m^T - \operatorname{div}(m^T D_p H(x, Du^T)) = 0 \text{ in } (0, T) \times \mathbb{R}^d \\ (iii) & m^T(0) = m_0, u^T(x, T) = u^f(x) \text{ in } \mathbb{R}^d \end{array} \right. \quad (1)$$

and the ergodic first order mean field game system

$$\left\{ \begin{array}{ll} (i) & \bar{\lambda} + H(x, D\bar{u}) = F(x, \bar{m}) \text{ in } \mathbb{R}^d \\ (ii) & -\operatorname{div}(\bar{m} D_p H(x, D\bar{u})) = 0 \text{ in } \mathbb{R}^d \\ (iii) & \int_Q \bar{u} dx = 0, \quad \int_Q \bar{m} dx = 1 \end{array} \right. \quad (2)$$

Let us recall that mean field games have been introduced simultaneously by Lasry and Lions [8, 9, 10] and by Huang, Caines and Malhamé [7] to analyze large population stochastic differential games. In (1), the scalar unknowns u^T, m^T are defined on $[0, T] \times \mathbb{R}^d$ and F is a coupling between the two equations. The function u^T can be understood as the value function—for a typical and small player—of a finite horizon optimal control problem in which the density m^T of the other players enters as a data. The optimal feedback of this small player is then given by $-D_p H(x, Du(x))$. When all players play according to this rule, their density $m^T = m^T(t, x)$ evolves in time by equation (1)-(ii). The ergodic problem (2) has similar interpretation, expect that now the optimal control problem is of ergodic type: the unknowns are the ergodic constant $\bar{\lambda}$, the value function \bar{u} of the ergodic problem and \bar{m} which is an associate invariant measure.

In analogy with the case of optimal control problems, it is expected that, as $T \rightarrow +\infty$, the solution of the finite horizon system (1) somehow converges to the solution of the ergodic system (2). For second order mean field game systems (i.e., systems corresponding to stochastic control

*Ceremade, Université Paris-Dauphine, Place du Maréchal de Lattre de Tassigny, 75775 Paris cedex 16 (France)

problems with a nondegenerate diffusion) this kind of behavior has been first discussed in [11] and then developed and sharpened in [1, 2]. Results in the discrete setting were also obtained in [6]. Typically it is proved in the above mentioned papers that $u^T(0, \cdot)/T$ converges to $\bar{\lambda}$ while (a rescaled version of) m^T converges to the invariant measure \bar{m} . The precise meaning of the convergence depends on the coupling F , which can be of local or nonlocal nature: when F is local (i.e., $F(x, m(t)) = \bar{F}(x, m(t, x))$ depends on the value $m(t, x)$ of the density of $m(t)$), the convergence holds in Lebesgue spaces. When the coupling F is of nonlocal nature and smoothing, the convergence is uniform. The main result of [1, 2] is an exponential convergence rate when the coupling—and the diffusion—are “strong enough”.

Here we consider the same issue for first order mean field games in which the coupling F is nonlocal and smoothing. We also assume that all functions are periodic in space and set $\mathbb{T}^d = \mathbb{R}^d/\mathbb{Z}^d$. In the uncoupled case $F = 0$, this problem has been the object of a lot of attention in the recent years under the name of weak-KAM theory (see in particular the pioneering works [3, 4, 14, 15] and the monograph [5]). Compared to the second order setting, several interesting issues arise: even for $F = 0$, one cannot expect the ergodic system to have a *unique* solution. As a consequence, in the presence of the coupling, *it is not even clear that the ergodic constant $\bar{\lambda}$ is unique*. There is also a strong difficulty to give a meaning to (2)-(ii): indeed, as a solution of a classical Hamilton-Jacobi equation, the map \bar{u} is at most Lipschitz continuous (actually semiconcave); on another hand, the ergodic measure cannot be expected to have a density (again, this is what happens in general for $F = 0$): as a consequence *the term $\bar{m}D_p H(x, D\bar{u})$ is a priori ill-defined*. To overcome these difficulties, we use two tools: the first one is a typical regularity property arising in weak-KAM theory [3, 4, 5]: the measure \bar{m} happens to concentrate on the so-called projected Mather-set, in which the derivative of \bar{u} exists and is uniquely defined (independently of the solution \bar{u}). This allows to give a meaning to the term $\bar{m}D_p H(x, D\bar{u})$ and provides a key tool for showing the existence of solutions to (2). As for the uniqueness issue, we introduce a weak coercivity condition on the coupling: namely we assume that there is a constant $\bar{c} > 0$ such that, for any pair of measures m_1, m_2 ,

$$\int_{\mathbb{T}^d} (F(x, m_1) - F(x, m_2)) d(m_1 - m_2) \geq \bar{c} \int_{\mathbb{T}^d} (F(x, m_1) - F(x, m_2))^2 dx$$

This condition—which is quite natural in the context of mean field game theory (see Example 1.1 below)—entails the uniqueness of the ergodic constant. Then, using energy estimates, we prove our main result concerning the convergence of $u^T(0, \cdot)/T$ to $\bar{\lambda}$: there is a constant C such that

$$\sup_{t \in [0, T]} \left\| \frac{u^T(t, \cdot)}{T} - \bar{\lambda} \left(1 - \frac{t}{T} \right) \right\|_{\infty} \leq \frac{C}{T^{\frac{1}{2}}}.$$

As for the convergence of m^T , we have little information due to the lack of uniqueness of the ergodic measure \bar{m} . However, the coupling $F(\cdot, \bar{m})$ turns out to be unique and therefore it seems the good quantity to look at: indeed we have the following estimate:

$$\int_0^T \|F(\cdot, m^T(t)) - F(\cdot, \bar{m})\|_{\infty} dt \leq CT^{\frac{1}{2}}.$$

Note that this inequality means that $F(\cdot, m^T(t))$ must be close to $F(\cdot, \bar{m})$ on a large amount of time.

The paper is organized as follows: in a first part we introduce the notations and state the assumptions used throughout the paper. Then we study the ergodic mean field game system.

In the last section we prove our convergence result. In appendix we recall the main steps of the proof for the well-posedness of (1).

Acknowledgement: We wish to thank Yves Achdou for fruitful discussions.

This work has been partially supported by the Commission of the European Communities under the 7-th Framework Programme Marie Curie Initial Training Networks Project SADCO, FP7-PEOPLE-2010-ITN, No 264735, and by the French National Research Agency ANR-10-BLAN 0112 and ANR-12-BS01-0008-01.

1 Preliminaries

Throughout this note, we work in the periodic setting: we denote by \mathbb{T}^d the torus $\mathbb{R}^d/\mathbb{Z}^d$, by $P(\mathbb{T}^d)$ the set of Borel probability measures on \mathbb{T}^d , and by $P(\mathbb{T}^d \times \mathbb{R}^d)$ the set of Borel probability measures on $\mathbb{T}^d \times \mathbb{R}^d$. Both sets \mathbb{T}^d and $P(\mathbb{T}^d \times \mathbb{R}^d)$ are endowed with the weak-* convergence. Let us recall that $P(\mathbb{T}^d)$ is compact for this topology. It will be convenient to put a metric on $P(\mathbb{T}^d)$ (which metricizes the weak-* topology): we shall work with the Monge-Wasserstein distance defined, for any $\mu, \nu \in P(\mathbb{T}^d)$, by

$$\mathbf{d}_1(\mu, \nu) = \sup_h \left\{ \int_{\mathbb{T}^d} h d(\mu - \nu) \right\} \quad (3)$$

where the supremum is taken over all the maps $h : \mathbb{T}^d \rightarrow \mathbb{R}$ which are 1-Lipschitz continuous.

The maps H and F are periodic in the space arguments: $H : \mathbb{T}^d \times \mathbb{R}^d \rightarrow \mathbb{R}$ while $F : \mathbb{T}^d \times P(\mathbb{T}^d) \rightarrow \mathbb{R}$. In the same way, the initial and terminal conditions m_0 and u^f —which are fixed throughout the paper, are periodic in space: $u^f : \mathbb{T}^d \rightarrow \mathbb{R}$ is supposed to be of class \mathcal{C}^2 while m^f belongs to $P(\mathbb{T}^d)$ is assumed to have a bounded density ($m^f \in L^\infty$).

We now state our key assumptions on the data: these conditions are valid throughout the paper. The map F is supposed to be regularizing:

$$\text{The map } m \rightarrow F(\cdot, m) \text{ is Lipschitz continuous from } P(\mathbb{T}^d) \text{ to } \mathcal{C}^2(\mathbb{T}^d). \quad (4)$$

In particular, as $P(\mathbb{T}^d)$ is compact, there is $\bar{C} > 0$ such that

$$\sup_{m \in P(\mathbb{T}^d)} \|F(\cdot, m)\|_{\mathcal{C}^2} \leq \bar{C} \quad (5)$$

As explained in the introduction, our key assumption is the following weak coercivity for the coupling: there is a constant $\bar{c} > 0$ such that, for any $m_1, m_2 \in P(\mathbb{T}^d)$,

$$\int_{\mathbb{T}^d} (F(x, m_1) - F(x, m_2)) d(m_1 - m_2) \geq \bar{c} \int_{\mathbb{T}^d} (F(x, m_1) - F(x, m_2))^2 dx. \quad (6)$$

An example of map F satisfying (4) and (6) is given below. Concerning the Hamiltonian, we suppose that H is of class \mathcal{C}^2 on $\mathbb{T}^d \times \mathbb{R}^d$ and quadratic-like in the second variable:

$$H \in \mathcal{C}^2(\mathbb{T}^d \times \mathbb{R}^d) \text{ and } \frac{1}{\bar{C}} I_d \leq D_{pp}^2 H(x, p) \leq \bar{C} I_d \quad \forall (x, p) \in \mathbb{T}^d \times \mathbb{R}^d. \quad (7)$$

Let us recall that, under assumptions (4) on F and (7) on H , for any time horizon T , there is a unique solution (u^T, m^T) to the mean field game system (1): by a solution, we mean that u^T is a Lipschitz continuous viscosity solution to (1)-(i) while $m^T \in L^\infty((0, T) \times \mathbb{T}^d)$ is a solution

of (1)-(ii) in the sense of distribution (see [10] and Theorem 4.1 in appendix). Recall also that the map $t \rightarrow m^T(t)$ is weakly continuous as a measure on \mathbb{T}^d . In particular, for any continuous map $\phi : \mathbb{T}^d \rightarrow \mathbb{R}$, the integral $\int_{\mathbb{T}^d} \phi(x) m^T(t, x)$ is continuous in $t \in [0, T]$.

Example 1.1. Assume that $F : \mathbb{T}^d \times P(\mathbb{T}^d) \rightarrow \mathbb{R}$ is of the form

$$F(x, m) = (\bar{F}(\cdot, m \star \xi(\cdot)) \star \xi)(x)$$

where $\xi : \mathbb{R}^d \rightarrow \mathbb{R}$ is a smooth, even kernel with compact support and where $\bar{F} : \mathbb{T}^d \times \mathbb{R} \rightarrow \mathbb{R}$ is a smooth map for which there is a constant $c > 0$ with

$$c \leq \frac{\partial \bar{F}(x, z)}{\partial z} \leq \frac{1}{c} \quad \forall (x, z) \in \mathbb{T}^d \times \mathbb{R}. \quad (8)$$

Then F satisfies conditions (4) and (6).

Proof. The Lipschitz continuity assumption (4) is straightforward from the smoothness assumption on \bar{F} . Let us now check (6). We have, on the one hand,

$$\begin{aligned} & \int_{\mathbb{T}^d} (F(x, m_1) - F(x, m_2)) d(m_1 - m_2) \\ &= \int_{\mathbb{T}^d} [\bar{F}(\cdot, m_1 \star \xi) - \bar{F}(\cdot, m_2 \star \xi)] \star \xi d(m_1 - m_2) \\ &= \int_{\mathbb{T}^d} [\bar{F}(\cdot, m_1 \star \xi) - \bar{F}(\cdot, m_2 \star \xi)] (m_1 \star \xi - m_2 \star \xi) \end{aligned}$$

which, in view of our growth condition on \bar{F} , implies that

$$\int_{\mathbb{T}^d} (F(x, m_1) - F(x, m_2)) d(m_1 - m_2) \geq c \int_{\mathbb{T}^d} (m_1 \star \xi - m_2 \star \xi)^2 \quad (9)$$

On another hand,

$$\begin{aligned} & \int_{\mathbb{T}^d} (F(x, m_1) - F(x, m_2))^2 dx \\ &= \int_{\mathbb{T}^d} [(\bar{F}(\cdot, m_1 \star \xi) - \bar{F}(\cdot, m_2 \star \xi)) \star \xi]^2 dx \\ &\leq \|(\bar{F}(\cdot, m_1 \star \xi) - \bar{F}(\cdot, m_2 \star \xi)) \star \xi\|_\infty^2 \\ &\leq \|\xi\|_{L^2(\mathbb{R}^d)}^2 \|(\bar{F}(\cdot, m_1 \star \xi) - \bar{F}(\cdot, m_2 \star \xi))\|_{L^2(\mathbb{T}^d)}^2 \\ &\leq \|\xi\|_{L^2(\mathbb{R}^d)}^2 c^{-2} \int_{\mathbb{T}^d} (m_1 \star \xi - m_2 \star \xi)^2 dx \end{aligned}$$

where we have used the second inequality in the right-hand side of assumption (8) in the last line. Using (9) we deduce that F satisfies (6) with $\bar{c} = c^3 \|\xi\|_{L^2(\mathbb{R}^d)}^{-2}$. \square

2 The ergodic problem

In this section, we show that the ergodic system (2) is well-posed.

Definition 2.1. We say that the triple $(\bar{\lambda}, \bar{u}, \bar{m})$ is a solution of (2) if \bar{u} is Lipschitz continuous viscosity solution of (2)-(i), if $D\bar{u}(x)$ exists for \bar{m} -a.e. $x \in \mathbb{T}^d$ and if (2)-(ii) is satisfied in the sense of distribution.

Remark 2.2. Because of the regularity of H and F , the map \bar{u} is semiconcave (see, for instance, Theorem 3.3 of [12]). In particular, $D\bar{u}$ is continuous at \bar{m} -a.e. $x \in \mathbb{T}^d$. Note also that $\bar{m}D\bar{u}$ is a vector measure.

Here is our main result concerning system (2).

Theorem 2.3. Under assumption (4) and (7), there is at least one solution of the ergodic problem (2). If, moreover, assumption (6) holds, the ergodic constant is unique: more precisely, if $(\bar{\lambda}_1, \bar{u}_1, \bar{m}_1)$ and $(\bar{\lambda}_2, \bar{u}_2, \bar{m}_2)$ are two solutions of (2), then $\bar{\lambda}_1 = \bar{\lambda}_2$ and $F(\cdot, \bar{m}_1) = F(\cdot, \bar{m}_2)$.

Proof. The existence of the solution relies on several aspects of weak-KAM theory, as developed by Fathi in [3, 4, 5]. Let L be the Fenchel conjugate of H with respect to the last variable:

$$L(x, v) = \sup_{p \in \mathbb{R}^d} \langle p, v \rangle - H(x, p) \quad \text{for } (x, v) \in \mathbb{T}^d \times \mathbb{R}^d.$$

In view of our assumptions, L is of class \mathcal{C}^2 and uniformly convex. Given $m \in P(\mathbb{T}^d)$, we consider the set E^m of measures η on $\mathbb{T}^d \times \mathbb{R}^d$ which are invariant under the Lagrangian flow $\phi_t^m = (\gamma(t), \dot{\gamma}(t))$ defined by

$$\begin{cases} -\frac{d}{dt} D_v [L(x, \dot{x}) + F(x, m)] + D_x [L(x, \dot{x}) + F(x, m)] = 0 \\ x(0) = x, \dot{x}(0) = v \end{cases}$$

The set $M(m)$ of the minimizers of the map

$$\eta \rightarrow \int_{\mathbb{T}^d \times \mathbb{R}^d} L(x, v) + F(x, m) \, d\eta(x, v)$$

over E^m is nonempty, compact and convex subset of $P(\mathbb{T}^d \times \mathbb{R}^d)$. If $\pi : \mathbb{T}^d \times \mathbb{R}^d \rightarrow \mathbb{T}^d$ denotes the canonical projection, we finally set $C(m) = \{\pi\# \eta, \eta \in M(m)\}$. Then $C(m)$ is a convex, compact nonempty subset of $P(\mathbb{T}^d)$. From the continuity assumption (4) on F and the coercivity of L , the set-valued map $m \rightarrow C(m)$ has a compact graph. Using Kakutani fixed point theorem, one deduce then that C has a fixed point: there is $\bar{m} \in P(\mathbb{T}^d)$ such that $\bar{m} \in C(\bar{m})$. Let $\bar{\eta} \in M(\bar{m})$ be such that $\pi\#\bar{\eta} = \bar{m}$.

Let $(\bar{\lambda}, \bar{u})$ be such that \bar{u} is a continuous, periodic viscosity solution of

$$\bar{\lambda} + H(x, D\bar{u}) = F(x, \bar{m}) \quad \text{in } \mathbb{T}^d.$$

Since the Hamiltonian is coercive and satisfies our smoothness conditions, \bar{u} is Lipschitz continuous and semiconcave. Following Fathi ([3], Corollary 2),

$$\begin{aligned} \bar{\lambda} &= \min_{\eta \in E^{\bar{m}}} \int_{\mathbb{T}^d \times \mathbb{R}^d} L(x, v) + F(x, \bar{m}) \, d\eta(x, v) \\ &= \int_{\mathbb{T}^d \times \mathbb{R}^d} L(x, v) + F(x, \bar{m}) \, d\bar{\eta}(x, v) \end{aligned}$$

Moreover $D\bar{u}$ exists everywhere on the support of \bar{m} and is Lipschitz continuous on this support ([3], Proposition 3). It is also known that the canonical map $\pi : \text{Spt}(\bar{\eta}) \rightarrow \text{Spt}(\bar{m})$ is one-to-one

and its inverse is given by $x \rightarrow (x, D_p H(x, D\bar{u}(x)))$ on $\text{Spt}(\bar{m})$. In particular, the first component γ_t^x of the flow ϕ_t satisfies

$$\frac{d}{dt}\gamma_t^x = D_p H(\gamma_t^x, D\bar{u}(\gamma_t^x)) \text{ for } x \in \text{Spt}(\bar{m}).$$

This implies that equality $-\text{div}(\bar{m}D\bar{u}) = 0$ holds in \mathbb{T}^d : indeed, as $\bar{\eta}$ is invariant under the flow $\phi_t^{\bar{m}}$, \bar{m} is invariant under the flow γ_t^x and we have, for any test function $f \in \mathcal{C}^\infty(\mathbb{T}^d)$,

$$\begin{aligned} 0 &= \frac{d}{dt} \int_{\mathbb{T}^d} f(\gamma_t^x) d\bar{m}(x) = \int_{\mathbb{T}^d} \langle Df(\gamma_t^x), D_p H(\gamma_t^x, D\bar{u}(\gamma_t^x)) \rangle d\bar{m}(x) \\ &= \int_{\mathbb{T}^d} \langle Df(y), D_p H(y, D\bar{u}(y)) \rangle d\bar{m}(y). \end{aligned}$$

We now show that uniqueness holds. Let $(\bar{\lambda}_1, \bar{u}_1, \bar{m}_1)$ and $(\bar{\lambda}_2, \bar{u}_2, \bar{m}_2)$ be two solutions of (2). Let $\epsilon > 0$, $\xi : \mathbb{R}^d \rightarrow \mathbb{R}$ be a smooth, nonnegative, symmetric kernel with a support contained in the unit ball and of integral one. We set $\xi^\epsilon(x) = \frac{1}{\epsilon^d} \xi(x/\epsilon)$ and, for $i = 1, 2$, $m_i^\epsilon = \xi_\epsilon \star \bar{m}_i$ and $V_i^\epsilon = \frac{\xi^\epsilon \star (\bar{m}_i D_p H(\cdot, D\bar{u}_i))}{m_i^\epsilon}$. Then $-\text{div}(m_i^\epsilon V_i^\epsilon) = 0$ in \mathbb{T}^d . We multiply this equality by $(\bar{u}_1 - \bar{u}_2)$, integrate by parts, and subtract the resulting formulas to get:

$$\int_{\mathbb{T}^d} \langle D(\bar{u}_1 - \bar{u}_2), m_1^\epsilon V_1^\epsilon - m_2^\epsilon V_2^\epsilon \rangle = 0.$$

Therefore

$$\begin{aligned} 0 &= \int_{\mathbb{T}^d} \langle D(\bar{u}_1 - \bar{u}_2), \xi^\epsilon \star (\bar{m}_1 D_p H(\cdot, D\bar{u}_1) - \bar{m}_2 D_p H(\cdot, D\bar{u}_2)) \rangle \\ &= \int_{\mathbb{T}^d} \langle D(\bar{u}_1 - \bar{u}_2), m_1^\epsilon D_p H(x, D\bar{u}_1) - m_2^\epsilon D_p H(x, D\bar{u}_2) \rangle + R_\epsilon \end{aligned} \tag{10}$$

where (the double integral being on $\mathbb{R}^d \times \mathbb{T}^d$)

$$\begin{aligned} R_\epsilon &= \iint \xi^\epsilon(x-y) \langle D(\bar{u}_1 - \bar{u}_2)(x), (D_p H(y, D\bar{u}_1(y)) - D_p H(x, D\bar{u}_1(x))) \rangle \bar{m}_1(dy) dx \\ &\quad - \iint \xi^\epsilon(x-y) \langle D(\bar{u}_1 - \bar{u}_2)(x), (D_p H(y, D\bar{u}_2(y)) - D_p H(x, D\bar{u}_2(x))) \rangle \bar{m}_2(dy) dx \\ &= \iint \xi(z) \langle D(\bar{u}_1 - \bar{u}_2)(y + \epsilon z), (D_p H(y, D\bar{u}_1(y)) - D_p H(y + \epsilon z, D\bar{u}_1(y + \epsilon z))) \rangle \bar{m}_1(dy) dz \\ &\quad - \iint \xi(z) \langle D(\bar{u}_1 - \bar{u}_2)(y + \epsilon z), (D_p H(y, D\bar{u}_2(y)) - D_p H(y + \epsilon z, D\bar{u}_2(y + \epsilon z))) \rangle \bar{m}_2(dy) dz \end{aligned}$$

Since \bar{u}_1 and \bar{u}_2 are Lipschitz continuous, we get

$$\begin{aligned} |R_\epsilon| &\leq \\ &C \iint \xi(z) |D_p H(y, D\bar{u}_1(y)) - D_p H(y + \epsilon z, D\bar{u}_1(y + \epsilon z))| \bar{m}_1(dy) dz \\ &\quad + C \iint \xi(z) |D_p H(y, D\bar{u}_2(y)) - D_p H(y + \epsilon z, D\bar{u}_2(y + \epsilon z))| \bar{m}_2(dy) dz \end{aligned}$$

Following Remark 2.2, the maps $D\bar{u}_i$ are continuous at \bar{m}_i -a.e. $y \in \mathbb{R}^d$ for $i = 1, 2$. Then Lebesgue Theorem implies that $R_\epsilon \rightarrow 0$ as $\epsilon \rightarrow 0$.

Next we multiply the equality satisfied by the \bar{u}_i by $(m_1^\epsilon - m_2^\epsilon)$, integrate in space (recalling that the m_i^ϵ are probability measures) and subtract to get

$$\int_{\mathbb{T}^d} (m_1^\epsilon - m_2^\epsilon) [H(x, D\bar{u}_1) - H(x, D\bar{u}_2) - F(x, \bar{m}_1) + F(x, \bar{m}_2)] dx = 0 .$$

We combine the above equality with (10) and obtain

$$\begin{aligned} \int_{\mathbb{T}^d} \langle D(\bar{u}_1 - \bar{u}_2), m_1^\epsilon D_p H(x, D\bar{u}_1) - m_2^\epsilon D_p H(x, D\bar{u}_2) \rangle \\ - (m_1^\epsilon - m_2^\epsilon) [H(x, D\bar{u}_1) - H(x, D\bar{u}_2) - F(x, \bar{m}_1) + F(x, \bar{m}_2)] dx = -R_\epsilon . \end{aligned}$$

Let us set (for $i = 1, 2$) $H_i = H(x, D\bar{u}_i)$ and $DH_i = D_p H(x, D\bar{u}_i)$. Then, following Lasry-Lions classical computation [8, 9, 10] the above formula can be rearranged as

$$\begin{aligned} \int_{\mathbb{T}^d} m_1^\epsilon [H_2 - H_1 - \langle DH_1, D(\bar{u}_2 - \bar{u}_1) \rangle] + m_2^\epsilon [H_1 - H_2 - \langle DH_2, D(\bar{u}_1 - \bar{u}_2) \rangle] \\ + \int_{\mathbb{T}^d} (m_1^\epsilon - m_2^\epsilon)(F(x, \bar{m}_1) - F(x, \bar{m}_2)) = -R_\epsilon . \end{aligned}$$

Using the convexity of H , the terms $H_2 - H_1 - \langle DH_1, D(\bar{u}_2 - \bar{u}_1) \rangle$ and $H_1 - H_2 - \langle DH_2, D(\bar{u}_1 - \bar{u}_2) \rangle$ are nonnegative. So

$$\int_{\mathbb{T}^d} (m_1^\epsilon - m_2^\epsilon)(F(x, \bar{m}_1) - F(x, \bar{m}_2)) \leq -R_\epsilon .$$

We let $\epsilon \rightarrow 0$ to get

$$\int_{\mathbb{T}^d} (F(x, \bar{m}_1) - F(x, \bar{m}_2)) d(\bar{m}_1 - \bar{m}_2) \leq 0 ,$$

which in turn implies that

$$\int_{\mathbb{T}^d} (F(x, \bar{m}_1) - F(x, \bar{m}_2))^2 dx \leq 0$$

thanks to assumption (6). So $F(x, \bar{m}_1) = F(x, \bar{m}_2)$. Then $(\bar{\lambda}_1, \bar{u}_1)$ and $(\bar{\lambda}_2, \bar{u}_2)$ are two solutions of an ergodic problem for the same Hamilton-Jacobi equation. This is known to entail that $\bar{\lambda}_1 = \bar{\lambda}_2$ (see, for instance, [13]). \square

Example 2.4. Assume for instance that H is quadratic:

$$H(x, p) = \frac{1}{2}|p|^2 - V(x) ,$$

where $V : \mathbb{R}^d \rightarrow \mathbb{R}$ is a smooth, periodic map. Let $(\bar{\lambda}, \bar{u}, \bar{m})$ be a solution of (2). Then, following [13],

$$\bar{\lambda} = - \min_{x \in \mathbb{T}^d} \{V(x) + F(x, \bar{m})\}$$

and \bar{m} is supported in the set $\operatorname{argmin}_{x \in \mathbb{T}^d} \{V(x) + F(x, \bar{m})\}$. Moreover, $D\bar{u} = 0$ in this set.

3 Convergence

Let us set $v^T(s, x) = u^T(sT, x)$ and $\nu^T(s, x) = m^T(sT, x)$ for $(s, x) \in [0, 1] \times \mathbb{T}^d$. Our aim is to prove the uniform convergence of $\frac{v^T(s, \cdot)}{T}$ to $\bar{\lambda}(1-s)$, where $\bar{\lambda}$ is the unique ergodic constant associated to the problem (2).

Theorem 3.1. *Under assumptions (4), (7) and (6), the following estimates hold:*

$$\sup_{s \in [0, 1]} \left\| \frac{v^T(s, \cdot)}{T} - \bar{\lambda}(1-s) \right\|_{\infty} \leq \frac{C}{T^{\frac{1}{2}}}$$

and

$$\int_0^1 \|F(\cdot, \nu^T(s)) - F(\cdot, \bar{m})\|_{\infty} ds \leq \frac{C}{T^{\frac{1}{2}}}.$$

In particular, there is a uniform convergence of the map $(s, x) \rightarrow \frac{v^T(s, x)}{T}$ to the map $s \rightarrow \bar{\lambda}(1-s)$ as $T \rightarrow +\infty$.

In order to prove Theorem 3.1, we need some uniform estimates:

Lemma 3.2. *The map u^T is uniformly (with respect to T) Lipschitz continuous.*

Proof. Throughout the proof C denotes a constant which varies with the data and may change from line to line. Since Du^T is bounded (with a bound which depends a priori on T) and H is of class \mathcal{C}^2 , the map $t \rightarrow m^T(t)$ is Lipschitz continuous for the Monge-Wasserstein distance (see Lemma 4.2 below). We denote by L_T its Lipschitz constant and by C_0 be the Lipschitz continuity modulus of F given by assumption (4). We claim that

$$\|\partial_t u\|_{\infty} \leq \|F\|_{\infty} + \|H(\cdot, Du^f)\|_{\infty} + C_0 L_T T. \quad (11)$$

To prove (11), we first note that, as the terminal condition u^f being of class \mathcal{C}^2 , the maps

$$(t, x) \rightarrow u^f(x) \pm \left(\|F\|_{\infty} + \|H(\cdot, Du^f)\|_{\infty} \right) (T - t)$$

are respectively super- (for $+$) and sub- (for $-$) solutions of equation (1)-(i) with terminal condition u^f . Since u is a solution of this equation, we get by comparison

$$\|u(T - h, \cdot) - u^f\|_{\infty} \leq \left(\|F\|_{\infty} + \|H(\cdot, Du^f)\|_{\infty} \right) h \quad \forall h \in (0, T). \quad (12)$$

On another hand, since the left-hand side of equation (1)-(i) is independent of time while the right-hand side is Lipschitz continuous in time, with a Lipschitz constant bounded above by $C_0 L_T$, the maps

$$(t, x) \rightarrow u(t - h, x) \pm \left(\|u(T - h, \cdot) - u^f\|_{\infty} + C_0 L_T h (T - t) \right)$$

are respectively a super (for $+$) and subsolution (for $-$) of the equation satisfied by u . Using again comparison, we have

$$|u(t, x) - u(t - h, x)| \leq \|u(T - h, \cdot) - u^f\|_{\infty} + C_0 L_T h (T - t) \quad \forall (t, x) \in (h, T) \times \mathbb{T}^d.$$

Plugging (12) into this last inequality then gives (11).

Using the coercivity of H in (1)-(i), we obtain therefore

$$\|Du^T\|_\infty^2 \leq C(L_T + 1). \quad (13)$$

As H is of class \mathcal{C}^2 , we have then

$$\|D_p H(\cdot, Du^T(\cdot, \cdot))\|_\infty \leq C(L_T + 1)^{\frac{1}{2}}.$$

Then we use Lemma 4.2, which states that m^T is Lipschitz continuous for the Monge-Wasserstein distance with a Lipschitz constant bounded above by $\|D_p H(\cdot, Du^T(\cdot, \cdot))\|_\infty$. So

$$L_T \leq C(L_T + 1)^{\frac{1}{2}}.$$

This shows that L_T is bounded uniformly in T and, in view of (11) and (13) that u^T is Lipschitz continuous with a constant independent of T \square

Lemma 3.3 (Energy estimate). *If (u^T, m^T) is as above and $(\bar{\lambda}, \bar{u}, \bar{m})$ is a solution of the ergodic problem (2), then*

$$\int_0^T \int_{\mathbb{T}^d} (F(x, m^T(t)) - F(x, \bar{m})) d(m^T(t) - \bar{m}) dt \leq C$$

Proof. We use the same kind of argument as for the uniqueness part of Theorem 2.3. Let $\epsilon > 0$, ξ and $\xi^\epsilon(x) = \frac{1}{\epsilon^d} \xi(x/\epsilon)$ be as before. We set $m^\epsilon = \xi_\epsilon \star m$ and $V^\epsilon = \frac{\xi^\epsilon \star (\bar{m} D_p H(\cdot, D\bar{u}))}{m^\epsilon}$, so that $-\operatorname{div}(m^\epsilon V^\epsilon) = 0$ in \mathbb{T}^d . We multiply this equality by $(u^T(t) - \bar{u})$ and integrate on $(0, T) \times \mathbb{T}^d$:

$$\begin{aligned} 0 &= \int_0^T \int_{\mathbb{T}^d} \langle D(u^T(t) - \bar{m}), m^\epsilon V^\epsilon \rangle \\ &= \int_0^T \int_{\mathbb{T}^d} \langle D(u^T(t) - \bar{m}), m^\epsilon D_p H(x, D\bar{u}) \rangle + R_\epsilon \end{aligned} \quad (14)$$

where, as in the proof of Theorem 2.3, $R_\epsilon \rightarrow 0$ as $\epsilon \rightarrow 0$. Since $m^T(t)$ and m^ϵ are probability measures, we also have, in view of (2)-(i):

$$\int_0^T \int_{\mathbb{T}^d} (m^T(t) - m^\epsilon)(H(x, D\bar{u}) - F(x, \bar{m})) = 0. \quad (15)$$

From (1)-(ii) we have

$$\partial_t(m^T - m^\epsilon) - \operatorname{div}(m^T D_p H(x, Du^T)) = 0.$$

Multiplying this equality by the Lipschitz map $u^T - \bar{u}$ and integrating in time-space gives, thanks to (1)-(i),

$$\begin{aligned} 0 &= \int_{\mathbb{T}^d} \left[(u^f - \bar{u})(m^T(T) - m^\epsilon) - (u^T(0) - \bar{u})(m_0 - m^\epsilon) \right] \\ &\quad + \int_0^T \int_{\mathbb{T}^d} -(H(x, Du^T(t)) - F(x, m^T(t)))(m^T(t) - m^\epsilon) \\ &\quad + \int_0^T \int_{\mathbb{T}^d} \langle D(u^T(t) - \bar{u}), m^T(t) D_p H(x, u^T(t)) \rangle \end{aligned} \quad (16)$$

Note that the first integral is bounded uniformly with respect to T because

$$\left| \int_{\mathbb{T}^d} (u^f - \bar{u})(m^T(T) - m^\epsilon) \right| \leq 2(\|u^f\|_\infty + \|\bar{u}\|_\infty)$$

while, since m_0 and m^ϵ are probability measures,

$$\begin{aligned} & \left| \int_{\mathbb{T}^d} (u^T(0) - \bar{u})(m_0 - m^\epsilon) \right| \\ & \leq \left| \int_{\mathbb{T}^d} (u^T(0) - \int_{\mathbb{T}^d} u^T(0))(m_0 - m^\epsilon) \right| + \left| \int_{\mathbb{T}^d} \bar{u}(m_0 - m^\epsilon) \right| \\ & \leq 2\|Du^T(0)\|_\infty + 2\|\bar{u}\|_\infty \end{aligned}$$

where Du^T is uniformly bounded. Putting together (14), (15), (16) and rearranging as in the proof of Theorem 2.3 we obtain

$$\begin{aligned} & \int_0^T \int_{\mathbb{T}^d} m^T(t) [\bar{H} - H(t) - \langle DH(t), D(\bar{u} - u^T) \rangle] \\ & + \int_0^T \int_{\mathbb{T}^d} m^\epsilon [H(t) - \bar{H} - \langle D\bar{H}, D(u^T(t) - \bar{u}) \rangle] \\ & + \int_0^T \int_{\mathbb{T}^d} (m^T(t) - m^\epsilon)(F(x, m^T(t)) - F(x, \bar{m})) \leq C + R_\epsilon \end{aligned}$$

where we have set $H(t) = H(x, Du^T(t, x))$, $DH(t) = D_p H(x, Du^T(t, x))$, $\bar{H} = H(x, D\bar{u}(x))$ and $D\bar{H} = D_p H(x, D\bar{u}(x))$. The first two terms being nonnegative, we get the desired result by letting $\epsilon \rightarrow 0$. \square

Proof of Theorem 3.1. Recall the notations $v^T(s, x) = u^T(sT, x)$ and $\nu^T(s, x) = m^T(sT, x)$ for $(s, x) \in [0, 1] \times \mathbb{T}^d$. According to Lemma 3.2, we have

$$\int_0^1 \int_{\mathbb{T}^d} (F(x, \nu^T(s)) - F(x, \bar{m})) d(\nu^T(s) - \bar{m}) \leq \frac{C}{T}.$$

From assumption (6) this implies that

$$\int_0^1 \int_{\mathbb{T}^d} (F(x, \nu^T(s)) - F(x, \bar{m}))^2 dx ds \leq \frac{C}{T},$$

and, using the uniform regularity of the map F and Hölder inequality,

$$\begin{aligned} \int_0^1 \|F(\cdot, \nu^T(s)) - F(\cdot, \bar{m})\|_\infty ds & \leq C \int_0^1 \|F(\cdot, \nu^T(s)) - F(\cdot, \bar{m})\|_2 ds \\ & \leq \frac{C}{T^{\frac{1}{2}}}. \end{aligned} \tag{17}$$

Note that the map v^T solves

$$-\frac{\partial_s v^T}{T} + H(x, Dv^T) = F(x, \nu^T), \quad v^T(1, x) = u^f(x), \tag{18}$$

while the map $w^T(s, x) = \bar{u}(x) + T\bar{\lambda}(1 - s)$ solves

$$-\frac{\partial_s w^T}{T} + H(x, Dw^T) = F(x, \bar{m}), \quad w^T(1, x) = \bar{u}(x).$$

From standard estimates in viscosity solutions we deduce that, for any $t \in [0, 1]$,

$$\begin{aligned} \|v^T(t, \cdot) - w^T(t, \cdot)\|_\infty & \leq \|v^T(1, \cdot) - w^T(1, \cdot)\|_\infty \\ & + T \int_t^1 \|F(\cdot, \nu^T(s)) - F(\cdot, \bar{m})\|_\infty ds. \end{aligned}$$

So

$$\begin{aligned} \left\| \frac{v^T(t, \cdot)}{T} - \bar{\lambda}(1-t) \right\|_{\infty} &\leq \frac{\|u^f\|_{\infty} + 2\|\bar{u}\|_{\infty}}{T} + \int_t^T \|F(\cdot, \nu^T(s)) - F(\cdot, \bar{m})\|_{\infty} ds \\ &\leq \frac{C}{T^{\frac{1}{2}}}. \end{aligned}$$

□

4 Appendix: proof of the existence and uniqueness result

The following result is stated in [10]. For convenience of the reader we recall the main ingredients of proof.

Theorem 4.1 ([10]). *Let H and F satisfy conditions (4) and (7). Then equation (1) has a solution. If moreover the following inequality holds:*

$$\int_{\mathbb{T}^d} (F(x, m_1) - F(x, m_2)) d(m_1 - m_2) \geq 0 \quad \forall m_1, m_2 \in P(\mathbb{T}^d),$$

then the solution of (1) is unique.

Proof. The proof is based on a vanishing viscosity argument. Let $\epsilon > 0$ and $(u^{\epsilon}, m^{\epsilon})$ be the solution to

$$\begin{cases} (i) & -\partial_t u^{\epsilon} - \epsilon \Delta u^{\epsilon} + H(x, Du^{\epsilon}) = F(x, m^{\epsilon}(t)) \text{ in } (0, T) \times \mathbb{R}^d \\ (ii) & \partial_t m^{\epsilon} - \epsilon \Delta m^{\epsilon} - \operatorname{div}(m^{\epsilon} D_p H(x, Du^{\epsilon})) = 0 \text{ in } (0, T) \times \mathbb{R}^d \\ (iii) & m^T(0) = m_0, u^{\epsilon}(x, T) = u^f(x) \text{ in } \mathbb{R}^d \end{cases} \quad (19)$$

By standard regularity results for parabolic equations and fixed point arguments, it is not difficult to check that system (19) has at least one classical solution. Moreover, as u^f is of class \mathcal{C}^2 and the Hamiltonian satisfies (7), the map u^{ϵ} is semiconcave with a semiconcavity argument independent of ϵ . In particular u^{ϵ} is uniformly Lipschitz continuous. Let us now show that m^{ϵ} is uniformly bounded. For this we note that

$$\begin{aligned} \operatorname{div}(m^{\epsilon} D_p H(x, Du^{\epsilon})) &= \langle Dm^{\epsilon}, D_p H(x, Du^{\epsilon}) \rangle + m^{\epsilon} \operatorname{Tr} (D_{xp}^2 H(x, Du^{\epsilon}) + D_{pp}^2 H(x, Du^{\epsilon}) D^2 u^{\epsilon}) \\ &\leq \langle Dm^{\epsilon}, D_p H(x, Du^{\epsilon}) \rangle + C m^{\epsilon} \end{aligned}$$

because $D_{xp}^2 H(x, Du^{\epsilon})$ is bounded thanks to the regularity of H and the uniform Lipschitz continuity of u^{ϵ} , and $\operatorname{Tr} (D_{pp}^2 H(x, Du^{\epsilon}) D^2 u^{\epsilon})$ is bounded above because $D_{pp}^2 H$ is positive and u^{ϵ} is uniformly semiconcave. So m^{ϵ} is a subsolution of the transport equation

$$\partial_t m^{\epsilon} - \epsilon \Delta m^{\epsilon} - \langle Dm^{\epsilon}, D_p H(x, Du^{\epsilon}) \rangle - C m^{\epsilon} = 0 \text{ in } (0, T) \times \mathbb{R}^d.$$

By maximum principle we get $\|m^{\epsilon}\|_{\infty} \leq \|m_0\|_{\infty} e^{CT}$. Next we claim that the map $t \rightarrow m^{\epsilon}(t)$ is uniformly Hölder continuous: indeed, if we multiply (19)-(ii) by m^{ϵ} and integrate, we get:

$$\epsilon \int_0^T \int_{\mathbb{T}^d} |Dm^{\epsilon}|^2 \leq \int_{\mathbb{T}^d} |m^2(T) - m^2(0)| + \int_0^T \int_{\mathbb{T}^d} m^{\epsilon} |Dm^{\epsilon}| |D_p H(x, Du^{\epsilon})|.$$

As m^ϵ and $D_p H(x, m^\epsilon)$ are uniformly bounded, this implies that: $\epsilon \left(\int_0^T \int_{\mathbb{T}^d} |Dm^\epsilon|^2 \right)^{\frac{1}{2}} \leq C$. Then, for any smooth test function φ and for any $0 \leq t_1 \leq t_2 \leq T$, we have by (19)-(ii) that

$$\int_{\mathbb{T}^d} \varphi(m^\epsilon(t_2) - m^\epsilon(t_1)) = - \int_{t_1}^{t_2} \int_{\mathbb{T}^d} \epsilon \langle Dm^\epsilon, D\phi \rangle + m^\epsilon \langle D_p H(x, Du), D\phi \rangle \leq C(t_2 - t_1)^{\frac{1}{2}} \|D\varphi\|_\infty$$

because Du^ϵ and m^ϵ are uniformly bounded. Taking the supremum over all 1-Lipschitz map φ gives $\mathbf{d}_1(m(t_1), m(t_2)) \leq C(t_2 - t_1)$.

Because of the bounds on (u^ϵ, m^ϵ) , we can assume that (up subsequences) u^ϵ converges uniformly to some Lipschitz continuous map u . On another hand m^ϵ converges in L^∞ -weak* and in $\mathcal{C}^0([0, T], P(\mathbb{T}^d))$ to some $m \in L^\infty \cap \mathcal{C}^0([0, T], P(\mathbb{T}^d))$. In particular, $m(0) = m_0$. Using the continuity assumption (4), we also have that $F(\cdot, m^\epsilon(\cdot))$ converges uniformly to $F(\cdot, m(\cdot))$. By standard viscosity solutions arguments, we can conclude to the convergence of u^ϵ to the unique solution of

$$\begin{cases} -\partial_t u + H(x, Du) = F(x, m(t)) & \text{in } (0, T) \times \mathbb{R}^d \\ u(x, T) = u^f(x) & \text{in } \mathbb{R}^d \end{cases}$$

Next we turn to the limit of m^ϵ : for a fixed test function $\varphi \in \mathcal{C}_c^\infty((0, T) \times \mathbb{T}^d)$, we have

$$\int_0^T \int_{\mathbb{T}^d} (-\partial_t \varphi - \epsilon \Delta \varphi + \langle D_p H(x, Du_n), D\varphi(t, x) \rangle) m_n(t, x) = 0$$

where $D_p H(x, Du_n)$ is bounded and converges a.e. to $D_p H(x, Du)$ while m^ϵ converges weakly* to m . So we get as $\epsilon \rightarrow +\infty$,

$$\int_0^T \int_{\mathbb{T}^d} (-\partial_t \varphi(t, x) + \langle D_p H(x, Du), D\varphi(t, x) \rangle) m(t, x) = 0,$$

which shows that m is a solution of the continuity equation (1)-(ii). In conclusion, the pair (u, m) solves (1). The uniqueness for this system is established in full details in [10], so we omit the proof. \square

We complete the paper by a standard estimate on the continuity equation:

$$\partial_t m + \operatorname{div}(mb) = 0 \text{ in } (0, T) \times \mathbb{T}^d \quad (20)$$

Lemma 4.2. *Assume that $b : (0, T) \times \mathbb{T}^d \rightarrow \mathbb{R}^d$ is a Borel vector field with $\|b\|_\infty < +\infty$. If m satisfies (20), then m is Lipschitz continuous as a map from $[0, T]$ to $P(\mathbb{T}^d)$, with a Lipschitz constant bounded above by $\|b\|_\infty$.*

Proof. Fix $0 < t_1 < t_2 < T$ and let $h \in \mathcal{C}_c^\infty(\mathbb{T}^d)$ be 1-Lipschitz continuous. Let $\epsilon > 0$ small and

$$\varphi_\epsilon(t, x) = \begin{cases} (t - t_1)h(x)/\epsilon & \text{if } t \in [t_1, t_1 + \epsilon] \\ h(x) & \text{if } t \in [t_1 + \epsilon, t_2 - \epsilon] \\ (t_2 - t)h(x)/\epsilon & \text{if } t \in [t_2 - \epsilon, t_2] \\ 0 & \text{otherwise} \end{cases}$$

As

$$\int_0^T \int_{\mathbb{R}^d} (-\partial_t \varphi_\epsilon + \langle m, D\varphi_\epsilon \rangle) m = 0,$$

we have

$$\int_{t_1}^{t_1+\epsilon} \int_{\mathbb{R}^d} \frac{hm}{\epsilon} + \int_{t_1+\epsilon}^{t_2-\epsilon} \int_{\mathbb{R}^d} \langle b, Dh \rangle m + \int_{t_2-\epsilon}^{t_2} \int_{\mathbb{R}^d} -\frac{hm}{\epsilon} = o(1).$$

Letting $\epsilon \rightarrow 0$ gives for a.e. $0 < t_1 < t_2 < T$:

$$\int_{\mathbb{R}^d} h d(m(t_1) - m(t_2)) + \int_{t_1}^{t_2} \int_{\mathbb{R}^d} \langle b, Dh \rangle m = 0.$$

So

$$\int_{\mathbb{R}^d} h d(m(t_1) - m(t_2)) \leq \|b\|_{\infty} \|Dh\|_{\infty} |t_2 - t_1|.$$

Taking the sup over h gives then:

$$\mathbf{d}_1(m(t_1), m(t_2)) \leq \|b\|_{\infty} |t_2 - t_1|.$$

□

References

- [1] Cardaliaguet P., Lasry J.-M., Lions P.-L., Porretta A. *Long time average of mean field games*. Networks and Heterogeneous Media 7 (2012), no. 2, 279-301.
- [2] Cardaliaguet P., Lasry J.-M., Lions P.-L., Porretta A. *Long time average of mean field games with a nonlocal coupling*. Preprint.
- [3] Fathi A. *Théorème KAM faible et théorie de Mather sur les systèmes lagrangiens*. C. R. Acad. Sci. Paris Sér. I Math., 324(9):1043–1046, 1997.
- [4] Fathi A. *Solutions KAM faibles conjuguées et barrières de Peierls*. C. R. Acad. Sci. Paris Sér. I Math., 325(6):649–652, 1997.
- [5] Fathi A. *Weak KAM Theorem and Lagrangian Dynamics*. Cambridge University Press, to appear.
- [6] Gomes, D.A., Mohr, J., Souza, R. *Discrete time, finite state space mean field games*. J. Math. Pures Appl. (9) 93 (2010), no. 3, 308-328.
- [7] Huang, M., Caines, P.E., Malhamé, R.P. (2006). *Large population stochastic dynamic games: closed-loop McKean-Vlasov systems and the Nash certainty equivalence principle*. Communication in information and systems. Vol. 6, No. 3, pp. 221-252.
- [8] Lasry, J.-M., Lions, P.-L. *Jeux à champ moyen. I. Le cas stationnaire*. C. R. Math. Acad. Sci. Paris 343 (2006), no. 9, 619-625.
- [9] Lasry, J.-M., Lions, P.-L. *Jeux à champ moyen. II. Horizon fini et contrôle optimal*. C. R. Math. Acad. Sci. Paris 343 (2006), no. 10, 679-684.
- [10] Lasry, J.-M., Lions, P.-L. *Mean field games*. Jpn. J. Math. 2 (2007), no. 1, 229–260.
- [11] Lions, P.L. In Cours au Collège de France. www.college-de-france.fr.
- [12] Lions, P.L. (1982) Generalized solution of Hamilton-Jacobi equations. Pitman, London.

- [13] Lions P.-L., Papanicolau G. and Varadhan S.R.S. *Homogenization of Hamilton-Jacobi equation*. Unpublished preprint, 1987.
- [14] Namah G. and Roquejoffre J.-M. *Remarks on the long time behaviour of the solutions of Hamilton-Jacobi equations*. Comm. Partial Differential Equations, 24 (5-6) (1999), 883-893.
- [15] Roquejoffre J.-M. *Comportement asymptotique des solutions d'équations de Hamilton-Jacobi monodimensionnelles*. C. R. Acad. Sci. Paris Sér. I Math., 326 (12) (1998), 185-189.