TRANSSERIES AND TODOROV-VERNAEVE'S ASYMPTOTIC FIELDS

MATTHIAS ASCHENBRENNER AND ISAAC GOLDBRING

ABSTRACT. We study the relationship between fields of transseries and residue fields of convex subrings of non-standard extensions of the real numbers. This was motivated by a question of Todorov and Vernaeve, answered in this paper.

In this note we answer a question by Todorov and Vernaeve (see, e.g., [35]) concerning the relationship between the field of logarithmic-exponential series from [14] and the residue field of a certain convex subring of a non-standard extension of the real numbers, introduced in [34] in connection with a non-standard approach to Colombeau's theory of generalized functions. The answer to this question can almost immediately be deduced from well-known (but non-trivial) results about o-minimal structures. It should therefore certainly be familiar to logicians working in this area, but perhaps less so to those in non-standard analysis, and hence may be worth recording.

We begin by explaining the question of Todorov-Vernaeve. Let ${}^*\mathbb{R}$ be a non-standard extension of \mathbb{R} . Given $X \subseteq \mathbb{R}^m$ we denote the non-standard extension of X by *X , and given also a map $f \colon X \to \mathbb{R}^n$, by abuse of notation we denote the non-standard extension of f to a map ${}^*X \to {}^*\mathbb{R}^n$ by the same symbol f.

Let $\mathcal O$ be a convex subring of ${}^*\mathbb R$. Then $\mathcal O$ is a valuation ring of ${}^*\mathbb R$, with maximal ideal

$$o := \{ x \in {}^*\mathbb{R} : x = 0, \text{ or } x \neq 0 \text{ and } x^{-1} \notin \mathcal{O} \}.$$

We denote the residue field \mathcal{O}/\mathfrak{o} of \mathcal{O} by $\widehat{\mathcal{O}}$, with natural surjective morphism

$$x \mapsto \widehat{x} := x + \rho \colon \mathcal{O} \to \widehat{\mathcal{O}}.$$

The ordering of ${}^*\mathbb{R}$ induces an ordering of \widehat{O} making \widehat{O} an ordered field and $x \mapsto \widehat{x}$ order-preserving. By standard facts from real algebra [26], \widehat{O} is real closed. Residue fields of convex subrings of ${}^*\mathbb{R}$ are called "asymptotic fields" in [34] (although this terminology is already used with a different meaning elsewhere [2]). The collection of convex subrings of ${}^*\mathbb{R}$ is linearly ordered by inclusion, and the smallest convex subring of ${}^*\mathbb{R}$ is

$${}^*\mathbb{R}_{\text{fin}} = \{x \in {}^*\mathbb{R} : |x| \leqslant n \text{ for some } n\},\$$

with maximal ideal

$${}^*\mathbb{R}_{\inf} = \left\{ x \in {}^*\mathbb{R} : |x| \leqslant \frac{1}{n} \text{ for all } n > 0 \right\}.$$

The inclusions $\mathbb{R} \to {}^*\mathbb{R}_{\text{fin}} \to \mathcal{O}$ give rise to a field embedding $\mathbb{R} \to \widehat{\mathcal{O}}$, by which we identify \mathbb{R} with a subfield of $\widehat{\mathcal{O}}$. In the case $\mathcal{O} = {}^*\mathbb{R}_{\text{fin}}$ we have $\widehat{\mathcal{O}} = \mathbb{R}$, and \widehat{x} is the standard part of $x \in {}^*\mathbb{R}_{\text{fin}}$, also denoted in the following by $\operatorname{st}(x)$.

 $\rm May~2012$

Let now $\xi \in {}^*\mathbb{R}$ with $\xi > \mathbb{R}$ and let \mathcal{E} be the smallest convex subring of ${}^*\mathbb{R}$ containing all iterated exponentials $\xi, \exp \xi, \exp \xi, \dots$ of ξ , that is,

$$\mathcal{E} = \{ x \in {}^*\mathbb{R} : |x| \leqslant \exp_n(\xi) \text{ for some } n \},$$

where $\exp_0(\xi) = \xi$ and $\exp_n(\xi) = \exp(\exp_{n-1}(\xi))$ for n > 0. The maximal ideal of \mathcal{E} is

$$e = \left\{ x \in {}^*\mathbb{R} : |x| \leqslant \frac{1}{\exp_n(\xi)} \text{ for all } n \right\},$$

with residue field $\widehat{\mathcal{E}} = \mathcal{E}/e$. Note that the definition of $\widehat{\mathcal{E}}$ depends on the choice of ${}^*\mathbb{R}$ and ξ , which is suppressed in our notation; in [34, 35], $\widehat{\mathcal{E}}$ is denoted by $\widehat{\mathcal{E}}_{\varrho}$ where $\varrho = 1/\xi$.

By an **exponential field** we mean an ordered field K equipped with an exponential function on K, i.e., an isomorphism $f \mapsto \exp(f)$ between the ordered additive group of K and the ordered multiplicative group $K^{>0}$ of positive elements of K. We often write e^f instead of $\exp(f)$, and the inverse of \exp is denoted by $\log \colon K^{>0} \to K$. It is shown in [35] (see also Section 4 below) that there are mutually inverse group morphisms $\exp \colon \widehat{\mathcal{L}} \to \widehat{\mathcal{L}}^{>0}$ and $\log \colon \widehat{\mathcal{L}}^{>0} \to \widehat{\mathcal{L}}$ with

$$\exp(\widehat{f}) = \widehat{\exp(f)}, \quad \log(\widehat{g}) = \widehat{\log(g)} \qquad \text{for all } f,g \in \mathcal{E}, \, \widehat{g} > 0.$$

Thus $(\widehat{\mathcal{L}}, \exp)$ is an exponential field containing the real exponential field (\mathbb{R}, \exp) as an exponential subfield.

Let now $\mathbb{R}[[x^{\mathbb{Z}}]]$ be the field of Laurent series in descending powers of x with coefficients in \mathbb{R} , made into an ordered field such that $x > \mathbb{R}$. Then $\mathbb{R}[[x^{\mathbb{Z}}]]$ does not admit an exponential function, but one can embed $\mathbb{R}[[x^{\mathbb{Z}}]]$ into a large real closed ordered field, whose elements are formal series (with monomials coming from some ordered abelian group extending $x^{\mathbb{Z}}$), which does carry an exponential function extending the exponential function on \mathbb{R} . There are several ways of performing such a construction, leading to different (non-isomorphic) exponential fields of **transseries**. One such construction leads to the exponential field $\mathbb{R}[[x^{\mathbb{R}}]]^{\mathrm{LE}}$ of logarithmic-exponential series (or LE-series), introduced in [5, 16] and further studied in [14, 24, 25]. Another construction results in a properly larger exponential field, first defined in [24] (and also employed in [3]), which is sometimes called the field of exponential-logarithmic series (or EL-series) and denoted here by \mathbb{T} .

The following was asked in [35]:

Question. Is there an embedding $\mathbb{R}[[x^{\mathbb{R}}]]^{LE} \to \widehat{\mathcal{E}}$ (of fields) which is the identity on \mathbb{R} ?

Our aim in this note is to give a positive answer to this question, under a sensible extra hypothesis on ${}^*\mathbb{R}$; in fact, we show that then the structure on $\widehat{\mathcal{E}}$ is even richer than suggested by the question above:

Theorem 0.1. Suppose ${}^*\mathbb{R}$ is \aleph_1 -saturated. Then there exists an elementary embedding $\mathbb{T} \to \widehat{\mathcal{E}}$ of exponential fields which is the identity on \mathbb{R} and sends x to ξ .

Note that the hypothesis on $*\mathbb{R}$ is satisfied automatically if $*\mathbb{R}$ is obtained (as usual) as an ultrapower of \mathbb{R} . Theorem 0.1 is a special case of a general embedding result stated and proved in Section 3 below. As a consequence of this result, the embedding $\mathbb{T} \to \widehat{\mathcal{E}}$ in Theorem 0.1 can be chosen to additionally respect the natural

structure on \mathbb{T} , respectively $\widehat{\mathcal{E}}$, coming from the restricted real analytic functions; see Theorem 3.7. This embedding can further be extended to an embedding, also respecting the restricted analytic structure, of the maximal immediate extension of \mathbb{T} into $\widehat{\mathcal{E}}$ (where \mathbb{T} is equipped with the valuation whose valuation ring is the convex hull of \mathbb{R} in \mathbb{T}); see Corollary 3.8. We should mention that fields of transseries, even larger than \mathbb{T} , are constructed in [24, 33], which also carry exponential functions which (presumably) make them elementary extensions of (\mathbb{R} , exp) of countable character (see Section 2.1), so the embedding theorem in Section 3 could be applied to produce embeddings of such fields of transseries into $\widehat{\mathcal{E}}$. (For another application of our embedding result, to surreal numbers, see Corollary 3.9 below.)

The ordered field \mathbb{T} has more structure than that dealt with in Theorem 3.7. For example, it comes equipped with a natural derivation $\frac{d}{dx}$ making it an H-field in the sense of [1] (see, e.g., [3]). By Lemma 2.6 and the remarks following Lemma 3.3 below, (\mathbb{R}, \exp) is an elementary substructure of both $(\widehat{\mathcal{L}}, \exp)$ and (\mathbb{T}, \exp) , so if $\widehat{\mathcal{L}} \neq \mathbb{R}$ then also $(\widehat{\mathcal{L}}, \exp, \mathbb{R}) \equiv (\mathbb{T}, \exp, \mathbb{R})$ by [11]. Thus, if the exponential field $(\widehat{\mathcal{L}}, \exp)$ equipped with a predicate for its subfield \mathbb{R} happens to be splendid in the sense of [23, Section 10.1] (e.g., if it is saturated), then there also exists a derivation ∂ on $\widehat{\mathcal{L}}$ making this ordered field an H-field with constant field \mathbb{R} , such that $(\widehat{\mathcal{L}}, \exp, \mathbb{R}, \partial)$ is elementarily equivalent to $(\mathbb{T}, \exp, \mathbb{R}, \frac{d}{dx})$. This raises the following question, not addressed in the present paper:

Question. Suppose ${}^*\mathbb{R}$ is \aleph_1 -saturated. Is there a derivation on $\widehat{\mathcal{L}}$ making this ordered field an H-field with constant field \mathbb{R} , and an elementary embedding $\mathbb{T} \to \widehat{\mathcal{L}}$ of exponential differential fields which is the identity on \mathbb{R} and sends x to ξ ?

Organization of the paper. We begin by recalling the construction of the exponential field \mathbb{T} in Section 1. In Section 2 we then show an embedding statement (Lemma 2.4) which is used in the following Section 3 to prove the main theorem stated above. Section 4 finally contains some remarks on the functions on residue fields of convex subrings of $*\mathbb{R}$ induced by real C^{∞} -functions.

Conventions and notations. We let m and n range over the set $\mathbb{N} := \{0, 1, 2, \dots\}$ of natural numbers. We use "ordered set" synonymously with "linearly ordered set." Let S be an ordered set. We let $S_{\pm\infty}$ be the set $S \cup \{+\infty, -\infty\}$, where $+\infty$ and $-\infty$ are distinct and not in S, equipped with the extension of the ordering on S to $S_{\pm\infty}$ satisfying $-\infty < s < +\infty$ for all $s \in S$, and we also let S_{∞} be the ordered subset $S \cup \{+\infty\}$ of $S_{\pm\infty}$. An interval in S is a subset of S of the form

$$(a,b) = \{x \in S : a < x < b\}$$
 $(a,b \in S_{\pm \infty}, a < b).$

If S' is an ordered set extending S, then for $a, b \in S_{\pm \infty}$ with a < b we also let

$$(a,b)_{S'} := \{x \in S' : a < x < b\}.$$

We equip S with the order topology, with a subbasis of open sets given by the intervals, and we equip each cartesian power S^n of S with the corresponding product topology.

Acknowledgements. The first-named author was partially supported by NSF grant DMS-0969642 and the second-named author was partially supported by NSF grant DMS-1007144. Both authors would like to thank Dave Marker for the argument in the example in Section 4.

1. Constructing \mathbb{T}

In this section we explain the construction of the transseries field \mathbb{T} .

1.1. Well-based series. Let \mathfrak{M} be an ordered abelian group, written multiplicatively, with identity 1. We refer to the elements of \mathfrak{M} as monomials, write the ordering on \mathfrak{M} as \preccurlyeq , and put $\mathfrak{m} \prec \mathfrak{n}$ if $\mathfrak{m} \preccurlyeq \mathfrak{n}$ and $\mathfrak{m} \neq \mathfrak{n}$, for $\mathfrak{m}, \mathfrak{n} \in \mathfrak{M}$. Let C be a field. A well-based series with coefficients in C and monomials from \mathfrak{M} is a mapping $f: \mathfrak{M} \to C$ whose support

$$\operatorname{supp} f := \left\{ \mathfrak{m} \in \mathfrak{M} : f(\mathfrak{m}) \neq 0 \right\}$$

is **well-based**, that is, there exists no infinite sequence $\mathfrak{m}_1, \mathfrak{m}_2, \ldots$ of monomials in supp f with $\mathfrak{m}_1 \prec \mathfrak{m}_2 \prec \cdots$. We put $f_{\mathfrak{m}} = f(\mathfrak{m})$, and we usually write f as a formal sum

$$f=\sum_{\mathfrak{m}\in\mathfrak{M}}f_{\mathfrak{m}}\mathfrak{m}.$$

We denote the set of well-based series with coefficients in C and monomials from \mathfrak{M} by $C[[\mathfrak{M}]]$. It was first noted by Hahn (1907) that $C[[\mathfrak{M}]]$ is a field with respect to the natural addition and multiplication of well-based series:

$$f+g=\sum_{\mathfrak{m}\in\mathfrak{M}}(f_{\mathfrak{m}}+g_{\mathfrak{m}})\,\mathfrak{m}, \qquad f\cdot g=\sum_{\mathfrak{m}\in\mathfrak{M}}\left(\sum_{\mathfrak{u}\cdot\mathfrak{v}=\mathfrak{m}}f_{\mathfrak{u}}\cdot g_{\mathfrak{v}}\right)\mathfrak{m}.$$

We call $C[[\mathfrak{M}]]$ the **field of well-based series** (or the **Hahn field**) with coefficients in C and monomials from \mathfrak{M} . It contains C as a subfield, identifying $c \in C$ with the series $f \in C[[\mathfrak{M}]]$ such that $f_1 = c$ and $f_{\mathfrak{m}} = 0$ for $\mathfrak{m} \neq 1$. Given $f \in C[[\mathfrak{M}]]$ we call $f_1 \in C$ the **constant term** of f.

Example. Let R be an ordered subgroup of the ordered additive group \mathbb{R} , and let $\mathfrak{M}=x^R$ be a multiplicative copy of R, with order-preserving isomorphism $r\mapsto x^r\colon R\to x^R$. Then $C[[\mathfrak{M}]]=C[[x^R]]$ is a Hahn field with coefficients in C and monomials of the form $x^r,\,r\in R$. Taking $R=\mathbb{Z}$ we obtain the field of formal Laurent series in descending powers of x with coefficients in C.

The support of any non-zero $f \in C[[\mathfrak{M}]]$, being well-based, has a maximal element (with respect to \leq), called the **dominant monomial**

$$\mathfrak{d}(f) = \max \operatorname{supp} f$$

of f. We also set $\mathfrak{d}(0) := 0$, and extend \leq to a linear ordering on $\{0\} \cup \mathfrak{M}$ by declaring $0 \leq \mathfrak{m}$ for $\mathfrak{m} \in \mathfrak{M}$. This linear ordering is extended to a binary relation on $C[[\mathfrak{M}]]$ by

$$f \preccurlyeq g : \iff \mathfrak{d}(f) \preccurlyeq \mathfrak{d}(g).$$

Every $f \in C[[\mathfrak{M}]]$ can be decomposed as

$$f = f^{\succ} + f_1 + f^{\prec}$$

where $f_1 \in C$ is the coefficient of $1 \in \mathfrak{M}$ in f, and

$$f^{\succ} = \sum_{\mathfrak{m} \succ 1} f_{\mathfrak{m}} \mathfrak{m}$$
 (infinite part of f),
 $f^{\prec} = \sum_{\mathfrak{m} \prec 1} f_{\mathfrak{m}} \mathfrak{m}$ (infinitesimal part of f).

This gives rise to a decomposition of $C[[\mathfrak{M}]]$ into a direct sum of C-vector spaces:

$$C[[\mathfrak{M}]] = C[[\mathfrak{M}]]^{\succ} \oplus C \oplus C[[\mathfrak{M}]]^{\prec}$$

where

$$C[[\mathfrak{M}]]^{\succ} := \big\{ f \in C[[\mathfrak{M}]] : \mathfrak{m} \succ 1 \text{ for all } \mathfrak{m} \in \operatorname{supp} f \big\},$$

$$C[[\mathfrak{M}]]^{\prec} := \big\{ f \in C[[\mathfrak{M}]] : \mathfrak{m} \prec 1 \text{ for all } \mathfrak{m} \in \operatorname{supp} f \big\}.$$

Let Γ be an additively written copy of \mathfrak{M} , with isomorphism $\mathfrak{m} \mapsto v\mathfrak{m} \colon \mathfrak{M} \to \Gamma$, equipped with the ordering \leqslant making this isomorphism decreasing: $\mathfrak{m} \leqslant \mathfrak{n} \iff v\mathfrak{m} \geqslant v\mathfrak{n}$, for all $\mathfrak{m}, \mathfrak{n} \in \mathfrak{M}$. Then the map

$$v \colon C[[\mathfrak{M}]] \to \Gamma_{\infty}, \qquad vf = \begin{cases} v(\mathfrak{d}(f)) & \text{if } f \neq 0 \\ \infty & \text{if } f = 0 \end{cases}$$

is a valuation on the field $C[[\mathfrak{M}]]$, with valuation ring $\mathcal{O} := C[[\mathfrak{M}]]^{\prec} = C \oplus C[[\mathfrak{M}]]^{\prec}$, whose maximal ideal is $\sigma := C[[\mathfrak{M}]]^{\prec}$. The map sending $f \in \mathcal{O}$ to its constant term f_1 is a surjective ring morphism $\mathcal{O} \to C$, with kernel σ , and hence induces an isomorphism $\mathcal{O}/\sigma \to C$ between the residue field of \mathcal{O} and the coefficient field C.

1.2. Ordering of well-based series. In the following we are mainly interested in the case where C is equipped with an ordering making C an ordered field. Then we make $C[[\mathfrak{M}]]$ into an ordered field extension of C as follows: for $0 \neq f \in C[[\mathfrak{M}]]$ define

$$f > 0 \quad :\iff \quad f_{\mathfrak{d}(f)} > 0.$$

Note that for $f, g \in C[[\mathfrak{M}]]$,

$$f \preccurlyeq g \iff |f| \leqslant c|g| \text{ for some } c \in C^{>0}.$$

The valuation ring \mathcal{O} of $C[[\mathfrak{M}]]$ is a convex subring of $C[[\mathfrak{M}]]$.

Suppose \mathfrak{M}_1 and \mathfrak{M}_2 are subgroups of \mathfrak{M} with \mathfrak{M}_1 convex in \mathfrak{M} , $\mathfrak{M} = \mathfrak{M}_1 \cdot \mathfrak{M}_2$, and $\mathfrak{M}_1 \cap \mathfrak{M}_2 = \{1\}$. Then we have an ordered field isomorphism

$$f = \sum_{\mathfrak{m} \in \mathfrak{M}} f_{\mathfrak{m}} \mathfrak{m} \mapsto \sum_{\mathfrak{m}_2 \in \mathfrak{M}_2} \left(\sum_{\mathfrak{m}_1 \in \mathfrak{M}_1} f_{\mathfrak{m}_1 \mathfrak{m}_2} \mathfrak{m}_1 \right) \mathfrak{m}_2 \colon \quad C[[\mathfrak{M}]] \to (C[[\mathfrak{M}_1]])[[\mathfrak{M}_2]]$$

which is the identity on C. We identify $C[[\mathfrak{M}]]$ and $(C[[\mathfrak{M}_1]])[[\mathfrak{M}_2]]$ via this isomorphism whenever convenient.

1.3. Analytic structure on Hahn fields. Let \mathfrak{M} be a multiplicatively written ordered abelian group and $K = \mathbb{R}[[\mathfrak{M}]]$. Every analytic function $f: (a,b) \to \mathbb{R}$ on an interval, where $a,b \in \mathbb{R}_{\pm \infty}$, a < b, extends naturally to $\widehat{f}: (a,b)_K \to K$, where

$$(a,b)_K = \big\{g \in K : a < g < b\big\} = \big\{c + \varepsilon : c \in (a,b), \ \varepsilon \in K^{\prec 1}\big\},$$

given by

$$\widehat{f}(c+\varepsilon) := \sum_{n=0}^{\infty} \frac{f^{(n)}(c)}{n!} \varepsilon^n \qquad \text{for } c \in (a,b) \text{ and } \varepsilon \in K^{\prec 1}.$$

For this, one needs to show that the infinite sum on the right-hand side makes sense in $K = \mathbb{R}[[\mathfrak{M}]]$; see, e.g., [14, p. 64]. For example, the real exponential function $c \mapsto e^c \colon \mathbb{R} \to \mathbb{R}^{>0}$ extends to the function

$$c + \varepsilon \mapsto \exp(c + \varepsilon) = e^{c + \varepsilon} := e^c \cdot \sum_{n=0}^{\infty} \frac{\varepsilon^n}{n!}$$
 $(c \in \mathbb{R} \text{ and } \varepsilon \in K^{\prec 1})$ (1)

from $K^{\leq 1} = \mathbb{R} \oplus K^{\leq 1}$ to $K^{>0}$. For $f, g \in K^{\leq 1}$, we have

$$\exp(f) \geqslant 1 \Leftrightarrow f \geqslant 0$$
, $\exp(f) \geqslant f + 1$, and $\exp(f + g) = \exp(f) \exp(g)$.

Thus exp is injective with image

$$K^{>0,\asymp 1} = \big\{g \in K : g > 0, \ \mathfrak{d}(g) = 1\big\}$$

and inverse

$$\log: K^{>0, \approx 1} \to K^{\leqslant 1}$$

given by

$$\log g := \log c + \log(1+\varepsilon)$$
 for $g = c(1+\varepsilon), c \in \mathbb{R}^{>0}, \varepsilon \prec 1$,

where $\log c$ is the usual natural logarithm of the positive real number c and

$$\log(1+\varepsilon) := \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n} \varepsilon^n.$$

In a similar way, every real-valued function $f: I^n \to \mathbb{R}^n$, analytic on an open neighborhood of the cube $I^n = [-1,1]^n$ in \mathbb{R}^n , extends naturally to a function \widehat{f} on the corresponding cube $I(K)^n$ in K^n (taking values in K).

One drawback of $K = \mathbb{R}[[\mathfrak{M}]]$ is that this ordered field does not support a (total) exponential function if $\mathfrak{M} \neq \{1\}$, as shown in [28]. Nonetheless, one can extend $x^{\mathbb{Z}}$ to a large ordered multiplicative group \mathfrak{T} and $\mathbb{R}[[x^{\mathbb{Z}}]]$ to a real closed *subfield* \mathbb{T} of the well-based series field $\mathbb{R}[[\mathfrak{T}]]$, such that $\mathfrak{T} \subseteq \mathbb{T}^{>0}$ inside $\mathbb{R}[[\mathfrak{T}]]$, and such that the usual exponential function on \mathbb{R} extends to an exponential function on \mathbb{T} . In the following we outline the construction of such an exponential field \mathbb{T} .

1.4. The exponential field \mathbb{T} . We begin by introducing the multiplicative group \mathfrak{L} of logarithmic monomials: this is the multiplicatively written vector space

$$\mathfrak{L} = \left\{ \ell_0^{\alpha_0} \ell_1^{\alpha_1} \cdots \ell_n^{\alpha_n} : (\alpha_0, \dots, \alpha_n) \in \mathbb{R}^{n+1}, \ n \in \mathbb{N} \right\}$$

over the field \mathbb{R} with basis $(\ell_n)_{n\geqslant 0}$. We equip \mathfrak{L} with the unique ordering \preccurlyeq making it an ordered vector space over the ordered field \mathbb{R} such that $\ell_n \succ \ell_{n+1}^m$ for each m. We let $\mathbb{L} := \mathbb{R}[[\mathfrak{L}]]$ be the ordered field of **logarithmic transseries** and let $\exp: \mathbb{L}^{\preccurlyeq} \to \mathbb{L}^{>0, \asymp 1}$ be as defined in (1). Note that the inverse $\log: \mathbb{L}^{>0, \asymp 1} \to \mathbb{L}^{\preccurlyeq}$ of exp extends to a strictly increasing group homomorphism $\log: \mathbb{L}^{>0} \to \mathbb{L}$ by

$$\log(\mathfrak{m} \cdot g) := \log \mathfrak{m} + \log g \qquad \text{for } \mathfrak{m} \in \mathfrak{L}, \, g \in \mathbb{L}^{>0, \times 1}, \tag{2}$$

where for $\mathfrak{m} = \ell_0^{\alpha_0} \ell_1^{\alpha_1} \cdots \ell_n^{\alpha_n} \in \mathfrak{L}, (\alpha_0, \ldots, \alpha_n) \in \mathbb{R}^n$, we set

$$\log \mathfrak{m} := \sum_{i=0}^{n} \alpha_{i} \ell_{i+1} \in \mathbb{R}[\mathfrak{L}].$$

So writing $x := \ell_0$, we have $\ell_1 = \log x, \ell_2 = \log \log x, \ldots$, and in general, ℓ_n is the nth iterated logarithm of x.

Now let $\mathfrak{T}_0 := \mathfrak{L}$, $K_0 := \mathbb{L}$. Since there is no reasonable way to define $\exp f$ as an element of K_0 for $f \in K_0^{\succ}$, we enlarge $K_0 = \mathbb{R}[[\mathfrak{T}_0]]$ to a bigger series field $K_1 = \mathbb{R}[[\mathfrak{T}_1]]$ such that $\exp f \in K_1$ for all $f \in K_0^{\succ}$: simply take a multiplicative copy $\exp(K_0^{\succ})$ of the ordered additive subgroup K_0^{\succ} of K_0 , with order-preserving isomorphism

$$f \mapsto \exp f \colon K_0^{\succ} \to \exp(K_0^{\succ}),$$

and form the direct product of multiplicative groups

$$\mathfrak{T}_1 := \exp(K_0^{\succ}) \cdot \mathfrak{T}_0.$$

Order \mathfrak{T}_1 lexicographically: for $f \in K_0^{\succ}$, $\mathfrak{m} \in \mathfrak{T}_0$, put

$$\exp(f) \cdot \mathfrak{m} \geq 1 \qquad \iff \qquad f > 0 \text{ or } (f = 0 \text{ and } \mathfrak{m} \geq 1 \text{ in } \mathfrak{T}_0).$$

The natural identification of \mathfrak{T}_0 with an ordered subgroup of \mathfrak{T}_1 makes K_0 an ordered subfield of K_1 . Define $\exp g \in K_1^{>0}$ for $g \in K_0$ by

$$\exp(f + c + \varepsilon) := \exp(f) \cdot e^{c + \varepsilon} \qquad (f \in K_0^{\succ}, \ c \in \mathbb{R}, \ \varepsilon \in K_0^{\prec}),$$

with $e^{c+\varepsilon}$ as in (1). Now K_1 has the same defect as K_0 : there is no reasonable way to define $\exp f$ as an element of K_1 for $f \in K_1$ with $\mathfrak{d}(f) \succ \mathfrak{T}_0$. In order to add the exponentials of such elements to K_1 , enlarge K_1 to a field K_2 just as K_0 was enlarged to K_1 . More generally, consider a tuple (K, A, B, \log) where

- (1) K is an ordered field;
- (2) A and B are additive subgroups of K with $K = A \oplus B$ and B convex in K;
- (3) $\log \colon K^{>0} \to K$ is a strictly increasing homomorphism (the inverse of which we denote by $\exp \colon \log(K^{>0}) \to K$) such that $B \subseteq \log(K^{>0})$.

We call such a quadruple (K, A, B, \log) a **pre-logarithmic ordered field**. So (K_0, A_0, B_0, \log_0) with $A_0 := K_0^{\succ}$, $B_0 := K_0^{\preccurlyeq}$ and $\log_0 \colon K_0^{>0} \to K_0$ given by (2) is a pre-logarithmic ordered field. Given a pre-logarithmic ordered field (K, A, B, \log) , define a pre-logarithmic ordered field (K', A', B', \log') as follows: Take a multiplicative copy $\exp(A)$ of the ordered additive group A with order-preserving isomorphism

$$\exp_A : A \to \exp(A),$$

and put

$$K' := K[[\exp(A)]], \quad A' := (K')^{\succ}, \quad B' := K^{\preccurlyeq} = K \oplus (K')^{\prec},$$

and define $\log' : (K')^{>0} \to K'$ by

$$\log' (f \cdot \exp_A(a) \cdot (1+\varepsilon)) := \log f + a + \log(1+\varepsilon)$$

for $f \in K$, $a \in A$, $\varepsilon \in (K')^{\prec}$. Note that \log' extends \log and $K \subseteq \log'((K')^{>0})$. Moreover, if $K = \mathbb{R}[[\mathfrak{M}]]$ for some multiplicative ordered abelian group \mathfrak{M} , then $K' = \mathbb{R}[[\mathfrak{M}']]$ where $\mathfrak{M}' = \exp(A) \cdot \mathfrak{M}$, ordered lexicographically. Inductively, set

$$(K_{n+1}, A_{n+1}, B_{n+1}, \log_{n+1}) := (K'_n, A'_n, B'_n, \log'_n).$$

Then $K_n = \mathbb{R}[[\mathfrak{T}_n]]$ with $\mathfrak{T}_n = \mathfrak{L} \cdot \exp(A_{n-1} \oplus \cdots \oplus A_0)$, and we put

$$\mathbb{T} := \bigcup_n \mathbb{R}[[\mathfrak{T}_n]], \qquad \mathfrak{T} := \bigcup_n \mathfrak{T}_n.$$

Thus $\mathbb{T} \subseteq \mathbb{R}[[\mathfrak{T}]]$ as ordered fields. We let $\log\colon \mathbb{T}^{>0} \to \mathbb{T}$ be the common extension of all the \log_n . The map \log is a strictly increasing group isomorphism $\mathbb{T}^{>0} \to \mathbb{T}$, so its inverse $\exp\colon \mathbb{T} \to \mathbb{T}^{>0}$ is an exponential function on \mathbb{T} . It is well-known that the exponential field $\mathbb{R}[[x^\mathbb{R}]]^{\mathrm{LE}}$ of LE-series embeds into \mathbb{T} in a natural way. (See [24], and also [29].) However, this embedding is not onto, since, e.g., the series $\sum_n \frac{1}{\ell_n} = \frac{1}{\ell_0} + \frac{1}{\ell_1} + \cdots$ is an element of \mathbb{L} , but is not an LE-series (since iterated logarithms ℓ_n of arbitrary "depth" n appear in it).

2. Character, and an Embedding Result

In this section we define a cardinal invariant of an ordered set (M, <), which we call character, and which is more meaningful than the cardinality of M when one is concerned with realizing cuts in M. We demonstrate this by proving an embedding criterion for models of o-minimal theories (Lemma 2.4) used in the proof of Theorem 3.7. We then recall some important examples of o-minimal structures.

2.1. Character. Let (M, <) be an ordered set. Recall that the cofinality of (M, <), denoted by $\operatorname{cf}(M, <)$, or $\operatorname{cf}(M)$ for brevity, is the minimal cardinality of a cofinal subset of M. Recall that M always has a well-ordered cofinal subset of cardinality $\operatorname{cf}(M)$. It is also well-known that if A is an ordered subset of M which is cofinal in M, then $\operatorname{cf}(A) = \operatorname{cf}(M)$. Dually, the coinitiality of (M, <), denoted by $\operatorname{ci}(M, <) = \operatorname{ci}(M)$, is the cofinality of (M, >).

A **cut** in M is a subset C of M which is closed downward in M, i.e., if $x \in M$ satisfies x < c for some $c \in C$, then $x \in C$. Given a cut C in M, an element x in an ordered set extending M is said to **realize** the cut C if $C < x < M \setminus C$. For every element x in an ordered set extending M with $x \notin M$, x realizes the cut

$$M^{< x} = \{ c \in M : c < x \}$$

in M, which we call the **cut of** x **in** M. The **character** of a cut C in M is the pair $(\operatorname{cf}(C),\operatorname{ci}(M\setminus C))$. We define the **character** of the ordered set (M,<), denoted by $\operatorname{ch}(M,<)$ or simply by $\operatorname{ch}(M)$, to be the supremum of $\operatorname{cf}(C) + \operatorname{ci}(M\setminus C)$, as C ranges over all cuts in M. If A is an ordered subset of M then $\operatorname{ch}(A) \leq \operatorname{ch}(M)$.

Every (reverse) well-ordered subset of the ordered set of real numbers is countable, so $ch(\mathbb{R}) = \aleph_0$. The following was shown by Esterle [19] (see also [14, p. 73]):

Lemma 2.1. Let \mathfrak{M} be an ordered abelian group. Suppose each (reverse) well-ordered subset of \mathfrak{M} is countable. Then each (reverse) well-ordered subset of $\mathbb{R}[[\mathfrak{M}]]$ is countable.

From this we easily obtain:

Corollary 2.2. Each (reverse) well-ordered subset of \mathbb{T} is countable. In particular, $\operatorname{ch}(\mathbb{T}) = \aleph_0$.

Proof. If the ordered set M is the union of countably many ordered subsets, each of whose (reverse) well-ordered subsets is countable, then each (reverse) well-ordered subset of M is countable. Thus we only need to show that each (reverse) well-ordered subset of the ordered subfield $\mathbb{R}[[\mathfrak{T}_n]]$ in the construction of \mathbb{T} is countable. This follows easily by induction on n, using the lemma above.

Let α be an ordinal and (M,<) be an ordered set. Then (M,<) is called an η_{α} -set if for all subsets A, B of M with A < B and $|A \cup B| < \aleph_{\alpha}$, there is an element x of M with A < x < B. Note that if the ordered set (M,<) is dense without endpoints, then (M,<) is an η_{α} -set iff (M,<), viewed as a structure in the language $\{<\}$, is \aleph_{α} -saturated (a consequence of the theory of (M,<) admitting quantifier elimination). We note the following obvious fact:

Lemma 2.3. Let N be an η_{α} -set extending M, where $\aleph_{\alpha} > \operatorname{ch}(M)$. Then each cut in M is realized by an element of N.

2.2. An embedding criterion. In the following we let \mathcal{L} be a first-order language containing a binary relation symbol <. An \mathcal{L} -structure $\mathbf{M} = (M, <, \ldots)$ expanding a dense linearly ordered set (M, <) without endpoints is said to be **o-minimal** if each subset of M which is definable in \mathbf{M} is a finite union of intervals (a, b) (where $a, b \in M_{\pm \infty}$) and singletons $\{c\}$ $(c \in M)$. (Here and in the rest of this paper, "definable" means "definable, possibly with parameters.") By [27], every structure elementarily equivalent to an o-minimal \mathcal{L} -structure is also o-minimal, and in this case, the complete theory $\mathrm{Th}(\mathbf{M})$ of \mathbf{M} is called o-minimal. We refer to [8] for a summary of basic facts from the theory of o-minimal structures used in this note.

Let now T be a complete o-minimal \mathcal{L} -theory and let $\mathbf{M} \models T$. We recall that the o-minimality assumption implies that given an elementary extension \mathbf{N} of \mathbf{M} , there is a one-to-one correspondence between the type of an element $x \in N \setminus M$ over M and the cut of x in M. In particular, if $\kappa > |\mathcal{L}|$, then \mathbf{M} is κ -saturated iff its underlying ordered set (M, <) is κ -saturated. We also recall that given a subset A of M, there is a prime model of T over A, i.e., a model \mathbf{N} of T with $A \subseteq N$ such that each elementary map $A \to N'$, where $\mathbf{N}' \models T$, extends to an elementary embedding $\mathbf{N} \to \mathbf{N}'$. This prime model of T over A is unique up to isomorphism over A. (See [31, Theorem 5.1].) Given an elementary extension $\mathbf{N} = (N, <, \ldots)$ of \mathbf{M} and $A \subseteq N$, we let $\mathbf{M}\langle A \rangle$ denote the prime model of T over $A \cup M$, taken as an elementary substructure of \mathbf{N} (implicitly understood from context). We also write $\mathbf{M}\langle x \rangle$ instead of $\mathbf{M}\langle \{x\}\rangle$. If also $\mathbf{M} \preccurlyeq \mathbf{N}'$, and $x \in N \setminus M$ and $x' \in N' \setminus M$ realize the same cut in M, then there is a (unique) isomorphism $\mathbf{M}\langle x \rangle \to \mathbf{M}\langle x' \rangle$ which is the identity on M and sends x to x'.

We can now easily show:

Lemma 2.4. Let $\mathbf{M} = (M, <, ...)$ and $\mathbf{N} = (N, <, ...)$ be models of T and suppose that (N, <) is κ^+ -saturated, where $\kappa = \operatorname{ch}(M)$. Then every elementary embedding of an elementary substructure of \mathbf{M} into \mathbf{N} extends to an elementary embedding of \mathbf{M} into \mathbf{N} .

Proof. Let $A \leq M$ and let $h: A \to N$ be an elementary embedding. By Zorn we may assume that h has no extension to an elementary embedding of an elementary substructure of M, properly containing A, into N. Suppose $M \neq A$ and take $x \in M \setminus A$ arbitrary. By κ^+ -saturation of (N, <) the image $h(A^{< x})$ of the cut of x in A is realized in N, say by $y \in N$. (Lemma 2.3.) Then the extension of h to a map $A \cup \{x\} \to N$ with $x \mapsto y$ is elementary, and hence further extends to an elementary embedding $A(x) \to N$, contradicting the maximality of A.

Together with the existence of prime models, this yields an improvement, for ominimal theories, of the general fact that κ -saturated models of complete first-order theories are κ -universal:

Corollary 2.5. Let $N \models T$ be κ -saturated, where κ is an infinite cardinal. Then every model M of T with $\operatorname{ch}(M) < \kappa$ elementarily embeds into N.

2.3. Some examples of o-minimal expansions of the ordered field of reals. All the examples of o-minimal structures that we will need are obtained as expansions of the ordered field $\overline{\mathbb{R}} = (\mathbb{R}, <, 0, 1, +, \times)$ of real numbers in a language extending the language of ordered rings. (The structure $\overline{\mathbb{R}}$ itself is o-minimal as a consequence of Tarski's quantifier elimination theorem. Its elementary theory is axiomatized by the axioms for real closed ordered fields.) We remark that to

any such o-minimal expansion \mathbf{R} of $\overline{\mathbb{R}}$, Miller's Growth Dichotomy Theorem [30] applies: either \mathbf{R} is **polynomially bounded** (i.e., for every definable function $f: \mathbb{R} \to \mathbb{R}$ there is some n such that $|f(x)| \leq x^n$ for sufficiently large x), or the exponential function $\exp: \mathbb{R} \to \mathbb{R}$ is 0-definable in \mathbf{R} . (For example, $\overline{\mathbb{R}}$ is polynomially bounded.) One calls \mathbf{R} exponentially bounded if for every definable function $f: \mathbb{R} \to \mathbb{R}$ there is some n such that $|f(x)| \leq \exp_n(x)$ for sufficiently large x. (At present, all known o-minimal expansion of the ordered field of real numbers are exponentially bounded.) Note that polynomial boundedness and exponential boundedness are part of the elementary theory of \mathbf{R} ; this allows us to speak of an o-minimal theory extending the theory of real closed ordered fields being polynomially bounded or exponentially bounded.

- 2.3.1. The ordered field of reals with restricted analytic functions. A restricted analytic function is a function $\mathbb{R}^n \to \mathbb{R}$ which is given on $I^n = [-1,1]^n$ by a power series in n variables with real coefficients converging on a neighborhood of I^n , and which is 0 outside of I^n . The structure $\mathbb{R}_{\rm an}$ is the expansion of $\overline{\mathbb{R}}$ by the restricted analytic functions $\mathbb{R}^n \to \mathbb{R}$, for varying n. The structure $\mathbb{R}_{\rm an}$ was shown to be o-minimal in [9] and model-complete in [6]; $\mathbb{R}_{\rm an}$ is also polynomially bounded. A complete axiomatization of $T_{\rm an} := \mathrm{Th}(\mathbb{R}_{\rm an})$ is given in [12], where it is also shown that if \mathfrak{M} is a divisible ordered abelian group, then the ordered field $\mathbb{R}[[\mathfrak{M}]]$, expanded to a structure in the language $\mathcal{L}_{\rm an}$ of $\mathbb{R}_{\rm an}$ as indicated in Section 1.3 above, is an elementary extension of $\mathbb{R}_{\rm an}$. It follows that \mathbb{T} , viewed as an $\mathcal{L}_{\rm an}$ -structure in the natural way, is a model of $T_{\rm an}$ (since it is obtained as the union of the increasing elementary chain of $\mathcal{L}_{\rm an}$ -structures $\mathbb{R}[[\mathfrak{T}_n]]$) and hence an elementary extension of $\mathbb{R}_{\rm an}$.
- 2.3.2. The exponential field of reals. The ordered field of real numbers $(\overline{\mathbb{R}}, \exp)$ augmented by the real exponential function was shown to be model-complete and o-minimal by Wilkie [37].
- 2.3.3. The exponential field of reals with restricted analytic functions. This is the expansion $\mathbb{R}_{\mathrm{an},\mathrm{exp}}$ of \mathbb{R}_{an} by the real exponential function. In [15] this structure was shown to be model-complete and o-minimal, by a generalization of Wilkie's proof for $(\overline{\mathbb{R}},\mathrm{exp})$. A simpler proof, and a complete axiomatization of $T_{\mathrm{an,exp}} := \mathrm{Th}(\mathbb{R}_{\mathrm{an,exp}})$, is given in [12]. In fact, $T_{\mathrm{an,exp}}$ is axiomatized by T_{an} together with (the universal closures of) the following statements about exp:

```
(E1) \exp(x+y) = \exp(x) \cdot \exp(y);
```

- (E2) $x < y \rightarrow \exp(x) < \exp(y)$;
- (E3) $x > 0 \rightarrow \exists y \exp(y) = x;$
- $(E4_n)$ $x > n^2 \to \exp(x) > x^n$ (for each n > 0);
- (E5) $-1 \le x \le 1 \to e(x) = \exp(x)$, where e is the function symbol of \mathcal{L}_{an} which represents the restricted analytic function $e: \mathbb{R} \to \mathbb{R}$ with $e(x) = e^x$ for $x \in [-1, 1]$.

Based on this axiomatization, in [13, Corollary 2.8] it was shown that $\mathbb{R}[[x^{\mathbb{R}}]]^{LE}$ is a model of $T_{\text{an,exp}}$ (and consequently, that $T_{\text{an,exp}}$ is exponentially bounded). We observe here in a similar way:

Lemma 2.6. $\mathbb{T} \models T_{\text{an,exp}}$.

Proof. We already noted that $\mathbb{T} \models T_{an}$. Clearly the exponential function on \mathbb{T} satisfies (E1)–(E3) by construction.

We prove $(E4_n)$. Suppose n > 0 and $f \in \mathbb{T}$ is such that $f > n^2$. We need $\exp(f) > f^n$. Take m such that $f \in \mathbb{R}[[\mathfrak{T}_m]]$. Write $f = g + c + \varepsilon$, with $g \in \mathbb{R}[[\mathfrak{T}_m]]^{\succ}$, $c \in \mathbb{R}$, and $\varepsilon \in \mathbb{R}[[\mathfrak{T}_m]]^{\prec}$. We first suppose that g = 0. We then have $\mathfrak{d}(\exp(f)) = 1$ with leading coefficient e^c , while $\mathfrak{d}(f^n) = 1$ with leading coefficient c^n . Since $c \geqslant n^2$, we have $e^c > c^n$ and thus $\exp(f) > f^n$. We now suppose that $g \neq 0$. Then $\mathfrak{d}(\exp(f)) = \exp(g) \cdot \mathfrak{d}(\exp(c + \varepsilon)) \succ \mathfrak{d}(f^n)$ since $\mathfrak{d}(f^n) \in \mathfrak{T}_m$. Thus, $\exp(f) > f^n$ if and only if the leading coefficient of $\exp(f)$ is positive; however, this leading coefficient is the coefficient of $\mathfrak{d}(\exp(c + \varepsilon))$ in $\exp(c + \varepsilon)$, which is positive.

Next we prove (E5). Suppose that $f \in \mathbb{T}$ with $-1 \leq f \leq 1$. Take m such that $f \in \mathbb{R}[[\mathfrak{T}_m]]$. Then $f \in \mathbb{R}[[\mathfrak{T}_m]]^{\prec}$, say $f = c + \varepsilon$, with $c \in \mathbb{R}$ and $\varepsilon \in \mathbb{R}[[\mathfrak{T}_m]]^{\prec}$. Then $e(f) = \exp(f) = e^c \sum_{n=0}^{\infty} \frac{\varepsilon^n}{n!}$.

3. T-Convexity

In this section we let $\mathbf{R} = (\mathbb{R}, <, 0, 1, +, \times, \dots)$ be an o-minimal expansion of the ordered field of real numbers in a language \mathcal{L} extending the language of ordered rings. "Definable" will always mean "definable in \mathbf{R} ." Note that the \mathcal{L} -reduct * \mathbf{R} of * \mathbb{R} is an elementary extension of \mathbf{R} . We let $T = \operatorname{Th}(\mathbf{R})$. Given a 0-definable set $X \subseteq \mathbb{R}^n$ and $\mathbf{S} \models T$, we let $X(\mathbf{S})$ be the subset of S^n defined by the same \mathcal{L} -formula as X in \mathbf{R} . We first recall the definition of T-convexity from [11] and some fundamental facts concerning this notion, and then give the proof of the main theorem from the introduction. In the last subsection we give another application of the embedding lemma from Section 2.

3.1. Definition and basic properties of T-convex subrings. Let $f: X \to \mathbb{R}$ be a 0-definable function, where $X \subseteq \mathbb{R}^n$. We say that a convex subring O of \mathbb{R} is closed under f if $f(X(\mathbb{R}) \cap O^n) \subseteq O$. A convex subring of \mathbb{R} is called T-convex if it is closed under all 0-definable continuous functions $\mathbb{R} \to \mathbb{R}$. Although this definition only talks about one-variable functions, every T-convex subring of \mathbb{R} is automatically closed under all 0-definable continuous functions $\mathbb{R}^n \to \mathbb{R}$ for all n > 0 [11, (2.9)]. In fact:

Lemma 3.1. Every T-convex subring of \mathbb{R} is closed under all 0-definable continuous functions $X \to \mathbb{R}$ where $X \subseteq \mathbb{R}^n$ is open or closed.

Proof. By the definable version of the Tietze Extension Theorem (see [4, Lemma 6.6] or [7, Chapter 8]), each 0-definable continuous function $X \to \mathbb{R}$ on a closed set $X \subseteq \mathbb{R}^n$ has an extension to a 0-definable continuous function $\mathbb{R}^n \to \mathbb{R}$. This implies the lemma for 0-definable continuous functions whose domain is a closed subset of \mathbb{R}^n , and this in turn yields the lemma also for 0-definable continuous functions whose domain is open: if $X \subseteq \mathbb{R}^n$, $X \neq \mathbb{R}^n$, is open then $X = \bigcup_{s>0} X_s$ where

$$X_s:=\left\{x\in\mathbb{R}^n: \mathrm{dist}(x,\mathbb{R}^n\setminus X)\geqslant s\right\}$$
 is closed, for $s\in\mathbb{R}^{>0}.$

Examples. If T is polynomially bounded then every convex subring of ${}^*\mathbb{R}$ is T-convex. If T is non-polynomially bounded but exponentially bounded, then a convex subring O of ${}^*\mathbb{R}$ is T-convex iff O is closed under $\exp: \mathbb{R} \to \mathbb{R}$. In particular, the convex subring \mathcal{E} of ${}^*\mathbb{R}$, defined in the introduction, is $T_{\mathrm{an,exp}}$ -convex, since it is closed under \exp (and is, indeed, the smallest convex subring of ${}^*\mathbb{R}$ containing \mathbb{R} and the infinite element ξ which is closed under \exp).

It is an easy consequence of the Monotonicity Theorem for o-minimal structures that the convex hull in ${}^*\mathbb{R}$ of an elementary substructure of ${}^*\boldsymbol{R}$ is a T-convex subring of ${}^*\boldsymbol{R}$. A strong converse of this observation is shown in [11], stated in the theorem below. Note that if O is a convex subring of ${}^*\mathbb{R}$ and R' is a subring of O which is a field, then the composition $R' \to O \to \widehat{O}$ is injective, and this map is bijective iff $O = \mathbf{R}' + o$; in this case $o = \{x \in {}^*\mathbb{R} : |x| < (R')^{>0}\}$, so O is the convex hull of O in O in O in O in O is the convex hull of O in O in O in O in O is the convex hull of O in O in O in O is the convex hull of O in O in O in O is the convex hull of O in O in O in O is the convex hull of O in O in O is the convex hull of O in O in O is the convex hull of O in O in O is the convex hull of O in O in O in O is the convex hull of O in O in O is the convex hull of O in O in O in O in O is the convex hull of O in O in O in O in O in O is the convex hull of O in O in O in O in O in O in O is the convex hull of O in O in

Theorem 3.2 (van den Dries-Lewenberg [11]). Let O be a T-convex subring of *R . Then:

- (1) an elementary substructure \mathbf{R}' of ${}^*\mathbf{R}$ contained in O is maximal among the elementary substructures of ${}^*\mathbf{R}$ contained in O iff $O = \mathbf{R}' + o$ (and hence, by Zorn, there is some $\mathbf{R}' \leq {}^*\mathbf{R}$ with $O = \mathbf{R}' + o$):
- (2) if \mathbf{R}' and \mathbf{R}'' are both maximal with respect to being elementary substructures of ${}^*\mathbf{R}$ contained in O, then there is a unique isomorphism $h \colon \mathbf{R}' \to \mathbf{R}''$ such that $\widehat{h(x)} = \widehat{x}$ for all $x \in \mathbf{R}'$.

In the following we fix a T-convex subring O of *R . By part (1) of the theorem above, we can make \widehat{O} into a model of T as follows: take an R' which is maximal among elementary substructures of *R contained in O, and make \widehat{O} into an \mathcal{L} -structure such that the restriction of the residue map $x \mapsto \widehat{x} \colon O \to \widehat{O}$ is an isomorphism $R' \to \widehat{O}$. By part (2) of the above theorem, this expansion of \widehat{O} to a model of T is independent of the choice of R'. In the following we will always consider \widehat{O} as a model of T in this way.

The following is [11, (2.20)]:

Lemma 3.3. Let $f: \mathbb{R}^n \to \mathbb{R}$ be a 0-definable continuous function. Then for all $x, y \in O^n$ we have: $x - y \in o^n \implies f(x) - f(y) \in o$.

In particular, if f is as in the previous lemma, and \widehat{f} denotes the function $\widehat{O}^n \to \widehat{O}$ defined by the same formula in the \mathcal{L} -structure \widehat{O} as f in \mathbb{R} , then by the lemma, we have $\widehat{f}(\widehat{x}) = \widehat{f(x)}$ for all $x \in O^n$. Here and below, for n > 0 and $x = (x_1, \dots, x_n) \in O^n$ we set $\widehat{x} = (\widehat{x_1}, \dots, \widehat{x_n}) \in \widehat{O}^n$.

It was noted in [34, 35] that (as a consequence of the Mean Value Theorem) the exponential function on ${}^*\mathbb{R}$ induces a function on $\widehat{\mathcal{E}}$; that is, there is a function $\widehat{\exp}:\widehat{\mathcal{E}}\to\widehat{\mathcal{E}}$ such that $\widehat{\exp}(\widehat{x})=\widehat{\exp}(x)$ for all $x\in\mathcal{E}$. By the discussion above, applied to $\mathbf{R}=(\overline{\mathbb{R}},\exp)$, we now see that this function $\widehat{\exp}$ agrees with the interpretation of the function symbol exp of the language \mathcal{L}_{\exp} of $(\overline{\mathbb{R}},\exp)$ in the \mathcal{L}_{\exp} -structure $\widehat{\mathcal{E}}$ (which is a model of T_{\exp}). For $T=T_{\operatorname{an}}$, the interpretations of the function symbols for restricted analytic functions in the $\mathcal{L}_{\operatorname{an}}$ -structure $\widehat{\mathcal{O}}$ are induced, in a similar way, by the interpretations of those symbols in ${}^*\mathbf{R}$. This is an immediate consequence of a variant of Lemma 3.3, which we formulate and prove below.

For this, suppose that $f: X \to \mathbb{R}$ is a 0-definable continuous function, where $X \subseteq \mathbb{R}^n$ is open. Let $\widehat{f}: X(\widehat{O}) \to \widehat{O}$ be the function defined by the same formula that defines f in \mathbf{R} . Then for $a \in X(\mathbf{R}')$ we have $\widehat{f}(\widehat{a}) = \widehat{f(a)}$. Recall that by Lemma 3.1, O is closed under f.

Lemma 3.4. Let $a \in X(\mathbf{R}')$ and $b \in \mathcal{O}^n$ such that $a - b \in \mathcal{O}^n$. Then $b \in X(^*\mathbf{R})$ and $f(a) - f(b) \in \mathcal{O}$.

Proof. Take $\delta \in (R')^{>0}$ such that $\mathbf{R}' \models \forall x (|x-a| < \delta \to x \in X)$. Since \mathbf{R}' is an elementary substructure of ${}^*\mathbf{R}$ and $|a-b| < \delta$, we have $b \in X({}^*\mathbf{R})$. Now fix $\varepsilon \in (R')^{>0}$. We need to prove that $|f(a)-f(b)| < \varepsilon$. Take $\delta \in (R')^{>0}$ such that $\mathbf{R}' \models \forall x (|x-a| < \delta \to |f(x)-f(a)| < \varepsilon)$. Since $\mathbf{R}' \preccurlyeq {}^*\mathbf{R}$ and $|a-b| < \delta$, we obtain $|f(a)-f(b)| < \varepsilon$.

Now suppose that $b \in \mathcal{O}^n$ is such that $\widehat{b} \in X(\widehat{\mathcal{O}})$. Take $a \in (R')^n$ such that $\widehat{a} = \widehat{b}$. Then $a \in X(\mathbf{R'})$, and $b \in X(^*\mathbf{R})$ and $f(a) - f(b) \in o$ by the previous lemma. Consequently, we see that $\widehat{f}(\widehat{b}) = \widehat{f}(\widehat{a}) = \widehat{f(b)}$.

In Section 4 below we show that not only each restricted analytic function on \mathbb{R}^n , but *each* restricted C^{∞} -function on \mathbb{R}^n induces a function on \widehat{O}^n . However, in general, the connection between this induced function and its original seems less tight. (See the question at the end of Section 4.)

3.2. **Proof of the main theorem.** We are now ready to give a proof of the main theorem from the introduction. We first observe:

Lemma 3.5. Let S be an elementary extension of R, and let S' be an elementary extension of S such that S is cofinal in S'. Let $h: S \to {}^*R$ be an elementary embedding such that $h(S) \subseteq O$. Suppose the ordered set ${}^*\mathbb{R}$ is κ^+ -saturated where $\kappa = \operatorname{ch}(S')$. Then h extends to an elementary embedding $S' \to {}^*R$ whose image is contained in O.

Proof. By Lemma 2.4, h extends to an elementary embedding $S' \to {}^*R$, also denoted by h. Since S is cofinal in S' and $h(S) \subseteq O$, we have $h(S') \subseteq O$.

In connection with the following proposition we note that [11, (2.13)] shows that if $\xi \in O$ with $\xi > \mathbb{R}$, then $\mathbf{R}\langle \xi \rangle \subseteq O$.

Proposition 3.6. Let S be an elementary extension of R and $x \in S$, $x > \mathbb{R}$, such that $R\langle x \rangle$ is cofinal in S. Suppose the ordered set ${}^*\mathbb{R}$ is κ^+ -saturated, where $\kappa = \operatorname{ch}(S)$. Then for each $\xi \in O$, $\xi > \mathbb{R}$, there is an elementary embedding $S \to {}^*R$ which is the identity on \mathbb{R} and sends x to ξ , and whose image is contained in O (and hence there is an elementary embedding $S \to \widehat{O}$ which is the identity on \mathbb{R} and sends x to $\widehat{\xi}$).

Proof. Let $\xi \in O$ with $\xi > \mathbb{R}$. By o-minimality, take an isomorphism $h \colon \mathbf{R}\langle x \rangle \to \mathbf{R}\langle \xi \rangle$ with h(r) = r for all $r \in \mathbb{R}$ and $h(x) = \xi$. By the remark preceding the proposition and the previous lemma, h extends to an elementary embedding $\mathbf{S} \to {}^*\mathbf{R}$, also denoted by h, with $h(S) \subseteq O$. By Zorn, take an elementary substructure \mathbf{R}' of ${}^*\mathbf{R}$, maximal subject to the conditions $h(S) \subseteq R' \subseteq O$. The residue map $O \to \widehat{O}$ now restricts to an isomorphism $\mathbf{R}' \to \widehat{O}$ of \mathcal{L} -structures, and pre-composition with h yields the desired elementary embedding $\mathbf{S} \to \widehat{O}$.

The previous proposition immediately yields a more precise form of Theorem 0.1 from the introduction:

Theorem 3.7. Suppose the ordered set ${}^*\mathbb{R}$ is \aleph_1 -saturated and let O be a convex subring of ${}^*\mathbb{R}$ which is closed under exp. Then for each $\xi > \mathbb{R}$ in O there exists an embedding $\mathbb{T} \to \widehat{O}$ of $\mathcal{L}_{\mathrm{an,exp}}$ -structures over \mathbb{R} with $x \mapsto \xi$.

Proof. We apply the above material to $\mathbf{R} = \mathbb{R}_{\mathrm{an,exp}}$ (so $T = T_{\mathrm{an,exp}}$). By the remark following Lemma 3.1, each convex subring of ${}^*\mathbb{R}$ which is closed under exp is a T-convex subring of ${}^*\mathbf{R}$. By Corollary 2.2, the ordered set \mathbb{T} has countable character. Moreover, by construction of \mathbb{T} , the sequence $(\exp_n(x))$ of iterated exponentials of x is cofinal in \mathbb{T} . Hence the theorem follows from the previous proposition. \square

By Lemmas 2.1 and 3.5 and (the proof of) the previous theorem, we obtain:

Corollary 3.8. Suppose the ordered set \mathbb{R} is \aleph_1 -saturated and let O be a convex subring of \mathbb{R} which is closed under exp. Then there is an embedding of \mathcal{L}_{an} -structures $\mathbb{R}[[\mathfrak{T}]] \to \widehat{O}$ over \mathbb{R} which restricts to an embedding $\mathbb{T} \to \widehat{O}$ of $\mathcal{L}_{an,exp}$ -structures with $x \mapsto \xi$.

We finish this section with another application of the embedding lemma from Section 2, to Conway's surreal numbers.

3.3. Embedding \mathbb{T} into the surreals. The surreal numbers form a (proper) class \mathbf{No} equipped with a linear ordering, extending the ordered class of all ordinal numbers, and also containing \mathbb{R} as an ordered subset in a natural way. This ordered class comes with natural algebraic operations making it a real closed ordered field extension of $\overline{\mathbb{R}}$. The remarkable characteristic property of the ordered field \mathbf{No} is that it is the homogeneous universal ordered field: every ordered field whose universe is a set embeds into \mathbf{No} , and any isomorphism between subfields of \mathbf{No} whose universes are sets extends to an automorphism of \mathbf{No} . We refer to [21] for the construction and basic properties of the class \mathbf{No} . We recall in particular that each surreal number has a length, which is an ordinal, and that the collection $\mathbf{No}(\lambda)$ of surreal numbers of length less than a given ordinal λ forms a set, with $\mathbb{R} \subseteq \mathbf{No}(\omega+1)$. Also, the ordinal ω , viewed as a surreal, is larger than every real number, viewed as a surreal: $\omega > \mathbb{R}$.

Recall that an ε -number is an ordinal λ with the property that $\omega^{\lambda} = \lambda$. For example, every uncountable cardinal is an ε -number. The smallest ε -number is $\varepsilon_0 = \sup\{\omega, \omega^{\omega}, \omega^{\omega^{\omega}}, \ldots\}$. If λ is an ε -number, then $\mathbf{No}(\lambda)$ is a subfield of \mathbf{No} . It was already noted by Kruskal (see [21, Chapter 10]) that the exponential function on \mathbb{R} extends to an exponential function on the ordered field \mathbf{No} . In [10], van den Dries and Ehrlich show that if λ is an ε -number then $\mathbf{No}(\lambda)$ is closed under exponentiation and under taking logarithms of positive elements, and the exponential field $(\mathbf{No}(\lambda), \exp)$ is an elementary extension of $(\overline{\mathbb{R}}, \exp)$. In fact, [10] also shows that in this case, the exponential field $(\mathbf{No}(\lambda), \exp)$ can be expanded to an $\mathcal{L}_{\mathrm{an,exp}}$ -structure making $\mathbf{No}(\lambda)$ an elementary extension of $\mathbb{R}_{\mathrm{an,exp}}$. We now obtain a complement to [18, Theorem 19]:

Corollary 3.9. There is an embedding of \mathcal{L}_{an} -structures $\mathbb{R}[[\mathfrak{T}]] \to \mathbf{No}(\omega_1)$ over \mathbb{R} which restricts to an embedding $\mathbb{T} \to \mathbf{No}(\omega_1)$ of $\mathcal{L}_{an,exp}$ -structures with $x \mapsto \omega$.

Proof. Let $\mathbb{R}\langle x \rangle$ denote the elementary $\mathcal{L}_{\mathrm{an,exp}}$ -substructure of \mathbb{T} generated by x over \mathbb{R} , and similarly let $\mathbb{R}\langle \omega \rangle$ denote the elementary $\mathcal{L}_{\mathrm{an,exp}}$ -substructure of $\mathbf{No}(\omega_1)$ generated by ω over \mathbb{R} . (Note that $\mathbb{R}\langle x \rangle \subseteq \mathbb{R}[[x^{\mathbb{R}}]]^{\mathrm{LE}}$ and $\mathbb{R}\langle \omega \rangle \subseteq \mathbf{No}(\varepsilon_0)$.) By ominimality, we have an isomorphism $h \colon \mathbb{R}\langle x \rangle \to \mathbb{R}\langle \omega \rangle$ of $\mathcal{L}_{\mathrm{an,exp}}$ -structures with h(r) = r for each $r \in \mathbb{R}$ and $h(x) = \omega$. The underlying ordered set of $\mathbf{No}(\omega_1)$ is \aleph_1 -saturated (see [17, Lemma 1] and proof of [18, Theorem 17]). The claim now follows from Lemmas 2.1 and 2.4.

This corollary leads to a number of natural questions which we do not pursue here, e.g.: what is the smallest ε -number λ such that there is an embedding $\mathbb{T} \to \mathbf{No}(\lambda)$ of $\mathcal{L}_{\mathrm{an,exp}}$ -structures with $x \mapsto \omega$?

4. Differentiability in Residue Fields, and a Question

As in the introduction we let O be a convex subring of ${}^*\mathbb{R}$, with maximal ideal o, residue field $\widehat{O} = O/o$, and residue morphism $x \mapsto \widehat{x} \colon O \to \widehat{O}$. In this last section of the paper we show that each restricted C^{∞} -function $\mathbb{R}^n \to \mathbb{R}$ extends to a restricted C^{∞} -function $\widehat{O}^n \to \widehat{O}$ in a natural way.

Notation. For $X \subseteq \mathbb{R}^n$ we let

$$\mu(X) := \left\{ x \in ({}^*\mathbb{R}_{\operatorname{fin}})^n : \operatorname{st}(x) \in X \right\}$$

be the monad of X, and we let

$$\widehat{\mu}(X) := \{\widehat{x} : x \in \mu(X)\} \subseteq \widehat{O}^n$$

be the image of $\mu(X)$ under $x \mapsto \hat{x}$. So for example, let

$$B = (a_1, b_1) \times \cdots \times (a_n, b_n)$$
 where $a_i, b_i \in \mathbb{R}$ with $a_i < b_i$ (3)

be an open box in \mathbb{R} ; then

$$\mu(B) = (a_1, b_1)_{*\mathbb{R}_{\text{fin}}} \times \cdots \times (a_n, b_n)_{*\mathbb{R}_{\text{fin}}}, \quad \widehat{\mu}(B) = [a_1, b_1]_{\widehat{O}} \times \cdots \times [a_n, b_n]_{\widehat{O}}.$$

We first show that each infinitely differentiable real-valued function defined on an open set $X \subseteq \mathbb{R}^n$ extends in a natural way to an infinitely differentiable function on the interior int $\widehat{\mu}(X)$ of $\widehat{\mu}(X)$ (taking values in \widehat{O}). Given $x, y \in {}^*\mathbb{R}^n$ we denote by

$$[x,y] := \{tx + (1-t)y : t \in [0,1]_{\mathbb{R}}\}\$$

the line segment in \mathbb{R} between x and y.

Let first $f: B \to \mathbb{R}$ be C^1 , where B is as in (3).

Lemma 4.1. Let $x \in \mu(B)$. Then $f(x) \in {}^*\mathbb{R}_{fin}$, and if in addition $y \in \mu(B)$ and $x - y \in \sigma^n$, then $f(x) - f(y) \in \sigma$.

Proof. We have $f(x) \approx f(\operatorname{st}(x))$, hence $f(x) \in {}^*\mathbb{R}_{\operatorname{fin}}$. Now let $y \in \mu(B)$ and $x - y \in \sigma^n$. By the Mean Value Theorem, there is $z \in [x,y]$ such that $f(y) - f(x) = f'(z) \cdot (y-x)$. However, $\operatorname{st}(z) = \operatorname{st}(x) \in B$, and f' is continuous, so $f'(z) \approx f'(\operatorname{st}(z))$ and hence $f(y) - f(x) = f'(z) \cdot (y-x)$ where $f'(z) \in ({}^*\mathbb{R}_{\operatorname{fin}})^n$ and $y - x \in \sigma^n$. Thus $f(y) - f(x) \in \sigma$ as required.

The previous lemma allows us to define a function

$$\widehat{f}\colon \widehat{\mu}(B)\to \widehat{\mathcal{O}}, \qquad \widehat{f}(\widehat{x}):=\widehat{f(x)} \quad \text{ for each } x\in \mu(B).$$

Next we show:

Lemma 4.2. The function \hat{f} is continuous.

Proof. Fix $x \in \mu(B)$. Let $\varepsilon \in O$ with $\varepsilon > o$. We need to find $\delta \in O$ with $\delta > o$ such that for all $y \in \mu(B)$ such that $|x - y| < \delta$ we have $|f(x) - f(y)| < \varepsilon$. For this, we may assume that $\varepsilon \in {}^*\mathbb{R}_{\inf}$. Take a closed box $C \subseteq B$ whose interior contains $\operatorname{st}(x)$, and $M \in \mathbb{N}^{>0}$ such that $|f'(y)| \leq M$ for all $y \in B$. We claim that $\delta = \frac{\varepsilon}{M}$ works. Indeed, let $y \in \mu(B)$ satisfy $|x - y| < \delta$. Take $z \in [x, y]$ such that $f(y) - f(x) = f'(z) \cdot (y - x)$. Then $y - x \in ({}^*\mathbb{R}_{\inf})^n$ and thus $\operatorname{st}(z) = \operatorname{st}(x)$, so $z \in {}^*C$ and $|f'(z)| \leq M$. Hence $|f(y) - f(x)| \leq |f'(z)| \cdot |y - x| < M \cdot \delta = \varepsilon$.

Lemma 4.3. Suppose f is C^2 . Then the restriction of \widehat{f} to int $\widehat{\mu}(B)$ is differentiable with derivative \widehat{f}' .

Proof. Fix $x \in \mu(B)$. Let $e_1, \ldots, e_n \in \mathbb{R}^n$ be the standard basis vectors of \mathbb{R} , and fix $i \in \{1, \ldots, n\}$. Let $\varepsilon \in {}^*\mathbb{R}_{inf}$ with $\varepsilon > o$; we need to find $\delta \in \mathcal{O}$ with $\delta > o$ such that for each $h \in {}^*\mathbb{R}$ with $0 < |h| < \delta$ we have

$$\left|\frac{f(x+he_i)-f(x)}{h}-\frac{\partial f}{\partial x_i}(x)\right|<\varepsilon.$$

Again, let C be a closed box contained in B such that $\operatorname{st}(x)$ is in the interior of C. Let $M \in \mathbb{N}^{>0}$ be such that $|(\frac{\partial f}{\partial x_i})'(y)| \leq M$ for all $y \in C$. We claim that $\delta = \frac{\varepsilon}{M}$ works. Suppose $h \in {}^*\mathbb{R}$, $0 < |h| < \delta$. Take $y \in [x, x + h]$ such that $\frac{f(x+he_i)-f(x)}{h} = \frac{\partial f}{\partial x_i}(y)$. We need $|\frac{\partial f}{\partial x_i}(y) - \frac{\partial f}{\partial x_i}(x)| < \varepsilon$. Take $z \in [x,y]$ such that $\frac{\partial f}{\partial x_i}(y) - \frac{\partial f}{\partial x_i}(x) = (\frac{\partial f}{\partial x_i})'(z) \cdot (y - x)$. Since $\operatorname{st}(z) = \operatorname{st}(x)$ we have $z \in {}^*C$ and thus $|(\frac{\partial f}{\partial x_i})'(z)| \cdot |y - x| < M \cdot \delta = \varepsilon$, as required.

Corollary 4.4. Let $N \in \mathbb{N}$. If f is C^{N+1} , then $\widehat{f} \upharpoonright \operatorname{int} \widehat{\mu}(B)$ is C^N . In particular, if f is C^{∞} , then $\widehat{f} \upharpoonright \operatorname{int} \widehat{\mu}(B)$ is C^{∞} .

Proof. By induction on N, where the case N=0 holds by Lemma 4.2. Now suppose that f is C^{N+2} . Then $f' \colon B \to \mathbb{R}^n$ is C^{N+1} , so $\widehat{f'} \upharpoonright \inf \widehat{\mu}(B)$ is C^N by induction. But $\widehat{f'} = (\widehat{f})'$ on int $\widehat{\mu}(B)$, by Lemma 4.3, so $\widehat{f} \upharpoonright \inf \widehat{\mu}(B)$ is C^{N+1} .

Let X be an arbitrary open subset of \mathbb{R}^n and let $f: X \to \mathbb{R}$ be C^1 . Then X is a union of open boxes B of the form (3), and $\widehat{\mu}(X)$ is then the union of the corresponding sets $\widehat{\mu}(B)$. Hence by the above, we can define a function

$$\widehat{f} \colon \widehat{\mu}(X) \to \widehat{\mathcal{O}}, \qquad \widehat{f}(\widehat{x}) := \widehat{f(x)} \quad \text{ for each } x \in \mu(X),$$

and then the analogues of Lemmas 4.2 and 4.3, and Corollary 4.4 hold (with X in place of B).

Let now I := [-1, 1], and for an ordered field extension K of \mathbb{R} write

$$I(K) := \{x \in K : -1 \le x \le 1\}.$$

Let C^{∞} be the collection of all functions $I^n \to \mathbb{R}$, for varying n, which extend to a C^{∞} -function on a neighborhood of I^n , and let $C \subseteq C^{\infty}$. A **restricted** C-function is a function $\mathbb{R}^n \to \mathbb{R}$ which on I^n agrees with a function in C and which is 0 on $\mathbb{R}^n \setminus I^n$. Let L_C be the language of ordered rings augmented by an n-ary function symbol for each restricted C-function $\mathbb{R}^n \to \mathbb{R}$; we use the same letter to denote the function from C and its corresponding function symbol. We may expand the ordered field \widehat{O} to an L_C -structure by interpreting each function symbol f, where $f: \mathbb{R}^n \to \mathbb{R}$ is a restricted C-function, by the function $f^{\widehat{O}}: \widehat{O}^n \to \widehat{O}$ given by

$$f^{\widehat{O}}(\widehat{x}) = \begin{cases} \widehat{f}(\widehat{x}) & \text{if } \widehat{x} \in I(\widehat{O})^n, \\ 0 & \text{otherwise.} \end{cases}$$

Note that this definition makes sense by the discussion above, since f agrees with a C^{∞} -function on an open neighborhood of I^n . Also note that \mathbb{R} is the underlying set of a substructure $\mathbb{R}_{\mathcal{C}}$ of the $\mathcal{L}_{\mathcal{C}}$ -structure $\widehat{\mathcal{O}}$.

Question. Are there natural conditions on \mathcal{C} which ensure that the $\mathcal{L}_{\mathcal{C}}$ -structure $\mathbb{R}_{\mathcal{C}}$ is an elementary substructure of $\widehat{\mathcal{O}}$?

One such natural condition is that $\mathbb{R}_{\mathcal{C}}$ be o-minimal: in this case, \widehat{O} can be made into an elementary extension of $\mathbb{R}_{\mathcal{C}}$ as explained in the previous section, and $f^{\widehat{O}}$ then agrees with the interpretation of the function symbol f in this $\mathcal{L}_{\mathcal{C}}$ -structure, by Lemma 3.4 and the discussion surrounding it. Example of families \mathcal{C} which make $\mathbb{R}_{\mathcal{C}}$ o-minimal are, of course, the family consisting of all restrictions to the unit cubes I^n of analytic functions on neighborhorhoods of I^n (so $\mathbb{R}_{\mathcal{C}} = \mathbb{R}_{\mathrm{an}}$), or the family of all restrictions to the unit cubes of C^{∞} -functions associated to a given Denjoy-Carleman class [32]. Also, Le Gal [20] and Grigoriev [22] have shown that the expansion of the real field by a generic restricted C^{∞} -function is o-minimal.

In general, however, $\mathbb{R}_{\mathcal{C}}$ is not an elementary substructure of $\widehat{\mathcal{O}}$, as was pointed out to us by Dave Marker:

Example. Suppose that \mathbb{N} is definable in $\mathbb{R}_{\mathcal{C}}$, say by the $\mathcal{L}_{\mathcal{C}}$ -formula $\varphi(x)$, possibly involving parameters. (This hypothesis is satisfied if \mathcal{C} contains a restricted C^{∞} -function with an infinite discrete zero set which is in semialgebraic bijection with \mathbb{N} , such as the restricted C^{∞} -function s with s(0) = 0 and $s(x) = e^{-1/x^2} \sin(1/x)$ for $x \in [-1,1] \setminus \{0\}$.) Suppose further that \mathcal{O} is the convex hull of $\mathbb{R}[\xi]$ in ${}^*\mathbb{R}$, where $\xi \in {}^*\mathbb{R}$ is positive infinite. We then claim that $\mathbb{R}_{\mathcal{C}}$ is not an elementary substructure of $\widehat{\mathcal{O}}$. Indeed, suppose that $\widehat{\mathcal{O}} \models \varphi(\alpha) \land \alpha > \xi$. Let $\psi(x,y)$ define the graph of the function $x \mapsto 2^x \colon \mathbb{N} \to \mathbb{N}$ in $\mathbb{R}_{\mathcal{C}}$. Then if $\widehat{\mathcal{O}} \models \psi(\alpha,\beta)$, then by elementarity, we have $\widehat{\mathcal{O}} \models \beta > \widehat{\xi}^n$, for each n, contradicting the fact that $\widehat{\xi}, \widehat{\xi}^2, \ldots$ is cofinal in $\widehat{\mathcal{O}}$.

Of course, once we can define the set \mathbb{N} in $\mathbb{R}_{\mathcal{C}}$, we immediately define all projective sets; in particular, we define all C^{∞} -functions, making the preceding example a very wild one. On the other hand, even in this situation, we sometimes obtain an existentially closed substructure:

Lemma 4.5. Suppose ${}^*\mathbb{R}$ is \mathfrak{c}^+ -saturated. Then there is a convex subring \mathcal{O} of ${}^*\mathbb{R}$ with $\mathbb{R}_{\text{fin}} \subseteq \mathcal{O} \subseteq {}^*\mathbb{R}$, such that $\mathbb{R}_{\mathcal{C}^{\infty}} \preceq_1 \widehat{\mathcal{O}}$.

This follows from results in [36]; we recall some definitions and basic results from that paper.

Let $\mathcal{L}_{\operatorname{src}}$ be the expansion of the language of rings by an n-ary function symbol f for every continuous function $f \colon \mathbb{R}^n \to \mathbb{R}$ (for varying n). A **super real closed ring** is an $\mathcal{L}_{\operatorname{src}}$ -structure which expands a commutative ring with 1, with the interpretations of +, \cdot and 0, 1 compatible with the interpretations of the functions symbols associated to the corresponding continuous functions $\mathbb{R}^2 \to \mathbb{R}$ and $\mathbb{R}^0 \to \mathbb{R}$, respectively, and satisfying the $\mathcal{L}_{\operatorname{src}}$ -sentences

$$\forall x (\mathrm{id}(x) = x)$$
 (where id: $\mathbb{R} \to \mathbb{R}$ is the identity function)

and

$$\forall x_1 \cdots \forall x_n \bigg(f \big(g_1(x_1), \dots, g_n(x_n) \big) = \big(f \circ (g_1, \dots, g_n) \big) (x_1, \dots, x_n) \bigg),$$

for all continuous functions $f: \mathbb{R}^n \to \mathbb{R}$ and $g_i: \mathbb{R}^{m_i} \to \mathbb{R}$ (i = 1, ..., n). The class of super real closed rings is a variety (in the sense of universal algebra).

Clearly the field \mathbb{R} can be expanded to an \mathcal{L}_{src} -structure in a natural way. Let A be a super real closed ring. Then A can be viewed as an extension of \mathbb{R} in a unique way (identifying each $r \in \mathbb{R}$ with the interpretation of the constant function r in A), and if A is an integral domain, then \mathbb{R} is existentially closed in A [36, Corollary 5.6].

A prime ideal P of A is said to be Υ -radical if it is the kernel of a morphism $A \to B$ of $\mathcal{L}_{\mathrm{src}}$ -structures, for some super real closed ring B. (See [36, Sections 3 and 6] for an explanation of this terminology.) In this case, there is a unique expansion of the integral domain A/P to a super real closed ring such that the residue map $A \to A/P$ is a morphism of $\mathcal{L}_{\mathrm{src}}$ -structures. Every maximal ideal of A is Υ -radical [36, Theorem 6.14]. Given a subset C of A, there exists a smallest super real closed subring of A containing C; the cardinality of this super real closed subring of A equals $|C| + \mathfrak{c}$. The convex hull of a super real closed subring of A is super real closed [36, Corollary 9.2, (i)].

We can now give:

Proof of Lemma 4.5. We view ${}^*\mathbb{R}$ as a super real closed ring in the natural way. Let ξ be a positive infinite element of ${}^*\mathbb{R}$ and let O be the convex hull of the smallest super real closed subring of ${}^*\mathbb{R}$ containing ξ . Then $\mathbb{R}_{\text{fin}} \subsetneq O$ (since $\xi \in O$) and $O \subsetneq {}^*\mathbb{R}$ (by \mathfrak{c}^+ -saturation of ${}^*\mathbb{R}$). The maximal ideal o of O is Υ -radical; extend the residue field $\widehat{O} = O/o$ to a super real closed ring such that the residue map $O \to \widehat{O}$ is a morphism of \mathcal{L}_{src} -structures. Then \mathbb{R} , viewed as super real closed ring, is existentially closed in \widehat{O} . Note that the ordering of \mathbb{R} is quantifier-free definable in the super real closed ring \mathbb{R} , as $x \geqslant 0 \Longleftrightarrow |x| = x$ for each $x \in \mathbb{R}$. Every restricted C^{∞} -function $f \colon \mathbb{R}^n \to \mathbb{R}$ is quantifier-free definable in the \mathcal{L}_{src} -structure \mathbb{R} (since $f \upharpoonright I^n$ has an extension to a continuous function $\mathbb{R}^n \to \mathbb{R}$, by Tietze Extension); moreover, in the super real closed ring \widehat{O} , the same quantifier-free formula defines the function $f^{\widehat{O}}$ introduced above. All of this now easily implies that $\mathbb{R}_{C^{\infty}} \preccurlyeq_1 \widehat{O}$. \square

It might be interesting to isolate a tameness property of $\mathbb{R}_{\mathcal{C}}$, weaker than ominimality, which guarantees $\mathbb{R}_{\mathcal{C}} \preccurlyeq \widehat{O}$ (in the language $\mathcal{L}_{\mathcal{C}}$) for *every* convex subring O of ${}^*\mathbb{R}$.

References

- M. Aschenbrenner, L. van den Dries, H-fields and their Liouville extensions, Math. Z. 242 (2002), 543–588.
- M. Aschenbrenner, L. van den Dries, J. van der Hoeven, Towards a model theory of transseries, preprint (2011).
- 3. _____, Differentially algebraic gaps, Selecta Math. 11 (2005), 247–280.
- M. Aschenbrenner, A. Fischer, Definable versions of theorems by Kirszbraun and Helly, Proc. London Math. Soc. 102 (2011), 468–502.
- 5. B. Dahn, P. Göring, Notes on exponential-logarithmic terms, Fund. Math. 127 (1986), 45-50.
- J. Denef, L. van den Dries, p-adic and real subanalytic sets, Ann. of Math. (2) 128 (1988), no. 1, 79–138.
- L. van den Dries, Tame Topology and O-Minimal Structures, London Math. Soc. Lecture Note Series, vol. 248, Cambridge University Press, Cambridge (1998).
- 8. ______, o-minimal structures, in: W. Hodges et al. (eds.), Logic: From Foundations to Applications (Staffordshire, 1993), pp. 137–185, Oxford Science Publications, The Clarendon Press, Oxford University Press, New York, 1996.
- 9. _____, A generalization of the Tarski-Seidenberg theorem, and some nondefinability results, Bull. Amer. Math. Soc. 15 (1986), 189–193.
- L. van den Dries, P. Ehrlich, Fields of surreal numbers and exponentiation, Fund. Math. 167 (2001), no. 2, 173–188.
- 11. L. van den Dries, A. Lewenberg, T-convexity and tame extensions, J. Symbolic Logic 60 (1995), no. 1, 74–102.
- L. van den Dries, A. Macintyre, D. Marker, The elementary theory of restricted analytic fields with exponentiation, Ann. Math. 140 (1994), 183–205.

- Logarithmic-exponential power series, J. London Math. Soc. 56 (1997), 417–434.
 Logarithmic-exponential series, Ann. Pure Appl. Logic 111 (2001), 61–113.
- 15. L. van den Dries, C. Miller, On the real exponential field with restricted analytic functions, Israel J. Math. 85 (1994), 19-56.
- 16. J. Écalle, Introduction aux Fonctions Analysables et Preuve Constructive de la Conjecture de Dulac, Actualités Mathématiques, Hermann, Paris (1992).
- 17. P. Ehrlich, An alternative construction of Conway's ordered field No, Algebra Universalis 25 (1988), no. 1, 7–16.
- _____, The absolute arithmetic continuum and the unification of all numbers great and small, Bull. Symbolic Logic 18 (2012), no. 1, 1-45.
- 19. J. Esterle, Solution d'un problème d'Erdős, Gillman et Henriksen et application à l'étude des homomorphismes de C(K), Acta Math. Acad. Sci. Hungar. 30 (1977), no. 1-2, 113–127.
- O. Le Gal, A generic condition implying o-minimality for restricted C^{∞} -functions, Ann. Fac. Sci. Toulouse Math. (6) 19 (2010), no. 3-4, 479-492.
- 21. H. Gonshor, An Introduction to the Theory of Surreal Numbers, London Mathematical Society Lecture Note Series, vol. 110, Cambridge University Press, Cambridge, 1986.
- 22. A. Grigoriev, On o-minimality of extensions of the real field by restricted generic smooth functions, preprint, available online at arXiv.org:math/0506109 (2005).
- 23. W. Hodges, Model Theory, Encyclopedia of Mathematics and its Applications, vol. 42, Cambridge University Press, Cambridge, 1993.
- 24. J. van der Hoeven, Asymptotique Automatique, Thèse, École Polytechnique, Paris (1997).
- , Transseries and Real Differential Algebra, Lecture Notes in Math., vol. 1888, Springer-Verlag, Berlin, 2002.
- 26. M. Knebusch, M. Wright, Bewertungen mit reeller Henselisierung, J. Reine Angew. Math. **286/287** (1976), 314–321.
- 27. J. Knight, A. Pillay, C. Steinhorn, Definable sets in ordered structures, II, Trans. Amer. Math. Soc. **295** (1986), no. 2, 593–605.
- 28. F.-V. Kuhlmann, S. Kuhlmann, S. Shelah, Exponentiation in power series fields, Proc. Amer. Math. Soc. 125 (1997), 3177–3183.
- 29. S. Kuhlmann, M. Tressl, Comparison of exponential-logarithmic and logarithmic-exponential series, preprint (2011).
- 30. C. Miller, Exponentiation is hard to avoid, Proc. Amer. Math. Soc. 122 (1994), 257-259.
- 31. A. Pillay, C. Steinhorn, Definable sets in ordered structures, I, Trans. Amer. Math. Soc. 295 (1986), 565-592.
- 32. J.-P. Rolin, P. Speissegger, A. J. Wilkie, Quasianalytic Denjoy-Carleman classes and ominimality, J. Amer. Math. Soc. 16 (2003), no. 4, 751-777.
- 33. M. Schmeling, Corps de Transséries, Ph.D. thesis, Université Paris VII (2001).
- 34. T. Todorov, H. Vernaeve, Asymptotic fields, generalized power series and special Colombeau algebras of generalized functions, manuscript (2012).
- _, Generalized power series and non-standard analysis: between model theory and NSA, slides of a talk given at York University (August 2009), available online at http://maths.york.ac.uk/www/sites/default/files/Todorov-slides.pdf
- 36. M. Tressl, Super real closed rings, Fund. Math. 194 (2007), no. 2, 121–177.
- 37. A. Wilkie, Model completeness results for expansions of the ordered field of real numbers by restricted Pfaffian functions and the exponential function, J. Amer. Math. Soc. 9 (1996), 1051 - 1094.

E-mail address: matthias@math.ucla.edu

Department of Mathematics, University of California, Los Angeles, Box 951555, Los Angeles, CA 90095-1555, U.S.A.

E-mail address: isaac@math.ucla.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, LOS ANGELES, BOX 951555, LOS Angeles, CA 90095-1555, U.S.A.