CHAINS OF SATURATED MODELS IN AECS

WILL BONEY AND SEBASTIEN VASEY

ABSTRACT. We study when a union of saturated models is saturated in the framework of tame abstract elementary classes (AECs) with amalgamation. We prove:

Theorem 0.1. If K is a tame AEC with amalgamation satisfying a natural definition of superstability (which follows from categoricity in a high-enough cardinal), then for all high-enough λ :

- (1) The union of an increasing chain of λ -saturated models is λ -saturated.
- (2) There exists a type-full good λ-frame with underlying class the saturated models of size λ.
- (3) There exists a unique limit model of size λ .

Our proofs use independence calculus and a generalization of averages to this non first-order context.

Contents

1.	Introduction	1
2.	Using independence calculus: the stable case	4
3.	Using independence calculus: the superstable case	10
4.	Averages	11
5.	Translating to AECs	20
6.	On superstability in AECs	25
Ref	References	

1. INTRODUCTION

Determining when a union of λ -saturated models is λ -saturated is an important dividing line in first-order model theory. Recall that Harnik and Shelah have shown:

Date: March 21, 2022

AMS 2010 Subject Classification: Primary 03C48. Secondary: 03C47, 03C52, 03C55, 03E55.

Key words and phrases. Abstract elementary classes; Forking; Independence calculus; Classification theory; Stability; Superstability; Tameness; Saturated models; Limit models; Averages; Stability theory inside a model.

This material is based upon work done while the first author was supported by the National Science Foundation under Grant No. DMS-1402191 and the second author was supported by the Swiss National Science Foundation under Grant No. 155136.

Fact 1.1 ([Har75], III.3.11 in [She90] for the case $\lambda \leq |T|$). Let T be a first-order theory.

- If T is superstable, then any increasing union of λ -saturated models is λ -saturated.
- If T is stable, then any increasing union of λ -saturated models of cofinality at least $|T|^+$ is λ -saturated.

A converse was later proven by Albert and Grossberg [AG90, Theorem 13]. Fact 1.1 can be used to prove:

Fact 1.2 (The saturation spectrum theorem, VIII.4.7 in [She90]). Let T be a stable first-order theory. Then T has a saturated model of size λ if and only if [T is stable in λ or $\lambda = \lambda^{<\lambda} + |D(T)|$].

Although not immediately evident from the statement, the proof of Fact 1.1 relies on the heavy machinery of forking and averages.

While the saturation spectrum theorem has been generalized to homogeneous model theory (see [She75, 1.13] or [GL02, 5.9]), to the best of our knowledge no explicit generalization of Fact 1.1 has been published in this context (Shelah asserts it without proof in [She75, 1.15]). Grossberg [Gro91] has proven a version of Fact 1.1 in the framework of stability theory inside a model. The proof uses averages but relies on a strong negation of the order property. Makkai and Shelah [MS90, 4.18] have given a generalization in the class of models of an $L_{\kappa,\omega}$ sentence where κ is a strongly compact cardinal. The proof uses independence calculus.

One can ask whether Fact 1.1 can also be generalized to abstract elementary classes (AECs), a general framework for classification theory introduced in [She87] (see [Gro02] for an introduction to AECs). In [She09a, I.5.39], Shelah proves a generalization of the superstable case of Fact 1.1 to "definable-enough" AECs with countable Löwenheim-Skolem number, using the weak continuum hypothesis.

In chapter II of [She09a], Shelah starts with a (weakly successful) good λ -frame (a local notion of superstability) on an abstract elementary class (AEC) K and wants to show that a union of saturated models is saturated in K_{λ^+} . For this purpose, he introduces a restriction \leq^* of the ordering that allows him to prove the result for \leq^* -increasing chains (II.7.7 there). Restricting the ordering of the AEC is somewhat artificial and one can ask what happens in the general case, and also if λ^+ is replaced by an arbitrary cardinal. Moreover, Shelah's methods to obtain a weakly successful good λ -frame typically use categoricity in two successive cardinals and the weak continuum hypothesis¹.

In [She99], Shelah had previously proven that a union of λ -saturated models is λ -saturated, for K an AEC with amalgamation, joint embedding, and no maximal models categorical in a successor $\lambda' > \lambda$ (see [Bal09, Chapter 15] for a writeup), but left the case $\lambda \geq \lambda'$ (or λ' not a successor) unexamined.

¹See for example [She09a, II.3.7]. Shelah also shows how to build a good frame in ZFC from more model-theoretic hypotheses in [She09a, IV.4.10], but he has to change the class and it is not clear his frame is weakly successful.

In this paper, we replace the local model-theoretic assumptions of Shelah with *global* ones, including *tameness*, a locality notion for types introduced by Grossberg and VanDieren [GV06]. We take advantage of recent developments in the study of forking in tame AECs (especially by Boney and Grossberg [BG] and Vasey [Vas16c, Vas16a]) to generalize Fact 1.1 to tame abstract elementary classes with amalgamation. Our main result is:

Corollary 3.5. Assume K is a $(< \kappa)$ -tame AEC with amalgamation. If $\kappa = \beth_{\kappa} > LS(K)$ and K is categorical in some cardinal strictly above κ , then for all $\lambda > 2^{\kappa}$, $K^{\lambda-\text{sat}}$ (the class of λ -saturated models of K) is an AEC with $LS(K^{\lambda-\text{sat}}) = \lambda$.

Notice that if $K^{\lambda\text{-sat}}$ is an AEC, then any increasing union of λ -saturated models is λ -saturated. Thus, in contrast to Shelah's [She99] result, we obtain a *global* theorem that holds for all high-enough λ and not just those under the categoricity cardinal. Furthermore categoricity at a successor is not assumed. We can also replace the categoricity by various notions of superstability defined in terms of the local character for independence notions such as coheir or splitting. In fact, we can combine this result with the construction of a good frame in [Vas16a] to obtain the theorem in the abstract:

Theorem 6.1. If K is a tame AEC with amalgamation satisfying a natural definition of superstability (see Definition 5.12), then for all high-enough λ , there exists a *unique* limit model of size λ .

This proves an eventual version of a statement appearing in early versions of [GVV16] (see the discussion in Section 6).

It is very convenient to have $K^{\lambda\text{-sat}}$ an AEC, as saturated models are typically better behaved than arbitrary ones. This is crucial for example in Shelah's upward transfer of frames in [She09a, Chapter II], and is also used in [Vas16a] to build an ω -successful good frame (and later a global independence notion). We also prove a result for the strictly stable case:

Theorem 5.16. Let K be a κ -tame AEC with amalgamation, $\kappa \geq \mathrm{LS}(K)$, stable in some cardinal above κ . Then there exists $\chi_0 \leq \lambda_0 < \beth_{(2^{\kappa})^+}$ such that whenever $\lambda \geq \lambda_0$ is such that $\mu^{<\chi_0} < \lambda$ for all $\mu < \lambda$, the union of an increasing chain of λ -saturated models of cofinality at least χ_0 is λ -saturated.

One caveat here is the introduction of cardinal arithmetic: in stable, first-order theories, all large enough λ satisfy that the long-enough union of λ -saturated models is λ -saturated (Fact 1.1), while here we have to add the condition that λ is χ_0 -closed. When dealing with compact classes (or even just ($< \omega$)-tame classes), the map $\lambda \mapsto \lambda^{<\omega}$ that takes the size of a set to the number of formulas with parameters in that set can be used freely. Even in the work of Makkai and Shelah [MS90], where κ is strongly compact and the class is ($< \kappa$)-tame, this map is $\lambda \mapsto \lambda^{<\kappa}$, which is constant on most cardinals (those with cofinality at least κ) by a result of Solovay. However, in our context of ($< \kappa$)-tameness for $\kappa > \omega$ but not strongly compact, this function can be much wilder (following the Galois Morleyization, we view Galois types of size less than κ as formulas; see Definition 5.1). Thus, we need to introduce assumptions that this map is well-behaved. Using various tricks, we can bypass these assumptions in the superstable case but are unable to do so in the stable case. For example in Theorem 5.16, the cardinal arithmetic assumption can be replaced by "K is stable in μ for unboundedly many $\mu < \lambda$ ", which is always true in case K is superstable.

We use two main methods: The first method is pure independence calculus, relying on a well-behaved independence relation (coheir), whose existence in our context is proven in [BG, Vas16c]. This works well in the superstable case if we define superstability in terms of coheir (called strong superstability in [Vas16a]) but we do not know how to make it work for weaker definitions of superstability (such as superstability defined in terms of splitting, a more classical definition implicit for example in [GVV16]). The second method is the use of syntactic averages, developed by Shelah in [She09b, Chapter V]. We end up proving a result on chains of saturated models in the framework of stability theory inside a model and then translate to AECs using *Galois Morleyization*, introduced in [Vas16c]. This method allows us to use superstability defined in terms of splitting. The two methods give incomparable results: in case we know that K is $(< \kappa)$ -tame, with $\kappa = \beth_{\kappa} > \mathrm{LS}(K)$, the first gives better Hanf numbers than the second. However if we know that K is $\mathrm{LS}(K)$ -tame, then we get better bounds using the second method, since we do not need to work above a fixed point of the beth function.

The paper is organized as follows. Section 2 gives the argument using independence calculus culminating in Theorems 2.15 and 3.4. Both of these arguments work just using forking relations, drawing inspiration from Makkai and Shelah, rather than the classical first-order argument using averages. Section 4 develops averages in our context based on earlier work of Shelah and culminates in the more local Theorem 4.27. Section 5 translates the local result to AECs and Section 6 proves consequences such as the uniqueness of limit models from superstability.

This paper was written while the second author was working on a Ph.D. thesis under the direction of Rami Grossberg at Carnegie Mellon University. He would like to thank Professor Grossberg for his guidance and assistance in his research in general and in this work specifically.

2. Using independence calculus: the stable case

We assume that the reader is familiar with the basics of AECs, as presented in [Bal09] or [Gro]. We will use the notation from [Vas16c]. In particular, $gtp(\bar{b}/A; N)$ denotes the Galois type of \bar{b} over A as computed in N.

We will use the following set-theoretic notation:

Notation 2.1. For κ an infinite cardinal, write κ_r for the least regular cardinal greater than or equal to κ . That is, κ_r is κ^+ if κ is singular and κ otherwise. Also let κ^- be κ if κ is limit, or the unique κ_0 such that $\kappa = \kappa_0^+$ if κ is a successor. We will often use the "Hanf function" (from [Bal09, Chapter 14]): for λ an infinite cardinal, let $h(\lambda) := \beth_{(2^{\lambda})^+}$. Also let $h^*(\lambda) := h(\lambda^-)$.

All throughout this section, we assume:

Hypothesis 2.2.

(1) K is an AEC with amalgamation, joint embedding, and arbitrarily large models. We work inside a monster model \mathfrak{C} .

- (2) $\operatorname{LS}(K) < \kappa = \beth_{\kappa}$.
- (3) K is $(< \kappa)$ -tame.
- (4) K is stable (in some cardinal above κ).

We will use the independence notion of coheir for AECs, introduced in [BG].

Definition 2.3 (Coheir). Define a tertiary relation \downarrow by $\downarrow(M, A, B)$ if and only if:

- (1) $M \leq \mathfrak{C}$, M is κ -saturated, and $A, B \subseteq |\mathfrak{C}|$.
- (2) For any $\bar{a} \in {}^{<\kappa}A$ and $B_0 \subseteq |M| \cup B$ of size less than κ , there exists $\bar{a}' \in {}^{<\kappa}|M|$ such that $gtp(\bar{a}/B_0) = gtp(\bar{a}'/B_0)$ (here, the Galois types are computed inside \mathfrak{C}).

We write $A \downarrow B$ instead of $\downarrow (M, A, B)$. We will also say that $gtp(\bar{a}/B)$ is a $(< \kappa)$ coheir over \overline{M} when ran $(\overline{a}) \downarrow B$ (it is straightforward to check that this does not Mdepend on the choice of \bar{a}).

The following locality cardinals will play an important role:

Definition 2.4. Let α be a cardinal.

- (1) Let $\bar{\kappa}_{\alpha}(\downarrow)$ be the minimal cardinal $\mu \geq |\alpha|^{+} + \kappa^{+}$ such that for any $M \leq \mathfrak{C}$ that is κ -saturated, any $A \subseteq |\mathfrak{C}|$ with $|A| \leq \alpha$, there exists $M_0 \leq M$ in $K_{\leq \mu}$ with $A \downarrow M$. When μ does not exist, we set $\bar{\kappa}_{\alpha_0}(\mathfrak{i}) = \infty$. M_0
- (2) Let $\kappa_{\alpha}(\downarrow)$ be the minimal cardinal $\mu \geq |\alpha|^{+} + \aleph_{0}$ such that for any regular $\delta \geq \mu$, any increasing chain $\langle M_i : i < \delta \rangle$ in K and any A of size at most α , there exists $i < \delta$ such that $A \downarrow_{M_i} \bigcup_{i < \delta} M_i$. When μ does not exist, we set $\kappa_{\alpha_0}(\mathfrak{i}) = \infty$. For K^* a subclass of K, we similarly define $\kappa_{\alpha}(\downarrow \upharpoonright K^*)$, where in addition we require that $M_i \in K^*$ for all $i < \delta$ (we will use this when K^* is a class of saturated models).

Remark 2.5. For any cardinal α , we always have that $\kappa_{\alpha}(\downarrow) \leq \bar{\kappa}_{\alpha}(\downarrow)$.

Fact 2.6. Under Hypothesis 2.2, \downarrow satisfies the following properties:

- (1) <u>Invariance</u>: If f is an automorphism of \mathfrak{C} and $A \underset{M}{\downarrow} B$, then $f[A] \underset{f[M]}{\downarrow} B$.
- (2) <u>Monotonicity</u>: Assume $A \underset{M}{\downarrow} B$, then:
 - (a) Left and right monotonicity: If $A_0 \subseteq A$, $B_0 \subseteq B$, then $A_0 \underset{M}{\downarrow} B_0$.
 - (b) Base monotonicity: If $M \leq M' \leq \mathfrak{C}$, $|M'| \subseteq B$, and M' is κ -saturated, then $A \downarrow B$. $\bar{M'}$
- (3) Left and right normality: If $A \underset{M}{\downarrow} B$, then $AM \underset{M}{\downarrow} BM$.
- (4) <u>Symmetry</u>: $A \downarrow B$ if and only if $B \downarrow A$. (5) <u>Strong transitivity</u>: If $M_0 \leq \mathfrak{C}$, $M_1 \leq \mathfrak{C}$, $A \downarrow M_1$, and $A \downarrow B$, then $A \downarrow B$ M_0 (note that we do *not* assume that $M_0 \leq M_1^{\circ}$).
- (6) Uniqueness for types of length one: If $M \leq M'$, $p,q \in gS(M')$ are both $(< \kappa)$ -coheir over M and $p \upharpoonright M = q \upharpoonright M$, then p = q.

(7) <u>Set local character</u>: For any α , $\bar{\kappa}_{\alpha}(\downarrow) \leq ((\alpha+2)^{<\kappa_r})^+$.

Moreover K is stable in all $\mu \geq \kappa$ such that $\mu = \mu^{2^{<\kappa_r}}$.

Remark 2.7. We will not use the exact definition of coheir, just that it satisfies the conclusion of Fact 2.6.

Remark 2.8. Strong transitivity will be used in the proof that the relation \downarrow (Definition 2.9) is transitive, see Proposition 2.12. We do not know if transitivity would suffice.

For what comes next, it will be convenient if we could say that $A \bigcup B$ and $M \leq N$ implies $A \downarrow B$. By base monotonicity, this holds if $|N| \subseteq B$ but in general this is not part of our assumptions (and in practice this need not hold). Thus we will close \downarrow under this property. This is where we depart from [MS90]; there the authors used that the singular cardinal hypothesis holds above a strongly compact to prove the result corresponding to our Lemma 3.2. Here we need to be more clever.

Definition 2.9. $A \stackrel{\cdot}{\underset{C}{\to}} B$ means that there exists $M_0 \leq \mathfrak{C}, |M_0| \subseteq C$ such that $A \underset{M_0}{\downarrow} B.$

Remark 2.10. \downarrow need not satisfy the normality property from Fact 2.6.

In what follows, we will apply the definition of κ_{α} and $\bar{\kappa}_{\alpha}$ (Definition 2.4) to other independence relations than coheir.

Definition 2.11. We write $A \underset{C}{\overset{*}{\downarrow}} [B]^1$ to mean that $A \underset{C}{\overset{*}{\downarrow}} b$ for all $b \in B$. Similarly define $[A]^1 \underset{C}{\overset{*}{\downarrow}} B$. Let $(\overset{*}{\downarrow})^1$ denote the relation defined by $A(\underset{C}{\overset{*}{\downarrow}})^1 B$ if and only if $A \stackrel{*}{\underset{C}{\downarrow}} [B]^1$. For α a cardinal, let $\bar{\kappa}^1_{\alpha} = \bar{\kappa}^1_{\alpha} (\stackrel{*}{\downarrow}) := \bar{\kappa}_{\alpha} ((\stackrel{*}{\downarrow})^1)$.

Note that $A \stackrel{*}{\underset{C}{\downarrow}} B$ implies $[A]^1 \stackrel{*}{\underset{C}{\downarrow}} B$ by monotonicity.

Proposition 2.12.

- (1) \downarrow satisfies invariance, monotonicity, symmetry, and strong right transitivity (see Fact 2.6).
- (2) For all α , $\bar{\kappa}_{\alpha}(\stackrel{*}{\downarrow}) = \bar{\kappa}_{\alpha}(\stackrel{*}{\downarrow})$, $\kappa_{\alpha}(\stackrel{*}{\downarrow}) = \kappa_{\alpha}(\stackrel{+}{\downarrow})$. (3) $\stackrel{*}{\downarrow}$ has strong base monotonicity: If $A \stackrel{*}{\downarrow}_{C} B$ and $C \subseteq C'$, then $A \stackrel{*}{\downarrow}_{C'} B$.
- (4) If $A \underset{M}{\downarrow} B$, then $A \underset{M}{\overset{*}{\downarrow}} B$.
- (5) If $A \underset{M}{\overset{*}{\downarrow}} B$ and M is κ -saturated such $|M| \subseteq B$, then $A \underset{M}{\downarrow} B$.
- (6) For all α , $\bar{\kappa}^1_{\alpha}(\overset{*}{\downarrow}) \leq \bar{\kappa}_{\alpha}(\downarrow)$.

 $\mathbf{6}$

Proof. All quickly follow from the definition. As an example, we prove that \downarrow has strong right transitivity. Assume $A \stackrel{*}{\underset{M_0}{\downarrow}} M_1$ and $A \stackrel{*}{\underset{M_1}{\downarrow}} B$. Then there exists $M'_0 \leq M_0$ and $M'_1 \leq M_1$ such that $A \stackrel{+}{\underset{M'_0}{\downarrow}} M_1$ and $A \stackrel{+}{\underset{M'_1}{\downarrow}} B$. By monotonicity for \downarrow , $A \stackrel{+}{\underset{M'_0}{\downarrow}} M'_1$. By strong right transitivity for \downarrow , $A \stackrel{+}{\underset{M'_0}{\downarrow}} B$. Thus M'_0 witnesses $A \stackrel{*}{\underset{M_0}{\downarrow}} B$. \Box

Proposition 2.13. Assume $\langle M_i : i < \delta \rangle$, $\langle N_i : i < \delta \rangle$ are increasing chains of κ -saturated models, A is a set. If $A \underset{M_i}{\downarrow} N_i$ for all $i < \delta$ and $\kappa_{|A|}(\downarrow) \leq \mathrm{cf}(\delta)$, then $A \underset{M_{\delta}}{\downarrow} N_{\delta}$, where² $M_{\delta} := \bigcup_{i < \delta} M_i$ and $N_{\delta} := \bigcup_{i < \delta} N_i$.

Proof. Without loss of generality, $\delta = cf(\delta)$. By definition of $\kappa_{|A|}(\downarrow)$, there exists $i < \delta$ such that $A \underset{N_i}{\downarrow} N_{\delta}$, so $A \underset{N_i}{\overset{*}{\downarrow}} N_{\delta}$. By strong right transitivity for $\overset{*}{\downarrow}$, $A \underset{M_i}{\overset{*}{\downarrow}} N_{\delta}$. By strong base monotonicity, $A \underset{M_{\delta}}{\overset{*}{\downarrow}} N_{\delta}$.

As already discussed, the reason we use $\stackrel{\circ}{\downarrow}$ is that we want to generalize [MS90, 4.17] to our context. In their proof, Makkai and Shelah use that cardinal arithmetic behaves nicely above a strongly compact, and we cannot make use of this fact here. Thus we are only able to prove this lemma for $\stackrel{*}{\downarrow}$ instead of \downarrow (see Lemma 3.2).

Fortunately, this turns out to be enough. The reader can also think of \downarrow as a trick to absorb some quantifiers.

The next lemma imitates [MS90, 4.18].

Lemma 2.14. Let $\lambda_0 \geq \kappa_r$ be regular, let $\lambda > \lambda_0$ be regular such that K is stable in unboundedly-many cardinals below λ and let $\langle M_i : i < \delta \rangle$ be an increasing chain with $M_i \lambda$ -saturated for all $i < \delta$. Assume that $\kappa_1(\downarrow \upharpoonright K^{\lambda_0-\text{sat}}) \leq \text{cf}(\delta)$.


If $\bar{\kappa}^1_{<\lambda}(\overset{*}{\downarrow}) \leq \lambda$, then $M_{\delta} := \bigcup_{i < \delta} M_i$ is λ -saturated.

Proof. Without loss of generality, $\delta = cf(\delta)$. Let $A \subseteq |M_{\delta}|$ have size less than λ . If $\lambda \leq \delta$, then $A \subseteq |M_i|$ for some $i < \delta$ and so any type over A is realized in $M_i \subseteq |M_{\delta}|$. Now assume without loss of generality that $\lambda > \delta$. We need to show every Galois type over A is realized in M_{δ} . Let $\mu := \lambda_0 + \delta$. Note that $\mu = cf(\mu) < \lambda$. First, we build an array of λ_0 -saturated models $\langle N_i^{\alpha} \in K_{<\lambda} : i < \delta, \alpha < \mu \rangle$ such that:

- (1) For all $i < \delta$, $\langle N_i^{\alpha} : \alpha < \mu \rangle$ is increasing.
- (2) For all $\alpha < \mu$, $\langle N_i^{\alpha} : i < \delta \rangle$ is increasing.
- (3) For all $i < \delta$ and all $\alpha < \mu$, $N_i^{\alpha} \le M_i$.
- (4) $A \subseteq \bigcup_{i < \delta} |N_i^0|.$
- (5) For all $\alpha < \mu$ and all $i < \delta$, $\bigcup_{i < \delta} N_i^{\alpha} \bigcup_{N_i^{\alpha+1}}^* [M_i]^1$.

²Note that M_{δ} and N_{δ} need not be κ -saturated.

For $\alpha < \mu$, write $N_{\delta}^{\alpha} := \bigcup_{i < \delta} N_i^{\alpha}$ and for $i \leq \delta$, write $N_i^{\mu} := \bigcup_{\alpha < \mu} N_i^{\alpha}$. The following is a picture of the array constructed.



This is enough: Note that for $i < \delta$, N_i^{μ} is λ_0 -saturated and has size less than λ (since $\lambda > \mu$ and λ is regular). Note also that since $\delta \le \mu < \lambda$, N_{δ}^{μ} has size less than λ (but we do *not* claim that it is λ_0 -saturated).

 $\begin{array}{ll} \textbf{Claim: For all } i < \delta, \ N^{\mu}_{\delta} \ \bigcup_{N^{\mu}_{i}}^{*} [M_{i}]^{1}. \\ \textbf{Proof of claim: Fix } i < \delta \ \text{and let } a \in M_{i}. \ \text{Fix } j < \delta. \ \text{By (5), monotonicity, and} \\ \text{symmetry, } a \ \bigcup_{N^{\alpha+1}_{i}}^{*} N^{\alpha}_{j} \ \text{for all } \alpha < \mu. \ \text{By Proposition 2.13 applied to} \ \langle N^{\alpha+1}_{i} : \alpha < \mu \rangle \\ \text{and } \langle N^{\alpha}_{j} : \alpha < \mu \rangle, \ a \ \bigcup_{N^{\mu}_{i}}^{*} N^{\mu}_{j} \ (\text{note that } \mu = \text{cf}(\mu) \ge \delta \ge \kappa_{1}(\downarrow)). \ \text{Since } j \ \text{was} \\ \text{arbitrary, we can apply Proposition 2.13 again with the constantly } N^{\mu}_{i} \ \text{sequence} \\ \text{and } \langle N^{\mu}_{j} : j < \delta \rangle \ (\text{note that } \delta = \text{cf}(\delta) \ge \kappa_{1}(\downarrow)) \ \text{to get that } a \ \bigcup_{N^{\mu}_{i}}^{*} N^{\mu}_{\delta}. \ \text{By symmetry,} \\ N^{\mu}_{i} \ \bigcup_{N^{\mu}_{i}}^{*} a, \ \text{as desired.} \\ \end{array}$

Now let $p \in \mathrm{gS}(A)$. By (4), $A \subseteq N_{\delta}^{\mu}$ so we can extend p to some $q \in \mathrm{gS}(N_{\delta}^{\mu})$. Since $\delta \geq \kappa_1(\downarrow)$, we can find $i < \delta$ such that q does not fork over N_i^{μ} . Since $N_i^{\mu} \leq M_i$, M_i is λ -saturated, and $||N_i^{\mu}|| < \lambda$, we can find $a \in M_i$ realizing $q \upharpoonright N_i^{\mu}$. Since by the claim $N_{\delta}^{\mu} \bigcup_{N_i^{\mu}}^* [M_i]^1$, we can use symmetry to conclude $a \bigcup_{N_i^{\mu}}^* N_{\delta}^{\mu}$, and hence (Proposition 2.12(5)) $a \bigcup_{N_i^{\mu}}^* N_{\delta}^{\mu}$. By uniqueness for types of length one, a must realize q, so in particular a realizes p. This concludes the proof that M_{δ} is λ -saturated.

This is possible: We define $\langle N_i^{\alpha} : i < \delta \rangle$ by induction on α . For a fixed $i < \delta$, choose any $N_i^0 \leq M_i$ in $K_{<\lambda}$ that contains $A \cap |M_i|$ and is λ_0 -saturated (this is possible since K is stable in unboundedly-many cardinals below λ). For $\alpha < \mu$ limit and $i < \delta$, pick any $N_i^{\alpha} \leq M_i$ containing $\bigcup_{\beta < \alpha} N_i^{\beta}$ which is in $K_{<\lambda}$ and λ_0 -saturated (this is possible for the same reason as in the base case). Now assume $\alpha = \beta + 1 < \mu$, and N_i^{β} has been defined for $i < \delta$. Define N_i^{α} by induction on i. Assume N_j^{α} has been defined for all j < i. Pick N_i^{α} containing $\bigcup_{j < i} N_j^{\alpha}$ which is in $K_{<\lambda}$, is λ_0 -saturated, and satisfies $N_i^{\alpha} \leq M_i$ and (5). This is possible by strong base monotonicity and definition of $\bar{\kappa}_{<\lambda}^1$.

Below, we give a more natural formulation of the hypotheses.

Theorem 2.15. Let $\lambda > \kappa$. Let $\langle M_i : i < \delta \rangle$ be an increasing chain with M_i λ -saturated for all $i < \delta$. If:

(1) $\operatorname{cf}(\delta) \geq \kappa_1(\downarrow)$; and (2) $\chi^{2^{<\kappa_r}} < \lambda$ for all $\chi < \lambda$,

then $\bigcup_{i < \delta} M_i$ is λ -saturated.

Proof. Let $M_{\delta} := \bigcup_{i < \delta} M_i$. Note that $\lambda > \kappa_r$: since $\lambda > \kappa$, $\lambda \ge \kappa^+$ and if $\lambda = \kappa^+$ then $\kappa^{<\kappa} < \lambda$ so $\kappa = \kappa^{<\kappa}$ hence κ is regular: $\kappa_r = \kappa$.

Let $\chi < \lambda$ be such that $\chi^+ > \kappa_r$. We show that M_{δ} is χ^+ -saturated. By hypothesis, $\chi^{2^{<\kappa_r}} < \lambda$, so replacing χ by $\chi^{<\kappa_r}$ if necessary, we might as well assume that $\chi = \chi^{2^{<\kappa_r}}$. We check that χ^+ satisfies the conditions of Lemma 2.14 (with λ_0 there standing for κ_r here) as λ there. By assumption, χ^+ is regular and $\chi^+ > \kappa_r$. Also, K is stable in unboundedly-many cardinals below χ^+ because by the moreover part of Fact 2.6, K is stable in χ .

Now by Proposition 2.12(6), $\bar{\kappa}_{\chi}^{1}(\stackrel{*}{\downarrow}) \leq \bar{\kappa}_{\chi}(\downarrow)$. By Fact 2.6, $\bar{\kappa}_{\chi}(\downarrow) \leq (\chi^{<\kappa_{r}})^{+} = \chi^{+}$. Thus $\bar{\kappa}_{\chi}^{1}(\stackrel{*}{\downarrow}) \leq \chi^{+}$, as needed.

Thus Lemma 2.14 applies and so M_{δ} is χ^+ -saturated. Since $\chi < \lambda$ was arbitrary, M_{δ} is λ -saturated.

For the next corollaries to AECs, we repeat our hypotheses.

Corollary 2.16. Let K be an AEC with amalgamation. Let $\kappa = \beth_{\kappa} > LS(K)$ be such that K is $(< \kappa)$ -tame. Assume that K is stable in some cardinal greater than or equal to κ and let $\langle M_i : i < \delta \rangle$ be an increasing chain of λ -saturated models. If:

- (1) $\operatorname{cf}(\delta) > 2^{<\kappa_r}$.
- (2) $\chi^{2 < \kappa_r} < \lambda$ for all $\chi < \lambda$.

Then $\bigcup_{i < \delta} M_i$ is λ -saturated.

Proof. Without loss of generality, $\delta = cf(\delta) < \lambda$. Also without loss of generality, K has joint embedding (otherwise, partition it into disjoint classes, each of which has joint embedding), and arbitrarily large models (since K has a model of cardinality

 $\kappa = \beth_{\kappa} > LS(K)$). Therefore Hypothesis 2.2, and hence the conclusion of Fact 2.6, hold.

Note (Remark 2.5) that $\kappa_1(\downarrow) \leq \bar{\kappa}_1(\downarrow) \leq (2^{<\kappa_r})^+$. Now use Theorem 2.15. \Box

3. Using independence calculus: the superstable case

Next we show that in the *superstable* case we can remove the cardinal arithmetic condition (2) in Corollary 2.16.

Hypothesis 3.1. Same as in the previous section: Hypothesis 2.2.

In the proof of Theorem 2.15, we estimated $\bar{\kappa}^1_{\alpha}(\downarrow)$ using $\bar{\kappa}_{\alpha}(\downarrow)$. Using superstability, we can prove a better bound. This is adapted from [MS90, 4.17].

Lemma 3.2. Assume that $\kappa_1(\downarrow) = \aleph_0$ and K is stable in all $\lambda \ge \kappa_r$. Then for any cardinal α , $\bar{\kappa}_{\alpha}^1(\downarrow) \le \bar{\kappa}_{\kappa_r}(\downarrow) + \alpha^+$.

Proof. Let A have size α and N be a κ -saturated model. We show by induction on α that there exists an $M \leq N$ with $||M|| < \mu := \bar{\kappa}_{\kappa_r}(\bot) + \alpha^+$ and $A \bigcup_{M}^* [N]^1$. Note that $\mu > \kappa_r$.

If $\alpha \leq \kappa_r$, then apply the definition of $\bar{\kappa}_{\kappa_r}(\downarrow)$ to get a $M \leq N$ with $||M|| < \bar{\kappa}_{\kappa_r}(\downarrow)$, $A \downarrow_M^* N$, which is more than what we need.

Now, assume $\alpha > \kappa_r$, and that the result has been proven for all $\alpha_0 < \alpha$. Closing A to a κ -saturated model (using the stability assumptions) if necessary, we can assume without loss of generality that A is a κ -saturated model. Let $\langle A_i : i < \alpha \rangle$ be an increasing resolution of A such that A_i is κ -saturated in $K_{<\alpha}$ for all $i < \alpha$. Now define an increasing chain $\langle M_i : i < \alpha \rangle$ such that for all $i < \alpha$:

(1) $M_i \in K_{<\mu}$ and M_i is κ -saturated. (2) $M_i \leq N$. (3) $A_i \downarrow_{M_i}^* [N]^1$.

<u>This is possible</u>: For $i < \alpha$, use the induction hypothesis to find $M_i \leq N$ such that $A_i \underset{M_i}{\overset{*}{\downarrow}} [N]^1$ and $||M_i|| < \mu$. By strong base monotonicity of $\overset{*}{\downarrow}$ and the closure assumption, we can assume that M_i contains $\bigcup_{i < i} M_i$.

This is enough: Let $M \in K_{<\mu}$ be κ -saturated and contain $\bigcup_{i < \alpha} M_i$. We claim that $A \downarrow_M^* [N]^1$. Let $a \in N$. By symmetry, it is enough to see $a \downarrow_M^* A$. This follows from strong base monotonicity and Proposition 2.13 applied to $\langle M_i : i < \alpha \rangle$ and $\langle A_i : i < \alpha \rangle$ since $\kappa_1(\downarrow) = \aleph_0 \leq \operatorname{cf}(\alpha)$ by Proposition 2.12(2) and the hypothesis. \Box

Remark 3.3. The heavy use of strong base monotonicity in the above proof was the reason for introducing $\overset{*}{\bigcup}$.

Theorem 3.4. Let $\lambda_0 \geq \kappa_r$ be regular. Assume that $\kappa_1(\downarrow \upharpoonright K^{\lambda_0\text{-sat}}) = \aleph_0$ and K is stable in all $\lambda \geq \lambda_0$. Let $\lambda \geq \bar{\kappa}_{\kappa_r}(\downarrow) + \lambda_0^+$.

Let $\langle M_i : i < \delta \rangle$ be an increasing chain with $M_i \lambda$ -saturated for all $i < \delta$. Then $M_{\delta} := \bigcup_{i < \delta} M_i$ is λ -saturated.

Proof. Let $\chi < \lambda$ be such that $\chi^+ \geq \bar{\kappa}_{\kappa_r}(\downarrow) + \lambda_0^+$. We claim that χ^+ satisfies the hypotheses of Lemma 2.14 (as λ there). Indeed by Lemma 3.2, $\bar{\kappa}_{\chi}^1(\downarrow) \leq \bar{\kappa}_{\kappa_r}(\downarrow) + \chi^+ = \chi^+$.

Thus Lemma 2.14 applies: M_{δ} is χ^+ -saturated. Since $\chi < \lambda$ was arbitrary, M_{δ} is λ -saturated.

For the next corollary to AECs, we drop our hypotheses.

Corollary 3.5. Let K be an AEC with amalgamation and no maximal models. Let $\kappa = \beth_{\kappa} > \mathrm{LS}(K)$ be such that K is $(< \kappa)$ -tame. If K is categorical in some cardinal strictly above κ , then for all $\lambda > 2^{\kappa}$, $K^{\lambda\text{-sat}}$ is an AEC with Löwenheim-Skolem number λ .

Proof. Using categoricity and amalgamation, it is easy to check that K has joint embedding. Let $\lambda_0 := \kappa^+$. By [Vas16a, 10.8,10.16], K is stable in all $\mu \geq \kappa$ and $\kappa_1(\downarrow \upharpoonright K^{\lambda_0\text{-sat}}) = \aleph_0$. In particular, Hypothesis 2.2 holds. Remembering (Fact 2.6) that $\bar{\kappa}_{\kappa_r}(\downarrow) \leq \kappa_r^{<\kappa_r} \leq 2^{\kappa}$, we obtain the result from Theorem 3.4 (to show that $\mathrm{LS}(K^{\lambda\text{-sat}}) = \lambda$, imitate the proof of [She90, III.3.12]).

4. Averages

In this section, we write in the framework of stability theory inside a model:

Hypothesis 4.1.

- (1) κ is an infinite cardinal.
- (2) L is a $(< \kappa)$ -ary language.
- (3) \mathcal{N} is a fixed *L*-structure.
- (4) We work inside \mathcal{N} .
- (5) Hypotheses 4.2 and 4.4, see the discussion below.

Midway through, we will also assume Hypothesis 4.22.

We use the same notation and convention as [Vas16c, Section 2]: although we may forget to say it, we always work with quantifier-free $L_{\kappa,\kappa}$ formulas and types (so the arity of all the variables inside a given formula is less than κ). Also, since we work inside \mathcal{N} , everything is defined relative to \mathcal{N} . For example $\operatorname{tp}(\bar{c}/A)$ means $\operatorname{tp}_{qL_{\kappa,\kappa}}(\bar{c}/A;\mathcal{N})$, the quantifier-free $L_{\kappa,\kappa}$ -type of \bar{c} over A, and saturated means saturated in \mathcal{N} . Similarly, we write $\models \phi[\bar{b}]$ instead of $\mathcal{N} \models \phi[\bar{b}]$. By "type", we mean a member of $S^{<\infty}(A)$ for some set A. Whenever we mention a set of formulas (meaning a possibly incomplete type), we mean a $(L_{\kappa,\kappa}$ -quantifier free) set of formulas that is satisfiable by an element in \mathcal{N} . Unless otherwise noted, the letters \bar{a} , \bar{b} , \bar{c} denote tuples of elements of length less than κ . The letters A, B, C, will denote subsets of \mathcal{N} . We say $\langle A_i : i < \delta \rangle$ is increasing if $A_i \subseteq A_j$ for all $i < j < \delta$.

We say \mathcal{N} is α -stable in λ if $|S^{\alpha}(A)| \leq \lambda$ for all A with $|A| \leq \lambda$ (the default value is for α is 1). We say \mathcal{N} has the order property of length χ if there exists a (quantifierfree) formula $\phi(\bar{x}, \bar{y})$ and elements $\langle \bar{a}_i : i < \chi \rangle$ of the same arity (less than κ) such that for $i, j < \chi, \models \phi[\bar{a}_i, \bar{a}_j]$ if and only if i < j.

Boldface letters like \mathbf{I} , \mathbf{J} will always denote sequences of tuples of the same arity (less than κ). We will use the corresponding non-boldface letter to denote the linear ordering indexing the sequence (writing for example $\mathbf{I} = \langle \bar{a}_i : i \in I \rangle$, where I is a linear order). We sometimes treat such sequences as sets of tuples, writing statements like $\bar{a} \in \mathbf{I}$, but then we are really looking at the range of the sequence. To avoid potential mistakes, we do not necessarily assume that the elements of \mathbf{I} are all distinct although it should always hold in cases of interest. We write $|\mathbf{I}|$ for the cardinality of the range, i.e. the number of distinct elements in \mathbf{I} . We will sometimes use the interval notation on linear order. For example, if I is a linear order and $i \in I$, $[i, \infty)_I := \{j \in I \mid j \geq i\}$.

As the reader will see, this section builds on earlier work of Shelah from [She09b, Chapter V.A]. Note that Shelah works in an arbitrary logic. We work only with quantifier-free $L_{\kappa,\kappa}$ -formulas in order to be concrete and because this is the case we are interested in to translate the syntactic results to AECs.

The reader may wonder what the right notion of submodel is in this context. We could simply say that it is "subset" but this does not quite work when translating to AECs. Thus we fix a set of subsets of \mathcal{N} that by definition will be the substructures of \mathcal{N} . We require that this set satisfies some axioms akin to those of AECs. This can be taken to be the full powerset if one is not interested in doing an AEC translation.

Hypothesis 4.2. $S \subseteq \mathcal{P}(|\mathcal{N}|)$ is a fixed set of subsets of \mathcal{N} satisfying:

- (1) Closure under chains: If $\langle A_i : i < \delta \rangle$ is an increasing sequence of members of S, then $\bigcup_{i < \delta} A_i$ is in S.
- (2) Löwenheim-Skolem axiom: If $A \subseteq B$ are sets and $B \in S$, there exists $A' \in S$ such that $A \subseteq A' \subseteq B$ and $|A'| \leq (|L|+2)^{<\kappa} + |A|$.

We exclusively use the letters M and N to denote elements of S and call such elements *models*. We pretend they are *L*-structures and write |M| and |N| for their universe and ||M|| and ||N|| for their cardinalities.

Remark 4.3. An element M of S is *not* required to be an *L*-structure. Note however that if it is κ -saturated for types of length less than κ (see below), then it will be one.

We also need to discuss the definition of saturated: define M to be λ -saturated for types of length α if for any $A \subseteq |M|$ of size less than λ , any $p \in S^{\alpha}(A)$ is realized in M. Similarly define λ -saturated for types of length less than α . Now in the framework we are working in, μ -saturated for types of length less than κ seems to be the right notion, so we say that M is μ -saturated if it is μ -saturated for types of length less than κ . Unfortunately it is not clear that it is equivalent to μ -saturated for types of length one (or length less than ω), even when $\mu > \kappa$. However [She09a, II.1.14] (the "model-homogeneity = saturativity" lemma) tells us that in case \mathcal{N} comes from an AEC, then this is the case. Thus we will make the following additional assumption. Note that it is possible to work without it, but then everywhere below "stability" must be replaced by "($< \kappa$)-stability".

Hypothesis 4.4. If $\mu > (|L|+2)^{<\kappa}$, then whenever M is μ -saturated for types of length one, it is μ -saturated (for types of length less than κ).

Our goal in this section is to use Shelah's notion of average in this framework to prove a result about chains of saturated models. Recall:

Definition 4.5 (Definition V.A.2.6 in [She09b]). For **I** a sequence, χ an infinite cardinal such that $|\mathbf{I}| \geq \chi$, and A a set, define $\operatorname{Av}_{\chi}(\mathbf{I}/A)$ to be the set of formulas $\phi(\bar{x})$ over A so that the set $\{\bar{b} \in \mathbf{I} \mid \models \neg \phi[\bar{b}]\}$ has size less than χ .

Note that if $|\mathbf{I}| \geq \chi$ (say all the elements of \mathbf{I} have the same arity α) and $\phi(\bar{x})$ is a formula with $\ell(\bar{x}) = \alpha$, then at most one of ϕ , $\neg \phi$ is in $\operatorname{Av}_{\chi}(\mathbf{I}/A)$. Thus the average is not obviously contradictory, but we do not claim that there is an element in \mathcal{N} realizing it. Also, $\operatorname{Av}_{\chi}(\mathbf{I}/A)$ might be empty. However, we give conditions below (see Fact 4.13 and Theorem 4.21) where it is in fact complete (i.e. exactly one of ϕ and $\neg \phi$ is in the average).

The next lemma is a simple counting argument allowing us to find such an element:

Lemma 4.6. Let **I** be a sequence with $|\mathbf{I}| \geq \chi$ and let A be a set. Let $p := \operatorname{Av}_{\chi}(\mathbf{I}/A)$. Assume that

$$|\mathbf{I}| > \chi + \min((|A| + |L| + 2)^{<\kappa}, |S^{\ell(p)}(A)|)$$

Then there exists $\bar{b} \in \mathbf{I}$ realizing p.

Proof. Assume first that the minimum is realized by $(|A|+|L|+2)^{<\kappa}$. By definition of the average, for every every formula $\phi(\bar{x}) \in p$, $\mathbf{J}_{\phi} := \{\bar{b} \in \mathbf{I} \mid \models \neg \phi[\bar{b}]\}$ has size less than χ . Let $\mathbf{J} := \bigcup_{\phi \in p} \mathbf{J}_{\phi}$. Note that $|\mathbf{J}| \leq \chi + (|A|+|L|+2)^{<\kappa}$ and by definition any $\bar{b} \in \mathbf{I} \setminus \mathbf{J}$ realizes p.

Now assume that the minimum is realized by $|S^{\ell(p)}(A)|$. Let $\mu := \chi + |S^{\ell(p)}(A)|$. By the pigeonhole principle, there exists $\mathbf{I}_0 \subseteq \mathbf{I}$ of size μ^+ such that $\bar{c}, \bar{c}' \in \mathbf{I}_0$ implies $q := \operatorname{tp}(\bar{c}/A) = \operatorname{tp}(\bar{c}'/A)$. We claim that $p \subseteq q$, which is enough: any $\bar{b} \in \mathbf{I}_0$ realizes p. If not, there exists $\phi(\bar{x}) \in p$ such that $\neg \phi(\bar{x}) \in q$. By definition of the average, fewer than χ -many elements of \mathbf{I} satisfy $\neg \phi(\bar{x})$. However, $\neg \phi(\bar{x})$ is in q which means that it is realized by all the elements of \mathbf{I}_0 and $|\mathbf{I}_0| = \mu^+ > \chi$, a contradiction. \Box

We now recall the definition of splitting and study how it interacts with averages.

Definition 4.7. A set of formulas p splits over A if there exists $\phi(\bar{x}, \bar{b}) \in p$ and \bar{b}' with $\operatorname{tp}(\bar{b}'/A) = \operatorname{tp}(\bar{b}/A)$ and $\neg \phi(\bar{x}, \bar{b}') \in p$.

The following result is classical:

Lemma 4.8 (Uniqueness for nonsplitting). Let $A \subseteq |M| \subseteq B$. Assume p, q are complete sets of formulas (say in the variable \bar{x} , with $\ell(\bar{x}) < \kappa$) over B that do not split over A and M is $|A|^+$ -saturated. If $p \upharpoonright M = q \upharpoonright M$, then p = q.

Proof. Let $\phi(\bar{x}, \bar{b}) \in p$ with $\bar{b} \in B$. We show $\phi(\bar{x}, \bar{b}) \in q$ and the converse is symmetric. By saturation³, find $\bar{b}' \in M$ such that $\operatorname{tp}(\bar{b}'/A) = \operatorname{tp}(\bar{b}/A)$. Since p does not split over A, $\phi(\bar{x}, \bar{b}') \in p$. Since $p \upharpoonright M = q \upharpoonright M$, $\phi(\bar{x}, \bar{b}') \in q$. Since again q does not split, $\phi(\bar{x}, \bar{b}) \in q$.

We would like to study when the average is a nonsplitting extension. This is the purpose of the next definition.

Definition 4.9. I is χ -based on A if for any B, $Av_{\chi}(I/B)$ does not split over A.

The next lemma tells us that any sequence is based on a set of small size.

Lemma 4.10 (IV.1.23(2) in [She09a]). If **I** is a sequence and $\mathbf{J} \subseteq \mathbf{I}$ has size at least χ , then **I** is χ -based on **J**.

Proof. Let *B* be a set. Let $p := \operatorname{Av}_{\chi}(\mathbf{I}/B)$. Note that $p \subseteq \operatorname{Av}_{\chi}(\mathbf{J}/B)$. Let $\bar{b}, \bar{b}' \in B$ be such that $\operatorname{tp}(\bar{b}/\mathbf{J}) = \operatorname{tp}(\bar{b}'/\mathbf{J})$. Assume $\phi(\bar{x}, \bar{b}) \in p$. Then since $p \subseteq \operatorname{Av}_{\chi}(\mathbf{J}/B)$, let $\bar{a} \in \mathbf{J}$ be such that $\models \phi[\bar{a}, \bar{b}]$. Since $\bar{a} \in \mathbf{J}, \models \phi[\bar{a}, \bar{b}']$. Since there are at least χ -many such \bar{a} 's, $\neg \phi(\bar{x}, \bar{b}') \notin p$.

We know that *at most* one of ϕ , $\neg \phi$ is in the average. It is very desirable to have that *exactly* one is in, i.e. that the average is a *complete* type. This is the purpose of the next definition. Recall from the beginning of this section that **I** always denotes a sequence of elements of the same arity less than κ .

Definition 4.11 (V.A.2.1 in [She09b]). A sequence **I** is said to be χ -convergent if $|\mathbf{I}| \geq \chi$ and for any set A, $\operatorname{Av}_{\chi}(\mathbf{I}/A)$ is a complete type over A. That is, whenever $\phi(\bar{x})$ is a formula with $\ell(\bar{x})$ equal to the arity of all the elements of **I**, then we have that exactly one of ϕ or $\neg \phi$ is in $\operatorname{Av}_{\chi}(\mathbf{I}/A)$.

Remark 4.12 (Monotonicity). If **I** is χ -convergent, $\mathbf{J} \subseteq \mathbf{I}$, and $|\mathbf{J}| \ge \chi' \ge \chi$, then for any set A, $\operatorname{Av}_{\chi}(\mathbf{I}/A) = \operatorname{Av}_{\chi'}(\mathbf{J}/A)$. In particular, **J** is χ' -convergent.

Recall [She90, III.1.7(1)] that if T is a first-order stable theory and \mathbf{I} is an infinite sequence of indiscernibles (in its monster model), then \mathbf{I} is \aleph_0 -convergent. The proof relies heavily on the compactness theorem. We would like a replacement of the form "if \mathcal{N} has some stability and \mathbf{I} is nice, then it is convergent. The next result is key. It plays the same role as the ability to extract indiscernible subsequences in first-order stable theories.

Fact 4.13 (The convergent set existence theorem: V.A.2.8 in [She09b]). Let $\chi_0 \geq (|L|+2)^{<\kappa}$ be such that \mathcal{N} does not have the order property of length χ_0^+ . Let μ be an infinite cardinal such that $\mu = \mu^{\chi_0} + 2^{2^{\chi_0}}$.

Let I be a sequence with $|I| = \mu^+$. Then there is $J \subseteq I$ of size μ^+ which is χ_0 -convergent.

³Note that we are really using saturation for types of length less than κ here.

However having to extract a subsequence every time is too much for us. One issue is with the cardinal arithmetic condition on μ : what if we have a sequence of length μ^+ when μ is a singular cardinal of low cofinality? We work toward proving a more constructive result: *Morley* sequences (defined below) are always convergent. The parameters represent respectively a bound on the size of A, the degree of saturation of the models, and the length of the sequence. They will be assigned default values in Hypothesis 4.22.

Definition 4.14. We say $\langle \bar{a}_i : i \in I \rangle \frown \langle N_i : i \in I \rangle$ is a (χ_0, χ_1, χ_2) -Morley sequence for p over A if:

- (1) $\chi_0 \leq \chi_1 \leq \chi_2$ are infinite cardinals, I is a linear order, A is a set, $p(\bar{x})$ is a set of formulas with parameters and $\ell(\bar{x}) < \kappa$, and there is $\alpha < \kappa$ such that for all $i \in I$, $\bar{a}_i \in {}^{\alpha}\mathcal{N}$.
- (2) For all $i \in I$, $A \subseteq |N_i|$ and $|A| < \chi_0$.
- (3) $\langle N_i : i \in I \rangle$ is increasing, and each N_i is χ_1 -saturated.
- (4) For all $i \in I$, \bar{a}_i realizes⁴ $p \upharpoonright N_i$ and for all j > i in I, $\bar{a}_i \in {}^{\alpha}N_j$.
- (5) i < j in I implies $\bar{a}_i \neq \bar{a}_j$.
- (6) $|I| \ge \chi_2$.
- (7) For all i < j in I, $\operatorname{tp}(\bar{a}_i/N_i) = \operatorname{tp}(\bar{a}_j/N_i)$.
- (8) For all $i \in I$, $\operatorname{tp}(\bar{a}_i/N_i)$ does not split over A.

When p or A is omitted, we mean "for some p or A". We call $\langle N_i : i \in I \rangle$ the witnesses to $\mathbf{I} := \langle \bar{a}_i : i \in I \rangle$ being Morley, and when we omit them we simply mean that $\mathbf{I} \cap \langle N_i : i \in I \rangle$ is Morley for some witnesses $\langle N_i : i \in I \rangle$.

Remark 4.15 (Monotonicity). Let $\langle \bar{a}_i : i \in I \rangle \frown \langle N_i : i \in I \rangle$ be (χ_0, χ_1, χ_2) -Morley for p over A. Let $\chi'_0 \ge \chi_0, \chi'_1 \le \chi_1$, and $\chi'_2 \le \chi_2$. Let $I' \subseteq I$ be such that $|I'| \ge \chi'_2$, then $\langle \bar{a}_i : i \in I' \rangle \frown \langle N_i : i \in I' \rangle$ is $(\chi'_0, \chi'_1, \chi'_2)$ -Morley for p over A.

Remark 4.16. By the proof of [She90, I.2.5], a Morley sequence is indiscernible (this will not be used).

The next result tells us how to build Morley sequences inside a given model:

Lemma 4.17. Let $A \subseteq |M|$ and let $\chi \ge (|L|+2)^{<\kappa}$ be such that $|A| \le \chi$. Let $p \in S^{\alpha}(M)$ be nonalgebraic (that is, $a_i \notin |M|$ for all $i < \alpha$ for any \bar{a} realizing p) such that p does not split over A, and let $\mu > \chi$. If:

- (1) M is μ^+ -saturated.
- (2) \mathcal{N} is stable in μ .

Then there exists $\langle \bar{a}_i : i < \mu^+ \rangle \land \langle N_i : i < \mu^+ \rangle$ inside M which is (χ^+, χ^+, μ^+) -Morley for p over A.

Proof. We build $\langle \bar{a}_i : i < \mu^+ \rangle$ and $\langle N_i : i < \mu^+ \rangle$ increasing such that for all $i < \mu^+$:

- (1) $A \subseteq |N_0|$.
- (2) $|N_i| \subseteq |M|$.
- (3) $||N_i|| \le \mu$.
- (4) N_i is χ^+ -saturated.

⁴Note that dom(p) might be smaller than N_i .

(5) $\bar{a}_i \in {}^{\alpha}N_{i+1}$.

(6) \bar{a}_i realizes $p \upharpoonright N_i$.

This is enough by definition of a Morley sequence (note that for all $i < \mu^+$, $\bar{a}_i \notin {}^{\alpha}N_i$ by nonalgebraicity of p, so $\bar{a}_i \neq \bar{a}_j$ for all j < i).

This is possible: assume inductively that $\langle \bar{a}_j : j < i \rangle \land \langle N_j : j < i \rangle$ has been defined. Pick $N_i \subseteq M$ which is χ^+ -saturated, has size $\leq \mu$, and contains $A \cup \bigcup_{j < i} N_j$. Such an N_i exists: simply build an increasing chain $\langle M_k : k < \chi^+ \rangle$ with $M_0 := A \cup \bigcup_{j < i} N_j$, $||M_k|| \leq \mu$, and M_k realizing all elements of $S(\bigcup_{k' < k} M_{k'})$ (this is where we use stability in μ). Then $N_i := \bigcup_{k < \chi^+} M_k$ is as desired (we are using Hypothesis 4.4 to deduce that it is χ^+ -saturated for types of length less than κ). Now pick $\bar{a}_i \in {}^{\alpha}M$ realizing $p \upharpoonright N_i$ (exists by saturation of M).

Before proving that Morley sequences are convergent (Theorem 4.21), we prove several useful lemmas:

Lemma 4.18. Let $\mathbf{I} := \langle \bar{a}_i : i \in I \rangle$ be (χ_0, χ_1, χ) -Morley, as witnessed by $\langle N_i : i \in I \rangle$. $I \rangle$. Let $i \in I$ be such that $[j, \infty)_I$ has size at least χ . Then $\operatorname{Av}_{\chi}(\mathbf{I}/N_i) \subseteq \operatorname{tp}(\bar{a}_i/N_i)$.

Proof. Let $\phi(\bar{x})$ be a formula over N_i with $\ell(\bar{x}) = \ell(\bar{a}_i)$. Assume $\phi(\bar{x}) \in \operatorname{Av}_{\chi}(\mathbf{I}/N_i)$. By definition of average and assumption there exists $j \in [i, \infty)$ such that $\models \phi[\bar{a}_j]$. By (7) in Definition 4.14, $\models \phi[\bar{a}_i]$ so $\phi(\bar{x}) \in \operatorname{tp}(\bar{a}_i/N_i)$.

Lemma 4.19. Let *I* be a linear order and let $\chi < |I|$ be infinite. Then there exists $i \in I$ such that both $(-\infty, i]_I$ and $[i, \infty)_I$ have size at least χ .

Proof. Without loss of generality, $|I| = \chi^+$. Let $I_0 := \{i \in I \mid |(-\infty, i]_I| < \chi\}$ and let $I_1 := \{i \in I \mid |[i, \infty)_I| < \chi\}$. Assume the conclusion of the lemma fails. Then $I_0 \cup I_1 = I$. Thus either $|I_0| = \chi^+$ or $|I_1| = \chi^+$. Assume that $|I_0| = \chi^+$, the proof in case $|I_1| = \chi^+$ is symmetric. Let $\delta := \operatorname{cf}(I_0)$ and let $\langle a_\alpha \in I_0 : \alpha < \delta \rangle$ be a cofinal sequence. If $\delta < \chi^+$, then, since $I_0 = \bigcup_{\alpha < \delta} (-\infty, a_\alpha)_I$ has size χ^+ , there is $\alpha < \delta$ such that $|(-\infty, a_\alpha)_I| = \chi^+$. If $\delta \ge \chi^+$, then $|(-\infty, a_\chi)_I| \ge \chi$. Either of these contradict the definition of I_0 .

Lemma 4.20. Let **I** be (χ^+, χ^+, χ^+) -Morley over A (for some type). If **I** is χ -convergent, then **I** is χ -based on A.

Proof. Let $\mathbf{I} := \langle \bar{a}_i : i \in I \rangle$ and let $\langle N_i : i \in I \rangle$ witness that \mathbf{I} is χ -Morley over A. By assumption, $|I| \ge \chi^+$, so let $i \in I$ be as given by Lemma 4.19: both $(-\infty, i]_I$ and $[i, \infty)_I$ have size at least χ . By Lemma 4.10 and the definition of i, we can find $A' \subseteq |N_i|$ containing A of size at most χ such that \mathbf{I} is χ -based on A'.

Let $p := \operatorname{Av}_{\chi}(\mathbf{I}/\mathcal{N})$. Assume for a contradiction that p splits over A and pick witnesses such that $\phi(\bar{x}, \bar{b}), \neg \phi(\bar{x}, \bar{b}') \in p$ and $\operatorname{tp}(\bar{b}/A') = \operatorname{tp}(\bar{b}'/A')$. Note that $p \upharpoonright N_i = \operatorname{tp}(\bar{a}_i/N_i)$ by convergence and Lemma 4.18. Since N_i is χ^+ -saturated, we can find $\bar{b}'' \in {}^{<\kappa}|N_i|$ such that $\operatorname{tp}(\bar{b}''/A') = \operatorname{tp}(\bar{b}/A')$. Now either $\phi(\bar{x}, \bar{b}'') \in p$ or $\neg \phi(\bar{x}, \bar{b}'') \in p$. If $\phi(\bar{x}, \bar{b}'') \in p$, then $\phi(\bar{x}, \bar{b}''), \neg \phi(\bar{x}, \bar{b})$ witness that p splits over Aand if $\neg \phi(\bar{x}, \bar{b}'') \in q$, then $\phi(\bar{x}, \bar{b}), \neg \phi(\bar{x}, \bar{b}'')$ witness the splitting. Either way, we can replace \bar{b} or \bar{b}' by \bar{b}'' . So (swapping the role of \bar{b} and \bar{b}' if necessary), assume without loss of generality that $\bar{b}'' = \bar{b}$ (so $\bar{b} \in {}^{<\kappa}|N_i|$).

16

By definition of a Morley sequence, $p \upharpoonright N_i$ does not split over A, so $\bar{b}' \notin \langle \kappa | N_i |$. Let $p'_i := p \upharpoonright N_i \cup \{\phi(\bar{x}, \bar{b}), \phi(\bar{x}, \bar{b}')\}$. We claim that p'_i does not split over A: if it does, since $\phi(\bar{x}, \bar{b}')$ is the only formula of p'_i with parameters outside of N_i , the splitting must be witnessed by $\phi(\bar{x}, \bar{c}), \neg \phi(\bar{x}, \bar{c}')$, and one of them must be outside N_i , so $\bar{c} = \bar{b}'$. Now $\operatorname{tp}(\bar{b}/A) = \operatorname{tp}(\bar{b}'/A) = \operatorname{tp}(\bar{c}'/A)$, and we have $\bar{c}' \in \langle \kappa | N_i |$ so by nonsplitting of $p \upharpoonright N_i$, also $\neg \phi(\bar{x}, \bar{b}) \in p \upharpoonright N_i$. This is a contradiction since we know $\phi(\bar{x}, \bar{b}) \in p \upharpoonright N_i$.

Now, since **I** is χ -based on A', p does not split over A' and by monotonicity p'_i also does not split over A'. Now use the proof of Lemma 4.8 (with $M = N_i$) to get a contradiction.

We are now ready to prove the relationship between Morley and convergent:

Theorem 4.21. Let $\chi_0 \geq (|L|+2)^{<\kappa}$ be such that \mathcal{N} does not have the order property of length χ_0^+ . Let $\chi := (2^{2^{\chi_0}})^+$.

If **I** is a $(\chi_0^+, \chi_0^+, \chi)$ -Morley sequence, then **I** is χ -convergent.

Proof. Write $\mathbf{I} = \langle \bar{a}_i : i \in I \rangle$ and let $\langle N_i : i \in I \rangle$ witness that it is Morley for p over A.

Assume for a contradiction that \mathbf{I} is *not* χ -convergent. Then there exists a formula $\phi(\bar{x})$ (over \mathcal{N}) and linear orders $I_{\ell} \subseteq I$, $\ell = 0, 1$ such that $|I_{\ell}| = \chi$ and $i \in I_{\ell}$ implies⁵ $\models \phi^{\ell}[\bar{a}_i]$. By Fact 4.13, we can assume without loss of generality that $\mathbf{I}_{\ell} := \langle \bar{a}_i : i \in I_{\ell} \rangle$ is χ_0 -convergent. By Lemma 4.20 (with χ_0 here standing for χ there), \mathbf{I}_{ℓ} is χ_0 -based on A for $\ell = 0, 1$. Let $p_{\ell} := \operatorname{Av}_{\chi_0}(\mathbf{I}_{\ell}/\mathcal{N})$. Since \mathbf{I}_{ℓ} is χ_0 -based on A, p_{ℓ} does not split over A. By Lemma 4.19, pick $i_{\ell} \in I$ so that $|(i, \infty)_{I_{\ell}}| \geq \chi_0$ for $\ell = 0, 1$. let $i := \min(i_0, i_1)$. By Lemma 4.18 and convergence, $p_{\ell} \upharpoonright N_{i_{\ell}} = \operatorname{tp}(a_{i_{\ell}}/N_{i_{\ell}})$ so $p_{\ell} \upharpoonright N_i = \operatorname{tp}(a_{i_{\ell}}/N_i) = \operatorname{tp}(a_i/N_i)$, so $p_0 \upharpoonright N_i = p_1 \upharpoonright N_i$. By assumption, N_i is χ_0^+ -saturated. By uniqueness for nonsplitting (Lemma 4.8), $p_0 = p_1$. However $\phi(\bar{x}) \in p_0$ while $\neg \phi(\bar{x}) \in p_1$, contradiction.

From now on we assume:

Hypothesis 4.22.

- (1) $\chi_0 \ge (|L|+2)^{<\kappa}$ is an infinite cardinal.
- (2) \mathcal{N} does not have the order property of length χ_0^+ .
- (3) $\chi := (2^{2^{\chi_0}})^+$.
- (4) The default parameters for Morley sequences are (χ⁺₀, χ⁺, χ⁺), and the default parameter for averages and convergence is χ. That is, Morley means (χ⁺₀, χ⁺, χ⁺)-Morley, convergent means χ-convergent, Av(I/A) means Av_χ(I/A), and based means χ-based.

Note that Theorem 4.21 and Hypothesis 4.22 imply that any Morley sequence is convergent. Moreover by Lemma 4.20, any Morley sequence over A is based on A. We will use this freely.

⁵Where ϕ^0 stands for ϕ , ϕ^1 for $\neg \phi$.

Before studying chains of saturated models, we generalize Lemma 4.20 to independence notions that are very close to splitting (the reason has to do with the translation to AECs (Section 5)):

Definition 4.23. A splitting-like notion is a binary relation R(p, A), where $p \in S^{<\infty}(B)$ for some set B and $A \subseteq B$, satisfying the following properties:

- (1) Monotonicity: If $A \subseteq A' \subseteq B_0 \subseteq B$, $p \in S^{<\infty}(B)$, and R(p, A), then $R(p \upharpoonright B_0, A')$.
- (2) Weak uniqueness: If $A \subseteq |M| \subseteq B$, M is $(|A| + (|L| + 2)^{<\kappa})^+$ -saturated, and for $\ell = 1, 2, q_{\ell} \in S^{<\infty}(B), R(q_{\ell}, A)$, and $q_1 \upharpoonright M = q_2 \upharpoonright M$, then $q_1 = q_2$.
- (3) R extends nonsplitting: If $p \in S^{<\infty}(B)$ does not split over $A \subseteq B$, then R(p, A).

We also say "p does not R-split over A" instead of R(p, A).

Remark 4.24. If R(p, A) holds if and only if p does not split over A, then R is a splitting-like notion: monotonicity is easy to check and R is nonsplitting. Weak uniqueness is Lemma 4.8.

Lemma 4.25. Let R be a splitting-like notion. Let $p \in S^{<\kappa}(B)$ be such that p does not R-split over $A \subseteq B$ with $|A| \leq \chi_0$.

Let $\mathbf{I} := \langle \bar{a}_i : i \in I \rangle \frown \langle N_i : i \in I \rangle$ be Morley for p over A. If $|\bigcup_{i \in I} N_i| \subseteq B$, then $\operatorname{Av}(\mathbf{I}/B) = p$.

Proof. Since **I** is Morley, **I** is convergent. By Lemma 4.20, **I** is based on A. Thus we have that $\operatorname{Av}(\mathbf{I}/B)$ does not split over A, so it does not R-split over A. Let $i \in I$ be such that $|(i, \infty)_I| \geq \chi$ (use Lemma 4.19). Then $\operatorname{Av}(\mathbf{I}/N_i) = \operatorname{tp}(\bar{a}_i/N_i) = p \upharpoonright N_i$ by Lemma 4.18. By the weak uniqueness axiom of splitting-like relations (with N_i here standing for M there), $\operatorname{Av}(\mathbf{I}/B) = p$.

To construct Morley sequences, we will also use:

Fact 4.26.

- (1) If $\mu = \mu^{\chi_0} + 2^{2^{\chi_0}}$, then \mathcal{N} is $(< \kappa)$ -stable in μ .
- (2) Let M be χ_0^+ -saturated. Then for any $p \in S^{<\kappa}(M)$, there exists $A \subseteq |M|$ of size at most χ_0 such that p does not split over A.

Proof. The first result is [She09b, V.A.1.19]. The second follows from [She09b, V.A.1.12]: one only has to observe that the condition between M and \mathcal{N} there holds when M is χ_0^+ -saturated.

We can now get a (completely local) result on unions of saturated models.

Theorem 4.27. Assume:

- (1) $\lambda > \chi^+$ is such that \mathcal{N} is stable in μ for unboundedly many $\mu < \lambda$.
- (2) $\langle M_i : i < \delta \rangle$ is increasing and for all $i < \delta$, M_i is λ -saturated. Write $M_{\delta} := \bigcup_{i < \delta} M_i$.

(3) For any $q \in S(M_{\delta})$, there exists a splitting-like notion $R, i < \delta$ and $A \subseteq |M_i|$ of size at most χ_0 such that q does not R-split over A.

Then M_{δ} is λ -saturated.

Proof. By Hypothesis 4.4, it is enough to check that M_{δ} is λ -saturated for types of length one. Let $p \in S(B)$, $B \subseteq |M_{\delta}|$ have size less than λ . Let q be an extension of p to $S(M_{\delta})$. If q is algebraic, then p is realized inside M_{δ} so assume without loss of generality that q is not algebraic. By assumption, there exists a splitting-like notion R, $i < \delta$ and $A \subseteq |M_i|$ such that p does not R-split over A and $|A| \leq \chi_0$. Without loss of generality, i = 0. Now M_0 is χ_0^+ -saturated so (by Fact 4.26) there exists $A' \subseteq |M_0|$ of size at most χ_0 such that $q \upharpoonright M_0$ does not split over A'. By making A larger if necessary, we can assume A = A'.

Pick $\mu < \lambda$ such that $\mu \ge \chi^+ + |B|$ and \mathcal{N} is stable in μ . Such a μ exists by the hypothesis on λ . By Lemma 4.17, there exists a sequence I of length μ^+ which is Morley for $q \upharpoonright M_0$ over A, with the witnesses living inside M_0 . Thus I is also Morley for q over A.

By Lemma 4.25, $\operatorname{Av}(\mathbf{I}/M_{\delta}) = q$, and so in particular $\operatorname{Av}(\mathbf{I}/B) = q \upharpoonright B = p$. By Lemma 4.6, p is realized by an element of $\mathbf{I} \subseteq |M_0| \subseteq |M_{\delta}|$, as needed.

The condition (3) in Theorem 4.27 is useful in case we know that the local character cardinal for chains κ_{α} is significantly lower than the local character cardinal for sets $\bar{\kappa}_{\alpha}$. This is the case when a superstability-like condition holds. If we do not care about the local character cardinal for chains, we can state a version of Theorem 4.27 without condition (3).

Corollary 4.28. Assume:

- (1) $\lambda > \chi^+$ is such that $\mu^{\chi_0} < \lambda$ for all $\mu < \lambda$. (2) $\langle M_i : i < \delta \rangle$ is increasing and for all $i < \delta$, M_i is λ -saturated.

If $cf(\delta) \ge \chi_0^+$, then $\bigcup_{i < \delta} M_i$ is λ -saturated.

Proof. Fix $\alpha < \kappa$. By Fact 4.26(1), \mathcal{N} is α -stable in μ for any $\mu < \lambda$ with $\mu^{\chi_0} = \mu$ and $\mu \geq \chi$. By hypothesis, there are unboundedly many such μ 's.

Let $M_{\delta} := \bigcup_{i < \delta} M_i$. By an easy argument using the cofinality condition on δ , M_{δ} is χ_0^+ -saturated. By Fact 4.26(2), for any $p \in S^{<\kappa}(M_{\delta})$, there exists $A \subseteq |M_{\delta}|$ of size $\leq \chi_0$ such that p does not split over A. By the cofinality assumption on δ , we can find $i < \delta$ such that $A \subseteq |M_i|$. Now apply Theorem 4.27 and get the result. \Box

Remark 4.29. The proof shows that we can still replace (1) with " $\lambda > \chi^+$ is such that \mathcal{N} is stable in μ for unboundedly many $\mu < \lambda^{"}$.

We end this section with the following interesting variation: the cardinal arithmetic condition on λ is improved, and we do not even need that the M_i 's be λ -saturated, only that they realize enough types from the previous M_j 's.

Theorem 4.30. Assume:

(1) $\lambda > \chi$ is such that $\mu^{<\kappa} < \lambda$ for all $\mu < \lambda$ (or such that \mathcal{N} is stable in μ for unboundedly many $\mu < \lambda$).

- (2) M is such that for any $q \in S(M)$ there exists $\langle M_i : i < \delta \rangle$ strictly increasing so that:
 - (a) $\delta \geq \lambda$ is a limit ordinal.
 - (b) $M = \bigcup_{i < \delta} M_i$
 - (c) For all $i < \delta$, M_i is χ^+ -saturated and M_{i+1} realizes $q \upharpoonright M_i$.
 - (d) There exists a splitting-like notion R, $i < \delta$ and $A \subseteq |M_i|$ of size at most χ_0 such that q does not R-split over A.

Then M is λ -saturated.

Proof. By Hypothesis 4.4, it is enough to check that M is λ -saturated for types of length one. Let $p \in S(B)$, $B \subseteq |M|$ have size less than λ . Let q be an extension of p to S(M). If q is algebraic, then p is realized inside M, so assume q is not algebraic. Let $\langle M_i : i < \delta \rangle$ be as given by (2) for q. Let R be a splitting-like notion for which there is $i < \delta$ and $A \subseteq |M_i|$ such that q does not R-split over A and $|A| \le \chi_0$. Without loss of generality, i = 0.

Let $\mu := (\chi + |B|)^{<\kappa}$ (or take $\mu < \lambda$ such that $\mu \ge \chi + |B|$ and \mathcal{N} is stable in μ). Note that $\mu < \lambda$. For $i < \mu^+$, let $a_i \in |M_{i+1}|$ realize $q \upharpoonright M_i$. By cofinality considerations, $\bigcup_{i < \mu^+} M_i$ is χ^+ -saturated. By Fact 4.26, there exists $i < \mu^+$ and $A' \subseteq |M_i|$ such that $q \upharpoonright \bigcup_{i < \mu^+} M_i$ does not split over A'. By some renaming we can assume without loss of generality that A' = A. It is now easy to check that $\mathbf{I} := \langle a_i : i < \mu^+ \rangle$ is Morley for q over A, as witnessed by $\langle M_i : i < \mu^+ \rangle$.

By Lemma 4.25, $\operatorname{Av}(\mathbf{I}/M) = q$, and so in particular $\operatorname{Av}(\mathbf{I}/B) = q \upharpoonright B = p$. By Lemma 4.6, p is realized by an element of $\mathbf{I} \subseteq |M_0| \subseteq |M|$, as needed. \Box

5. TRANSLATING TO AECS

To translate the result of the previous section to AECs, we will use the *Galois Morleyization* of an AEC, a tool introduced in [Vas16c]: Essentially, we expand the language of the AEC with a symbol for each Galois type. With enough tameness, Galois types then become syntactic.

Definition 5.1 (3.3 in [Vas16c]). Let K be an AEC and let κ be an infinite cardinal. Define an (infinitary) expansion \hat{L} of L(K) by adding a relation symbol R_p of arity $\ell(p)$ for each $p \in \mathrm{gS}^{<\kappa}(\emptyset)$. Expand each $N \in K$ to a \hat{L} -structure \hat{N} by specifying that for each $\bar{a} \in \hat{N}$, $R_p^{\hat{N}}(\bar{a})$ holds exactly when $\mathrm{gtp}(\bar{a}/\emptyset; N) = p$. We write $\hat{K}^{<\kappa}$ for \hat{K} . We call $\hat{K}^{<\kappa}$ the $(<\kappa)$ -Galois Morleyization of K.

Remark 5.2. Let K be an AEC and κ be an infinite cardinal. Then $|L(\widehat{K}^{<\kappa})| \le |gS^{<\kappa}(\emptyset)| + |L| \le 2^{<(\kappa + LS(K)^+)}$.

Fact 5.3 (3.16 in [Vas16c]). Let K be a $(<\kappa)$ -tame AEC, and let $M \leq N_{\ell}, a_{\ell} \in |N_{\ell}|, \ell = 1, 2$. Then $\operatorname{gtp}(a_1/M; N_1) = \operatorname{gtp}(a_2/M; N_2)$ if and only if $\operatorname{fp}_{q\widehat{L}_{\kappa,\kappa}}(a_1/M; \widehat{N}_1) = \operatorname{tp}_{q\widehat{L}_{\kappa,\kappa}}(a_2/M; \widehat{N}_2)$.

20

⁶Recall that $\operatorname{tp}_{q\widehat{L}_{\kappa,\kappa}}$ stands for quantifier-free $L_{\kappa,\kappa}$ -type.

Moreover the left to right direction does not need tameness: if $M \leq N_{\ell}$, $\bar{a}_{\ell} \in {}^{<\infty}|N_{\ell}|, \ \ell = 1, 2$, and $\operatorname{gtp}(\bar{a}_1/M; N_1) = \operatorname{gtp}(\bar{a}_2/M; N_2)$, then $\operatorname{tp}_{q\widehat{L}_{\kappa,\kappa}}(\bar{a}_1/M; \widehat{N}_1) = \operatorname{tp}_{q\widehat{L}_{\kappa,\kappa}}(\bar{a}_2/M; \widehat{N}_2)$.

Note that this implies in particular that (if K is $(< \kappa)$ -tame and has amalgamation) the Galois version of saturation and stability coincide with their syntactic analog in $\hat{K}^{<\kappa}$. There is also a nice correspondence between the syntactic version of the order property defined at the beginning of Section 4 and Shelah's semantic version [She99, 4.3]:

Definition 5.4. Let α and μ be cardinals and let K be an AEC. A model $M \in K$ has the α -order property of length μ if there exists $\langle \bar{a}_i : i < \mu \rangle$ inside M with $\ell(\bar{a}_i) = \alpha$ for all $i < \mu$, such that for any $i_0 < j_0 < \mu$ and $i_1 < j_1 < \mu$, $gtp(\bar{a}_{i_0}\bar{a}_{j_0}/\emptyset; N) \neq gtp(\bar{a}_{j_1}\bar{a}_{i_1}/\emptyset; N)$.

M has the $(<\alpha)$ -order property of length μ if it has the β -order property of length μ for some $\beta < \alpha$. *M* has the order property of length μ if it has the α -order property of length μ for some α .

K has the α -order of length μ if some $M \in K$ has it. K has the order property if it has the order property for every length.

Fact 5.5 (4.4 in [Vas16c]). Let K be an AEC. Let $\widehat{K} := \widehat{K}^{<\kappa}$. If $\widehat{N} \in \widehat{K}$ has the (syntactic) order property of length χ , then N has the (Galois) ($< \kappa$)-order property of length χ . Conversely, if $\chi \geq 2^{<(\kappa + \mathrm{LS}(K)^+)}$ and N has the (Galois) ($< \kappa$)-order property of length $(2^{\chi})^+$, then \widehat{N} has the (syntactic) order property of length χ .

We will use Facts 5.3 and 5.5 freely in this section. We will also use the following results about stability and the order property:

Fact 5.6 (4.5 and 4.13 in [Vas16c]). Let K be an $(< \kappa)$ -tame AEC with amalgamation. The following are equivalent:

- (1) K is stable in some $\lambda \ge \kappa + \mathrm{LS}(K)$.
- (2) There exists $\mu \leq \lambda_0 < h^*(\kappa + \mathrm{LS}(K)^+)$ (see Notation 2.1) such that K is stable in any $\lambda \geq \lambda_0$ with $\lambda = \lambda^{<\mu}$.
- (3) K does not have the order property.
- (4) There exists $\chi < h^*(\kappa + \mathrm{LS}(K)^+)$ such that K does not have the $(< \kappa)$ -order property of length χ .

It remains to find an independence notion to satisfy condition (3) in Theorem 4.27. The splitting-like notion R there will be given by the following:

Definition 5.7. Let K be an AEC and let κ be an infinite cardinal. For $p \in$ $gS^{<\infty}(B;N)$ and $A \subseteq B$, say p κ -explicitly does not split over A if whenever p = $gtp(\bar{c}/B;N)$, for any $\bar{b}, \bar{b}' \in {}^{<\kappa}B$, if $gtp(\bar{b}/A;N) = gtp(\bar{b}'/A;N)$, then $tp_{q\widehat{L}_{\kappa,\kappa}}(\bar{c}\bar{b}/A;N) =$ $tp_{q\widehat{L}_{\kappa,\kappa}}(\bar{c}\bar{b}'/A;N)$, where $\widehat{L} = L(\widehat{K}^{<\kappa})$.

Remark 5.8. This is closely related to explicit nonsplitting defined in [BGKV16, 3.13]. The definition there is that p explicitly does not split if and only if it κ -explicitly does not split for all κ . When K is fully ($< \kappa$)-tame and short (see [Bon14b, 3.3]), this is equivalent to just asking for p to κ -explicitly not split.

Remark 5.9 (Syntactic invariance). Let $\widehat{K} := \widehat{K}^{<\kappa}$. Assume $\operatorname{tp}_{\widehat{qL}_{\kappa,\kappa}}(\overline{c}/B; N) =$ $\operatorname{tp}_{\widehat{aler}}(\overline{c}'/B;N)$ and $\operatorname{gtp}(\overline{c}/B;N)$ κ -explicitly does not split over $A \subseteq B$. Then $gtp(\bar{c}'/B; N)$ κ -explicitly does not split over A.

We will use the following definition of an independence relation, which appears implicitly in [Vas16b, 4.8].

Definition 5.10. Let K be an AEC with amalgamation and let $\lambda \geq LS(K)$ be such that K is λ -tame and stable in λ . For $M \leq N$ with $M \lambda^+$ -saturated, we say that $p \in gS(N)$ does not λ -fork over M if there exists $M_0 \in K_\lambda$ such that $M_0 \leq M$ and p does not λ -split over M_0 (that is [She99, I.3.2], whenever N_{ℓ} , $\ell = 1, 2$, are of size λ such that $M_0 \leq N_\ell \leq N$ and $f: N_1 \cong_{M_0} N_2$, we have that $f(p \upharpoonright N_1) = p \upharpoonright N_2$.

We write $a \stackrel{\lambda-nf}{\underset{M}{\downarrow}} N$ to say that gtp(a/N) does not λ -fork over M and will apply the definition of the properties Definition 2.4 and Fact 2.6 to it.

Fact 5.11 (§4 and §5 in [Vas16b]). Let K be an AEC with amalgamation and let $\lambda \geq LS(K)$ be such that K is λ -tame, stable in λ , and has no maximal models in λ.

Then λ -nonforking satisfies invariance, monotonicity, transitivity (i.e. if $M_1 \leq M_2 \leq$ M_3 are such that M_1 and M_2 are λ^+ -saturated, $p \in gS(M_3)$, p does not λ -fork over M_2 , $p \upharpoonright M_2$ does not λ -fork over M_1 , then p does not λ -fork over M_1), and uniqueness. Moreover $\bar{\kappa}_1(\downarrow^{\lambda-\mathrm{nf}} \upharpoonright K^{\lambda^+-\mathrm{sat}}) = \lambda^{++}$.

We recall the definition of superstability from [Vas16a, 10.1] using local character of nonsplitting. Note that it coincides with the first-order definition (see [Vas16a, 10.9) and is equivalent to the definition implicit in [GVV16, Vas16b] and explicit in [Gro02, 7.12].

Definition 5.12 (Superstability). Let K be an AEC.

- (1) For $M, N \in K$, say N is universal over M if and only if $M \leq N$ and whenever we have $M' \ge M$ such that ||M'|| = ||M||, then there exists $f:M'\xrightarrow[M]{} N.$
- (2) K is λ -superstable if:
 - (a) $LS(K) \leq \lambda$ and $K_{\lambda} \neq \emptyset$.
 - (b) K_{λ} has amalgamation, joint embedding, and no maximal models.
 - (c) K is stable in λ .
 - (d) K has no long splitting chains in λ : for any limit $\delta < \lambda^+$ and increasing continuous $\langle M_i : i \leq \delta \rangle$ in K_{λ} with M_{i+1} universal over M_i for all $i < \delta$, and any $p \in gS(M_{\delta})$, there exists $i < \delta$ such that p does not λ -split over M_i .

Note that superstability implies local character of λ -forking, and superstability transfers up assuming tameness:

Fact 5.13. Let K be an AEC with amalgamation that is λ -tame and λ -superstable.

- (1) [Vas16b, 4.11] $\kappa_1(\stackrel{\lambda-\mathrm{nf}}{\downarrow} \upharpoonright K^{\lambda^+\mathrm{-sat}}) = \aleph_0.$ (2) K is λ' -superstable for all $\lambda' \ge \lambda$.

The next result imitates [BGKV16, 5.6]:

Lemma 5.14. Let K be an AEC with amalgamation and let $\lambda \geq LS(K)$ be such that K is λ -tame, stable in λ , and has no maximal models in λ . Let $\kappa \leq \lambda^+$.

Let $M \leq N$ be given with $M \lambda^+$ -saturated. Let $p \in gS(N)$. If p does not λ -fork over M, then $p \kappa$ -explicitly does not split over A.

Proof. By definition of λ -nonforking, there exists $M_0 \leq M$ of size λ such that pdoes not λ -split over M_0 . We will show that p explicitly does not κ -split over M_0 which is enough by base monotonicity of explicit κ -nonsplitting.

Work inside a monster model \mathfrak{C} and write $p = \operatorname{gtp}(c/N)$. Let $\overline{b}, \overline{b'} \in {}^{<\kappa}|N|$ be such that $gtp(b/M_0) = gtp(b'/M_0)$. Let f be an automorphism of \mathfrak{C} fixing M_0 such that $f(\bar{b}) = \bar{b}'$. By invariance, f(p) does not λ -split over M_0 . Now using uniqueness of λ -splitting (see [Van06, I.4.12]), $f(p \upharpoonright M_0 \overline{b}) = p \upharpoonright M_0 \overline{b}'$. The result follows.

The next technical lemma captures the essence of our translation:

Lemma 5.15. Let K be a $(< \kappa)$ -tame AEC with amalgamation and no maximal models. Let χ_0 be such that:

- (1) $\chi_0 \ge 2^{<(\kappa + \mathrm{LS}(K)^+)}$. (2) K does not have the $(< \kappa)$ -order property of length χ_0^+ .

Set $\chi := (2^{2^{\chi_0}})^+$. Let λ be such that:

- (1) $\lambda > \chi^+$.
- (2) K is stable in μ for unboundedly many $\mu < \lambda$.

Let $\theta := \kappa_1(\bigcup^{\chi\text{-nf}} \upharpoonright K^{\chi^+\text{-sat}})$. Then:

- (1) If $\langle M_i : i < \delta \rangle$ is an increasing chain of λ -saturated models and $cf(\delta) \ge \theta$, then $\bigcup_{i < \delta} M_i$ is λ -saturated.
- (2) If $M \in K$ is such that for any $q \in gS(M)$ there exists $\langle M_i : i < \delta \rangle$ strictly increasing so that:
 - (a) $\delta \geq \lambda$ and $cf(\delta) \geq \theta$.
 - (b) $M = \bigcup_{i < \delta} M_i$.
 - (c) For all $i < \delta$, M_i is χ^+ -saturated and M_{i+1} realizes $q \upharpoonright M_i$. Then M is λ -saturated.

Proof. We prove the first statement. The proof of the second is analogous but uses Theorem 4.30 instead of Theorem 4.27. Set $M_{\delta} := \bigcup_{i < \delta} M_i$. Let $N \geq M_{\delta}$ be such that N realizes all types in $gS^{<\kappa}(M_{\delta})$. We check that M_{δ} is λ -saturated in N. Let $\widehat{K} := \widehat{K}^{<\kappa}$ be the $(<\kappa)$ -Galois Morleyization of K. Let $\mathcal{N} := \widehat{N}$. By $(<\kappa)$ -tameness, it is enough to show that \widehat{M}_{δ} is (syntactically) λ -saturated in \mathcal{N} . Work inside \mathcal{N} in the language of \widehat{K} . We also let $\mathcal{S} := \{|M| \mid M \leq N\}$. Note that \mathcal{S} satisfies Hypothesis 4.2.

First observe that Hypothesis 4.22 holds as (Remark 5.2) $|L(\hat{K})| \leq 2^{\langle (\kappa + LS(K)^+)}$, so χ_0 has all the required properties. Also, Hypothesis 4.4 holds by [She09a, II.1.14].

Note that K is stable in χ by Fact 4.26(1). By hypothesis, $\lambda > \chi^+$. We want to use Theorem 4.27, and it remains to check that (3) there holds.

For $A \subseteq B$ and $p \in S^{<\infty}(B)$, define the relation R(p,A) to hold if and only if $p = \operatorname{tp}(\bar{c}/B)$ and $\operatorname{gtp}(\bar{c}/B; N)$ κ -explicitly does not split over A. Note that this is well-defined by Remark 5.9. We want to check that this is a splitting-like notion (Definition 4.23). By definition of κ -explicit nonsplitting, if $p \in S^{<\infty}(B)$ does not split over $A \subseteq B$, then R(p, A). Also, it is easy to check that R satisfies the monotonicity axiom. It remains to check the weak uniqueness axiom. So let $M \text{ be } \mu := \left(|A| + (|L(\widehat{K})| + 2)^{<\kappa} \right)^+ \text{-saturated}, A \subseteq |M| \subseteq B, \text{ and for } \ell = 1, 2,$ $q_{\ell} \in S^{<\infty}(B)$, $R(q_{\ell}, A)$ and $q_1 \upharpoonright M = q_2 \upharpoonright M$. Note that M is also μ -saturated in the Galois sense (by tameness and Remark 3). Thus we can imitate the proof of Lemma 4.8, using Galois saturation instead of syntactic saturation to get \bar{b}' satisfying $gtp(\overline{b}'/A) = gtp(\overline{b}/A)$ (instead of just $tp(\overline{b}'/A) = tp(\overline{b}'/A)$ as there). The definition of κ -explicit nonsplitting then makes the proof go through.

Now let $q \in gS(M_{\delta})$. By definition of θ , there exists $i < \delta$ such that q does not χ -fork over M_i . Now by set local character, there exists $M \leq M_i$ of size χ^+ such that $q \upharpoonright M_i$ does not χ -fork over M. By transitivity, q does not χ -fork over M. By Lemma 5.14, working syntactically inside \mathcal{N} , q does not R-split over M. Thus (3) holds. Therefore M_{δ} is λ -saturated, as desired. \square

We obtain the following result on chains of saturated models in stable AECs:

Theorem 5.16. Let K be a $(< \kappa)$ -tame AEC with amalgamation, $\kappa > LS(K)^+$. If K is stable, then there exists $\chi_0 \leq \lambda_0 < h^*(\kappa)$ (see Notation 2.1) satisfying the following property:

If $\lambda \geq \lambda_0$ is such that $\mu^{\chi_0} < \lambda$ for all $\mu < \lambda$ (or just that K is stable in μ for unboundedly many $\mu < \lambda$, then whenever $\langle M_i : i < \delta \rangle$ is an increasing chain of λ -saturated models with $cf(\delta) \geq \lambda_0$, we have that $\bigcup_{i < \delta} M_i$ is λ -saturated.

Proof. Using Fact 5.6, pick $\chi_0 \leq \mu_0 < h^*(\kappa)$ such that:

- (1) $\chi_0^+ \ge 2^{<(\kappa + \operatorname{LS}(K)^+)} + \kappa^+$. (2) K is stable in any $\mu \ge \mu_0$ with $\mu = \mu^{\chi_0}$. (3) K does not have the $(<\kappa)$ -order property of length χ_0^+ .

Now set $\lambda_0 := (2^{2^{\chi_0}})^{+3}$ and apply Lemma 5.15.

The statement becomes much nicer in superstable AECs:

Theorem 5.17. Let K be a $(< \kappa)$ -tame AEC with amalgamation, $LS(K)^+ \leq \kappa$. Let $\mu \geq LS(K)$ be a cardinal with $\mu^+ \geq \kappa$ and assume that K is μ -superstable. Then there exists $\lambda_0 < h^*(\kappa) + \mu^{++}$ with $\lambda_0 > \mu$ such that for any $\lambda \ge \lambda_0$:

- (1) $K^{\lambda-\text{sat}}$ is an AEC with $LS(K^{\lambda-\text{sat}}) = \lambda$.
- (2) If $M \in K^{\lambda_0\text{-sat}}$ is such that for any $q \in gS(M)$ there exists $\langle M_i : i < \lambda \rangle$ a resolution of M in K^{λ_0 -sat such that $q \upharpoonright M_i$ is realized in M_{i+1} for all $i < \lambda$, then $M \in K^{\lambda-\text{sat}}$.

Proof.

- (1) We first show that any increasing union of λ -saturated models is saturated. Let $\lambda_{00} < h^*(\kappa)$ be as given by the proof of Theorem 5.16 and let $\lambda_0 := \lambda_{00} + \mu^+$. By Fact 5.13, $\kappa_1(\downarrow \ \mid K^{\lambda_0-\text{sat}}) = \aleph_0$. Now apply Lemma 5.15 (note that by Fact 5.13, K is stable in any $\mu' \ge \mu$). To see that $\text{LS}(K^{\lambda-\text{sat}}) = \lambda$, imitate the proof of [She90, Theorem III.3.12].
- (2) Similar: use the second conclusion of Lemma 5.15.

6. ON SUPERSTABILITY IN AECS

In the introduction to [She09a], Shelah points out the importance of finding a definition of superstability for AECs. He also remarks (p. 19) that superstability in AECs suffers from "schizophrenia": definitions that are equivalent in the first-order case might not be equivalent in AECs. In this section, we point out that Definition 5.12 implies several other candidate definitions of superstability. Recall from Fact 5.13 that Definition 5.12 implies that the class is stable on a tail of cardinals. We will focus on five other definitions:

- (1) For every high-enough λ , the union of any increasing chain of λ -saturated models is λ -saturated. This is the focus of this paper and is equivalent to first-order superstability by [AG90, Theorem 13].
- (2) The existence of a saturated model of size λ for every high-enough λ . In first-order, this is an equivalent definition of superstability by the saturation spectrum theorem (Fact 1.2).
- (3) The existence of a superlimit model of size λ for every high-enough λ . This is the definition of superstability listed by Shelah in [She09a, N.2.4]. Recall that a model $M \in K_{\lambda}$ is *superlimit* if it is universal, has an isomorphic proper extension in K_{λ} , and whenever $\langle M_i : i < \delta \rangle$ is increasing in K_{λ} , $\delta < \lambda^+$, and $M_i \cong M$ for all $i < \delta$, then $\bigcup_{i < \delta} M_i \cong M$.
- (4) The existence of a good λ-frame on a subclass of saturated models (e.g. for every high-enough λ). Recall that a good frame is essentially a forking-like notion for types of length one (see [She09a, II.2.1] for the formal definition). Good frames are *the* central notion in [She09a] and are described by Shelah as a "bare bone" definition of superstability.
- (5) The uniqueness of limit models of size λ for every high-enough λ : Recall that a model M is (λ, δ) -limit over M_0 if $M_0 \leq M$ are in K_{λ} , $\delta < \lambda^+$ is a limit ordinal and there exists $\langle M_i : i \leq \delta \rangle$ increasing continuous such that $M_{\delta} = M$ and $i < \delta$ implies $M_i <_{\text{univ}} M_{i+1}$ (recall Definition 5.12). We say K_{λ} has uniqueness of limit models if for any $M_0 \in K_{\lambda}$, any limit $\delta_1, \delta_2 < \lambda^+$, any M_{ℓ} which are (λ, δ_{ℓ}) -limit over M_0 are isomorphic over M_0 . Uniqueness of limit models is central in [She99, SV99, Van06, Van13] and is further examined in [GVV16] (Theorem 6.1 there proves that the condition is equivalent to first-order superstability). These papers all prove the uniqueness under a categoricity (or no Vaughtian pair) assumption. In [She09a, II.4.8], uniqueness of limit models is proven from a good frame (see also [Bon14a, 9.2] for a detailed writeup). This is used in [BG] to get

eventual uniqueness of limit models from categoricity, but the authors have to make an extra assumption (the extension property for coheir).

Note that some easy implications between these definitions are already known (see for example [Dru13, 2.3.12]). We now show that assuming amalgamation and tameness, if K is superstable, then all five of these conditions hold. This gives an eventual version of [Dru13, Conjecture 4.2.5]. It also shows how to build a good frame without relying on categoricity (as opposed to all previous constructions, see [She09a, II.3.7], [Vas16b, 7.3], or [Vas16a, 10.16]).

Theorem 6.1. If K is a μ -tame, μ -superstable AEC with amalgamation, then there exists $\lambda_0 < h(\mu)$ such that for all $\lambda \ge \lambda_0$:

- (1) The union of any increasing chain of λ -saturated models in K is λ -saturated.
- (2) K has a saturated model of size λ .
- (3) K has a superlimit model of size λ .
- (4) There exists a type-full good λ -frame with underlying class $K_{\lambda}^{\lambda-\text{sat}}$.
- (5) K_{λ} has uniqueness of limit models.

Proof. Note that by Fact 5.13, $K_{\geq\mu}$ has no maximal models, joint embedding, and is stable in every cardinal. Let $\lambda_0 < h(\mu)$ be as given by Theorem 5.17 and let $\lambda \geq \lambda_0$. Then $K^{\lambda\text{-sat}}$ is an AEC with $\mathrm{LS}(K^{\lambda\text{-sat}}) = \lambda$. Thus (1) and (2) hold. If M is the saturated model of size λ , then it is easy to check that M is superlimit: it is universal as $K_{\geq\mu}$ has joint embedding, it has a saturated proper extension of size λ since $\mathrm{LS}(K^{\lambda\text{-sat}}) = \lambda$, and any increasing chain of saturated models in K_{λ} of length less than λ^+ has a saturated union. Thus (3) holds. To see (4), use [Vas16a, 10.8(2c)].

We are now ready to prove (5). As observed above, a good frame implies uniqueness of limit models. Thus $K_{\lambda}^{\lambda-\text{sat}}$ has uniqueness of limit models. It follows that K_{λ} has uniqueness of limit models: Let M_{ℓ} be (λ, δ_{ℓ}) -limit over M_0 , $\ell = 1, 2$. Pick $M'_0 \geq M_0$ in $K_{\lambda}^{\lambda-\text{sat}}$. By universality, M_{ℓ} is also (λ, δ_{ℓ}) -limit over some copy of M'_0 , so after some renaming we can assume without loss of generality that $M_0 = M'_0$. For $\ell = 1, 2$, build $\langle M'_i : i \leq \delta_{\ell} \rangle$ increasing continuous such that for all $i < \delta_{\ell}$, $M'_i \in K_{\lambda}^{\lambda-\text{sat}}$ and $M'_i <_{\text{univ}} M'_{i+1}$. This is easy to do and by a back and forth argument, $M_{\ell} \cong_{M_0} M'_{\delta_{\ell}}$. By uniqueness of limit models in $K^{\lambda-\text{sat}}$, $M'_{\delta_1} \cong_{M_0} M'_{\delta_2}$.

References

- [AG90] Michael H. Albert and Rami Grossberg, *Rich models*, The Journal of Symbolic Logic 55 (1990), no. 3, 1292–1298.
- [Bal09] John T. Baldwin, Categoricity, University Lecture Series, vol. 50, American Mathematical Society, 2009.
- [BG] Will Boney and Rami Grossberg, Forking in short and tame AECs, Annals of Pure and Applied Logic, To appear. URL: http://arxiv.org/abs/1306.6562v11.
- [BGKV16] Will Boney, Rami Grossberg, Alexei Kolesnikov, and Sebastien Vasey, Canonical forking in AECs, Annals of Pure and Applied Logic 167 (2016), no. 7, 590–613.
- [Bon14a] Will Boney, Tameness and extending frames, Journal of Mathematical Logic 14 (2014), no. 2, 1450007.
- [Bon14b] _____, Tameness from large cardinal axioms, The Journal of Symbolic Logic 79 (2014), no. 4, 1092–1119.

- [Dru13] Fred Drueck, Limit models, superlimit models, and two cardinal problems in abstract elementary classes, Ph.D. thesis, 2013, Available online. URL: http://homepages.math.uic.edu/~drueck/thesis.pdf.
- [GL02] Rami Grossberg and Olivier Lessmann, Shelah's stability spectrum and homogeneity spectrum in finite diagrams, Archive for Mathematical Logic 41 (2002), no. 1, 1-31. [Gro] Rami Grossberg, A course in model theory I, A book in preparation.
- [Gro91] _, On chains of relatively saturated submodels of a model without the order property, The Journal of Symbolic Logic 56 (1991), 124-128.
- [Gro02] Classification theory for abstract elementary classes, Contemporary Mathematics 302 (2002), 165-204.
- [GV06] Rami Grossberg and Monica VanDieren, Galois-stability for tame abstract elementary classes, Journal of Mathematical Logic 6 (2006), no. 1, 25-49.
- [GVV16] Rami Grossberg, Monica VanDieren, and Andrés Villaveces, Uniqueness of limit models in classes with amalgamation, Mathematical Logic Quarterly 62 (2016), 367-382.
- [Har75] Victor Harnik, On the existence of saturated models of stable theories, Proceedings of the American Mathematical Society 52 (1975), 361–367.
- Michael Makkai and Saharon Shelah, Categoricity of theories in $L_{\kappa,\omega}$, with κ a com-[MS90] pact cardinal, Annals of Pure and Applied Logic 47 (1990), 41-97.
- [She75] Saharon Shelah, The lazy model theoretician's guide to stability, Logique et Analyse **18** (1975), 241–308.
- , Classification of non elementary classes II. Abstract elementary classes, Clas-[She87] sification Theory (Chicago, IL, 1985) (John T. Baldwin, ed.), Lecture Notes in Mathematics, vol. 1292, Springer-Verlag, 1987, pp. 419-497.
- [She90]_, Classification theory and the number of non-isomorphic models, 2nd ed., Studies in logic and the foundations of mathematics, vol. 92, North-Holland, 1990.
- [She99] , Categoricity for abstract classes with amalgamation, Annals of Pure and Applied Logic 98 (1999), no. 1, 261-294.
- [She09a] , Classification theory for abstract elementary classes, Studies in Logic: Mathematical logic and foundations, vol. 18, College Publications, 2009.
- _, Classification theory for abstract elementary classes 2, Studies in Logic: [She09b] Mathematical logic and foundations, vol. 20, College Publications, 2009.
- [SV99] Saharon Shelah and Andrés Villaveces, Toward categoricity for classes with no maximal models, Annals of Pure and Applied Logic **97** (1999), 1–25.
- [Van06] Monica VanDieren, Categoricity in abstract elementary classes with no maximal models, Annals of Pure and Applied Logic 141 (2006), 108-147.
- [Van13] _, Erratum to "Categoricity in abstract elementary classes with no maximal models" [Ann. Pure Appl. Logic 141 (2006) 108-147], Annals of Pure and Applied Logic 164 (2013), no. 2, 131–133.
- [Vas16a] Sebastien Vasey, Building independence relations in abstract elementary classes, Annals of Pure and Applied Logic 167 (2016), no. 11, 1029-1092.
- [Vas16b] ., Forking and superstability in tame AECs, The Journal of Symbolic Logic 81 (2016), no. 1, 357-383.
- [Vas16c] ., Infinitary stability theory, Archive for Mathematical Logic 55 (2016), 567-592.

E-mail address: wboney@math.harvard.edu

URL: http://math.harvard.edu/~wboney/

DEPARTMENT OF MATHEMATICS, HARVARD UNIVERSITY, CAMBRIDGE, MASSACHUSETTS, USA

E-mail address: sebv@cmu.edu

URL: http://math.cmu.edu/~svasey/

DEPARTMENT OF MATHEMATICAL SCIENCES, CARNEGIE MELLON UNIVERSITY, PITTSBURGH, PENN-SYLVANIA, USA