n-Dimensional Optical Orthogonal Codes, Bounds and **Optimal Constructions**

T. L. Alderson *1

¹University of New Brunswick, Canada

August 12, 2018

Abstract

We generalized to higher dimensions the notions of optical orthogonal codes. We establish uper bounds on the capacity of general n-dimensional OOCs, and on specific types of ideal codes (codes with zero off-peak autocorrelation). The bounds are based on the Johnson bound, and subsume many of the bounds that are typically applied to codes of dimension three or less. We also present two new constructions of ideal codes; one furnishes an infinite family of optimal codes for each dimension $n \ge 2$, and another which provides an asymptotically optimal family for each dimension $n \ge 2$. The constructions presented are based on certain point-sets in finite projective spaces of dimension k over GF(q) denoted PG(k,q).

Introduction 1

A (1-dimensional) $(n, w, \lambda_a, \lambda_c)$ optical orthogonal code (OOC) is a family of binary sequences (codewords) of length n, and constant Hamming weight w satisfying the following two conditions:

- (off-peak auto-correlation property) for any codeword $c = (c_0, c_1, \ldots, c_{n-1})$ and for any integer $1 \le t \le n-1$, we have $\sum_{i=1}^{n-1} c_i c_{i+t} \le \lambda_a$,
- (cross-correlation property) for any two distinct codewords c, c' and for any integer $0 \le t \le$ n-1

$$n-1$$
, we have $\sum_{i=0} c_i c'_{i+t} \leq \lambda_c$,

where each subscript is reduced modulo n.

An $(N, w, \lambda_a, \lambda_c)$ OOC with $\lambda_a = \lambda_c$ is denoted an (N, w, λ) OOC. The number of codewords is the size of the code. For fixed values of N, w, λ_a and λ_c , the largest size of an $(N, w, \lambda_a, \lambda_c)$ -OOC is denoted $\Phi(N, w, \lambda_a, \lambda_c)$. An $(N, w, \lambda_a, \lambda_c)$ -OOC of size $\Phi(N, w, \lambda_a, \lambda_c)$ is said to be optimal.

A family of $(N, w, \lambda_a, \lambda_c)$ OOCs is called asymptotically optimal if

$$\lim_{N \to \infty} \frac{|C|}{\Phi(C)} = 1.$$
(1)

^{*}The author acknowledges the support of the NSERC of Canada Discovery Grant program.

Since the work of Salehi *et. al.* [24] [23], OOCs have been employed within optical code division multiple access (OCDMA) networks. OCDMA networks are widely employed due to their strong performance with multiple users. They are ideally suited for bursty, asynchronous, concurrent traffic. In applications, optimal OOCs facilitate the largest possible number of asynchronous users to transmit information efficiently and reliably. In order to maintain low correlation values the code length must increase quite rapidly with the number of users, reducing bandwidth utilization.

The 1-D OOCs spread the input data bits only in the time domain. Technologies such as wavelength-division-multiplexing (WDM) and dense-WDM enable the spreading of codewords in both space and time [21], or in wave-length and time [15]. Hence, codewords may be considered as $\Lambda \times T$ (0,1)-matrices. These codes are referred to in the literature as multiwavelength, multiple-wavelength, wavelength-time hopping, and 2-dimensional OOCs (2-D OOCs).

This addition of another dimension allows codes with off-peak autocorrelation zero and thereby improving the OCDMA performance in comparison with 1-D OCDMA. For optimal constructions of 2-D OOC's see [7, 13, 17], and for asymptotically optimal constructions see [18, 19, 26, 27, 28]. Later, a third dimension was added which gave an increase the code size and the performance of the code [11, 2]. In 3-D OCDMA the optical pulses are spread in three domains space, wave-length, and time, with codes referred to as *space/wavelength/time spreading* codes, or 3-D OOC. In [8], coherent fibre-optic communication systems are discussed, whereby both quadratures and both polarizations of the electromagnetic field are used, resulting in a four-dimensional signal space.

In the present work we carry these developments to the next natural stage, introducing constructions and bounds on *n*-dimensional OOCs, for all $n \ge 1$. In section 1.1 we introduce *n*-dimensional OOCs. We develop some upper bounds on these codes based on the Johnson Bound. We also develop bounds on higher dimensional ideal codes ($\lambda_a = 0$). In Section 3 we present two new constructions of ideal codes; one infinite family of optimal codes, and another which is asymptotically optimal.

1.1 *n*-D OOCs and Bounds

Denote by $(\Lambda_1 \times \Lambda_2 \cdots \times \Lambda_{n-1} \times T, w, \lambda_a, \lambda_c)$ an *n* dimensional Optical Orthogonal Code (*n*-D OOC) with constant weight *w*, *i*'th spreading length Λ_i , $1 \leq i \leq n-1$, and time-spreading length *T*. Each codeword may be considered as an *n*-dimensional $\Lambda_1 \times \Lambda_2 \cdots \times \Lambda_{n-1} \times T$ binary array. The off-peak autocorrelation, and cross correlation of an $(\Lambda_1 \times \Lambda_2 \cdots \times \Lambda_{n-1} \times T, w, \lambda_a, \lambda_c)$ *n*-D OOC have the following properties.

• (off-peak auto-correlation property) for any codeword $A = (a_{i_1,i_2,...,i_n})$ and for any integer $1 \le t \le T - 1$, we have

$$\sum_{i_1=0}^{\Lambda_1-1} \sum_{i_2=0}^{\Lambda_2-1} \cdots \sum_{i_{n-1}=0}^{\Lambda_{n-1}-1} \sum_{i_n=1}^{T-1} a_{i_1,i_2,\dots,i_n} a_{i_1,i_2,\dots,i_n+t} \le \lambda_a,$$

• (cross-correlation property) for any two distinct codewords $A = (a_{i_1,i_2,...,i_n}), B = (b_{i_1,i_2,...,i_n})$ and for any integer $0 \le t \le T - 1$, we have

$$\sum_{i_1=0}^{\Lambda_1-1} \sum_{i_2=0}^{\Lambda_2-1} \cdots \sum_{i_{n-1}=0}^{\Lambda_{n-1}-1} \sum_{i_n=1}^{T-1} a_{i_1,i_2,\dots,i_n} b_{i_1,i_2,\dots,i_n+t} \le \lambda_c,$$

where each subscript is reduced modulo T. In the case that $\lambda_a = \lambda_c$, C is denoted an $(\Lambda_1 \times \Lambda_2 \cdots \times \Lambda_{n-1} \times T, w, \lambda)$ OOC.

We note that taking all but t - 1 of the Λ_i 's to be 1 results in a t-dimensional OOC. As with other OOCs we shall take minimal correlation values to be most desirable. Codes satisfying $\lambda_a = 0$ will be said to be *ideal*.

As it is of interest to construct codes with as large cardinality as possible, we now discuss some upper bounds on the size of codes. We shall require the following notation. By an $(N, w, \lambda)_{m+1}$ code, we denote a code of length N, with constant weight w, and maximum Hamming correlation (the number of non-zero agreements between the two codewords) of λ over an alphabet (containing zero) of size m + 1. For binary codes (m = 1) the subscript 2 is typically dropped. Let $A(N, w, \lambda)_{m+1}$ denote the maximum size of an $(N, w, \lambda)_{m+1}$ -code. In [2], the following bound is established.

Theorem 1.1 ([2], Johnson Bound Non-binary).

$$A(N, w, \lambda)_{m+1} \leq \left\lfloor \frac{mN}{w} \left\lfloor \frac{m(N-1)}{w-1} \left\lfloor \cdots \left\lfloor \frac{m(N-\lambda)}{w-\lambda} \right\rfloor \right\rfloor \cdots \right\rfloor.$$

If $w^2 > mN\lambda$ then

$$A(N, w, \lambda)_{m+1} \leq \min \left\{ mN, \left\lfloor \frac{mN(w-\lambda)}{w^2 - mN\lambda} \right\rfloor \right\}.$$

From the Johnson Bound for constant weight codes it follows [9] that

$$\Phi(N, w, \lambda) \le J(N, w, \lambda) = \left\lfloor \frac{1}{w} \left\lfloor \frac{N-1}{w-1} \left\lfloor \frac{N-2}{w-2} \left\lfloor \cdots \left\lfloor \frac{N-\lambda}{w-\lambda} \right\rfloor \right\rfloor \cdots \right\rfloor$$
(2)

$$:= \lfloor f(N, w, \lambda) \rfloor . \tag{3}$$

We note that the first bound in Theorem 1.1 may also be found in [18].

Observe that by choosing a fixed linear ordering of the array entries, each codeword from a $(\Lambda_1 \times \Lambda_2 \cdots \times \Lambda_{n-1} \times T, w, \lambda)$ *n*-D OOC *C* can be viewed as a binary constant weight (w)code of length $N = \Lambda_1 \Lambda_2 \cdots \Lambda_{n-1} T$. Moreover, by including the *T* distinct cyclic shifts of each codeword we obtain a corresponding constant weight binary code of size $T \cdot |C|$. It follows that

$$|C| \le \left\lfloor \frac{A(N, w, \lambda)}{T} \right\rfloor \tag{4}$$

From the equation (4) and Theorem 1.1 we obtain the following bounds for n-D OOCs.

Theorem 1.2 (Johnson Bound for *n*-D OOCs). If C is an $(\Lambda_1 \times \Lambda_2 \cdots \times \Lambda_{n-1} \times T, w, \lambda)$ OOC, then

$$\Phi(C) \le J(\Lambda_1 \times \Lambda_2 \cdots \times \Lambda_{n-1} \times T, w, \lambda)$$
(5)

$$= \left\lfloor \frac{N}{Tw} \left\lfloor \frac{N-1}{w-1} \left\lfloor \cdots \left\lfloor \frac{N-\lambda}{w-\lambda} \right\rfloor \right\rfloor \cdots \right\rfloor$$
(6)

$$:= \lfloor f(\Lambda_1 \times \Lambda_2 \cdots \times \Lambda_{n-1} \times T, w, \lambda) \rfloor, \qquad (7)$$

where $N = \Lambda_1 \Lambda_2 \cdots \Lambda_{n-1} T$. If $w^2 > N\lambda$ then

$$\Phi(C) \le \min\left\{\frac{N}{T}, \left\lfloor\frac{\frac{N}{T}(w-\lambda)}{w^2 - N\lambda}\right\rfloor\right\}.$$
(8)

We note that the bounds in Theorem 1.2 subsume the Johnson type bounds on 1, 2, and 3-dimensional codes, such as those found in [7, 9, 20]. Moreover, we can see from the theorem, that in a certain sense, maximum capacity is more intrinsically linked to the time spreading length than to the other dimensions.

Corollary 1.3. If $N = \Lambda_1 \cdot \Lambda_2 \cdots \Lambda_{s-1} \cdot T = \Lambda'_1 \cdot \Lambda'_2 \cdots \Lambda'_{t-1} \cdot T$ where $s, t \ge 1$ then

$$J(\Lambda_1 \times \dots \times \Lambda_{s-1} \times T, w, \lambda) = J(\Lambda'_1 \times \dots \times \Lambda'_{t-1} \times T, w, \lambda)$$
(9)

Some easy arithmetic gives the following.

Lemma 1.4. If $N = \Lambda_1 \cdot \Lambda_2 \cdots \Lambda_{n-1} \cdot T$, then

$$\frac{N}{T} \cdot J(N, w, \lambda) \le J(\Lambda_1 \times \dots \times \Lambda_{n-1} \times T, w, \lambda)$$
(10)

$$\leq \frac{N}{T} \cdot J(N, w, \lambda) + \frac{N}{T} - 1 \tag{11}$$

In particular, if $f(N, w, \lambda) - J(N, w, \lambda) < \frac{T}{N}$ (such as the case in which $f(N, w, \lambda)$ is integral) then

$$\frac{N}{T} \cdot J(N, w, \lambda) = J(\Lambda_1 \times \dots \times \Lambda_{n-1} \times T, w, \lambda).$$
(12)

Corollary 1.5. Let $N = \Lambda_1 \cdot \Lambda_2 \cdots \Lambda_{n-1} \cdot T$ where $T = \Lambda_n \cdot T'$. If $f(\Lambda_1 \times \cdots \times \Lambda_{n-1} \times T, w, \lambda) - J(\Lambda_1 \times \cdots \times \Lambda_{n-1} \times T, w, \lambda) < \frac{1}{\Lambda_n}$, then

$$\Lambda_n \cdot J(\Lambda_1 \times \dots \times \Lambda_{n-1} \times T, w, \lambda) = J(\Lambda_1 \times \dots \times \Lambda_{n-1} \Lambda_n \times T', w, \lambda)$$
(13)

$$= J(\Lambda_1 \times \dots \times \Lambda_{n-1} \times \Lambda_n \times T', w, \lambda).$$
(14)

As observed in [2] for 3-dimensional OOCs, an *n*-D OOC *C* with $\lambda_a = 0$ can be viewed as a constant weight (*w*) code of length $\frac{N}{T} = \Lambda_1 \Lambda_2 \cdots \Lambda_{n-1}$ over an alphabet of size T + 1 containing zero. By including the *T* distinct cyclic shifts of each codeword we obtain a corresponding constant weight code of size $T \cdot |C|$.

It follows that

$$|C| \le \left\lfloor \frac{A(\frac{N}{T}, w, \lambda)_{T+1}}{T} \right\rfloor.$$
(15)

From Theorem 1.1 and the equation (15) we obtain the following bound for ideal n-D OOCs.

Theorem 1.6. [Johnson Bound for Ideal n-D OOC] Let C be an $(\Lambda_1 \times \cdots \times \Lambda_{n-1} \times T, w, 0, \lambda_c)$ OOC, then

$$\Phi(C) \leq J(Ideal) = \left\lfloor \frac{N}{Tw} \left\lfloor \frac{N-T}{w-1} \left\lfloor \frac{N-2T}{w-2} \left\lfloor \cdots \left\lfloor \frac{N-\lambda T}{w-\lambda_c} \right\rfloor \right\rfloor \cdots \right\rfloor$$
(16)

where $N = \Lambda_1 \cdot \Lambda_2 \cdots \Lambda_{n-1} \cdot T$. In particular, if C has (maximal) weight $w = \frac{N}{T}$, then $\Phi(C) \leq T^{\lambda}$.

Note that the bound (16) is tight in certain cases, see *e.g.* the codes constructed in [17].

1.2 Ideal Codes and Sections

Suppose A is a codeword from an n-dimensional $(\Lambda_1 \times \Lambda_2 \times \cdots \times \Lambda_{n-1} \times T, w, \lambda_a, \lambda_c)$ OOC. For any fixed $i, 1 \leq i \leq n-1$, a Λ_i plane of A may be considered as an (n-1)-dimensional array. Such a plane is called a Λ_i section, or an *i*-section of A.




Figure 1: Two sections of a 3-D, $\Lambda \times S \times T$ (= $\Lambda_1 \times \Lambda_2 \times T$) codeword. Figure (b) is a 2-section, whereas (a) is a 1-section.

For $i \neq j$, the intersection of an *i*-section and a *j*-section is a section of degree 2, denoted an (i, j)-section. A section of degree $t \geq 3$ is defined in the analogous way, denoted an (i_1, i_1, \ldots, i_t) -section.

One way to ensure an *n*-D OOC is ideal, is to restrict the code to having at most one pulse per *i*-section, for some fixed *i*. Such a code is said to be AMOPS(i). For 2-D OOCs these are the At Most One Pulse Per Wavelength (AMOPW) codes [17, 13, 7]. For 3-D codes these are At-Most-One-Pulse-per-Plane (AMOPP) codes [25, 6].

If an *n*-D OOC, *C*, is restricted to having at most one pulse per (i_1, i_2, \ldots, i_j) -section, where $1 \leq j \leq n-1$, then *C* will be ideal, and is said to have *at most one pulse per section of degree j*, and is denoted an AMOPS (i_1, i_2, \ldots, i_j) code. If such a code has exactly one pulse per (i_1, i_2, \ldots, i_j) -section, then it is said to have a *single pulse per section of degree j*, and is denoted an SPS (i_1, i_2, \ldots, i_j) code. An ideal *n*-dimensional OOC is necessarily AMOPS $(1, 2, \ldots, n-1)$. It is readily seen that an AMOPS (i_1, i_2, \ldots, i_j) corresponds to a constant weight 1-dimensional code of length $m = \Lambda_{i_1} \cdot \Lambda_{i_2} \cdots \Lambda_{i_j}$ over an alphabet of size $\frac{N}{m} + 1$ (containing zero). Consequently, we obtain the following bounds on AMOPS codes.

Theorem 1.7. [Johnson Bound for AMOPS codes]

Let C be an (ideal) $(\Lambda_1 \times \cdots \times \Lambda_{n-1} \times T, w, 0, \lambda)$ -AMOPS (i_1, i_2, \dots, i_j) OOC, where $j \ge 1$ then

$$\Phi(C) \leq J(AMOPS)$$

$$= \left\lfloor \frac{N}{Tw} \left\lfloor \frac{N\left(1 - \frac{1}{M}\right)}{w - 1} \left\lfloor \frac{N\left(1 - \frac{2}{M}\right)}{w - 2} \left\lfloor \cdots \left\lfloor \frac{N\left(1 - \frac{\lambda}{M}\right)}{w - \lambda} \right\rfloor \right\rfloor \cdots \right\rfloor$$

$$\leq J(Ideal)$$
(17)

where $N = \Lambda_1 \cdot \Lambda_2 \cdots \Lambda_{n-1} \cdot T$, and $M = \Lambda_{i_1} \cdot \Lambda_{i_2} \cdots \Lambda_{i_j}$. In the extremal case where w = M, the bound (17) simplifies to $\frac{N^{\lambda+1}}{TM^{\lambda+1}}$.

In particular, if C is an $(\Lambda_1 \times \cdots \times \Lambda_{n-1} \times T, w, 0, \lambda)$ -AMOPS(i) OOC, then

$$\Phi(C) \le \left\lfloor \frac{N}{Tw} \left\lfloor \frac{N\left(1 - \frac{1}{\Lambda_i}\right)}{w - 1} \left\lfloor \frac{N\left(1 - \frac{2}{\Lambda_i}\right)}{w - 2} \left\lfloor \cdots \left\lfloor \frac{N\left(1 - \frac{\lambda}{\Lambda_i}\right)}{w - \lambda} \right\rfloor \right\rfloor \cdots \right\rfloor$$
(18)

where $N = \Lambda_1 \cdot \Lambda_2 \cdots \Lambda_{n-1} \cdot T$. In the extremal case where $w = \Lambda_i$, the bound (18) simplifies to $T^{\lambda} \prod_{j \neq i} \Lambda_j^{\lambda+1}$.

The bound (17) is tight in certain cases, see *e.g.* the codes constructed in [2, 7, 12, 13, 14, 25]. We also note that the bound (17) reduces to the bound in Theorem 1.6 when j = n - 1.

2 Iterative Constructions of Optimal n-D OOCs

Suppose C is a $(\Lambda \times T, w, \lambda_a, \lambda_c)$ 2-D OOC where $\Lambda = \Lambda_1 \cdot \Lambda_2$. Each codeword in C can be considered as a $\Lambda \times T$ array. Let $X \in C$ where $X = (x_{i,j})$. We may construct a corresponding 3-D $\Lambda_1 \times \Lambda_2 \times T$ codeword $Y_x = (y_{i,j,k}), 0 \le i < \Lambda_1, 0 \le j < \Lambda_2, 0 \le k < T$, where

$$y_{i,j,k} = c_{i+j\Lambda_1,k}.$$
(19)

It is readily verified that $C' = \{Y_x \mid x \in C\}$ is a $(\Lambda_1 \times \Lambda_2 \times T, w, \lambda_a, \lambda_c)$ 3-D OOC with |C'| = |C|. Inductively we arrive at the following.

Lemma 2.1. Let $\Lambda = \Lambda_1 \cdot \Lambda_2 \cdots \Lambda_{s-1}$. There exists an $(\Lambda \times T, w, \lambda_a, \lambda_c)$ 2-D OOC with capacity M if and only if there exists an $(\Lambda_1 \times \Lambda_2 \cdots \times \Lambda_{s-1} \times T, w, \lambda_a, \lambda_c)$ s-D OOC with capacity M.

An *n*-D OOC meeting any of the Johnson-type bounds established in the previous sections is referred to as a *J*-optimal code. With reference to Lemma 2.1 along with Corollary 1.3 we observe that a higher dimensional OOC with time spreading length T obtained from a J-optimal lower dimensional OOC by factoring the Λ_i 's will always be J-optimal. For example, each of the optimal codes in Table 1 give rise to optimal codes of dimension 4 or more.

Table 1: J-optimal ideal $(\Lambda_1 \times \Lambda_2 \times T)$ 3D OOCs. Unless stated otherwise, $\lambda_c = 1$.

Conditions	Type	Capacity	Reference
$w = \Lambda_1 \leq p$ for all p dividing $\Lambda_2 T$	SPS(1)	$\Lambda^2 T$	[12]
$w = q + 1 = \Lambda_1, \Lambda_2 = q > 3, T = p > q$	SPS(1)	$\Lambda^2 T$	[14]
$w = 4 = \Lambda_1 \le \Lambda_2 = q, T \ge 2$	SPS(1)	$\Lambda^2 T$	[14]
$w = 3 = \Lambda_1, \Lambda_2$ and T have the same parity	SPS(1)	$\Lambda^2 T$	[25]
$w = 3, \Lambda T(S-1)$ even, $\Lambda T(S-1)S \equiv 0 \mod 3$, and $S \equiv 0, 1 \mod 4$ if $T \equiv 2 \mod 4$ and Λ	AMOPS(1)	$\frac{\Lambda^2 T(S^2 - S)}{6}$	[25]
is odd.			
$w = \Lambda_1 \Lambda_2 \le p$ for all p dividing T	Ideal	$\Lambda^2 T$	[12]
$w = q, \Lambda ST = q^k - 1, T = q - 1$	Ideal	$\left\lfloor \frac{\Lambda S}{q} \left\lfloor \frac{T(\Lambda S - 1)}{q - 1} \right\rfloor$	[2]
$w = q^2, \Lambda S = q^2 + 1, T = q + 1, \lambda_c = q - 1$	Ideal	ΛS	[2]

p a prime, q a prime power	$, \theta(k,q) =$	$\frac{q^{k+1}-1}{a-1}$
------------------------------	-------------------	-------------------------

On the other hand, a J-optimal *n*-D OOC may correspond to an *s*-D OOC with s < n that is strictly asymptotically optimal. For example, from the bound (17), we see that a J-optimal $(5 \times 5 \times 5, 5, 0, 1)$ -AMOPS(1) OOC has capacity 125, whereas a J-optimal $(25 \times 5, 5, 0, 1)$ -AMOPS(1) OOC has capacity 150.

Corollary 2.2. Let $\Lambda = \Lambda_1 \cdot \Lambda_2 \cdots \Lambda_{n-1}$ be a positive integral factorisation.

- 1. If there exists a (resp. asymptotically) J-optimal $(\Lambda \times T, w, \lambda_a, \lambda_c)$ 2-D OOC then there exists a (resp. asymptotically) J-optimal $(\Lambda_1 \times \Lambda_2 \cdots \times \Lambda_{n-1} \times T, w, \lambda_a, \lambda_c)$ n-D OOC.
- 2. If there exists a (resp. asymptotically) J-optimal $(\Lambda_1 \times \Lambda_2 \cdots \times \Lambda_{n-1} \times T, w, \lambda_a, \lambda_c)$ n-D OOC, then there exists a $(\Lambda \times T, w, \lambda_a, \lambda_c)$ 2-D OOC which is at least asymptotically J-optimal.

Theorem 2.3. Let C be an $(\Lambda_1 \times \Lambda_2 \times \cdots \times \Lambda_{n-1} \times T, w, \lambda_a, \lambda_c)$ n-D OOC, $n \ge 1$. For any positive integral factorization $T = T_1 \cdot T_2$, there exists an $(T_1\Lambda_1 \times \Lambda_2 \times \cdots \times \Lambda_{n-1} \times T, w, \lambda'_a, \lambda'_c)$ n-D OOC, C' with $\lambda'_a \le \lambda_a, \lambda'_c \le \max\{\lambda_a, \lambda_c\}$, and $|C'| = T_1 \cdot |C|$.

Proof. For n = 1, 2 see Theorems 3 and 5 in [5]. The result then follows from Lemma 2.1.

There are many constructions of optimal 1-dimensional OOCs. From the Theorem 2.3 we see that in some cases optimal 1-dimensional OOCs give optimal *n*-D OOCs.

Corollary 2.4. Let C be an (N, w, λ) OOC with $N = \Lambda_1 \cdot \Lambda_2 \cdots \Lambda_{n-1} \cdot T$.

- 1. If C is J-optimal and $f(N, w, \lambda) J(N, w, \lambda) < \frac{T}{N}$, then a J-optimal $((\Lambda_1 \times \Lambda_2 \times \cdots \times \Lambda_{n-1} \times T, w, \lambda) n$ -D OOC exists.
- 2. If C is a member of a J-optimal (or asymptotically J-optimal) family then a family of $(\Lambda_1 \times \Lambda_2 \times \cdots \times \Lambda_{n-1} \times T, w, \lambda)$ n-D OOCs exists which is (at least) asymptotically optimal.

Proof. Follows from Theorem 2.3, (taking n = 1), Lemma 1.4, and the bounds in Theorem 1.2.

In [9], by considering orbits of lines in finite projective spaces, it is shown that for any prime power q, an infinite family of J-optimal $\left(\frac{q^{k+1-1}}{q-1}, q+1, 1\right)$ OOCs exits. From Corollary we now see that for any factorisation $\frac{q^{k+1-1}}{q-1} = \Lambda_1 \cdot \Lambda_2 \cdots \Lambda_{n-1} \cdot T$, an optimal $(\Lambda_1 \times \Lambda_2 \times \cdots \times \Lambda_{n-1} \times T, q+1, 1)$ n-D OOC exists.

For dimensions n > 1, we may also construct new optimal codes from others.

Corollary 2.5. Let C be an $(\Lambda_1 \times \Lambda_2 \times \cdots \times \Lambda_{n-1} \times T, w, \lambda)$ n-D OOC with $T = T_1 \cdot T_2$.

- 1. If C is J-optimal and $f(\Lambda_1 \times \Lambda_2 \times \cdots \times \Lambda_{n-1} \times T, w, \lambda) J(\Lambda_1 \times \Lambda_2 \times \cdots \times \Lambda_{n-1} \times T, w, \lambda) < \frac{1}{T_1}$ (in particular, if $f(\Lambda_1 \times \Lambda_2 \times \cdots \times \Lambda_{n-1} \times T, w, \lambda)$ is integral), then a J-optimal $(T_1\Lambda_1 \times \Lambda_2 \times \cdots \times \Lambda_{n-1} \times T, w, \lambda, w, \lambda)$ n-D OOC exists.
- 2. If C is a member of a J-optimal family, or an asymptotically J-optimal family then a family of $(\Lambda_1 \times \Lambda_2 \times \cdots \times \Lambda_{n-1} \times T, w, \lambda)$ n-D OOCs exists which is (at least) asymptotically optimal.

Proof. Follows from Theorem 2.3, Corollary 1.5, and the bounds in Theorem 1.2.

3 New optimal and asymptotically optimal codes

3.1 Preliminaries

Our techniques will rely heavily on the properties of finite projective and affine spaces. Such techniques have been used successfully in the construction of infinite families of optimal OOCs of one dimension, [9, 1, 16, 4, 3], two dimensions [7, 5], and three dimensions [6], [2]. We start with a brief overview of the necessary concepts. By PG(k,q) we denote the classical (or Desarguesian) finite projective geometry of dimension k and order q. PG(k,q) may be modeled with the affine (vector) space AG(k + 1, q) of dimension k + 1 over the finite field GF(q). Under this model, points of PG(k,q) correspond to 1-dimensional subspaces of AG(k,q), projective lines correspond to 2-dimensional affine subspaces, and so on. A *d*-flat Π in PG(k,q) is a subspace isomorphic to PG(d,q); if d = k - 1, the subspace Π is called a *hyperplane*. Elementary counting shows that the number of *d*-flats in PG(k,q) is given by the Gaussian coefficient

$$\begin{bmatrix} k+1\\ d+1 \end{bmatrix}_{q} = \frac{(q^{k+1}-1)(q^{k+1}-q)\cdots(q^{k+1}-q^{d})}{(q^{d+1}-1)(q^{d+1}-q)\cdots(q^{d+1}-q^{d})}.$$
(20)

In particular, the number of points of PG(k,q) is given by $\theta(k,q) = \frac{q^{k+1}-1}{q-1}$. We will use $\theta(k)$ to represent this number when q is understood to be the order of the field. Further, we shall denote by $\mathcal{L}(k)$ the number of lines in PG(k,q). For a point set A in PG(k,q) we shall denote by $\langle A \rangle$ the span of A, so $\langle A \rangle = PG(t,q)$ for some $t \leq k$.

A Singer group of PG(k, q) is a cyclic group of automorphisms acting sharply transitively on the points. The generator of such a group is known as a Singer cycle. Singer groups are known to exist in classical projective spaces of any order and dimension and their existence follows from that of primitive elements in a finite field.

Here, we make use of a Singer group that is most easily understood by modelling a finite projective space using a finite field. If we let β be a primitive element of $GF(q^{k+1})$, the points of $\Sigma = PG(k,q)$ can be represented by the field elements $\beta^0 = 1, \beta, \beta^2, \ldots, \beta^{n-1}$, where $n = \theta(k)$.

The non-zero elements of $GF(q^{k+1})$ form a cyclic group under multiplication. Multiplication by β induces an automorphism, or collineation, on the associated projective space PG(k,q) (see e.g. [22]). Denote by ϕ the collineation of Σ defined by $\beta^i \mapsto \beta^{i+1}$. The map ϕ clearly acts sharply transitively on the points of Σ .

As observed in [5], we can construct 2-dimensional codewords by considering orbits under some subgroup of G. Let $n = \theta(k) = \Lambda \cdot T$ where G is the Singer group of $\Sigma = PG(k,q)$. Since G is cyclic there exists an unique subgroup H of order T (H is the subgroup with generator ϕ^{Λ}).

Definition 1. Let Λ, T be integers such that $n = \theta(k) = \Lambda \cdot T$. For an arbitrary pointset S in $\Sigma = PG(k,q)$ we define the $\Lambda \times T$ incidence matrix $A = (a_{i,j}), 0 \le i \le \Lambda - 1, 0 \le j \le T - 1$ where $a_{i,j} = 1$ if and only if the point corresponding to $\beta^{i+\Lambda j}$ is in S.

If S is a pointset of Σ with corresponding $\Lambda \times T$ incidence matrix W of weight w, then ϕ^{Λ} induces a cyclic shift on the columns of W. For any such set S, consider its orbit $Orb_H(S)$ under the group H generated by ϕ^{Λ} . The set S has full H-orbit if $|Orb_H(S)| = T = \frac{n}{\Lambda}$ and short H-orbit otherwise. If S has full H-orbit then a representative member of the orbit and corresponding 2-dimensional codeword is chosen. The collection of all such codewords gives rise to a $(\Lambda \times T, w, \lambda_a, \lambda_c)$ 2-D OOC, where λ_a is determined by

$$\max_{1 \le i < j \le T} \left\{ |\phi^{\Lambda \cdot i}(S) \cap \phi^{\Lambda \cdot j}(S)| \right\}$$

and λ_c is determined by

$$\max_{1\leq i,j\leq \ T}\left\{|\phi^{\Lambda\cdot i}(S)\cap\phi^{\Lambda\cdot j}(S')|\right\}$$

3.2 Construction

Let $\Sigma = PG(k,q)$ where $G = \langle \phi \rangle$ is the Singer group of Σ as in the previous section. Our work will rely on the following results about orbits of flats.

Theorem 3.1 (Rao [22], Drudge[10]). In $\Sigma = PG(k,q)$, there exists a short *G*-orbit of *d*-flats if and only if $gcd(k+1, d+1) \neq 1$. In the case that d+1 divides k+1 there is a short orbit Swhich partitions the points of Σ (i.e. constitutes a *d*-spread of Σ). There is precisely one such orbit, and the *G*-stabilizer of any $\Pi \in S$ is $Stab_G(\Pi) = \langle \phi^{\frac{\theta(k)}{\theta(d)}} \rangle$.

3.2.1 Construction 1

For our first construction we mimic the methods of [2], whereby codewords correspond to lines that are not contained in any element of a d-spread of Σ .

For $d \geq 1$, let k > 1 such that d + 1 divides k + 1. Let $G = \langle \phi \rangle$ be the Singer group of $\Sigma = PG(k, q)$, as detailed above, and let S be the *d*-spread determined (as in Theorem 3.1) by G, where say $Stab_G(S) = H = \langle \phi^{\Lambda} \rangle$, where $\Lambda = \frac{\theta(k)}{\theta(d)}$.

Let ℓ be a line not contained in any spread element (a *d*-flat in S), and let A be the $\Lambda \times \theta(d)$ projective incidence array corresponding to ℓ . Observe that ℓ has a full H-orbit. H acts sharply transitively on the points of each spread element. It follows that A, considered as a $\Lambda \times \theta(d)$ codeword, satisfies $\lambda_a = 0$. For each such line ℓ , we choose a representative element of it's H-orbit, and include its corresponding incidence array as a codeword. The aggregate of these codewords gives an ideal ($\Lambda \times \theta(d), q + 1, 0, 1$)-3D OOC, C. Elementary counting gives

$$|C| = \frac{\mathcal{L}(k) - \mathcal{L}(d) \cdot \frac{\theta(k)}{\theta(d)}}{\theta(d)}$$

= $\frac{\theta(k)\theta(k-1)}{\theta(d)(q+1)} - \frac{\theta(d-1)\theta(k)}{\theta(d)(q+1)}$
= $\frac{\theta(k)}{\theta(d)(q+1)} \left[\theta(k-1) - \theta(d-1)\right].$ (21)

Comparing (21) with the bound in Theorem 1.6 shows these codes to be optimal.

Theorem 3.2. For d+1 a proper divisor of k+1, there exists a *J*-optimal $\left(\frac{\theta(k)}{\theta(d)} \times \theta(d), q+1, 0, 1\right)$ 2-D OOC.

With the observation that $\frac{\theta(k)}{\theta(d)} = \theta(m-1, q^{d+1})$, we have shown the following.

Corollary 3.3. For $d \ge 1$, m > 1, and for any positive integral factorisation $\Lambda_1 \cdot \Lambda_2 \cdots \Lambda_{n-1} = \theta(m-1, q^{d+1})$, there exists a J-optimal (ideal) $(\Lambda_1 \times \Lambda_2 \times \cdots \times \Lambda_{n-1} \times \theta(d), q+1, 0, 1)$ n-D OOC.

The following table will perhaps place this construction in context. Each of the optimal $\Lambda \times T$ constructions described in the table gives rise to optimal higher dimensional OOCs, with dimensions limited by the number of distinct factors in Λ .

Table 2: J-optimal families of ideal 2-D OOCs that give rise to higher dimensional optimal codes.

Parameters	Conditions	Reference			
${\bf Codes \ with} \ \lambda = {\bf 1}$					
$(\Lambda imes p, \Lambda, 0, 1)$	$\Lambda \leq p,$	[13]			
$\left(\theta(k,q^2) \times (q+1), q+1, 0, 1\right)$	$k \ge 1$	[7]			
$(\theta(k,q) \times (q-1), q, 0, 1)$	$k \ge 1$	[7]			
$((2^n+1) \times \theta(k,2), 2^n, 0, 1)$	$k \ge 1, n \ge 2$	[7]			
$\left(\frac{\theta(k)}{\theta(d)} \times \theta(d), q+1, 0, 1\right), d < k, d+1 k+1,$	Ideal	Theorem 3.2			
${\bf Codes \ with} \ \lambda \geq {\bf 2}$					
$(\Lambda imes p, \Lambda, 0, \lambda_c)$	$\Lambda \le p, \lambda_c \ge 1$	[17]			
$((q^n+1) \times \theta(k,q), q^n, 0, q-1)$	$k\geq 1,n\geq 2$	[7]			

(p prime, q a prime power)

3.2.2 Construction 2

In our second construction, codewords correspond to conics, and lines in $\Sigma = PG(3,q)$. An *m*-arc in PG(2,q) is a collection of m > 2 points such that no 3 points are incident with a common line. In PG(2,q), a (non-degenerate) conic is a (q+1)-arc. Elementary counting shows that this arc is complete (of maximal size) when q is odd. The (q+2)-arcs (hyperovals) exist in PG(2,q) if q is even and they are necessarily complete. Conics are a special case of the so called normal rational curves. We will be interested in the existence of large collections of arcs pairwise intersecting in at most two points. From Theorem 8 of [1], and its proof, we obtain the following.

Theorem 3.4 ([1]). In $\Pi = PG(2,q)$ there exists a family \mathcal{F} of conics, pairwise intersecting in at most 2 points, where $|\mathcal{F}| = q^3 - q^2$. Moreover, there is a distinguished line ℓ in Π disjoint from each member of \mathcal{F} .

Let $G = \langle \phi \rangle$ be the Singer group as above, and let S be the 1-spread determined (as in Theorem 3.1) by G where say $Stab_G(S) = H = \langle \phi^{\Lambda} \rangle$ where $\Lambda = \frac{\theta(3)}{q+1} = q^2 + 1$. Through each line ℓ of S, choose a plane $\pi(\ell)$. As the members of S are disjoint, each such plane

Through each line ℓ of S, choose a plane $\pi(\ell)$. As the members of S are disjoint, each such plane contains precisely one member of S (and therefore meets q^2 further members of S in precisely one point). As H acts sharply transitively on the points of each line in S, each such plane has full Horbit. A dimension argument shows that any two elements in the H-orbit of $\pi(\ell)$ meet precisely in ℓ . In each $\pi(\ell)$, let $\mathcal{F}(\ell)$ be a family of conics as in Theorem 3.4. Denote by $\mathcal{F} = \cup \mathcal{F}(\ell)$, where the union is taken over all spread lines.

Let $C \in \mathcal{F}$ be a conic, and let A be the $(q^2 + 1) \times (q + 1)$ incidence array corresponding to ℓ . From the above, it follows that A, considered as a codeword, satisfies $\lambda_a = 0$. For each such conic, choose a representative element of it's H-orbit, and include its corresponding incidence array as a codeword. The aggregate of these codewords gives an ideal $(q^2+1\times q+1, q+1, 0, 2)$ -2D OOC, C_1 . Note that $\lambda_c = 2$ follows from the fact that two conics in \mathcal{F} are either coplanar, and therefore meet in at most two points, or are not coplanar, in which case their intersection lies on the line common to the two planes.

Note that as in Construction 1, the *H*-orbits of non-spread lines of Σ correspond to an ideal $(q^2 + 1 \times q + 1, q + 1, 0, 1)$ -2D OOC, C_2 . Since a line and a conic meet in at most two points, we have $C = C_1 \cup C_2$ is an ideal $(q^2 + 1 \times q + 1, q + 1, 0, 2)$ -2D OOC. Moreover

$$|C| = (q^2 + 1) \cdot (q^3 - q^2) + \frac{\mathcal{L}(3) - (q^2 + 1)}{q + 1} = q(q^2 + 1)(q^2 - q + 1)$$
(22)

Comparing 22 to the bound in Theorem 1.6 shows C to be asymptotically optimal.

Theorem 3.5. For q a prime power, there exists an asymptotically optimal $(q^2 + 1 \times q + 1, q + 1, 0, 1)$ 2-D OOC.

Corollary 3.6. For any positive integral factorisation $\Lambda_1 \cdot \Lambda_2 \cdots \Lambda_{n-1} = q^2 + 1$, there exists an asymptotically optimal (ideal) $(\Lambda_1 \times \Lambda_2 \times \cdots \times \Lambda_{n-1} \times q + 1, q + 1, 0, 1)$ n-D OOC.

4 Conclusion

Here, we have generalized to higher dimensions the notions of optical orthogonal codes. We establish bounds on general *n*-dimensional OOCs, as well as specific types of ideal codes. The bounds presented here subsume many of the existing bounds appearing in the literature that are typically applied to codes of dimension three or less. We present two new constructions of ideal codes; one furnishes an infinite family of optimal codes for each dimension $n \ge 2$, and another which provides an asymptotically optimal family for each dimension $n \ge 2$.

References

- T. L. Alderson. Optical orthogonal codes and arcs in PG(d,q). Finite Fields Appl., 13(4):762-768, 2007.
- [2] T. L. Alderson. 3-dimensional optical orthogonal codes with ideal autocorrelation-bounds and optimal constructions. *Information Theory, IEEE Transactions on*, in press:1–7, 2017.
- [3] T. L. Alderson and K. E. Mellinger. Families of optimal OOCs with $\lambda = 2$. Information Theory, IEEE Transactions on, 54(8):3722–3724, Aug. 2008.
- [4] T. L. Alderson and Keith E. Mellinger. Constructions of optical orthogonal codes from finite geometry. SIAM J. Discrete Math., 21(3):785–793 (electronic), 2007.
- [5] T. L. Alderson and Keith E. Mellinger. 2-dimensional optical orthogonal codes from Singer groups. Discrete Appl. Math., 157(14):3008–3019, 2009.
- [6] Tim L. Alderson. New space/wavelength/time optical codes for ocdma. WSEAS Transactions on Communications, 16:35–42, 03 2017. Refereed Proceedings from conference at Cambridge.
- T.L. Alderson and Keith E. Mellinger. Spreads, arcs, and multiple wavelength codes. *Discrete Mathematics*, 311(13):1187 1196, 2011. Selected Papers from the 22nd British Combinatorial Conference.
- [8] A. Alvarado and E. Agrell. Four-dimensional coded modulation with bit-wise decoders for future optical communications. *Journal of Lightwave Technology*, 33(10):1993–2003, May 2015.
- [9] Fan R. K. Chung, Jawad A. Salehi, and Victor K. Wei. Optical orthogonal codes: design, analysis, and applications. *IEEE Trans. Inform. Theory*, 35(3):595–604, 1989.

- [10] Keldon Drudge. On the orbits of Singer groups and their subgroups. *Electron. J. Combin.*, 9(1):Research Paper 15, 10 pp. (electronic), 2002.
- [11] Robert M. Gagliardi and Antonio J. Mendez. Performance improvement with hybrid wdm and cdma optical communications. In Louis S. Lome, editor, *Wavelength Division Multiplexing Components*, volume 2690, pages 88–96. SPIE-Intl Soc Optical Eng, may 1996.
- [12] Sangin Kim, Kyungsik Yu, and N. Park. A new family of space/wavelength/time spread three-dimensional optical code for ocdma networks. *Journal of Lightwave Technology*, 18(4):502–511, April 2000.
- [13] W.C. Kwong and Guu-Chang Yang. Extended carrier-hopping prime codes for wavelengthtime optical code-division multiple access. *IEEE Trans. Commun.*, 52(7):1084–1091, July 2004.
- [14] X. Li, P. Fan, and K. W. Shum. Construction of space/wavelength/time spread optical code with large family size. *IEEE Communications Letters*, 16(6):893–896, June 2012.
- [15] A. J. Mendez and R. M. Gagliardi. Code division multiple access (CDMA) enhancement of wavelength division multiplexing (WDM) systems. In Proc. IEEE Int Communications ICC '95 Seattle, 'Gateway to Globalization' Conf, volume 1, pages 271–276 vol.1, June 1995.
- [16] Nobuko Miyamoto, Hirobumi Mizuno, and Satoshi Shinohara. Optical orthogonal codes obtained from conics on finite projective planes. *Finite Fields Appl.*, 10(3):405–411, 2004.
- [17] R. Omrani and P. Vijay Kumar. Improved constructions and bounds for 2-d optical orthogonal codes. ISIT 2005, Proc. Int. Symp. Inf. Theory, pages 127–131, Sept. 2005.
- [18] Reza Omrani, Petros Elia, and P. Vijay Kumar. New constructions and bounds for 2d optical orthogonal codes. In Tor Helleseth, Dilip V. Sarwate, Hong-Yeop Song, and Kyeongcheol Yang, editors, SETA, volume 3486 of Lecture Notes in Computer Science, pages 389–395. Springer, 2004.
- [19] Reza Omrani and P. Vijay Kumar. Codes for optical CDMA. In SETA, volume 4086 of Lecture Notes in Computer Science, pages 34–46. Springer, 2006.
- [20] J. Ortiz-Ubarri, O. Moreno, and A. Tirkel. Three-dimensional periodic optical orthogonal code for ocdma systems. In Proc. IEEE Information Theory Workshop (ITW), pages 170– 174, October 2011.
- [21] E. Park, A. J. Mendez, and E. M. Garmire. Temporal/spatial optical CDMA networksdesign, demonstration, and comparison with temporal networks. *IEEE Photonics Technology Letters*, 4(10):1160–1162, October 1992.
- [22] C. Radhakrishna Rao. Cyclical generation of linear subspaces in finite geometries. In Combinatorial Mathematics and its Applications (Proc. Conf., Univ. North Carolina, Chapel Hill, N.C., 1967), pages 515–535. Univ. North Carolina Press, Chapel Hill, N.C., 1969.
- [23] J. A. Salehi and C. A. Brackett. Code division multiple-access techniques in optical fiber networks. ii. systems performance analysis. *IEEE Transactions on Communications*, 37(8):834– 842, August 1989.
- [24] J.A. Salehi. Code-division multiple access techniques in optical fiber networks, part 1. fundamental principles. *IEEE Trans. Commun.*, 37:824–833, 1989.

- [25] Kenneth W. Shum. Optimal three-dimensional optical orthogonal codes of weight three. Des. Codes Cryptogr., 75(1):109–126, 2015.
- [26] Chengqian Xu and Yi Xian Yang. Algebraic constructions of asymptotically optimal twodimensional optical orthogonal codes. *Chinese Sci. Bull.*, 42(19):1659–1662, 1997.
- [27] Guu-Chang Yang and W.C. Kwong. Two-dimensional spatial signature patterns. Communications, IEEE Transactions on, 44(2):184–191, Feb 1996.
- [28] Guu-Chang Yang and W.C. Kwong. Performance comparison of multiwavelength cdma and wdma+cdma for fiber-optic networks. *Communications, IEEE Transactions on*, 45(11):1426–1434, November Nov 1997.