# CONVERGENCE AND OPTIMALITY OF THE ADAPTIVE MORLEY ELEMENT METHOD

JUN HU, ZHONGCI SHI, AND JINCHAO XU

ABSTRACT. This paper is devoted to the convergence and optimality analysis of the adaptive Morley element method for the fourth order elliptic problem. A new technique is developed to establish a quasi-orthogonality which is crucial for the convergence analysis of the adaptive nonconforming method. By introducing a new parameter-dependent error estimator and further establishing a discrete reliability property, sharp convergence and optimality estimates are then fully proved for the fourth order elliptic problem.

#### 1. INTRODUCTION

This paper is devoted to the study of adaptive nonconforming finite element methods for high order elliptic boundary value problems. The adaptive conforming finite element method for the second order elliptic problems has been a subject of extensive studies for many years since the pioneering work of Babuska and Rheinboldt [2], and its theory has become rather mature [32, 20, 27, 28, 6, 31, 30, 14]. For the nonconforming method, the a posteriori error theory of the second order elliptic problems has been studied only very recently [18, 19, 10, 9, 12, 11]; for the fourth order elliptic problem, only the a posteriori error estimate of the Morley element method can be found in the literature [22, 3, 33] and there have been no works on either convergence or optimality for any finite element methods for fourth order problems.

The main difficulty for the analysis of nonconforming finite element methods arises from the nonconformity of the discrete space and consequently the lack of the Galerkin-orthogonality which is a key ingredient for the convergence analysis of the adaptive conforming method of the second order elliptic problem [20, 27, 28, 30, 14]. For the nonconforming linear element of the Poisson equation, a quasi-orthogonality is established instead in [13] by using some special equivalency between the nonconforming linear element and the lowest order Raviart-Thomas element [25]. For the Morley element of the fourth order elliptic problem, however, it is unclear whether

Key words and phrases. A posteriori error estimator, the Morley element.

AMS Subject Classification: 65N30, 65N15, 35J25.

The first author was supported by the NSFC under Grant 10971005, and Foundation for the Author of National Excellent Doctoral Dissertation of PR China 200718, and partially supported by the Chinesisch-Deutsches Zentrum project GZ578. The third author was supported in part by NSF 0915153 and NSFC-10528102.

such type of equivalency still holds. We also note that the convergence (not to mention optimality) analysis of the adaptive conforming method is still missing for the fourth order elliptic problem in the literature.

This paper is devoted to the convergence and optimality analysis of the adaptive version of the Morley element [26, 29, 34]. Our analysis is based on an observation that a quasi-orthogonality can be obtained from a crucial local conservative property (that plays a critical role in a general study in [34]), of the Morley element method. Another ingredient is a new parameter dependent estimator which is introduced to analyze optimality of the adaptive nonconforming method. With the help of the discrete reliability which is established by introducing two interpolation operators between two nonconforming spaces, we show convergence and optimality of the adaptive algorithm.

The rest of the paper is organized as follows. In Section 2, we present the Kirchhoff plate problem and the Morley finite element method, and recall a posteriori error analysis due to [22]. In Section 3, we prove the quasi-orthogonality and then show reduction of some total error in Section 4 by introducing a new parameter-dependent estimator. To obtain optimality of the adaptive algorithm, we establish the discrete reliability in Section 5. Consequently, we show optimality of the adaptive Morley element method in Section 6. We give a brief comment on the extension of the theory to the Morley element method in three dimensions in Section 7. Also, we discuss the generalization to the nonconforming linear elements in both two and three dimensions therein. This extension gives an alternative analysis of the convergence result from [13]. The paper ends with Section 8 where we give the conclusion and some comments.

# 2. The Morley element for the Kirchhoff plate problem and an a posteriori error estimate

Let  $\Omega \subset \mathbb{R}^2$  be a bounded domain,  $\mathbb{E}$  the Young modulus, and  $\nu$  the Poisson ratio. For all  $2 \times 2$  symmetric matrices, the linear operator  $\mathcal{C}$  is defined by

$$\mathcal{C}\tau := \frac{\mathbb{E}}{12(1-\nu^2)} \left( (1-\nu)\tau + \nu \operatorname{tr}(\tau)I \right).$$

The bilinear form a(u, v) is defined by

(2.1) 
$$a(u,v) = (\mathcal{C}\nabla^2 u, \nabla^2 v)_{L^2(\Omega)}, \text{ for any } u, v \in W := H^2_0(\Omega),$$

where  $\nabla^2 u$  is the Hessian matrix of u. The corresponding energy norm is given by

(2.2) 
$$||u||_{\mathcal{C}}^2 := a(u, u) \text{ for any } u \in W,$$

which is equivalent to the usual norm  $|\cdot|_{H^2(\Omega)}$  for any  $u \in W$ .

We consider the Kirchhoff plate bending problem as follows: Given  $f \in L^2(\Omega)$ , find  $u \in W$  such that

(2.3) 
$$a(u,v) = (f,v)_{L^2(\Omega)} \quad \text{for all } v \in W.$$

We now present the Morley element. Suppose that  $\overline{\Omega}$  is covered exactly by a shape-regular triangulation  $\mathcal{T}_h$  consisting of triangles in 2D, see [16].  $\mathcal{E}_h$  is the set of all edges in  $\mathcal{T}_h$ ,  $\mathcal{E}_h(\Omega)$  is the set of interior edges, and  $\mathcal{E}(K)$  is the set of edges of any given element K in  $\mathcal{T}_h$ ;  $h_K = |K|^{1/2}$ , the size of the element  $K \in \mathcal{T}_h$ .  $\omega_K$  is the union of elements  $K' \in \mathcal{T}_h$  that share an edge with K, and  $\omega_e$  is the union of elements that share a common edge e. Given any edge  $e \in \mathcal{E}_h(\Omega)$  with the length  $h_e$  we assign one fixed unit normal  $\nu_e := (\nu_1, \nu_2)$  and tangential vector  $\tau_e := (-\nu_2, \nu_1)$ . For e on the boundary we choose  $\nu_e = \nu$  the unit outward normal to  $\Omega$ . Once  $\nu_e$  and  $\tau_e$  have been fixed on e, in relation to  $\nu_e$  one defines the elements  $K_- \in \mathcal{T}_h$  and  $K_+ \in \mathcal{T}_h$ , with  $e = K_+ \cap K_-$ . Given  $e \in \mathcal{E}_h(\Omega)$  and some  $\mathbb{R}^d$ -valued function v defined in  $\Omega$ , with d = 1, 2, we denote by  $[v] := (v|_{K_+})|_e - (v|_{K_-})|_e$  the jump of v across e.

The discrete space of the Morley finite element method is defined as follows [26, 29, 34]

(2.4)  
$$W_{h} := \{ v \in M_{2,h}, \int_{e} [\nabla_{h} v \cdot \nu_{e}] ds = 0 \text{ on } e \in \mathcal{E}_{h}(\Omega), \\ \text{and } \int_{e} \nabla v \cdot \nu_{e} ds = 0 \text{ on } e \in \mathcal{E}_{h} \cap \partial \Omega \},$$

where  $M_{2,h}$  is the space of piecewise polynomials of degree  $\leq 2$  over  $\mathcal{T}_h$  which are continuous at all the internal nodes and vanish at all the nodes on the boundary  $\partial\Omega$ , and  $\nabla_h$  the discrete gradient operator which is defined elementwise. We define

(2.5) 
$$a_h(u_h, v_h) := (\mathcal{C}\nabla_h^2 u_h, \nabla_h^2 v_h)_{L^2(\Omega)} \text{ for any } u_h, v_h \in W + W_h$$
$$\|u_h\|_{\mathcal{C}_h}^2 := a_h(u_h, u_h) \text{ for any } u_h \in W + W_h,$$

where the discrete Hessian operator  $\nabla_h^2$  is defined elementwise with respect to the triangulation  $\mathcal{T}_h$ .

We now consider the finite element discretization of (2.3) as follows: Find  $u_h \in W_h$  such that

(2.6) 
$$a_h(u_h, v_h) = (f, v_h)_{L^2(\Omega)} \quad \text{for all } v_h \in W_h$$

To recall the a posteriori error estimate for the Morley element, we first define an estimator on each element  $K \in \mathcal{T}_h$  as

(2.7) 
$$\eta_K = h_K^2 \|f\|_{L^2(K)} + \left(\sum_{e \in \partial K} h_K \|[\nabla_h^2 u_h \tau_e]\|_{L^2(e)}^2\right)^{1/2}.$$

For any  $S_h \subset \mathcal{T}_h$ , we define the estimator over  $S_h$  by

(2.8) 
$$\eta^2(u_h, S_h) := \sum_{K \in S_h} \eta_K^2.$$

In particular, for  $S_h = \mathcal{T}_h$ , we have

(2.9) 
$$\eta^2(u_h, \mathcal{T}_h) := \sum_{K \in \mathcal{T}_h} \eta_K^2.$$

We further define the oscillation  $\operatorname{osc}(f, \mathcal{T}_h)$  by

(2.10) 
$$\operatorname{osc}^{2}(f, \mathcal{T}_{h}) := \sum_{K \in \mathcal{T}_{h}} h_{K}^{4} \|f - f_{K}\|_{L^{2}(K)}^{2},$$

where  $f_K$  is the constant projection of f over K. For the estimator (2.9), we have the following reliability and efficiency whose proof can be found in [22].

**Lemma 2.1.** Let u be the solution of Problem (2.3), and  $u_h$  be the solution of Problem (2.6). Then,

(2.11) 
$$\|u - u_h\|_{\mathcal{C}_h} \lesssim \eta(u_h, \mathcal{T}_h) \lesssim \|u - u_h\|_{\mathcal{C}_h} + \operatorname{osc}(f, \mathcal{T}_h) .$$

Here and throughout the paper, we shall follow [35] to use the notation  $\leq$  and  $\approx$ . When we write

$$A_1 \lesssim B_1$$
, and  $A_2 \simeq B_2$ 

then there exist possible constants  $C_1$ ,  $c_2$  and  $C_2$  such that

$$A_1 \le C_1 B_1$$
, and  $c_2 B_2 \le A_2 \le C_2 B_2$ .

Given  $v \in H^2(\mathcal{T}_h) := \{v \in L^2(\Omega), v | K \in H^2(K), \text{ for any } K \in \mathcal{T}_h\}$ , we define the following residual

(2.12) 
$$\operatorname{Res}_{H}(v) = (f, v)_{L^{2}(\Omega)} - a_{h}(u_{H}, v), \text{ for any } v \in H^{2}(\mathcal{T}_{h}),$$

with  $u_H$  being the solution of the discrete problem (2.6) on  $\mathcal{T}_H$ , which is a nested and coarser mesh to  $\mathcal{T}_h$ ; namely,  $\mathcal{T}_h$  is some refinement of  $\mathcal{T}_H$ . It follows from the discrete problem (2.6) that

(2.13) 
$$\operatorname{Res}_{H}(v) = \operatorname{Res}_{H}(v - v_{H}), \text{ for any } v_{H} \in W_{H}.$$

**Lemma 2.2.** For any  $v \in W$ , it holds that

(2.14) 
$$|\operatorname{Res}_{H}(v)| \lesssim \left(\sum_{K \in \mathcal{T}_{H}} h_{K}^{4} \|f\|_{L^{2}(K)}^{2}\right)^{1/2} \|v\|_{\mathcal{C}} \text{ for any } v \in W.$$

The proof of the above lemma can be found in [33, 3, 22].

### 

## 3. QUASI-ORTHOGONALITY

In this section, we address one difficulty, namely the quasi-orthogonality, in the convergence analysis of the adaptive Morley element method. Our analysis is based on two interpolation operators: the canonical interpolation operator  $\Pi_h$  of the non-conforming space  $W_h$ , and the restriction operator  $I_H$  from the discrete space  $W_h$  on the mesh  $\mathcal{T}_h$  to the discrete space  $W_H$  on the nested and coarser mesh  $\mathcal{T}_H$  of  $\mathcal{T}_h$ .

Here and in what follows,  $\mathcal{N}_h$  denotes the set of nodes of the partition  $\mathcal{T}_h$ . We first define the canonical interpolation operator  $\Pi_h : W \to W_h$  by,

(3.1) 
$$(\Pi_h v)(P) = v(P), \int_e \nabla_h (\Pi_h v - v) \cdot \nu_e \, ds = 0 \text{ for any } v \in W, P \in \mathcal{N}_h, e \in \mathcal{E}_h.$$

5

**Lemma 3.1.** Let the interpolation operator  $\Pi_h$  be defined as in (3.1). Then,

(3.2) 
$$\int_{e} \nabla_{h} (v - \Pi_{h} v) \, ds = 0 \text{ for any } e \in \mathcal{E}_{h} \text{ and } v \in W,$$

(3.3) 
$$a_h(v - \Pi_h v, v_h) = 0$$
 for any  $v \in W, v_h$  piecewise quadratic,

(3.4) 
$$||v - \Pi_h v||_{L^2(K)} \lesssim h_K^2 |v|_{H^2(K)}$$
 for any  $K \in \mathcal{T}_h$  and  $v \in W$ .

The above properties are immediate from the definition of  $\Pi_h$ . Now we define the restriction interpolation operator  $I_H : W_h \to W_H$  by, for any  $v_h \in W_h$ , (3.5)

$$\begin{cases} (I_H v_h)(P) = v_h(P), P \in \mathcal{N}_H, \\ \int_e \frac{\partial (I_H v_h)}{\partial \nu_e} ds = \sum_{l=1}^\ell \int_{e_l} \frac{\partial v_h}{\partial \nu_e} ds, e \in \mathcal{E}_H \text{ with } e = e_1 \cup e_2 \dots \cup e_\ell \text{ and } e_i \in \mathcal{E}_h. \end{cases}$$

Before analyzing the properties of this interpolation, we state the following simple result.

**Lemma 3.2.** Let  $K_1, K_2 \in \mathcal{T}_h$  be two elements sharing a common edge e. If  $v_h \in W_h(K_1 \cup K_2)$  and  $\nabla_h^2 v_h = 0$ , then  $v_h \in P_1(K_1 \cup K_2)$ . Namely  $v_h$  is a polynomial of degree  $\leq 1$  over  $K_1 \cup K_2$ .

*Proof.* By the definition of  $W_h$ ,  $v_h$  is continuous on  $K_1 \cup K_2$ . Further  $\frac{\partial v_h}{\partial \nu_e}|_{K_1}$  and  $\frac{\partial v_h}{\partial \nu_e}|_{K_2}$  are two constant functions that must be equal since by the definition of  $W_h$  $\int_e [\frac{\partial v_h}{\partial \nu_e}] ds = 0$ . Thus v must belong to  $P_1(K_1 \cup K_2)$ .

The properties of the interpolation operator  $I_H$  are summarized in the following lemma.

**Lemma 3.3.** Let the interpolation operator  $I_H$  be defined as in (3.5). Then,

(3.6) 
$$\int_{e} \nabla_{h} (v_{h} - I_{H} v_{h}) \, ds = 0 \text{ for any } e \in \mathcal{E}_{H} \text{ and } v_{h} \in W_{h} \,,$$

(3.7) 
$$a_h(v_H, v_h - I_H v_h) = 0 \text{ for any } v_H \in W_H, v_h \in W_h,$$

(3.8) 
$$I_H v_h|_K = v_h|_K \text{ for any } K \in \mathcal{T}_h \cap \mathcal{T}_H \text{ and } v_h \in W_h,$$

(3.9) 
$$\|I_H v_h - v_h\|_{L^2(K)} \lesssim h_K^2 \|\nabla_h^2 v_h\|_{L^2(K)} \text{ for any } K \in \mathcal{T}_H \setminus \mathcal{T}_h \text{ and } v_h \in W_h.$$

*Proof.* The properties of (3.6), and (3.8) directly follow from the definition of the interpolation. We only need to prove (3.7) and the estimate (3.9).

We first define  $\sigma_H = \mathcal{C} \nabla_H^2 v_H$  to assert that

(3.10) 
$$\int_{e} \nabla_{h} (I - I_{H}) v_{h} \cdot \sigma_{H} \nu_{e} \, ds = 0 \text{ for any } e \in \mathcal{E}_{H}.$$

In fact, for  $e \in \mathcal{E}_H \setminus \mathcal{E}_h$ , this assertion follows from the fact that  $\sigma_H$  is a piecewise constant matrix with respect to  $\mathcal{T}_H$  and the definition of  $I_H$  in (3.5). For  $e \in \mathcal{E}_h \cap \mathcal{E}_H$ , the assertion follows from  $(I - I_H)v_h|_e = 0$ .

For the edge  $e \in \mathcal{E}_h$  which lies in the interior of some  $K \in \mathcal{T}_H$ , we can use the continuity of  $\int_e \nabla_h v_h \, ds$  over e and the fact  $\sigma_H$  is constant over K to show that

(3.11) 
$$\int_{e} [\nabla_h (I - I_H) v_h] \cdot \sigma_H \nu_e \, ds = 0$$

Whence, we integrate by parts and use (3.10) and (3.11) to conclude (3.7).

Now we turn to (3.9). In fact, both sides of (3.9) are semi-norms of the restriction  $W_h(K)$  of  $W_h$  on K. If the right hand side vanishes for some  $v_h \in W_h(K)$ , then  $v_h$  is a piecewise polynomial of degree  $\leq 1$  on K with respect to  $\mathcal{T}_h$ . It follows from Lemma 3.2 that  $v_h$  is a polynomial of degree  $\leq 1$  on K. Therefore the left hand side also vanishes for the same  $v_h$ . The desired result then follows from a scaling argument.

**Lemma 3.4.** (Quasi-orthogonality) Let  $\mathcal{T}_h$  be a refinement of  $\mathcal{T}_H$ , and  $u_h$  and  $u_H$  be the solutions of (2.6) on  $\mathcal{T}_h$  and  $\mathcal{T}_H$ , respectively. Then,

(3.12) 
$$|a_h(u_h - u_H, u - u_h)| \lesssim \sum_{K \in \mathcal{T}_H \setminus \mathcal{T}_h} h_K^2 ||f||_{L^2(K)} ||\nabla_h^2(u - u_h)||_{L^2(K)}$$

*Proof.* Let the interpolation operator  $\Pi_h$  be defined as in (3.1). Since  $\Pi_h$  is welldefined for any  $v_h \in W_h$  (in fact,  $\Pi_h v_h = v_h$ ) and  $a_h(u_h - u_H, (I - \Pi_h)(u - u_h)) = 0$ (by (3.3)), we have

(3.13) 
$$a_h(u_h - u_H, u - u_h) = a_h(u_h - u_H, \Pi_h(u - u_h)).$$

Let  $v_h = \prod_h (u - u_h)$  and the interpolation  $I_H v_h$  be defined as in (3.5). The combination of (2.6) and (2.12) leads to

(3.14) 
$$a_h(u_h - u_H, v_h) = (f, v_h)_{L^2(\Omega)} - a_h(u_H, v_h) = (f, (I - I_H)v_h)_{L^2(\Omega)} - a_h(u_H, (I - I_H)v_h).$$

By (3.8) and (3.9), we have

(3.15) 
$$|(f, (I - I_H)v_h)_{L^2(\Omega)}| \lesssim \sum_{K \in \mathcal{T}_H \setminus \mathcal{T}_h} h_K^2 ||f||_{L^2(K)} ||\nabla_h^2 v_h||_{L^2(K)}.$$

From (3.7) we have  $a_h(u_H, (I - I_H)v_h) = 0$ . Then, the desired result then follows from the triangle inequality and the approximation property of  $\Pi_h$ .

**Remark 3.5.** For the nonconforming  $P_1$  element of the Poisson equation, the quasiorthogonality was established in [13]. The analysis therein is based on some special equivalency between the nonconforming  $P_1$  and Raviart-Thomas elements. For the Stokes-like problem, the quasi-orthogonality of the nonconforming  $P_1$  element has been first proved in [24] based on some special relation of the nonconforming  $P_1$  and Raviart-Thomas elements. For the Morley element, it is unclear whether there exists similar equivalency or relation so far. **Remark 3.6.** This paper is a refined version of a technical report in 2009 [23], where it was the first time in the literature to make use of the conservative property of the nonconforming finite element space to analyze the quasi-orthogonality.

## 4. Reduction of a properly defined total error

In the rest of the paper, we shall establish convergence and optimality of our Adaptive Nonconforming Finite Element Method(ANFEM). Our analysis is based on two main ingredients: the strict reduction of some total error between two levels and the discrete reliability of the estimator. To this end, we shall first introduce a modified estimator  $\tilde{\eta}$  with a undetermined positive constant; we shall then borrow the concept of the total error of [14, 13] which contains the energy norm of the error and the scaled estimator  $\tilde{\eta}$ ; we shall finally show reduction of this total error. We shall establish the discrete reliability of the estimator in the next section.

Let us first define our adaptive algorithm. Given an initial shape regular triangulation  $\mathcal{T}_0$ , a right-hand side function  $f \in L^2(\Omega)$ , a tolerance  $\varepsilon > 0$ , and a parameter  $\theta \in (0, 1)$ . Hereafter, we shall replace the subscript h by an iteration counter called k.

Algorithm 4.1.  $[\mathcal{T}_N, u_N] = \mathbf{ANFEM}(\mathcal{T}_0, f, \varepsilon, \theta)$ 

 $\eta = \varepsilon \,, k = 0$ 

WHILE  $\eta \geq \varepsilon$ , DO

(1) Solve (2.6) on  $\mathcal{T}_k$ , to get the solution  $u_k$ .

(2) Compute the error estimator  $\eta = \eta(u_k, \mathcal{T}_k)$ .

(3) Mark the minimal element set  $\mathcal{M}_k$  such that

(4.1) 
$$\eta^2(u_k, \mathcal{M}_k) \ge \theta \, \eta^2(u_k, \mathcal{T}_k).$$

(4) Refine each triangle  $K \in \mathcal{M}_k$  by the newest vertex bisection and possible further refining to conformity to get  $\mathcal{T}_{k+1}$ .

k = k + 1.

## END WHILE

# $\mathcal{T}_N = \mathcal{T}_k.$ END ANFEM

In order to prove a strict reduction of some total error, we define the following modified estimator

(4.2) 
$$\tilde{\eta}^2(u_H, \mathcal{T}_H) := \sum_{K \in \mathcal{T}_H} \left( \beta_1 h_K^4 \| f \|_{L^2(K)}^2 + \eta_K^2 \right) \text{ with } \eta_K \text{ defined in (2.7)}$$

for some positive constant  $\beta_1$  to be determined later.

**Remark 4.1.** Note that, as we can see below, the modified error estimator  $\tilde{\eta}(u_H, \mathcal{T}_H)$  is only for the analysis, the final results concerning both convergence and optimality will be proved for Algorithm 4.1.

**Lemma 4.2.** Let  $\mathcal{T}_h$  be some refinement of  $\mathcal{T}_H$  with the bulk criterion (4.1), then there exist  $\rho > 0$  and a positive constant  $\beta \in (1 - \rho\theta, 1)$  such that

(4.3) 
$$\eta^2(u_H, \mathcal{T}_h) \le \beta \eta^2(u_H, \mathcal{T}_H) + (1 - \rho \theta - \beta) \eta^2(u_H, \mathcal{T}_H).$$

*Proof.* The result can be proved by following the idea in [14]. We give the details only for the readers' convenience. In fact, we have

(4.4) 
$$\eta^2(u_H, \mathcal{T}_h) = \eta^2(u_H, \mathcal{T}_H \cap \mathcal{T}_h) + \eta^2(u_H, \mathcal{T}_H \setminus \mathcal{T}_h).$$

For any  $K \in \mathcal{T}_H \setminus \mathcal{T}_h$ , we only need to consider the case where K is subdivided into  $K_1, K_2 \in \mathcal{T}_h$  with  $|K_1| = |K_2| = \frac{1}{2}|K|$ . By the definitions of  $h_K$  and  $\eta_K(u_H)$ , we have

(4.5) 
$$\sum_{i=1}^{2} \eta_{K_{i}}^{2}(u_{H}) := \sum_{i=1}^{2} \left( h_{K_{i}}^{2} \| f \|_{L^{2}(K_{i})} + \left( \sum_{\mathcal{E}_{h} \ni e \subset \partial K_{i}} h_{K_{i}} \| [\nabla_{H}^{2} u_{H} \tau_{e}] \|_{L^{2}(e)}^{2} \right)^{1/2} \right)^{2} \\ \leq \frac{1}{2^{1/2}} \eta_{K}^{2}(u_{H}) := \frac{1}{2^{1/2}} \left( h_{K}^{2} \| f \|_{L^{2}(K)} + \left( \sum_{\mathcal{E}_{H} \ni e \subset \partial K} h_{K} \| [\nabla_{H}^{2} u_{H} \tau_{e}] \|_{L^{2}(e)}^{2} \right)^{1/2} \right)^{2},$$

since  $[\nabla_H^2 u_H \tau_e] = 0$  over  $e = K_1 \cap K_2 \in \mathcal{E}_h$ . Consequently

(4.6) 
$$\sum_{K\in\mathcal{T}_H\setminus\mathcal{T}_h}\sum_{i=1}^2\eta_{K_i}^2(u_H)\leq\frac{1}{2^{1/2}}\eta^2(u_H,\mathcal{T}_H\setminus\mathcal{T}_h)\,,$$

and

(4.7) 
$$\eta^2(u_H, \mathcal{T}_h) \le \eta^2(u_H, \mathcal{T}_H) - \rho \eta^2(u_H, \mathcal{T}_H \setminus \mathcal{T}_h),$$

with  $\rho = 1 - \frac{1}{2^{1/2}}$ . Taking the positive parameter  $\beta$  with  $1 - \rho\theta < \beta < 1$ , the desired result follows by combining the above inequality and the bulk criterion (4.1).

**Lemma 4.3.** Let  $\mathcal{T}_h$  be some refinement of  $\mathcal{T}_H$ , then there exists  $\rho > 0$  such that

(4.8) 
$$\sum_{K \in \mathcal{T}_h} h_K^4 \|f\|_{L^2(K)}^2 \le \sum_{K \in \mathcal{T}_H} h_K^4 \|f\|_{L^2(K)}^2 - \rho \sum_{K \in \mathcal{T}_H \setminus \mathcal{T}_h} h_K^4 \|f\|_{L^2(K)}^2.$$

*Proof.* The proof immediately follows from the definition of the meshsize  $h_K$ .  $\Box$ 

**Lemma 4.4.** (Continuity of the estimator) Let  $u_h$  and  $u_H$  be the solutions to the discrete problem (2.6) on the meshes  $\mathcal{T}_h$  and  $\mathcal{T}_H$ , respectively. Given any positive constant  $\epsilon$ , there exists a positive constant  $\beta_2(\epsilon)$  dependent on  $\epsilon$  such that

(4.9) 
$$\eta^2(u_h, \mathcal{T}_h) \le (1+\epsilon)\eta^2(u_H, \mathcal{T}_h) + \frac{1}{\beta_2(\epsilon)} \|u_h - u_H\|_{\mathcal{C}_h}^2.$$

*Proof.* Given any  $K \in \mathcal{T}_h$ , it follows from the definitions of  $\eta_K(u_h)$  and  $\eta_K(u_H)$  in (4.5) that

(4.10)  
$$\left| \eta_{K}(u_{h}) - \eta_{K}(u_{H}) \right| = \left| \left( \sum_{\mathcal{E}_{h} \ni e \subset \partial K} h_{K} \| [\nabla_{h}^{2} u_{h} \tau_{e}] \|_{L^{2}(e)}^{2} \right)^{1/2} - \left( \sum_{\mathcal{E}_{h} \ni e \subset \partial K} h_{K} \| [\nabla_{H}^{2} u_{H} \tau_{e}] \|_{L^{2}(e)}^{2} \right)^{1/2} \right|$$
$$\leq \left( \sum_{\mathcal{E}_{h} \ni e \subset \partial K} h_{K} \| [\nabla_{h}^{2} (u_{h} - u_{H}) \tau_{e}] \|_{L^{2}(e)}^{2} \right)^{1/2}.$$

With  $e = K_1 \cap K_2 \in \mathcal{E}_h$ , we use the trace theorem and the fact that  $\nabla_h^2(u_h - u_H)$  is a piecewise constant matrix to get

$$(4.11) \\ \| [\nabla_h^2(u_h - u_H)\tau_e] \|_{L^2(e)} \le \| \nabla_h^2(u_h - u_H)\tau_e|_{K_1} \|_{L^2(e)} + \| \nabla_h^2(u_h - u_H)\tau_e|_{K_2} \|_{L^2(e)} \\ \lesssim h_K^{-1/2} \| \nabla_h^2(u_h - u_H) \|_{L^2(\omega_e)} ,$$

which gives

(4.12) 
$$|\eta_K(u_h) - \eta_K(u_H)| \lesssim ||\nabla_h^2(u_h - u_H)||_{L^2(\omega_K)}.$$

Applying the Young inequality with any positive constant  $\epsilon$  and summarizing over all elements in  $\mathcal{T}_h$  completes the proof of the lemma.

**Theorem 4.5.** Let u be the solution to the problem (2.3), and  $u_H$  and  $u_h$  be the solutions to the discrete problem (2.6) on the meshes  $\mathcal{T}_H$  and  $\mathcal{T}_h$ , respectively. Then, there exists positive constants  $\gamma_1$ ,  $\beta_1$ , and  $0 < \alpha < 1$  with

(4.13) 
$$\|u - u_h\|_{\mathcal{C}_h}^2 + \gamma_1 \tilde{\eta}^2(u_h, \mathcal{T}_h) \le \alpha(\|u - u_H\|_{\mathcal{C}_H}^2 + \gamma_1 \tilde{\eta}^2(u_H, \mathcal{T}_H)).$$

*Proof.* Let  $\delta$ ,  $\gamma_1$ , and  $\gamma_2$ , be three positive constants to be chosen later. Applying the Young inequality to Lemma 3.4 and adding the resulting estimate to the inequality (4.9) leads to

$$(1-\delta)\|u-u_{h}\|_{\mathcal{C}_{h}}^{2}+\gamma_{1}\eta^{2}(u_{h},\mathcal{T}_{h})+\gamma_{2}\sum_{K\in\mathcal{T}_{h}}h_{K}^{4}\|f\|_{L^{2}(K)}^{2}$$

$$(4.14) \qquad \leq \|u-u_{H}\|_{\mathcal{C}_{H}}^{2}+\gamma_{1}(1+\epsilon)\eta^{2}(u_{H},\mathcal{T}_{h})+(\frac{\gamma_{1}}{\beta_{2}(\epsilon)}-1)\|u_{h}-u_{H}\|_{\mathcal{C}_{h}}^{2}$$

$$+\gamma_{2}\sum_{K\in\mathcal{T}_{H}}h_{K}^{4}\|f\|_{L^{2}(K)}^{2}+(C_{1}(\delta)-\rho\gamma_{2})\sum_{K\in\mathcal{T}_{H}\setminus\mathcal{T}_{h}}h_{K}^{4}\|f\|_{L^{2}(K)}^{2},$$

with the positive constant  $\rho$  from (4.8). We note that the bound for  $\eta^2(u_H, \mathcal{T}_h)$  is given in Lemma 4.2. Hence

$$(1-\delta)\|u-u_{h}\|_{\mathcal{C}_{h}}^{2}+\gamma_{1}\eta^{2}(u_{h},\mathcal{T}_{h})+\gamma_{2}\sum_{K\in\mathcal{T}_{h}}h_{K}^{4}\|f\|_{L^{2}(K)}^{2}$$

$$\leq \|u-u_{H}\|_{\mathcal{C}_{H}}^{2}+\gamma_{1}((1-\rho\theta-\beta)(1+\epsilon)+\epsilon\beta)\eta^{2}(u_{H},\mathcal{T}_{H})+\gamma_{1}\beta\eta^{2}(u_{H},\mathcal{T}_{H})$$

$$(4.15) +(\frac{\gamma_{1}}{\beta_{2}(\epsilon)}-1)\|u_{h}-u_{H}\|_{\mathcal{C}_{h}}^{2}+(C_{1}(\delta)-\rho\gamma_{2})\sum_{K\in\mathcal{T}_{H}\setminus\mathcal{T}_{h}}h_{K}^{4}\|f\|_{L^{2}(K)}^{2}$$

$$+\gamma_{2}\sum_{K\in\mathcal{T}_{H}}h_{K}^{4}\|f\|_{L^{2}(K)}^{2},$$

with  $\rho$  and  $\beta$  from Lemma 4.2. In what follows we shall choose the parameters  $\alpha$ ,  $\beta$ ,  $\gamma_1$ ,  $\gamma_2$ , and  $\delta$  to achieve the reduction of the total error. We first set

(4.16) 
$$\gamma_2 = \frac{C_1(\delta)}{\rho}, \gamma_1 = \beta_2(\epsilon), \text{ and } \beta = (1 - \rho\theta)(1 + \epsilon)$$

which leads to

(4.17) 
$$(1-\delta)\|u-u_{h}\|_{\mathcal{C}_{h}}^{2}+\gamma_{1}\eta^{2}(u_{h},\mathcal{T}_{h})+\gamma_{2}\sum_{K\in\mathcal{T}_{h}}h_{K}^{4}\|f\|_{L^{2}(K)}^{2}$$
$$\leq \|u-u_{H}\|_{\mathcal{C}_{H}}^{2}+\gamma_{1}\beta\eta^{2}(u_{H},\mathcal{T}_{H})+\gamma_{2}\sum_{K\in\mathcal{T}_{H}}h_{K}^{4}\|f\|_{L^{2}(K)}^{2}.$$

We choose  $\epsilon$  to be small enough such that  $0 < \beta < 1$ . Let the positive constant  $\alpha$  with  $\beta < \alpha < 1$  be determined later, this gives

$$(1-\delta)\|u-u_{h}\|_{\mathcal{C}_{h}}^{2}+\gamma_{1}\eta^{2}(u_{h},\mathcal{T}_{h})+\gamma_{2}\sum_{K\in\mathcal{T}_{h}}h_{K}^{4}\|f\|_{L^{2}(K)}^{2}$$

$$\leq \alpha\left((1-\delta)\|u-u_{H}\|_{\mathcal{C}_{H}}^{2}+\gamma_{1}\eta^{2}(u_{H},\mathcal{T}_{H})+\gamma_{2}\sum_{K\in\mathcal{T}_{H}}h_{K}^{4}\|f\|_{L^{2}(K)}^{2}\right)$$

$$+(1-\alpha(1-\delta))\|u-u_{H}\|_{\mathcal{C}_{H}}^{2}+\gamma_{1}(\beta-\alpha)\eta^{2}(u_{H},\mathcal{T}_{H})$$

$$+\gamma_{2}(1-\alpha)\sum_{K\in\mathcal{T}_{H}}h_{K}^{4}\|f\|_{L^{2}(K)}^{2}.$$

Recalling the reliability of  $\eta(u_H, \mathcal{T}_H)$  with the reliability coefficient  $C_{Rel}$  in Lemma 2.1

(4.19) 
$$\|u - u_H\|_{\mathcal{C}_H}^2 \le C_{Rel} \eta^2(u_H, \mathcal{T}_H) ,$$

and the fact that

(4.20) 
$$\sum_{K \in \mathcal{T}_H} h_K^4 \|f\|_{L^2(K)}^2 \le \eta^2(u_H, \mathcal{T}_H).$$

Whence we derive as

$$(4.21) (1 - \alpha(1 - \delta)) \|u - u_H\|_{\mathcal{C}_H}^2 + \gamma_1(\beta - \alpha)\eta^2(u_H, \mathcal{T}_H) + \gamma_2(1 - \alpha) \sum_{K \in \mathcal{T}_H} h_K^4 \|f\|_{L^2(K)}^2 \\ \leq \left((1 - \alpha(1 - \delta))C_{Rel} + \gamma_1(\beta - \alpha) + \gamma_2(1 - \alpha)\right)\eta^2(u_H, \mathcal{T}_H),$$

provided that  $0 < \delta < 1$ . Then, the choice of  $\alpha = \frac{\gamma_1 \beta + \gamma_2 + C_{Rel}}{\gamma_1 + \gamma_2 + C_{Rel}(1 - \delta)} > \beta$  gives

(4.22) 
$$(1-\delta)\|u-u_{h}\|_{\mathcal{C}_{h}}^{2}+\gamma_{1}\eta^{2}(u_{h},\mathcal{T}_{h})+\gamma_{2}\sum_{K\in\mathcal{T}_{h}}h_{K}^{4}\|f\|_{L^{2}(K)}^{2}$$
$$\leq \alpha \left((1-\delta)\|u-u_{H}\|_{\mathcal{C}_{H}}^{2}+\gamma_{1}\eta^{2}(u_{H},\mathcal{T}_{H})+\gamma_{2}\sum_{K\in\mathcal{T}_{H}}h_{K}^{4}\|f\|_{L^{2}(K)}^{2}\right)$$

We choose such that  $0 < \delta < \min(\frac{\gamma_1(1-\beta)}{C_{Rel}}, 1)$  to assure that  $\alpha < 1$ . Finally, we take  $\beta_1 = \gamma_2/\gamma_1$  and redefine  $\gamma_1 = \gamma_1/(1-\delta)$  to end the proof.

# 5. DISCRETE RELIABILITY

This section is devoted to the discrete reliability of the estimator  $\eta(u_H, \mathcal{T}_H)$ . The analysis needs the prolongation operator  $I'_h : W_H \to W_h$  defined as follows. Given  $P \in \mathcal{N}_h$  and  $e \in \mathcal{E}_h$ , the nodal patch  $\omega_{P,H}$  of P and the edge patch  $\omega_{e,H}$  of e with respect to the mesh  $\mathcal{T}_H$  are defined by, respectively,

(5.1) 
$$\omega_{P,H} := \{ K \in \mathcal{T}_H, P \in \partial K \text{ or } P \text{ is in the interior of } K \}, \\ \omega_{e,H} := \{ K \in \mathcal{T}_H, e \subset \partial K \text{ or } e \text{ is in the interior of } K \}.$$

Define  $\xi_P = \operatorname{card}(\omega_{P,H})$  and  $\xi_e = \operatorname{card}(\omega_{e,H})$ . We define the prolongation interpolation  $I'_h v_H \in W_h$  by, for any  $v_H \in W_H$ ,

(5.2) 
$$\begin{cases} (I'_h v_H)(P) = \frac{1}{\xi_P} \sum_{K \in \omega_{P,H}} v_H|_K(P) \text{ for any } P \in \mathcal{N}_h, \\ \int_e \frac{\partial (I'_h v_H)}{\partial \nu_e} ds = \frac{1}{\xi_e} \sum_{K \in \omega_{e,H}} \int_e \frac{\partial (v_H|_K)}{\partial \nu_e} ds \text{ for any } e \in \mathcal{E}_h \end{cases}$$

**Lemma 5.1.** Let  $K_1, K_2 \in \mathcal{T}_H$  be two elements sharing a common edge e with two endpoints  $P_1$  and  $P_2$ . Suppose that  $v_H \in W_H$  and  $\nabla_H v_H$  is continuous over e. Then,  $v_H$  is continuous over e.

*Proof.* We can assume that the common edge e shared by  $K_1$  and  $K_2$  lies along the x-axis. Then, v can be expressed as

$$v_H|_{K_1} = a_0 + a_1 x + a_2 y + a_3 x y + a_4 x^2 + a_5 y^2$$
, and  $v_H|_{K_2} = b_0 + b_1 x + b_2 y + b_3 x y + b_4 x^2 + b_5 y^2$   
Since  $\frac{\partial v_H}{\partial x}$  is continuous over  $e$ , we have  $a_1 = b_1$  and  $a_4 = b_4$ . The continuity of  $\frac{\partial v_H}{\partial y}$   
over  $e$  gives  $a_2 = b_2$  and  $a_3 = b_3$ . Finally,  $v_H|_{K_1}(P_\ell) = v_H|_{K_2}(P_\ell), \ell = 1, 2$ , concludes  
 $a_0 = b_0$ . Therefore,  $v_H$  is continuous over  $e$ .

**Lemma 5.2.** Let the interpolation operator  $I'_h$  be defined as in (5.2). Then,

(5.3) 
$$\|\nabla_h^2 (I'_h v_H - v_H)\|_{L^2(\Omega)}^2 \lesssim \sum_{K \in \mathcal{T}_H \setminus \mathcal{T}_h} \sum_{e \subset \partial K} h_K \|[\nabla_H^2 v_H \tau_e]\|_{L^2(e)}^2$$
 for any  $v_H \in W_H$ .

Proof. It follows from the definition of  $I'_h$  (5.2) that  $I'_h v_H|_K = v_H|_K$  for any  $K \in \mathcal{T}_H \cap \mathcal{T}_h$ . Therefore, we only need to estimate  $\|\nabla_h^2(I'_h v_H - v_H)\|_{L^2(K)}$  for  $K \in \mathcal{T}_H \setminus \mathcal{T}_h$ . To prove the desired result, it is sufficient to show that (5.4)

$$\|\nabla_h^2(I_h'v_H - v_H)\|_{L^2(K)} \lesssim \sum_{e \subset \partial K} h_K \|[\nabla_H^2 v_H \tau_e]\|_{L^2(e)}^2 \text{ for any } v_H \in W_H \text{ and } K \in \mathcal{T}_H \setminus \mathcal{T}_h.$$

For any  $e \in \mathcal{E}_H$ ,  $\|[\nabla_H^2 v_H \tau_e]\|_{L^2(e)} = 0$  indicates that there are no jumps over e for all tangential components of  $\nabla_H^2 v_H$ , which in turn implies that  $\nabla_H v_H$  is continuous over e since  $\nabla_H v_H$  is average continuous over e. Since  $v_H$  is continuous at all the internal nodes, Lemma 5.1 proves that  $v_H$  is continuous over e. Whence,  $I'_h v_H|_K =$  $v_H|_K$  provided that  $\|[\nabla_H^2 v_H \tau_e]\|_{L^2(e)} = 0$  for any  $e \in \partial K \subset \mathcal{E}_H$ . Finally, the local quasi-uniformity of the mesh together with a scaling argument leads to the estimate (5.4).

**Remark 5.3.** An easy observation finds that the positive constant in (5.4) depends on the following ratio

(5.5) 
$$\mu = \max_{K \in \mathcal{T}_H \setminus \mathcal{T}_h} \max_{\mathcal{T}_h \ni T \subset K} \frac{h_K}{h_T}.$$

In the analysis of optimality of the adaptive finite element method, this dependence is not allowed since we only know that  $\mathcal{T}_h$  is some refinement of  $\mathcal{T}_H$  by the newest vertex bisection and the boundness of  $\mu$  is not guaranteed.

To overcome the above difficulty, we introduce a modified prolongation operator  $J_h$  which preserves the local projection property. We need the prolongation operator  $\Pi: W_H \to W_H^C$ , where  $W_H^C \subset W$  is some conforming finite element space over the mesh  $\mathcal{T}_H$ . Here we take  $W_H^C$  as the Hsieh-Clough-Tocher finite element space over the mesh  $\mathcal{T}_H$  [16, 7].

Let  $\mathcal{F}$  be any (global) degree of freedom of  $W_H^C$ , i.e.,  $\mathcal{F}$  is either the evaluation of a shape function or its first order derivatives at an interior node of  $\mathcal{T}_H$ , or the evaluation of the normal derivative of a shape function at a node on an interior edge. For  $v_H \in W_H$ , we define [8]

(5.6) 
$$\mathcal{F}(\Pi v_H) = \frac{1}{|\omega_{\mathcal{F}}|} \sum_{K \in \omega_{\mathcal{F}}} \mathcal{F}(v_H|_K)$$

where  $\omega_{\mathcal{F}}$  is the set of triangles in  $\mathcal{T}_H$  that share the degree of freedom  $\mathcal{F}$ , and  $|\omega_{\mathcal{F}}|$  is the number of elements of  $\omega_{\mathcal{F}}$ . Then a similar argument of [8] proves

(5.7) 
$$\|\nabla_{H}^{2}(v_{H} - \Pi v_{H})\|_{L^{2}(\Omega)}^{2} \lesssim \sum_{e \in \mathcal{E}_{H}} h_{e} \|[\nabla_{H}^{2}v_{H}\tau_{e}]\|_{L^{2}(e)}^{2}$$

Define

$$\Omega_{\mathcal{R}} := \operatorname{interior}(\bigcup \{ K : K \in \mathcal{T}_H \setminus \mathcal{T}_h, \}),$$

and

$$\Omega_{\mathcal{C}} := \operatorname{interior}(\bigcup \{ K : K \in \mathcal{T}_H \cap \mathcal{T}_h, \partial K \cap \partial \Omega_{\mathcal{R}} = \emptyset \}).$$

The main idea herein is to take the mixture of the prolongation operators  $I'_h$  and  $\Pi$ . More precisely, we use  $\Pi$  in the region  $\Omega_{\mathcal{R}}$  where the elements of  $\mathcal{T}_H$  are refined and take  $I'_h$  on  $\Omega_{\mathcal{C}}$ , and we define some mixture in the layer between them. This leads to the prolongation operator  $J_h: W_H \to W_h$  by

$$\mathcal{J}_h v_H = \begin{cases} \Pi_h \Pi v_H & \text{on } \Omega_{\mathcal{R}}, \\ I'_h v_H & \text{on } \Omega_{\mathcal{C}}, \\ v_{h,tr} & \text{on } \Omega \setminus (\Omega_{\mathcal{R}} \cup \Omega_{\mathcal{C}}), \end{cases}$$

where  $v_{h,tr}$  is defined by

(5.8)  
$$v_{h,tr}(P) = \begin{cases} (\Pi v_H)(P) \text{ if } P \in \partial \Omega_{\mathcal{R}}, \\ (I'_h v_H)(P) \text{ otherwise }, \end{cases} \text{ for } P \in \mathcal{N}_h, \\ \int_e \nabla_h v_{h,tr} \cdot \nu_e ds = \begin{cases} \int_e \nabla_h \Pi v_H \cdot \nu_e ds & \text{ if } e \subset \partial \Omega_{\mathcal{R}} \\ \int_e \nabla_h I'_h v_H \cdot \nu_e ds & \text{ otherwise} \end{cases} \text{ for } e \in \mathcal{E}_h \end{cases}$$

Define

$$\mathcal{M}_{H,h} := \{ K \in \mathcal{T}_H, \partial K \cap \overline{\cup(\mathcal{T}_H \setminus \mathcal{T}_h)} \neq \emptyset \}.$$

**Remark 5.4.** It follows immediately from regularity of the mesh  $\mathcal{T}_H$  that

$$#\mathcal{M}_{H,h} \leq \kappa #\mathcal{T}_H \setminus \mathcal{T}_h$$

for a positive constant  $\kappa \geq 1$  which is only dependent on the initial mesh  $\mathcal{T}_0$ .

Lemma 5.5. It holds true that

(5.9) 
$$\|\nabla_h^2 (J_h v_H - v_H)\|_{L^2(\Omega)}^2 \lesssim \sum_{K \in \mathcal{M}_{H,h}} \sum_{e \subset \partial K} h_K \|[\nabla_H^2 v_H \tau_e]\|_{L^2(e)}^2$$
 for any  $v_H \in W_H$ .

Proof. We only need to use the scaling argument like that in Lemma 5.2 in the layer  $\Omega \setminus (\Omega_{\mathcal{R}} \cup \Omega_{\mathcal{C}})$ . The desired result follows from the estimate (5.7) and the local projection property  $J_h v_H|_K = v_H|_K$  for  $K \in \Omega_{\mathcal{C}}$ .

**Lemma 5.6.** (Discrete reliability) It holds that

(5.10) 
$$\|u_h - u_H\|_{\mathcal{C}_h}^2 \lesssim \eta^2(u_H, \mathcal{M}_{H,h})$$

*Proof.* For any  $v_h \in W_h$ , we deduce from the discrete problem (2.6) that

(5.11) 
$$||u_h - u_H||^2_{\mathcal{C}_h} = a_h(u_h - u_H, u_h - v_h) + a_h(u_h - u_H, v_h - u_H) = J_1 + J_2,$$

where

$$J_1 = \text{Res}_H(u_h - v_h)$$
, and  $J_2 = a_h(u_h - u_H, v_h - u_H)$ .

Thanks to (2.13), (3.8) and (3.9), the residual  $J_1$  can be bounded by a similar argument for the term on the right hand-side of (3.14), which implies

(5.12) 
$$|J_1| = |\operatorname{Res}_H(u_h - v_h)| = |\operatorname{Res}_H((I - I_H)(u_h - v_h))| \lesssim \sum_{K \in \mathcal{M}_{H,h}} h_K^2 ||f||_{L^2(K)} ||\nabla_h^2(u_h - v_h)||_{L^2(K)}.$$

Since  $v_h \in W_h$  is arbitrary, we apply the Young and Cauchy-Schwarz inequalities in (5.11) to obtain that

(5.13) 
$$\|u_h - u_H\|_{\mathcal{C}_h}^2 \lesssim \sum_{K \in \mathcal{M}_{H,h}} h_K^4 \|f\|_{L^2(K)}^2 + \inf_{v_h \in W_h} \|v_h - u_H\|_{\mathcal{C}_h}^2.$$

Taking  $v_h = J_h u_H$  and applying (5.9) complete the proof of the Lemma.

We end this section by applying the previous discrete reliability to show a result indicating that the bulk criterion is in some sense a necessary condition for reduction of the energy norm between two levels.

**Lemma 5.7.** If  $\mathcal{T}_h$  is a refinement of  $\mathcal{T}_H$  such that the following reduction holds

(5.14) 
$$\|u - u_h\|_{\mathcal{C}_h}^2 + \operatorname{osc}^2(f, \mathcal{T}_h) \le \alpha'(\|u - u_H\|_{\mathcal{C}_H}^2 + \operatorname{osc}^2(f, \mathcal{T}_H)),$$

for some  $0 < \alpha' < 1$ , then there exists  $0 < \theta_* < 1$  such that

(5.15) 
$$\theta_*\eta^2(u_H,\mathcal{T}_H) \le \eta^2(u_H,\mathcal{M}_{H,h}).$$

*Proof.* We start with the following decomposition

$$(1 - \alpha')(\|u - u_H\|_{\mathcal{C}_H}^2 + \operatorname{osc}^2(f, \mathcal{T}_H))$$

$$(5.16) \qquad \leq \|u - u_H\|_{\mathcal{C}_H}^2 + \operatorname{osc}^2(f, \mathcal{T}_H) - \|u - u_h\|_{\mathcal{C}_h}^2 - \operatorname{osc}^2(f, \mathcal{T}_h)$$

$$= \|u_H - u_h\|_{\mathcal{C}_h}^2 + 2a_h(u - u_h, u_h - u_H) + \operatorname{osc}^2(f, \mathcal{T}_H) - \operatorname{osc}^2(f, \mathcal{T}_h).$$

By the discrete reliability of Lemma 5.6 with the coefficient  $C_{Drel}$ , we have

(5.17) 
$$\|u_h - u_H\|_{\mathcal{C}_h}^2 \le C_{Drel} \eta^2(u_H, \mathcal{M}_{H,h}) .$$

The quasi-orthogonality in Lemma 3.4 with the coefficient  $C_{QO}$  yields

(5.18) 
$$|2a_h(u-u_h, u_h-u_H)| \le 2C_{QO} ||u-u_h||_{\mathcal{C}_h} \Big(\sum_{K \in \mathcal{M}_{H,h}} h_K^4 ||f||_{L^2(K)}^2\Big)^{1/2}.$$

It follows from (5.14) that

(5.20)

(5.19) 
$$||u - u_h||_{\mathcal{C}_h} \le \sqrt{\alpha'} (||u - u_H||^2_{\mathcal{C}_H} + \operatorname{osc}^2(f, \mathcal{T}_H))^{1/2}$$

Therefore, we apply the Cauchy-Schwarz inequality to obtain

$$|2a_h(u - u_h, u_h - u_H)| \le \frac{1}{2}(1 - \alpha') \left( ||u - u_H||_{\mathcal{C}_H}^2 + \operatorname{osc}^2(f, \mathcal{T}_H) \right)$$

+ 
$$2(C_{QO})^2 \frac{\alpha'}{1-\alpha'} \sum_{K \in \mathcal{M}_{H,h}} h_K^4 ||f||_{L^2(K)}^2$$
.

Since it is obvious that

(5.21) 
$$|\operatorname{osc}^{2}(f, \mathcal{T}_{H}) - \operatorname{osc}^{2}(f, \mathcal{T}_{h})| \leq \eta^{2}(u_{H}, \mathcal{M}_{H,h}),$$

we combine (5.16)- (5.21) to prove the desired result by the parameter

$$\theta_* = \frac{(1-\alpha')^2 C_{Eff}}{2(2\alpha'(C_{QO})^2 + (1-\alpha')(C_{Drel}+1))}$$

with the efficiency constant  $C_{Eff}$  of the estimator  $\eta(u_H, \mathcal{T}_H)$  from Lemma 2.1.

## 6. Optimality

To analyze the optimality, we follow an idea commonly used in the adaptive finite element literature to introduce a nonlinear approximation class. First, we have the following quasi-optimality.

(6.1) 
$$\|u - u_H\|_{\mathcal{C}_H}^2 \simeq \inf_{v_H \in W_H} \|u - v_H\|_{\mathcal{C}_H}^2 + \kappa^2(u, \mathcal{T}_H),$$

where the consistency error term is given by

(6.2) 
$$\kappa(u, \mathcal{T}_H) = \sup_{v_H \in W_H} \frac{(f, v_H)_{L^2(\Omega)} - a_H(u, v_H)}{\|v_H\|_{\mathcal{C}_H}}$$

It follows from [21, Section 4.1] that

$$\kappa(u, \mathcal{T}_H) \lesssim \inf_{v_H \in W_H} \|u - v_H\|_{\mathcal{C}_H} + \operatorname{osc}(f, \mathcal{T}_H).$$

Therefore, we define

(6.3) 
$$\mathfrak{E}(N; u, f) := \inf_{\mathcal{T} \in \mathbb{T}_N} \inf_{v \in W_{\mathcal{T}}} \left( \|u - v\|_{\mathcal{C}_{\mathcal{T}}}^2 + \operatorname{osc}^2(f, \mathcal{T}) \right).$$

Finally, we choose the nonlinear approximation class as follows:

(6.4) 
$$\mathbb{A}_s := \{(u, f), |u, f|_s := \sup_{N > N_0} N^s \mathfrak{E}(N; u, f) < +\infty \}.$$

Compared to the adaptive conforming method for the second order elliptic problem [14, 30], we have not the following monotone convergence:

(6.5) 
$$\inf_{v_h \in W_h} \|u - v_h\|_{\mathcal{C}_h}^2 + \kappa^2 (u, \mathcal{T}_h)^2 \le \inf_{v_H \in W_H} \|u - v_H\|_{\mathcal{C}_H}^2 + \kappa^2 (u, \mathcal{T}_H),$$

where  $T_h$  is some refinement of  $T_H$ . However, it follows from the quasi-orthogonality in Lemma 3.4, the efficiency of the estimator in Lemma 2.1, and the Young inequality that

(6.6) 
$$\|u - u_h\|_{\mathcal{C}_h}^2 + \operatorname{osc}^2(f, \mathcal{T}_h) \le C_2(\|u - u_H\|_{\mathcal{C}_H}^2 + \operatorname{osc}^2(f, \mathcal{T}_H)).$$

**Theorem 6.1.** Let  $\mathcal{M}_k$  be a set of marked elements with minimal cardinality from Algorithm 4.1, u the solution of Problem (2.3), and  $(\mathcal{T}_k, W_k, u_k)$  the sequence of meshes, finite element spaces, and discrete solutions produced by the adaptive finiteelement methods with  $0 < \theta < \frac{C_{Eff}}{2(2(C_{QO})^2 + C_{Drel} + 1)}$ . Then, the following estimate holds:

(6.7) 
$$\#\mathcal{M}_k \lesssim (\alpha')^{-\frac{1}{s}} |u, f|_s^{\frac{1}{s}} (C_2)^{\frac{1}{s}} (\|u - u_k\|_{\mathcal{C}_k}^2 + \operatorname{osc}^2(f, \mathcal{T}_k))^{-\frac{1}{s}} \text{ for any } (u, f) \in \mathbb{A}_s,$$

where the parameter  $0 < \alpha' < 1$  is from Lemma 5.7.

*Proof.* We set  $\epsilon = \alpha'(C_2)^{-1} (||u - u_k||_{\mathcal{C}_k}^2 + \operatorname{osc}^2(f, \mathcal{T}_k))$  with  $0 < \alpha' < 1$ . Since  $(u, f) \in \mathbb{A}_s$ , there exists a  $\mathcal{T}_{\epsilon}$  of the refinement of  $\mathcal{T}_0$  and  $u_{\epsilon} \in W_{\mathcal{T}_{\epsilon}}$  with

(6.8)  $\#\mathcal{T}_{\epsilon} - \#\mathcal{T}_{0} \leq |u, f|_{s}^{1/s} \epsilon^{-1/s} \text{ and } \|u - u_{\epsilon}\|_{\mathcal{C}_{\mathcal{T}_{\epsilon}}}^{2} + \operatorname{osc}^{2}(f, \mathcal{T}_{\epsilon}) \leq \epsilon.$ 

The overlay  $\mathcal{T}_*$  of  $\mathcal{T}_{\epsilon}$  and  $\mathcal{T}_k$  is the smallest refinement of both  $\mathcal{T}_{\epsilon}$  and  $\mathcal{T}_k$ . Let  $u_*$  be the finite element solution of (2.6) on the mesh  $\mathcal{T}_*$ . Since  $\mathcal{T}_*$  is a refinement of  $\mathcal{T}_{\epsilon}$ , we use, (6.8), and (6.6) to obtain that

(6.9) 
$$\|u - u_*\|_{\mathcal{C}_{\mathcal{T}_*}}^2 + \operatorname{osc}^2(f, \mathcal{T}_*) \le C_2(\|u - u_\epsilon\|_{\mathcal{C}_{\mathcal{T}_\epsilon}}^2 + \operatorname{osc}^2(f, \mathcal{T}_\epsilon)) \\ \le C_2 \epsilon = \alpha'(\|u - u_k\|_{\mathcal{C}_k}^2 + \operatorname{osc}^2(f, \mathcal{T}_k)) .$$

We deduce from Lemma 5.7 that

(6.10) 
$$\theta_*\eta^2(u_k, \mathcal{T}_k) \le \eta^2(u_k, \mathcal{M}_{k,*}), \text{ for some } \theta_* \in (0, 1).$$

We note that the step (3) in Algorithm 4.1 with  $\theta \leq \theta_*$  chooses a subset of  $M_k \subset \mathcal{T}_k$  with minimal cardinality with the same property. Therefore

(6.11) 
$$\#\mathcal{M}_k \lesssim \#\mathcal{T}_* - \#\mathcal{T}_k \leq \#\mathcal{T}_\epsilon - \#\mathcal{T}_0$$

This together with the definition of  $\epsilon$  leads to

(6.12) 
$$\#\mathcal{M}_k \lesssim (\alpha')^{-\frac{1}{s}} |u, f|_s^{\frac{1}{s}} (C_2)^{\frac{1}{s}} (\|u - u_k\|_{\mathcal{C}_k}^2 + \operatorname{osc}^2(f, \mathcal{T}_k))^{-\frac{1}{s}},$$

which completes the proof.

**Theorem 6.2.** Let the marking step in Algorithm 4.1 select a set  $\mathcal{M}_k$  of marked elements with minimal cardinality, u the solution to Problem (2.6), and  $(\mathcal{T}_k, W_k, u_k)$ the sequence of meshes, finite element spaces, and discrete solutions produced by the adaptive finite-element methods with  $0 < \theta < \frac{C_{Eff}}{2(2(C_{OO})^2 + C_{Drel} + 1)}$ . Then, it holds that

(6.13) 
$$||u - u_N||^2_{\mathcal{C}_N} + \operatorname{osc}^2(f, \mathcal{T}_N) \lesssim |u, f|_s (\#\mathcal{T}_N - \#\mathcal{T}_0)^{-s}, \text{ for } (u, f) \in \mathbb{A}_s.$$

*Proof.* Let  $\mu = (\alpha')^{-\frac{1}{s}} |u, f|_s^{\frac{1}{s}} (C_2)^{\frac{1}{s}}$ . We use the result that  $\#\mathcal{T}_k - \#\mathcal{T}_0 \lesssim \sum_{j=0}^{k-1} \mathcal{M}_j$  from [31, 30] to obtain that

(6.14) 
$$\#\mathcal{T}_N - \#\mathcal{T}_0 \lesssim \sum_{j=0}^{N-1} \mathcal{M}_j \lesssim \mu \sum_{j=0}^{N-1} (\|u - u_j\|_{\mathcal{C}_j}^2 + \operatorname{osc}^2(f, \mathcal{T}_j))^{-\frac{1}{s}}.$$

It follows from the efficiency of the estimator that

(6.15) 
$$\|u - u_j\|_{\mathcal{C}_j}^2 + \operatorname{osc}^2(f, \mathcal{T}_j) \cong \tilde{\eta}^2(u_j, \mathcal{T}_j),$$

which gives

(6.16) 
$$\|u - u_j\|_{\mathcal{C}_j}^2 + \gamma_1 \tilde{\eta}^2(u_j, \mathcal{T}_j) \lesssim \|u - u_j\|_{\mathcal{C}_j}^2 + \operatorname{osc}^2(f, \mathcal{T}_j).$$

For any  $0 \leq j \leq N - 1$ , we use the convergence result from Theorem 4.5 to derive that

(6.17) 
$$\|u - u_N\|_{\mathcal{C}_N}^2 + \gamma_1 \tilde{\eta}^2(u_N, \mathcal{T}_N) \le \alpha^{(N-j)} (\|u - u_j\|_{\mathcal{C}_j}^2 + \gamma_1 \tilde{\eta}^2(u_j, \mathcal{T}_j)).$$

A combination of (6.14)-(6.17) yields

(6.18) 
$$\#\mathcal{T}_N - \#\mathcal{T}_0 \lesssim \mu(\|u - u_N\|_{\mathcal{C}_N}^2 + \operatorname{osc}^2(f, \mathcal{T}_N))^{-1/s} \sum_{j=1}^N \alpha^{j/s}.$$

Setting  $C_{\theta} = \alpha^{1/s} (1 - \alpha^{1/s})^{-1}$ , it is easy to prove that

(6.19) 
$$\sum_{j=1}^{N} \alpha^{j/s} \le C_{\theta}$$

Inserting this bound into (6.18) leads to

(6.20) 
$$\|u - u_N\|_{\mathcal{C}_N}^2 + \operatorname{osc}^2(f, \mathcal{T}_N) \lesssim |u, f|_s (\#\mathcal{T}_N - \#\mathcal{T}_0)^{-s}$$

which completes the proof.

## 7. The extensions of the theory

,

This section extends the theory to the Morley element in three dimensions and the nonconforming linear elements in both two and three dimensions.

7.1. The Morley element in three dimensions. Let  $\mathcal{T}_h$  be a decomposition of the domain  $\Omega \subset \mathbb{R}^3$  into simplicies. Given any face F, we let  $\nu_F$  denote its unit normal vector. The Morley element in three dimensions is defined and analyzed in [34], where the space reads

$$W_h := \{ v \in L^2(\Omega), v |_K \in P_2(K), K \in \mathcal{T}_h, \int_e [v] \, ds = 0 \text{ for any internal edge } e, \\ \int_e v \, ds = 0 \text{ for any boundary edge } e, \int_F [\nabla v \cdot \nu_F] dF = 0 \text{ for any} \\ \text{internal face } F, \text{ and } \int_F \nabla v \cdot \nu_F dF = 0 \text{ for any boundary face } F \}.$$

Define the estimator on each element  $K \in \mathcal{T}_h$  as

(7.2) 
$$\eta_K = h_K^2 \|f\|_{L^2(K)} + \left(\sum_{F \subset \partial K} h_F \|[\nabla_h^2 u_h \times \nu_F]\|_{L^2(F)}^2\right)^{1/2},$$

where  $\times$  denotes the usual tensor product. The estimator is defined by

(7.3) 
$$\eta^2(u_h, \mathcal{T}_h) := \sum_{K \in \mathcal{T}_h} \eta_K^2.$$

The following reliability and efficiency of the estimator were proved in [22].

**Lemma 7.1.** Let u be the solution to the fourth order elliptic problem with  $u|_{\partial\Omega} = \frac{\partial u}{\partial \nu}|_{\partial\Omega} = 0$  in three dimensions,  $u_h$  be the finite element solution corresponding to the discrete space  $W_h$  defined in (7.1). Then,

(7.4) 
$$\|u - u_h\|_{\mathcal{C}_h} \cong \eta_h$$

up to the oscillation  $\operatorname{osc}(f, \mathcal{T}_h)$ , where  $\|\cdot\|_{\mathcal{C}_h}$  and  $\operatorname{osc}(f, \mathcal{T}_h)$  are the three dimensional counterparts of the discrete energy norm in (2.5) and the oscillation in (2.10), respectively.

**Lemma 7.2.** Let  $K_1, K_2 \in \mathcal{T}_h$  be two elements sharing a common face F with three edges  $e_{\ell}$  and midpoints  $m_{\ell}$ ,  $\ell = 1, 2, 3$ , and v be a piecewise polynomial of degree  $\leq 1$  over  $K_1 \cup K_2$  such that

(7.5) 
$$v|_{K_1}(m_\ell) = v|_{K_2}(m_\ell), \ell = 1, 2, 3, \text{ and } \int_F [\frac{\partial v}{\partial \nu_F}] dF = 0.$$

Then, v is a polynomial of degree  $\leq 1$  over  $K_1 \cup K_2$ .

With these preparations, one can generalize the theories of the quasi-orthogonality of Lemma 3.4, error reduction of Theorem 4.5, the discrete reliability of Lemma 5.6, and the optimality of Theorem 6.2 to the Morley element method in three dimensions.

7.2. The nonconforming linear elements for second order elliptic problems. In this subsection, we let  $\mathcal{T}_h$  be a decomposition of the domain  $\Omega \subset \mathbb{R}^2$  or  $\Omega \subset \mathbb{R}^3$  into simplicies in both two and three dimensions. The nonconforming linear element spaces in both two and three dimensions is defined by, respectively,

$$W_h := \{ v \in L^2(\Omega), v |_K \in P_1(K), K \in \mathcal{T}_h, \int_e [v] \, ds = 0 \text{ for any internal edge } e, \\ \int_e v \, ds = 0 \text{ for any boundary edge } e \},$$
(7.7)

$$W_h := \{ v \in L^2(\Omega), v |_K \in P_1(K), K \in \mathcal{T}_h, \int_F [v] \, dF = 0 \text{ for any internal face } F, \}$$

$$\int_{F} v \, dF = 0 \text{ for any boundary face } F \}.$$

The continuous problems read: Given  $f \in L^2(\Omega)$ , find  $u \in H^1_0(\Omega)$  such that

(7.8) 
$$(\nabla u, \nabla v)_{L^2(\Omega)} = (f, v)_{L^2(\Omega)} \text{ for any } v \in H^1_0(\Omega)$$

The discrete problems read: Given  $f \in L^2(\Omega)$ , find  $u_h \in W_h$  such that

(7.9) 
$$(\nabla_h u_h, \nabla_h v_h)_{L^2(\Omega)} = (f, v_h)_{L^2(\Omega)} \text{ for any } v_h \in W_h.$$

The convergence of the adaptive nonconforming linear element methods was first analyzed in [13]. The theory in Sections 3-7 can be extended to this case. This extension gives another analysis of the convergence result from [13].

## 8. Conclusion and comments

In this paper, we carry out the convergence and optimality analysis of the Morley element for the fourth order elliptic equation. Moreover, we generalize the theory to the nonconforming linear elements. However, the analysis herein heavily depends on the conservative properties of these two classes of nonconforming elements and the fact that the discrete stress is a piecewise constant tensor. At the present time, it is unclear how to generalize these techniques to other nonconforming schemes of the fourth order elliptic problems.

#### References

- M. Ainsworth and R. Rankin. Robust a posteriori error estimation for the nonconforming Fortin-Soulie finite-element approximation. Math. Comp. 77(2008), pp. 1917–1939.
- [2] I. Babuska and W.C. Rheinboldt. Error estimates for adaptive finite element computations. SIAM. J. Numer. Anal. 15(1978), pp. 736–754.
- [3] L. Beirao da Veiga, J. Niiranen and R. Stenberg. A posteriori error estimates for the Morley plate bending element. Numer. Math. 106(2007), pp. 165–179.
- [4] P. Binev, W. Dahmen, R. DeVore, and P. Petrushev. Approximation classes for adaptive methods. Serdica Math. J. 28(2002), pp. 391–416.
- [5] P. Binev and R. DeVore. Fast computation in adaptive tree approximation. Numer. Math. 97(2004), pp. 193–217.
- [6] P. Binev, W. Dahmen, and R. DeVore. Adaptive finite-element methods with convergence rate. Numer. Math. 97(2004), pp. 219–268.
- [7] S. Brenner and L.R. Scott. The mathematical theory of finite element methods, Springer, New York, 1994.
- [8] S. C. Brenner and L.-Y. Sung. C<sup>0</sup> interior penalty methods for fourth order elliptic boundary value problems on polygonal domains, J. Sci. Comput. 22/23(2005), pp. 83–118.
- [9] C. Carstensen. A unifying theory of a posteriori finite element error control. Numer. Math. 100 (2005), pp. 617–637.
- [10] C. Carstensen, S. Bartels, and S. Jansche. A posteriori error estimates for nonconforming finite-element methods. Numer. Math. 92(2002), pp. 233–256.
- [11] C. Carstensen and J. Hu. A unifying theory of a posteriori error control for nonconforming finite-element methods. Numer. Math. 107(2007), pp. 473–502.
- [12] C. Carstensen, J. Hu and A. Orlando. Framework for the a posteriori error analysis of nonconforming finite elements, SIAM. J. Numer. Anal. 45(2007), pp. 68–82
- [13] C. Carstensen and R. H. W. Hoppe. Convergence analysis of an adaptive nonconforming finite-element method. Numer. Math. 103(2006), pp. 251–266.
- [14] J. Manuel Cascon, C. Kreuzer, R. H. Nochetto, and K. G. Siebert. Quasi-optimal convergence rate for an adaptive finite-element method. SIAM J. Numer. Anal. 46(2008), pp. 2524–2550.
- [15] L. Chen, M. Holst, and J. Xu. Convergence and optimality of adaptive mixed finite-element methods. Math. Comp. 78(2008), pp. 35–53.
- [16] P.G. Ciarlet. The Finite Element Method for Elliptic Problems. North-Holland, 1978; reprinted as SIAM Classics in Applied Mathematics, 2002.

- [17] P. Clément. Approximation by finite-element functions using local regularization. RAIRO Anal. Numér. 9(1975), pp. 77–84.
- [18] E. Dari, R. Duran, C. Padra. Error estimators for nonconforming finite-element approximations of the Stokes problem. Math. Comp. 64 (1995), pp. 1017–1033.
- [19] E. Dari, R. Duran, C. Padra, and V. Vampa. A posteriori error estimators for nonconforming finite-element methods. Math. Model. Numer. Anal. 30 (1996), pp. 385–400.
- [20] W. Dörfler. A convergent adaptive algorithm for Poisson's equation. SIAM J. Numer. Anal. 33(1996), pp. 1106–1124.
- [21] T. Gudi. A new error analysis for discontinous finite element methods for linear problems, Math. Comp. 79(2010), pp. 2169–2189.
- [22] J. Hu and Z. C. Shi. A new a posteriori error estimate for the Morley element. Numer. Math. 112(2009), pp. 25–40.
- [23] J. Hu, Z. C. Shi and J. C. Xu. Convergence and optimality of adaptive nonconforming methods for high-order differential equations, Research Report 19 (2009), School of Mathematical Sciences and Institute of Mathematics, Peking University.
- [24] J. Hu and J. C. Xu. Convergence of Adaptive Conforming and Nonconforming Finite Element Methods for the Perturbed Stokes Equation, Research Report 73(2007), School of Mathematical Sciences and Institute of Mathematics, Peking University. (Also available online: http://www.math.pku.edu.cn:8000/var/preprint/7297.pdf.)
- [25] L. Marini. An inexpensive method for the evaluation of the solution of the lowest order Raviart-Thomas mixed method. SIAM J. Numer. Anal. 22(1985), pp. 493–496.
- [26] L. S. D. Morley. The triangular equilibrium element in the solutions of plate bending problem. Aero.Quart.19(1968), pp 149–169.
- [27] P. Morin, R. Nochetto, and K. Siebert. Data oscillation and convergence of adaptive FEM. SIAM J. Numer. Anal. 38(2000), pp. 466–488.
- [28] P. Morin, R. H. Nochetto, and K. G. Siebert. Convergence of adaptive finite-element methods. SIAM Rev. 44(2002), pp. 631–658.
- [29] Z. C. Shi. Error estimates for the Morley element. Chin. J. Numer. Math. Appl. 12(1990), pp. 102–108.
- [30] R. Stevenson. Optimality of a standard adaptive finite element method. Found. Comput. Math. 7(2007), pp. 245–269.
- [31] R. Stevenson. The completion of locally refined simplicial partitions created by bisection. Math.Comp. 77(2007), pp. 227–241.
- [32] R. Verfürth. A review of a posteriori error estimation and adaptive mesh-refinement technique. Wiley-Teubner, New York, 1996.
- [33] M. Wang and S. Zhang. Local a priori and a posteriori error estimates of finite-elements for biharmonic equation. Research Report 13(2006), School of Mathematical Sciences and Institute of Mathematics, Peking University.
- [34] M. Wang and J. C. Xu. Minimal finite-element spaces for 2m-th order partial differential equations in  $\mathbb{R}^n$ . Research Report 29(2006), School of Mathematical Sciences and Institute of Mathematics, Peking University(submitted to Mathematics of Computation).
- [35] J. C. Xu. Iterative methods by space secomposition and subspace correction, SIAM Review, Vol. 34(1992), pp. 581–613.

CONVERGENCE AND OPTIMALITY OF ADAPTIVE NONCONFORMING METHODS 21

LMAM and School of Mathematical Sciences, Peking University, Beijing 100871, P. R. China

E-mail address: hujun@math.pku.edu.cn

INSTITUTE OF COMPUTATIONAL MATHEMATICS, CHINESE ACADEMY OF SCIENCES, BEIJING 100080, P. R. CHINA.

 $E\text{-}mail\ address:\ \mathtt{shi@lsec.cc.ac.cn}$ 

THE SCHOOL OF MATHEMATICAL SCIENCES, PEKING UNIVERSITY, AND DEPARTMENT OF MATHEMATICS, PENNSYLVANIA STATE UNIVERSITY, UNIVERSITY PARK, PA 16801 *E-mail address*: xu@math.psu.edu