# Uniform-in-time convergence of numerical methods for non-linear degenerate parabolic equations

Jérôme Droniou\* and Robert Eymard<sup>†</sup> March 23, 2020

#### Abstract

Gradient schemes is a framework that enables the unified convergence analysis of many numerical methods for elliptic and parabolic partial differential equations: conforming and non-conforming Finite Element, Mixed Finite Element and Finite Volume methods. We show here that this framework can be applied to a family of degenerate non-linear parabolic equations (which contain in particular the Richards', Stefan's and Leray-Lions' models), and we prove a uniform-in-time strong-in-space convergence result for the gradient scheme approximations of these equations. In order to establish this convergence, we develop several discrete compactness tools for numerical approximations of parabolic models, including a discontinuous Ascoli-Arzelà theorem and a uniform-in-time weak-in-space discrete Aubin-Simon theorem. The model's degeneracies, which occur both in the time and space derivatives, also requires us to develop a discrete compensated compactness result.

AMS Subject Classification: 65M12, 35K65, 46N40.

**Keywords**: gradient schemes, convergence analysis, degenerate parabolic equations, uniform convergence, discontinuous Ascoli-Arzelà theorem, discrete Aubin-Simon theorem, compensated compactness.

## 1 Introduction

## 1.1 Motivation

The following generic nonlinear parabolic model

$$\partial_{t}\beta(\overline{u}) - \operatorname{div}\left(\boldsymbol{a}(\boldsymbol{x},\nu(\overline{u}),\nabla\zeta(\overline{u}))\right) = f \quad \text{in } \Omega \times (0,T), 
\beta(\overline{u})(\boldsymbol{x},0) = \beta(u_{\text{ini}})(\boldsymbol{x}) \quad \text{in } \Omega, 
\zeta(\overline{u}) = 0 \quad \text{on } \partial\Omega \times (0,T),$$
(1)

where  $\beta$  and  $\zeta$  are non-decreasing,  $\nu$  is such that  $\nu' = \beta' \zeta'$  and a is a Leray-Lions operator, arises in various frameworks (see next section for precise hypotheses on the data). This model includes

1. The Richards model, setting  $\zeta(s) = s$ ,  $\nu = \beta$  and  $a(x, \nu(\overline{u}), \nabla \zeta(\overline{u})) = K(x, \beta(\overline{u})) \nabla \overline{u}$ , which describes the flow of water in a heterogeneous anisotropic underground medium,

<sup>\*</sup>School of Mathematical Sciences, Monash University, Victoria 3800, Australia. jerome.droniou@monash.edu.

†Université Paris-Est, Laboratoire d'Analyse et de Mathématiques Appliquées, UMR 8050, 5 boulevard Descartes,
Champs-sur-Marne 77454 Marne-la-Vallée Cedex 2, France. Robert.Eymard@univ-mlv.fr.

- 2. The Stefan model [8], setting  $\beta(s) = s$ ,  $\nu = \zeta$ ,  $a(x, \nu(\overline{u}), \nabla \zeta(\overline{u})) = K(x, \zeta(\overline{u})) \nabla \zeta(\overline{u})$ , which arises in the study of a simplified heat diffusion process in a melting medium,
- 3. The *p*-Laplace problem, setting  $\beta(s) = \zeta(s) = \nu(s) = s$  and  $\mathbf{a}(\mathbf{x}, \nu(\overline{u}), \nabla \zeta(\overline{u})) = |\nabla \overline{u}|^{p-2} \nabla \overline{u}$ , which is involved in the motion of glaciers [37] or flows of incompressible turbulent fluids through porous media [16].

General Leray–Lions operators  $a(x, s, \xi)$  have growth, monotony and coercivity properties (see (2f)–(2h) below) which ensure that  $-\text{div}(a(x, w, \nabla \cdot))$  maps  $W_0^{1,p}(\Omega)$  into  $W^{-1,p'}(\Omega)$ , and thanks to which this differential operator is viewed as a generalisation of the p-Laplace operator.

The numerical approximation of these models has been extensively studied in the literature – see the fundamental work on the Stefan's problem [48] and [51, 30] for some of its numerical approximations, [46, 33] for the Richards' problem, and [19, 23] and references therein for some studies of convergence of numerical methods for the Leray-Lions' problem. In [52], fully discrete implicit schemes are considered in 2D domains for the problem  $\partial_t e - \Delta u = f$ ,  $e \in \beta(u)$  with  $\beta$  a maximal monotone operator; error estimates are obtained and the results are relevant, e.g., for the Stefan problem and the porous medium equation.

More generally, studies have been carried out on numerical time-stepping approximations of non-linear abstract parabolic equations. In [43] the authors study the stability and convergence properties of linearised implicit methods for the time discretization of nonlinear parabolic equations in the general framework of Hilbert spaces. The time discretisation of nonlinear evolution equations in an abstract Banach space setting of analytic semigroups is studied in [38]; this setting covers fully nonlinear parabolic initial-boundary value problems with smooth coefficients. [3] deals with a general formulation for semi-discretisations of linear parabolic evolution problems in Hilbert spaces; this time-stepping formulation encompasses continuous and discontinuous Galerkin methods, as well as Runge Kutta methods. The study in [3] has been extended in [2] to semi-linear equations, i.e. with the addition of a right-hand side which is locally Lipschitz-continuous with respect to the unknown. In the same directions, we also quote [42, 44, 45, 49, 39] for Runge-Kutta time discretizations of linear and quasilinear parabolic equations (reaction-diffusion, Navier-Stokes equations, etc.). Multisteps methods have also been considered, see e.g. [50].

However, most of these studies are only applicable under regularity assumptions on the solution or data, and to semi-linear equations or semi-discretised schemes. None deals with as many non-linearities and degeneracies as in (1). Moreover, the results in these works mostly yield space-time averaged convergences, e.g. in  $L^2(\Omega \times (0,T))$ . Yet, the quantity of interest is often not  $\overline{u}$  on  $\Omega \times (0,T)$  but  $\overline{u}$  at a given time, for example t=T. Current numerical analyses therefore do not ensure that this quantity of interest is properly approximated by numerical methods.

The usual way to obtain pointwise-in-time approximation results for numerical schemes is to prove estimates in  $L^{\infty}(0,T;L^2(\Omega))$  on  $u-\overline{u}$ , where u is the approximated solution. Establishing such error estimates is however only feasible when uniqueness of the solution  $\overline{u}$  to (1) can be proved, which is the case for Richards' and Stefan's problems (with K only depending on x), but not for more complex non-linear parabolic problems as (1) or even p-Laplace problems. It moreover requires some regularity assumptions on  $\overline{u}$ , which clearly fail to hold for (1) (and simpler p-Laplace problems); indeed, because of the possible plateaux of  $\beta$  and  $\zeta$ , the solution's gradient can develop jumps.

The purpose of this article is to prove that, using Discrete Functional Analysis techniques (i.e. the translation to numerical analysis of nonlinear analysis techniques), an  $L^{\infty}(0,T;L^2(\Omega))$  convergence result can be established for numerical approximations of (1), without having to assume non-physical regularity assumptions on the data. Note that, although Richards' and Stefan's models are formally equivalent when  $\beta$  and  $\zeta$  are strictly increasing (consider  $\beta = \zeta^{-1}$  to pass from

one model to the other), they change nature when these functions are allowed to have plateaux. Stefan's model can degenerate to an ODE (if  $\zeta$  is constant on the range of the solution) and Richards' model can become a non-transient elliptic equation (if  $\beta$  is constant on this range). The innovative technique we develop in this paper is nonetheless generic enough to work directly on (1) and with a vast number of numerical methods.

That being said, a particular numerical framework must be selected to write precise equations and estimates. The framework we choose is that of gradient schemes, which has the double benefit of covering a vast number of numerical methods, and of having already been studied for many models – elliptic, parabolic, linear or non-linear, possibly degenerate, etc. – with various boundary conditions. The schemes or family of schemes included in the gradient schemes framework, and to which our results therefore directly apply, currently are:

- Galerkin methods, including conforming Finite Element schemes,
- finite element with mass lumping [12],
- the Crouzeix-Raviart non-conforming finite element, with or without mass lumping [14, 27],
- the Raviart-Thomas mixed finite elements [9],
- the vertex approximate gradient scheme [31],
- the hybrid mimetic mixed family [22], which includes mimetic finite differences [10], mixed finite volume [20] and the SUSHI scheme [29],
- the discrete duality finite volume scheme in dimension 2 [40, 5], and the CeVeFE-discrete duality finite volume scheme in dimension 3 [13],
- the multi-point flux approximation O-method [1, 25].

We refer the reader to [21, 23, 28, 34, 32] for more details. Let us finally emphasize that the unified convergence study of numerical schemes for Problem (1), which combines a general Leray–Lions operator and nonlinear functions  $\beta$  or  $\zeta$ , seems to be new even without the uniform-in-time convergence result.

The paper is organised as follows. In Section 1.2, we present the assumptions and the notion of weak solution for (1) and, in Section 1.3, we give an overview of the ideas involved in the proof of uniform-in-time convergence. This overview is given not in a numerical analysis context but in the context of a pure stability analysis of (1) with very little regularity on the data, for which the uniform-in-time convergence result also seems to be new. Section 2 presents the gradient schemes for our generic model (1). We give in Section 3 some preliminaries to the convergence study, in particular a crucial uniform-in-time weak-in-space discrete Aubin-Simon compactness result. Section 4 contains the complete convergence proof of gradient schemes for (1), including the uniform-intime convergence result. This proof is initially conducted under a simplifying assumption on  $\beta$ and  $\zeta$ . We demonstrate in Section 5 that, in the case  $p \geq 2$ , this assumption can be removed thanks to a discrete compensated compactness result. We also remark in this section that our results apply to the model considered in [52]. An appendix, Section 6, concludes the article with technical results, in particular a generalisation of the Ascoli-Arzelà compactness result to discontinuous functions and a characterisation of the uniform convergence of a sequence of functions; these results are critical to establishing our uniform-in-time convergence result. We believe that the Discrete Functional Analysis results we establish in order to study the approximations of (1) - in particular the discrete compensated compactness theorem (Theorem 5.4) - could be critical to the numerical analysis of other degenerate or coupled models of physical importance.

Note that the main results and their proofs have been sketched and illustrated by some numerical examples in [24], for  $a(x, \nu(\overline{u}), \nabla \zeta(\overline{u})) = \nabla \zeta(\overline{u})$  and  $\beta = \text{Id}$  or  $\zeta = \text{Id}$ .

### 1.2 Hypotheses and weak sense for the continuous problem

We consider the evolution problem (1) under the following hypotheses.

$$\Omega$$
 is an open bounded subset of  $\mathbb{R}^d$   $(d \in \mathbb{N}^*)$  and  $T > 0$ , (2a)

$$\zeta \in C^0(\mathbb{R})$$
 is non–decreasing, Lipschitz continuous with Lipschitz constant  $L_{\zeta} > 0$ ,  $\zeta(0) = 0$  and, for some  $M_0, M_1 > 0$ ,  $|\zeta(s)| \ge M_0 |s| - M_1$  for all  $s \in \mathbb{R}$ . (2b)

$$\beta \in C^0(\mathbb{R})$$
 is non-decreasing, Lipschitz continuous with Lipschitz constant  $L_{\beta} > 0$ , and  $\beta(0) = 0$ .

$$\forall s \in \mathbb{R}, \quad \nu(s) = \int_0^s \zeta'(q)\beta'(q)dq. \tag{2d}$$

$$\mathbf{a}: \Omega \times \mathbb{R} \times \mathbb{R}^d \to \mathbb{R}^d$$
 is a Carathéodory function (2e)

(i.e. a function such that, for a.e.  $x \in \Omega$ ,  $(s, \xi) \mapsto a(x, s, \xi)$  is continuous and, for any  $(s, \xi) \in \mathbb{R} \times \mathbb{R}^d$ ,  $x \mapsto a(x, s, \xi)$  is measurable) and, for some  $p \in (1, +\infty)$ ,

$$\exists \underline{a} \in (0, +\infty) : \mathbf{a}(\mathbf{x}, s, \boldsymbol{\xi}) \cdot \boldsymbol{\xi} \ge \underline{a} |\boldsymbol{\xi}|^p, \text{ for a.e. } \boldsymbol{x} \in \Omega, \ \forall s \in \mathbb{R}, \ \forall \boldsymbol{\xi} \in \mathbb{R}^d,$$
 (2f)

$$(\boldsymbol{a}(\boldsymbol{x}, s, \boldsymbol{\xi}) - \boldsymbol{a}(\boldsymbol{x}, s, \boldsymbol{\chi})) \cdot (\boldsymbol{\xi} - \boldsymbol{\chi}) \ge 0$$
, for a.e.  $\boldsymbol{x} \in \Omega$ ,  $\forall s \in \mathbb{R}$ ,  $\forall \boldsymbol{\xi}, \boldsymbol{\chi} \in \mathbb{R}^d$ , (2g)

$$\exists \overline{a} \in L^{p'}(\Omega), \exists \mu \in (0, +\infty) : 
|\boldsymbol{a}(\boldsymbol{x}, s, \boldsymbol{\xi})| \leq \overline{a}(\boldsymbol{x}) + \mu |\boldsymbol{\xi}|^{p-1}, \text{ for a.e. } \boldsymbol{x} \in \Omega, \ \forall s \in \mathbb{R}, \ \forall \boldsymbol{\xi} \in \mathbb{R}^d.$$
(2h)

We also assume, setting  $p' = \frac{p}{p-1}$  the dual exponent of the p previously introduced,

$$u_{\text{ini}} \in L^2(\Omega), \quad f \in L^{p'}(\Omega \times (0,T)).$$
 (2i)

We denote by  $R_{\beta}$  the range of  $\beta$  and define the pseudo-inverse function  $\beta_r: R_{\beta} \to \mathbb{R}$  of  $\beta$  by

$$\forall s \in R_{\beta}, \ \beta_{r}(s) = \begin{cases} \inf\{t \in \mathbb{R} \mid \beta(t) = s\} & \text{if } s > 0, \\ 0 & \text{if } s = 0, \\ \sup\{t \in \mathbb{R} \mid \beta(t) = s\} & \text{if } s < 0, \end{cases}$$

$$= \text{closest } t \text{ to } 0 \text{ such that } \beta(t) = s.$$
(3)

Since  $\beta(t)$  has the same sign as t, we have  $\beta_r \geq 0$  on  $R_\beta \cap \mathbb{R}^+$  and  $\beta_r \leq 0$  on  $R_\beta \cap \mathbb{R}^-$ . We then define  $B: R_\beta \to [0, \infty]$  by

$$B(z) = \int_0^z \zeta(\beta_r(s)) \, ds.$$

Since  $\beta_r$  is non-decreasing, this expression is always well-defined in  $[0,\infty)$ . The signs of  $\beta_r$  and  $\zeta$  ensure that B is non-decreasing on  $R_{\beta} \cap \mathbb{R}^+$  and non-increasing on  $R_{\beta} \cap \mathbb{R}^-$ , and therefore has limits (possibly  $+\infty$ ) at the endpoints of  $R_{\beta}$ . We can thus extend B as a function defined on  $\overline{R_{\beta}}$  with values in  $[0,+\infty]$ .

The precise notion of solution to (1) that we consider is the following:

$$\begin{cases}
\overline{u} \in L^{p}(0, T; L^{p}(\Omega)), \ \zeta(\overline{u}) \in L^{p}(0, T; W_{0}^{1,p}(\Omega)), \\
B(\beta(\overline{u})) \in L^{\infty}(0, T; L^{1}(\Omega)), \ \beta(\overline{u}) \in C([0, T]; L^{2}(\Omega) - \mathbf{w}), \partial_{t}\beta(\overline{u}) \in L^{p'}(0, T; W^{-1,p'}(\Omega)), \\
\beta(\overline{u})(\cdot, 0) = \beta(u_{\text{ini}}) \text{ in } L^{2}(\Omega), \\
\int_{0}^{T} \langle \partial_{t}\beta(\overline{u})(\cdot, t), \overline{v}(\cdot, t) \rangle_{W^{-1,p'}, W_{0}^{1,p}} dt \\
+ \int_{0}^{T} \int_{\Omega} a(x, \nu(\overline{u}(x, t)), \nabla \zeta(\overline{u})(x, t)) \cdot \nabla \overline{v}(x, t) dx dt = \int_{0}^{T} \int_{\Omega} f(x, t) \overline{v}(x, t) dx dt, \\
\forall \overline{v} \in L^{p}(0; T; W_{0}^{1,p}(\Omega)).
\end{cases} \tag{4}$$

where  $C([0,T];L^2(\Omega)-w)$  denotes the space of continuous functions  $[0,T] \mapsto L^2(\Omega)$  for the weak-\* topology of  $L^2(\Omega)$ . Here and in the following, we remove the mention of  $\Omega$  in the duality bracket  $\langle \cdot, \cdot \rangle_{W^{-1,p'},W_0^{1,p}} = \langle \cdot, \cdot \rangle_{W^{-1,p'}(\Omega),W_0^{1,p}(\Omega)}$ .

Remark 1.1 The derivative  $\partial_t \beta(\overline{u})$  is to be understood in the usual sense of distributions on  $\Omega \times (0,T)$ . Since the set  $\mathcal{T} = \{\sum_{i=1}^q \varphi_i(t)\gamma_i(\boldsymbol{x}) : q \in \mathbb{N}, \varphi_i \in C_c^{\infty}(0,T), \gamma_i \in C_c^{\infty}(\Omega)\}$  of tensorial functions in  $C_c^{\infty}(\Omega \times (0,T))$  is dense in  $L^p(0,T;W_0^{1,p}(\Omega))$ , one can ensure that this distribution derivative  $\partial_t \beta(\overline{u})$  belongs to  $L^{p'}(0,T;W^{-1,p'}(\Omega)) = (L^p(0,T;W_0^{1,p}(\Omega)))'$  by checking that the linear form

$$\varphi \in \mathcal{T} \mapsto \langle \partial_t \beta(\overline{u}), \varphi \rangle_{\mathcal{D}', \mathcal{D}} = -\int_0^T \int_{\Omega} \beta(\overline{u})(\boldsymbol{x}, t) \partial_t \varphi(\boldsymbol{x}, t) d\boldsymbol{x} dt$$

is continuous for the norm of  $L^p(0,T;W_0^{1,p}(\Omega))$ .

Note that the continuity property of  $\beta(\overline{u})$  in (4) is natural. Indeed, since  $\beta(\overline{u}) \in L^{\infty}(0,T;L^{2}(\Omega))$  (this comes from  $B(\beta(\overline{u})) \in L^{\infty}(0,T;L^{1}(\Omega))$  and (26)), the PDE in the sense of distributions shows that for any  $\varphi \in C_{c}^{\infty}(\Omega)$  the mapping  $T_{\varphi}: t \mapsto \langle \beta(\overline{u})(t), \varphi \rangle_{L^{2}}$  belongs to  $W^{1,1}(0,T) \subset C([0,T])$ . By density of  $C_{c}^{\infty}(\Omega)$  in  $L^{2}(\Omega)$  and the integrability properties of  $\beta(\overline{u})$ , we deduce that  $T_{\varphi} \in C([0,T])$  for any  $\varphi \in L^{2}(\Omega)$ , which precisely establishes the continuity of  $\beta(\overline{u}): [0,T] \to L^{2}(\Omega)$ -w. This notion of  $\beta(\overline{u})$  as a function continuous in time is nevertheless a subtle one. It is to be understood in the sense that the function  $(x,t) \mapsto \beta(\overline{u}(x,t))$  has an a.e. representative which is continuous  $[0,T] \mapsto L^{2}(\Omega)$ -w. In other words, there is a function  $Z \in C([0,T];L^{2}(\Omega)$ -w) such that  $Z(t)(x) = \beta(\overline{u}(x,t))$  for a.e.  $(x,t) \in \Omega \times (0,T)$ . We must however make sure, when dealing with pointwise value in time, to separate Z from  $\beta(\overline{u}(\cdot,\cdot))$  as  $\beta(\overline{u}(\cdot,t_{1}))$  may not make sense for a particular  $t_{1} \in [0,T]$ . That being said, in order to adopt a simple notation, we will denote by  $\beta(\overline{u})(\cdot,\cdot)$  the function Z, and by  $\beta(\overline{u})(\cdot,t)$  for a particular  $t_{1} \in [0,T]$ , and we will only write  $\beta(\overline{u})(x,t) = \beta(\overline{u}(x,t))$  for a.e.  $(x,t) \in \Omega \times (0,T)$ . Note that from this a.e. equality we can ensure

#### 1.3 General ideas for the uniform-in-time convergence result

that  $\beta(\overline{u})(\cdot,\cdot)$  takes its values in the closure  $\overline{R_{\beta}}$  of the range of  $\beta$ .

As explained in the introduction, the main innovative result of this article is the uniform-intime convergence result (Theorem 2.16 below). Although it's stated and proved in the context of numerical approximations of (1), we emphasize that the ideas underlying its proof are also applicable to theoretical analysis of PDEs. Let us informally present these ideas on the following continuous approximation of (1):

$$\partial_{t}\beta(\overline{u}_{\varepsilon}) - \operatorname{div}\left(\boldsymbol{a}_{\varepsilon}(\boldsymbol{x}, \nu(\overline{u}_{\varepsilon}), \nabla\zeta(\overline{u}_{\varepsilon}))\right) = f \quad \text{in } \Omega \times (0, T), 
\beta(\overline{u}_{\varepsilon})(\boldsymbol{x}, 0) = \beta(u_{\text{ini}})(\boldsymbol{x}) \quad \text{in } \Omega, 
\zeta(\overline{u}_{\varepsilon}) = 0 \quad \text{on } \partial\Omega \times (0, T)$$
(5)

where  $a_{\varepsilon}$  satisfies Assumptions (2e)–(2h) with constants not depending on  $\varepsilon$  and, as  $\varepsilon \to 0$ ,  $a_{\varepsilon} \to a$  locally uniformly with respect to  $(s, \xi)$ .

We want to show here how to deduce from averaged convergences a strong uniform-in-time convergence result. We therefore assume the following convergences (up to a subsequence as  $\varepsilon \to 0$ ), which are compatible with basic compactness results that can be obtained on  $(\overline{u}_{\varepsilon})_{\varepsilon}$  and also correspond to the initial convergences (18) on numerical approximations of (1):

$$\beta(\overline{u}_{\varepsilon}) \to \beta(\overline{u}) \text{ in } C([0,T]; L^{2}(\Omega)\text{-w}), \ \nu(\overline{u}_{\varepsilon}) \to \nu(\overline{u}) \text{ strongly in } L^{1}(\Omega \times (0,T)),$$

$$\zeta(\overline{u}_{\varepsilon}) \to \zeta(\overline{u}) \text{ weakly in } L^{p}(0,T; W_{0}^{1,p}(\Omega)),$$

$$\boldsymbol{a}_{\varepsilon}(\cdot, \nu(\overline{u}_{\varepsilon}), \nabla\zeta(\overline{u}_{\varepsilon})) \to \boldsymbol{a}(\cdot, \nu(\overline{u}), \nabla\zeta(\overline{u})) \text{ weakly in } L^{p'}(\Omega \times (0,T))^{d}.$$

$$(6)$$

We will prove from these convergences that, along the same subsequence,  $\nu(\overline{u}_{\varepsilon}) \to \nu(\overline{u})$  strongly in  $C([0,T]; L^2(\Omega))$ , which is our uniform-in-time convergence result.

We start by noticing that the weak-in-space uniform-in-time convergence of  $\beta(\overline{u}_{\varepsilon})$  gives, for any  $T_0 \in [0,T]$  and any family  $(T_{\varepsilon})_{\varepsilon>0}$  converging to  $T_0$  as  $\varepsilon \to 0$ ,  $\beta(\overline{u}_{\varepsilon})(T_{\varepsilon},\cdot) \to \beta(\overline{u})(T_0,\cdot)$  weakly in  $L^2(\Omega)$ . Classical strong-weak semi-continuity properties of convex functions (see Lemma 3.4) and the convexity of B (see Lemma 3.3) then ensure that

$$\int_{\Omega} B(\beta(\overline{u})(\boldsymbol{x}, T_0)) d\boldsymbol{x} \le \liminf_{\varepsilon \to 0} \int_{\Omega} B(\beta(\overline{u}_{\varepsilon})(\boldsymbol{x}, T_{\varepsilon})) d\boldsymbol{x}. \tag{7}$$

The second step is to notice that, by (2g) for  $a_{\varepsilon}$ ,

$$\int_0^{T_{\varepsilon}} \int_{\Omega} \left[ \boldsymbol{a}_{\varepsilon}(\cdot, \nu(\overline{u}_{\varepsilon}), \nabla \zeta(\overline{u}_{\varepsilon})) - \boldsymbol{a}_{\varepsilon}(\cdot, \nu(\overline{u}_{\varepsilon}), \nabla \zeta(\overline{u})) \right] \cdot \left[ \nabla \zeta(\overline{u}_{\varepsilon}) - \nabla \zeta(\overline{u}) \right] d\boldsymbol{x} dt \geq 0.$$

Developing this expression and using the convergences (6), we find that

$$\liminf_{\varepsilon \to 0} \int_{0}^{T_{\varepsilon}} \int_{\Omega} \boldsymbol{a}_{\varepsilon}(\cdot, \nu(\overline{u}_{\varepsilon}), \nabla \zeta(\overline{u}_{\varepsilon})) \cdot \nabla \zeta(\overline{u}_{\varepsilon})(\boldsymbol{x}, t) d\boldsymbol{x} dt \ge \int_{0}^{T_{0}} \int_{\Omega} \boldsymbol{a}(\cdot, \nu(\overline{u}), \nabla \zeta(\overline{u})) \cdot \nabla \zeta(\overline{u}) d\boldsymbol{x} dt. \tag{8}$$

We then establish the following formula:

$$\int_{\Omega} B(\beta(\overline{u}_{\varepsilon}(\boldsymbol{x}, T_{\varepsilon}))) d\boldsymbol{x} + \int_{0}^{T_{\varepsilon}} \int_{\Omega} \boldsymbol{a}_{\varepsilon}(\boldsymbol{x}, \nu(\overline{u}_{\varepsilon}(\boldsymbol{x}, t)), \nabla \zeta(\overline{u}_{\varepsilon})(\boldsymbol{x}, t)) \cdot \nabla \zeta(\overline{u}_{\varepsilon})(\boldsymbol{x}, t) d\boldsymbol{x} dt 
= \int_{\Omega} B(\beta(u_{\text{ini}}(\boldsymbol{x}))) d\boldsymbol{x} + \int_{0}^{T_{\varepsilon}} \int_{\Omega} f(\boldsymbol{x}, t) \zeta(\overline{u}_{\varepsilon})(\boldsymbol{x}, t) d\boldsymbol{x} dt. \quad (9)$$

This energy equation is formally obtained by multiplying (5) by  $\zeta(\overline{u}_{\varepsilon})$  and integrating by parts, using  $(B \circ \beta)' = \zeta \beta'$  (see Lemma 3.3); the rigorous justification of (9) is however quite technical – see Lemma 3.6 and Corollary 3.8. Thanks to (8), we can pass to the lim sup in (9) and we find, using the same energy equality with  $(\overline{u}, \boldsymbol{a}, T_0)$  instead of  $(\overline{u}_{\varepsilon}, \boldsymbol{a}_{\varepsilon}, T_{\varepsilon})$ ,

$$\limsup_{\varepsilon \to 0} \int_{\Omega} B(\beta(\overline{u}_{\varepsilon}(\boldsymbol{x}, T_{\varepsilon}))) d\boldsymbol{x} \le \int_{\Omega} B(\beta(\overline{u}(\boldsymbol{x}, T_{0}))) d\boldsymbol{x}.$$
(10)

Combined with (7), this shows that  $\int_{\Omega} B(\beta(\overline{u}_{\varepsilon}(\boldsymbol{x},T_{\varepsilon}))) d\boldsymbol{x} \to \int_{\Omega} B(\beta(\overline{u}(\boldsymbol{x},T_{0}))) d\boldsymbol{x}$ . A uniform convexity property of B (see (28)) then allows us to deduce that  $\nu(\overline{u}_{\varepsilon}(\cdot,T_{\varepsilon})) \to \nu(\overline{u}(\cdot,T_{0}))$  strongly in  $L^{2}(\Omega)$  and thus that  $\nu(\overline{u}_{\varepsilon}) \to \nu(\overline{u})$  strongly in  $C([0,T];L^{2}(\Omega))$  (see Lemma 6.4).

Remark 1.2 A close examination of this proof indicates that equality in the energy relation (9) is not required for  $\overline{u}_{\varepsilon}$ . An inequality  $\leq$  would be sufficient. This is particularly important in the context of numerical methods which may introduce additional numerical diffusion (for example due to an implicit-in-time discretisation) and therefore only provide an upper bound in this energy estimate, see (42). It is however essential that the limit solution  $\overline{u}$  satisfies the equivalent of (9) with an equal sign (or  $\geq$ ).

## 2 Gradient discretisations and gradient schemes

#### 2.1 Definitions

We give here a minimal presentation of gradient discretisations and gradient schemes, limiting ourselves to what is necessary to study the discretisation of (1). We refer the reader to [21, 31, 23] for more details.

A gradient scheme can be viewed as a general formulation of several discretisations of (1), that are based on a nonconforming approximation of the weak formulation of the problem. This approximation is constructed by using discrete space and mappings, the set of which are called a gradient discretisation.

# Definition 2.1 (Space-Time gradient discretisation for homogeneous Dirichlet boundary conditions)

We say that  $\mathcal{D} = (X_{\mathcal{D},0}, \Pi_{\mathcal{D}}, \nabla_{\mathcal{D}}, \mathcal{I}_{\mathcal{D}}, (t^{(n)})_{n=0,...,N})$  is a space-time gradient discretisation for homogeneous Dirichlet boundary conditions if

- 1. the set of discrete unknowns  $X_{\mathcal{D},0}$  is a finite dimensional real vector space,
- 2. the linear mapping  $\Pi_{\mathcal{D}}: X_{\mathcal{D},0} \to L^{\infty}(\Omega)$  is a piecewise constant reconstruction operator in the following sense: there exists a set I of degrees of freedom and a family  $(\Omega_i)_{i\in I}$  of disjoint subsets of  $\Omega$  such that  $X_{\mathcal{D},0} = \mathbb{R}^I$ ,  $\Omega = \bigcup_{i\in I} \Omega_i$  and, for all  $u = (u_i)_{i\in I} \in X_{\mathcal{D},0}$  and all  $i\in I$ ,  $\Pi_{\mathcal{D}}u = u_i$  on  $\Omega_i$ ,
- 3. the linear mapping  $\nabla_{\mathcal{D}}: X_{\mathcal{D},0} \to L^p(\Omega)^d$  gives a reconstructed discrete gradient. It must be chosen such that  $\|\nabla_{\mathcal{D}}\cdot\|_{L^p(\Omega)^d}$  is a norm on  $X_{\mathcal{D},0}$ ,
- 4.  $\mathcal{I}_{\mathcal{D}}: L^2(\Omega) \to X_{\mathcal{D},0}$  is a linear interpolation operator,
- 5.  $t^{(0)} = 0 < t^{(1)} < t^{(2)} < \ldots < t^{(N)} = T$ .

We then set  $\delta t^{(n+\frac{1}{2})} = t^{(n+1)} - t^{(n)}$  for  $n = 0, \dots, N-1$ , and  $\delta t_{\mathcal{D}} = \max_{n=0,\dots,N-1} \delta t^{(n+\frac{1}{2})}$ . We define the dual semi-norm  $|w|_{\star,\mathcal{D}}$  of  $w \in X_{\mathcal{D},0}$  by

$$|w|_{\star,\mathcal{D}} = \sup \left\{ \int_{\Omega} \Pi_{\mathcal{D}} w(\boldsymbol{x}) \Pi_{\mathcal{D}} z(\boldsymbol{x}) d\boldsymbol{x} : z \in X_{\mathcal{D},0}, ||\nabla_{\mathcal{D}} z||_{L^{p}(\Omega)^{d}} = 1 \right\}.$$
(11)

Remark 2.2 (Boundary conditions) Other boundary conditions can be seamlessly handled by gradient schemes, see [21].

Remark 2.3 (Nonlinear function of the elements of  $X_{\mathcal{D},0}$ ) Let  $\mathcal{D}$  be a gradient discretisation in the sense of Definition 2.1. For any  $\chi: \mathbb{R} \to \mathbb{R}$  and any  $u = (u_i)_{i \in I} \in X_{\mathcal{D},0}$ , we define  $\chi_I(u) \in X_{\mathcal{D},0}$  by  $\chi_I(u) = (\chi(u_i))_{i \in I}$ . As indicated by the subscript I, this definition depends on the choice of the degrees of freedom in  $X_{\mathcal{D},0}$ . That said, these degrees of freedom are usually canonical and the index I can be dropped. An important consequence of the fact that  $\Pi_{\mathcal{D}}$  is a piecewise constant reconstruction is the following:

$$\forall \chi : \mathbb{R} \mapsto \mathbb{R}, \ \forall u \in X_{\mathcal{D},0}, \quad \Pi_{\mathcal{D}}\chi(u) = \chi(\Pi_{\mathcal{D}}u). \tag{12}$$

It is customary to use the notations  $\Pi_{\mathcal{D}}$  and  $\nabla_{\mathcal{D}}$  also for space-time dependent functions. Moreover, we will need a notation for the jump-in-time of piecewise constant functions in time. Hence, if  $(v^{(n)})_{n=0,...,N} \subset X_{\mathcal{D},0}$ , we set

for a.e. 
$$\boldsymbol{x} \in \Omega$$
,  $\Pi_{\mathcal{D}} v(\boldsymbol{x}, 0) = \Pi_{\mathcal{D}} v^{(0)}(\boldsymbol{x})$  and,  $\forall n = 0, \dots, N-1$ ,  $\forall t \in (t^{(n)}, t^{(n+1)}]$ ,  $\Pi_{\mathcal{D}} v(\boldsymbol{x}, t) = \Pi_{\mathcal{D}} v^{(n+1)}(\boldsymbol{x})$ ,  $\nabla_{\mathcal{D}} v(\boldsymbol{x}, t) = \nabla_{\mathcal{D}} v^{(n+1)}(\boldsymbol{x})$  and  $\delta_{\mathcal{D}} v(t) = \delta_{\mathcal{D}}^{(n+\frac{1}{2})} v := \frac{v^{(n+1)} - v^{(n)}}{\delta t^{(n+\frac{1}{2})}} \in X_{\mathcal{D},0}$ . (13)

If  $\mathcal{D} = (X_{\mathcal{D},0}, \Pi_{\mathcal{D}}, \nabla_{\mathcal{D}}, \mathcal{I}_{\mathcal{D}}, (t^{(n)})_{n=0,\dots,N})$  is a space-time gradient discretisation in the sense of Definition 2.1, the associated gradient scheme for Problem (1) is obtained by replacing in this problem

the continuous space and mappings with their discrete ones. Using the notations in Remark 2.3, the implicit-in-time gradient scheme therefore consists in considering a sequence  $(u^{(n)})_{n=0,...,N} \subset X_{\mathcal{D},0}$  such that

$$\begin{cases}
 u^{(0)} = \mathcal{I}_{\mathcal{D}} u_{\text{ini}} \text{ and, for all } v = (v^{(n)})_{n=1,\dots,N} \subset X_{\mathcal{D},0}, \\
 \int_{0}^{T} \int_{\Omega} \left[ \Pi_{\mathcal{D}} \delta_{\mathcal{D}} \beta(u)(\boldsymbol{x},t) \Pi_{\mathcal{D}} v(\boldsymbol{x},t) + \boldsymbol{a}(\boldsymbol{x},\Pi_{\mathcal{D}} \nu(u)(\boldsymbol{x},t), \nabla_{\mathcal{D}} \zeta(u)(\boldsymbol{x},t)) \cdot \nabla_{\mathcal{D}} v(\boldsymbol{x},t) \right] d\boldsymbol{x} dt \\
 = \int_{0}^{T} \int_{\Omega} f(\boldsymbol{x},t) \Pi_{\mathcal{D}} v(\boldsymbol{x},t) d\boldsymbol{x} dt.
\end{cases}$$
(14)

Remark 2.4 (Time-stepping) Scheme (14) is implicit-in-time because of the choice, in the definitions of  $\Pi_{\mathcal{D}}$  and  $\nabla_{\mathcal{D}}$  in (13), of  $v^{(n+1)}$  when  $t \in (t^{(n)}, t^{(n+1)}]$ . As a consequence,  $u^{(n+1)}$  appears in  $\mathbf{a}(\mathbf{x}, \cdot, \cdot)$  in (14) for  $t \in (t^{(n)}, t^{(n+1)}]$ . Instead of a fully implicit method, we could as well consider a Crank-Nicolson scheme or any scheme between those two ( $\theta$ -scheme). This would consist in choosing  $\theta \in [\frac{1}{2}, 1]$  and in replacing these terms  $u^{(n+1)}$  with  $u^{(n+\theta)} = \theta u^{(n+1)} + (1-\theta)u^{(n)}$ . All results established here for (14) would hold for such a scheme. We refer the reader to the treatment done in [23] for the details.

### 2.2 Properties of gradient discretisations

In order to establish the convergence of the associated gradient schemes, sequences of spacetime gradient discretisations are required to satisfy four properties: *coercivity*, *consistency*, *limit-conformity* and *compactness*.

**Definition 2.5 (Coercivity)** If  $\mathcal{D}$  is a space-time gradient discretisation in the sense of Definition 2.1, the norm of  $\Pi_{\mathcal{D}}$  is denoted by

$$C_{\mathcal{D}} = \max_{v \in X_{\mathcal{D},0} \setminus \{0\}} \frac{||\Pi_{\mathcal{D}}v||_{L^{p}(\Omega)}}{||\nabla_{\mathcal{D}}v||_{L^{p}(\Omega)^{d}}}.$$

A sequence  $(\mathcal{D}_m)_{m\in\mathbb{N}}$  of space-time gradient discretisations in the sense of Definition 2.1 is said to be coercive if there exists  $C_P \geq 0$  such that, for any  $m \in \mathbb{N}$ ,  $C_{\mathcal{D}_m} \leq C_P$ .

**Definition 2.6 (Consistency)** If  $\mathcal{D}$  is a space-time gradient discretisation in the sense of Definition 2.1, we define

$$\forall \varphi \in L^{2}(\Omega) \cap W_{0}^{1,p}(\Omega), \ \widehat{S}_{\mathcal{D}}(\varphi) = \min_{w \in X_{\mathcal{D},0}} \left( ||\Pi_{\mathcal{D}}w - \varphi||_{L^{\max(p,2)}(\Omega)} + ||\nabla_{\mathcal{D}}w - \nabla\varphi||_{L^{p}(\Omega)^{d}} \right).$$
 (15)

A sequence  $(\mathcal{D}_m)_{m\in\mathbb{N}}$  of space-time gradient discretisations in the sense of Definition 2.1 is said to be consistent if

- for all  $\varphi \in L^2(\Omega) \cap W_0^{1,p}(\Omega)$ ,  $\widehat{S}_{\mathcal{D}_m}(\varphi) \to 0$  as  $m \to \infty$ ,
- for all  $\varphi \in L^2(\Omega)$ ,  $\Pi_{\mathcal{D}_m} \mathcal{I}_{\mathcal{D}_m} \varphi \to \varphi$  in  $L^2(\Omega)$  as  $m \to \infty$ , and
- $\delta t_{\mathcal{D}_m} \to 0$  as  $m \to \infty$ .

**Definition 2.7 (Limit-conformity)** If  $\mathcal{D}$  is a space-time gradient discretisation in the sense of Definition 2.1 and  $W^{\text{div},p'}(\Omega) = \{\varphi \in L^{p'}(\Omega)^d : \text{div}\varphi \in L^{p'}(\Omega)\}$ , we define

$$\forall \varphi \in W^{\operatorname{div},p'}(\Omega), \ W_{\mathcal{D}}(\varphi) = \max_{u \in X_{\mathcal{D},0} \setminus \{0\}} \frac{\left| \int_{\Omega} \left( \nabla_{\mathcal{D}} u(x) \cdot \varphi(x) + \Pi_{\mathcal{D}} u(x) \operatorname{div} \varphi(x) \right) dx \right|}{\| \nabla_{\mathcal{D}} u \|_{L^{p}(\Omega)^{d}}}. \tag{16}$$

A sequence  $(\mathcal{D}_m)_{m\in\mathbb{N}}$  of space-time gradient discretisations in the sense of Definition 2.1 is said to be limit-conforming if, for all  $\varphi \in W^{\operatorname{div},p'}(\Omega)$ ,  $W_{\mathcal{D}_m}(\varphi) \to 0$  as  $m \to \infty$ .

**Remark 2.8** The convergences  $\widehat{S}_{\mathcal{D}_m} \to 0$  on  $L^2(\Omega) \cap W_0^{1,p}(\Omega)$  and  $W_{\mathcal{D}_m} \to 0$  on  $W^{\mathrm{div},p'}(\Omega)$  only need to be checked on dense subsets of these spaces [21, 31].

**Definition 2.9 (Compactness)** If  $\mathcal{D}$  is a space-time gradient discretisation in the sense of Definition 2.1, we define

$$\forall \boldsymbol{\xi} \in \mathbb{R}^d, \ T_{\mathcal{D}}(\boldsymbol{\xi}) = \max_{v \in X_{\mathcal{D},0} \setminus \{0\}} \frac{||\Pi_{\mathcal{D}}v(\cdot + \boldsymbol{\xi}) - \Pi_{\mathcal{D}}v||_{L^p(\mathbb{R}^d)}}{||\nabla_{\mathcal{D}}v||_{L^p(\Omega)^d}},$$

where  $\Pi_{\mathcal{D}}v$  has been extended by 0 outside  $\Omega$ .

A sequence  $(\mathcal{D}_m)_{m\in\mathbb{N}}$  of space-time gradient discretisations is said to be compact if

$$\lim_{\boldsymbol{\xi} \to 0} \sup_{m \in \mathbb{N}} T_{\mathcal{D}_m}(\boldsymbol{\xi}) = 0.$$

We refer the reader to [23, 21] for a proof of the following lemma.

Lemma 2.10 (Regularity of the limit) Let  $(\mathcal{D}_m)_{m\in\mathbb{N}}$  be a sequence of space-time gradient discretisations, in the sense of Definition 2.1, that is coercive and limit-conforming in the sense of Definitions 2.5 and 2.7. Let, for any  $m \in \mathbb{N}$ ,  $v_m = (v_m^{(n)})_{n=0,\dots,N_m} \subset X_{\mathcal{D}_m,0}$  be such that, with the notations in (13),  $(\nabla_{\mathcal{D}_m} v_m)_{m\in\mathbb{N}}$  is bounded in  $L^p(\Omega \times (0,T))^d$ .

Then there exists  $v \in L^p(0,T;W_0^{1,p}(\Omega))$  such that, up to a subsequence as  $m \to \infty$ ,  $\Pi_{\mathcal{D}_m}v_m \to v$  weakly in  $L^p(\Omega \times (0,T))$  and  $\nabla_{\mathcal{D}_m}v_m \to \nabla v$  weakly in  $L^p(\Omega \times (0,T))^d$ .

#### 2.3 Main results

Uniform-in-time convergence of numerical solutions to schemes for parabolic equations starts with a weak convergence with respect to the space variable. This weak convergence is then used to prove a stronger convergence. We therefore first recall a standard definition related to the weak topology of  $L^2(\Omega)$  (we also refer the reader to Proposition 6.5 in the appendix for a classical characterisation of the weak topology of bounded sets in  $L^2(\Omega)$ ).

**Definition 2.11 (Uniform-in-time**  $L^2(\Omega)$ -weak convergence) Let  $\langle \cdot, \cdot \rangle_{L^2(\Omega)}$  denote the inner product in  $L^2(\Omega)$ , let  $(u_m)_{m \in \mathbb{N}}$  be a sequence of functions  $[0,T] \to L^2(\Omega)$  and let  $u : [0,T] \mapsto L^2(\Omega)$ . We say that  $(u_m)_{m \in \mathbb{N}}$  converges weakly in  $L^2(\Omega)$  uniformly on [0,T] to u if, for all  $\varphi \in L^2(\Omega)$ , as  $m \to \infty$  the sequence of functions  $t \in [0,T] \to \langle u_m(t), \varphi \rangle_{L^2(\Omega)}$  converges uniformly on [0,T] to the function  $t \in [0,T] \to \langle u(t), \varphi \rangle_{L^2(\Omega)}$ .

Our first theorem states weak or space-time averaged convergence properties of gradient schemes for (1). These results have already been established for Leray–Lions', Richards' and Stefan's models, see [23, 28, 32]. The convergence proof we provide afterwards however covers more non-linear model and is more compact than the previous proofs.

**Theorem 2.12 (Convergence of gradient schemes)** We assume (2) and we take a sequence  $(\mathcal{D}_m)_{m\in\mathbb{N}}$  of space-time gradient discretisations, in the sense of Definition 2.1, that is coercive, consistent, limit-conforming and compact (see Section 2.2). Then for any  $m \in \mathbb{N}$  there exists a solution  $u_m$  to (14) with  $\mathcal{D} = \mathcal{D}_m$ .

Moreover, if we assume that

$$(\forall s \in \mathbb{R}, \ \beta(s) = s) \quad or \quad (\forall s \in \mathbb{R}, \ \zeta(s) = s),$$
 (17)

then there exists a solution  $\overline{u}$  to (4) such that, up to a subsequence, the following convergences hold as  $m \to \infty$ :

$$\Pi_{\mathcal{D}_m}\beta(u_m) \to \beta(\overline{u}) \text{ weakly in } L^2(\Omega) \text{ uniformly on } [0,T] \text{ (see Definition 2.11)}, 
\Pi_{\mathcal{D}_m}\nu(u_m) \to \nu(\overline{u}) \text{ strongly in } L^1(\Omega \times (0,T)), 
\Pi_{\mathcal{D}_m}\zeta(u_m) \to \zeta(\overline{u}) \text{ weakly in } L^p(\Omega \times (0,T)), 
\nabla_{\mathcal{D}_m}\zeta(u_m) \to \nabla\zeta(\overline{u}) \text{ weakly in } L^p(\Omega \times (0,T))^d.$$
(18)

Remark 2.13 Since  $|\nu| \leq L_{\zeta}|\beta|$  and  $|\nu| \leq L_{\beta}|\zeta|$ , the  $L^{\infty}(0,T;L^{2}(\Omega))$  bound on  $\Pi_{\mathcal{D}_{m}}\beta(u_{m})$  and the  $L^{p}(\Omega \times (0,T))$  bound on  $\Pi_{\mathcal{D}_{m}}\zeta(u_{m})$  (see Lemma 4.1 and Definition 2.5) shows that the strong convergence of  $\Pi_{\mathcal{D}_{m}}\nu(u_{m})$  is also valid in  $L^{q}(0,T;L^{r}(\Omega))$  for any  $(q,r) \in [1,\infty) \times [1,2)$ , any  $(q,r) \in [1,p)^{2}$  and, of course, any space interpolated between these two cases.

**Remark 2.14** We do not assume the existence of a solution  $\overline{u}$  to the continuous problem, our convergence analysis will establish this existence.

**Remark 2.15** Assumption (17) covers Richards' and Stefan's models, as well as many other non-linear parabolic equations. As we prove in Section 5, this assumption is actually not required if  $p \geq 2$ . However, we first state and prove Theorem 2.12 under (17) in order to simplify the presentation. See also Remark 2.19.

The main innovation of this paper is the following theorem, which states the *uniform-in-time* strong-in-space convergence of numerical methods for fully non-linear degenerate parabolic equations with no regularity assumptions on the data.

Theorem 2.16 (Uniform-in-time convergence) Under Assumptions (2), let  $(\mathcal{D}_m)_{m\in\mathbb{N}}$  be a sequence of space-time gradient discretisations, in the sense of Definition 2.1, that is coercive, consistent, limit-conforming and compact (see Section 2.2). We assume that  $u_m$  is a solution to (14) with  $\mathcal{D} = \mathcal{D}_m$  that converges as  $m \to \infty$  to a solution  $\overline{u}$  of (4) in the sense (18). Then, as  $m \to \infty$ ,  $\Pi_{\mathcal{D}_m} \nu(u_m) \to \nu(\overline{u})$  strongly in  $L^{\infty}(0,T;L^2(\Omega))$ .

**Remark 2.17** Since the functions  $\Pi_{\mathcal{D}_m}\nu(u_m)$  are piecewise constant in time, their convergence in  $L^{\infty}(0,T;L^2(\Omega))$  is actually a uniform-in-time convergence (not "uniform a.e. in time").

The last theorem completes our convergence result by stating the strong space-time averaged convergence of the discrete gradients. Its proof is inspired by the study of gradient schemes for Leray–Lions operators made in [23].

Theorem 2.18 (Strong convergence of gradients) Under Assumptions (2), let  $(\mathcal{D}_m)_{m\in\mathbb{N}}$  be a sequence of space-time gradient discretisations, in the sense of Definition 2.1, that is coercive, consistent, limit-conforming and compact (see Section 2.2). We assume that  $u_m$  is a solution to (14) with  $\mathcal{D} = \mathcal{D}_m$  that converges as  $m \to \infty$  to a solution  $\overline{u}$  of (4) in the sense (18). We also assume that  $\mathbf{a}$  is strictly monotone in the sense:

$$(\boldsymbol{a}(\boldsymbol{x}, s, \boldsymbol{\xi}) - \boldsymbol{a}(\boldsymbol{x}, s, \boldsymbol{\chi})) \cdot (\boldsymbol{\xi} - \boldsymbol{\chi}) > 0, \text{ for a.e. } \boldsymbol{x} \in \Omega, \ \forall s \in \mathbb{R}, \ \forall \boldsymbol{\xi} \neq \boldsymbol{\chi} \in \mathbb{R}^d.$$
 (19)

Then, as  $m \to \infty$ ,  $\Pi_{\mathcal{D}_m} \zeta(u_m) \to \zeta(\overline{u})$  strongly in  $L^p(\Omega \times (0,T))$  and  $\nabla_{\mathcal{D}_m} \zeta(u_m) \to \nabla \zeta(\overline{u})$  strongly in  $L^p(\Omega \times (0,T))^d$ .

**Remark 2.19** Theorems 2.16 and 2.18 do not require the structural assumption (17); they only require that the convergences (18) hold.

## 3 Preliminaries

We establish here a few results which will be used in the analysis of the gradient scheme (14).

## 3.1 Uniform-in-time compactness for space-time gradient discretisations

Aubin-Simon compactness results roughly consist in establishing the compactness of a sequence of space-time functions from some strong bounds on the functions with respect to the space variable (typically, bounds in a Sobolev space with positive exponent) and some weaker bounds on their time derivatives (typically, bounds in a Sobolev space with a negative exponent, i.e. the dual of a Sobolev space with positive exponent). Several variants exist, including for piecewise constant-in-time functions appearing in the numerical approximation of parabolic equations [17, 11, 4, 36]. Although quite strong in space, the convergence results provided by these discrete versions of Aubin-Simon theorems are only averaged-in-time – i.e. in an  $L^p(0,T;E)$  space where E is a normed space.

Theorem 3.1 can be considered as a discrete form of an Aubin-Simon theorem, that establishes a uniform-in-time but weak-in-space compactness result. The corresponding convergence is therefore weaker than in Theorem 2.16, but it is a critical initial step for establishing the uniform-in-time strong-in-space convergence result. Given that the functions considered here are piecewise constant in time, it might be surprising to obtain a uniform-in-time convergence result; everything hinges on the fact that the jumps in time tend to vanish as the time step goes to zero. The proof of Theorem 3.1 is based on the results in Section 6, and in particular on the discontinuous Ascoli-Arzelà theorem stated and proved there.

#### Theorem 3.1 (Uniform-in-time weak-in-space discrete Aubin-Simon theorem)

Let T > 0 and take a sequence  $(\mathcal{D}_m)_{m \in \mathbb{N}} = (X_{\mathcal{D}_m,0}, \Pi_{\mathcal{D}_m}, \nabla_{\mathcal{D}_m}, \mathcal{I}_{\mathcal{D}_m}, (t_m^{(n)})_{n=0,...,N_m})_{m \in \mathbb{N}}$  of spacetime gradient discretisations, in the sense of Definition 2.1, that is consistent in the sense of Definition 2.6.

For any  $m \in \mathbb{N}$ , let  $v_m = (v_m^{(n)})_{n=0,...,N_m} \subset X_{\mathcal{D}_m,0}$ . If there exists q > 1 and C > 0 such that, for any  $m \in \mathbb{N}$ ,

$$||\Pi_{\mathcal{D}_m} v_m||_{L^{\infty}(0,T;L^2(\Omega))} \le C \quad and \quad \int_0^T |\delta_m v_m(t)|_{\star,\mathcal{D}_m}^q dt \le C, \tag{20}$$

then the sequence  $(\Pi_{\mathcal{D}_m} v_m)_{m \in \mathbb{N}}$  is relatively compact uniformly-in-time and weakly in  $L^2(\Omega)$ , i.e. it has a subsequence that converges in the sense of Definition 2.11.

Moreover, any limit of such a subsequence is continuous  $[0,T] \to L^2(\Omega)$  for the weak topology of  $L^2(\Omega)$ .

**Remark 3.2** The bound on  $|\delta_m v_m|_{\star,\mathcal{D}_m}$  is often a consequence of a numerical scheme satisfied by  $v_m$  and of a bound on  $||\nabla_{\mathcal{D}_m} v_m||_{L^p(\Omega \times (0,T))^d}$ , see the proof of Lemma 4.3 for example.

**Proof.** This result is a consequence of the discontinuous Ascoli-Arzelà theorem (Theorem 6.2) with K = [0, T] and E the ball of radius C in  $L^2(\Omega)$  endowed with the weak topology. We let  $(\varphi_l)_{l \in \mathbb{N}} \subset C_c^{\infty}(\Omega)$  be a dense sequence in  $L^2(\Omega)$  and equipp E with the metric (82) from these  $\varphi_l$  (see Proposition 6.5). The set E is metric compact and therefore complete, and the functions  $\Pi_{\mathcal{D}_m} v_m$  take their values in E. It remains to estimate  $d_E(v_m(s), v_m(s'))$ . In what follows, we drop the index m in  $\mathcal{D}_m$  for the sake of legibility.

Let us define the interpolant  $P_{\mathcal{D}}\varphi_l \in X_{\mathcal{D},0}$  by

$$P_{\mathcal{D}}\varphi_l = \underset{w \in X_{\mathcal{D},0}}{\operatorname{argmin}} \left( ||\Pi_{\mathcal{D}}w - \varphi_l||_{L^{\max(p,2)}(\Omega)} + ||\nabla_{\mathcal{D}}w - \nabla\varphi_l||_{L^p(\Omega)^d} \right). \tag{21}$$

For  $0 \le s \le s' \le T$ , by writing  $\Pi_{\mathcal{D}}v_m(s') - \Pi_{\mathcal{D}}v_m(s)$  as the sum of its jumps  $\delta t^{(n+\frac{1}{2})}\Pi_{\mathcal{D}}\delta_{\mathcal{D}}^{(n+\frac{1}{2})}v_m$  at the points  $(t^{(n)})_{n=n_1,\ldots,n_2}$  between s and s', the definition of  $|\cdot|_{\star,\mathcal{D}}$ , Hölder's inequality and Estimate (20) give

$$\left| \int_{\Omega} \left( \Pi_{\mathcal{D}} v_m(\boldsymbol{x}, s') - \Pi_{\mathcal{D}} v_m(\boldsymbol{x}, s) \right) \Pi_{\mathcal{D}} P_{\mathcal{D}} \varphi_l(\boldsymbol{x}) d\boldsymbol{x} \right|$$

$$= \left| \int_{t^{(n_1)}}^{t^{(n_2+1)}} \int_{\Omega} \Pi_{\mathcal{D}} \delta_{\mathcal{D}} v(t)(\boldsymbol{x}) \Pi_{\mathcal{D}} P_{\mathcal{D}} \varphi_l(\boldsymbol{x}) d\boldsymbol{x} dt \right| \leq C^{1/q} (t^{(n_2+1)} - t^{(n_1)})^{1/q'} ||\nabla_{\mathcal{D}} P_{\mathcal{D}} \varphi_l||_{L^p(\Omega)^d}. \tag{22}$$

By definition of  $P_{\mathcal{D}}$ , we have

$$||\Pi_{\mathcal{D}}P_{\mathcal{D}}\varphi_l - \varphi_l||_{L^2(\Omega)} \le \widehat{S}_{\mathcal{D}}(\varphi_l)$$

and

$$||\nabla_{\mathcal{D}} P_{\mathcal{D}} \varphi_l||_{L^p(\Omega)^d} \le \widehat{S}_{\mathcal{D}}(\varphi_l) + ||\nabla \varphi_l||_{L^p(\Omega)^d} \le C_{\varphi_l}$$

with  $C_{\varphi_l}$  not depending on  $\mathcal{D}$  (and therefore on m). Since  $t^{(n_2+1)} - t^{(n_1)} \leq |s' - s| + \delta t$  and  $(\Pi_{\mathcal{D}} v_m)_{m \in \mathbb{N}}$  is bounded in  $L^{\infty}(0, T; L^2(\Omega))$ , we deduce from (22) that

$$\left| \int_{\Omega} \left( \Pi_{\mathcal{D}} v_m(\boldsymbol{x}, s') - \Pi_{\mathcal{D}} v_m(\boldsymbol{x}, s) \right) \varphi_l(\boldsymbol{x}) d\boldsymbol{x} \right|$$

$$\leq \left| \int_{\Omega} \left( \Pi_{\mathcal{D}} v_m(\boldsymbol{x}, s') - \Pi_{\mathcal{D}} v_m(\boldsymbol{x}, s) \right) \Pi_{\mathcal{D}} P_{\mathcal{D}} \varphi_l(\boldsymbol{x}) d\boldsymbol{x} \right| + 2||\Pi_{\mathcal{D}} v_m||_{L^{\infty}(0, T; L^{2}(\Omega))} ||\Pi_{\mathcal{D}} P_{\mathcal{D}} \varphi_l - \varphi_l||_{L^{2}(\Omega)}$$

$$\leq 2C \widehat{S}_{\mathcal{D}}(\varphi_l) + C^{1/q} C_{\varphi_l} |s' - s|^{1/q'} + C^{1/q} C_{\varphi_l} \delta t^{1/q'}.$$

Plugged into the definition (82) of the distance in E, this shows that

$$\begin{aligned} d_E\Big(\Pi_{\mathcal{D}}v_m(s'), \Pi_{\mathcal{D}}v_m(s)\Big) \\ &\leq \sum_{l \in \mathbb{N}} \frac{\min(1, C^{1/q'}C_{\varphi_l}|s'-s|^{1/q'})}{2^l} + \sum_{l \in \mathbb{N}} \frac{\min(1, 2C\widehat{S}_{\mathcal{D}_m}(\varphi_l) + C^{1/q'}C_{\varphi_l} \delta t_m^{1/q'})}{2^l} \\ &=: \ \omega(s, s') + \delta_m. \end{aligned}$$

Using the dominated convergence theorem for series, we see that  $\omega(s,s') \to 0$  as  $s-s' \to 0$  and that  $\delta_m \to 0$  as  $m \to \infty$  (we invoke the consistency to establish that  $\lim_{m \to \infty} \widehat{S}_{\mathcal{D}_m}(\varphi_l) \to 0$  for any l). Hence, the assumptions of Theorem 6.2 are satisfied and the proof is complete.

#### 3.2 Technical results

We state here a family of technical lemmas, starting with a few properties on  $\nu$  and B.

Lemma 3.3 Under Assumptions (2) there holds

$$|\nu(a) - \nu(b)| \le L_{\beta}|\zeta(a) - \zeta(b)|,\tag{23}$$

$$(\nu(a) - \nu(b))^2 \le L_\beta L_\zeta(\zeta(a) - \zeta(b))(\beta(a) - \beta(b)). \tag{24}$$

The function B is convex continuous on  $\overline{R_{\beta}}$ , the function  $B \circ \beta : \mathbb{R} \to [0, \infty)$  is continuous,

$$\forall s \in \mathbb{R}, \quad B(\beta(s)) = \int_0^s \zeta(q)\beta'(q)\mathrm{d}q, \qquad (25)$$

$$\exists K_0, K_1, K_2 > 0 \text{ such that, } \forall s \in \mathbb{R}, \quad K_0 \beta(s)^2 - K_1 \le B(\beta(s)) \le K_2 s^2,$$
 (26)

$$\forall a \in \mathbb{R}, \ \forall S \in \overline{R_{\beta}}, \quad \zeta(a)(S - \beta(a)) \le B(S) - B(\beta(a)),$$
 (27)

and

$$\forall s, s' \in \mathbb{R}, \ (\nu(s) - \nu(s'))^2 \le 4L_{\beta}L_{\zeta} \left[ B(\beta(s)) + B(\beta(s')) - 2B\left(\frac{\beta(s) + \beta(s')}{2}\right) \right]. \tag{28}$$

#### Proof.

Inequality (23) is a straightforward consequence of the estimate  $\nu' = \zeta'\beta' \leq L_{\beta}\zeta'$ . Note that the same inequality also holds with  $\beta$  and  $\zeta$  swapped. Since these functions are non-decreasing, Inequality (24) follows from (23) and the similar inequality with  $\beta$  and  $\zeta$  swapped.

Since  $\beta$  is non-decreasing,  $\beta_r$  is also non-decreasing on  $R_{\beta}$  and therefore locally bounded on  $R_{\beta}$ . Hence, B is locally Lipschitz-continuous on  $R_{\beta}$ , with an a.e. derivative  $B' = \zeta(\beta_r)$ . B' is therefore non-decreasing and B is convex continuous on  $R_{\beta}$ , and thus also on  $\overline{R_{\beta}}$  by choice of its values at the endpoints of  $R_{\beta}$ .

To prove (25), we denote by  $P \subset R_{\beta}$  the countable set of plateaux values of  $\beta$ , i.e. the  $y \in \mathbb{R}$  such that  $\beta^{-1}(\{y\})$  is not reduced to a singleton. If  $s \notin \beta^{-1}(P)$  then  $\beta^{-1}(\{\beta(s)\})$  is the singleton  $\{s\}$  and therefore  $\beta_r(\beta(s)) = s$ . Moreover,  $\beta_r$  is continuous at  $\beta(s)$  and thus B is differentiable at  $\beta(s)$  with  $B'(\beta(s)) = \zeta(\beta_r(\beta(s))) = \zeta(s)$ . Since  $\beta$  is differentiable a.e., we deduce that, for a.e.  $s \notin \beta^{-1}(P)$ ,  $(B(\beta))'(s) = B'(\beta(s))\beta'(s) = \zeta(s)\beta'(s)$ . The set  $\beta^{-1}(P)$  is a union of intervals on which  $\beta$  and thus  $B(\beta)$  are locally constant; hence, for a.e. s in this set,  $(B(\beta))'(s) = 0$  and  $\zeta(s)\beta'(s) = 0$ . Hence, the locally Lipschitz-continuous functions  $B(\beta)$  and  $s \to \int_0^s \zeta(q)\beta'(q)dq$  have identical derivatives a.e. on  $\mathbb{R}$  and take the same value at s = 0. They are thus equal on  $\mathbb{R}$  and the proof of (25) is complete.

The continuity of  $B \circ \beta$  is an obvious consequence of (25). The second inequality in (26) can also be easily deduced from (25) by noticing that  $|\zeta(s)\beta'(s)| \leq L_{\zeta}L_{\beta}|s|$  (we can take  $K_2 = \frac{L_{\zeta}L_{\beta}}{2}$ ). To prove the first inequality in (26), we start by inferring from (2b) the existence of S > 0 such that  $|\zeta(q)| \geq \frac{M_0}{2}|q| \geq \frac{M_0}{2L_{\beta}}|\beta(q)|$  whenever  $|q| \geq S$ . We then write, for  $s \geq S$ ,

$$B(\beta(s)) = \int_0^S \zeta(q)\beta'(q)\mathrm{d}q + \int_S^s \zeta(q)\beta'(q)\mathrm{d}q \ge \frac{M_0}{2L_\beta} \int_S^s \beta(q)\beta'(q)\mathrm{d}q = \frac{M_0}{4L_\beta} \left(\beta(s)^2 - \beta(S)^2\right).$$

A similar inequality holds for  $s \leq -S$  (with  $\beta(-S)$  instead of  $\beta(S)$ ) and the first inequality in (26) therefore holds with  $K_0 = \frac{M_0}{4L_\beta}$  and  $K_1 = \frac{M_0}{4L_\beta} \max_{[-S,S]} \beta^2$ .

We now prove (27), which states that  $\zeta(a)$  belongs to the convex sub-differential of B at  $\beta(a)$ . We first start with the case  $S \in R_{\beta}$ , that is  $S = \beta(b)$  for some  $b \in \mathbb{R}$ . If  $\beta_r$  is continuous at  $\beta(a)$  then this inequality is an obvious consequence of the convexity of B since B is then differentiable at  $\beta(a)$  with  $B'(\beta(a)) = \zeta(\beta_r(\beta(a))) = \zeta(a)$ . Otherwise, a plain reasoning also does the job:

$$B(S) - B(\beta(a)) = B(\beta(b)) - B(\beta(a))$$

$$= \int_a^b \zeta(q)\beta'(q)dq = \int_a^b (\zeta(q) - \zeta(a))\beta'(q)dq + \zeta(a)(\beta(b) - \beta(a)) \ge \zeta(a)(S - \beta(a)),$$

the inequality coming from the fact that  $\beta' \geq 0$  and that  $\zeta(q) - \zeta(a)$  has the same sign as b - a when q is between a and b. The general case  $S \in \overline{R_{\beta}}$  is obtained by passing to the limit on  $b_n$  such that  $\beta(b_n) \to S$  and by using the fact that B has limits (possibly  $+\infty$ ) at the endpoints of  $R_{\beta}$ . Let us now take  $s, s' \in \mathbb{R}$ . Let  $\bar{s} \in \mathbb{R}$  be such that  $\beta(\bar{s}) = \frac{\beta(s) + \beta(s')}{2}$ . We notice that

$$B(\beta(s)) + B(\beta(s')) - 2B(\beta(\bar{s})) = \int_{\bar{s}}^{s} (\zeta(q) - \zeta(\bar{s}))\beta'(q)dq + \int_{\bar{s}}^{s'} (\zeta(q) - \zeta(\bar{s}))\beta'(q)dq.$$
 (29)

We then notice that  $|\zeta(q) - \zeta(\bar{s})| \ge \frac{1}{L_{\beta}} |\nu(q) - \nu(\bar{s})|$  and  $\beta'(q) \ge \beta'(q) \frac{\zeta'(q)}{L_{\zeta}} = \frac{\nu'(q)}{L_{\zeta}}$ . If  $\tilde{s} = s$  or s', since  $\zeta(q) - \zeta(\bar{s})$  has the same sign as  $\tilde{s} - \bar{s}$  for all q between  $\bar{s}$  and  $\tilde{s}$ , we can write

$$\int_{\bar{s}}^{\tilde{s}} (\zeta(q) - \zeta(\bar{s})) \beta'(q) dq \ge \frac{1}{L_{\beta} L_{\zeta}} \int_{\bar{s}}^{\tilde{s}} \nu'(q) (\nu(q) - \nu(\bar{s})) dq = \frac{1}{2L_{\beta} L_{\zeta}} (\nu(\tilde{s}) - \nu(\bar{s}))^{2}.$$
(30)

Estimate (28) follows from (29), (30) and the inequality  $(\nu(s) - \nu(s'))^2 \le 2(\nu(s) - \nu(\bar{s}))^2 + 2(\nu(s') - \nu(\bar{s}))^2$ .

The next lemma is an easy consequence of Fatou's lemma and the fact that strongly lower semi-continuous convex functions are also weakly lower semi-continuous. We all the same provide its short proof.

**Lemma 3.4** Let I be a closed interval of  $\mathbb{R}$  and let  $H: I \to (-\infty, \infty]$  be a convex continuous function (continuity for possible infinite values, at the endpoints of I, corresponding to H having limits at these endpoints). We denote by  $L^2(\Omega; I)$  the convex set of functions in  $L^2(\Omega)$  with values in I. Let  $v \in L^2(\Omega; I)$  and let  $(v_m)_{m \in \mathbb{N}}$  be a sequence of functions in  $L^2(\Omega; I)$  that converges weakly to v in  $L^2(\Omega)$ . Then

$$\int_{\Omega} H(v(\boldsymbol{x})) d\boldsymbol{x} \leq \liminf_{m \to \infty} \int_{\Omega} H(v_m(\boldsymbol{x})) d\boldsymbol{x}.$$

#### Proof.

For  $w \in L^2(\Omega; I)$  we set  $\Phi(w) = \int_{\Omega} H(w(x)) dx$ . Since H is convex, it is greater than a linear functional and  $\Phi(w)$  is thus well defined in  $(-\infty, \infty]$ . Moreover, if  $w_k \to w$  strongly in  $L^2(\Omega; I)$  then, up to a subsequence,  $w_k \to w$  a.e. on  $\Omega$  and therefore  $H(w_k) \to H(w)$  a.e. on  $\Omega$ . Thanks to the linear lower bound of H, we can apply Fatou's lemma to see that  $\Phi(w) \leq \liminf_{k \to \infty} \Phi(w_k)$ . Hence,  $\Phi$  is lower semi-continuous for the strong topology of  $L^2(\Omega; I)$ . Since  $\Phi$  (like H) is convex, we deduce that this lower semi-continuity property is also valid for the weak topology of  $L^2(\Omega; I)$ , see [26]. The result of the lemma is just the translation of this weak lower semi-continuity of  $\Phi$ .

The last technical result is a consequence of the Minty trick. It has been proved and used in the  $L^2$  case in [28, 21], but we need here an extension to the non-Hilbertian case.

**Lemma 3.5 (Minty's trick)** Let  $H \in C^0(\mathbb{R})$  be a nondecreasing function. Let  $(X, \mu)$  be a measurable set with finite measure and let  $(u_n)_{n\in\mathbb{N}} \subset L^p(X)$ , with p>1, satisfy

- 1. there exists  $u \in L^p(X)$  such that  $(u_n)_{n \in \mathbb{N}}$  converges weakly to u in  $L^p(X)$ ;
- 2.  $(H(u_n))_{n\in\mathbb{N}}\subset L^1(X)$  and there exists  $w\in L^1(X)$  such that  $(H(u_n))_{n\in\mathbb{N}}$  converges strongly to w in  $L^1(X)$ ;

Then w = H(u) a.e. on X.

#### Proof.

For k, l > 0 we define the truncation at levels -l and k by  $T_{k,l}(s) = \max(-l, \min(s, k))$  and we let  $T_k = T_{k,k}$ . Since H is non-decreasing, there exists sequences  $(h_k)_{k \in \mathbb{N}}$  and  $(m_k)_{k \in \mathbb{N}}$  that tend to  $+\infty$  as  $k \to \infty$  and such that  $H(T_k(s)) = T_{h_k,m_k}(H(s))$ . Thus,  $H(T_k(u_n)) \to T_{h_k,m_k}(w)$  in  $L^1(X)$  as  $n \to \infty$ . Given that  $(H(T_k(u_n)))_{n \in \mathbb{N}}$  remains bounded in  $L^\infty(X)$ , its convergence to  $T_{h_k,m_k}(w)$  also holds in  $L^{p'}(X)$ .

Using fact that  $H \circ T_k$  is non-decreasing, we write for any  $g \in L^p(X)$ 

$$\int_{X} (H(T_k(u_n)) - H(T_k(g)))(u_n - g) d\mu \ge 0.$$

By strong convergence of  $H(T_k(u_n))$  in  $L^{p'}(X)$  and weak convergence of  $u_n$  in  $L^p(X)$ , as well as the fact that  $H \circ T_k$  is bounded, we can take the limit of this expression as  $n \to \infty$  and we find

$$\int_{X} (T_{h_k, m_k}(w) - H(T_k(g)))(u - g) d\mu \ge 0.$$
(31)

We then use Minty's trick. We pick a generic  $\varphi \in L^p(X)$ , apply (31) to  $g = u - t\varphi$ , divide by t and let  $t \to \pm 0$  (using the dominated convergence theorem and the fact that  $H \circ T_k$  is continuous and bounded) to find

$$\int_X (T_{h_k,m_k}(w) - H(T_k(u)))\varphi d\mu = 0.$$

Selecting  $\varphi = \text{sign}(T_{h_k,m_k}(w) - H(T_k(u)))$ , we deduce that  $T_{h_k,m_k}(w) = H(T_k(u))$  a.e. on X. Letting  $k \to \infty$ , we conclude that w = H(u) a.e. on X.

## 3.3 Integration-by-parts for the continuous solution

The last series of preliminary results are properties on the solution to (4), all based on the following integration-by-parts property. This property, used in the proof of Theorems 2.12 and 2.16, enables us to compute the value of the linear form  $\partial_t \beta(\overline{u}) \in L^{p'}(0,T;W^{-1,p'}(\Omega))$  on the function  $\zeta(\overline{u}) \in L^p(0,T;W^{1,p}(\Omega))$ . Because of the lack of regularity on  $\overline{u}$  and the double non-linearity ( $\beta$  and  $\zeta$ ), justifying this integration-by-parts is however not straightforward at all...

**Lemma 3.6** Let us assume (2b) and (2c). Let  $v: \Omega \times (0,T) \mapsto \mathbb{R}$  be measurable such that  $\zeta(v) \in L^p(0,T;W_0^{1,p}(\Omega)), \ B(\beta(v)) \in L^\infty(0,T;L^1(\Omega)), \ \beta(v) \in C([0,T];L^2(\Omega)-w)$  and  $\partial_t \beta(v) \in L^{p'}(0,T;W^{-1,p'}(\Omega)).$  Then  $t \in [0,T] \to \int_{\Omega} B(\beta(v)(\boldsymbol{x},t)) d\boldsymbol{x} \in [0,\infty)$  is continuous and, for all  $t_1,t_2 \in [0,T]$ ,

$$\int_{t_1}^{t_2} \langle \partial_t \beta(v)(t), \zeta(v(\cdot, t)) \rangle_{W^{-1, p'}, W_0^{1, p}} dt = \int_{\Omega} B(\beta(v)(\boldsymbol{x}, t_2)) d\boldsymbol{x} - \int_{\Omega} B(\beta(v)(\boldsymbol{x}, t_1)) d\boldsymbol{x}.$$
(32)

**Remark 3.7** Similarly to the discussion at the end of Section 1.2, we notice that it is important to keep in mind the separation between  $\beta(v(\cdot,\cdot))$  and its continuous representative  $\beta(v)(\cdot,\cdot)$ .

### Proof.

Without loss of generality, we assume that  $0 \le t_1 < t_2 \le T$ .

**Step 1**: truncation, extension and approximation of  $\beta(v)$ .

We define  $\overline{\beta(v)}: \mathbb{R} \to L^2(\Omega)$  by setting

$$\overline{\beta(v)}(t) = \begin{cases} \beta(v)(t) & \text{if } t \in [t_1, t_2], \\ \beta(v)(t_1) & \text{if } t \leq t_1, \\ \beta(v)(t_2) & \text{if } t \geq t_2. \end{cases}$$

By the continuity property of  $\beta(v)$ , this definition makes sense and gives  $\overline{\beta(v)} \in C(\mathbb{R}; L^2(\Omega)\text{-w})$  such that  $\partial_t \overline{\beta(v)} = \mathbf{1}_{(t_1,t_2)} \partial_t \beta(v) \in L^{p'}(\mathbb{R}; W^{-1,p'}(\Omega))$  where **1** is the characteristic function (no Dirac masses have been introduced at  $t = t_1$  or  $t = t_2$ ). This regularity of  $\partial_t \overline{\beta(v)}$  ensures that the function  $D_h \overline{\beta(v)} : \mathbb{R} \mapsto W^{-1,p'}(\Omega)$  defined by

$$\forall t \in \mathbb{R} , \ D_h \overline{\beta(v)}(t) = \frac{1}{h} \int_{t}^{t+h} \partial_t \overline{\beta(v)}(s) ds = \frac{\overline{\beta(v)}(t+h) - \overline{\beta(v)}(t)}{h}$$
(33)

tends to  $\partial_t \overline{\beta(v)}$  in  $L^{p'}(\mathbb{R}; W^{-1,p'}(\Omega))$  as  $h \to 0$ .

Step 2: we prove that  $||B(\overline{\beta(v)}(t))||_{L^1(\Omega)} \le ||B(\beta(v))||_{L^{\infty}(0,T;L^1(\Omega))}$  for all  $t \in \mathbb{R}$  (not only for a.e. t).

Let  $t \in [t_1, t_2]$ . Since  $\beta(v)(\cdot, \cdot) = \beta(v(\cdot, \cdot))$  a.e. on  $\Omega \times (t_1, t_2)$ , there exists a sequence  $t_n \to t$  such that  $\beta(v)(\cdot, t_n) = \beta(v(\cdot, t_n))$  in  $L^2(\Omega)$  and  $||B(\beta(v)(\cdot, t_n))||_{L^1(\Omega)} \le ||B(\beta(v))||_{L^{\infty}(0,T;L^1(\Omega))}$  for all n. As  $\beta(v) \in C([0,T];L^2(\Omega)-w)$ , we have  $\beta(v)(\cdot, t_n) \to \beta(v)(\cdot, t)$  weakly in  $L^2(\Omega)$ . We then use the convexity of B and Lemma 3.4 to write, thanks to our choice of  $t_n$ ,

$$\int_{\Omega} B(\beta(v)(\boldsymbol{x},t)) d\boldsymbol{x} \leq \liminf_{n \to \infty} \int_{\Omega} B(\beta(v)(\boldsymbol{x},t_n)) d\boldsymbol{x} \leq ||B(\beta(v))||_{L^{\infty}(0,T;L^{1}(\Omega))}$$

and the proof is complete for  $t \in [t_1, t_2]$ . The result for  $t \le t_1$  or  $t \ge t_2$  is obvious since  $\overline{\beta(v)}(t)$  is then either  $\beta(v)(t_1)$  or  $\beta(v)(t_2)$ .

**Step 3**: We prove that for all  $\tau \in \mathbb{R}$  and a.e.  $t \in (t_1, t_2)$ ,

$$\langle \overline{\beta(v)}(\tau) - \beta(v)(t), \zeta(v(\cdot,t)) \rangle_{W^{-1,p'}, W_0^{1,p}} \le \int_{\Omega} B(\overline{\beta(v)}(\boldsymbol{x},\tau)) - B(\beta(v)(\boldsymbol{x},t)) d\boldsymbol{x}. \tag{34}$$

If we could just replace the duality product  $W^{-1,p'}-W_0^{1,p}$  with an  $L^2$  inner product, this formula would be a straightforward consequence of (27). The problem is that nothing ensures that  $\zeta(v)(t) \in L^2(\Omega)$  for a.e. t.

We first notice that  $\overline{\beta(v)}(\tau) - \beta(v)(t) = \int_t^{\tau} \partial_t \overline{\beta(v)}(s) ds$  belongs to  $W^{-1,p'}(\Omega)$  so the left-hand side of (34) makes sense provided that t is chosen such that  $\zeta(v(\cdot,t)) \in W_0^{1,p}(\Omega)$  (which we do from here on). To deal with the fact that  $\zeta(v(\cdot,t))$  does not necessarily belong to  $L^2(\Omega)$ , we replace it with a truncation. As in the proof of Lemma 3.5, we introduce  $T_{k,l}(s) = \max(-l, \min(s, k))$  and we let  $T_k = T_{k,k}$ . By the monotony assumption (2b) on  $\zeta$  we see that there exists sequences  $(r_k)_{k \in \mathbb{N}}$  and  $(l_k)_{k \in \mathbb{N}}$  that tend to  $+\infty$  as  $k \to +\infty$  and such that  $\zeta(T_k(v(\cdot,t))) = T_{r_k,l_k}(\zeta(v(\cdot,t)))$ . Hence,  $\zeta(T_k(v(\cdot,t))) \in W_0^{1,p}(\Omega)$  and converges, as  $k \to \infty$ , to  $\zeta(v(\cdot,t))$  in  $W_0^{1,p}(\Omega)$ . We can therefore write

$$\langle \overline{\beta(v)}(\tau) - \beta(v)(t), \zeta(v(\cdot,t)) \rangle_{W^{-1,p'}, W_0^{1,p}} = \lim_{k \to \infty} \langle \overline{\beta(v)}(\tau) - \beta(v)(t), \zeta(T_k(v(\cdot,t))) \rangle_{W^{-1,p'}, W_0^{1,p}} 
= \lim_{k \to \infty} \int_{\Omega} \left[ \overline{\beta(v)}(\boldsymbol{x}, \tau) - \beta(v(\boldsymbol{x}, t)) \right] \zeta(T_k(v(\boldsymbol{x}, t))) d\boldsymbol{x}, \quad (35)$$

the replacement of the duality product by an  $L^2(\Omega)$  inner product being justified since  $\overline{\beta(v)}(\tau) - \beta(v)(t)$  and  $\zeta(T_k(v(\cdot,t)))$  both belong to  $L^2(\Omega)$ . We also used that, for a.e.  $t \in (t_1,t_2)$ ,  $\beta(v)(\cdot,t) = \beta(v(\cdot,t))$  a.e. on  $\Omega$ ; hence (35) is valid for a.e.  $t \in (t_1,t_2)$ .

We then write  $\beta(v(\boldsymbol{x},t)) = \beta(T_k(v(\boldsymbol{x},t))) + [\beta(v(\boldsymbol{x},t)) - \beta(T_k(v(\boldsymbol{x},t)))]$  and apply (27) with  $S = \beta(v)(\boldsymbol{x},\tau)$  and  $a = T_k(v(\boldsymbol{x},t))$  to find

$$\begin{split} \int_{\Omega} \left[ \overline{\beta(v)}(\boldsymbol{x},\tau) - \beta(v(\boldsymbol{x},t)) \right] \zeta(T_k(v(\boldsymbol{x},t))) \mathrm{d}\boldsymbol{x} \\ &= \int_{\Omega} \left[ \overline{\beta(v)}(\boldsymbol{x},\tau) - \beta(T_k(v(\boldsymbol{x},t))) \right] \zeta(T_k(v(\boldsymbol{x},t))) \mathrm{d}\boldsymbol{x} \\ &- \int_{\Omega} \left[ \beta(v(\boldsymbol{x},t)) - \beta(T_k(v(\boldsymbol{x},t))) \right] \zeta(T_k(v(\boldsymbol{x},t))) \mathrm{d}\boldsymbol{x} \\ &\leq \int_{\Omega} B(\overline{\beta(v)}(\boldsymbol{x},\tau)) - B(\beta(T_k(v(\boldsymbol{x},t)))) \mathrm{d}\boldsymbol{x} - \int_{\Omega} \left[ \beta(v(\boldsymbol{x},t)) - \beta(T_k(v(\boldsymbol{x},t))) \right] \zeta(T_k(v(\boldsymbol{x},t))) \mathrm{d}\boldsymbol{x}. \end{split}$$

By the monotony of  $\beta$ , the sign of  $\zeta$  and by studying the cases  $v(x,t) \geq k$ ,  $-k \leq v(x,t) \leq k$  and  $v(\boldsymbol{x},t) \leq -k$ , we notice that the last integrand is everywhere non-negative. We can therefore write

$$\int_{\Omega} \left[ \overline{\beta(v)}(\boldsymbol{x},\tau) - \beta(v(\boldsymbol{x},t)) \right] \zeta(T_k(v(\boldsymbol{x},t))) d\boldsymbol{x} \le \int_{\Omega} B(\overline{\beta(v)}(\boldsymbol{x},\tau)) - B(\beta(T_k(v(\boldsymbol{x},t)))) d\boldsymbol{x}.$$

We then use the continuity of  $B \circ \beta$  and Fatou's lemma to deduce

$$\begin{split} \limsup_{k \to \infty} \int_{\Omega} \left[ \overline{\beta(v)}(\boldsymbol{x}, \tau) - \beta(v(\boldsymbol{x}, t)) \right] \zeta(T_k(v(\boldsymbol{x}, t))) \mathrm{d}\boldsymbol{x} \\ & \leq \int_{\Omega} B(\overline{\beta(v)}(\boldsymbol{x}, \tau)) \mathrm{d}\boldsymbol{x} - \liminf_{k \to \infty} \int_{\Omega} B(\beta(T_k(v(\boldsymbol{x}, t)))) \mathrm{d}\boldsymbol{x} \\ & \leq \int_{\Omega} B(\overline{\beta(v)}(\boldsymbol{x}, \tau)) \mathrm{d}\boldsymbol{x} - \int_{\Omega} B(\beta(v(\boldsymbol{x}, t))) \mathrm{d}\boldsymbol{x} \end{split}$$

which, combined with (35), concludes the proof of (34) (recall that t has been chosen such that  $\beta(v(\cdot,t)) = \beta(v)(\cdot,t)$  a.e. on  $\Omega$ ).

**Step 4**: proof of the formula Since  $\mathbf{1}_{(t_1,t_2)}\zeta(v) \in L^p(\mathbb{R};W_0^{1,p}(\Omega))$  and  $D_h\overline{\beta(v)} \to \partial_t\overline{\beta(v)}$  in  $L^{p'}(\mathbb{R};W^{-1,p'}(\Omega))$  as  $h \to 0$ , we have

$$\int_{t_1}^{t_2} \langle \partial_t \beta(v)(t), \zeta(v(\cdot,t)) \rangle_{W^{-1,p'}, W_0^{1,p}} dt = \int_{\mathbb{R}} \langle \partial_t \overline{\beta(v)}(t), \mathbf{1}_{(t_1, t_2)}(t) \zeta(v(\cdot, t)) \rangle_{W^{-1,p'}, W_0^{1,p}} dt$$

$$= \lim_{h \to 0} \int_{\mathbb{R}} \langle D_h \overline{\beta(v)}(t), \mathbf{1}_{(t_1, t_2)}(t) \zeta(v(\cdot, t)) \rangle_{W^{-1,p'}, W_0^{1,p}} dt$$

$$= \lim_{h \to 0} \frac{1}{h} \int_{t_1}^{t_2} \langle \overline{\beta(v)}(t+h) - \overline{\beta(v)}(t), \zeta(v(\cdot, t)) \rangle_{W^{-1,p'}, W_0^{1,p}} dt. \quad (36)$$

We then use (34) for a.e.  $t \in (t_1, t_2)$  to obtain, for h small enough such that  $t_1 + h < t_2$ ,

$$\frac{1}{h} \int_{t_{1}}^{t_{2}} \langle \overline{\beta(v)}(t+h) - \overline{\beta(v)}(t), \zeta(v(\cdot,t)) \rangle_{W^{-1,p'},W_{0}^{1,p}} dt$$

$$\leq \frac{1}{h} \int_{t_{1}}^{t_{2}} \int_{\Omega} B(\overline{\beta(v)}(\boldsymbol{x},t+h)) - B(\overline{\beta(v)}(\boldsymbol{x},t)) d\boldsymbol{x} dt$$

$$= \frac{1}{h} \int_{t_{2}}^{t_{2}+h} \int_{\Omega} B(\overline{\beta(v)}(\boldsymbol{x},t)) d\boldsymbol{x} dt - \frac{1}{h} \int_{t_{1}}^{t_{1}+h} \int_{\Omega} B(\overline{\beta(v)}(\boldsymbol{x},t)) d\boldsymbol{x} dt$$

$$= \int_{\Omega} B(\beta(v)(\boldsymbol{x},t_{2})) d\boldsymbol{x} - \frac{1}{h} \int_{t_{1}}^{t_{1}+h} \int_{\Omega} B(\beta(v)(\boldsymbol{x},t)) d\boldsymbol{x} dt. \tag{37}$$

We used the estimate in Step 2 to justify the separation of the integrals in (37). We now take the lim sup as  $h \to 0$  of this inequality, using again Step 2 to see that  $B(\beta(v)(\cdot,t_2))$  is integrable and therefore take its integral out of the lim sup. Coming back to (36) we obtain

$$\int_{t_{1}}^{t_{2}} \langle \partial_{t} \beta(v)(t), \zeta(v(\cdot,t)) \rangle_{W^{-1,p'}, W_{0}^{1,p}} dt$$

$$\leq \int_{\Omega} B(\beta(v)(\boldsymbol{x}, t_{2})) d\boldsymbol{x} - \liminf_{h \to 0} \frac{1}{h} \int_{t_{1}}^{t_{1}+h} \int_{\Omega} B(\beta(v)(\boldsymbol{x}, t)) d\boldsymbol{x} dt. \quad (38)$$

Since  $\beta(v) \in C([0,T]; L^2(\Omega)\text{-w})$ , as  $h \to 0$  we have  $\frac{1}{h} \int_{t_1}^{t_1+h} \beta(v)(t) dt \to \beta(v)(t_1)$  weakly in  $L^2(\Omega)$ . Hence, the convexity of B, Lemma 3.4 and Jensen's inequality give

$$\int_{\Omega} B(\beta(v)(\boldsymbol{x}, t_1)) d\boldsymbol{x} \leq \liminf_{h \to 0} \int_{\Omega} B\left(\frac{1}{h} \int_{t_1}^{t_1 + h} \beta(v)(\boldsymbol{x}, t) dt\right) d\boldsymbol{x} \\
\leq \liminf_{h \to 0} \int_{\Omega} \frac{1}{h} \int_{t_1}^{t_1 + h} B(\beta(v)(\boldsymbol{x}, t)) dt d\boldsymbol{x}.$$

Plugged into (38), this inequality shows that (32) holds with  $\leq$  instead of =. The reverse inequality is obtained by reversing the time. We consider  $\widetilde{v}(t) = v(t_1 + t_2 - t)$ . Then  $\zeta(\widetilde{v})$ ,  $B(\beta(\widetilde{v}))$  and  $\beta(\widetilde{v})$  have the same properties as  $\zeta(v)$ ,  $B(\beta(v))$  and  $\beta(v)$ , and  $\beta(\widetilde{v})$  takes values  $\beta(v)(t_1)$  at  $t = t_2$  and  $\beta(v)(t_2)$  at  $t = t_1$ . Applying (32) with " $\leq$ " instead of "=" to  $\widetilde{v}$  and using the fact that  $\partial_t \beta(\widetilde{v})(t) = -\partial_t \beta(v)(t_1 + t_2 - t)$ , we obtain (32) with " $\geq$ " instead of "=" and the proof of (32) is complete.

The continuity of  $t \in [0,T] \mapsto \int_{\Omega} B(\beta(v)(\boldsymbol{x},t)) d\boldsymbol{x}$  is straightforward from (32) as the left-hand side of this relation is continuous with respect to  $t_1$  and  $t_2$ .

The following corollary states continuity properties and an essential formula on the solution to (4).

Corollary 3.8 Under Assumptions (2a)–(2i), if  $\overline{u}$  is a solution of (4) then:

- 1. the function  $t \in [0,T] \mapsto \int_{\Omega} B(\beta(\overline{u})(\boldsymbol{x},t)) d\boldsymbol{x} \in [0,\infty)$  is continuous and bounded,
- 2. for any  $T_0 \in [0, T]$ ,

$$\int_{\Omega} B(\beta(\overline{u})(\boldsymbol{x}, T_0)) d\boldsymbol{x} + \int_{0}^{T_0} \int_{\Omega} \boldsymbol{a}(\boldsymbol{x}, \nu(\overline{u}(\boldsymbol{x}, t)), \nabla \zeta(\overline{u})(\boldsymbol{x}, t)) \cdot \nabla \zeta(\overline{u})(\boldsymbol{x}, t) d\boldsymbol{x} dt 
= \int_{\Omega} B(\beta(u_{\text{ini}}(\boldsymbol{x}))) d\boldsymbol{x} + \int_{0}^{T_0} \int_{\Omega} f(\boldsymbol{x}, t) \zeta(\overline{u})(\boldsymbol{x}, t) d\boldsymbol{x} dt, \quad (39)$$

3.  $\nu(\overline{u})$  is continuous  $[0,T] \to L^2(\Omega)$ .

Remark 3.9 The continuity of  $\nu(\overline{u})$  has to be understood in the same sense as the continuity of  $\beta(\overline{u})$ , that is  $\nu(\overline{u})$  is a.e. on  $\Omega \times (0,T)$  equal to a continuous function  $[0,T] \to L^2(\Omega)$ . We use in particular the notation  $\nu(\overline{u})(\cdot,\cdot)$  for the continuous-in-time representative of  $\nu(\overline{u}(\cdot,\cdot))$ , similarly to the way we denote the continuous-in-time representative of  $\beta(\overline{u}(\cdot,\cdot))$ .

#### Proof.

The continuity of  $t \in [0,T] \mapsto \int_{\Omega} B(\beta(\overline{u})(\boldsymbol{x},t)) d\boldsymbol{x} \in [0,\infty)$  and Formula (39) are straightforward consequences of Lemma 3.6 with  $v = \overline{u}$  and using (4) with  $\overline{v} = \zeta(\overline{u})$ . Note that the bound on  $\int_{\Omega} B(\beta(\overline{u})(\boldsymbol{x},t)) d\boldsymbol{x}$  can be seen as a consequence of (39), or from Step 2 in the proof of Lemma 3.6.

Let us prove the strong continuity of  $\nu(\overline{u}): [0,T] \mapsto L^2(\Omega)$ . Let  $\mathcal{T}$  be the set of  $\tau \in [0,T]$  such that  $\beta(\overline{u}(\cdot,\tau)) = \beta(\overline{u})(\cdot,\tau)$  a.e. on  $\Omega$ , and let  $(s_l)_{l\in\mathbb{N}}$  and  $(t_k)_{k\in\mathbb{N}}$  be two sequences in  $\mathcal{T}$  that converge to the same value s. Invoking (28) we can write

$$\int_{\Omega} (\nu(\overline{u}(\boldsymbol{x}, s_{l})) - \nu(\overline{u}(\boldsymbol{x}, t_{k})))^{2} d\boldsymbol{x} \leq 4L_{\beta}L_{\zeta} \left( \int_{\Omega} B(\beta(\overline{u})(\boldsymbol{x}, s_{l})) d\boldsymbol{x} + \int_{\Omega} B(\beta(\overline{u})(\boldsymbol{x}, t_{k})) d\boldsymbol{x} \right) \\
- 8L_{\beta}L_{\zeta} \int_{\Omega} B\left( \frac{\beta(\overline{u})(\boldsymbol{x}, s_{l}) + \beta(\overline{u})(\boldsymbol{x}, t_{k})}{2} \right) d\boldsymbol{x}. \quad (40)$$

Since  $\frac{\beta(\overline{u})(\cdot,s_l)+\beta(\overline{u})(\cdot,t_k)}{2} \to \beta(\overline{u})(\cdot,s)$  weakly in  $L^2(\Omega)$  as  $l,k\to\infty$ , Lemma 3.4 gives

$$\int_{\Omega} B(\beta(\overline{u})(\boldsymbol{x},s)) d\boldsymbol{x} \leq \liminf_{l,k \to \infty} \int_{\Omega} B\left(\frac{\beta(\overline{u})(\boldsymbol{x},s_l) + \beta(\overline{u})(\boldsymbol{x},t_k)}{2}\right) d\boldsymbol{x}.$$

Taking the lim sup as  $l, k \to \infty$  of (40) and using the continuity of  $t \mapsto \int_{\Omega} B(\beta(\overline{u})(x, t)) dx$  thus shows that

$$||\nu(\overline{u}(\cdot, s_l)) - \nu(\overline{u}(\cdot, t_k))||_{L^2(\Omega)} \to 0 \quad \text{as } l, k \to \infty.$$
 (41)

The existence of an a.e. representative of  $\nu(\overline{u}(\cdot,\cdot))$  which is continuous  $[0,T] \mapsto L^2(\Omega)$  is a direct consequence of this convergence. Let  $s \in [0,T]$  and  $(s_l)_{l \in \mathbb{N}} \subset \mathcal{T}$  that converges to s. Applied with  $t_k = s_k$ , (41) shows that  $(\nu(\overline{u}(\cdot,s_l)))_{l \in \mathbb{N}}$  is a Cauchy sequence in  $L^2(\Omega)$  and therefore that  $\lim_{l \to \infty} \nu(\overline{u}(\cdot,s_l))$  exists in  $L^2(\Omega)$ . Moreover, (41) shows that this limit, that we denote by  $\nu(\overline{u})(\cdot,s)$ , does not depend on the sequence in  $\mathcal{T}$  that converges to s. Whenever  $s \in \mathcal{T}$ , the choice  $t_k = s$  in (41) shows that  $\nu(\overline{u})(\cdot,s) = \nu(\overline{u}(\cdot,s))$  a.e. on  $\Omega$ , and  $\nu(\overline{u})(\cdot,\cdot)$  is therefore equal to  $\nu(\overline{u}(\cdot,\cdot))$  a.e. on  $\Omega \times (0,T)$ .

It remains to establish that  $\nu(\overline{u})$  thus defined is continuous  $[0,T] \mapsto L^2(\Omega)$ . For any  $(\tau_r)_{r \in \mathbb{N}} \subset [0,T]$  that converges to  $\tau \in [0,T]$ , we can pick  $s_r \in \mathcal{T} \cap (\tau_r - \frac{1}{r}, \tau_r + \frac{1}{r})$  and  $t_r \in \mathcal{T} \cap (\tau - \frac{1}{r}, \tau + \frac{1}{r})$  such that

$$||\nu(\overline{u})(\cdot,\tau_r) - \nu(\overline{u}(\cdot,s_r))||_{L^2(\Omega)} \le \frac{1}{r}, \quad ||\nu(\overline{u})(\cdot,\tau) - \nu(\overline{u}(\cdot,t_r))||_{L^2(\Omega)} \le \frac{1}{r}.$$

We therefore have

$$||\nu(\overline{u})(\cdot,\tau_r) - \nu(\overline{u})(\cdot,\tau)||_{L^2(\Omega)} \le \frac{2}{r} + ||\nu(\overline{u}(\cdot,s_r)) - \nu(\overline{u}(\cdot,t_r))||_{L^2(\Omega)}.$$

This proves by (41) with l = k = r that  $\nu(\overline{u})(\cdot, \tau_r) \to \nu(\overline{u})(\cdot, \tau)$  in  $L^2(\Omega)$  as  $r \to \infty$ , and the proof is complete.

## 4 Proof of the convergence theorems

#### 4.1 Estimates on the approximate solution

As usual in the study of numerical methods for PDE with strong non-linearities or without regularity assumptions on the data, everything starts with a priori estimates.

Lemma 4.1 ( $L^{\infty}(0,T;L^2(\Omega))$ ) estimate and discrete  $L^p(0,T;W_0^{1,p}(\Omega))$  estimate) Under Assumptions (2), let  $\mathcal{D}$  be a space-time gradient discretisation in the sense of Definition 2.1. Let u be a solution to Scheme (14).

Then, for any  $T_0 \in (0,T]$ , denoting by k = 1, ..., N the index such that  $T_0 \in (t^{(k-1)}, t^{(k)}]$  we have

$$\int_{\Omega} B(\Pi_{\mathcal{D}}\beta(u)(\boldsymbol{x}, T_{0})) d\boldsymbol{x} + \int_{0}^{T_{0}} \int_{\Omega} \boldsymbol{a}(\boldsymbol{x}, \Pi_{\mathcal{D}}\nu(u)(\boldsymbol{x}, t), \nabla_{\mathcal{D}}\zeta(u)(\boldsymbol{x}, t)) \cdot \nabla_{\mathcal{D}}\zeta(u)(\boldsymbol{x}, t) d\boldsymbol{x} dt 
\leq \int_{\Omega} B(\Pi_{\mathcal{D}}\beta(\mathcal{I}_{\mathcal{D}}u_{\text{ini}})(\boldsymbol{x})) d\boldsymbol{x} + \int_{0}^{t^{(k)}} \int_{\Omega} f(\boldsymbol{x}, t) \Pi_{\mathcal{D}}\zeta(u)(\boldsymbol{x}, t) d\boldsymbol{x} dt.$$
(42)

Consequently, there exists  $C_1 > 0$  only depending on p,  $L_{\beta}$ ,  $C_P \geq C_{\mathcal{D}}$  (see Definition 2.5),  $C_{\text{ini}} \geq \|\Pi_{\mathcal{D}}\mathcal{I}_{\mathcal{D}}u_{\text{ini}}\|_{L^2(\Omega)}$ , f,  $\underline{a}$  and the constants  $K_0$ ,  $K_1$  and  $K_2$  in (26) such that

$$\|\Pi_{\mathcal{D}}B(\beta(u))\|_{L^{\infty}(0,T;L^{1}(\Omega))} \leq C_{1}, \ \|\nabla_{\mathcal{D}}\zeta(u)\|_{L^{p}(\Omega\times(0,T))^{d}} \leq C_{1}$$

$$and \ \|\Pi_{\mathcal{D}}\beta(u)\|_{L^{\infty}(0,T;L^{2}(\Omega))} \leq C_{1}.$$

$$(43)$$

**Proof.** By using (12) and (27) we notice that for any n = 0, ..., N-1 and any  $t \in (t^{(n)}, t^{(n+1)}]$ 

$$\Pi_{\mathcal{D}}\delta_{\mathcal{D}}\beta(u)(t)\Pi_{\mathcal{D}}\zeta(u^{(n+1)}) = \frac{1}{\check{\alpha}^{(n+\frac{1}{2})}} \left(\beta(\Pi_{\mathcal{D}}u^{(n+1)}) - \beta(u^{(n)})\right) \zeta(\Pi_{\mathcal{D}}u^{(n+1)}) 
\geq \frac{1}{\check{\alpha}^{(n+\frac{1}{2})}} \left(B(\Pi_{\mathcal{D}}\beta(u^{(n+1)})) - B(\Pi_{\mathcal{D}}\beta(u^{(n)}))\right).$$

Hence, with  $v = (\zeta(u^{(1)}), \dots, \zeta(u^{(k)}), 0, \dots, 0) \subset X_{D,0}$  in (14) we find

$$\int_{\Omega} B(\Pi_{\mathcal{D}}\beta(u)(\boldsymbol{x}, t^{(k)})) d\boldsymbol{x} + \int_{0}^{t^{(k)}} \int_{\Omega} \boldsymbol{a}(\boldsymbol{x}, \Pi_{\mathcal{D}}\nu(u)(\boldsymbol{x}, t), \nabla_{\mathcal{D}}\zeta(u)(\boldsymbol{x}, t)) \cdot \nabla_{\mathcal{D}}\zeta(u)(\boldsymbol{x}, t) d\boldsymbol{x} dt 
\leq \int_{\Omega} B(\Pi_{\mathcal{D}}\beta(u^{(0)})(\boldsymbol{x})) d\boldsymbol{x} + \int_{0}^{t^{(k)}} f(\boldsymbol{x}, t) \Pi_{\mathcal{D}}\zeta(u)(\boldsymbol{x}, t) d\boldsymbol{x} dt.$$
(44)

Equation (42) is a straightforward consequence of this estimate, of the relation  $\beta(u)(\cdot, T_0) = \beta(u)(\cdot, t^{(k)})$  (see (13)) and of the fact that the integrand involving  $\boldsymbol{a}$  is nonnegative on  $[T_0, t^{(k)}]$ . By using Young's inequality  $ab \leq \frac{1}{p}a^p + \frac{1}{p'}b^{p'}$ , we can write

$$\int_{0}^{t^{(k)}} \int_{\Omega} f(\boldsymbol{x}, t) \Pi_{\mathcal{D}} \zeta(u)(\boldsymbol{x}, t) d\boldsymbol{x} dt 
\leq \frac{2^{1/(p-1)} C_{\mathcal{D}}^{p'}}{(p\underline{a})^{1/(p-1)} p'} \|f\|_{L^{p'}(\Omega \times (0, t^{(k)}))}^{p'} + \frac{\underline{a}}{2 C_{\mathcal{D}}^{p}} \|\Pi_{\mathcal{D}} \zeta(u)\|_{L^{p}(\Omega \times (0, t^{(k)}))}^{p}$$

and the first two estimates in (43) therefore follow from (44), (26), the coercivity assumption (2f) on  $\boldsymbol{a}$  and the definition 2.5 of  $C_{\mathcal{D}}$ . The estimate on  $\Pi_{\mathcal{D}}\beta(u) = \beta(\Pi_{\mathcal{D}}u)$  in  $L^{\infty}(0,T;L^{2}(\Omega))$  is a consequence of the estimate on  $B(\beta(\Pi_{\mathcal{D}}u))$  in  $L^{\infty}(0,T;L^{1}(\Omega))$  and of (26).

Corollary 4.2 (Existence of a solution to the gradient scheme) Under Assumptions (2), if  $\mathcal{D}$  is a gradient discretisation in the sense of Definition 2.1 then there exists at least a solution to the gradient scheme (14).

**Proof.** We endow  $E = \{(u^{(n)})_{n=1,\dots,N} : u^{(n)} \in X_{\mathcal{D},0} \text{ for all } n\}$  with the dot product "·" coming from the degrees of freedom I (see Remark 2.3), and we denote by  $|\cdot|$  the corresponding norm. Let  $T: E \mapsto E$  be such that, for all  $u, v \in E$ ,

$$T(u) \cdot v = \int_0^T \int_{\Omega} \left[ \Pi_{\mathcal{D}} \delta_{\mathcal{D}} \beta(u)(\boldsymbol{x}, t) \Pi_{\mathcal{D}} v(\boldsymbol{x}, t) + \boldsymbol{a}(\boldsymbol{x}, \Pi_{\mathcal{D}} \nu(u)(\boldsymbol{x}, t), \nabla_{\mathcal{D}} \zeta(u)(\boldsymbol{x}, t)) \cdot \nabla_{\mathcal{D}} v(\boldsymbol{x}, t) \right] d\boldsymbol{x} dt,$$

where  $\delta_{\mathcal{D}}^{(\frac{1}{2})}\beta(u)$  is defined by setting  $u^{(0)} = \mathcal{I}_{\mathcal{D}}u_{\text{ini}}$ . Set  $f_E \in E$  such that, for all  $v \in E$ ,  $f_E \cdot v = \int_0^T \int_\Omega f(\boldsymbol{x},t) \Pi_{\mathcal{D}}v(\boldsymbol{x},t) d\boldsymbol{x} dt$ . A solution to (14) is an element  $u \in E$  such that  $T(u) = f_E$ . The continuity and growth properties of  $\beta$ ,  $\zeta$  and  $\boldsymbol{a}$  clearly show that T is continuous  $E \mapsto E$ , so we can prove that  $T(u) = f_E$  has has a solution by establishing that, for R large enough,  $d(T, B(R), f_E) \neq 0$  where d is the Brouwer topological degree [15] and B(R) is the open ball of radius R in E.

Following the reasoning used to prove (42), the coercivity property (2f) on a and the equivalence of all norms on E give  $C_2$  and  $C_3$  not depending on  $u \in E$  such that

$$T(u) \cdot \zeta(u) \ge \underline{a} ||\nabla_{\mathcal{D}} \zeta(u)||_{L^p(\Omega)^d}^p - ||B(\Pi_{\mathcal{D}} \beta(\mathcal{I}_{\mathcal{D}} u_{\mathrm{ini}}))||_{L^1(\Omega)} \ge C_2 |u|^p - C_3.$$

From the choice of the dot product on E and Assumption (2b) on  $\zeta$ , we have  $|\zeta(v)| \leq L_{\zeta}|v|$  and  $\zeta(v) \cdot v \geq C_4|v|^2 - C_5$ , with  $C_4 > 0$  and  $C_5$  not depending on  $v \in E$ . Let us consider the homotopy  $h(\rho, u) = \rho T(u) + (1 - \rho)u$  between T and Id, and assume that u is a solution to  $h(\rho, u) = f_E$  for some  $\rho \in [0, 1]$ . We have if  $|u| \geq 1$ 

$$|f_E|L_{\zeta}|u| \ge f_E \cdot \zeta(u) = \rho T(u) \cdot \zeta(u) + (1-\rho)u \cdot \zeta(u)$$
  
 
$$\ge \rho C_2|u|^p - \rho C_3 + (1-\rho)C_4|u|^2 - (1-\rho)C_5 \ge \min(C_2, C_4)|u|^{\min(p,2)} - C_3 - C_5.$$

Hence, if we select R > 1 such that  $|f_E|L_{\zeta}R < \min(C_2, C_4)R^{\min(p,2)} - C_3 - C_5$ , which is possible since  $\min(p,2) > 1$ , no solution to  $h(\rho,u) = f_E$  can lie on  $\partial B(R)$ . The invariance by homotopy of the topological degree then gives  $d(T,B(R),f_E) = d(\operatorname{Id},B(R),f_E)$ , and this last degree is equal to 1 if we select R such that  $f_E \in B(R)$ . The proof is complete.

## Lemma 4.3 (Estimate on the dual semi-norm of the discrete time derivative)

Under Assumptions (2), let  $\mathcal{D}$  be a space-time gradient discretisation in the sense of Definition 2.1. Let u be a solution to Scheme (14). Then there exists  $C_6$  only depending on p,  $L_{\beta}$ ,  $C_P \geq C_{\mathcal{D}}$ ,  $C_{\text{ini}} \geq \|\Pi_{\mathcal{D}} I_{\mathcal{D}} u_{\text{ini}}\|_{L^2(\Omega)}$ , f,  $\underline{a}$ ,  $\mu$ ,  $\overline{a}$ , T and the constants  $K_0$ ,  $K_1$  and  $K_2$  in (26) such that

$$\int_0^T |\delta_{\mathcal{D}}\beta(u)(t)|_{\star,\mathcal{D}}^{p'} dt \le C_6.$$
(45)

**Proof.** Let us take a generic  $v = (v^{(n)})_{n=1,...,N} \subset X_{\mathcal{D},0}$  as a test function in Scheme (14). We have, thanks to Assumption (2h) on  $\boldsymbol{a}$ ,

$$\int_{0}^{T} \int_{\Omega} \Pi_{\mathcal{D}} \delta_{\mathcal{D}} \beta(u)(\boldsymbol{x}, t) \Pi_{\mathcal{D}} v(\boldsymbol{x}, t) d\boldsymbol{x} dt \leq \int_{0}^{T} \int_{\Omega} (\overline{a}(\boldsymbol{x}) + \mu |\nabla_{\mathcal{D}} \zeta(u)(\boldsymbol{x}, t)|^{p-1}) |\nabla_{\mathcal{D}} v(\boldsymbol{x}, t)| d\boldsymbol{x} dt + \int_{0}^{T} \int_{\Omega} f(\boldsymbol{x}, t) \Pi_{\mathcal{D}} v(\boldsymbol{x}, t) d\boldsymbol{x} dt.$$

Using Hölder's inequality, Definition 2.5 and Estimates (43), this leads to the existence of  $C_7 > 0$  only depending on p,  $L_{\beta}$ ,  $C_P$ ,  $C_{\rm ini}$ , f,  $\underline{a}$ ,  $\overline{a}$ ,  $\mu$  and  $K_0$ ,  $K_1$  and  $K_2$  such that

$$\int_0^T \int_{\Omega} \Pi_{\mathcal{D}} \delta_{\mathcal{D}} \beta(u)(\boldsymbol{x}, t) \Pi_{\mathcal{D}} v(\boldsymbol{x}, t) d\boldsymbol{x} dt \leq C_7 \|\nabla_{\mathcal{D}} v\|_{L^p(0, T; L^p(\Omega))^d}.$$

The proof of (45) is completed by selecting  $v = (|\delta_{\mathcal{D}}^{(n+\frac{1}{2})}\beta(u)|_{\star,\mathcal{D}}^{p'-1}z^{(n)})_{n=1,\ldots,N}$  with  $(z^{(n)})_{n=1,\ldots,N} \subset X_{\mathcal{D},0}$  such that, for any  $n=1,\ldots,N,$   $z^{(n)}$  realises the supremum in (11) with  $w=\delta_{\mathcal{D}}^{(n+\frac{1}{2})}\beta(u)$ .

## Lemma 4.4 (Estimate on the time translates of $\nu(u)$ )

Under Assumptions (2), let  $\mathcal{D}$  be a space-time gradient discretisation in the sense of Definition 2.1. Let u be a solution to Scheme (14). Then there exists  $C_8$  only depending on p,  $L_{\beta}$ ,  $L_{\zeta}$ ,  $C_P \geq C_{\mathcal{D}}$ ,  $C_{\text{ini}} \geq \|\Pi_{\mathcal{D}} I_{\mathcal{D}} u_{\text{ini}}\|_{L^2(\Omega)}$ , f,  $\underline{a}$ ,  $\mu$ ,  $\overline{a}$ , T and  $K_0$ ,  $K_1$  and  $K_2$  in (26) such that

$$\|\Pi_{\mathcal{D}}\nu(u)(\cdot,\cdot+\tau) - \Pi_{\mathcal{D}}\nu(u)(\cdot,\cdot)\|_{L^2(\Omega\times(0,T-\tau))}^2 \le C_8(\tau+\delta t), \quad \forall \tau \in (0,T).$$
(46)

**Proof.** Let  $\tau \in (0,T)$ . Thanks to (24), we can write

$$\int_{\Omega \times (0, T - \tau)} \left( \Pi_{\mathcal{D}} \nu(u)(\boldsymbol{x}, t + \tau) - \Pi_{\mathcal{D}} \nu(u)(\boldsymbol{x}, t) \right)^{2} d\boldsymbol{x} dt \le L_{\beta} L_{\zeta} \int_{0}^{T - \tau} A(t) dt, \tag{47}$$

where

$$A(t) = \int_{\Omega} \left( \Pi_{\mathcal{D}} \zeta(u)(\boldsymbol{x}, t + \tau) - \Pi_{\mathcal{D}} \zeta(u)(\boldsymbol{x}, t) \right) \left( \Pi_{\mathcal{D}} \beta(u)(\boldsymbol{x}, t + \tau) - \Pi_{\mathcal{D}} \beta(u)(\boldsymbol{x}, t) \right) d\boldsymbol{x}.$$

For  $s \in (0, T)$ , we define  $n(s) \in \{0, \dots, N-1\}$  such that  $t^{(n(s))} < s \le t^{(n(s)+1)}$ . Taking  $t \in (0, T-\tau)$ , we may write

$$A(t) = \int_{\Omega} \left( \Pi_{\mathcal{D}} \zeta(u^{(n(t+\tau)+1)})(\boldsymbol{x}) - \Pi_{\mathcal{D}} \zeta(u^{(n(t)+1)})(\boldsymbol{x}) \right) \left( \sum_{n=n(t)+1}^{n(t+\tau)} \delta^{(n+\frac{1}{2})} \Pi_{\mathcal{D}} \delta^{(n+\frac{1}{2})}_{\mathcal{D}} \beta(u)(\boldsymbol{x}) \right) d\boldsymbol{x}.$$

We then use the definition (11) of the discrete dual semi-norm to infer

$$A(t) \le \sum_{n=n(t)+1}^{n(t+\tau)} \delta t^{(n+\frac{1}{2})} \left\| \left| \nabla_{\mathcal{D}} \left[ \zeta(u^{(n(t+\tau)+1)}) - \zeta(u^{(n(t)+1)}) \right] \right| \right|_{L^{p}(\Omega)^{d}} \left| \delta_{\mathcal{D}}^{(n+\frac{1}{2})} \beta(u) \right|_{\star,\mathcal{D}}.$$
(48)

We apply the triangular inequality on the first norm in this right-hand side, Young's inequality and we integrate over  $t \in (0, T - \tau)$  to get

$$\int_{0}^{T-\tau} A(t)dt \le \mathcal{A}_{\tau} + \mathcal{A}_{0} + \mathcal{B}$$
(49)

with, for s = 0 or  $s = \tau$ ,

$$\mathcal{A}_{s} = \frac{1}{p} \int_{0}^{T-\tau} \sum_{n=n(t)+1}^{n(t+\tau)} \delta t^{(n+\frac{1}{2})} ||\nabla_{\mathcal{D}}\zeta(u^{(n(t+s)+1)})||_{L^{p}(\Omega)^{d}}^{p} dt \le \frac{C_{1}^{p}}{p} (\tau + \delta t)$$
 (50)

and

$$\mathcal{B} = \frac{2}{p'} \int_0^{T-\tau} \sum_{n=n(t)+1}^{n(t+\tau)} \delta t^{(n+\frac{1}{2})} |\delta_{\mathcal{D}}^{(n+\frac{1}{2})} \beta(u)|_{\star,\mathcal{D}}^{p'} dt \le \frac{2C_6}{p'} \tau.$$
 (51)

In (50), the quantity  $\mathcal{A}_s$  has been estimated by using (84) in Lemma 6.6 and the estimate on  $\nabla_{\mathcal{D}}\zeta(u)$  in (43). In (51),  $\mathcal{B}$  has been estimated by applying (83) in Lemma 6.6 and by using the bound (45) on  $\int_0^T |\delta_{\mathcal{D}}\beta(u)(t)|_{\star,\mathcal{D}}^{p'} dt$ . The proof is completed by gathering (47), (49), (50) and (51).

#### 4.2 Proof of Theorem 2.12

Step 1 Application of compactness results.

Thanks to Theorem 3.1 and Estimates (43) and (45), we first extract a subsequence such that  $(\Pi_{\mathcal{D}_m}\beta(u_m))_{m\in\mathbb{N}}$  converges weakly in  $L^2(\Omega)$  uniformly on [0,T] (in the sense of Definition 2.11) to some function  $\overline{\beta}\in C([0,T];L^2(\Omega)-w)$  which satisfies  $\overline{\beta}(\cdot,0)=\beta(u_{\rm ini})$  in  $L^2(\Omega)$ . Using again Estimates (43) and applying Lemma 2.10, we extract a further subsequence such that, for some  $\overline{\zeta}\in L^p(0,T;W_0^{1,p}(\Omega))$ ,  $\Pi_{\mathcal{D}_m}\zeta(u_m)\to \overline{\zeta}$  weakly in  $L^p(\Omega\times(0,T))$  and  $\nabla_{\mathcal{D}_m}\zeta(u_m)\to \nabla\overline{\zeta}$  weakly in  $L^p(\Omega\times(0,T))^d$ . Estimates (43), Definition 2.5 and the growth assumption (2b) on  $\zeta$  show that  $(\Pi_{\mathcal{D}_m}u_m)_{m\in\mathbb{N}}$  is bounded in  $L^p(\Omega\times(0,T))$  and we can therefore assume, up to a subsequence, that it converges weakly to some  $\overline{u}$  in this space.

We then prove, by means of the Kolmogorov theorem, that  $(\Pi_{\mathcal{D}_m}\nu(u_m))_{m\in\mathbb{N}}$  is relatively compact in  $L^1(\Omega \times (0,T))$ . We first remark that  $|\nu(a) - \nu(b)| \leq L_{\beta}|\zeta(a) - \zeta(b)|$ , which implies, using Estimate (43) and Definition 2.9 with  $v = \zeta(u_m)$ ,

$$||\Pi_{\mathcal{D}_m}\nu(u_m)(\cdot + \boldsymbol{\xi}, \cdot) - \Pi_{\mathcal{D}_m}\nu(u_m)(\cdot, \cdot)||_{L^p(\mathbb{R}^d \times (0,T))} \le L_\beta C_1 T_{\mathcal{D}_m}(\boldsymbol{\xi})$$
(52)

where  $\Pi_{\mathcal{D}_m}\nu(u_m)$  has been extended by 0 outside  $\Omega$ , and  $\lim_{\xi\to 0}\sup_{m\in\mathbb{N}}T_{\mathcal{D}_m}(\xi)=0$ . This takes care of the space translates. Let us now turn to the time translates. Invoking Lemma 4.4 and, to control the time translates at both ends of [0,T], the fact that  $\Pi_{\mathcal{D}_m}\beta(u_m)$  – and therefore also  $\Pi_{\mathcal{D}_m}\nu(u_m)$  since  $|\nu| \leq L_{\zeta}|\beta|$  – remains bounded in  $L^{\infty}(0,T;L^2(\Omega))$ , we can write for any  $M \in \mathbb{N}$ 

$$\sup_{m\in\mathbb{N}} ||\Pi_{\mathcal{D}_m}\nu(u_m)(\cdot,\cdot+\tau) - \Pi_{\mathcal{D}_m}\nu(u_m)(\cdot,\cdot)||^2_{L^2(\Omega\times(0,T))}$$

$$\leq \max\left(\max_{m\leq M}||\Pi_{\mathcal{D}_m}\nu(u_m)(\cdot,\cdot+\tau) - \Pi_{\mathcal{D}_m}\nu(u_m)(\cdot,\cdot)||_{L^2(\Omega\times(0,T))}^2; C_9\left(\tau + \sup_{m>M}\delta t_m\right)\right), \quad (53)$$

where  $C_9$  does not depend on m or  $\tau$ , and the functions have been extended by 0 outside (0,T). Since each  $||\Pi_{\mathcal{D}_m}\nu(u_m)(\cdot,\cdot+\tau)-\Pi_{\mathcal{D}_m}\nu(u_m)||^2_{L^2(\Omega\times(0,T))}$  tends to 0 as  $\tau\to 0$  and since  $\delta t_m\to 0$  as  $m\to\infty$ , taking in that order the limsup as  $\tau\to0$  and the limit as  $M\to\infty$  of (53) shows that the left-hand side of this inequality tends to 0 as  $\tau \to 0$ , as required. Hence, Kolmogorov's theorem shows that, up to extraction of another subsequence,  $\Pi_{\mathcal{D}_m}\nu(u_m)\to \overline{\nu}$  in  $L^1(\Omega\times(0,T))$ .

Let us now identify these limits  $\overline{\beta}$ ,  $\overline{\zeta}$  and  $\overline{\nu}$ . Under the first case in the structural hypothesis (17), we have  $\beta = \mathrm{Id}$ , and therefore  $\overline{\beta} = \overline{u} = \beta(\overline{u})$  and  $\nu = \zeta$ . The strong convergence of  $\Pi_{\mathcal{D}_m}\nu(u_m)=\Pi_{\mathcal{D}_m}\zeta(u_m)$  to  $\overline{\nu}=\overline{\zeta}$  allows us to apply Lemma 3.5 to see that  $\overline{\zeta}=\zeta(\overline{u})$  and  $\overline{\nu} = \nu(\overline{u})$ . Exchanging the roles of  $\beta$  and  $\zeta$ , we see that  $\overline{\beta} = \beta(\overline{u})$ ,  $\overline{\zeta} = \zeta(\overline{u})$  and  $\overline{\nu} = \nu(\overline{u})$  still hold in the second case of (17). We notice that this is the only place where we use this structural assumption (17) on  $\beta, \zeta$ .

Using the growth assumption (2h) on a and Estimates (43), upon extraction of another subsequence we can also assume that  $\mathbf{a}(\cdot, \Pi_{\mathcal{D}_m} \nu(u_m), \nabla_{\mathcal{D}_m} \zeta(u_m))$  has a weak limit in  $L^{p'}(\Omega \times (0,T))^d$ , which we denote by A.

Finally, for any  $T_0 \in [0, T]$ , since  $\Pi_{\mathcal{D}_m} \beta(u_m(\cdot, T_0)) \to \beta(\overline{u})(\cdot, T_0)$  weakly in  $L^2(\Omega)$ , Lemma 3.4 gives

$$\int_{\Omega} B(\beta(\overline{u})(\boldsymbol{x}, T_0)) d\boldsymbol{x} \le \liminf_{m \to \infty} \int_{\Omega} B(\beta(\Pi_{\mathcal{D}_m} u_m)(\boldsymbol{x}, T_0)) d\boldsymbol{x}.$$
 (54)

With (43), this shows that  $B(\beta(\overline{u})) \in L^{\infty}(0,T;L^{1}(\Omega))$ .

**Step 2** Passing to the limit in the scheme.

We drop the indices m for legibility reasons. Let  $\varphi \in C^1_c(-\infty,T)$  and let  $w \in W^{1,p}_0(\Omega) \cap L^2(\Omega)$ . We introduce  $v = (\varphi(t^{(n-1)})P_{\mathcal{D}}w)_{n=1,...,N}$  as a test function in (14), with  $P_{\mathcal{D}}$  defined by (21). We get  $T_1^{(m)} + T_2^{(m)} = T_3^{(m)}$  with

$$T_1^{(m)} = \sum_{n=0}^{N-1} \varphi(t^{(n)}) \delta t^{(n+\frac{1}{2})} \int_{\Omega} \Pi_{\mathcal{D}} \delta_{\mathcal{D}}^{(n+\frac{1}{2})} \beta(u)(\boldsymbol{x}) \Pi_{\mathcal{D}} P_{\mathcal{D}} w(\boldsymbol{x}) d\boldsymbol{x},$$

$$T_2^{(m)} = \sum_{n=0}^{N-1} \varphi(t^{(n)}) \delta t^{(n+\frac{1}{2})} \int_{\Omega} \boldsymbol{a} \left( \boldsymbol{x}, \Pi_{\mathcal{D}} \nu(u^{(n+1)}), \nabla_{\mathcal{D}} \zeta(u^{(n+1)})(\boldsymbol{x}) \right) \cdot \nabla_{\mathcal{D}} P_{\mathcal{D}} w(\boldsymbol{x}) \mathrm{d}\boldsymbol{x},$$

and

$$T_3^{(m)} = \sum_{n=0}^{N-1} \varphi(t^{(n)}) \int_{t^{(n)}}^{t^{(n+1)}} \int_{\Omega} f(\boldsymbol{x}, t) \Pi_{\mathcal{D}} P_{\mathcal{D}} w(\boldsymbol{x}) d\boldsymbol{x} dt.$$

Using discrete integrate-by-parts to transform the terms  $\varphi(t^{(n)})(\Pi_{\mathcal{D}}\beta(u^{(n+1)}) - \Pi_{\mathcal{D}}\beta(u^{(n)}))$  appearing in  $T_1^{(m)}$  into  $(\varphi(t^{(n)}) - \varphi(t^{(n+1)}))\Pi_{\mathcal{D}}\beta(u^{(n+1)})$ , we have

$$T_1^{(m)} = -\int_0^T \varphi'(t) \int_{\Omega} \Pi_{\mathcal{D}} \beta(u)(\boldsymbol{x}, t) \Pi_{\mathcal{D}} P_{\mathcal{D}} w(\boldsymbol{x}) d\boldsymbol{x} dt - \varphi(0) \int_{\Omega} \Pi_{\mathcal{D}} \beta(u^{(0)})(\boldsymbol{x}) \Pi_{\mathcal{D}} P_{\mathcal{D}} w(\boldsymbol{x}) d\boldsymbol{x}.$$

Setting  $\varphi_{\mathcal{D}}(t) = \varphi(t^{(n)})$  for  $t \in (t^{(n)}, t^{(n+1)})$ , we have

$$T_{2}^{(m)} = \int_{0}^{T} \varphi_{\mathcal{D}}(t) \int_{\Omega} \boldsymbol{a} \left( \boldsymbol{x}, \Pi_{\mathcal{D}} \nu(u)(\boldsymbol{x}, t), \nabla_{\mathcal{D}} \zeta(u)(\boldsymbol{x}, t) \right) \cdot \nabla_{\mathcal{D}} P_{\mathcal{D}} w(\boldsymbol{x}) d\boldsymbol{x} dt$$

$$T_{3}^{(m)} = \int_{0}^{T} \varphi_{\mathcal{D}}(t) \int_{\Omega} f(\boldsymbol{x}, t) \Pi_{\mathcal{D}} P_{\mathcal{D}} w(\boldsymbol{x}) d\boldsymbol{x} dt.$$

Since  $\varphi_{\mathcal{D}} \to \varphi$  uniformly on [0,T],  $\Pi_{\mathcal{D}}P_{\mathcal{D}}w \to w$  in  $L^p(\Omega) \cap L^2(\Omega)$  and  $\nabla_{\mathcal{D}}P_{\mathcal{D}}w \to \nabla w$  in  $L^p(\Omega)^d$ , we may let  $m \to \infty$  in  $T_1^{(m)} + T_2^{(m)} = T_3^{(m)}$  to see that  $\overline{w}$  satisfies

$$\begin{cases}
\overline{u} \in L^{p}(\Omega \times (0,T)), \ \zeta(\overline{u}) \in L^{p}(0,T;W_{0}^{1,p}(\Omega)), \ B(\beta(\overline{u})) \in L^{\infty}(0,T;L^{1}(\Omega)), \\
\beta(\overline{u}) \in C([0,T];L^{2}(\Omega)\text{-w}), \ \beta(\overline{u})(\cdot,0) = \beta(u_{\text{ini}}), \\
-\int_{0}^{T} \varphi'(t) \int_{\Omega} \beta(\overline{u}(\boldsymbol{x},t))w(\boldsymbol{x})d\boldsymbol{x}dt - \varphi(0) \int_{\Omega} \beta(u_{\text{ini}}(\boldsymbol{x}))w(\boldsymbol{x})d\boldsymbol{x} \\
+\int_{0}^{T} \varphi(t) \int_{\Omega} \boldsymbol{A}(\boldsymbol{x},t) \cdot \nabla w(\boldsymbol{x})d\boldsymbol{x}dt = \int_{0}^{T} \varphi(t) \int_{\Omega} f(\boldsymbol{x},t)w(\boldsymbol{x})d\boldsymbol{x}dt, \\
\forall w \in W_{0}^{1,p}(\Omega) \cap L^{2}(\Omega), \ \forall \varphi \in C_{c}^{\infty}(-\infty,T).
\end{cases} (55)$$

Note that the regularity properties on  $\overline{u}$ ,  $\zeta(\overline{u})$ ,  $\beta(\overline{u})$  and  $B(\beta(\overline{u}))$  have been established in Step 1. Linear combinations of this relation show that (55) also holds with  $\varphi(t)w(x)$  replaced by a tensorial functions in  $C_c^{\infty}(\Omega \times (0,T))$ . This proves that  $\partial_t \beta(\overline{u}) \in L^{p'}(0,T;W^{-1,p'}(\Omega))$  (see Remark 1.1). Using the density of tensorial functions in  $L^p(0,T;W^{1,p}(\Omega))$  [18], we then see that  $\overline{u}$  satisfies

$$\int_{0}^{T} \langle \partial_{t} \beta(\overline{u})(\cdot, t), \overline{v}(\cdot, t) \rangle_{W^{-1, p'}, W_{0}^{1, p}} dt 
+ \int_{0}^{T} \int_{\Omega} \mathbf{A}(\mathbf{x}, t) \cdot \nabla \overline{v}(\mathbf{x}, t) d\mathbf{x} dt = \int_{0}^{T} \int_{\Omega} f(\mathbf{x}, t) \overline{v}(\mathbf{x}, t) d\mathbf{x} dt, \quad \forall \overline{v} \in L^{p}(0, T; W_{0}^{1, p}(\Omega)).$$
(56)

**Step 3** Proof that  $\overline{u}$  is a solution to (4).

It only remains to show that

$$\mathbf{A}(\mathbf{x},t) = \mathbf{a}(\mathbf{x},\nu(\overline{u})(\mathbf{x},t),\nabla\zeta(\overline{u})(\mathbf{x},t)) \text{ for a.e. } (\mathbf{x},t) \in \Omega \times (0,T).$$
 (57)

We take  $T_0 \in [0,T]$ , write (42) with  $\mathcal{D} = \mathcal{D}_m$  and take the lim sup as  $m \to \infty$ . We notice that the  $t^{(k)} =: T_m$  from Lemma 4.1 converges to  $T_0$  as  $m \to \infty$ . Hence, by using the convergence  $\Pi_{\mathcal{D}_m} \mathcal{I}_{\mathcal{D}_m} u_{\text{ini}} \to u_{\text{ini}}$  in  $L^2(\Omega)$  (consistency of  $(\mathcal{D}_m)_{m \in \mathbb{N}}$ ), and the continuity and quadratic growth of  $B \circ \beta$  (upper bound in (26)), we obtain

$$\limsup_{m \to \infty} \int_{0}^{T_{0}} \int_{\Omega} \boldsymbol{a}(\boldsymbol{x}, \Pi_{\mathcal{D}_{m}} \nu(u_{m})(\boldsymbol{x}, t), \nabla_{\mathcal{D}_{m}} \zeta(u_{m})(\boldsymbol{x}, t)) \cdot \nabla_{\mathcal{D}_{m}} \zeta(u_{m})(\boldsymbol{x}, t) d\boldsymbol{x} dt$$

$$\leq \int_{\Omega} B(\beta(u_{\text{ini}})(\boldsymbol{x})) d\boldsymbol{x} + \int_{0}^{T_{0}} \int_{\Omega} f(\boldsymbol{x}, t) \zeta(\overline{u})(\boldsymbol{x}, t) d\boldsymbol{x} dt - \liminf_{m \to \infty} \int_{\Omega} B(\beta(\Pi_{\mathcal{D}_{m}} u_{m})(\boldsymbol{x}, T_{0})) d\boldsymbol{x}. \quad (58)$$

We take  $\overline{v} = \zeta(\overline{u})\mathbf{1}_{[0,T_0]}$  in (56) and apply Lemma 3.6 to get

$$\int_{\Omega} B(\beta(\overline{u})(\boldsymbol{x}, T_0)) d\boldsymbol{x} - \int_{\Omega} B(\beta(\overline{u})(\boldsymbol{x}, 0)) d\boldsymbol{x} 
+ \int_{0}^{T_0} \int_{\Omega} \boldsymbol{A}(\boldsymbol{x}, t) \cdot \nabla \zeta(\overline{u})(\boldsymbol{x}, t) d\boldsymbol{x} dt = \int_{0}^{T_0} \int_{\Omega} f(\boldsymbol{x}, t) \zeta(\overline{u})(\boldsymbol{x}, t) d\boldsymbol{x} dt.$$

This relation, combined with (58) and using (54), shows that

$$\lim_{m \to \infty} \int_{0}^{T_{0}} \int_{\Omega} \boldsymbol{a}(\boldsymbol{x}, \Pi_{\mathcal{D}_{m}} \nu(u_{m})(\boldsymbol{x}, t), \nabla_{\mathcal{D}_{m}} \zeta(u_{m})(\boldsymbol{x}, t)) \cdot \nabla_{\mathcal{D}_{m}} \zeta(u_{m})(\boldsymbol{x}, t) d\boldsymbol{x} dt \\
\leq \int_{0}^{T_{0}} \int_{\Omega} \boldsymbol{A}(\boldsymbol{x}, t) \cdot \nabla \zeta(\overline{u})(\boldsymbol{x}, t) d\boldsymbol{x} dt. \quad (59)$$

It is now possible to apply Minty's trick. Consider for  $G \in L^p(\Omega \times (0,T))^d$  the following relation, stemming from the monotony (2g) of a:

$$\int_{0}^{T_{0}} \int_{\Omega} \left[ \boldsymbol{a}(\cdot, \Pi_{\mathcal{D}_{m}} \nu(u_{m}), \nabla_{\mathcal{D}_{m}} \zeta(u_{m})) - \boldsymbol{a}(\cdot, \Pi_{\mathcal{D}_{m}} \nu(u_{m}), \boldsymbol{G}) \right] \cdot \left[ \nabla_{\mathcal{D}_{m}} \zeta(u_{m}) - \boldsymbol{G} \right] d\boldsymbol{x} dt \ge 0.$$
 (60)

By strong convergence of  $\Pi_{\mathcal{D}_m}\nu(u_m)$  to  $\nu(\overline{u})$  in  $L^1(\Omega\times(0,T))$  and Assumptions (2e), (2h) on  $\boldsymbol{a}$ , we see that  $\boldsymbol{a}(\cdot,\Pi_{\mathcal{D}_m}\nu(u_m),\boldsymbol{G})\to\boldsymbol{a}(\cdot,\nu(\overline{u}),\boldsymbol{G})$  strongly in  $L^{p'}(\Omega\times(0,T))^d$ . The development of (60) gives a sum of four terms, the first one being the integral in the left-hand side of (59) and the other three being integrals of products of weakly and strongly converging sequences. We can thus take the  $\limsup$  of (60) with  $T_0=T$  to find

$$\int_0^T \int_{\Omega} \left[ \boldsymbol{A}(\boldsymbol{x},t) - \boldsymbol{a}(\boldsymbol{x},\nu(\overline{u})(\boldsymbol{x},t),\boldsymbol{G}(\boldsymbol{x},t)) \right] \cdot \left[ \nabla \zeta(\overline{u})(\boldsymbol{x},t) - \boldsymbol{G}(\boldsymbol{x},t) \right] \mathrm{d}\boldsymbol{x} \mathrm{d}t \geq 0.$$

Application of Minty's method [47] (i.e. taking  $G = \nabla \zeta(\overline{u}) + r\varphi$  for  $\varphi \in L^p(\Omega \times (0,T))^d$  and letting  $r \to 0$ ) then shows that (57) holds and concludes the proof that  $\overline{u}$  satisfies (4).

#### 4.3 Proof of Theorem 2.16

Let  $T_0 \in [0,T]$  and  $(T_m)_{m\geq 1}$  be a sequence in [0,T] that converges to  $T_0$ . By setting  $T_0 = T_m$  and  $G = \nabla \zeta(\overline{u})$  in the developed form of (60), by taking the infimum limit (thanks to the strong convergence of  $\mathbf{a}(\cdot, \Pi_{\mathcal{D}_m}\nu(u_m), \nabla\zeta(\overline{u}))$ ) and by using (57), we find

$$\lim_{m \to \infty} \int_{0}^{T_{m}} \int_{\Omega} \boldsymbol{a}(\boldsymbol{x}, \Pi_{\mathcal{D}_{m}} \nu(u_{m})(\boldsymbol{x}, t), \nabla_{\mathcal{D}_{m}} \zeta(u_{m})(\boldsymbol{x}, t)) \cdot \nabla_{\mathcal{D}_{m}} \zeta(u_{m})(\boldsymbol{x}, t) d\boldsymbol{x} dt$$

$$\geq \int_{0}^{T_{0}} \int_{\Omega} \boldsymbol{a}(\boldsymbol{x}, \nu(\overline{u})(\boldsymbol{x}, t), \nabla \zeta(\overline{u})(\boldsymbol{x}, t)) \cdot \nabla \zeta(\overline{u})(\boldsymbol{x}, t) d\boldsymbol{x} dt. \quad (61)$$

We then write (42) with  $T_m$  instead of  $T_0$  and we take the  $\limsup$  as  $m \to \infty$ . We notice that the  $t^{(k)}$  such that  $T_m \in (t^{(k-1)}, t^{(k)}]$  converges to  $T_0$  as  $m \to \infty$ . Thanks to (61) and (39) we obtain

$$\limsup_{m \to \infty} \int_{\Omega} B(\beta(\Pi_{\mathcal{D}_m} u_m(\boldsymbol{x}, T_m))) d\boldsymbol{x} \le \int_{\Omega} B(\beta(\overline{u})(\boldsymbol{x}, T_0)) d\boldsymbol{x}.$$
 (62)

By Lemma 6.4, the uniform-in-time weak convergence of  $\beta(\Pi_{\mathcal{D}_m}u_m)$  to  $\beta(\bar{u})$  and the continuity of  $\beta(\bar{u}):[0,T]\to L^2(\Omega)$ -w, we have  $\beta(\Pi_{\mathcal{D}_m}u_m)(T_m)\to\beta(\bar{u})(T_0)$  weakly in  $L^2(\Omega)$  as  $m\to\infty$ .

Therefore, for any  $(s_m)_{m\in\mathbb{N}}$  converging to  $T_0$ ,  $\frac{1}{2}(\beta(\Pi_{\mathcal{D}_m}u_m(T_m)) + \beta(\overline{u})(s_m)) \to \beta(\overline{u})(T_0)$  weakly in  $L^2(\Omega)$  as  $m \to \infty$  and Lemma 3.4 gives, by convexity of B,

$$\int_{\Omega} B(\beta(\overline{u})(\boldsymbol{x}, T_0)) d\boldsymbol{x} \le \liminf_{m \to \infty} \int_{\Omega} B\left(\frac{\beta(\Pi_{\mathcal{D}_m} u_m(\boldsymbol{x}, T_m)) + \beta(\overline{u})(\boldsymbol{x}, s_m)}{2}\right) d\boldsymbol{x}. \tag{63}$$

Property (28) of B and the two inequalities (62) and (63) allow us to conclude the proof. Let  $(s_m)_{m\in\mathbb{N}}$  be a sequence in  $\mathcal{T}$  (see proof of Corollary 3.8) that converges to  $T_0$ . Then  $\nu(\overline{u}(\cdot, s_m)) \to \nu(\overline{u})(\cdot, T_0)$  in  $L^2(\Omega)$  as  $m \to \infty$ . Using (28), we get

$$\begin{split} \|\nu(\Pi_{\mathcal{D}_{m}}u_{m}(\cdot,T_{m})) - \nu(\overline{u})(\cdot,T_{0})\|_{L^{2}(\Omega)}^{2} \\ &\leq 2\|\nu(\Pi_{\mathcal{D}_{m}}u_{m}(\cdot,T_{m})) - \nu(\overline{u}(\cdot,s_{m}))\|_{L^{2}(\Omega)}^{2} + 2\|\nu(\overline{u}(\cdot,s_{m})) - \nu(\overline{u})(\cdot,T_{0})\|_{L^{2}(\Omega)}^{2} \\ &\leq 8L_{\beta}L_{\zeta} \int_{\Omega} \left[B(\beta(\Pi_{\mathcal{D}_{m}}u_{m}(\boldsymbol{x},T_{m}))) + B(\beta(\overline{u}(\boldsymbol{x},s_{m})))\right] d\boldsymbol{x} \\ &- 16L_{\beta}L_{\zeta} \int_{\Omega} B\left(\frac{\beta(\Pi_{\mathcal{D}_{m}}u_{m}(\boldsymbol{x},T_{m})) + \beta(\overline{u}(\boldsymbol{x},s_{m}))}{2}\right) d\boldsymbol{x} \\ &+ 2\|\nu(\overline{u}(\cdot,s_{m})) - \nu(\overline{u})(\cdot,T_{0})\|_{L^{2}(\Omega)}^{2}. \end{split}$$

We then take the  $\limsup$  as  $m \to \infty$  of this expression. Thanks to (62) and the continuity of  $t \in [0,T] \mapsto \int_{\Omega} B(\beta(\overline{u})(\boldsymbol{x},t)) \mathrm{d}\boldsymbol{x} \in [0,\infty)$  (see Corollary 3.8), the first term in the right-hand side has a finite  $\limsup$ , bounded above by  $16L_{\beta}L_{\zeta}\int_{\Omega} B(\beta(\overline{u})(\boldsymbol{x},T_0)) \mathrm{d}\boldsymbol{x}$ . We can therefore split the  $\limsup$  of this right-hand side without risking writing  $\infty - \infty$  and we get, thanks to (63),

$$\limsup_{m \to \infty} \|\nu(\Pi_{\mathcal{D}_m} u_m(\cdot, T_m)) - \nu(\overline{u})(\cdot, T_0)\|_{L^2(\Omega)}^2 \le 0.$$

Thus,  $\nu(\Pi_{\mathcal{D}_m}u_m(\cdot,T_m)) \to \nu(\overline{u})(T_0)$  strongly in  $L^2(\Omega)$ . By Lemma 6.4 and the continuity of  $\nu(\overline{u}): [0,T] \mapsto L^2(\Omega)$  stated in Corollary 3.8, this concludes the proof of the convergence of  $\nu(\Pi_{\mathcal{D}_m}u_m)$  to  $\nu(\overline{u})$  in  $L^{\infty}(0,T;L^2(\Omega))$ .

Remark 4.5 Since  $\beta(\Pi_{\mathcal{D}_m}u_m)(T_m) \to \beta(\bar{u})(T_0)$  weakly in  $L^2(\Omega)$  as  $m \to \infty$ , Lemma 3.4 shows that  $\int_{\Omega} B(\beta(\bar{u})(\boldsymbol{x}, T_0)) d\boldsymbol{x} \leq \liminf_{m \to \infty} \int_{\Omega} B(\beta(\Pi_{\mathcal{D}_m}u_m)(\boldsymbol{x}, T_m)) d\boldsymbol{x}$ . Combined with (62), this gives

$$\lim_{m \to \infty} \int_{\Omega} B(\beta(\Pi_{\mathcal{D}_m} u_m(\boldsymbol{x}, T_m))) d\boldsymbol{x} = \int_{\Omega} B(\beta(\overline{u})(\boldsymbol{x}, T_0)) d\boldsymbol{x}.$$
 (64)

Item 1 in Corollary 3.8 and Lemma 6.4 therefore show that the functions  $\int_{\Omega} B(\beta(\Pi_{\mathcal{D}_m} u_m(\boldsymbol{x},\cdot))) d\boldsymbol{x}$  converges uniformly on [0,T] to  $\int_{\Omega} B(\beta(\overline{u})(\boldsymbol{x},\cdot)) d\boldsymbol{x}$ .

## 4.4 Proof of Theorem 2.18

By taking the lim sup as  $m \to \infty$  of (42) for  $u_m$  with  $T_0 = T$ , and by using (64) (with  $T_m \equiv T$ ) and the continuous integration-by-parts formula (39), we find

$$\limsup_{m \to \infty} \int_0^T \int_{\Omega} \boldsymbol{a}(\boldsymbol{x}, \Pi_{\mathcal{D}_m} \nu(u_m)(\boldsymbol{x}, t), \nabla_{\mathcal{D}_m} \zeta(u_m)(\boldsymbol{x}, t)) \cdot \nabla_{\mathcal{D}_m} \zeta(u_m)(\boldsymbol{x}, t) d\boldsymbol{x} dt \\
\leq \int_0^{T_0} \int_{\Omega} \boldsymbol{a}(\boldsymbol{x}, \nu(\overline{u})(\boldsymbol{x}, t), \nabla \zeta(\overline{u})(\boldsymbol{x}, t)) \cdot \nabla \zeta(\overline{u})(\boldsymbol{x}, t) d\boldsymbol{x} dt.$$

Combined with (61), this shows that

$$\lim_{m \to \infty} \int_{0}^{T} \int_{\Omega} \boldsymbol{a}(\boldsymbol{x}, \Pi_{\mathcal{D}_{m}} \nu(u_{m})(\boldsymbol{x}, t), \nabla_{\mathcal{D}_{m}} \zeta(u_{m})(\boldsymbol{x}, t)) \cdot \nabla_{\mathcal{D}_{m}} \zeta(u_{m})(\boldsymbol{x}, t) d\boldsymbol{x} dt$$

$$= \int_{0}^{T_{0}} \int_{\Omega} \boldsymbol{a}(\boldsymbol{x}, \nu(\overline{u})(\boldsymbol{x}, t), \nabla \zeta(\overline{u})(\boldsymbol{x}, t)) \cdot \nabla \zeta(\overline{u})(\boldsymbol{x}, t) d\boldsymbol{x} dt. \quad (65)$$

Let us define

$$f_m = [\boldsymbol{a}(\boldsymbol{x}, \Pi_{\mathcal{D}_m} \nu(u_m), \nabla_{\mathcal{D}_m} \zeta(u_m)) - \boldsymbol{a}(\boldsymbol{x}, \Pi_{\mathcal{D}_m} \nu(u_m)(\cdot, t), \nabla \zeta(\overline{u}))] \cdot [\nabla_{\mathcal{D}_m} \zeta(u_m) - \nabla \zeta(\overline{u})] \ge 0.$$

By developing this expression and using (65), (57) and (18), we see that  $\int_0^T \int_\Omega f_m(\boldsymbol{x},t) \mathrm{d}\boldsymbol{x} \mathrm{d}t \to 0$  as  $m \to \infty$ . This shows that  $f_m \to 0$  in  $L^1(\Omega \times (0,T))$  and therefore a.e. up to a subsequence. We can then reason as in [23], using the strict monotony (19) of  $\boldsymbol{a}$ , the coercivity assumption (2f) and Vitali's theorem, to deduce that  $\nabla_{\mathcal{D}_m} \zeta(u_m) \to \nabla \zeta(\overline{u})$  strongly in  $L^p(\Omega \times (0,T))^d$  as  $m \to \infty$ .

## 5 Removal of the assumption " $\beta = Id$ or $\zeta = Id$ "

We show here that all previous results are actually true without the structural assumption (17) – i.e. without assuming that  $\beta = \operatorname{Id}$  or  $\zeta = \operatorname{Id}$  – provided that the range of p is slightly restricted. The main theorem in this section is the following convergence result.

**Theorem 5.1** Under Assumptions (2), let  $(\mathcal{D}_m)_{m\in\mathbb{N}}$  be a sequence of space-time gradient discretisations, in the sense of Definition 2.1, that is coercive, consistent, limit-conforming and compact (see Section 2.2). Let, for any  $m \in \mathbb{N}$ ,  $u_m$  be a solution to (14) with  $\mathcal{D} = \mathcal{D}_m$ , provided by Theorem 2.12.

If  $p \geq 2$  then there exists a solution  $\overline{u}$  to (4) such that, up to a subsequence,

- the convergences in (18) hold,
- $\Pi_{\mathcal{D}_m}\nu(u_m) \to \nu(\overline{u})$  strongly in  $L^{\infty}(0,T;L^2(\Omega))$  as  $m \to \infty$ ,
- under the strict monotony assumption on  $\mathbf{a}$  (i.e. (19)), as  $m \to \infty$  we have  $\Pi_{\mathcal{D}_m} \zeta(u_m) \to \zeta(\overline{u})$  strongly in  $L^p(\Omega \times (0,T))$  and  $\nabla_{\mathcal{D}_m} \zeta(u_m) \to \nabla \zeta(\overline{u})$  strongly in  $L^p(\Omega \times (0,T))^d$ .

#### Proof.

We only need to prove the first conclusion of the theorem, i.e. that the convergences (18) hold. Theorems 2.16 and 2.18 then provide the last two conclusions. The difference with respect to Theorem 2.12 is the removal, here, of the structural assumption (17). The only place in the proof of Theorem 2.12 where this assumption was used is in Step 1, to identify the limits  $\overline{\beta}$ ,  $\overline{\zeta}$  and  $\overline{\nu}$  of  $\Pi_{\mathcal{D}_m}\beta(u_m)$ ,  $\Pi_{\mathcal{D}_m}\zeta(u_m)$  and  $\Pi_{\mathcal{D}_m}\nu(u_m)$ . We will show that these limits can still be identified without assuming (17).

Set  $\mu = \beta + \zeta$ , let  $\overline{\mu} = \overline{\beta} + \overline{\zeta}$  and fix a measurable  $\overline{u}$  such that  $(\mu + \nu)(\overline{u}) = \overline{\mu} + \overline{\nu}$ . The existence of such a  $\overline{u}$  is ensured by Assumptions (2b) and (2c). Indeed, these assumptions show that the range of  $\mu + \nu$  is  $\mathbb{R}$  and therefore that the pseudo-reciprocal  $(\mu + \nu)_r$  of  $\mu + \nu$  (defined as in (3)) has domain  $\mathbb{R}$ ; this allows us to set, for example,  $\overline{u} = (\mu + \nu)_r(\overline{\mu} + \overline{\nu})$ . Let us now prove that, for such a function  $\overline{u}$ , we have  $\overline{\beta} = \beta(\overline{u})$ ,  $\overline{\zeta} = \zeta(\overline{u})$  and  $\overline{\nu} = \nu(\overline{u})$ .

By using estimates (52) and (53), Kolmogorov's compactness theorem shows that the convergence of  $\Pi_{\mathcal{D}_m}\nu(u_m)$  towards  $\overline{\nu}$  is actually strong in  $L^2(\Omega\times(0,T))$  (we use  $p\geq 2$  here). Since  $\mu(\Pi_{\mathcal{D}_m}u_m)=\beta(\Pi_{\mathcal{D}_m}u_m)+\zeta(\Pi_{\mathcal{D}_m}u_m)\to\overline{\beta}+\overline{\zeta}=\overline{\mu}$  weakly in  $L^2(\Omega\times(0,T))$ , we can apply Lemma 5.6 with  $\varphi\equiv 1,\ w_m=\Pi_{\mathcal{D}_m}u_m,\ w=\overline{u}$  and  $(\mu,\nu)$  instead of  $(\beta,\zeta)$  to deduce that  $\overline{\nu}=\nu(\overline{u})$  and  $\overline{\mu}=\mu(\overline{u})$ . The second of these relations translates into  $\overline{\beta}+\overline{\zeta}=(\beta+\zeta)(\overline{u})$ .

We now turn to identifying  $\overline{\beta}$  and  $\overline{\zeta}$ . Lemmas 4.1 and 4.3 show that  $\beta_m = \beta(u_m)$  and  $\zeta_m = \zeta(u_m)$  satisfy the assumptions of the discrete compensated compactness theorem 5.4 below (we use  $p \geq 2$  here). Hence,  $\Pi_{\mathcal{D}_m}\beta(u_m)\Pi_{\mathcal{D}_m}\zeta(u_m) \to \overline{\beta}\,\overline{\zeta}$  in the sense of measures on  $\Omega \times (0,T)$ . Since we already established that  $(\beta + \zeta)(\overline{u}) = \overline{\beta} + \overline{\zeta}$ , we can therefore apply Lemma 5.6 with  $\varphi \equiv 1$ ,  $w_m = \Pi_{\mathcal{D}_m}u_m$  and  $w = \overline{u}$ . This gives  $\overline{\beta} = \beta(\overline{u})$  and  $\overline{\zeta} = \zeta(\overline{u})$  a.e. on  $\Omega \times (0,T)$ , as required.

To summarise, the limits of  $\Pi_{\mathcal{D}_m}\beta(u_m)$ ,  $\Pi_{\mathcal{D}_m}\zeta(u_m)$  and  $\Pi_{\mathcal{D}_m}\nu(u_m)$  have been identified as  $\beta(\overline{u})$ ,  $\zeta(\overline{u})$  and  $\nu(\overline{u})$  for some  $\overline{u}$ . Since  $\zeta(\overline{u}) = \overline{\zeta} \in L^p(\Omega \times (0,T))$ , the growth assumptions (2b) on  $\zeta$  ensure that  $\overline{u} \in L^p(\Omega \times (0,T))$ . We can then take over the proof of Theorem 2.12 from after the usage of (17), using the  $\overline{u}$  we just found instead of the one defined as the weak limit of  $\Pi_{\mathcal{D}_m}u_m$ . This allows us to conclude that  $\overline{u}$  is a solution to (4), and that the convergences in (18) hold.

**Remark 5.2** It is not proved that  $\overline{u}$  is a weak limit of  $\Pi_{\mathcal{D}_m} u_m$ . Such a limit is not stated in (18) and is not necessarily expected for the model (1), in which the quantities of interest (physically relevant when this PDE models a natural phenomenon) are  $\beta(\overline{u})$ ,  $\zeta(\overline{u})$  and  $\nu(\overline{u})$ .

Remark 5.3 (Maximal monotone operator) Hypotheses (2b) and (2c) imply that the operator T defined by the graph  $\mathcal{G}(T) = \{(\zeta(s), \beta(s)), s \in \mathbb{R}\}$  is a maximal monotone operator with domain  $\mathbb{R}$ , such that  $0 \in T(0)$ . Indeed, assume that x, y satisfy  $(\zeta(s) - x)(\beta(s) - y) \geq 0$  for all  $s \in \mathbb{R}$ . Then, letting  $w \in \mathbb{R}$  be such that

$$\frac{\beta(w) + \zeta(w)}{2} = \frac{x+y}{2},\tag{66}$$

we have  $(\zeta(w)-x)(\beta(w)-y)=-(\frac{\zeta(w)-\beta(w)}{2}-\frac{x-y}{2})^2\geq 0$ . This implies  $\frac{\zeta(w)-\beta(w)}{2}=\frac{x-y}{2}$  which, combined with (66), gives  $x=\zeta(w)$  and  $y=\beta(w)$  and hence  $(x,y)\in\mathcal{G}(T)$ .

Reciprocally, for any maximal monotone operator T from  $\mathbb{R}$  to  $\mathbb{R}$  such that  $0 \in T(0)$ , one can find  $\zeta$  and  $\beta$  satisfying (2b) and (2c), and such that  $\mathcal{G}(T) = \{(\zeta(s), \beta(s)), s \in \mathbb{R}\}$ . Indeed, for all  $(x,y) \in \mathcal{G}(T)$  and  $(x',y') \in \mathcal{G}(T)$  satisfying x+y=x'+y', since  $(x-x')(y-y') \geq 0$  we have x=x' and y=y'. We can therefore define  $\zeta$  and  $\beta$  by: for all  $(x,y) \in \mathcal{G}(T)$ ,  $x=\zeta(\frac{x+y}{2})$  and  $y=\beta(\frac{x+y}{2})$ . We observe that these functions are nondecreasing and Lipschitz-continuous with constant 2, and that  $\zeta + \beta = 2\mathrm{Id}$ .

Hence, Theorem 5.1 applies to the model considered in [52], but provides convergence results for much more general equations and various numerical methods in any space dimension.

We now state the two key results that allowed us to remove Assumption (17) if  $p \ge 2$ . The first one is a discrete version of a compensated compactness result in [41]. The second is a Minty-like result, useful to identify weak non-linear limits.

We note that Theorem 5.4 states a more general convergence result than needed for the proof of Theorem 5.1 (which only requires  $\varphi \equiv 1$ ). We nevertheless state the general form in order to obtain the genuine discrete equivalent of the result in [41]. We also believe that this discrete compensated compactness theorem will find many more applications in the numerical analysis of degenerate or coupled parabolic models. We also refer to [6] for another transposition to the discrete setting of a compensated compactness result.

Theorem 5.4 (Discrete compensated compactness) We take T > 0,  $p \ge 2$  and a sequence  $(\mathcal{D}_m)_{m \in \mathbb{N}} = (X_{\mathcal{D}_m,0},\Pi_{\mathcal{D}_m},\nabla_{\mathcal{D}_m},\mathcal{I}_{\mathcal{D}_m},(t_m^{(n)})_{n=0,...,N_m})_{m \in \mathbb{N}}$  of space-time gradient discretisations, in the sense of Definition 2.1, that is consistent and compact in the sense of Definitions 2.6 and 2.9.

For any  $m \in \mathbb{N}$ , let  $\beta_m = (\beta_m^{(n)})_{n=0,\dots,N_m} \subset X_{\mathcal{D}_m,0}$  and  $\zeta_m = (\zeta_m^{(n)})_{n=0,\dots,N_m} \subset X_{\mathcal{D}_m,0}$  be such that

- the sequences  $(\int_0^T |\delta_m \beta_m(t)|_{\star,\mathcal{D}_m})_{m \in \mathbb{N}}$  and  $(||\nabla_{\mathcal{D}_m} \zeta_m||_{L^2(0,T;L^p(\Omega)^d)})_{m \in \mathbb{N}}$  are bounded,
- as  $m \to \infty$ ,  $\Pi_{\mathcal{D}_m} \beta_m \to \overline{\beta}$  and  $\Pi_{\mathcal{D}_m} \zeta_m \to \overline{\zeta}$  weakly in  $L^2(\Omega \times (0,T))$ .

Then  $(\Pi_{\mathcal{D}_m}\beta_m)(\Pi_{\mathcal{D}_m}\zeta_m) \to \overline{\beta}\overline{\zeta}$  in the sense of measures on  $\Omega \times (0,T)$ , that is, for all  $\varphi \in C(\overline{\Omega} \times [0,T])$ ,

$$\lim_{m \to \infty} \int_0^T \int_{\Omega} \Pi_{\mathcal{D}_m} \beta_m(\boldsymbol{x}, t) \Pi_{\mathcal{D}_m} \zeta_m(\boldsymbol{x}, t) \varphi(\boldsymbol{x}, t) d\boldsymbol{x} dt = \int_0^T \int_{\Omega} \overline{\beta}(\boldsymbol{x}, t) \overline{\zeta}(\boldsymbol{x}, t) \varphi(\boldsymbol{x}, t) d\boldsymbol{x} dt.$$
 (67)

#### Proof.

The idea is to reduce to the case where  $\Pi_{\mathcal{D}_m}\zeta_m$  is a tensorial function, in order to separate the space and time variables and make use of the compactness of  $\Pi_{\mathcal{D}_m}\zeta_m$  and  $\Pi_{\mathcal{D}_m}\beta_m$  with respect to each of these variables. Note that the technique we use here apparently provides a new proof for the continuous equivalent of this compensated compactness result.

Step 1: reduction of  $\Pi_{\mathcal{D}_m}\zeta_m$  to tensorial functions.

Let us take  $\delta > 0$  and let us consider a covering  $(A_k^{\delta})_{k=1,...,K}$  of  $\Omega$  in disjoint cubes of length  $\delta$ . Let  $R_{\delta}: L^2(\Omega) \to L^2(\Omega)$  be the operator defined by:

$$\forall g \in L^2(\Omega), \ \forall k = 1, \dots, K, \ \forall \boldsymbol{x} \in A_k^{\delta} \cap \Omega \ : \ R_{\delta}g(\boldsymbol{x}) = \frac{1}{\operatorname{meas}(A_k^{\delta})} \int_{A_k^k} g(\boldsymbol{y}) \mathrm{d}\boldsymbol{y},$$

where g has been extended by 0 outside  $\Omega$ . Let  $\boldsymbol{x} \in A_k^{\delta} \cap \Omega$ . Using Jensen's inequality, the fact that  $\operatorname{meas}(A_k^{\delta}) = \delta^d$  and the change of variable  $\boldsymbol{y} \in A_k^{\delta} \mapsto \boldsymbol{\xi} = \boldsymbol{y} - \boldsymbol{x} \in (-\delta, \delta)^d$ , we can write

$$|R_{\delta}g(\boldsymbol{x}) - g(\boldsymbol{x})|^2 \le \delta^{-d} \int_{A_k^{\delta}} |g(\boldsymbol{y}) - g(\boldsymbol{x})|^2 d\boldsymbol{y} \le \delta^{-d} \int_{(-\delta,\delta)^d} |g(\boldsymbol{x} + \boldsymbol{\xi}) - g(\boldsymbol{x})|^2 d\boldsymbol{\xi}.$$

Integrating over  $x \in A_k^{\delta}$  and summing over k = 1, ..., K gives

$$||R_{\delta}g - g||_{L^{2}(\Omega)}^{2} \leq \delta^{-d} \int_{(-\delta,\delta)^{d}} \int_{\mathbb{R}^{d}} |g(\boldsymbol{x} + \boldsymbol{\xi}) - g(\boldsymbol{x})|^{2} d\boldsymbol{x} d\boldsymbol{\xi}$$

$$\leq 2^{d} \sup_{\boldsymbol{\xi} \in (-\delta,\delta)^{d}} \int_{\mathbb{R}^{d}} |g(\boldsymbol{x} + \boldsymbol{\xi}) - g(\boldsymbol{x})|^{2} d\boldsymbol{x}. \tag{68}$$

The compactness of  $(\mathcal{D}_m)_{m\in\mathbb{N}}$  (Definition 2.9) and the fact that  $p\geq 2$  give  $\epsilon(\boldsymbol{\xi})$  such that  $\epsilon(\boldsymbol{\xi})\to 0$  as  $\boldsymbol{\xi}\to 0$  and, for all  $m\in\mathbb{N}$  and all  $v\in X_{\mathcal{D}_m,0}$ ,

$$||\Pi_{\mathcal{D}_m} v(\cdot + \boldsymbol{\xi}) - \Pi_{\mathcal{D}_m} v||_{L^2(\mathbb{R}^d)}^2 \le \epsilon(\boldsymbol{\xi}) ||\nabla_{\mathcal{D}_m} v||_{L^p(\Omega)^d}^2.$$

Combining this with (68) and using the bound on  $||\nabla_{\mathcal{D}_m}\zeta_m||_{L^2(0,T;L^p(\Omega)^d)}$  shows that

$$||R_{\delta}\Pi_{\mathcal{D}_{m}}\zeta_{m} - \Pi_{\mathcal{D}_{m}}\zeta_{m}||_{L^{2}(\Omega\times(0,T))} \leq C \sup_{|\boldsymbol{\xi}|_{\infty}\leq\delta} \sqrt{\epsilon(\boldsymbol{\xi})} =: \omega(\delta)$$
(69)

where C does not depend on m, and  $\omega(\delta) \to 0$  as  $\delta \to 0$ . Note that a similar estimate holds with  $\Pi_{\mathcal{D}_m} \zeta_m$  replaced with  $\overline{\zeta}$  since  $\overline{\zeta} \in L^2(\Omega \times (0,T))$ .

If we respectively denote by  $\mathcal{A}_m(\Pi_{\mathcal{D}_m}\zeta_m)$  and  $\mathcal{A}(\overline{\zeta})$  the integrals in the left-hand side and right-hand side (67), then since  $(\Pi_{\mathcal{D}_m}\beta_m)_{m\in\mathbb{N}}$  is bounded in  $L^2(\Omega\times(0,T))$  we have by (69)

$$|\mathcal{A}_m(\Pi_{\mathcal{D}_m}\zeta_m) - \mathcal{A}(\overline{\zeta})| \le C\omega(\delta) + |\mathcal{A}_m(R_\delta\Pi_{\mathcal{D}_m}\zeta_m) - \mathcal{A}(R_\delta\overline{\zeta})|. \tag{70}$$

Let us assume that we can prove that, for a fixed  $\delta$ ,

$$\mathcal{A}_m(R_\delta \Pi_{\mathcal{D}_m} \zeta_m) \to \mathcal{A}(R_\delta \overline{\zeta}) \text{ as } m \to \infty.$$
 (71)

Then (70) gives  $\limsup_{m\to\infty} |\mathcal{A}_m(\Pi_{\mathcal{D}_m}\zeta_m) - \mathcal{A}(\overline{\zeta})| \leq C\omega(\delta)$ . Letting  $\delta \to 0$  in this inequality gives  $\mathcal{A}_m(\Pi_{\mathcal{D}_m}\zeta_m) \to \mathcal{A}(\overline{\zeta})$  as wanted. Hence, we only need to prove (71). The definition of  $R_\delta$  shows that

$$R_{\delta}g = \sum_{k=1}^{K} \frac{1}{\operatorname{meas}(A_k^{\delta})} \mathbf{1}_{A_k^{\delta}}[g]_{A_k^{\delta}},$$

where  $\mathbf{1}_{A_k^{\delta}}$  is the characteristic function of  $A_k^{\delta}$  and  $[g]_A = \int_A g(\mathbf{x}) d\mathbf{x}$ . Hence, (71) follows if we can prove that for any measurable set A

$$\lim_{m \to \infty} \int_0^T \int_{\Omega} \Pi_{\mathcal{D}_m} \beta_m(\boldsymbol{x}, t) [\Pi_{\mathcal{D}_m} \zeta_m]_A(t) \varphi(t, \boldsymbol{x}) \mathbf{1}_A(\boldsymbol{x}) d\boldsymbol{x} dt$$

$$= \int_0^T \int_{\Omega} \overline{\beta}(\boldsymbol{x}, t) [\overline{\zeta}]_A(t) \varphi(t, \boldsymbol{x}) \mathbf{1}_A(\boldsymbol{x}) d\boldsymbol{x} dt \quad (72)$$

where for  $g \in L^2(\Omega \times (0,T))$  we set  $[g]_A(t) = \int_A g(t, \boldsymbol{y}) d\boldsymbol{y}$ .

#### Step 2: further reductions.

We now reduce  $\varphi$  to a tensorial function and  $\mathbf{1}_A$  to a smooth function. It is well-known that there exists tensorial functions  $\varphi_r = \sum_{l=1}^{L_r} \theta_{l,r}(t) \gamma_{l,r}(\boldsymbol{x})$ , with  $\theta_{l,r} \in C^{\infty}([0,T])$  and  $\gamma_{l,r} \in C^{\infty}(\overline{\Omega})$ , such that  $\varphi_r \to \varphi$  uniformly on  $\Omega \times (0,T)$  as  $r \to \infty$ . Moreover, there exists  $\rho_r \in C_c^{\infty}(\Omega)$  such that  $\rho_r \to \mathbf{1}_A$  in  $L^2(\Omega)$  as  $r \to \infty$ .

Hence, as  $r \to \infty$  the function  $(t, \boldsymbol{x}) \mapsto \varphi_r(t, \boldsymbol{x})\rho_r(\boldsymbol{x})$  converges in  $L^{\infty}(0, T; L^2(\Omega))$  to the function  $(t, \boldsymbol{x}) \mapsto \varphi(t, \boldsymbol{x})\mathbf{1}_A(\boldsymbol{x})$ . Since the sequence of functions  $(t, \boldsymbol{x}) \mapsto \Pi_{\mathcal{D}_m}\beta_m(t, \boldsymbol{x})[\Pi_{\mathcal{D}_m}\zeta_m]_A(t)$  is bounded in  $L^1(0, T; L^2(\Omega))$  (notice that  $([\Pi_{\mathcal{D}_m}\zeta_m]_A)_{m\in\mathbb{N}}$  is bounded in  $L^2(0, T)$  since  $(\Pi_{\mathcal{D}_m}\zeta_m)_{m\in\mathbb{N}}$  is bounded in  $L^2(\Omega \times (0, T))$ ), a reasoning similar to the one used in Step 1 shows that we only need to prove (72) with  $\varphi(t, \boldsymbol{x})\mathbf{1}_A(\boldsymbol{x})$  replaced with  $\varphi_r(t, \boldsymbol{x})\rho_r(\boldsymbol{x})$  for a fixed r.

need to prove (72) with  $\varphi(t, \boldsymbol{x})\mathbf{1}_A(\boldsymbol{x})$  replaced with  $\varphi_r(t, \boldsymbol{x})\rho_r(\boldsymbol{x})$  for a fixed r. We have  $\varphi_r(t, \boldsymbol{x})\rho_r(\boldsymbol{x}) = \sum_{l=1}^{L_r} \theta_{l,r}(t)(\gamma_{l,r}\rho_r)(\boldsymbol{x})$  and  $\gamma_{l,r}\rho_r \in C_c^{\infty}(\Omega)$ . Hence, (72) with  $\varphi(t, \boldsymbol{x})\mathbf{1}_A(\boldsymbol{x})$  replaced with  $\varphi_r(t, \boldsymbol{x})\rho_r(\boldsymbol{x})$  will follow if we can establish that for any  $\theta \in C_c^{\infty}([0, T])$ , any  $\psi \in C_c^{\infty}(\Omega)$  and any measurable set A

$$\lim_{m \to \infty} \int_0^T \int_{\Omega} \theta(t) \Pi_{\mathcal{D}_m} \beta_m(\boldsymbol{x}, t) [\Pi_{\mathcal{D}_m} \zeta_m]_A(t) \psi(\boldsymbol{x}) d\boldsymbol{x} dt = \int_0^T \int_{\Omega} \theta(t) \overline{\beta}(\boldsymbol{x}, t) [\overline{\zeta}]_A(t) \psi(\boldsymbol{x}) d\boldsymbol{x} dt.$$
(73)

#### **Step 3**: proof of (73).

We now use the estimate on  $\delta_m \beta_m$  to conclude. We write

$$\int_{0}^{T} \int_{\Omega} \theta(t) \Pi_{\mathcal{D}_{m}} \beta_{m}(\boldsymbol{x}, t) [\Pi_{\mathcal{D}_{m}} \zeta_{m}]_{A}(t) \psi(\boldsymbol{x}) d\boldsymbol{x} dt = \int_{0}^{T} \theta(t) [\Pi_{\mathcal{D}_{m}} \zeta_{m}]_{A}(t) F_{m}(t)$$
(74)

with  $F_m(t) = \int_{\Omega} \Pi_{\mathcal{D}_m} \beta_m(\boldsymbol{x}, t) \psi(\boldsymbol{x}) d\boldsymbol{x}$ . It is clear from the weak convergence of  $\Pi_{\mathcal{D}_m} \zeta_m$  that  $[\Pi_{\mathcal{D}_m} \zeta_m]_A \to [\overline{\zeta}]_A$  weakly in  $L^2(0,T)$ . Hence, if we can prove that  $F_m \to F := \int_{\Omega} \overline{\beta}(\boldsymbol{x}, \cdot) \psi(\boldsymbol{x}) d\boldsymbol{x}$  strongly in  $L^2(0,T)$ , we can pass to the limit in (74) and obtain (73). Since  $F_m$  weakly converges to F in  $L^2(0,T)$  (thanks to the weak convergence of  $\Pi_{\mathcal{D}_m} \beta_m$  in  $L^2(\Omega \times (0,T))$ ), we only have to prove that  $(F_m)_{m \in \mathbb{N}}$  is relatively compact in  $L^2(0,T)$ .

We introduce the interpolant  $P_{\mathcal{D}_m}$  defined by (21) and we define  $G_m$  as  $F_m$  with  $\psi$  replaced with  $\Pi_{\mathcal{D}_m} P_{\mathcal{D}_m} \psi$ . We then have

$$|F_m(t) - G_m(t)| \le ||\Pi_{\mathcal{D}_m} \beta_m(\cdot, t)||_{L^2(\Omega)} S_{\mathcal{D}_m}(\psi).$$

The consistency of  $(\mathcal{D}_m)_{m\in\mathbb{N}}$  thus shows that

$$F_m - G_m \to 0$$
 strongly in  $L^2(0,T)$  as  $m \to \infty$ . (75)

We now study the strong convergence of  $G_m$ . This function is, like  $\Pi_{\mathcal{D}_m}\beta_m$ , piecewise constant on (0,T) and, by definition of  $|\cdot|_{\star,\mathcal{D}_m}$ , its discrete derivative satisfies

$$|\delta_m G_m(t)| \le |\delta_m \beta_m(t)|_{\star, \mathcal{D}_m} ||\nabla_{\mathcal{D}_m} P_{\mathcal{D}_m} \psi||_{L^p(\Omega)^d}.$$

Since  $||\nabla_{\mathcal{D}_m} P_{\mathcal{D}_m} \psi||_{L^p(\Omega)^d} \leq S_{\mathcal{D}_m}(\psi) + ||\nabla \psi||_{L^p(\Omega)^d}$  is bounded uniformly with respect to m, the assumption on  $\delta_m \beta_m$  proves that  $(||\delta_m G_m||_{L^1(0,T)})_{m \in \mathbb{N}}$  is bounded. We have  $||\delta_m G_m||_{L^1(0,T)} = |G_m|_{BV(0,T)}$ , and  $(\Pi_{\mathcal{D}_m} \beta_m)_{m \in \mathbb{N}}$  is bounded in  $L^2(\Omega \times (0,T))$ ; hence,  $(G_m)_{m \in \mathbb{N}}$  is bounded in  $BV(0,T) \cap L^2(0,T)$  and therefore relatively compact in  $L^2(0,T)$  (see [7, Theorem 10.1.4]). Combined with (75), this shows that  $(F_m)_{m \in \mathbb{N}}$  is relatively compact in  $L^2(0,T)$  and concludes the proof.

Remark 5.5 If we assume that  $(\Pi_{\mathcal{D}_m}\beta_m)_{m\in\mathbb{N}}$  is bounded in  $L^{\infty}(0,T;L^2(\Omega))$  and that, for some q>1,  $(\int_0^T |\delta_m\beta_m(t)|_{\star,\mathcal{D}_m}^q)_{m\in\mathbb{N}}$  is bounded, then Step 3 becomes a trivial consequence of Theorem 3.1. Indeed, this theorem shows that  $(\Pi_{\mathcal{D}_m}\beta_m)_{m\in\mathbb{N}}$  is relatively compact uniformly-in-time and weakly in  $L^2(\Omega)$ , which translates into the relative compactness of  $(F_m)_{m\in\mathbb{N}}$  in  $L^{\infty}(0,T)$ .

**Lemma 5.6** Let V be a non-empty measurable subset of  $\mathbb{R}^N$ ,  $N \geq 1$ . Let  $\beta, \zeta \in C^0(\mathbb{R})$  be two nondecreasing functions such that  $\beta(0) = \zeta(0) = 0$ . We assume that there exists a sequence  $(w_m)_{m \in \mathbb{N}}$  of measurable functions on V, and two functions  $\overline{\beta}, \overline{\zeta} \in L^2(V)$  such that:

- $\beta(w_m) \to \overline{\beta}$  and  $\zeta(w_m) \to \overline{\zeta}$  weakly in  $L^2(V)$ ,
- there exists  $\varphi \in L^{\infty}(V)$  such that  $\varphi > 0$  a.e. on V and

$$\lim_{m \to \infty} \int_{V} \varphi(z) \beta(w_m(z)) \zeta(w_m(z)) dz = \int_{V} \varphi(z) \overline{\beta}(z) \overline{\zeta}(z) dz.$$
 (76)

Then, for any measurable function w such that  $(\beta + \zeta)(w) = \overline{\beta} + \overline{\zeta}$  a.e. in V, we have

$$\overline{\beta} = \beta(w) \text{ and } \overline{\zeta} = \zeta(w) \text{ a.e. in } V.$$
 (77)

**Proof.** We first notice that  $\beta(w)$  and  $\zeta(w)$  belong to  $L^2(V)$  since they have the same sign and therefore verify  $|\beta(w)| + |\zeta(w)| = |\overline{\beta} + \overline{\zeta}| \in L^2(V)$ . Using the fact that  $\beta$  and  $\zeta$  are non-decreasing, we can write

$$\int_{V} \varphi(\boldsymbol{z}) \left[ \beta(w_{m}(\boldsymbol{z})) - \beta(w(\boldsymbol{z})) \right] \left[ \zeta(w_{m}(\boldsymbol{z})) - \zeta(w(\boldsymbol{z})) \right] d\boldsymbol{z} \geq 0.$$

Letting  $m \to \infty$  in the above inequality, and using the convergences of  $\beta(w_m)$ ,  $\zeta(w_m)$  and (76), we obtain

$$\int_{V} \varphi(z) \left[ \overline{\beta}(z) - \beta(w(z)) \right] \left[ \overline{\zeta}(z) - \zeta(w(z)) \right] dz \ge 0.$$
 (78)

We then remark that  $\overline{\beta} + \overline{\zeta} = \beta(w) + \zeta(w)$  gives  $\beta(w) = \frac{\overline{\beta} + \overline{\zeta}}{2} + \left(\frac{\beta - \zeta}{2}\right)(w)$  and  $\zeta(w) = \frac{\overline{\beta} + \overline{\zeta}}{2} - \left(\frac{\beta - \zeta}{2}\right)(w)$ . Hence, (78) leads to

$$-\int_{V}\varphi(\boldsymbol{z})\left[\frac{\overline{\beta}-\overline{\zeta}}{2}(\boldsymbol{z})-\left(\frac{\beta-\zeta}{2}\right)(w(\boldsymbol{z}))\right]^{2}\mathrm{d}\boldsymbol{z}\geq0.$$

Since  $\varphi$  is almost everywhere strictly positive on V, we deduce that  $\frac{\overline{\beta}-\overline{\zeta}}{2}=\frac{\beta(w)-\zeta(w)}{2}$  a.e. in V, and (77) follows from  $\frac{\overline{\beta}+\overline{\zeta}}{2}=\frac{\beta(w)+\zeta(w)}{2}$ .

## 6 Appendix: uniform-in-time compactness results for timedependent problems

We establish in this appendix some generic results, unrelated to the framework of gradient schemes, that form the starting point for our uniform-in-time convergence results.

Solutions of numerical schemes for parabolic equations are usually piecewise constant, and therefore not continuous, in time. As their jumps nevertheless tend to become small as the time step goes to 0, it is possible to establish uniform-in-time convergence properties using a generalisation to non-continuous functions of the classical Ascoli-Arzelà theorem.

**Definition 6.1** If  $(K, d_K)$  and  $(E, d_E)$  are metric spaces, we denote by  $\mathcal{F}(K, E)$  the space of functions  $K \to E$  endowed with the uniform metric  $d_{\mathcal{F}}(v, w) = \sup_{s \in K} d_E(v(s), w(s))$  (note that this metric may take infinite values).

**Theorem 6.2** (discontinuous Ascoli-Arzelà's theorem) Let  $(K, d_K)$  be a compact metric space,  $(E, d_E)$  be a complete metric space and  $(\mathcal{F}(K, E), d_{\mathcal{F}})$  be as in Definition 6.1. Let  $(v_m)_{m \in \mathbb{N}}$  be a sequence in  $\mathcal{F}(K, E)$  such that there exists a function  $\omega : K \times K \to [0, \infty]$  and a sequence  $(\delta_m)_{m \in \mathbb{N}} \subset [0, \infty)$  satisfying

$$\lim_{d_K(s,s')\to 0} \omega(s,s') = 0, \quad \lim_{m\to\infty} \delta_m = 0,$$

$$\forall (s,s') \in K^2, \ \forall m \in \mathbb{N}, \ d_E(v_m(s),v_m(s')) \le \omega(s,s') + \delta_m.$$
(79)

We also assume that, for all  $s \in K$ ,  $\{v_m(s) : m \in \mathbb{N}\}$  is relatively compact in  $(E, d_E)$ . Then  $(v_m)_{m \in \mathbb{N}}$  is relatively compact in  $(\mathcal{F}(K, E), d_{\mathcal{F}})$  and any adherence value of  $(v_m)_{m \in \mathbb{N}}$  in this space is continuous  $K \to E$ .

**Proof.** Let us first notice that the last conclusion of the theorem, i.e. that any adherence value v of  $(v_m)_{m\in\mathbb{N}}$  in  $\mathcal{F}(K,E)$  is continuous, is trivially obtained by passing to the limit in (79), which shows that the modulus of continuity of v is bounded above by  $\omega$ .

The proof of the compactness result is an easy generalisation of the proof of the classical Ascoli-Arzelà theorem. We start by taking a countable dense subset  $\{s_l : l \in \mathbb{N}\}$  in K (the existence of this set is ensured since K is compact metric). Since each set  $\{v_m(s_l) : m \in \mathbb{N}\}$  is relatively compact in E, by diagonal extraction we can select a subsequence of  $(v_m)_{m \in \mathbb{N}}$ , denoted the same way, such that, for any  $l \in \mathbb{N}$ ,  $(v_m(s_l))_{m \in \mathbb{N}}$  converges in E. We then proceed to show that  $(v_m)_{m \in \mathbb{N}}$  is a Cauchy sequence in  $(\mathcal{F}(K, E), d_{\mathcal{F}})$ . Since this space is complete, this will prove that this sequence converges in this space, which will complete the proof.

Let  $\varepsilon > 0$  and, using (79), take  $\rho > 0$  and  $M \in \mathbb{N}$  such that  $\omega(s, s') \leq \varepsilon$  whenever  $d_K(s, s') \leq \rho$  and  $\delta_m \leq \varepsilon$  whenever  $m \geq M$ . Select a finite set  $\{s_{l_1}, \ldots, s_{l_N}\}$  such that any  $s \in K$  is within distance  $\rho$  of a  $s_{l_i}$ . Then for any  $m, m' \geq M$ 

$$d_{E}(v_{m}(s), v_{m'}(s)) \leq d_{E}(v_{m}(s), v_{m}(s_{l_{i}})) + d_{E}(v_{m}(s_{l_{i}}), v_{m'}(s_{l_{i}})) + d_{E}(v_{m'}(s_{l_{i}}), v_{m'}(s))$$

$$\leq \omega(s, s_{l_{i}}) + \delta_{m} + d_{E}(v_{m}(s_{l_{i}}), v_{m'}(s_{l_{i}})) + \omega(s, s_{l_{i}}) + \delta_{m'}$$

$$\leq 4\varepsilon + d_{E}(v_{m}(s_{l_{i}}), v_{m'}(s_{l_{i}})).$$

Since  $\{(v_m(s_{l_i}))_{m\in\mathbb{N}}: i=1,\ldots,N\}$  forms a finite number of converging sequences in E, we can find  $M'\geq M$  such that, for all  $m,m'\geq M'$  and all  $i=1,\ldots,N,$   $d_E(v_m(s_{l_i}),v_{m'}(s_{l_i}))\leq \varepsilon$ . This shows that, for all  $m,m'\geq M'$  and all  $s\in K,$   $d_E(v_m(s),v_{m'}(s))\leq 5\varepsilon$  and concludes the proof that  $(v_m)_{m\in\mathbb{N}}$  is a Cauchy sequence in  $(\mathcal{F}(K,E),d_{\mathcal{F}})$ .

Remark 6.3 Conditions (79) are usually the most practical when  $(v_m)_{m\in\mathbb{N}}$  are piecewise constantin-time solutions to numerical schemes (see e.g. the proof of Theorem 3.1). Here,  $\omega$  is expected to measure the size of the cumulated jumps of  $v_m$  between s and s', and  $\delta_m$  accounts for boundary effects which may occur in the small time intervals containing s and s'. It is easy to see that (79) can be replaced with

$$d_E(v_m(s), v_m(s')) \to 0$$
, as  $m \to \infty$  and  $d_K(s, s') \to 0$  (80)

(under this condition, the proof can be carried out by selecting  $M \in \mathbb{N}$  and  $\rho > 0$  such that  $d_E(v_m(s), v_m(s')) \leq \varepsilon$  whenever  $m \geq M$  and  $d_K(s, s') \leq \rho$ ). It turns out that (80) is actually a necessary and sufficient condition for the theorem's conclusions to hold true.

The following lemma states an equivalent condition for the uniform convergence of functions, which proves extremely useful to establish uniform-in-time convergence of numerical schemes for parabolic equations when no smoothness is assumed on the data.

**Lemma 6.4** Let  $(K, d_K)$  be a compact metric space,  $(E, d_E)$  be a metric space and  $(\mathcal{F}(K, E), d_{\mathcal{F}})$  as in Definition 6.1. Let  $(v_m)_{m \in \mathbb{N}}$  be a sequence in  $\mathcal{F}(K, E)$  and  $v : K \mapsto E$  be continuous. Then  $v_m \to v$  for  $d_{\mathcal{F}}$  if and only if, for any  $s \in K$  and any sequence  $(s_m)_{m \in \mathbb{N}} \subset K$  converging to s for  $d_K$ , we have  $v_m(s_m) \to v(s)$  for  $d_E$ .

**Proof.** If  $v_m \to v$  for  $d_{\mathcal{F}}$  then for any sequence  $(s_m)_{m \in \mathbb{N}}$  converging to s

$$d_E(v_m(s_m), v(s)) \le d_E(v_m(s_m), v(s_m)) + d_E(v(s_m), v(s)) \le d_F(v_m, v) + d_E(v(s_m), v(s)).$$

The right-hand side tends to 0 by definition of  $v_m \to v$  for  $d_{\mathcal{F}}$  and by continuity of v, which shows that  $v_m(s_m) \to v(s)$  for  $d_E$ .

Let us now prove the converse by contradiction. If  $(v_m)_{m\in\mathbb{N}}$  does not converge to v for  $d_{\mathcal{F}}$  then there exists  $\varepsilon > 0$  and a subsequence  $(v_{m_k})_{k\in\mathbb{N}}$ , such that, for any  $k \in \mathbb{N}$ ,  $\sup_{s \in K} d_E(v_{m_k}(s), v(s)) \ge \varepsilon$ . We can then find a sequence  $(r_k)_{k\in\mathbb{N}} \subset K$  such that, for any  $k \in \mathbb{N}$ ,

$$d_E(v_{m_k}(r_k), v(r_k)) \ge \varepsilon/2. \tag{81}$$

K being compact, up to another subsequence denoted the same way, we can assume that  $r_k$  converges as  $k \to \infty$  to some s in K. It is then trivial to construct a sequence  $(s_m)_{m \in \mathbb{N}}$  converging to s and such that  $s_{m_k} = r_k$  (just take  $s_m = s$  when m is not an  $m_k$ ). We then have  $v_m(s_m) \to v(s)$  in E and, by continuity of  $v, v(s_m) \to v(s)$  in E. This shows that  $d_E(v_m(s_m), v(s_m)) \to 0$ , which contradicts (81) and concludes the proof.

The next result is classical. Its short proof is recalled for the reader's convenience.

**Proposition 6.5** Let E be a closed bounded ball in  $L^2(\Omega)$  and let  $(\varphi_l)_{l\in\mathbb{N}}$  be a dense sequence in  $L^2(\Omega)$ . Then, on E, the weak topology of  $L^2(\Omega)$  is the topology given by the metric

$$d_E(v, w) = \sum_{l \in \mathbb{N}} \frac{\min(1, |\langle v - w, \varphi_l \rangle_{L^2(\Omega)}|)}{2^l}.$$
 (82)

Moreover, a sequence of functions  $u_m:[0,T]\to E$  converges uniformly-in-time to  $u:[0,T]\to E$  for the weak topology of  $L^2(\Omega)$  (see Definition 2.11) if and only if, as  $m\to\infty$ ,  $d_E(u_m,u):[0,T]\to [0,\infty)$  converges uniformly to 0.

**Proof.** The sets  $E_{\varphi,\varepsilon} = \{v \in E : |\langle v, \varphi \rangle_{L^2(\Omega)}| < \varepsilon\}$ , for  $\varphi \in L^2(\Omega)$  and  $\varepsilon > 0$ , define a basis of neighborhoods of 0 for the weak  $L^2(\Omega)$  topology on E, and a basis of neighborhoods of any other point is obtained by translation. If R is the radius of the ball E then for any  $\varphi \in L^2(\Omega)$ ,  $l \in \mathbb{N}$  and  $v \in E$  we have

$$|\langle v, \varphi \rangle_{L^2(\Omega)}| \le R||\varphi - \varphi_l||_{L^2(\Omega)} + |\langle v, \varphi_l \rangle_{L^2(\Omega)}|.$$

By density of  $(\varphi_l)_{l\in\mathbb{N}}$  we can select  $l\in\mathbb{N}$  such that  $||\varphi-\varphi_l||_{L^2(\Omega)}<\varepsilon/(2R)$  and we then see that  $E_{\varphi_l,\varepsilon/2}\subset E_{\varphi,\varepsilon}$ . Hence, a basis of neighborhoods of 0 in E for the weak  $L^2(\Omega)$  is also given by  $(E_{\varphi_l,\varepsilon})_{l\in\mathbb{N}}$   $\varepsilon>0$ .

From the definition of  $d_E$  we see that, for any  $l \in \mathbb{N}$ ,  $\min(1, |\langle v, \varphi_l \rangle_{L^2(\Omega)}|) \leq 2^l d_E(0, v)$ . If  $d_E(0, v) < 2^{-l}$  this shows that  $|\langle v, \varphi_l \rangle_{L^2(\Omega)}| \leq 2^l d_E(0, v)$  and therefore that

$$B_{d_E}(0, \min(2^{-l}, \varepsilon 2^{-l})) \subset E_{\varphi_l, \varepsilon}.$$

Hence, any neighborhood of 0 in E for the  $L^2(\Omega)$  weak topology is a neighborhood of 0 for  $d_E$ . Conversely, for any  $\varepsilon > 0$ , selecting  $N \in \mathbb{N}$  such that  $\sum_{l \geq N+1} 2^{-l} < \varepsilon/2$  gives, from the definition (82) of  $d_E$ ,

$$\bigcap_{l=1}^{N} E_{\varphi_{l},\varepsilon/4} \subset B_{d_{E}}(0,\varepsilon).$$

Hence, any ball for  $d_E$  centered at 0 is a neighborhood of 0 for the  $L^2(\Omega)$  weak topology. Since  $d_E$  and the  $L^2(\Omega)$  weak neighborhoods are invariant by translation, this concludes the proof that this weak topology is identical to the topology generated by  $d_E$ .

The conclusion on weak uniform convergence of sequences of functions follows from the preceding result, and more precisely by noticing that all previous inclusions are, when applied to  $u_m(t) - u(t)$ , uniform with respect to  $t \in [0, T]$ .

The following lemma has been initially established in [35, Proposition 9.3].

#### Lemma 6.6

Let  $(t^{(n)})_{n\in\mathbb{Z}}$  be a strictly increasing sequence of real values such that  $\delta t^{(n+\frac{1}{2})} := t^{(n+1)} - t^{(n)}$  is uniformly bounded by  $\delta t > 0$ ,  $\lim_{n \to -\infty} t^{(n)} = -\infty$  and  $\lim_{n \to \infty} t^{(n)} = \infty$ . For all  $t \in \mathbb{R}$ , we denote by n(t) the element  $n \in \mathbb{Z}$  such that  $t \in (t^{(n)}, t^{(n+1)}]$ . Let  $(a^{(n)})_{n \in \mathbb{Z}}$  be a family of non negative real numbers with a finite number of non zero values. Then

$$\int_{\mathbb{R}} \sum_{n=n(t)+1}^{n(t+\tau)} (\delta^{(n+\frac{1}{2})} a^{(n+1)}) dt = \tau \sum_{n \in \mathbb{Z}} (\delta^{(n+\frac{1}{2})} a^{(n+1)}), \quad \forall \tau > 0,$$
 (83)

and

$$\int_{\mathbb{R}} \left( \sum_{n=n(t)+1}^{n(t+\tau)} \delta t^{(n+\frac{1}{2})} \right) a^{n(t+s)+1} dt \le (\tau + \delta t) \sum_{n \in \mathbb{Z}} \delta t^{(n+\frac{1}{2})} a^{(n+1)}, \quad \forall \tau > 0, \ \forall s \in \mathbb{R}.$$
 (84)

#### Proof.

Let us define  $\chi$  by  $\chi(t, n, \tau) = 1$  if  $t^{(n)} \in [t, t + \tau)$ , otherwise  $\chi(t, n, \tau) = 0$ . We have

$$\int_{\mathbb{R}} \sum_{n=n(t)+1}^{n(t+\tau)} (\check{\alpha}^{(n+\frac{1}{2})} a^{(n+1)}) dt = \int_{\mathbb{R}} \sum_{n \in \mathbb{Z}} (\check{\alpha}^{(n+\frac{1}{2})} a^{(n+1)} \chi(t, n, \tau)) dt$$
$$= \sum_{n \in \mathbb{Z}} \left( \check{\alpha}^{(n+\frac{1}{2})} a^{(n+1)} \int_{\mathbb{R}} \chi(t, n, \tau) dt \right).$$

Since  $\int_{\mathbb{R}} \chi(t, n, \tau) dt = \int_{t^{(n)} - \tau}^{t^{(n)}} dt = \tau$ , Relation (83) is proved. We now turn to the proof of (84). We define  $\widetilde{\chi}$  by  $\widetilde{\chi}(n, t) = 1$  if n(t) = n, otherwise  $\widetilde{\chi}(n, t) = 0$ . We have

$$\int_{\mathbb{R}} \left( \sum_{n=n(t)+1}^{n(t+\tau)} \delta t^{(n+\frac{1}{2})} \right) a^{(n(t+s)+1)} dt = \int_{\mathbb{R}} \left( \sum_{n=n(t)+1}^{n(t+\tau)} \delta t^{(n+\frac{1}{2})} \right) \sum_{m \in \mathbb{Z}} a^{(m+1)} \widetilde{\chi}(m,t+s) dt,$$

which yields

$$\int_{\mathbb{R}} \left( \sum_{n=n(t)+1}^{n(t+\tau)} \delta t^{(n+\frac{1}{2})} \right) a^{(n(t+s)+1)} dt = \sum_{m \in \mathbb{Z}} a^{(m+1)} \int_{t^{(m)}-s}^{t^{(m+1)}-s} \left( \sum_{n=n(t)+1}^{n(t+\tau)} \delta t^{(n+\frac{1}{2})} \right) dt. \tag{85}$$

Since

$$\sum_{n=n(t)+1}^{n(t+\tau)} \delta t^{(n+\frac{1}{2})} = \sum_{n\in\mathbb{Z},\ t< t^{(n)} < t+\tau} (t^{(n+1)} - t^{(n)}) \le \tau + \delta t,$$

we deduce from (85) that

$$\int_{\mathbb{R}} \left( \sum_{n=n(t)+1}^{n(t+\tau)} \delta t^{(n+\frac{1}{2})} \right) a^{(n(t+s)+1)} dt \le (\tau + \delta t) \sum_{m \in \mathbb{Z}} a^{(m+1)} \int_{t^{(m)}-s}^{t^{(m+1)}-s} dt 
= (\tau + \delta t) \sum_{m \in \mathbb{Z}} a^{(m+1)} \delta t^{(m+\frac{1}{2})},$$

which is exactly (84).

**Acknowledgements**: The authors would like to thank Clément Cancès for fruitful discussions on discrete compensated compactness theorems.

#### References

- [1] I. Aavatsmark, T. Barkve, O. Boe, and T. Mannseth. Discretization on non-orthogonal, quadrilateral grids for inhomogeneous, anisotropic media. *J. Comput. Phys.*, 127(1):2–14, 1996.
- [2] G. Akrivis, C. Makridakis, and R. H. Nochetto. Optimal order a posteriori error estimates for a class of Runge-Kutta and Galerkin methods. *Numer. Math.*, 114(1):133–160, 2009.

- [3] G. Akrivis, C. Makridakis, and R. H. Nochetto. Galerkin and Runge-Kutta methods: unified formulation, a posteriori error estimates and nodal superconvergence. *Numer. Math.*, 118(3):429–456, 2011.
- [4] H. Amann. Compact embeddings of vector-valued Sobolev and Besov spaces. *Glas. Mat. Ser. III*, 35(55)(1):161–177, 2000. Dedicated to the memory of Branko Najman.
- [5] B. Andreianov, F. Boyer, and F. Hubert. Discrete duality finite volume schemes for Leray-Lions-type elliptic problems on general 2D meshes. *Numer. Methods Partial Differential Equa*tions, 23(1):145–195, 2007.
- [6] B. Andreianov, C. Cancès, and A. Moussa. A nonlinear time compactness result and applications to discretization of degenerate parabolic-elliptic PDEs. HAL: hal-01142499. Submitted, 2015.
- [7] H. Attouch, G. Buttazzo, and G. Michaille. Variational analysis in Sobolev and BV spaces, volume 6 of MPS/SIAM Series on Optimization. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA; Mathematical Programming Society (MPS), Philadelphia, PA, 2006.
- [8] M. Bertsch, P. De Mottoni, and L. Peletier. The Stefan problem with heating: appearance and disappearance of a mushy region. *Trans. Amer. Math. Soc*, 293:677–691, 1986.
- [9] F. Brezzi and M. Fortin. Mixed and hybrid finite element methods, volume 15 of Springer Series in Computational Mathematics. Springer-Verlag, New York, 1991.
- [10] F. Brezzi, K. Lipnikov, and V. Simoncini. A family of mimetic finite difference methods on polygonal and polyhedral meshes. *Math. Models Methods Appl. Sci.*, 15(10):1533–1551, 2005.
- [11] X. Chen, A. Jüngel, and J.-G. Liu. A note on Aubin-Lions-Dubinskiĭ lemmas. *Acta Appl. Math.*, 133:33–43, 2014.
- [12] P. Ciarlet. The finite element method. In P. G. Ciarlet and J.-L. Lions, editors, *Part I*, Handbook of Numerical Analysis, III. North-Holland, Amsterdam, 1991.
- [13] Y. Coudière and F. Hubert. A 3d discrete duality finite volume method for nonlinear elliptic equations. SIAM Journal on Scientific Computing, 33(4):1739–1764, 2011.
- [14] M. Crouzeix and P.-A. Raviart. Conforming and nonconforming finite element methods for solving the stationary Stokes equations. I. Rev. Française Automat. Informat. Recherche Opérationnelle Sér. Rouge, 7(R-3):33-75, 1973.
- [15] K. Deimling. Nonlinear functional analysis. Springer-Verlag, Berlin, 1985.
- [16] J. Diaz and F. de Thelin. On a nonlinear parabolic problem arising in some models related to turbulent flows. SIAM J. Math. Anal., 25(4):1085–1111, 1994.
- [17] M. Dreher and A. Jüngel. Compact families of piecewise constant functions in  $L^p(0,T;B)$ . Nonlinear Anal., 75(6):3072–3077, 2012.
- [18] J. Droniou. Intégration et espaces de sobolev à valeurs vectorielles. Polycopiés de l'Ecole Doctorale de Mathématiques-Informatique de Marseille, available at http://www-gm3.univ-mrs.fr/polys, 2001.
- [19] J. Droniou. Finite volume schemes for fully non-linear elliptic equations in divergence form. ESAIM: Mathematical Modelling and Numerical Analysis, 40(6):1069, 2006.

- [20] J. Droniou and R. Eymard. A mixed finite volume scheme for anisotropic diffusion problems on any grid. *Numer. Math.*, 105(1):35–71, 2006.
- [21] J. Droniou, R. Eymard, T. Gallouët, C. Guichard, and R. Herbin. Gradient schemes for elliptic and parabolic problems. 2015. In preparation.
- [22] J. Droniou, R. Eymard, T. Gallouët, and R. Herbin. A unified approach to mimetic finite difference, hybrid finite volume and mixed finite volume methods. *Math. Models Methods Appl. Sci.*, 20(2):265–295, 2010.
- [23] J. Droniou, R. Eymard, T. Gallouët, and R. Herbin. Gradient schemes: a generic framework for the discretisation of linear, nonlinear and nonlocal elliptic and parabolic equations. *Math. Models Methods Appl. Sci.* (M3AS), 23(13):2395–2432, 2013.
- [24] J. Droniou, R. Eymard, and C. Guichard. Uniform-in-time convergence of numerical schemes for Richards' and Stefan's models. In *Finite Volumes for Complex Applications VII*. Springer, 2014.
- [25] M. G. Edwards and C. F. Rogers. Finite volume discretization with imposed flux continuity for the general tensor pressure equation. *Comput. Geosci.*, 2(4):259–290, 1998.
- [26] I. Ekeland and R. Témam. Convex analysis and variational problems, volume 28 of Classics in Applied Mathematics. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, english edition, 1999. Translated from the French.
- [27] A. Ern and J.-L. Guermond. *Theory and practice of finite elements*, volume 159. Springer, Berlin, 2004.
- [28] R. Eymard, P. Feron, T. Gallouët, R. Herbin, and C. Guichard. Gradient schemes for the Stefan problem. *International Journal On Finite Volumes*, 10s, 2013.
- [29] R. Eymard, T. Gallouët, and R. Herbin. Discretization of heterogeneous and anisotropic diffusion problems on general nonconforming meshes SUSHI: a scheme using stabilization and hybrid interfaces. *IMA J. Numer. Anal.*, 30(4):1009–1043, 2010.
- [30] R. Eymard, T. Gallouët, D. Hilhorst, and Y. Naït Slimane. Finite volumes and nonlinear diffusion equations. *RAIRO Modél. Math. Anal. Numér.*, 32(6):747–761, 1998.
- [31] R. Eymard, C. Guichard, and R. Herbin. Small-stencil 3d schemes for diffusive flows in porous media. M2AN, 46:265–290, 2012.
- [32] R. Eymard, C. Guichard, R. Herbin, and R. Masson. Gradient schemes for two-phase flow in heterogeneous porous media and Richards equation. ZAMM Z. Angew. Math. Mech., 94(7-8):560–585, 2014.
- [33] R. Eymard, M. Gutnic, and D. Hilhorst. The finite volume method for Richards equation. *Comput. Geosci.*, 3(3-4):259–294, 1999.
- [34] R. Eymard and R. Herbin. Gradient scheme approximations for diffusion problems. Finite Volumes for Complex Applications VI Problems & Perspectives, pages 439–447, 2011.
- [35] R. Eymard, R. Herbin, and A. Michel. Mathematical study of a petroleum-engineering scheme. M2AN Math. Model. Numer. Anal., 37(6):937–972, 2003.

- [36] T. Gallouët and J.-C. Latché. Compactness of discrete approximate solutions to parabolic PDEs—application to a turbulence model. *Commun. Pure Appl. Anal.*, 11(6):2371–2391, 2012.
- [37] R. Glowinski and J. Rappaz. Approximation of a nonlinear elliptic problem arising in a non-newtonian fluid flow model in glaciology. *M2AN Math. Model. Numer. Anal.*, 37(1):175–186, 2003.
- [38] C. González, A. Ostermann, C. Palencia, and M. Thalhammer. Backward Euler discretization of fully nonlinear parabolic problems. *Math. Comp.*, 71(237):125–145, 2002.
- [39] J. Gwinner and M. Thalhammer. Full discretisations for nonlinear evolutionary inequalities based on stiffly accurate Runge-Kutta and hp-finite element methods. Found. Comput. Math., 14(5):913–949, 2014.
- [40] F. Hermeline. Approximation of diffusion operators with discontinuous tensor coefficients on distorted meshes. Computer methods in applied mechanics and engineering, 192(16):1939–1959, 2003.
- [41] A. V. Kazhikhov. Recent developments in the global theory of two-dimensional compressible Navier-Stokes equations, volume 25 of Seminar on Mathematical Sciences. Keio University, Department of Mathematics, Yokohama, 1998.
- [42] C. Lubich and A. Ostermann. Runge-Kutta methods for parabolic equations and convolution quadrature. *Math. Comp.*, 60(201):105–131, 1993.
- [43] C. Lubich and A. Ostermann. Linearly implicit time discretization of non-linear parabolic equations. *IMA J. Numer. Anal.*, 15(4):555–583, 1995.
- [44] C. Lubich and A. Ostermann. Runge-Kutta approximation of quasi-linear parabolic equations. *Math. Comp.*, 64(210):601–627, 1995.
- [45] C. Lubich and A. Ostermann. Runge-Kutta time discretization of reaction-diffusion and Navier-Stokes equations: nonsmooth-data error estimates and applications to long-time behaviour. *Appl. Numer. Math.*, 22(1-3):279–292, 1996. Special issue celebrating the centenary of Runge-Kutta methods.
- [46] E. Maitre. Numerical analysis of nonlinear elliptic-parabolic equations. *M2AN Math. Model. Numer. Anal.*, 36(1):143–153, 2002.
- [47] G. Minty. On a monotonicity method for the solution of non-linear equations in Banach spaces. *Proceedings of the National Academy of Sciences of the United States of America*, 50(6):1038, 1963.
- [48] R. H. Nochetto and C. Verdi. Approximation of degenerate parabolic problems using numerical integration. SIAM J. Numer. Anal., 25(4):784–814, 1988.
- [49] A. Ostermann and M. Thalhammer. Convergence of Runge-Kutta methods for nonlinear parabolic equations. *Appl. Numer. Math.*, 42(1-3):367–380, 2002. Ninth Seminar on Numerical Solution of Differential and Differential-Algebraic Equations (Halle, 2000).
- [50] A. Ostermann, M. Thalhammer, and G. Kirlinger. Stability of linear multistep methods and applications to nonlinear parabolic problems. *Appl. Numer. Math.*, 48(3-4):389–407, 2004. Workshop on Innovative Time Integrators for PDEs.

- [51] I. S. Pop. Numerical schemes for degenerate parabolic problems. In *Progress in industrial mathematics at ECMI 2004*, volume 8 of *Math. Ind.*, pages 513–517. Springer, Berlin, 2006.
- [52] J. Rulla and N. J. Walkington. Optimal rates of convergence for degenerate parabolic problems in two dimensions. SIAM J. Numer. Anal., 33(1):56–67, 1996.