

Mathematical Analysis of Robustness of Two-Level Domain Decomposition Methods with respect to Inexact Coarse Solves

F. Nataf¹

¹Laboratoire J.L. Lions, UPMC, CNRS UMR7598, Equipe LJL-INRIA Alpines,
frederic.nataf@sorbonne-universite.fr, Paris, France

August 4, 2020

Contents

1	Introduction	2
2	Basic definitions	3
3	Inexact Coarse Solves for GenEO	5
4	Inexact Coarse Solves for GenEO2	10
	Auxiliary results on GEVP	13
5	Annex	17

Abstract

Convergence of domain decomposition methods rely heavily on the efficiency of the coarse space used in the second level. The GenEO coarse space has been shown to lead to a robust two-level Schwarz preconditioner which scales well over multiple cores [19, 2]. The robustness is due to its good approximation properties for problems with highly heterogeneous material parameters. It is available in the finite element packages FreeFem++ [9], Feel++ [17], Dune [1] and is implemented as a standalone library in HPDDM [10] and as such is available as well as a PETSc preconditioner. But the coarse component of the preconditioner can ultimately become a bottleneck if the number of subdomains is very large and exact solves are used. It is therefore interesting to consider the effect of inexact coarse solves. In this paper, robustness of GenEO methods is analyzed with respect to inexact coarse solves. Interestingly, the GenEO-2 method introduced in [7] has to be modified in order to be able to prove its robustness in this context.

1 Introduction

Convergence of domain decomposition methods rely heavily on the efficiency of the coarse space used in the second level, see [15, 22, 16] and references therein. These methods are based on two ingredients: a coarse space (CS) and a correction formula (see e.g. [21]). The GenEO coarse space introduced in [19] has been shown to lead to a robust two-level Schwarz preconditioner which scales well over multiple cores. The robustness is due to its good approximation properties for problems with highly heterogeneous material parameters. This approach is closely related to [4]. We refer to the introduction of [19] for more details on the differences and similarities between both approaches. Here we will mainly work with a slight modification of the GenEO CS introduced in [2] for the additive Schwarz method (see e.g. [22]) and the GenEO-2 CS introduced in [7] for the P.L. Lions algorithm [11]. These variants are easier to implement and in practice have similar performances although they may lead to a larger CS. More details are given in Annex 5 where we explain how to adapt the framework of [2] to the GenEO CS of [19].

We focus in this paper on a modification of the coarse component of the correction formula. Indeed, the coarse component of the preconditioner can ultimately become a bottleneck if the number of subdomains is very large and exact solves are used. It is therefore interesting to consider the effect of inexact coarse solves on the robustness. We show that the additive Schwarz method is naturally robust. Interestingly, the GenEO-2 method introduced in [7] has to be modified in order to be able to prove its robustness in this context. In the context of domain decomposition methods, the robustness of the BDDC w.r.t. inexact coarse solves has been studied in [23, 24] and in [13]. We focus here on GenEO methods. Compared to works on multilevel methods such as [25, 3] which are concerned with Schwarz multilevel methods where the coarse space is obtained by a coarse grid discretisation of the elliptic problem, we explicitly state robustness results of the two level method with respect to inexact coarse solves when the coarse space is obtained by the solution of local generalized eigenvalue problems. Moreover, we are not concerned only with Schwarz methods but also with P.L. Lions algorithm.

The general framework of our work is the following. Let M^{-1} be a one level preconditioner further enhanced by a second level correction based on a rectangular matrix Z whose columns are a basis of a coarse space V_0 . The coarse space correction is

$$Z(Z^T A Z)^{-1} Z^T, \quad (1)$$

and the coarse operator is defined by

$$E := Z^T A Z. \quad (2)$$

Let M^{-1} denote a one-level preconditioner, the hybrid two-level method is defined by:

$$M_2^{-1} := Z E^{-1} Z^T + (I - Z E^{-1} Z^T A) M^{-1} (I - A Z E^{-1} Z^T),$$

see the balancing domain decomposition method by J. Mandel [12]. This formula also appeared in an unpublished work by Schnabel [18], see [5] for more details on the connections between these works.

We consider Geneo methods, where the coarse space V_0 spanned by the columns of Z is built from solving generalized eigenvalue problems (GEVP) in the subdomains. Recall that these GEVP solves are purely parallel tasks with no communication involved. This part of the preconditioner setup is not penalizing parallelism. Actually, in strong scaling experiments where the number of degrees of freedom of subdomains is smaller and smaller, the elapsed time taken by these tasks will decrease. Thus, this task scales strongly. On the other hand, as the size of matrix $Z^T A Z$ typically increases linearly with the number of subdomains, the solving of the corresponding linear systems for instance with a LU factorization becomes a bottleneck in two-level domain decomposition methods. It is therefore interesting to estimate the robustness of the modified two-level method when in (2) the operator E is approximated by some operator \tilde{E} :

$$\boxed{\tilde{E} \simeq E},$$

since it paves the way to inexact coarse solves or to three or more level methods. Operator \tilde{E} may be obtained in many ways: approximate LU factorizations (e.g. ILU(k), ILU- ϵ or single precision factorization), Sparse Approximate Inverse, Krylov subspace recycling methods, multigrid methods and of course domain decomposition methods. In the latter case, we would have a multilevel method. Note that our results are expressed in terms of the spectral properties of $E\tilde{E}^{-1}$ so that an approximation method for which such results exist is preferable.

More precisely, formula (2) is modified and the preconditioner we study is defined by:

$$\tilde{M}_2^{-1} := Z \tilde{E}^{-1} Z^T + (I - Z \tilde{E}^{-1} Z^T A) M^{-1} (I - A Z \tilde{E}^{-1} Z^T).$$

and throughout the paper we make

Assumption 1.1 *The operator \tilde{E} is symmetric positive definite.*

2 Basic definitions

The problem to be solved is defined via a variational formulation on a domain $\Omega \subset \mathbb{R}^d$ for $d \in \mathbb{N}$:

$$\text{Find } u \in V \text{ such that } : a_\Omega(u, v) = l(v), \quad \forall v \in V,$$

where V is a Hilbert space of functions from Ω with real values. The problem we consider is given through a symmetric positive definite bilinear form a_Ω that is defined in terms of an integral over any open set $\omega \subset \Omega$. Typical examples are the heterogeneous diffusion equation (\mathbf{K} is a diffusion tensor)

$$a_\omega(u, v) := \int_\omega \mathbf{K} \nabla u \cdot \nabla v \, dx,$$

or the elasticity system (\mathbf{C} is the fourth-order stiffness tensor and $\varepsilon(\mathbf{u})$ is the strain tensor of a displacement field \mathbf{u}):

$$a_\omega(\mathbf{u}, \mathbf{v}) := \int_\omega \mathbf{C} : \varepsilon(\mathbf{u}) : \varepsilon(\mathbf{v}) \, dx.$$

The problem is discretized by a finite element method. Let \mathcal{N} denote the set of degrees of freedom and $(\phi_k)_{k \in \mathcal{N}}$ be a finite element basis on a mesh \mathcal{T}_h . Let $A \in \mathbb{R}^{\#\mathcal{N} \times \#\mathcal{N}}$ be the associated finite element matrix, $A_{kl} := a_\Omega(\phi_l, \phi_k)$, $k, l \in \mathcal{N}$. For some given right hand side $\mathbf{F} \in \mathbb{R}^{\#\mathcal{N}}$, we have to solve a linear system in \mathbf{U} of the form

$$A\mathbf{U} = \mathbf{F}.$$

Domain Ω is decomposed into N (overlapping or non overlapping) subdomains $(\Omega_i)_{1 \leq i \leq N}$ so that all subdomains are a union of cells of the mesh \mathcal{T}_h . This decomposition induces a natural decomposition of the set of indices \mathcal{N} into N subsets of indices $(\mathcal{N}_i)_{1 \leq i \leq N}$:

$$\mathcal{N}_i := \{k \in \mathcal{N} \mid \text{meas}(\text{supp}(\phi_k) \cap \Omega_i) > 0\}, \quad 1 \leq i \leq N. \quad (3)$$

For all $1 \leq i \leq N$, let R_i be the restriction matrix from $\mathbb{R}^{\#\mathcal{N}}$ to the subset $\mathbb{R}^{\#\mathcal{N}_i}$ and D_i be a diagonal matrix of size $\#\mathcal{N}_i \times \#\mathcal{N}_i$, so that we have a partition of unity at the algebraic level,

$$\sum_{i=1}^N R_i^T D_i R_i = I, \quad (4)$$

where $I \in \mathbb{R}^{\#\mathcal{N} \times \#\mathcal{N}}$ is the identity matrix.

We also define for all subdomains $1 \leq j \leq N$, \tilde{A}^j , the $\#\mathcal{N}_j \times \#\mathcal{N}_j$ matrix defined by

$$\mathbf{V}_j^T \tilde{A}^j \mathbf{U}_j := a_{\Omega_j} \left(\sum_{l \in \mathcal{N}_j} \mathbf{U}_{jl} \phi_l, \sum_{l \in \mathcal{N}_j} \mathbf{V}_{jl} \phi_l \right), \quad \mathbf{U}_j, \mathbf{V}_j \in \mathbb{R}^{\#\mathcal{N}_j}. \quad (5)$$

When the bilinear form a results from the variational solve of a Laplace problem, the previous matrix corresponds to the discretization of local Neumann boundary value problems. For this reason we will call it “Neumann” matrix even in a more general setting.

We also make use of two numbers k_0 and k_1 related to the domain decomposition. Let

$$k_0 := \max_{1 \leq i \leq N} \# \{j \mid R_j A R_i^T \neq 0\} \quad (6)$$

be the maximum multiplicity of the interaction between subdomains plus one. Let k_1 be the maximal multiplicity of subdomains intersection, i.e. the largest integer m such that there exists m different subdomains whose intersection has a non zero measure.

Let \tilde{P}_0 be defined as:

$$\boxed{\tilde{P}_0 := Z \tilde{E}^{-1} Z^T A}, \quad (7)$$

the operator \tilde{P}_0 is thus an approximation to the A -orthogonal projection on V_0

$$\boxed{P_0 := Z E^{-1} Z^T A}$$

which corresponds to an exact coarse solve.

Note that although \tilde{P}_0 is not a projection it has the same kernel and range as P_0 :

Lemma 2.1 *We have*

$$\ker P_0 = \ker \tilde{P}_0 = V_0^{A\perp} \quad \text{and} \quad \text{Im } P_0 = \text{Im } \tilde{P}_0 = V_0,$$

where $V_0^{A\perp}$ is the vector space A -orthogonal to V_0 , that is when $\mathbb{R}^{\#\mathcal{N}}$ is endowed with the scalar product induced by A : $(x, y)_A := (x, Ay)$.

Proof First note that the kernel of \tilde{P}_0 contains $\ker Z^T A$. On the other hand, we have:

$$\tilde{P}_0 x = Z \tilde{E}^{-1} Z^T A x = 0 \Rightarrow (Z \tilde{E}^{-1} Z^T A x, A x) = (\tilde{E}^{-1} Z^T A x, Z^T A x) = 0.$$

Since \tilde{E} is SPD, it means that $Z^T A x = 0$, that is $x \in \ker Z^T A$. We have thus $\ker \tilde{P}_0 = \ker Z^T A$. Note that

$$Z^T A x = 0 \Leftrightarrow \forall y \quad (A x, Z y) = 0 \Leftrightarrow x \in V_0^{A\perp}.$$

As for the image of \tilde{P}_0 , since the last operation in its definition is the multiplication by the matrix Z we have $\text{Im } \tilde{P}_0 \subset V_0$. Conversely, let $y \in V_0$, there exists β such that $y = Z \beta$. It is easy to check that $y = \tilde{P}_0 (Z (Z A Z)^{-1} \tilde{E} \beta)$. Thus, $\text{Im } \tilde{P}_0 = V_0$.

The same arguments hold if \tilde{E} is replaced by E . Thus, \tilde{P}_0 and P_0 have the same kernel and image. \blacksquare

3 Inexact Coarse Solves for GenEO

The GenEO coarse space was introduced in [19] and its slight modification is defined as follows, see [2]:

Definition 3.1 (Generalized Eigenvalue Problem for GenEO) *For each subdomain $1 \leq j \leq N$, we introduce the generalized eigenvalue problem*

$$\begin{aligned} \text{Find } (\mathbf{V}_{jk}, \tau_{jk}) \in \mathbb{R}^{\#\mathcal{N}_j} \setminus \{0\} \times \mathbb{R} \text{ such that} \\ D_j R_j A R_j^T D_j \mathbf{V}_{jk} = \tau_{jk} \tilde{A}^j \mathbf{V}_{jk}. \end{aligned} \quad (8)$$

Let $\tau > 0$ be a user-defined threshold, we define $V_{\text{genEO}}^\tau \subset \mathbb{R}^{\#\mathcal{N}}$ as the vector space spanned by the family of vectors $(R_j^T D_j \mathbf{V}_{jk})_{\tau_{jk} > \tau, 1 \leq j \leq N}$ corresponding to eigenvalues larger than τ .

Let $\tilde{\pi}_j$ be the projection from $\mathbb{R}^{\#\mathcal{N}_j}$ on $\text{Span}\{\mathbf{V}_{jk} | \tau_{jk} > \tau\}$ parallel to $\text{Span}\{\mathbf{V}_{jk} | \tau_{jk} \leq \tau\}$.

In this section, Z denotes a rectangular matrix whose columns are a basis of the coarse space V_{genEO}^τ defined in Definition (3.1). The dimension of Z is $\#\mathcal{N} \times \#\mathcal{N}_0$. The GenEO preconditioner with inexact coarse solve reads:

$$M_{\text{GenEOACS}}^{-1} := Z \tilde{E}^{-1} Z^T + (I - \tilde{P}_0) \left(\sum_{i=1}^N R_i^T (R_i A R_i^T)^{-1} R_i \right) (I - \tilde{P}_0^T). \quad (9)$$

The study the spectrum of $M_{\text{GenEOACS}}^{-1} A$ is based on the Fictitious Space lemma which is recalled here, see [14] for the original paper and [6] for a modern presentation.

Lemma 3.1 (Fictitious Space Lemma, Nepomnyaschikh 1991) *Let H and H_D be two Hilbert spaces, with the scalar products denoted by (\cdot, \cdot) and $(\cdot, \cdot)_D$. Let the symmetric positive bilinear forms $a : H \times H \rightarrow \mathbb{R}$ and $b : H_D \times H_D \rightarrow \mathbb{R}$, generated by the s.p.d. operators $A : H \rightarrow H$ and $B : H_D \rightarrow H_D$, respectively (i.e. $(Au, v) = a(u, v)$ for all $u, v \in H$ and $(Bu_D, v_D)_D = b(u_D, v_D)$ for all $u_D, v_D \in H_D$). Suppose that there exists a linear operator $\mathcal{R} : H_D \rightarrow H$ that satisfies the following three assumptions:*

(i) \mathcal{R} is surjective.

(ii) Continuity of \mathcal{R} : there exists a positive constant c_R such that

$$a(\mathcal{R}u_D, \mathcal{R}u_D) \leq c_R \cdot b(u_D, u_D) \quad \forall u_D \in H_D. \quad (10)$$

(iii) Stable decomposition: there exists a positive constant c_T such that for all $u \in H$ there exists $u_D \in H_D$ with $\mathcal{R}u_D = u$ and

$$c_T \cdot b(u_D, u_D) \leq a(\mathcal{R}u_D, \mathcal{R}u_D) = a(u, u). \quad (11)$$

We introduce the adjoint operator $\mathcal{R}^* : H \rightarrow H_D$ by $(\mathcal{R}u_D, u) = (u_D, \mathcal{R}^*u)_D$ for all $u_D \in H_D$ and $u \in H$.

Then, we have the following spectral estimate

$$c_T \cdot a(u, u) \leq a(\mathcal{R}B^{-1}\mathcal{R}^*Au, u) \leq c_R \cdot a(u, u), \quad \forall u \in H \quad (12)$$

which proves that the eigenvalues of operator $\mathcal{R}B^{-1}\mathcal{R}^*A$ are bounded from below by c_T and from above by c_R .

Loosely speaking, the first assumption corresponds to equation (2.3), page 36 of [22] where the global Hilbert space is assumed to satisfy a decomposition into subspaces. The second assumption is related to Assumptions 2.3 and 2.4, page 40 of [22]. The third assumption corresponds to the Stable decomposition Assumption 2.2 page 40 of [22].

In order to apply this lemma to the preconditioned operator $M_{GenEOACS}^{-1}A$, we introduce Hilbert spaces H and H_D as follows:

$$H := \mathbb{R}^{\#\mathcal{N}}$$

endowed with the bilinear form $a(\mathbf{U}, \mathbf{U}) := (A\mathbf{U}, \mathbf{U})$ and

$$H_D := \mathbb{R}^{\#\mathcal{N}_0} \times \prod_{i=1}^N \mathbb{R}^{\#\mathcal{N}_i}$$

endowed with the following bilinear form

$$\begin{aligned} \tilde{b} : H_D \times H_D &\longrightarrow \mathbb{R} \\ ((\mathbf{U}_0, (\mathbf{U}_i)_{1 \leq i \leq N}), (\mathbf{V}_0, (\mathbf{V}_i)_{1 \leq i \leq N})) &\longmapsto (\tilde{E}\mathbf{U}_0, \mathbf{V}_0) + (R_i A R_i^T \mathbf{U}_i, \mathbf{V}_i). \end{aligned} \quad (13)$$

We denote by $\tilde{B} : H_D \rightarrow H_D$ the operator such that $(\tilde{B}u_D, v_D)_D = \tilde{b}(u_D, v_D)$ for all $u_D, v_D \in H_D$.

Let $\tilde{\mathcal{R}} : H_D \rightarrow H$ be defined by

$$\tilde{\mathcal{R}}(\mathcal{U}) := Z\mathbf{U}_0 + (I - \tilde{P}_0) \sum_{i=1}^N R_i^T \mathbf{U}_i, \quad (14)$$

where $\mathcal{U} := (\mathbf{U}_0, (\mathbf{U}_i)_{1 \leq i \leq N})$. Recall that if we had used an exact coarse space solve, we would have introduced:

$$\mathcal{R}(\mathcal{U}) := Z \mathbf{U}_0 + (I - P_0) \sum_{i=1}^N R_i^T \mathbf{U}_i. \quad (15)$$

Note that we have

$$\tilde{\mathcal{R}}(\mathcal{U}) = \mathcal{R}(\mathcal{U}) + (P_0 - \tilde{P}_0) \sum_{i=1}^N R_i^T \mathbf{U}_i.$$

It can be checked that $M_{GenEOACS}^{-1} = \tilde{\mathcal{R}} \tilde{B}^{-1} \tilde{\mathcal{R}}^T$, see (9). In order to apply the fictitious space Lemma 3.1, three assumptions have to be checked.

- $\tilde{\mathcal{R}}$ is onto.

Let $\mathbf{U} \in H$, we have $\mathbf{U} = \tilde{P}_0 \mathbf{U} + (I - \tilde{P}_0) \mathbf{U}$. By Lemma 2.1, $\tilde{P}_0 \mathbf{U} \in V_0$ so that there exists $\mathbf{U}_0 \in \mathbb{R}^{\#\mathcal{N}_0}$ such that $\tilde{P}_0 \mathbf{U} = Z \mathbf{U}_0$. Owing to (4), we have

$$\mathbf{U} = Z \mathbf{U}_0 + (I - \tilde{P}_0) \sum_{i=1}^N R_i^T D_i R_i \mathbf{U} = \tilde{\mathcal{R}}(\mathbf{U}_0, (D_i R_i \mathbf{U})_{1 \leq i \leq N})$$

- Continuity of $\tilde{\mathcal{R}}$

We have to estimate a constant c_R such that for all $\mathcal{U} = (\mathbf{U}_0, (\mathbf{U}_i)_{1 \leq i \leq N}) \in H$, we have:

$$a(\tilde{\mathcal{R}}(\mathcal{U}), \tilde{\mathcal{R}}(\mathcal{U})) \leq c_R \tilde{b}(\mathcal{U}, \mathcal{U}).$$

Let δ be some positive number. Using that the image of $P_0 - \tilde{P}_0$ is a -orthogonal to the image of $I - P_0$, Cauchy-Schwarz inequality and the a -orthogonality of the projection $I - P_0$, we have:

$$\begin{aligned} a(\tilde{\mathcal{R}}(\mathcal{U}), \tilde{\mathcal{R}}(\mathcal{U})) &= \|\mathcal{R}(\mathcal{U}) + (P_0 - \tilde{P}_0) \sum_{i=1}^N R_i^T \mathbf{U}_i\|_A^2 \\ &= \|\mathcal{R}(\mathcal{U})\|_A^2 + 2a(Z \mathbf{U}_0 + (I - P_0) \sum_{i=1}^N R_i^T \mathbf{U}_i, (P_0 - \tilde{P}_0) \sum_{i=1}^N R_i^T \mathbf{U}_i) \\ &\quad + \|(P_0 - \tilde{P}_0) \sum_{i=1}^N R_i^T \mathbf{U}_i\|_A^2 \\ &= \|\mathcal{R}(\mathcal{U})\|_A^2 + 2a(Z \mathbf{U}_0, (P_0 - \tilde{P}_0) \sum_{i=1}^N R_i^T \mathbf{U}_i) \\ &\quad + \|(P_0 - \tilde{P}_0) \sum_{i=1}^N R_i^T \mathbf{U}_i\|_A^2 \\ &\leq \|\mathcal{R}(\mathcal{U})\|_A^2 + \delta \|Z \mathbf{U}_0\|_A^2 + \frac{1}{\delta} \|(P_0 - \tilde{P}_0) \sum_{i=1}^N R_i^T \mathbf{U}_i\|_A^2 \\ &\quad + \|(P_0 - \tilde{P}_0) \sum_{i=1}^N R_i^T \mathbf{U}_i\|_A^2 \\ &\leq \|Z \mathbf{U}_0\|_A^2 + \|\sum_{i=1}^N R_i^T \mathbf{U}_i\|_A^2 + \delta \|Z \mathbf{U}_0\|_A^2 \\ &\quad + (1 + \frac{1}{\delta}) \|(P_0 - \tilde{P}_0) \sum_{i=1}^N R_i^T \mathbf{U}_i\|_A^2 \\ &\leq (1 + \delta) \|Z \mathbf{U}_0\|_A^2 + (1 + \|P_0 - \tilde{P}_0\|_A^2 (1 + \frac{1}{\delta})) \|\sum_{i=1}^N R_i^T \mathbf{U}_i\|_A^2 \\ &\leq (1 + \delta) \lambda_{\max}(E \tilde{E}^{-1}) (\tilde{E} \mathbf{U}_0, \mathbf{U}_0) + (1 + \|P_0 - \tilde{P}_0\|_A^2 (1 + \frac{1}{\delta})) k_0 \sum_{i=1}^N \|R_i^T \mathbf{U}_i\|_A^2 \\ &\leq \max \left((1 + \delta) \lambda_{\max}(E \tilde{E}^{-1}), [1 + \|P_0 - \tilde{P}_0\|_A^2 (1 + \frac{1}{\delta})] k_0 \right) \tilde{b}(\mathcal{U}, \mathcal{U}). \end{aligned}$$

It is possible to minimize over δ the factor in front of $\tilde{b}(\mathcal{U}, \mathcal{U})$ using the

Lemma 3.2 *Let c, d, α and β be positive constant, we have*

$$\min_{\delta > 0} \max(c + \alpha\delta, d + \beta\delta^{-1}) = \frac{d + c + \sqrt{(d - c)^2 + 4\alpha\beta}}{2}.$$

Proof The optimal value for δ corresponds to the equality $c + \alpha\delta = d + \beta\delta^{-1}$. ■

Let

$$\epsilon_A := \|P_0 - \tilde{P}_0\|_A = \|Z^T((Z^T A Z)^{-1} - \tilde{E}^{-1})Z^T A\|_A, \quad (16)$$

the formula of Lemma 3.2 yields

$$c_R := \frac{k_0(1 + \epsilon_A^2) + \lambda_{\max}(E\tilde{E}^{-1}) + \sqrt{(k_0(1 + \epsilon_A^2) - \lambda_{\max}(E\tilde{E}^{-1}))^2 + 4\lambda_{\max}(E\tilde{E}^{-1})k_0\epsilon_A^2}}{2}. \quad (17)$$

Actually, ϵ_A can be expressed in term of the minimal eigenvalue of $E\tilde{E}^{-1}$.

Lemma 3.3 *Other formula for ϵ_A :*

$$\epsilon_A = \sup_{\mathbf{U}_0 \in \mathbb{R}^{\#\mathcal{N}_0}} \frac{(E(E^{-1} - \tilde{E}^{-1})E\mathbf{U}_0, \mathbf{U}_0)}{(E\mathbf{U}_0, \mathbf{U}_0)} = \max(|1 - \lambda_{\min}(E\tilde{E}^{-1})|, |1 - \lambda_{\max}(E\tilde{E}^{-1})|).$$

Proof Since $P_0 - \tilde{P}_0$ is A -symmetric, its norm is also given by

$$\epsilon_A = \sup_{\mathbf{U}} \frac{|((P_0 - \tilde{P}_0)\mathbf{U}, \mathbf{U})_A|}{\|\mathbf{U}\|_A^2}$$

We can go further by using the fact that P_0 is a A -orthogonal and that P_0 and \tilde{P}_0 have the same kernels and images:

$$\begin{aligned} \epsilon_A &= \sup_{\mathbf{U}} \frac{|((P_0 - \tilde{P}_0)(P_0 \mathbf{U} + (I - P_0)\mathbf{U}), P_0 \mathbf{U} + (I - P_0)\mathbf{U})_A|}{\|P_0 \mathbf{U}\|_A^2 + \|(I - P_0)\mathbf{U}\|_A^2} \\ &= \sup_{\mathbf{U}} \frac{|((P_0 - \tilde{P}_0)P_0 \mathbf{U}, P_0 \mathbf{U})_A|}{\|P_0 \mathbf{U}\|_A^2 + \|(I - P_0)\mathbf{U}\|_A^2} = \sup_{\mathbf{U}} \frac{|((P_0 - \tilde{P}_0)P_0 \mathbf{U}, P_0 \mathbf{U})_A|}{\|P_0 \mathbf{U}\|_A^2} \\ &= \sup_{\mathbf{U} \in V_0} \frac{|((P_0 - \tilde{P}_0)\mathbf{U}, \mathbf{U})_A|}{\|\mathbf{U}\|_A^2} = \sup_{\mathbf{U}_0 \in \mathbb{R}^{\#\mathcal{N}_0}} \frac{|(E(E^{-1} - \tilde{E}^{-1})E\mathbf{U}_0, \mathbf{U}_0)|}{(E\mathbf{U}_0, \mathbf{U}_0)} \\ &= \sup_{\mathbf{U}_0 \in \mathbb{R}^{\#\mathcal{N}_0}} \left| 1 - \frac{(\tilde{E}^{-1}E\mathbf{U}_0, E\mathbf{U}_0)}{(E\mathbf{U}_0, \mathbf{U}_0)} \right|. \end{aligned}$$

This means that formula (17) for c_R can be expressed explicitly in terms of k_0 and of the minimal and maximal eigenvalue of $\tilde{E}^{-1}E$. ■

- Stable decomposition

Let $\mathbf{U} \in H$ be decomposed as follows:

$$\begin{aligned} \mathbf{U} &= P_0 \mathbf{U} + (I - P_0)\mathbf{U} = P_0 \mathbf{U} + (I - P_0) \sum_{j=1}^N R_j^T D_j R_j \mathbf{U} \\ &= P_0 \mathbf{U} + (I - P_0) \sum_{j=1}^N R_j^T D_j (I - \tilde{\pi}_j) R_j \mathbf{U} + \underbrace{(I - P_0) \sum_{j=1}^N R_j^T D_j \tilde{\pi}_j R_j \mathbf{U}}_{=0} \\ &= \underbrace{P_0 \mathbf{U} + (\tilde{P}_0 - P_0) \sum_{j=1}^N R_j^T D_j (I - \tilde{\pi}_j) R_j \mathbf{U}}_{:= F\mathbf{U} \in V_0} + (I - \tilde{P}_0) \sum_{j=1}^N R_j^T D_j (I - \tilde{\pi}_j) R_j \mathbf{U}. \end{aligned}$$

Let $\mathbf{U}_0 \in \mathbb{R}^{\#\mathcal{N}_0}$ be such that $Z\mathbf{U}_0 = F\mathbf{U}$, we choose the following decomposition:

$$\mathbf{U} = \tilde{\mathcal{R}}(\mathbf{U}_0, (D_j(I - \tilde{\pi}_j)R_j\mathbf{U})_{1 \leq j \leq N}).$$

The stable decomposition consists in estimating a constant $c_T > 0$ such that:

$$c_T [(\tilde{E}\mathbf{U}_0, \mathbf{U}_0) + \sum_{j=1}^N (R_j A R_j^T D_j(I - \tilde{\pi}_j)R_j\mathbf{U}, D_j(I - \tilde{\pi}_j)R_j\mathbf{U})] \leq a(\mathbf{U}, \mathbf{U}). \quad (18)$$

Since the second term in the left hand side is the same as in the exact coarse solve method, we have (see [2], page 177, Lemma 7.15):

$$\sum_{j=1}^N (R_j A R_j^T D_j(I - \tilde{\pi}_j)R_j\mathbf{U}, D_j(I - \tilde{\pi}_j)R_j\mathbf{U}) \leq k_1 \tau a(\mathbf{U}, \mathbf{U}). \quad (19)$$

We now focus on the first term of the left hand side of (18). Let δ be some positive number, using again (19), the following auxiliary result holds:

$$\begin{aligned} \|F\mathbf{U}\|_A^2 &\leq (1 + \delta) \|P_0\mathbf{U}, P_0\mathbf{U}\|_A^2 \\ &\quad + (1 + \frac{1}{\delta}) \|(P_0 - \tilde{P}_0) \sum_{j=1}^N R_j^T D_j(I - \tilde{\pi}_j)R_j\mathbf{U}\|_A^2 \\ &\leq (1 + \delta) (A\mathbf{U}, \mathbf{U}) \\ &\quad + (1 + \frac{1}{\delta}) \|(P_0 - \tilde{P}_0)\|_A^2 \|\sum_{j=1}^N R_j^T D_j(I - \tilde{\pi}_j)R_j\mathbf{U}\|_A^2 \\ &\leq (1 + \delta) (A\mathbf{U}, \mathbf{U}) \\ &\quad + (1 + \frac{1}{\delta}) \|(P_0 - \tilde{P}_0)\|_A^2 k_0 \sum_{j=1}^N \|R_j^T D_j(I - \tilde{\pi}_j)R_j\mathbf{U}\|_A^2 \\ &\leq \left(1 + \delta + (1 + \frac{1}{\delta}) \|(P_0 - \tilde{P}_0)\|_A^2 k_0 k_1 \tau\right) a(\mathbf{U}, \mathbf{U}) \end{aligned}$$

The best possible value for δ is

$$\delta := \epsilon_A \sqrt{k_0 k_1 \tau}.$$

Hence, we have:

$$(Z^T A Z \mathbf{U}_0, \mathbf{U}_0) = \|F\mathbf{U}\|_A^2 \leq (1 + \epsilon_A \sqrt{k_0 k_1 \tau})^2 a(\mathbf{U}, \mathbf{U}). \quad (20)$$

Thus, we have:

$$\begin{aligned} (\tilde{E}\mathbf{U}_0, \mathbf{U}_0) &= (\tilde{E} E^{-1/2} E^{1/2} \mathbf{U}_0, E^{-1/2} E^{1/2} \mathbf{U}_0) = (E^{-1/2} \tilde{E} E^{-1/2} E^{1/2} \mathbf{U}_0, E^{1/2} \mathbf{U}_0) \\ &\leq \lambda_{\max}(E^{-1/2} \tilde{E} E^{-1/2}) (E^{1/2} \mathbf{U}_0, E^{1/2} \mathbf{U}_0) \\ &= \lambda_{\max}(E^{-1} \tilde{E}) (Z^T A Z \mathbf{U}_0, \mathbf{U}_0) \\ &\leq \lambda_{\max}(E^{-1} \tilde{E}) (1 + \epsilon_A \sqrt{k_0 k_1 \tau})^2 a(\mathbf{U}, \mathbf{U}). \end{aligned}$$

This last estimate along with (19) prove that in (18), it is possible to take

$$c_T = \frac{\lambda_{\min}(E \tilde{E}^{-1})}{(1 + \epsilon_A \sqrt{k_0 k_1 \tau})^2 + \lambda_{\min}(E \tilde{E}^{-1}) k_1 \tau}. \quad (21)$$

Overall, with c_T given by (21) and c_R by (17), we have proved the following spectral estimate:

$$c_T \leq \lambda(M_{GenEOACS}^{-1} A) \leq c_R. \quad (22)$$

Constants c_T and c_R are stable with respect to ϵ_A and the spectrum of $E \tilde{E}^{-1}$ so that (22) proves the stability of preconditioner $M_{GenEOACS}^{-1}$ w.r.t. inexact solves.

4 Inexact Coarse Solves for GenEO2

The GenEO-2 coarse space construction was introduced in [8, 7], see [2] also § 7.7, page 186. It is motivated by domain decomposition methods for which the local solves are not necessarily Dirichlet solves e.g. discretization of Robin boundary value problems, see [20]. We have not been able to prove the robustness of the GenEO-2 coarse space with respect to inexact coarse solves when used in the original GenEO-2 preconditioner (40), see remark 4.3. For this reason, we study here a slight modification of the preconditioner, eq. (28), for which we prove robustness. The more intricate analysis of GenEO2 compared to the one of GenEO is related to the differences between the Schwarz and P.L. Lions algorithms themselves. Indeed, in the Schwarz method, Assumption (ii) of the fictitious space lemma 3.1 comes almost for free even for a one level method whereas Assumption (iii) (stable decomposition) can only be fulfilled with a two level method. In P.L. Lions algorithm neither of the two assumptions are satisfied by the one level method. This is reflected in the fact that the proofs for GenEO2 are more intricate than for GenEO.

For all subdomains $1 \leq i \leq N$, let B_i be a matrix of size $\#\mathcal{N}_i \times \#\mathcal{N}_i$, which comes typically from the discretization of boundary value local problems using optimized transmission conditions or Neumann boundary conditions. Recall that by construction matrix $D_i R_i A R_i^T D_i$ is symmetric positive-semi definite and we make the extra following assumption:

Assumption 4.1 *For all subdomains $1 \leq i \leq N$, matrix B_i is symmetric positive semi-definite and either of the two conditions holds*

- B_i is definite,
- $B_i = \tilde{A}^i$ and $D_i R_i A R_i^T D_i$ is definite.

We first consider the case where B_i is definite. The other case will be treated in Remark 4.4. We recall the coarse space defined in [8, 7, 2]. Let γ and τ be two user defined thresholds. We introduce two generalized eigenvalue problems which by Assumption 4.1 are regular.

Definition 4.1 (Generalized Eigenvalue Problem for the lower bound)

For each subdomain $1 \leq j \leq N$, we introduce the generalized eigenvalue problem

$$\begin{aligned} \text{Find } (\mathbf{V}_{jk}, \lambda_{jk}) \in \mathbb{R}^{\#\mathcal{N}_j \setminus \{0\}} \times \mathbb{R} \text{ such that} \\ \tilde{A}^j \mathbf{V}_{jk} = \lambda_{jk} B_j \mathbf{V}_{jk} . \end{aligned} \quad (23)$$

Let $\tau > 0$ be a user-defined threshold and $\tilde{\pi}_j$ be the projection from $\mathbb{R}^{\#\mathcal{N}_j}$ on $V_{j\tau} = \text{Span}\{\mathbf{V}_{jk} | \lambda_{jk} < \tau\}$ parallel to $\text{Span}\{\mathbf{V}_{jk} | \lambda_{jk} \geq \tau\}$. We define $V_{j, \text{genEO}}^\tau \subset \mathbb{R}^{\#\mathcal{N}}$ as the vector space spanned by the family of vectors $(R_j^T D_j \mathbf{V}_{jk})_{\lambda_{jk} < \tau}$ corresponding to eigenvalues smaller than τ . Let V_{genEO}^τ be the vector space spanned by the collection over all subdomains of vector spaces $(V_{j, \text{genEO}}^\tau)_{1 \leq j \leq N}$.

Definition 4.2 (Generalized Eigenvalue Problem for the upper bound)

For each subdomain $1 \leq i \leq N$, we introduce the generalized eigenvalue problem

$$\begin{aligned} \text{Find } (\mathbf{U}_{ik}, \mu_{ik}) \in \mathbb{R}^{\#\mathcal{N}_i \setminus \{0\}} \times \mathbb{R} \text{ such that} \\ D_i R_i A R_i^T D_i \mathbf{U}_{ik} = \mu_{ik} B_i \mathbf{U}_{ik} . \end{aligned} \quad (24)$$

Let $\gamma > 0$ be a user-defined threshold, we define $V_{i, \text{geneo}}^\gamma \subset \mathbb{R}^{\#\mathcal{N}}$ as the vector space spanned by the family of vectors $(R_i^T D_i \mathbf{U}_{ik})_{\mu_{ik} > \gamma}$ corresponding to eigenvalues larger than γ . Let V_{geneo}^γ be the vector space spanned by the collection over all subdomains of vector spaces $(V_{j, \text{geneo}}^\gamma)_{1 \leq j \leq N}$.

Now, let ξ_i denote the B_i -orthogonal projection from $\mathbb{R}^{\#\mathcal{N}_i}$ on

$$V_{i, \gamma} := \text{Span} \{ \mathbf{U}_{ik} \mid \gamma < \mu_{ik} \}$$

parallel to

$$W_{i, \gamma} := \text{Span} \{ \mathbf{U}_{ik} \mid \gamma \geq \mu_{ik} \}.$$

The coarse space V_0 built from the above generalized eigenvalues is defined as the following sum:

$$V_0 := V_{\text{geneo}}^\tau + V_{\text{geneo}}^\gamma.$$

It is spanned by the columns of a full rank rectangular matrix $Z = R_0^T$ with $\#\mathcal{N}_0$ columns. Projection P_0 and its approximation \tilde{P}_0 are defined by the same formula as above, see (7).

We have the following

Lemma 4.1 *For $1 \leq j \leq N$, let us introduce the B_j -orthogonal projection p_j from $\mathbb{R}^{\#\mathcal{N}_j}$ on*

$$V_{j, \tau\gamma} := V_{j, \tau} \oplus V_{j, \gamma}.$$

Then for all $\mathbf{U}_j \in \mathbb{R}^{\#\mathcal{N}_j}$, we have:

$$\tau (B_j(I - p_j)\mathbf{U}_j, (I - p_j)\mathbf{U}_j) \leq (\tilde{A}_j \mathbf{U}_j, \mathbf{U}_j).$$

Moreover, for all $\mathbf{U} \in \mathbb{R}^{\#\mathcal{N}}$, we have:

$$\tau \sum_{j=1}^N (B_j(I - p_j)R_j \mathbf{U}, (I - p_j)R_j \mathbf{U}) \leq k_1 a(\mathbf{U}, \mathbf{U}).$$

Proof Let $\mathbf{U}_j \in \mathbb{R}^{\#\mathcal{N}_j}$, we have:

$$\begin{aligned} (B_j(I - \tilde{\pi}_j)\mathbf{U}_j, (I - \tilde{\pi}_j)\mathbf{U}_j) &= (B_j(I - p_j + (p_j - \tilde{\pi}_j))\mathbf{U}_j, (I - p_j + (p_j - \tilde{\pi}_j))\mathbf{U}_j) \\ &= \|(I - p_j)\mathbf{U}_j\|_{B_j}^2 + \|(p_j - \tilde{\pi}_j)\mathbf{U}_j\|_{B_j}^2 \\ &\quad + 2 \underbrace{(B_j(I - p_j)\mathbf{U}_j, (p_j - \tilde{\pi}_j)\mathbf{U}_j)}_{=0 \text{ since } \tilde{\pi}_j \mathbf{U}_j \in V_{j, \tau} \subset V_{j, \tau\gamma}} \\ &\geq \|(I - p_j)\mathbf{U}_j\|_{B_j}^2 = (B_j(I - p_j)\mathbf{U}_j, (I - p_j)\mathbf{U}_j). \end{aligned}$$

Since we have by Lemma 7.6, page 167 in [2] :

$$\tau (B_j(I - \tilde{\pi}_j)\mathbf{U}_j, (I - \tilde{\pi}_j)\mathbf{U}_j) \leq (\tilde{A}_j \mathbf{U}_j, \mathbf{U}_j),$$

the conclusion follows by summation over all subdomains. \blacksquare

The definition of the stable preconditioner is based on a pseudo inverse of B_i that we introduce now. Let b_{W_i} denote the restriction of b_i to $W_{i, \gamma} \times W_{i, \gamma}$ where $W_{i, \gamma}$ is endowed with the Euclidean scalar product:

$$\begin{aligned} b_{W_i} : W_{i, \gamma} \times W_{i, \gamma} &\longrightarrow \mathbb{R} \\ (\mathbf{U}_i, \mathbf{V}_i) &\mapsto b_i(\mathbf{U}_i, \mathbf{V}_i). \end{aligned} \tag{25}$$

By Riesz representation theorem, there exists a unique isomorphism $B_{W_i} : W_{i\gamma} \longrightarrow W_{i\gamma}$ into itself so that for all $\mathbf{U}_i, \mathbf{V}_i \in W_{i\gamma}$, we have:

$$b_{W_i}(\mathbf{U}_i, \mathbf{V}_i) = (B_{W_i} \mathbf{U}_i, \mathbf{V}_i).$$

The inverse of B_{W_i} will be denoted by B_i^\dagger and is given by the following formula

$$B_i^\dagger = (I - \xi_i) B_i^{-1}. \quad (26)$$

In order to check this formula, we have to show that $B_{W_i}(I - \xi_i)B_i^{-1}y = y$ for all $y \in W_{i\gamma}$. Let $z \in W_{i\gamma}$, using the fact that $I - \xi_i$ is the b_i -orthogonal projection on $W_{i\gamma}$, we have:

$$(B_{W_i}(I - \xi_i)B_i^{-1}y, z) = b_i((I - \xi_i)B_i^{-1}y, z) = b_i(B_i^{-1}y, z) = (y, z). \quad (27)$$

Since this equality holds for any $z \in W_{i\gamma}$, this proves that $B_{W_i}(I - \xi_i)B_i^{-1}y = y$.

We study now the preconditioner given by:

Definition 4.3 (Preconditioner $M_{GenEO2ACS}^{-1}$) Let q_i denote the orthogonal projection (for the Euclidean scalar product) from $\mathbb{R}^{\#N_i}$ onto $W_{i\gamma}$. We define the preconditioner $M_{GenEO2ACS}^{-1}$ as follows:

$$M_{GenEO2ACS}^{-1} := Z \tilde{E}^{-1} Z^T + (I - \tilde{P}_0) \left(\sum_{i=1}^N R_i^T D_i q_i B_i^\dagger q_i D_i R_i \right) (I - \tilde{P}_0^T). \quad (28)$$

Remark 4.1 In order to write an explicit form for the projection q_i , we denote by $Z_{i\gamma}$ a rectangular matrix whose columns are a basis for $V_{i\gamma}$. Let $\mathbf{U}_i \in \mathbb{R}^{\#N_i}$ be a vector we want to project. The projection $q_i \mathbf{U}_i$ is the solution to the constrained minimization problem:

$$\min_{\mathbf{W}_i \in W_{i\gamma}} \frac{1}{2} \|\mathbf{W}_i - \mathbf{U}_i\|^2 = \min_{\mathbf{W}_i | (B_i Z_{i\gamma})^T \mathbf{W}_i = 0} \frac{1}{2} \|\mathbf{W}_i - \mathbf{U}_i\|^2$$

Using the Lagrange multiplier technique, we introduce $\lambda \in \mathbb{R}^{\dim(V_{i\gamma})}$ and the optimality conditions read:

$$\begin{aligned} (q_i \mathbf{U}_i - \mathbf{U}_i) + B_i Z_{i\gamma} \lambda &= 0 \\ (B_i Z_{i\gamma})^T q_i \mathbf{U}_i &= 0. \end{aligned}$$

The vector λ must satisfy

$$-(B_i Z_{i\gamma})^T \mathbf{U}_i + (B_i Z_{i\gamma})^T B_i Z_{i\gamma} \lambda = 0.$$

Finally, an explicit formula for the projection q_i is:

$$q_i = I - B_i Z_{i\gamma} ((B_i Z_{i\gamma})^T B_i Z_{i\gamma})^{-1} Z_{i\gamma}^T B_i^T.$$

Thus applying q_i amounts to solving concurrently in each subdomain a small linear system of size the number of local eigenvectors contributing to the coarse space.

Remark 4.2 Note that $q_i B_i^\dagger$ is actually equal to B_i^\dagger but its presence shows the symmetry of the preconditioner.

We can now define the abstract framework for the preconditioner. Let H_D be defined by

$$H_D := \mathbb{R}^{\#\mathcal{N}_0} \times \Pi_{i=1}^N W_{i\gamma}$$

endowed with the following bilinear form arising from local SPD matrices $(B_i)_{1 \leq i \leq N}$

$$\begin{aligned} \tilde{b} : H_D \times H_D &\longrightarrow \mathbb{R} \\ (\mathcal{U}, \mathcal{V}) &\longmapsto b(\mathcal{U}, \mathcal{V}) := (\tilde{E}\mathbf{U}_0, \mathbf{V}_0) + \sum_{i=1}^N (B_i \mathbf{U}_i, \mathbf{V}_i) \end{aligned} \quad (29)$$

We denote by $\tilde{B} : H_D \rightarrow H_D$ the operator such that $(\tilde{B}u_D, v_D)_D = \tilde{b}(u_D, v_D)$ for all $u_D, v_D \in H_D$.

Let $\tilde{\mathcal{R}} : H_D \rightarrow H$ be defined using operator \tilde{P}_0 (see eq. (7)):

$$\tilde{\mathcal{R}}(\mathcal{U}) := Z \mathbf{U}_0 + (I - \tilde{P}_0) \sum_{i=1}^N R_i^T D_i \mathbf{U}_i. \quad (30)$$

Recall that if we had used an exact coarse space solve, we would have introduced:

$$\mathcal{R}(\mathcal{U}) := Z \mathbf{U}_0 + (I - P_0) \sum_{i=1}^N R_i^T D_i \mathbf{U}_i. \quad (31)$$

Note that we have

$$\tilde{\mathcal{R}}(\mathcal{U}) = \mathcal{R}(\mathcal{U}) + (P_0 - \tilde{P}_0) \sum_{i=1}^N R_i^T D_i \mathbf{U}_i.$$

It can be checked that the resulting preconditioner with inexact coarse solve $M_{GenEO2ACS}^{-1}$ (Eq. (28)) satisfies $M_{GenEO2ACS}^{-1} = \tilde{\mathcal{R}} \tilde{B}^{-1} \tilde{\mathcal{R}}^T$. Indeed, we have:

$$\tilde{\mathcal{R}}^T \mathbf{V} = (Z^T \mathbf{V}, (q_i D_i R_i (I - \tilde{P}_0^T) \mathbf{V})_{1 \leq i \leq N})$$

Auxiliary results on GEVP Beware, in this paragraph, A and B have nothing to do with the global problem to be solved:

Lemma 4.2 Let A be a symmetric positive semi definite matrix and B be a symmetric positive definite matrix. We consider the generalized eigenvalue problem:

$$A\mathbf{U} = \lambda B\mathbf{U}.$$

The generalized eigenvectors and eigenvalues are denoted by $(\mathbf{U}_k, \lambda_k)_{k \geq 1}$. Let τ be a positive number. We define

$$V_\tau := \text{Span}\{\mathbf{U}_k \mid \lambda_k < \tau\}.$$

Let W be any linear subspace. We denote by p the B -orthogonal projection on $V_\tau + W$.

Then, for all \mathbf{U} we have the following estimate:

$$\tau (B(I-p)\mathbf{U}, (I-p)\mathbf{U}) \leq (A(I-p)\mathbf{U}, (I-p)\mathbf{U}). \quad (32)$$

Similarly, let γ be a positive number. We define

$$V_\gamma := \text{Span}\{\mathbf{U}_k \mid \lambda_k > \gamma\}.$$

Let W be any linear subspace. We denote by q the B -orthogonal projection on $V_\gamma + W$.

Then, for all \mathbf{U} we have the following estimate:

$$(A(I - q)\mathbf{U}, (I - q)\mathbf{U}) \leq \gamma (B(I - q)\mathbf{U}, (I - q)\mathbf{U}). \quad (33)$$

Proof We have using $V_\tau \subset V_\tau + W$:

$$\tau \leq \min_{\mathbf{U} \in V_\tau^{B\perp}} \frac{(A\mathbf{U}, \mathbf{U})}{(B\mathbf{U}, \mathbf{U})} \leq \min_{\mathbf{U} \in (V_\tau + W)^{B\perp}} \frac{(A\mathbf{U}, \mathbf{U})}{(B\mathbf{U}, \mathbf{U})}.$$

For all \mathbf{U} , the vector $(I - p)\mathbf{U}$ is B -orthogonal to $V_\tau + W$ and this ends the proof of (32). The proof of (33) follows similarly from

$$\gamma \geq \max_{\mathbf{U} \in V_\gamma^{B\perp}} \frac{(A\mathbf{U}, \mathbf{U})}{(B\mathbf{U}, \mathbf{U})}.$$

■

In order to apply the fictitious space Lemma 3.1 to the study of the preconditioner (28), three assumptions have to be checked.

- $\tilde{\mathcal{R}}$ is onto.

Let $\mathbf{U} \in H$, we have

$$\begin{aligned} \mathbf{U} &= \tilde{P}_0 \mathbf{U} + (I - \tilde{P}_0) \mathbf{U} \\ &= \tilde{P}_0 \mathbf{U} + (I - \tilde{P}_0) \sum_{i=1}^N R_i^T D_i R_i \mathbf{U} \\ &= \tilde{P}_0 \mathbf{U} + (I - \tilde{P}_0) \sum_{i=1}^N R_i^T D_i \xi_i R_i \mathbf{U} + (I - \tilde{P}_0) \sum_{i=1}^N R_i^T D_i (I - \xi_i) R_i \mathbf{U} \\ &= \tilde{P}_0 \mathbf{U} + \underbrace{(P_0 - \tilde{P}_0) \sum_{i=1}^N R_i^T D_i \xi_i R_i \mathbf{U}}_{:=F\mathbf{U}} + \underbrace{(I - P_0) \sum_{i=1}^N R_i^T D_i \xi_i R_i \mathbf{U}}_{=0} \\ &\quad + (I - \tilde{P}_0) \sum_{i=1}^N R_i^T D_i (I - \xi_i) R_i \mathbf{U}. \end{aligned}$$

Let us consider the last equality. Since $F\mathbf{U}$ is the sum of two terms that belong to \mathbf{V}_0 there exists \mathbf{U}_0 such that $Z\mathbf{U}_0 = F\mathbf{U}$. The third term is zero since $\sum_{i=1}^N R_i^T D_i \xi_i R_i \mathbf{U} \in V_0$. Note also that $(I - \xi_i) R_i \mathbf{U} \in W_{i\gamma}$. Therefore, we have

$$\mathbf{U} = \tilde{\mathcal{R}}(\mathbf{U}_0, ((I - \xi_i) R_i \mathbf{U})_{1 \leq i \leq N}).$$

- Continuity of $\tilde{\mathcal{R}}$

We have to estimate a constant c_R such that for all $\mathcal{U} = (\mathbf{U}_0, (\mathbf{U}_i)_{1 \leq i \leq N}) \in H_D$ we have:

$$\begin{aligned} a(\tilde{\mathcal{R}}(\mathcal{U}), \tilde{\mathcal{R}}(\mathcal{U})) &\leq c_R \tilde{b}(\mathcal{U}, \mathcal{U}) \\ &= c_R [(\tilde{E}\mathbf{U}_0, \mathbf{U}_0) + \sum_{i=1}^N (B_i \mathbf{U}_i, \mathbf{U}_i)]. \end{aligned}$$

Note that using $(I - \xi_i)\mathbf{U}_i = \mathbf{U}_i$ (recall that $\mathbf{U}_i \in W_{i\gamma}$), we have:

$$\begin{aligned}
\tilde{\mathcal{R}}(\mathcal{U}) &= Z\mathbf{U}_0 + (I - \tilde{P}_0) \sum_{i=1}^N R_i^T D_i \mathbf{U}_i \\
&= Z\mathbf{U}_0 + (P_0 - \tilde{P}_0) \sum_{i=1}^N R_i^T D_i \mathbf{U}_i + (I - P_0) \sum_{i=1}^N R_i^T D_i \mathbf{U}_i \\
&= \underbrace{Z\mathbf{U}_0 + (P_0 - \tilde{P}_0) \sum_{i=1}^N R_i^T D_i (I - \xi_i)\mathbf{U}_i}_{\in V_0} + (I - P_0) \sum_{i=1}^N R_i^T D_i (I - \xi_i)\mathbf{U}_i
\end{aligned}$$

We have thus the following estimate using the A -orthogonality of $I - P_0$:

$$\begin{aligned}
a(\tilde{\mathcal{R}}(\mathcal{U}), \tilde{\mathcal{R}}(\mathcal{U})) &= \|Z\mathbf{U}_0 + (P_0 - \tilde{P}_0) \sum_{i=1}^N R_i^T D_i (I - \xi_i)\mathbf{U}_i \\
&\quad + (I - P_0) \sum_{i=1}^N R_i^T D_i (I - \xi_i)\mathbf{U}_i\|_A^2 \\
&= \|Z\mathbf{U}_0 + (P_0 - \tilde{P}_0) \sum_{i=1}^N R_i^T D_i (I - \xi_i)\mathbf{U}_i\|_A^2 \\
&\quad + \|(I - P_0) \sum_{i=1}^N R_i^T D_i (I - \xi_i)\mathbf{U}_i\|_A^2 \\
&\leq (1 + \delta) \|Z\mathbf{U}_0\|_A^2 + (1 + \frac{1}{\delta}) \|(P_0 - \tilde{P}_0) \sum_{i=1}^N R_i^T D_i (I - \xi_i)\mathbf{U}_i\|_A^2 \\
&\quad + \|\sum_{i=1}^N R_i^T D_i (I - \xi_i)\mathbf{U}_i\|_A^2 \\
&\leq (1 + \delta) (E\mathbf{U}_0, \mathbf{U}_0) + k_0 \sum_{i=1}^N \|R_i^T D_i (I - \xi_i)\mathbf{U}_i\|_A^2 \\
&\quad + (1 + \frac{1}{\delta}) \|(P_0 - \tilde{P}_0)\|_A^2 k_0 \sum_{i=1}^N \|R_i^T D_i (I - \xi_i)\mathbf{U}_i\|_A^2 \\
&\leq (1 + \delta) \lambda_{\max}(E\tilde{E}^{-1}) (\tilde{E}\mathbf{U}_0, \mathbf{U}_0) \\
&\quad + k_0 \gamma (1 + (1 + \frac{1}{\delta}) \|(P_0 - \tilde{P}_0)\|_A^2) \sum_{i=1}^N (B_i (I - \xi_i)\mathbf{U}_i, (I - \xi_i)\mathbf{U}_i) \\
&\leq \max((1 + \delta) \lambda_{\max}(E\tilde{E}^{-1}), k_0 \gamma (1 + (1 + \frac{1}{\delta}) \epsilon_A^2)) \tilde{b}(\mathcal{U}, \mathcal{U}).
\end{aligned}$$

Based on Lemma 3.2, we can optimize the value of δ and take

$$c_R := \frac{k_0 \gamma (1 + \epsilon_A^2) + \lambda_{\max}(E\tilde{E}^{-1}) + \sqrt{(k_0 \gamma (1 + \epsilon_A^2) - \lambda_{\max}(E\tilde{E}^{-1}))^2 + 4\lambda_{\max}(E\tilde{E}^{-1})k_0\gamma\epsilon_A^2}}{2}. \quad (34)$$

- Stable decomposition

The stable decomposition estimate is based on using projections p_j defined in Lemma 4.1. Let $\mathbf{U} \in H$ be decomposed as follows:

$$\begin{aligned}
\mathbf{U} &= P_0\mathbf{U} + (I - P_0) \sum_{j=1}^N R_j^T D_j (I - p_j) R_j \mathbf{U} + \underbrace{(I - P_0) \sum_{j=1}^N R_j^T D_j p_j R_j \mathbf{U}}_{=0} \\
&= \underbrace{P_0\mathbf{U} + (\tilde{P}_0 - P_0) \sum_{j=1}^N R_j^T D_j (I - p_j) R_j \mathbf{U}}_{:= F\mathbf{U} \in V_0} + (I - \tilde{P}_0) \sum_{j=1}^N R_j^T D_j (I - p_j) R_j \mathbf{U}.
\end{aligned}$$

We define \mathbf{U}_0 be such that $Z\mathbf{U}_0 = F\mathbf{U}$. We have that $(I - p_j)R_j \mathbf{U}$ is B_j -orthogonal to $V_{\gamma j} + V_{\tau j}$ and thus to $V_{\gamma j}$. This means that $(I - p_j)R_j \mathbf{U} \in W_{\gamma j}$ and that we can choose the following decomposition:

$$\mathbf{U} = \tilde{\mathcal{R}}(\mathbf{U}_0, ((I - p_j)R_j \mathbf{U})_{1 \leq j \leq N}).$$

The stability of the decomposition consists in estimating a constant $c_T > 0$ such that :

$$c_T [(\tilde{E} \mathbf{U}_0, \mathbf{U}_0) + \sum_{j=1}^N (B_j(I - p_j) R_j \mathbf{U}, (I - p_j) R_j \mathbf{U})] \leq a(\mathbf{U}, \mathbf{U}). \quad (35)$$

Using Lemma 4.1, we have

$$\tau \sum_{j=1}^N (B_j(I - p_j) R_j \mathbf{U}, (I - p_j) R_j \mathbf{U}) \leq k_1 a(\mathbf{U}, \mathbf{U}). \quad (36)$$

We now focus on the first term of the left hand side of (35). Let δ be some positive number, the following auxiliary result will be useful:

$$\begin{aligned} \|F\mathbf{U}\|_A^2 &\leq (1 + \delta) \|P_0 \mathbf{U}, P_0 \mathbf{U}\|_A^2 \\ &\quad + (1 + \tfrac{1}{\delta}) \|(P_0 - \tilde{P}_0) \sum_{j=1}^N R_j^T D_j (I - p_j) R_j \mathbf{U}\|_A^2 \\ &\leq (1 + \delta) (A\mathbf{U}, \mathbf{U}) \\ &\quad + (1 + \tfrac{1}{\delta}) \|(P_0 - \tilde{P}_0)\|_A^2 \|\sum_{j=1}^N R_j^T D_j (I - p_j) R_j \mathbf{U}\|_A^2 \\ &\leq (1 + \delta) a(\mathbf{U}, \mathbf{U}) \\ &\quad + (1 + \tfrac{1}{\delta}) \|(P_0 - \tilde{P}_0)\|_A^2 k_0 \sum_{j=1}^N \|R_j^T D_j (I - p_j) R_j \mathbf{U}\|_A^2 \\ &\leq (1 + \delta) a(\mathbf{U}, \mathbf{U}) \\ &\quad + (1 + \tfrac{1}{\delta}) \|(P_0 - \tilde{P}_0)\|_A^2 k_0 \gamma \sum_{j=1}^N (B_j(I - p_j) R_j \mathbf{U}, (I - p_j) R_j \mathbf{U}) \\ &\leq ((1 + \delta) + (1 + \tfrac{1}{\delta}) \|(P_0 - \tilde{P}_0)\|_A^2 k_0 \gamma \tau^{-1} k_1) a(\mathbf{U}, \mathbf{U}) \end{aligned}$$

where we have used Lemma 4.2 (33) (applied with A replaced by $D_j R_j A R_j^T D_j$ and B by B_j) for the one before last estimate and Lemma 4.1 for the last estimate.

The optimal value for δ yields:

$$\|F\mathbf{U}\|_A^2 \leq (1 + \epsilon_A \sqrt{k_0 k_1 \gamma \tau^{-1}})^2 a(\mathbf{U}, \mathbf{U}). \quad (37)$$

We have

$$\begin{aligned} (\tilde{E} \mathbf{U}_0, \mathbf{U}_0) &\leq \lambda_{\max}(E^{-1} \tilde{E})(E \mathbf{U}_0, \mathbf{U}_0) = \lambda_{\max}(E^{-1} \tilde{E})(A Z \mathbf{U}_0, Z \mathbf{U}_0) \\ &= \lambda_{\max}(E^{-1} \tilde{E}) \|F\mathbf{U}\|_A^2. \end{aligned}$$

so that with (37), this yields:

$$(\tilde{E} \mathbf{U}_0, \mathbf{U}_0) \leq \lambda_{\max}(E^{-1} \tilde{E}) (1 + \epsilon_A \sqrt{k_0 k_1 \gamma \tau^{-1}})^2 a(\mathbf{U}, \mathbf{U}).$$

Finally, in (35) we can take :

$$c_T := \frac{1}{\lambda_{\max}(E^{-1} \tilde{E}) (1 + \epsilon_A \sqrt{k_0 k_1 \gamma \tau^{-1}})^2 + k_1 \tau^{-1}}. \quad (38)$$

Overall, with c_T given by (38) and c_R by (34), we have proved the following spectral estimate:

$$c_T \leq \lambda(M_{GenEO2ACS}^{-1} A) \leq c_R. \quad (39)$$

Constants c_T and c_R are stable with respect to ϵ_A and the spectrum of $E\tilde{E}^{-1}$ so that (39) proves the stability of preconditioner $M_{GenEO2ACS}^{-1}$ (28) w.r.t. inexact solves.

Remark 4.3 *Had we taken the GenEO-2 algorithm introduced in [7] and modified only the coarse space solves:*

$$\tilde{M}_{GenEO,2}^{-1} = Z \tilde{E}^{-1} Z^T + (I - \tilde{P}_0) \left(\sum_{i=1}^N R_i^T D_i B_i^{-1} D_i R_i \right) (I - \tilde{P}_0^T), \quad (40)$$

the estimate for the upper bound of the preconditioned system would be for arbitrary $\delta > 0$

$$\lambda_{max} \leq \max(1 + \delta, k_0 \gamma + (1 + \frac{1}{\delta}) \epsilon_A^2 k_0 \max_{1 \leq i \leq N} \lambda_{max}(B_i^{-1} D_i R_i A R_i^T D_i)^2)$$

and would depend on the product of ϵ_A with the largest eigenvalue of the local operators $B_i^{-1} D_i R_i A R_i^T D_i$. This last term can be very large and we were not able to guarantee robustness with respect to approximate coarse solves.

Remark 4.4 *If for some subdomain i , $1 \leq i \leq N$, $B_i = \tilde{A}_i$ and \tilde{A}_i is symmetric positive semi-definite and $D_i R_i A R_i^T D_i$ is SPD, the eigenvalue problem (23) will not contribute to the coarse space. More precisely, the contribution of the subdomain to the coarse space involves (24) and will be $R_i^T D_i \ker(\tilde{A}_i) \oplus V_{i,genEO}^\gamma$. Also in Definition 4.3, B_i^\dagger is the pseudo inverse of B_i where $W_{i\gamma}$ is the image of B_i which is orthogonal to $\ker(\tilde{A}_i)$.*

5 Annex

We explain here how to adapt the GenEO coarse space as defined in [2] so that it will behave as the one defined in [19]. In one sentence, it consists simply in computing the local components of the coarse space on the subdomain extended by one layer (or more).

More precisely, we start from a domain decomposition $(\Omega_i)_{1 \leq i \leq N}$ and inherited indices decomposition $(\mathcal{N}_i)_{1 \leq i \leq N}$ as defined in the present article. Let us denote with a tilde all the quantities related to the subdomains $\Omega_{\tilde{i}}$ obtained by extending by one (or more) layers of cells subdomains Ω_i , see fig. 5. Similarly to [2], we define

$$\mathcal{N}_{\tilde{i}} := \{k \in \mathcal{N} \mid \text{meas}(\text{Supp}(\phi_k) \cap \Omega_{\tilde{i}}) > 0\}.$$

Since $\Omega_i \subset \Omega_{\tilde{i}}$, we have

$$\mathcal{N}_i \subset \mathcal{N}_{\tilde{i}}.$$

Also from the partition of unity on the original decomposition, we can define a partition of unity on $(\mathcal{N}_{\tilde{i}})_{1 \leq i \leq N}$ inherited from the one on $(\mathcal{N}_i)_{1 \leq i \leq N}$ by defining diagonal matrices $(D_{\tilde{i}})_{1 \leq i \leq N}$ in the following manner:

$$(D_{\tilde{i}})_{kk} := \begin{cases} (D_i)_{kk} & \text{if } k \in \mathcal{N}_i \\ 0 & \text{if } k \in \mathcal{N}_{\tilde{i}} \setminus \mathcal{N}_i \end{cases}.$$




Figure 1: A subdomain and its extension by one layer

We have clearly a partition of unity:

$$I = \sum_{i=1}^N R_i^T D_i R_i.$$

Also since for all subdomains $1 \leq i \leq N$, the entries of D_i are zero on the added layers, we have the following equality:

$$D_i R_i = R_i R_i^T D_i R_i.$$

The coarse space is built by first introducing the Neumann matrices A_i^{Neu} on subdomains Ω_i for $1 \leq i \leq N$, as in [19], so that we have:

$$\sum_{i=1}^N (A_i^{Neu} R_i \mathbf{U}, R_i \mathbf{U}) \leq \tilde{k}_1 (A \mathbf{U}, \mathbf{U}),$$

where \tilde{k}_1 is the maximum multiplicity of the intersections of subdomains Ω_i . Let V_{ik} be the eigenvectors of the following generalized eigenvalue problem:

$$D_i R_i A R_i^T D_i V_{ik} = \lambda_{ik} A_i^{Neu} V_{ik}.$$

Note that for λ_{ik} not equal to 1, the eigenvectors are harmonic for the interior degrees of freedom since for these points the left and right matrices have identical entries for the corresponding lines. Thus, for $\lambda_{ik} \neq 1$, we might as well zero the lines corresponding the interior degrees of freedom and keep only the entries of the degrees of freedom in the overlap. This GEVP is thus also of the type GenEO.

For a user-defined parameter τ , let us define the coarse space as follows:

$$V_0 := \text{Span}\{R_i^T D_i V_{ik} \mid 1 \leq i \leq N, \lambda_{ik} > \tau\},$$

and a rectangular matrix $Z \in \mathbb{R}^{\#\mathcal{N} \times \#\mathcal{N}_0}$ whose columns are a basis of V_0 where \mathcal{N}_0 is a set of indices whose cardinal is the dimension of the vector space V_0 . We also define local projections $(\pi_i)_{1 \leq i \leq N}$ on $\text{Span}\{V_{ik} \mid \lambda_{ik} > \tau\}$ parallel to $\text{Span}\{V_{ik} \mid \lambda_{ik} \leq \tau\}$.

We have then a stable decomposition. Indeed, let $\mathbf{U} \in \mathbb{R}^{\#\mathcal{N}}$,

$$\mathbf{U} = \sum_{i=1}^N R_i^T D_i R_i \mathbf{U} = \sum_{i=1}^N R_i^T R_i R_i^T D_i (R_i \mathbf{U} - \pi_i R_i \mathbf{U}) + \sum_{i=1}^N R_i^T R_i R_i^T D_i \pi_i R_i \mathbf{U}.$$

The last term is clearly in V_0 so that there exists $\mathbf{U}_0 \in \mathbb{R}^{\mathcal{N}_0}$ such that

$$Z\mathbf{U}_0 = \sum_{i=1}^N R_i^T R_i R_i^T D_i \pi_i R_i \mathbf{U} = \sum_{i=1}^N R_i^T D_i \pi_i R_i \mathbf{U} \in V_0.$$

Let us define

$$\mathbf{U}_i := R_i R_i^T D_i (R_i \mathbf{U} - \pi_i R_i \mathbf{U}).$$

This decomposition is stable since

$$\begin{aligned} \sum_{i=1}^N (A R_i^T \mathbf{U}_i, R_i^T \mathbf{U}_i) &= \sum_{i=1}^N (A R_i^T R_i R_i^T D_i (R_i \mathbf{U} - \pi_i R_i \mathbf{U}), R_i^T R_i R_i^T D_i (R_i \mathbf{U} - \pi_i R_i \mathbf{U})) \\ &= \sum_{i=1}^N (A R_i^T D_i (R_i \mathbf{U} - \pi_i R_i \mathbf{U}), R_i^T D_i (R_i \mathbf{U} - \pi_i R_i \mathbf{U})) \\ &\leq \tau \sum_{i=1}^N (A_i^{Neu} R_i \mathbf{U}, R_i \mathbf{U}) \leq \tau \tilde{k}_1 (A \mathbf{U}, \mathbf{U}). \end{aligned}$$

Note also that Assumption 2.1 of [19] is automatically satisfied in the finite element framework chosen here whereas Assumptions 3.12 and 3.13 are not needed here since our construction is simpler.

References

- [1] Markus Blatt, Ansgar Burchardt, Andreas Dedner, Christian Engwer, Jorrit Fahlke, Bernd Flemisch, Christoph Gersbacher, Carsten Gräser, Felix Gruber, Christoph Grüninger, et al. The distributed and unified numerics environment, version 2.4. *Archive of Numerical Software*, 4(100):13–29, 2016.
- [2] Victorita Dolean, Pierre Jolivet, and Frédéric Nataf. *An Introduction to Domain Decomposition Methods: algorithms, theory and parallel implementation*. SIAM, 2015.
- [3] Maksymilian Dryja, Marcus V. Sarkis, and Olof B. Widlund. Multilevel Schwarz methods for elliptic problems with discontinuous coefficients in three dimensions. *Numer. Math.*, 72(3):313–348, 1996.
- [4] Yalchin Efendiev, Juan Galvis, Raytcho Lazarov, and Joerg Willems. Robust domain decomposition preconditioners for abstract symmetric positive definite bilinear forms. *ESAIM Math. Model. Numer. Anal.*, 46(5):1175–1199, 2012.
- [5] R. M. Gower, D. Goldfarb, and P. Richtàrik. Stochastic block bfgs: Squeezing more curvature out of data. Technical report, Arxiv, 2016.
- [6] M. Griebel and P. Oswald. On the abstract theory of additive and multiplicative Schwarz algorithms. *Numer. Math.*, 70(2):163–180, 1995.
- [7] R. Haferssas, P. Jolivet, and F. Nataf. An Additive Schwarz Method Type Theory for Lions’s Algorithm and a Symmetrized Optimized Restricted Additive Schwarz Method. *SIAM J. Sci. Comput.*, 39(4):A1345–A1365, 2017.

- [8] Ryadh Haferssas, Pierre Jolivet, and Frédéric Nataf. A robust coarse space for optimized Schwarz methods: SORAS-GenEO-2. *C. R. Math. Acad. Sci. Paris*, 353(10):959–963, 2015.
- [9] F. Hecht. New development in Freefem++. *J. Numer. Math.*, 20(3-4):251–265, 2012.
- [10] Pierre Jolivet and Frédéric Nataf. Hpddm: High-Performance Unified framework for Domain Decomposition methods, MPI-C++ library. <https://github.com/hpddm/hpddm>, 2014.
- [11] Pierre-Louis Lions. On the Schwarz alternating method. III: a variant for nonoverlapping subdomains. In Tony F. Chan, Roland Glowinski, Jacques Périaux, and Olof Widlund, editors, *Third International Symposium on Domain Decomposition Methods for Partial Differential Equations, held in Houston, Texas, March 20-22, 1989*, Philadelphia, PA, 1990. SIAM.
- [12] Jan Mandel. Balancing domain decomposition. *Comm. on Applied Numerical Methods*, 9:233–241, 1992.
- [13] Jan Mandel, Bedřich Sousedík, and Clark R. Dohrmann. *On Multilevel BDDC*, pages 287–294. Springer Berlin Heidelberg, Berlin, Heidelberg, 2008.
- [14] Sergey V. Nepomnyaschikh. Mesh theorems of traces, normalizations of function traces and their inversions. *Sov. J. Numer. Anal. Math. Modeling*, 6:1–25, 1991.
- [15] Roy A. Nicolaides. Deflation of conjugate gradients with applications to boundary value problems. *SIAM J. Numer. Anal.*, 24(2):355–365, 1987.
- [16] Clemens Pechstein and Clark R. Dohrmann. A unified framework for adaptive BDDC. *Electron. Trans. Numer. Anal.*, 46:273–336, 2017.
- [17] C. Prud’homme. A Domain Specific Embedded Language in c++ for automatic differentiation, projection, integration and variational formulations. *Scientific Programming*, 14(2):81–110, 2006.
- [18] R. B. Schnabel. Quasi-Newton Methods using Multiple Secant Equations. Tech. rep. Computer Science Technical Reports 247-83, Paper 244, Colorado University-Computer Science, 1983.
- [19] Nicole Spillane, Victorita Dolean, Patrice Hauret, Frédéric Nataf, Clemens Pechstein, and Robert Scheichl. Abstract robust coarse spaces for systems of PDEs via generalized eigenproblems in the overlaps. *Numer. Math.*, 126(4):741–770, 2014.
- [20] Amik St-Cyr, Martin J. Gander, and Stephen J. Thomas. Optimized Multiplicative, Additive, and Restricted Additive Schwarz Preconditioning. *SIAM J. Sci. Comput.*, 29(6):2402–2425 (electronic), 2007.
- [21] J.M. Tang, R. Nabben, C. Vuik, and Y.A. Erlangga. Comparison of two-level preconditioners derived from deflation, domain decomposition and multigrid methods. *Journal of Scientific Computing*, 39(3):340–370, 2009.

- [22] Andrea Toselli and Olof Widlund. *Domain Decomposition Methods - Algorithms and Theory*, volume 34 of *Springer Series in Computational Mathematics*. Springer, 2005.
- [23] Xuemin Tu. Three-level BDDC in three dimensions. *SIAM J. Sci. Comput.*, 29(4):1759–1780, 2007.
- [24] Xuemin Tu. A three-level BDDC algorithm for a saddle point problem. *Numer. Math.*, 119(1):189–217, 2011.
- [25] Xuejun Zhang. Multilevel Schwarz methods. *Numer. Math.*, 63(4):521–539, 1992.