

On Smooth Orthogonal and Octilinear Drawings: Relations, Complexity and Kandinsky Drawings^{*}

Michael A. Bekos, Henry Förster, Michael Kaufmann

Wilhelm-Schickhard-Institut für Informatik, Universität Tübingen, Germany

Abstract. We study two variants of the well-known orthogonal drawing model: (i) the smooth orthogonal, and (ii) the octilinear. Both models form an extension of the orthogonal, by supporting one additional type of edge segments (circular arcs and diagonal segments, respectively).

For planar graphs of max-degree 4, we analyze relationships between the graph classes that can be drawn bendless in the two models and we also prove NP-hardness for a restricted version of the bendless drawing problem for both models. For planar graphs of higher degree, we present an algorithm that produces bi-monotone smooth orthogonal drawings with at most two segments per edge, which also guarantees a linear number of edges with exactly one segment.

1 Introduction

Orthogonal graph drawing is an intensively studied and well established model for drawing graphs. As a result, several efficient algorithms providing good aesthetics and good readability have been proposed over the years, see e.g., [7,17,28,34]. In such drawings, each vertex corresponds to a point on the Euclidean plane and each edge is drawn as a sequence of axis-aligned line segments; see Fig. 1.

Several research directions build upon this successful model. We focus on two models that have recently received attention: (i) the *smooth orthogonal* [4], in which every edge is a sequence of axis-aligned segments and circular arc segments with common axis-aligned tangents (i.e., quarter, half or three-quarter arc segments), and (ii) the *octilinear* [2], in which every edge is a sequence of axis-aligned and diagonal (at $\pm 45^\circ$) segments.

Observe that both models extend the orthogonal by allowing one more type of edge-segments. The former was introduced with the aim of combining the artistic appeal of *Lombardi drawings* [12,14] with the clarity of the orthogonal drawings. The latter, on the other hand, is primarily motivated by metro-map and map schematization applications (see, e.g., [24,30,31,33]). Note that in the orthogonal and in the smooth orthogonal models, each edge may enter a vertex using one out of four available (axis-aligned) directions, called *ports*. Thus both models support graphs of max-degree 4. In the octilinear model, each vertex has eight available ports and therefore one can draw graphs of max-degree 8.

^{*} This work is supported by DFG grant Ka812/17-1.




Fig. 1: Different drawings of a planar graph of max-degree 4: (a) straight-line, (b) orthogonal 3-drawing, (c) octilinear 2-drawing, and (d) smooth orthogonal 2-drawing.

For readability purposes, usually in such drawings one seeks to minimize the *edge complexity* [10,26], i.e., the maximum number of segments used for representing any edge. Also, when the input is a planar graph, one seeks for a corresponding planar drawing. Note that drawings with edge complexity 1 are also called *bendless*. We refer to drawings with edge complexity k as k -drawings; thus, by definition, orthogonal k -drawings have at most $k - 1$ bends per edge.

Known results. There exists a plethora of results for each of the aforementioned models; here we list existing results for drawings with low edge complexity.

- All planar graphs of max-degree 4, except for the octahedron, admit orthogonal 3-drawings; the octahedron is orthogonal 4-drawable [7,28]. Minimizing the number of bends over all embeddings of a planar graph of max-degree 4 is \mathcal{NP} -hard [21]. For a given planar embedding, however, finding a planar orthogonal drawing with minimum number of bends can be done in polynomial time by an approach, called *topology-shape-metrics* [34], that is based on min-cost flow computations and works in three phases. Initially, a planar embedding is computed if not specified by the input. In the next phase, the angles and the bends of the drawing are computed, yielding an *orthogonal representation*. In the last phase, the actual coordinates for the vertices and bends are computed.
- All planar graphs of max-degree 4 (including the octahedron) admit smooth orthogonal 2-drawings. Note that not all planar graphs of max-degree 4 allow for bendless smooth orthogonal drawings [4], and that such drawings may require exponential area [1]. Bendless smooth orthogonal drawings are possible only for subclasses, e.g., for planar graphs of max-degree 3 [3] and for outerplanar graphs of max-degree 4 [1]. It is worth mentioning that the complexity of the problem, whether a planar graph of max-degree 4 admits a bendless smooth orthogonal drawing, has not been settled (it is conjectured to be \mathcal{NP} -hard [1]).
- All planar graphs of max-degree 8 admit octilinear 3-drawings [27], while planar graphs of max-degree 4 or 5 allow for octilinear 2-drawings [2]. Bendless octilinear drawings are always possible for planar graphs of max-degree 3 [22]. Note that deciding whether an embedded planar graph of max-degree 8 admits a bendless octilinear drawing is \mathcal{NP} -hard [30]. It is not, however, known whether this negative result applies for planar graphs of max-degree 4 or whether these graphs allow for a decision algorithm (in fact, there exist planar graphs of max-degree 4 that do not admit bendless octilinear drawings [5]).




Fig. 2: Different inclusion-relationships: For $k \geq 1$, SC_k and $8C_k$ correspond to the classes of graphs admitting smooth orthogonal and octilinear k -drawings, respectively.

Our contribution Motivated by the fact that usually one can “easily” convert an octilinear drawing of a planar graph of max-degree 4 to a corresponding smooth orthogonal one (e.g., by replacing diagonal edge segments with quarter circular arc segments; see Figs. 1c-1d for an example), and vice versa, we study in Section 2 inclusion-relationships between the graph-classes that admit such drawings. In Section 3, we show that it is \mathcal{NP} -hard to decide whether an embedded planar graph of max-degree 4 admits a bendless smooth orthogonal or a bendless octilinear drawing, in the case where the angles between any two edges incident to a common vertex and the shapes of all edges are specified as part of the input (e.g., as in the last step of the topology-shape-metrics approach [34]). Our proof is a step towards settling the complexities of both decision problems in their general form. Inspired from the *Kandinsky model* (see, e.g., [6,9,17]) for drawing planar graphs of arbitrary degree in an orthogonal style, we present in Section 4 two drawing algorithms that yield bi-monotone smooth orthogonal drawings of good quality. The first yields drawings of smaller area, which can also be transformed to octilinear with bends at 135° . The second yields larger drawings but guarantees that at most $2n - 5$ edges are drawn with two segments. We conclude in Section 5 with open problems.

Preliminaries For graph theoretic notions refer to [23]. For definitions on planar graphs, we point the reader to [10,26]. We also assume familiarity with standard graph drawing techniques, such as the *canonical ordering* [18,25] and the *shift-method* by de Fraysseix, Pach and Pollack [18]; see App. A for more details.

2 Relationships between Graph Classes

In this section, we consider relationships between the classes of graphs that admit smooth orthogonal k -drawings and octilinear k -drawings, $k \geq 1$, denoted as SC_k and $8C_k$, respectively. Our findings are also summarized in Fig. 2.

By definition, $SC_1 \subseteq SC_2$ and $8C_1 \subseteq 8C_2 \subseteq 8C_3$ hold. Since each planar graph of max-degree 8 admits an octilinear 3-drawing [27], class $8C_3$ coincides with the class of planar graphs of max-degree 8. Similarly, class SC_2 coincides with the class of planar graph of max-degree 4, as these graphs admit




Fig. 3: Illustrations for the proof of Theorem 1.

smooth orthogonal 2-drawings [1]. This also implies that $SC_2 \subseteq 8C_2$, since each planar graph of max-degree 4 admits an octilinear 2-drawing [2]. The relationship $8C_2 \neq 8C_3$ follows from [2], where it was proven that there exist planar graphs of max-degree 6 that do not admit octilinear 2-drawings. The relationship $SC_2 \neq 8C_2$ follows from [5], where it was shown that there exist planar graphs of max-degree 5 that admit octilinear 2-drawings and no octilinear 1-drawings, and the fact that planar graphs of max-degree 5 cannot be drawn in the smooth orthogonal model. The octahedron graph admits neither a bendless smooth orthogonal drawing [4] nor a bendless octilinear drawing [5]. However, since it is of max-degree 4, it admits 2-drawings in both models [1,2]. Hence, it belongs to $8C_2 \cap SC_2 \setminus (8C_1 \cup SC_1)$. To prove that $8C_1 \setminus SC_2 \neq \emptyset$, observe that a caterpillar whose spine vertices are of degree 8 clearly admits an octilinear 1-drawing, however, due to its degree it does not admit a smooth orthogonal.

To complete the discussion of the relationships of Fig. 2, we have to show that SC_1 and $8C_1$ are incomparable. This is the most interesting part of our proof, as usually one can “easily” convert a bendless octilinear drawing of a planar graph of max-degree 4 to a corresponding bendless smooth orthogonal one (e.g., by replacing diagonal segments with quarter circular arcs), and vice versa; see, e.g., Figs. 1c-1d. Since the endpoints of each edge of a bendless smooth orthogonal or octilinear drawing are along a line with slope 0, 1, -1 or ∞ , such conversions are in principle possible. Two difficulties that might arise are to preserve planarity and to guarantee that no two edges enter a vertex using the same port. Clearly, however, there exist infinitely many (even 4-regular) planar graphs that admit both drawings in both models; see Fig. 3 and App. B for more details.

Theorem 1. *There is an infinitely large family of 4-regular planar graphs that admit both bendless smooth orthogonal and bendless octilinear drawings.*

In the next two theorems we show that SC_1 and $8C_1$ are incomparable.

Theorem 2. *There is an infinitely large family of 4-regular planar graphs that admit bendless smooth orthogonal drawings but no bendless octilinear drawings.*

Proof. Consider the planar graph C of Fig. 4a, which is drawn bendless smooth orthogonal. We claim that C admits no bendless octilinear drawing. If one substitutes its degree-2 vertex (denoted by c in Fig. 4a) by an edge connecting its




Fig. 4: Illustrations for the proof of Theorem 2.

two neighbors, then the resulting graph is triconnected, which admits an unique embedding (up to the choice of its outerface; see Figs. 4a-4b). Now, observe that the outerface of any octilinear drawing of graph C (if any) has length at most 5 (Constraint 1). In addition, each vertex of this outerface (except for c , which is of degree 2) must have two ports pointing in the interior of this drawing, because every vertex of C is of degree 4 except for c . This implies that the angle formed by any two consecutive edges of this outerface is at most 225° , except for the pair of edges incident to c (Constraint 2). But if we want to satisfy both constraints, then at least one edge of this outerface must be drawn with a bend; see Fig. 4c. Hence, graph C does not admit a bendless octilinear drawing.

Based on graph C , for each $k \in \mathbb{N}_0$ we construct a 4-regular planar graph G_k consisting of $k + 2$ biconnected components C_1, \dots, C_{k+2} arranged in a chain; see Fig. 4d for the case $k = 1$. Clearly, G_k admits a bendless smooth orthogonal drawing for any k . Since the end-components of the chain (i.e., C_1 and C_{k+2}) are isomorphic to C , G_k does not admit a bendless octilinear drawing for any k . \square

Theorem 3. *There is an infinitely large family of 4-regular planar graphs that admit bendless octilinear drawings but no bendless smooth orthogonal drawings.*

Proof (sketch). Consider the planar graph B of Fig. 5a, which is drawn bendless octilinear. Graph B has two separation pairs (i.e., $\{t_1, t_2\}$ and $\{p_1, p_2\}$ in Fig. 5a).




Fig. 5: Illustrations for the proof of Theorem 3.

Based on graph B , for each $k \in \mathbb{N}_0$ we construct a 4-regular planar graph G_k consisting of $2k + 4$ copies of B arranged in a cycle; see Fig. 5b where each copy of B is drawn as a gray-shaded parallelogram. By construction, G_k admits a bendless octilinear drawing for any k . By planarity at least one copy of graph B must be embedded with the outerface of Fig. 5a. However, if we require the outerface of B to be the one of Fig. 5a, then all possible planar embeddings of B are isomorphic to the one of Fig. 5a. We exploit this property in App. B to show that B does not admit a bendless smooth orthogonal drawing with this outerface. The detailed proof is based on an exhaustive consideration of all bendless smooth orthogonal drawings of subgraphs of B , which we incrementally augment by adding more vertices to them. Thus, for any k , graph G_k does not admit a bendless smooth orthogonal drawing. \square

3 \mathcal{NP} -hardness Results

In this section, we study the complexity of the bendless smooth orthogonal and octilinear drawing problems. As a first step towards addressing the complexity of both problems for planar graphs of max-degree 4 in general, here we make an additional assumption. We assume that the input, apart from an embedding, also specifies a *smooth orthogonal* or an *octilinear representation*, which are defined analogously to the orthogonal ones: (i) the angles between consecutive edges incident to a common vertex in the cyclic order around it (given by the planar embedding) are specified, and (ii) the *shape* of each edge (e.g., straight-line, or quarter-circular arc) is also specified. In other words, we assume that our input is analogous to the one of the last step of the topology-shape-metrics approach [34].

Theorem 4. *Given a planar graph G of max-degree 4 and a smooth orthogonal representation \mathcal{R} , it is \mathcal{NP} -hard to decide whether G admits a bendless smooth orthogonal drawing preserving \mathcal{R} .*

Proof. Our reduction is from the well-known 3-SAT problem [20]. Given a formula φ in conjunctive normal form, we construct a graph G_φ and a smooth orthogonal representation \mathcal{R}_φ , such that G_φ admits a bendless smooth orthogonal drawing Γ_φ preserving \mathcal{R}_φ if and only if φ is satisfiable; see also Fig. 6.

The main ideas of our construction are: (i) specific straight-line edges in Γ_φ transport *information* encoded in their length, (ii) rectangular faces of Γ_φ propagate the edge length of one side to its opposite, and (iii) for a face composed of two straight-line edges and a quarter circle arc, the straight-line edges are of same length, which allows us to change the *direction* in which the information “flows”.

Variable gadget. For each variable x of φ , we introduce a gadget; see Figs. 7a-7b. The bold-drawn quarter circle arc ensures that the sum of the edge lengths to its left is the same as the sum of the edge lengths to its bottom (refer to the edges with gray endvertices). As “input” the gadget gets three edges of unit length $\ell(u)$. This ensures that $\ell(x) + \ell(\bar{x}) = 3 \cdot \ell(u)$ holds for the “output literals” x and \bar{x} , where $\ell(x)$ and $\ell(\bar{x})$ denote the lengths of two edges representing x and \bar{x} .




Fig. 6: Drawing Γ_φ for $\varphi = (a \vee b \vee c) \wedge (\bar{a} \vee \bar{b} \vee c)$ and the assignment $a = \text{false}$ and $b = c = \text{true}$.




Fig. 7: The (a)-(b) variable, and (c)-(d) the parity gadgets; gray-colored arrows show the information “flow”.

To introduce our concept, assume that the lengths of all straight-line edges are integral and at least 1. If we could require $\ell(u) = 1$, then $\ell(x), \ell(\bar{x}) \in \{1, 2\}$. This would allow us to encode the assignment $x = \text{true}$ with $\ell(x) = 2$ and $\ell(\bar{x}) = 1$, and the assignment $x = \text{false}$ with $\ell(x) = 1$ and $\ell(\bar{x}) = 2$. However, if we cannot avoid, e.g., that $\ell(u) = 2$, then the variable gadget would not prevent us from setting $\ell(x) = \ell(\bar{x}) = 3$, which means that x and \bar{x} are “half-true”. We solve this issue by the so-called *parity gadget*, that allows us to relax the integral constraint and to ensure that $\ell(x), \ell(\bar{x}) \in \{\ell(u) + \varepsilon, 2\ell(u) - \varepsilon\}$, for $0 < \varepsilon < \ell(u)$.

Parity gadget. For each variable x of φ , G_φ has a gadget (see Figs. 7c-7d), which results in overlaps in Γ_φ , if the values of $\ell(x)$ and $\ell(\bar{x})$ do not differ significantly. The central part of this gadget is a “vertical gap” of width $3 \cdot \ell(u)$ (shaded in gray in Figs. 7c-7d) with two blocks of vertices (triangular- and square-shaped in Figs. 7c-7d) pointing inside the gap. Each block defines two square-shaped faces and three faces of length 3, each formed by two straight-line edges and a quarter circle arc. Depending on the choice of $\ell(x)$ and $\ell(\bar{x})$, one of the blocks may be located above the other. If $\ell(x) \approx \ell(\bar{x})$, however, we can observe that the two blocks are not far enough apart from each other, which leads to overlaps. Using elementary geometry, we prove in App. C that overlaps can be avoided if and only if $|\ell(\bar{x}) - \ell(x)| > \sqrt{3}/2 \cdot \ell(u) \approx 0.866 \cdot \ell(u)$, which implies: that $\ell(x), \ell(\bar{x}) \in (0, 1.067 \cdot \ell(u)] \cup [1.933 \cdot \ell(u), 3)$, i.e., $\varepsilon < 0.067 \cdot \ell(u)$.

Clause gadget. For each clause of φ with literals a, b and c , we introduce a gadget, which is illustrated in Fig. 8a. The bold-drawn quarter circle arc of Fig. 8a compares two sums of information. From the righthand side, four edges of unit




Fig. 8: Different gadgets; gray-colored arrows show the information “flow”.

length “enter” the arc. Observe that there is also a *free edge* (marked with an asterisk in Fig. 8a), which also contributes to the sum but can be stretched independently of any other edge. Hence, the sum of edge lengths on the righthand side of this arc is $> 4 \cdot \ell(u)$. The three literals “enter” at the bottom; the sum here is $\ell(a) + \ell(b) + \ell(c)$. Combining both, we obtain that $\ell(a) + \ell(b) + \ell(c) > 4 \cdot \ell(u)$ must hold. This implies that not all a , b and c can be **false**, since in this case $\ell(a) + \ell(b) + \ell(c) = 3 \cdot (\ell(u) + \varepsilon) < 4 \cdot \ell(u)$.

Auxiliary gadgets. The *crossing gadget* just consists of a rectangle and is used to allow two flows of information to cross each other; see Fig. 8b. The *copy gadget* takes an information and creates three copies of this information; see Fig. 8c. This is because both quarter circular arcs of the copy gadget must have the same radius in the presence of the half circular arc of the copy gadget. Finally, the *unit length gadget* is a single edge, which we assume to be of length $\ell(u)$.

We now describe our construction; see Fig. 6: G_φ contains one unit length gadget, which is copied several times using the copy gadget (the number of copies depends linearly on the number of variables ν and clauses μ of φ). For each variable of φ , G_φ has a variable gadget and a parity gadget, each of which is connected to different copies of the unit length gadget. For each clause of φ , G_φ has a clause gadget, which has four connections to different copies of the unit length gadget. We compute \mathcal{R}_φ as follows. We place the variable gadget of each variable x above and to the left of its parity gadget and we connect the output literals of the variable gadget of x with its parity gadget through a copy gadget. We place the variable and the parity gadgets of the i -th variable below and to the right of the corresponding ones of the $(i - 1)$ -th variable. We place each clause gadget to the right of the sketch constructed so far, so that the gadget of the i -th clause is to the right of the $(i - 1)$ -th clause. This allows us to connect copies of the output literals of the variable gadget of each variable with the clause gadgets that contain it, so that all possible crossings (which are resolved using the crossing gadget) appear above the clause gadgets. More precisely, if a clause contains a literal of the i -th variable, we have a crossing with the literals of all variables with indices $(i + 1)$ to ν . Hence, for each clause we add $O(\nu)$ crossing and three copy gadgets. Note that all copy gadgets of the unit length gadget lie below all variable, parity, and clause gadgets. The obtained representation \mathcal{R}_φ conforms with the one of Fig. 6. The construction can be done in $O(\nu\mu)$ time.

To complete the proof, assume that G_φ admits a bendless smooth orthogonal drawing Γ_φ preserving \mathcal{R}_φ . For each variable x of φ , we set x to **true** if and only if $\ell(x) \geq 1.933 \cdot \ell(u)$. Since for each clause $(a \vee b \vee c)$ of φ we have that $\ell(a) + \ell(b) + \ell(c) > 4 \cdot \ell(u)$, at least one of a, b and c must be **true**. Hence, φ admits a truth assignment. For the opposite direction, based on a truth assignment of φ , we can set, e.g., $\ell(x) = 1.95$ and $\ell(\bar{x}) = 1.05$ for each variable x , assuming that $\ell(u) = 1$. Then, arranging the variable and the clause gadgets of G_φ as in Fig. 6 yields a bendless smooth orthogonal drawing Γ_φ preserving \mathcal{R}_φ . \square

Remark 1. The special case of our problem, in which circular arcs are not present, is known as *HV-rectilinear planarity testing* [29]. As opposed to our problem, HV-rectilinear planarity testing is polynomial-time solvable in the fixed embedding setting [13] (and becomes \mathcal{NP} -hard in the variable embedding setting [11]).

Theorem 5. *Given a planar graph G of max-degree 4 and an octilinear representation \mathcal{R} , it is \mathcal{NP} -hard to decide whether G admits a bendless octilinear drawing preserving \mathcal{R} .*

Proof (sketch). Except for the parity gadget, we can adjust to the octilinear model simply by replacing arcs with diagonal segments; for details see App. A. In this case the parity gadget guarantees $|\ell(x) - \ell(\bar{x})| > 5/6 \cdot \ell(u) \approx 0.833 \cdot \ell(u)$, which implies that $\varepsilon < 0.084 \cdot \ell(u)$. \square

4 Bi-Monotone Drawings

In this section, we study variants of the *Kandinsky* drawing model [6,9,17], which forms an extension of the orthogonal model to graphs of degree greater than 4. In this model, the vertices are represented as squares, placed on a *coarse grid*, with multiple edges attached to each side of them aligned on a *finer grid*.

The Kandinsky model allows for natural extensions to both smooth orthogonal and octilinear models. We are aware of only one preliminary result in this direction: A linear time drawing algorithm is presented in [4] for the production of smooth orthogonal 2-drawings for planar graphs of arbitrary degree in quadratic area, in which all vertices are on a line ℓ and the edges are drawn either as half circles (above or below ℓ), or as two consecutive half circles one above and one below ℓ (i.e., the latter ones are of complexity 2, but they are at most $n - 2$).

For an input maximal planar graph G (of arbitrary degree), our goal is to construct a smooth orthogonal (or an octilinear) 2-drawing for G with the following aesthetic benefits over the aforementioned drawing algorithm: (i) the vertices are distributed evenly over the drawing area, and (ii) each edge is *bi-monotone* [19], i.e., *xy-monotone*. We achieve our goal at the cost of slightly more edges drawn with complexity 2 or at the cost of increased drawing area (but still polynomial).

Our first approach is a modification of the *shift-method* [18]. Based on a canonical order $\pi = (v_1, \dots, v_n)$ of G , we construct a planar smooth orthogonal 2-drawing Γ of G in the Kandinsky model, as follows. We place v_1, v_2 and v_3 at $(0, 0)$, $(2, 0)$ and $(1, 1)$. Hence, we can draw (v_1, v_2) as a horizontal segment, and




Fig. 9: Illustration of the contour condition (left) and the placement of v_k (right).

each of (v_1, v_3) and (v_2, v_3) as a quarter circular arc. We also color (v_1, v_3) blue and (v_2, v_3) green. For $k = 4, \dots, n$, assume that a smooth orthogonal 2-drawing Γ_{k-1} of the subgraph G_{k-1} of G induced by v_1, \dots, v_{k-1} has been constructed, in which each edge of the outerface C_{k-1} of Γ_{k-1} is drawn as a quarter circular arc, whose endvertices are on a line with slope ± 1 , except for edge (v_1, v_2) , which is drawn as a horizontal segment (called *contour condition* in the shift-method; see Fig. 9). Each of v_1, \dots, v_{k-1} is also associated with a so-called *shift-set*, which for v_1, v_2 and v_3 are singletons containing only themselves.

Let w_1, \dots, w_p be the vertices of C_{k-1} from left to right in Γ_{k-1} , where $w_1 = v_1$ and $w_p = v_2$. Let (w_ℓ, \dots, w_r) , $1 \leq \ell < r \leq p$, be the neighbors of v_k from left to right along C_{k-1} in Γ_{k-1} . As in the shift-method, our algorithm first translates each vertex in $\cup_{i=1}^\ell S(w_i)$ one unit to the left and each vertex in $\cup_{i=r}^p S(w_i)$ one unit to the right, where $S(v)$ is the shift-set of $v \in V$. During this translation, $(w_\ell, w_{\ell+1})$ and (w_{r-1}, w_r) acquire a horizontal segment each (see the bold edges of Fig. 9). We place v_k at the intersection of line L_ℓ with slope $+1$ through w_ℓ with line L_r with slope -1 through w_r (dotted in Fig. 9) and we set the shift-set of v_k to $\{v_k\} \cup \cup_{i=\ell+1}^{r-1} S(w_i)$, as in the shift-method. We draw each of (w_ℓ, v_k) and (v_k, w_r) as a quarter circular arc. For $i = \ell + 1, \dots, r - 1$, (w_i, v_k) has a vertical line-segment that starts from w_i and ends either at L_ℓ or L_r and a quarter circle arc from the end of the previous segment to v_k . Hence, the contour condition is satisfied. We color (w_ℓ, v_k) blue, (v_k, w_r) green and the remaining edges of v_k red; see also [15,32]. Observe that each blue and green edge consists of a quarter circular arc and a horizontal segment (that may have zero length), while a red edge consists of a vertical segment and a quarter circular arc (that may have zero radius). We are now ready to state our first theorem; the analogous of Theorem 6 for the octilinear model is shown in App. D.

Theorem 6. *A maximal planar n -vertex graph admits a bi-monotone planar smooth orthogonal 2-drawing in the Kandinsky model, which requires $O(n^2)$ area and can be computed in $O(n)$ time.*

Proof. Bi-monotonicity follows by construction. The time complexity follows from [8]. Planarity is proven by induction. Drawing Γ_3 is planar by construction. Assuming that Γ_{k-1} is planar, we observe that no two edges incident to v_k cross in Γ_k . Also, these edges do not cross edges of Γ_{k-1} . Since the radii of the arcs of the edges incident to vertices that are shifted remain unchanged and since edges incident to vertices in the shift-sets retain their shape, drawing Γ_k is planar. \square

We reduce the number of edges drawn with complexity 2 in two steps. (S.1) We stretch the drawing horizontally (by employing appropriate vertical cuts; refer to App. A) to eliminate the vertical segments of all red edges with a circular arc segment of non-zero radius. (S.2) We stretch the drawing vertically, to guarantee that the edges of a spanning tree (i.e., $n - 1$) are drawn with complexity 1.

For Step S.1, we assume that each blue and green edge has a horizontal segment (that may be of zero length). Consider a red edge (u, v) with a vertical segment of length δ and assume w.l.o.g. that u is to the right and above v ; see Fig. 18. If we shift u by δ units to the right, then (u, v) can be drawn as a quarter circular arc. If the shift is by more than δ units, then a horizontal segment is needed. Since all edges incident to u that are drawn below u enter u from its left or from its right side, the shift of u cannot introduce crossings between them.

We eliminate the vertical segments of all red edges with a circular arc segment of non-zero radius, as follows. As long as there exist such edges, we choose the one, call it (u, v) , whose vertical segment has the largest length δ , and assume that u is to the right and above v . We eliminate the vertical segment of (u, v) using a vertical cut L at $x(u) - \varepsilon$, for small $\varepsilon > 0$. Since L crosses several edges, shifting all vertices to the right of L by δ to the right has the following effects. By the choice of (u, v) , the vertical segments of all red edges crossed by L are eliminated; note that this may introduce new horizontal segments. The horizontal segment of each blue and green edge crossed by L is elongated by δ . Both imply that no edge crossings are introduced. Hence, by the termination of our algorithm all edges with vertical segments are of complexity 1.

Step 1 ensures that the x -distance of adjacent vertices is at least as large as their y -distance (unless they are connected by vertical edges). Based on this property, in Step 2 we compute new y -coordinates for the vertices in the sequence of the canonical ordering π , keeping their x -coordinates unchanged. First, we set $y(v_1) = y(v_2) = 0$. For each $k = 3, \dots, n$, we set $y(v_k) = \max_{w \in \{w_\ell, \dots, w_r\}} \{y(w) + \max\{\Delta_x(v_k, w), 1\}\}$, where w_ℓ, \dots, w_r are the neighbors of v_k in Γ_{k-1} , i.e., v_k is placed above w_ℓ, \dots, w_r in Γ_{k-1} , such that one of its edges (the one of the maximum; call it (v_k, w^*)) is drawn with complexity 1; as a quarter circle arc or as a vertical edge depending on whether the x -distance of v_k and w^* is non-zero or not. Since (v_k, w^*) is the edge that must be stretched the most in order to ensure that it is drawn with complexity 1, for all other edges incident to v_k in G_k , the y -distance of their endpoints is at least as large as their corresponding x -distance. Hence, they are drawn as vertical segments followed by quarter circular arcs (that may have zero radius). We are now ready to state our second theorem.

Theorem 7. *A maximal planar n -vertex graph admits a bi-monotone planar smooth orthogonal 2-drawing with at least $n - 1$ edges with complexity 1 in the Kandinsky model, which requires $O(n^4)$ area and can be computed in $O(n^2)$ time.*

Proof (sketch). For $k = 3, \dots, n$, vertex v_k is incident to an edge drawn with complexity 1 in Step 2. Since (v_1, v_2) is drawn as a horizontal segment, at least $n - 1$ edges have complexity 1. Planarity is proven by induction; the main invariant is that all edges on $C_k \setminus \{(v_1, v_2)\}$ have a quarter circular arc and possibly a vertical segment. Time and area requirements are shown in App. D. \square

5 Conclusions

In this paper, we continued the study on smooth orthogonal and octilinear drawings. Our \mathcal{NP} -hardness proofs are a first step towards settling the complexity of both drawing problems. We conjecture that the former is \mathcal{NP} -hard, even in the case where only the planar embedding is specified by the input. For the latter, it is of interest to know if it remains \mathcal{NP} -hard even for planar graphs of max-degree 4 or if these graphs allow for a decision algorithm. Our drawing algorithms guarantee bi-monotone 2-drawings with a certain number of complexity-1 edges for maximal planar graphs. Improvements on this number or generalizations to triconnected or simply connected planar graphs are of importance.

Acknowledgements. The authors would like to thank Patrizio Angelini and Martin Gronemann for useful discussions.

References

1. Alam, M.J., Bekos, M.A., Kaufmann, M., Kindermann, P., Kobourov, S.G., Wolff, A.: Smooth orthogonal drawings of planar graphs. In: Pardo, A., Viola, A. (eds.) LATIN. LNCS, vol. 8392, pp. 144–155. Springer (2014)
2. Bekos, M.A., Gronemann, M., Kaufmann, M., Krug, R.: Planar octilinear drawings with one bend per edge. *J. Graph Algorithms Appl.* 19(2), 657–680 (2015)
3. Bekos, M.A., Gronemann, M., Pupyrev, S., Raftopoulou, C.N.: Perfect smooth orthogonal drawings. In: Bourbakis, N.G., Tsihrintzis, G.A., Virvou, M. (eds.) IISA. pp. 76–81. IEEE (2014)
4. Bekos, M.A., Kaufmann, M., Kobourov, S.G., Symvonis, A.: Smooth orthogonal layouts. *J. Graph Algorithms Appl.* 17(5), 575–595 (2013)
5. Bekos, M.A., Kaufmann, M., Krug, R.: On the total number of bends for planar octilinear drawings. In: Kranakis, E., Navarro, G., Chávez, E. (eds.) LATIN. LNCS, vol. 9644, pp. 152–163. Springer (2016)
6. Bertolazzi, P., Di Battista, G., Didimo, W.: Computing orthogonal drawings with the minimum number of bends. *IEEE Trans. Computers* 49(8), 826–840 (2000)
7. Biedl, T.C., Kant, G.: A better heuristic for orthogonal graph drawings. *Comput. Geom.* 9(3), 159–180 (1998)
8. Chrobak, M., Payne, T.H.: A linear-time algorithm for drawing a planar graph on a grid. *Inf. Process. Lett.* 54(4), 241–246 (1995)
9. Di Battista, G., Didimo, W., Patrignani, M., Pizzonia, M.: Orthogonal and quasi-upward drawings with vertices of prescribed size. In: Kratochvíl, J. (ed.) *Graph Drawing*. LNCS, vol. 1731, pp. 297–310. Springer (1999)
10. Di Battista, G., Eades, P., Tamassia, R., Tollis, I.G.: *Graph Drawing: Algorithms for the Visualization of Graphs*. Prentice-Hall (1999)
11. Didimo, W., Liotta, G., Patrignani, M.: On the complexity of hv-rectilinear planarity testing. In: Duncan, C.A., Symvonis, A. (eds.) *Graph Drawing*. LNCS, vol. 8871, pp. 343–354. Springer (2014)
12. Duncan, C.A., Eppstein, D., Goodrich, M.T., Kobourov, S.G., Nöllenburg, M.: Lombardi drawings of graphs. *J. Graph Algorithms Appl.* 16(1), 85–108 (2012)
13. Durocher, S., Felsner, S., Mehrabi, S., Mondal, D.: Drawing hv-restricted planar graphs. In: Pardo, A., Viola, A. (eds.) LATIN. LNCS, vol. 8392, pp. 156–167. Springer (2014)

14. Eppstein, D.: Planar lombardi drawings for subcubic graphs. In: Didimo, W., Patrignani, M. (eds.) *Graph Drawing*. LNCS, vol. 7704, pp. 126–137. Springer (2012)
15. Felsner, S.: *Geometric Graphs and Arrangements*. Advanced Lectures in Mathematics, Vieweg (2004)
16. Fößmeier, U., Heß, C., Kaufmann, M.: On improving orthogonal drawings: The 4M-algorithm. In: Whitesides, S. (ed.) *Graph Drawing*. LNCS, vol. 1547, pp. 125–137. Springer (1998)
17. Fößmeier, U., Kaufmann, M.: Drawing high degree graphs with low bend numbers. In: Brandenburg, F. (ed.) *Graph Drawing*. LNCS, vol. 1027, pp. 254–266. Springer (1995)
18. de Fraysseix, H., Pach, J., Pollack, R.: How to draw a planar graph on a grid. *Combinatorica* 10(1), 41–51 (1990)
19. Fulek, R., Pelsmajer, M.J., Schaefer, M., Stefankovic, D.: Hanani-tutte and monotone drawings. In: Kolman, P., Kratochvíl, J. (eds.) *WG*. LNCS, vol. 6986, pp. 283–294. Springer (2011)
20. Garey, M.R., Johnson, D.S.: *Computers and Intractability: A Guide to the Theory of NP-Completeness*. W. H. Freeman (1979)
21. Garg, A., Tamassia, R.: On the computational complexity of upward and rectilinear planarity testing. *SIAM J. Comput.* 31(2), 601–625 (2001)
22. Giacomo, E.D., Liotta, G., Montecchiani, F.: The planar slope number of subcubic graphs. In: Pardo, A., Viola, A. (eds.) *LATIN*. LNCS, vol. 8392, pp. 132–143. Springer (2014)
23. Harary, F.: *Graph theory*. Addison-Wesley (1991)
24. Hong, S., Merrick, D., do Nascimento, H.A.D.: Automatic visualisation of metro maps. *J. Vis. Lang. Comput.* 17(3), 203–224 (2006)
25. Kant, G.: Drawing planar graphs using the canonical ordering. *Algorithmica* 16(1), 4–32 (1996)
26. Kaufmann, M., Wagner, D. (eds.): *Drawing Graphs, Methods and Models*, LNCS, vol. 2025. Springer (2001)
27. Keszegh, B., Pach, J., Pálvölgyi, D.: Drawing planar graphs of bounded degree with few slopes. *SIAM J. Discrete Math.* 27(2), 1171–1183 (2013)
28. Liu, Y., Morgana, A., Simeone, B.: A linear algorithm for 2-bend embeddings of planar graphs in the two-dimensional grid. *Discrete Applied Mathematics* 81(1-3), 69–91 (1998)
29. Manuch, J., Patterson, M., Poon, S., Thachuk, C.: Complexity of finding non-planar rectilinear drawings of graphs. In: Brandes, U., Cornelsen, S. (eds.) *Graph Drawing*. LNCS, vol. 6502, pp. 305–316. Springer (2010)
30. Nöllenburg, M.: *Automated Drawing of Metro Maps*. Master’s thesis, Fakultät für Informatik, Universität Karlsruhe (TH) (Aug 2005)
31. Nöllenburg, M., Wolff, A.: Drawing and labeling high-quality metro maps by mixed-integer programming. *IEEE Trans. Vis. Comput. Graph.* 17(5), 626–641 (2011)
32. Schnyder, W.: Embedding planar graphs on the grid. In: Johnson, D.S. (ed.) *SODA*. pp. 138–148. SIAM (1990)
33. Stott, J.M., Rodgers, P., Martinez-Ovando, J.C., Walker, S.G.: Automatic metro map layout using multicriteria optimization. *IEEE Trans. Vis. Comput. Graph.* 17(1), 101–114 (2011)
34. Tamassia, R.: On embedding a graph in the grid with the minimum number of bends. *SIAM J. Comput.* 16(3), 421–444 (1987)

Appendix

A Preliminary Notions and Definitions

Unless otherwise specified, we consider simple undirected graphs. Let $G = (V, E)$ be a graph. We denote by n and m the number of vertices and edges of G . We say that G has *max-degree* Δ , if G has no vertex with degree larger than Δ .

A planar drawing Γ of G partitions the plane into connected regions, called *faces*; the unbounded one is called *outerface*. A (*topological*) *planar embedding* \mathcal{E} of G is an equivalence class of planar drawings that define the same set of faces and outerface. Embedding \mathcal{E} can equivalently be defined by the cyclic orders of the edges incident to each vertex (also called *combinatorial embedding*). Given a drawing Γ , a *vertical cut* is a vertical line, which crosses only horizontal edge segments of the drawing and splits it into two parts; a left and right one [16]. If one shifts the right part to the right, while keeping the left part in place, the result has no crossings. A *horizontal cut* is defined analogously.

The *canonical order* for maximal planar graphs [18] is formally defined as follows. Let $G = (V, E)$ be a maximal planar graph and let $\pi = (v_1, \dots, v_n)$ be a permutation of V . Assume that edges (v_1, v_2) , (v_2, v_n) and (v_1, v_n) form the outerface of G . For $k = 1, \dots, n$, let G_k be the subgraph induced by $\cup_{i=1}^k v_i$ and denote by C_k the outerface of G_k . Then, π is a *canonical ordering* of G if for each $k = 2, \dots, n$ the following hold: (i) G_k is biconnected, (ii) all neighbors of P_k in G_{k-1} are on C_{k-1} , and (iii) all vertices of P_k with $2 \leq k < n$ have at least one neighbor in P_j for some $j > k$. A canonical ordering of a maximal planar graph can be computed in linear time [25].

The *shift-method* [18] is a well-known linear-time algorithm, which constructs a planar drawing Γ of a maximal planar graph $G = (V, E)$ on a grid of quadratic area, based on a canonical order π of G as follows. It places v_1, v_2 and v_3 at points $(0, 0)$, $(2, 0)$ and $(1, 1)$. For $k = 4, \dots, n$, assume that a planar drawing Γ_{k-1} of G_{k-1} has been constructed in which each edge of C_{k-1} is drawn as a straight-line segment with slope ± 1 , except for the edge (v_1, v_2) , which is drawn as a horizontal line segment (*contour condition*; see Fig. 10a) and that each of the vertices v_1, \dots, v_{k-1} has been associated with a so-called *shift-set*, which for v_1, v_2 and v_3 are singletons containing only themselves. Let w_1, \dots, w_p be the




Fig. 10: Illustration of the shift-method by de Fraysseix, Pach and Pollack [18].

vertices of C_{k-1} from left to right in Γ_{k-1} , where $w_1 = v_1$ and $w_p = v_2$. Let also (w_ℓ, \dots, w_r) , with $1 \leq \ell < r \leq p$ be the neighbors of v_k from left to right along C_{k-1} in Γ_{k-1} . To avoid edge-overlaps, the algorithm first translates each vertex in $\cup_{i=1}^\ell S(w_i)$ one unit to the left and each vertex in $\cup_{i=r}^p S(w_i)$ one unit to the right, where $S(v)$ denotes the shift-set of $v \in V$. Then, the algorithm places v_k at the intersection of the line with slope $+1$ through w_ℓ with the line with slope -1 through w_r and sets the shift-set of v_k to $\{v_k\} \cup_{i=\ell+1}^{r-1} S(w_i)$; see Fig. 10b.

B Omitted Proofs from Section 2

Theorem 1. *There is an infinitely large family of 4-regular planar graphs that admit both bendless smooth orthogonal and bendless octilinear drawings.*

Proof. For each $k \in \mathbb{N}_+$ we describe a 4-regular planar graph $G_k = (V_k, E_k)$ with $20k$ vertices that admits both a bendless smooth orthogonal drawing and a bendless octilinear drawing; refer to Fig. 3 for the case $k = 2$. G_k has $4k$ subgraphs $W_{i,j}$ such that $1 \leq i \leq 2k$ and $j \in \{t, b\}$ (top and bottom). Graph $W_{i,j}$ consists of five vertices $c_{i,j}$, $n_{i,j}$, $w_{i,j}$, $e_{i,j}$, and $s_{i,j}$ (center, north, west, east, south, respectively), such that $W_{i,j}$ is a wheel on five vertices, i.e., $W_{i,j}$ consists of a center-vertex $c_{i,j}$ and a cycle $C_{i,j} = (n_{i,j}, w_{i,j}, s_{i,j}, e_{i,j})$, such that $c_{i,j}$ is connected to all vertices of cycle $C_{i,j}$.

All vertices except for $c_{i,j}$ already have degree three in $W_{i,j}$. So, we just have to describe the missing edges that make G_k 4-regular: For $1 \leq h \leq 2k-1$, $(e_{h,j}, w_{h+1,j}) \in E_k$ for $j \in \{t, b\}$; blue edges in Fig. 3. Also, $(w_{1,t}, w_{1,b}) \in E_k$ and $(e_{2k,t}, e_{2k,b}) \in E_k$; gray edges in Fig. 3. For $1 \leq h \leq 2k$, $(s_{h,t}, n_{h,b}) \in E_k$; red edges in Fig. 3. Finally, for $1 \leq h \leq k$, $(n_{2h-1,t}, n_{2h,t}) \in E_k$ and $(s_{2h-1,b}, s_{2h,b}) \in E_k$; green edges in Fig. 3. With those additional edges, G_k becomes 4-regular. Fig. 3 is a certificate that $G_k = (V_k, E_k)$ indeed admits both a bendless smooth orthogonal drawing and a bendless octilinear drawing. \square

Lemma 1. *The graph B of Fig. 5a does not admit a bendless smooth orthogonal drawing, when the outerface is fixed to (p_1, q_1, p_2, q_2) , and each of q_1 and q_2 have two unoccupied ports on the outerface.*

Proof. First, we discuss some structural properties of graph B . Observe that graph B contains a wheel W_5 on five vertices as a subgraph, which is induced by the vertices drawn as circles in Fig. 5a. Its center is vertex c and its rim consists of vertices w_1 , w_2 , w_3 , and w_4 . Vertices w_1 and w_2 form a triangular face with vertex t_1 ; vertices w_3 and w_4 form a triangular face with t_2 (vertices t_1 and t_2 are drawn as triangles in Fig. 5a). Observe that t_1 and t_2 form a separation pair and both are connected to vertices p_1 and p_2 (drawn as pentagons in Fig. 5a) forming two pentagonal faces $(p_1, t_1, w_1, w_4, t_2)$ and $(p_2, t_2, w_3, w_2, t_1)$. Observe that p_1 and p_2 also form a separation pair and are both connected to vertices q_1 and q_2 (drawn as squares in Fig. 5a) forming two quadrilateral faces (q_1, p_2, t_1, p_1) and (q_2, p_1, t_2, p_2) . Hence, B has two separation pairs and two vertices of degree 2 (i.e., q_1 and q_2). The remaining vertices of B are of degree exactly 4.




Fig. 11: All possible drawings of (a)-(b) a triangular face, and of (c)-(d) a wheel on five vertices, such that all unoccupied ports are on the outerface.

In order to show that B does not admit a bendless smooth orthogonal drawing, when the outerface is (p_1, q_1, p_2, q_2) , and each of q_1 and q_2 have two unoccupied ports on the outerface, we first observe the following: If we want to draw W_5 such that all of its unoccupied ports are on its outerface, then none of its four triangular faces must have an unoccupied port pointing in its interior. In the bendless smooth orthogonal model, there are only two possible drawings for a triangular face fulfilling this property, as shown in [1]; see Figs. 11a and 11b. This implies that W_5 admits only two bendless smooth orthogonal drawings such that all of its unoccupied ports are on its outerface; see Figs. 11c and 11d.

Next, we consider t_1 and t_2 . Since t_1 and t_2 define two triangular faces in the subgraph induced by W_5 , and t_1 and t_2 , similar as above, we can conclude that there are five different drawings of this graph; see Fig. 12. Note that t_1 and t_2 can be moved along diagonals of slope 1 in Figs. 12d and 12e.

In the following step, we will consider all candidate positions for placing p_1 and p_2 , which we can identify as follows: In a bendless smooth orthogonal drawing, both endpoints of an edge are located along a horizontal, vertical or diagonal line. Both p_1 and p_2 are neighbors of both t_1 and t_2 , for which we already defined their locations. If we consider all rays emanating from t_1 and t_2 with slopes $\{0, 1, -1, \infty\}$, then p_1 and p_2 must be located at an intersection of a ray emanating from t_1 and a ray emanating from t_2 ; see Fig. 13a. For each candidate position, we then try to draw the edges to t_1 and t_2 using one of the edge segments supported by the smooth orthogonal model. The resulting drawing is *valid* if and only if none of the following arises: (i) a vertex has an




Fig. 12: All possible drawings of the subgraph induced by W_5 , t_1 and t_2 , such that all unoccupied ports are on the outerface.




Fig. 13: Method used for identifying valid drawings for p_1 and p_2 : (a) Identification of candidate positions. (b)-(d) Cases of invalid drawings.




Fig. 14: All valid drawings of the subgraph induced by W_5 , t_1 , t_2 , and at least one of p_1 and p_2 .

unoccupied port not incident to the outerface; see Fig. 13b, (ii) a port is used twice; see Fig. 13c, (iii) an edge is not drawn planar; see Fig. 13d. Recall that for the cases shown in Figs. 12d and 12e, we have to take into account all relevant combinations of positions of t_1 and t_2 along the diagonals.

As a result of our analysis, we can conclude that the only valid drawings of the subgraph induced by W_5 , t_1 , t_2 and at least one of p_1 and p_2 are those shown in Fig. 14. Note that in the cases shown in Figs. 14b-14g, we can only place one of p_1 and p_2 . For the case shown in Fig. 14a we proceed by considering all candidate positions of q_1 and q_2 , as we did for p_1 and p_2 . As a result, we conclude that q_1 and q_2 cannot be added such that each of them has two unoccupied ports on the outerface, which completes the proof of this lemma. \square

C Omitted Proofs and Material from Section 3




Fig. 15: Illustration of the parity gadget when $\ell(x) \approx \ell(\bar{x})$.

Lemma 2. *For the smooth orthogonal model, the parity gadget of variable x of formula φ contains no edge crossings if and only if $|\ell(\bar{x}) - \ell(x)| > \frac{\sqrt{3}}{2}\ell(u)$, where $\ell(u)$ denotes the unit length.*

Proof. Refer to Fig. 16a, which gives a more detailed illustration of the vertical gap of the parity gadget. Consider the case where $x = \mathbf{false}$. The case where $x = \mathbf{true}$ is symmetric. If $x = \mathbf{false}$, we want the bottom arc cut by the dashed diagonal to be completely below the top arc cut by the dashed diagonal. Since we know that both arcs have radius $\ell(u)$, their centers (gray-colored in Fig. 16a) should be a distance of $> 2 \cdot \ell(u)$ apart from each other. This corresponds to the length of the dashed diagonal. We can compute the length of the dashed diagonal in dependence of $\lambda = \ell(\bar{x}) - \ell(x)$ as the dashed diagonal is part of a right triangle for which we know the remaining side lengths. Thus, we can use Pythagoras' theorem and compute the length of the diagonal which gives us $\lambda > \sqrt{3}/2 \cdot \ell(u) \approx 0.866 \cdot \ell(u)$. \square

The corresponding parity gadget for the octilinear model is illustrated in Fig. 16b.

Lemma 3. *For the octilinear model, the parity gadget of variable x of formula φ contains no edge crossings if and only if $|\ell(\bar{x}) - \ell(x)| > \frac{5}{6}\ell(u)$, where $\ell(u)$ denotes the unit length.*

Proof. Refer to Fig. 16b. Consider the case where $x = \mathbf{false}$. The case where $x = \mathbf{true}$ is symmetric. In comparison to the smooth orthogonal setting we subdivide several axis-aligned edges of unit edge length $\ell(u)$ into three equally sized smaller edges (see the edges with endpoints drawn as squares in the figure) of length $\frac{1}{3}\ell(u)$. Note that we can realize this by actually composing each $\ell(u)$ in the representation of three edges of size $\frac{1}{3}\ell(u)$ instead.




Fig. 16: Illustrations for the proofs of Lemmas 2 and 3.

Since we do not deal with circular arc segments here, the smallest distance $d > 0$ between both blocks is easy to compute; in particular it is located between the bottom diagonal of the upper block and the middle diagonal of the lower block which are parallel. For instance, d is the distance between the bottom endpoint of the bottom diagonal of the upper block v_b (gray squared-shaped vertex in Fig. 16b) and the point on the diagonal of the lower block that is vertically below v_b (gray cross in Fig. 16b) which is not a vertex). For both, we know the y -coordinates $2 \cdot \ell(x)$ and $2 \cdot \ell(\bar{x}) + 5/3 \cdot \ell(u)$, respectively. Hence, $d = 2 \cdot (\ell(\bar{x}) - \ell(x)) - 5/3 \cdot \ell(u) > 0$ which implies $\ell(\bar{x}) - \ell(x) > 5/6 \cdot \ell(u)$. \square

D Omitted Proofs and Material from Section 4

Theorem 8. *A maximal planar n -vertex graph admits a bi-monotone planar octilinear 2-drawing in the Kandinsky model, which requires $O(n^2)$ area and can be computed in $O(n)$ time. In addition, each bend is at 135° .*




Fig. 17: Illustration of the modified shift-method for the octilinear Kandinsky model.

Proof. The proof is rather simple. We can actually convert the layout computed for the smooth orthogonal model to octilinear by redrawing all quarter circular arcs of it as diagonal segments; see also Fig. 17b. This results in bends at 135° . Planarity follows from blue and green edges not passing through vertices by virtue of construction. \square




Fig. 18: Stretching an edge containing a vertical segment of length δ .

Theorem 7. *A maximal planar n -vertex graph admits a bi-monotone planar smooth orthogonal 2-drawing with at least $n - 1$ edges with complexity 1 in the Kandinsky model, which requires $O(n^4)$ area and can be computed in $O(n^2)$ time.*

Proof. The time complexity follows from the fact that in Steps 1 and 2 of our algorithm we may have to stretch the drawing a linear number of times.




Fig. 19: Proof of the area requirement.

For the area requirement, we first consider Step 1. Recall that in this step we stretch each red edge that is not a vertical line segment. Note that the red edges form a tree, which implies that we stretch at most $n - 1$ edges. If at some point we stretch due to a red edge (u, v) such that vertex u is to the right of v and above v , our algorithm guarantees that we will not stretch again due to another edge (u, w) with u to the right of w and above w . This implies that the blue and the green edges incident to u define a rectangle R that will not contain another edge that must be stretched later in the algorithm (refer to the gray colored rectangle in Fig. 19). If (u, v) is composed of a vertical segment of length δ and

a quarter circle arc of radius ρ , then the entire drawing is stretched by δ . This fixes the length of all segments necessary to compute the area of rectangle R ; see Fig. 19. In total, the area of R is $\mathcal{A}(R) = \frac{1}{2}\delta^2 + \rho\delta$. Obviously, $\mathcal{A}(R)$ is minimized when $\rho = 1$, for a constant δ , which reduces the formula to $\mathcal{A}(R) = \frac{1}{2}\delta^2 + \delta$, which grows quadratically in δ . Since the area of the drawing computed before the application of Step 1 is $O(n^2)$, the rectangles of red edges are on average of $O(n)$ area. Note that by planarity no two such rectangles overlap. Since the area of each rectangle grows quadratically in the length of the vertical segment of its associated red edge, the drawing requires the most total stretching, if each red edge is stretched by the same amount. In this case, $1/2\delta^2 + \delta = O(n)$, which implies that $\delta = O(\sqrt{n})$. Hence, the drawing obtained by Step 1 of our algorithm has width $O(n^{3/2})$.

For each vertex in Step 2 of our algorithm, we may increase the height of the drawing obtained by Step 1 by an amount that is bounded by the width of the initial drawing, i.e., $O(n^{3/2})$. This leads to a total height of $O(n^{5/2})$ and therefore a total area bound of $O(n^4)$. \square

E Example Run of our Drawing Algorithm

In this section, we describe an example run of our drawing algorithm from Section 4 on the octahedron graph. Figs. 20 and 21 illustrate Steps 1 and 2, respectively. In particular, Fig. 20a shows the output of our modification of the shift-method and the first vertical cut. Fig. 20b shows the result of the first horizontal stretching and the second vertical cut. Fig. 20c shows the result after the second horizontal stretching, which is also the output of Step 1.




Fig. 20: Example run of Step 1 of our drawing algorithm.

Fig. 21 illustrates how we assign new y -coordinates to the vertices in Step 2 of our algorithm. In particular, Fig. 21a shows how this is done for the first three vertices. Figs. 21b, 21c and 21d illustrate how the fourth, the fifth and the sixth vertex of the octahedron is attached to the drawing. The bold edges in each subfigure of Fig. 21 are the ones defining the maximum (denoted by (v_k, w^*)) in the description of the algorithm. The final drawing is the one of Fig. 21d.




Fig. 21: Example run of Step 2 of our drawing algorithm.