Spectral Operators of Matrices

Chao Ding^{*}, Defeng Sun[†], Jie Sun[‡] and Kim-Chuan Toh[§]

January 10, 2014

Abstract

The class of matrix optimization problems (MOPs) has been recognized in recent years to be a powerful tool by researchers far beyond the optimization community to model many important applications involving structured low rank matrices. This trend can be credited to some extent to the exciting developments in the emerging field of compressed sensing. The Löwner operator, which generates a matrix valued function by applying a single-variable function to each of the singular values of a matrix, has played an important role for a long time in solving matrix optimization problems. However, the classical theory developed for Löwner operators has become inadequate in these recent applications. The main objective of this paper is to provide some necessary theoretical foundations for designing numerical methods for solving the MOP. This goal is achieved by introducing and conducting a thorough study on a new class of matrix valued functions, coined as spectral operators of matrices. Several fundamental properties of spectral operators, including the well-definedness, continuity, directional differentiability, Fréchet-differentiability, locally Lipschitzian continuity, ρ -order B(ouligand)differentiability ($0 < \rho \le 1$), ρ -order G-semismooth ($0 < \rho \le 1$) and the characterization of Clarke's generalized Jacobian, are systematically studied.

AMS subject classifications: 90C25, 90C06, 65K05, 49J50, 49J52

OR/MS subject classifications: Primary: Mathematics/matrices; Secondary: Mathematics/functions **Key Words**: directional differentiability; Fréchet differentiability; matrix valued functions; proximal mappings; semismoothness; spectral operators

1 Introduction

Let $\mathbb{R}^{m \times n}$ and $\mathbb{C}^{m \times n}$ be the vector spaces of $m \times n$ real and complex matrices over the scalar field of real numbers \mathbb{R} , respectively. For any $X \in \mathbb{C}^{m \times n}$, we denote the conjugate transpose of X by $X^{\mathbb{T}}$. If $X \in \mathbb{R}^{m \times n}$, then $X^{\mathbb{T}}$ is just the transpose of X. We use $\mathbb{V}^{m \times n}$ to represent either the real Euclidean vector space $\mathbb{R}^{m \times n}$ or $\mathbb{C}^{m \times n}$ with the trace inner product $\langle X, Y \rangle := \operatorname{Re}(\operatorname{trace}(X^{\mathbb{T}}Y))$ for $X, Y \in \mathbb{V}^{m \times n}$ and its induced norm $\|\cdot\|$, where "Re" means the real part of a complex number. Without loss of generality, we assume that $m \leq n$ throughout this paper. For convenience, we also call $\mathbb{V}^{m \times n}$ a matrix space.

Let $\mathbb{S}^m \subseteq \mathbb{V}^{m \times m}$ be the real vector subspace of $m \times m$ real symmetric matrices or complex Hermitian matrices. For any given $Y \in \mathbb{S}^m$, we use $\lambda_1(Y) \ge \lambda_2(Y) \ge \ldots \ge \lambda_m(Y)$ to denote the eigenvalues of Y(all real and counting multiplicity) and use $\lambda(Y)$ to denote the vector of eigenvalues of Y. For any given $Z \in \mathbb{V}^{m \times n}$, we use $\sigma_1(Z) \ge \sigma_2(Z) \ge \ldots \ge \sigma_m(Z)$ to denote the singular values of Z (always nonnegative

^{*}National Center for Mathematics and Interdisciplinary Sciences, Chinese Academy of Sciences, Beijing, China. This work was initiated while C. Ding was with Department of Mathematics, National University of Singapore during 2007 to 2012. Email: dingchao@amss.ac.cn.

[†]Department of Mathematics and Risk Management Institute, National University of Singapore, Singapore. Email: matsundf@nus.edu.sg.

[‡]Department of Mathematics and Statistics, Curtin University, Australia. Email: sun.curtin@gmail.com.

[§]Department of Mathematics, National University of Singapore, Singapore. Email: mattohkc@nus.edu.sg.

and counting multiplicity) and use $\sigma(Z)$ to denote the vector of the singular values of Z. We use \mathbb{O}^p (p = m, n) to denote the set of $p \times p$ orthogonal matrices in $\mathbb{R}^{p \times p}$ if $\mathbb{V}^{m \times n} = \mathbb{R}^{m \times n}$ and the set of $p \times p$ unitary matrices in $\mathbb{C}^{p \times p}$ if $\mathbb{V}^{m \times n} = \mathbb{C}^{m \times n}$. For $X \in \mathbb{V}^{m \times m}$, diag(X) denotes the column vector consisting of all the diagonal entries of X being arranged from the first to the last and for $x \in \mathbb{R}^m$, Diag(x) denotes the m by m diagonal matrix whose *i*-th diagonal entry is $x_i, i = 1, \ldots, m$.

In this paper, we shall introduce and study a class of matrix valued functions, to be called spectral operators of matrices. This class of matrix valued functions frequently arise in various applications. Our first motivating application comes from matrix optimization problems (MOPs). Let \mathcal{X} be the vector space $\mathbb{V}^{m \times n}$ or \mathbb{S}^n . Suppose that $f : \mathcal{X} \to (-\infty, \infty]$ is a closed proper convex function. One simple class of MOPs just mentioned take the form of

min
$$f_0(X) + f(X)$$

s.t. $\mathcal{A}X = b, \quad X \in \mathcal{X},$ (1)

where $f_0: \mathcal{X} \to \mathbb{R}$ is a smooth function whose gradient is Lipschitz continuous, e.g., a linear function $f_0(\cdot) = \langle C, \cdot \rangle$ for some $C \in \mathcal{X}, \mathcal{A} : \mathcal{X} \to \mathbb{R}^p$ is a linear operator, and $b \in \mathbb{R}^p$ is a given vector. The above MOPs cover many problems as special cases. For example, by considering the particular case that $f \equiv \delta_{\mathbb{S}^m_+}$, the indicator function of the positive semidefinite matrix cone \mathbb{S}^m_+ , we can see that the extensively studied semidefinite programming (SDP) [53] is in the form of (1) with a linear function f_0 . MOPs also arise frequently from other applications such as the matrix norm approximation, matrix completion, rank minimization, graph theory, machine learning, etc [21, 54, 55, 43, 2, 3, 4, 6, 57, 9, 20, 33, 17, 24]. See [15] for more details.

The Karush-Kuhn-Tucker (KKT) condition of (1) can be written in the following form [45, Corollary 28.3.1]:

$$\begin{cases} \nabla f_0(X) - \mathcal{A}^* y + \Gamma = 0, \\ \mathcal{A}X - b = 0, \\ \Gamma \in \partial f(X). \end{cases}$$
(2)

Let $\psi_f : \mathcal{X} \to \mathbb{R}$ be the Moreau-Yosida regularization of the closed proper convex function f, i.e.,

$$\psi_f(X) := \min_{Y \in \mathcal{X}} \left\{ f(Y) + \frac{1}{2} \|Y - X\|^2 \right\}, \quad X \in \mathcal{X},$$
(3)

and $P_f(X)$ be the proximal mapping of f at X, the unique optimal solution to (3). It is well-known (see e.g., [46, Proposition 12.19]) that the mapping $P_f : \mathcal{X} \to \mathcal{X}$ is globally Lipschitz continuous on \mathcal{X} with modulus 1 and ψ_f is continuously differentiable on \mathcal{X} with $\nabla \psi_f(X) = X - P_f(X)$. From [36] (see also [45, Theorem 31.5]) we know that the KKT condition (2) is equivalent to the following system of Lipschitzian equations

$$\begin{bmatrix} \nabla f_0(X) - \mathcal{A}^* y + \Gamma \\ \mathcal{A}X - b \\ X - P_f(X + \Gamma) \end{bmatrix} = 0.$$

Thus, the study of MOPs depends crucially on the study of various differential properties of P_f . In [60, 7, 31], Newton-CG based proximal-point algorithms have been designed to solve large scale SDPs, matrix spectral norm approximation, and nuclear norm minimization problems, respectively. Those algorithms and their convergence analyzes all depend crucially on understanding the various differential properties of the associated proximal mappings P_f .

For any given $Z \in \mathbb{V}^{m \times n}$, let $\mathbb{O}^{m,n}(Z)$ denote the set of matrix pairs $(U, V) \in \mathbb{O}^m \times \mathbb{O}^n$ satisfying the singular value decomposition $Z = U[\Sigma(Z) \quad 0] V^{\mathbb{T}}$, where $\Sigma(Z)$ is an $m \times m$ diagonal matrix whose *i*-th diagonal entry is $\sigma_i(Z) \ge 0$. For any given $Y \in \mathbb{S}^m$, we use $\mathbb{O}^m(Y)$ to denote the set of matrices $P \in \mathbb{O}^m$ satisfying the eigenvalue decomposition $Y = P\Lambda(Y)P^{\mathbb{T}}$, where $\Lambda(Y)$ is an $m \times m$ diagonal matrix whose *i*-th diagonal entry is $\lambda_i(Y)$, a real number. Assume that the closed proper convex function $f : \mathcal{X} \to (-\infty, \infty]$ is unitarily invariant, i.e., for any $X \in \mathcal{X} \equiv \mathbb{V}^{m \times n}$, $U \in \mathbb{O}^m$ and $V \in \mathbb{O}^n$, $f(X) = f(U^{\mathbb{T}}XV)$, or for any

 $X \in \mathcal{X} \equiv \mathbb{S}^m, P \in \mathbb{O}^m, f(X) = f(P^{\mathbb{T}}XP)$. For example, for a given $k \in \{1, \ldots, m\}$, Ky Fan's k-norm matrix function $\|X\|_{(k)} = \sum_{i=1}^k \sigma_i(X), X \in \mathbb{V}^{m \times n}$ and the indicator function $\delta_{\mathbb{S}^m_+}$ are unitarily invariant. Recall that a function $\phi : \mathbb{R}^m \to (-\infty, \infty]$ is said to be symmetric if $\phi(x) = \phi(Qx) \forall x \in \mathbb{R}^m$ and any permutation matrix Q, and is said to be absolutely symmetric if $\phi(x) = \phi(Qx) \forall x \in \mathbb{R}^m$ and any signed permutation matrix Q, which has exactly one nonzero entry in each row and each column, and that entry being ± 1 . For the unitarily invariant function $f : \mathcal{X} \to (-\infty, \infty]$, we know from Lewis [26, Proposition 2.1] and Davis [11] that there exists an absolutely symmetric function $\theta : \mathbb{R}^m \to (-\infty, +\infty]$ such that $f(\cdot) \equiv \theta(\sigma(\cdot))$ if $\mathcal{X} \equiv \mathbb{V}^{m \times n}$ and a symmetric function $\theta : \mathbb{R}^m \to (-\infty, +\infty]$ such that $f(\cdot) \equiv \theta(\lambda(\cdot))$ if $\mathcal{X} \equiv \mathbb{V}^m$ and a symmetric function $\theta : \mathbb{R}^m \to (-\infty, +\infty]$ such that $f(\cdot) \equiv \theta(\lambda(\cdot))$ if $\mathcal{X} \equiv \mathbb{V}^m \times n$ and a symmetric function $\theta : \mathbb{R}^m \to (-\infty, +\infty]$ such that $f(\cdot) \equiv \theta(\lambda(\cdot))$ if $\mathcal{X} \equiv \mathbb{V}^m \times n$ and a symmetric function $\theta : \mathbb{R}^m \to (-\infty, +\infty]$ such that $f(\cdot) \equiv \theta(\lambda(\cdot))$ if $\mathcal{X} \equiv \mathbb{V}^m \times n$ and a symmetric function $\theta : \mathbb{R}^m \to (-\infty, +\infty]$ such that $f(\cdot) \equiv \theta(\lambda(\cdot))$ if $\mathcal{X} \equiv \mathbb{V}^m \times n$ and a symmetric function $\theta : \mathbb{R}^m \to (-\infty, +\infty)$ such that $f(\cdot) \equiv \theta(\lambda(\cdot))$ if $\mathcal{X} \equiv \mathbb{V}^m \times n$ and a symmetric function $\theta : \mathbb{R}^m \to (-\infty, +\infty)$ such that $f(\cdot) \equiv \theta(\lambda(\cdot))$ if $\mathcal{X} \equiv \mathbb{V}^m \times n$ and a symmetric function $\theta : \mathbb{R}^m \to (-\infty, +\infty)$ such that $f(\cdot) \equiv \theta(\lambda(\cdot))$ if $\mathcal{X} \equiv \mathbb{V}^m \times n$ and a symmetric function $\theta : \mathbb{R}^m \to (-\infty, +\infty)$ such that $f(\cdot) \equiv \theta(\lambda(\cdot))$ if $\mathcal{X} \equiv \mathbb{V}^m$, respectively. Furthermore, from [27, 26, 29], we know that the proximal mapping $P_f : \mathcal{X} \to \mathcal{X}$ can be written as

$$P_f(X) = \begin{cases} U \left[\text{Diag} \left(P_{\theta}(\sigma(X)) \right) & 0 \right] V^{\mathbb{T}} & \text{if } X \in \mathcal{X} \equiv \mathbb{V}^{m \times n}, \\ P \text{Diag} \left(P_{\theta}(\lambda(X)) \right) P^{\mathbb{T}} & \text{if } X \in \mathcal{X} \equiv \mathbb{S}^m, \end{cases}$$

where $(U, V) \in \mathbb{O}^{m,n}(X)$ if $X \in \mathbb{V}^{m \times n}$ and $P \in \mathbb{O}^m(X)$ if $X \in \mathbb{S}^m$. The proximal mapping P_f is a spectral operator with respect to the mixed symmetric mapping $P_{\theta} : \mathbb{R}^m \to \mathbb{R}^m$ (see Definition 2.1 in Section 2).

Proximal mappings of unitarily invariant proper closed convex functions belong to a class of matrix functions studied previously in two seminal papers by Lewis [27], and Lewis and Sendov [28]. In [27], Lewis defined a Hermitian matrix valued function by using the gradient mapping $g(\cdot) = \nabla \phi(\cdot) : \mathbb{R}^m \to \mathbb{R}^m$ \mathbb{R}^m of a symmetric function $\phi : \mathbb{R}^m \to (-\infty, \infty]$. The corresponding Hermitian matrix valued function $G : \mathbb{S}^m \to \mathbb{S}^m$ is defined by $G(Y) = \sum_{i=1}^m g_i(\lambda(Y)) p_i p_i^{\mathbb{T}}$, where $P \in \mathbb{O}^m(Y)$ and p_i is the *i*-th column of P. Lewis [27] proved that such a function G is well-defined, by using the "block-refineness" property of g. It was further shown by Lewis and Sendov in [28] that G is (continuously) differentiable at X if and only if q is (continuously) differentiable at $\lambda(X)$. Qi and Yang [39] proved that the locally Lipschitz continuous function G is (strongly) semismooth at X if and only if g is (strongly) semismooth at $\lambda(X)$. Note that if the function g has the form $g(y) = (h(y_1), \ldots, h(y_m)) \forall y \in \mathbb{R}^m$ for some given real valued functional $h : \mathbb{R} \to \mathbb{R}$, then the corresponding Hermitian matrix valued function G is called Löwner's (Hermitian) operator [32], which has been well-studied in the literature. See e.g., [8, 52] for more details. For the non-Hermitian case, by considering the gradient mapping $g(\cdot) = \nabla \phi(\cdot) : \mathbb{R}^m \to \mathbb{R}^m$ of an absolutely symmetric function $\phi : \mathbb{R}^m \to (-\infty, \infty]$, Lewis [26] studied the corresponding matrix valued function by $G(Z) = \sum_{i=1}^m g_i(\sigma(Z))u_iv_i^{\mathbb{T}}$ for $Z \in \mathbb{V}^{m \times n}$, where $(U, V) \in \mathbb{O}^{m,n}(Z)$ and u_i and v_i are the *i*-th column of Uand V, respectively. See also Lewis and Sendov [29] for more details. If the function g has the form g(z) = $(h(z_1),\ldots,h(z_m))^T$ $\forall z \in \mathbb{R}^m$ for some given real valued functional $h: \mathbb{R} \to \mathbb{R}$ satisfying h(0) = 0, then the corresponding non-Hermitian matrix valued function G is called Löwner's (non-Hermitian) operator in Yang's thesis [59]. Some important properties of Löwner's (non-Hermitian) operators have been studied by Yang in [59], including the well-definedness, (continuous) differentiability and (strongly) semismoothness.

Besides MOPs, the proximal mapping P_f has also played a crucial role in some recent applications. For instance, by employing its differentiability and the corresponding derivative formulas, one can derive the divergence expression of the proximal mapping P_f , which can be used to obtain the Stein unbiased risk estimator (SURE) [48] of the regularization solution of the matrix recovery problem involving the nuclear norm regularization. Candés et al. [5] provided a parameter selection method based on the SURE for the singular value thresholding (SVT) operator. See also [12] for more details. Although some partial work has been done on different cases, many fundamental problems of the proximal mapping are unsolved. For example, even in the Hermitian case, there still remain two important issues to be explored (i) an explicit formula for the directional derivative of G and (ii) the characterization of Clarke's generalized Jacobian of the general Hermitian matrix valued function G.

The spectral operators of matrices to be considered in this paper go much beyond proximal mappings. As a matter of fact, the spectral operators arising from applications may not even be the gradient mapping of any scalar valued (absolutely) symmetric function. Therefore, the theoretical results on the spectral operators obtained in this paper are not covered by the previous works just mentioned [27, 28, 39, 59]. For instance, such spectral operators have already been used in low-rank matrix completion problems with

fixed basis coefficients [34]. The problem of low-rank matrix completion aims to recover an unknown low-rank matrix from some under-sampled observations with or without noises. A basic approach to solve a low-rank matrix completion problem is to minimize the rank of a matrix subject to certain constraints consistent with the sampled observations. Since minimizing a rank function with constraints is generally NP-hard, a widely-used convex relaxation approach is to replace the rank function with the nuclear norm. For various theoretical breakthroughs along this direction, we refer the readers to [2, 3, 22, 23, 42, 43] and references therein. However, since for many situations, such as the correlation matrix completion in statistics and the density matrix completion in the quantum state tomography where the nuclear norm is a constant, the efficacy of the nuclear norm minimization approach for recovery is inadequate, Miao et al. [34] proposed a rank-corrected procedure to generate an estimator of high accuracy and low rank, in which non-traditional spectral operators play a pivotal role. A rank-correction term of the form $-\langle G(\tilde{X}), X \rangle$ was added to the nuclear norm penalized least squares model, where $\tilde{X} \in \mathbb{V}^{m \times n}$ is a given initial estimator and $G: \mathbb{V}^{m \times n} \to \mathbb{V}^{m \times n}$ is a matrix-valued function defined by

$$G(X) = U \left[\text{Diag} \left(g(\sigma(X)) \right) \quad 0 \right] V^{\mathbb{T}}, \quad X \in \mathbb{V}^{m \times n}$$

with $(U, V) \in \mathbb{O}^{m,n}(X)$, and $g : \mathbb{R}^m \to \mathbb{R}^m$ is given by

$$g_i(x) = h\left(\frac{x_i}{\|x\|_{\infty}}\right) \text{ if } x \in \mathbb{R}^m \setminus \{0\}, \quad g_i(0) = 0$$

$$\tag{4}$$

for some scalar valued function $h : \mathbb{R} \to \mathbb{R}$. For example, for given $\varepsilon, \tau > 0$, the following h was considered in [34]:

$$h(t) = \operatorname{sgn}(t)(1+\varepsilon^{\tau})\frac{|t|^{\tau}}{|t|^{\tau}+\varepsilon^{\tau}}, \quad t \in \mathbb{R}.$$
(5)

It can be checked that G is the spectral operator with respect to the absolutely symmetric mapping g (Definition 2.1). Note that for such a spectral operator G, there does not exist a function $\psi : \mathbb{V}^{m \times n} \to \mathbb{R}$ such that G is derived through the gradient mapping of ψ because the Jacobian of G at X, when it exists, is not self-adjoint. By using the rank-correction term, Miao et al. [34] established a non-asymptotic recovery error result and provided necessary and sufficient conditions for rank consistency. Various properties of spectral operators such as well-definedness and continuity play an important role in their study. More discussions on the rank-correction function can be found in Miao's PhD thesis [35].

Spectral operators of matrices can also be used in some other related areas such as in statistical shape analysis, which involves low rank matrices. For instance, in order to establish necessary and sufficient conditions on the existence of the extrinsic mean shape for the reflection shape space (see e.g., [1] for the definition) and to provide the corresponding explicit formula (which has important applications in biology, medicine, image analysis, archeology, etc (cf. [18])), very recently Ding and Qi [16] used the following matrix valued function $G : \mathbb{S}^m \to \mathbb{S}^m$ defined by

$$G(X) = P \operatorname{Diag}(g(\lambda(X))) P^{\mathbb{T}}, \quad X \in \mathbb{S}^m$$

with $P \in \mathbb{O}^m(X)$, and $g : \mathbb{R}^m \to \mathbb{R}^m$ being given by $g(x) = Q^{\mathbb{T}}p(x)$, where for $x \in \mathbb{R}^m$, Q is an m by m permutation matrix such that $Qx = x^{\downarrow}$, the vector of entries of x being arranged in the non-increasing order $x_1^{\downarrow} \ge \ldots \ge x_m^{\downarrow}$ and p(x) is the unique optimal solution to the following convex optimization problem

$$\min\left\{\frac{1}{2}\|y-x^{\downarrow}\|^2 \mid \sum_{i=1}^m y_i = 1, \ y_1 \ge \ldots \ge y_k \ge 0, \ y_{k+1} = \ldots = y_m = 0\right\},\$$

where $1 \leq k \leq m$ is a given integer to indicate the rank of a desired matrix. For a certain nonempty open set $\mathcal{N} \in \mathbb{S}^m$, e.g., $\mathcal{N} = \{X \in \mathbb{S}^m \mid \lambda_{k-1}(X) > \lambda_k(X) > \lambda_{k+1}(X)\}$, one can easily check that $g : \mathbb{R}^m \to \mathbb{R}^m$ is symmetric (see Definition 2.1) on $\lambda_{\mathcal{N}} := \{\lambda(X) \mid X \in \mathcal{N}\}$ and the defined matrix function G is a spectral operator on \mathcal{N} . The remaining parts of this paper are organized as follows. In Section 2, we give the definition of spectral operators of matrices and study their well-definedness. Some preliminary results on the differential properties of singular values and vectors of matrices are also given in this section. We study the continuity, directional and Fréchet-differentiability of spectral operators defined on the single matrix space $\mathbb{V}^{m \times n}$ in Section 3. More sophisticated differential properties such as Bouligand-differentiability and *G*-semismoothness of spectral operators are presented in Section 4. In Section 5, we study the spectral operators defined on the Cartesian product of several matrix spaces, and list the main results corresponding to those derived in Sections 3 and 4. We conclude our paper in the final section.

2 Spectral operators of matrices

In this section, we will first define spectral operators on the Cartesian product of several real or complex matrix spaces. The study of spectral operators under this general setting is not only useful but also necessary. In fact, spectral operators defined on the Cartesian product of several matrix spaces appear naturally in the study of the differentiability of spectral operators, even if they are only defined on a single matrix space (see Section 3.1). Moreover, the spectral operators used in many applications are defined on the Cartesian product of several matrix space. See e.g., [14, 58] for more details.

Let s be a positive integer and $0 \le s_0 \le s$ be a nonnegative integer. For given positive integers m_1, \ldots, m_s and n_{s_0+1}, \ldots, n_s , define the finite dimensional real vector space \mathcal{X} by

$$\mathcal{X} := \mathbb{S}^{m_1} \times \ldots \times \mathbb{S}^{m_{s_0}} \times \mathbb{V}^{m_{s_0+1} \times n_{s_0+1}} \times \ldots \times \mathbb{V}^{m_s \times n_s}$$

Without loss of generality, we assume that $m_k \leq n_k$, $k = s_0 + 1, \ldots, s$. For any $X = (X_1, \ldots, X_s) \in \mathcal{X}$, we have for $1 \leq k \leq s_0$, $X_k \in \mathbb{S}^{m_k}$ and $s_0 + 1 \leq k \leq s$, $X_k \in \mathbb{V}^{m_k \times n_k}$. Denote $m_0 := \sum_{k=1}^{s_0} m_k$ and $m = \sum_{k=s_0+1}^{s} m_k$. For any $X \in \mathcal{X}$, define $\kappa(X) \in \mathbb{R}^{m_0+m}$ by

$$\kappa(X) := (\lambda(X_1), \dots, \lambda(X_{s_0}), \sigma(X_{s_0+1}), \dots, \sigma(X_s))$$

Recall that a matrix $Q \in \mathbb{R}^{p \times p}$ is said to be a signed permutation matrix if Q has exactly one nonzero entry in each row and each column and that entry being ± 1 . Let \mathbb{P}^p and $\pm \mathbb{P}^p$ be the sets of all $p \times p$ permutation matrices and signed permutation matrices, respectively. For \mathcal{X} , define the set \mathcal{P} by

$$\mathcal{P} := \{ (Q_1, \dots, Q_s) \mid Q_k \in \mathbb{P}^{m_k}, \ 1 \le k \le s_0 \text{ and } Q_k \in \pm \mathbb{P}^{m_k}, \ s_0 + 1 \le k \le s \} .$$

Let $\mathbf{g} : \mathbb{R}^{m_0+m} \to \mathbb{R}^{m_0+m}$ be a given mapping. For any $\mathbf{x} = (\mathbf{x}_1, \dots, \mathbf{x}_s) \in \mathbb{R}^{m_0+m}$ with $\mathbf{x}_k \in \mathbb{R}^{m_k}$, rewrite $\mathbf{g}(\mathbf{x}) \in \mathbb{R}^{m_0+m}$ in the form $\mathbf{g}(\mathbf{x}) = (\mathbf{g}_1(\mathbf{x}), \dots, \mathbf{g}_s(\mathbf{x}))$ with $\mathbf{g}_k(\mathbf{x}) \in \mathbb{R}^{m_k}$ for $1 \le k \le s$.

Definition 2.1 The given mapping $\mathbf{g} : \mathbb{R}^{m_0+m} \to \mathbb{R}^{m_0+m}$ is said to be mixed symmetric, with respect to \mathcal{P} , at $\mathbf{x} = (\mathbf{x}_1, \dots, \mathbf{x}_s) \in \mathbb{R}^{m_0+m}$ with $\mathbf{x}_k \in \mathbb{R}^{m_k}$, if

$$\mathbf{g}(Q_1\mathbf{x}_1,\ldots,Q_s\mathbf{x}_s) = (Q_1\mathbf{g}_1(\mathbf{x}),\ldots,Q_s\mathbf{g}_s(\mathbf{x})) \quad \forall \ (Q_1,\ldots,Q_s) \in \mathcal{P}.$$
(6)

The mapping **g** is said to be mixed symmetric, with respect to \mathcal{P} , over a set $\mathcal{D} \subseteq \mathbb{R}^{m_0+m}$ if (6) holds for every $\mathbf{x} \in \mathcal{D}$. We call **g** a mixed symmetric mapping, with respect to \mathcal{P} , if (6) holds for every $\mathbf{x} \in \mathbb{R}^{m_0+m}$.

Note that for each $k \in \{1, \ldots, s\}$, the function value $\mathbf{g}_k(\mathbf{x}) \in \mathbb{R}^{m_0+m}$ is dependent on all $\mathbf{x}_1, \ldots, \mathbf{x}_s$. With causing no confusion, in later discussions we often drop "with respect to \mathcal{P} " from Definition 2.1. The following result on \mathbf{g} can be checked directly from the definition.

Proposition 2.1 Suppose that the mapping $\mathbf{g} : \mathbb{R}^{m_0+m} \to \mathbb{R}^{m_0+m}$ is mixed symmetric at $\mathbf{x} = (\mathbf{x}_1, \dots, \mathbf{x}_s) \in \mathbb{R}^{m_0+m}$ with $\mathbf{x}_k \in \mathbb{R}^{m_k}$. Then, for any $i, j \in \{1, \dots, m_k\}$,

$$(\mathbf{g}_k(\mathbf{x}))_i = (\mathbf{g}_k(\mathbf{x}))_j$$
 if $(\mathbf{x}_k)_i = (\mathbf{x}_k)_j$, $\forall 1 \le k \le s$

and

$$(\mathbf{g}_k(\mathbf{x}))_i = 0$$
 if $(\mathbf{x}_k)_i = 0$, $\forall s_0 + 1 \le k \le s$.

Let \mathcal{N} be a given nonempty set in \mathcal{X} . Define

$$\kappa_{\mathcal{N}} := \{ \kappa(X) \mid X \in \mathcal{N} \} .$$

Definition 2.2 Suppose that $\mathbf{g} : \mathbb{R}^{m_0+m} \to \mathbb{R}^{m_0+m}$ is mixed symmetric on κ_N . The spectral operator $G : \mathcal{N} \to \mathcal{X}$ with respect to \mathbf{g} is defined by

$$G(X) := (G_1(X), \dots, G_s(X)), \quad X = (X_1, \dots, X_s) \in \mathcal{N}$$

with

$$G_k(X) := \begin{cases} P_k \operatorname{Diag}(\mathbf{g}_k(\kappa(X))) P_k^{\mathbb{T}} & \text{if } 1 \le k \le s_0, \\ U_k \left[\operatorname{Diag}(\mathbf{g}_k(\kappa(X))) & 0 \right] V_k^{\mathbb{T}} & \text{if } s_0 + 1 \le k \le s \end{cases}$$

where $P_k \in \mathbb{O}^{m_k}(X_k), \ 1 \le k \le s_0, \ (U_k, V_k) \in \mathbb{O}^{m_k, n_k}(X_k), \ s_0 + 1 \le k \le s.$

Before studying the well-definedness of spectral operators, it is worth mentioning that for the case that $\mathcal{X} \equiv \mathbb{S}^m$ (or $\mathbb{V}^{m \times n}$) if **g** has the form $\mathbf{g}(y) = (h(y_1), \ldots, h(y_m)) \in \mathbb{R}^m$ with $y_i \in \mathbb{R}$ for some given scalar valued functional $h : \mathbb{R} \to \mathbb{R}$, then the corresponding spectral operator G is called Löwner operator by Sun and Sun [52] in recognitions of Löwner's original contribution on this topic in [32] (or the Löwner non-Hermitian operator by Yang in her thesis [59] if h(0) = 0).

2.1 The well-definedness

In order to show the well-definedness of spectral operators, we need the following two simple propositions. Let $\overline{Y} \in \mathbb{S}^m$ be given. Denote $\overline{\mu}_1 > \overline{\mu}_2 > \ldots > \overline{\mu}_r$ the distinct eigenvalues of \overline{Y} . Define the index sets

$$\alpha_l := \{i \mid \lambda_i(\overline{Y}) = \overline{\mu}_l, \ 1 \le i \le m\}, \quad l = 1, \dots, r.$$

Let $\Lambda(\overline{Y})$ be the $m \times m$ diagonal matrix whose *i*-th diagonal entry is $\lambda_i(\overline{Y})$. Then, the following elementary property on the eigenvalue decomposition of \overline{Y} can be checked directly.

Proposition 2.2 The matrix $Q \in \mathbb{O}^m$ satisfies $Q\Lambda(\overline{Y}) = \Lambda(\overline{Y})Q$ if and only if there exist $Q_l \in \mathbb{Q}^{|\alpha_l|}$, $l = 1, \ldots, r$ such that Q is a block diagonal matrix whose *l*-th diagonal block is Q_l , *i.e.*,

$$Q = \operatorname{Diag}(Q_1, Q_2, \ldots, Q_r).$$

Let $\overline{Z} \in \mathbb{V}^{m \times n}$ be given. We use $\overline{\nu}_1 > \overline{\nu}_2 > \ldots > \overline{\nu}_r > 0$ to denote the nonzero distinct singular values of \overline{Z} . Define

$$a_{l} := \{i \mid \sigma_{i}(\overline{Z}) = \overline{\nu}_{l}, \ 1 \le i \le m\}, \quad l = 1, \dots, r \quad \text{and} \quad b := \{i \mid \sigma_{i}(\overline{Z}) = 0, \ 1 \le i \le m\}.$$
(7)

The following observation can be derived easily. For the real case, the proof can be found in [30, Theorem 3.7], and the corresponding result for the complex case can be obtained similarly.

Proposition 2.3 Let $\overline{\Sigma} := \Sigma(\overline{Z})$. Then, $P \in \mathbb{O}^m$ and $W \in \mathbb{O}^n$ satisfy

$$P\begin{bmatrix} \overline{\Sigma} & 0 \end{bmatrix} = \begin{bmatrix} \overline{\Sigma} & 0 \end{bmatrix} W$$

if and only if there exist $Q \in \mathbb{O}^{|a|}$, $Q' \in \mathbb{O}^{|b|}$ and $Q'' \in \mathbb{O}^{n-|a|}$ such that

$$P = \begin{bmatrix} Q & 0 \\ 0 & Q' \end{bmatrix} \quad \text{and} \quad W = \begin{bmatrix} Q & 0 \\ 0 & Q'' \end{bmatrix},$$

where $|a| = |a_1| + \ldots + |a_r|$ and $Q = \text{Diag}(Q_1, Q_2, \ldots, Q_r)$ is a block diagonal matrix whose *l*-th diagonal block is $Q_l \in \mathbb{O}^{|a_l|}$.

By combining Propositions 2.2 and 2.3 with the mixed symmetric property of \mathbf{g} , we are able to obtain the following result on the well-definedness of spectral operators.

Theorem 2.1 Let $\mathbf{g} : \mathbb{R}^{m_0+m} \to \mathbb{R}^{m_0+m}$ be mixed symmetric on κ_N . Then the spectral operator $G : \mathcal{N} \to \mathcal{X}$ defined in Definition 2.2 with respect to \mathbf{g} is well-defined.

Proof. Let $X = (X_1, \ldots, X_s) \in \mathcal{N}$ be arbitrarily chosen with $X_k \in \mathbb{S}^{m_k}$ for $1 \leq k \leq s_0$ and $X_k \in \mathbb{V}^{m_k \times n_k}$ for $s_0 + 1 \leq k \leq s$. Let $\mathbf{x} = (\mathbf{x}_1, \ldots, \mathbf{x}_s) := \kappa(X)$ with $\mathbf{x}_k \in \mathbb{R}^{m_k}$. Then we know from Proposition 2.1 that for any $i, j \in \{1, \ldots, m_k\}$,

$$(\mathbf{g}_k(\mathbf{x}))_i = (\mathbf{g}_k(\mathbf{x}))_j$$
 if $(\mathbf{x}_k)_i = (\mathbf{x}_k)_j$, $\forall 1 \le k \le s$

and

$$(\mathbf{g}_k(\mathbf{x}))_i = 0$$
 if $(\mathbf{x}_k)_i = 0$, $\forall s_0 + 1 \le k \le s$,

which, together with Propositions 2.2 and 2.3, imply that the matrix G(X) is independent of the choices of $P_k \in \mathbb{O}^{m_k}(X_k)$, $1 \leq k \leq s_0$, $(U_k, V_k) \in \mathbb{O}^{m_k, n_k}(X_k)$, $s_0 + 1 \leq k \leq s$. That is, G is well defined at X. Since X is arbitrarily chosen from \mathcal{N} , the spectral operator G is well-defined on \mathcal{N} . \Box

2.2 Differential properties of singular values and vectors

In this subsection, we collect some useful preliminary results on the singular value decomposition (SVD) of matrices. Let $\overline{Z} \in \mathbb{V}^{m \times n}$ be given. Consider the following SVD of \overline{Z} :

$$\overline{Z} = \overline{U} \begin{bmatrix} \Sigma(\overline{Z}) & 0 \end{bmatrix} \overline{V}^{\mathbb{T}} = \overline{U} \begin{bmatrix} \Sigma(\overline{Z}) & 0 \end{bmatrix} \begin{bmatrix} \overline{V}_1 & \overline{V}_2 \end{bmatrix}^{\mathbb{T}} = \overline{U} \Sigma(\overline{Z}) \overline{V}_1^{\mathbb{T}},$$
(8)

where $\overline{U} \in \mathbb{O}^m$ and $\overline{V} = \begin{bmatrix} \overline{V}_1 & \overline{V}_2 \end{bmatrix} \in \mathbb{O}^n$ with $\overline{V}_1 \in \mathbb{V}^{n \times m}$ and $\overline{V}_2 \in \mathbb{V}^{n \times (n-m)}$. Define the index sets a and c by

$$a := \{i \mid \sigma_i(\overline{Z}) > 0, \ 1 \le i \le m\} \text{ and } c := \{m+1, \dots, n\}.$$
(9)

Let the index sets a_l , l = 1, ..., r and b be defined by (7). For each $i \in \{1, ..., m\}$, we also define $l_i(\overline{Z})$ to be the number of singular values which are equal to $\sigma_i(\overline{Z})$ but are ranked before i (including i), and $\tilde{l}_i(\overline{Z})$ to be the number of singular values which are equal to $\sigma_i(\overline{Z})$ but are ranked after i (excluding i), i.e., define $l_i(\overline{Z})$ and $\tilde{l}_i(\overline{Z})$ such that

$$\sigma_{1}(\overline{Z}) \geq \ldots \geq \sigma_{i-l_{i}(\overline{Z})}(\overline{Z}) > \sigma_{i-l_{i}(\overline{Z})+1}(\overline{Z}) = \ldots = \sigma_{i}(\overline{Z}) = \ldots = \sigma_{i+\tilde{l}_{i}(\overline{Z})}(\overline{Z})$$
$$> \sigma_{i+\tilde{l}_{i}(\overline{Z})+1}(\overline{Z}) \geq \ldots \geq \sigma_{m}(\overline{Z}).$$
(10)

In later discussions, when the dependence of l_i and \tilde{l}_i on \overline{Z} are clear from the context, we often drop \overline{Z} from these notations.

For any $Y \in \mathbb{V}^{m \times n}$, let Y_{ij} be the (i, j)-th entry of Y. For any $Y \in \mathbb{V}^{m \times n}$ and the given index sets $\mathcal{I} \subseteq \{1, \ldots, m\}$ and $\mathcal{J} \subseteq \{1, \ldots, n\}$, we use $Y_{\mathcal{J}}$ to denote the sub-matrix of Y obtained by removing all the columns of Y not in \mathcal{J} and use $Y_{\mathcal{I}\mathcal{J}}$ to denote the $|\mathcal{I}| \times |\mathcal{J}|$ sub-matrix of Y obtained by removing all the rows of Y not in \mathcal{I} and all the columns of Y not in \mathcal{J} . For notational convenience, we define two linear matrix operators $S : \mathbb{V}^{p \times p} \to \mathbb{S}^p$, $T : \mathbb{V}^{p \times p} \to \mathbb{V}^{p \times p}$ by

$$S(Y) := \frac{1}{2}(Y + Y^{\mathbb{T}}) \quad T(Y) := \frac{1}{2}(Y - Y^{\mathbb{T}}), \quad Y \in \mathbb{V}^{p \times p}.$$
 (11)

The following proposition can be derived directly from the directional differentiability (e.g., see [25, Theorem 7] and [56, Proposition 1.4]) of the eigenvalues of a Hermitian matrix. For more details, see [30, Section 5.1]. **Proposition 2.4** Suppose that $\overline{Z} \in \mathbb{V}^{m \times n}$ has the SVD (8). For any $\mathbb{V}^{m \times n} \ni H \to 0$, we have

$$\sigma_i(\overline{Z} + H) - \sigma_i(\overline{Z}) - \sigma'_i(\overline{Z}; H) = O(||H||^2), \quad i = 1, \dots, m,$$
(12)

where

$$\sigma_{i}'(\overline{Z};H) = \begin{cases} \lambda_{l_{i}} \left(S(\overline{U}_{a_{l}}^{\mathbb{T}} H \overline{V}_{a_{l}}) \right) & \text{if } i \in a_{l}, \ l = 1, \dots, r ,\\ \sigma_{l_{i}} \left(\left[\overline{U}_{b}^{\mathbb{T}} H \overline{V}_{b} \quad \overline{U}_{b}^{\mathbb{T}} H \overline{V}_{2} \right] \right) & \text{if } i \in b , \end{cases}$$

$$(13)$$

where for each $i \in \{1, \ldots, m\}$, l_i is defined in (10).

The following results are also needed for subsequent discussions. For the real case, the detailed proof can be found in [14, Proposition 7]. The results for the complex case can be derived in a similar manner.

Proposition 2.5 For any $\mathbb{V}^{m \times n} \ni H \to 0$, let $Z := \begin{bmatrix} \Sigma(\overline{Z}) & 0 \end{bmatrix} + H$. Suppose that $U \in \mathbb{O}^m$ and $V = \begin{bmatrix} V_1 & V_2 \end{bmatrix} \in \mathbb{O}^n$ with $V_1 \in \mathbb{V}^{n \times m}$ and $V_2 \in \mathbb{V}^{n \times (n-m)}$ satisfy

$$Z = \begin{bmatrix} \Sigma(\overline{Z}) & 0 \end{bmatrix} + H = U \begin{bmatrix} \Sigma(Z) & 0 \end{bmatrix} V^{\mathbb{T}} = U \begin{bmatrix} \Sigma(Z) & 0 \end{bmatrix} \begin{bmatrix} V_1 & V_2 \end{bmatrix}^{\mathbb{T}}.$$

Then, there exist $Q \in \mathbb{O}^{|a|}$, $Q' \in \mathbb{O}^{|b|}$ and $Q'' \in \mathbb{O}^{n-|a|}$ such that

$$U = \begin{bmatrix} Q & 0 \\ 0 & Q' \end{bmatrix} + O(||H||) \quad \text{and} \quad V = \begin{bmatrix} Q & 0 \\ 0 & Q'' \end{bmatrix} + O(||H||), \tag{14}$$

where $Q = \text{Diag}(Q_1, Q_2, \dots, Q_r), Q_l \in \mathbb{O}^{|a_l|}$. Furthermore, we have

$$\Sigma(Z)_{a_l a_l} - \Sigma(\overline{Z})_{a_l a_l} = Q_l^{\mathbb{T}} S(H_{a_l a_l}) Q_l + O(||H||^2), \quad l = 1, \dots, r,$$
(15)

$$\left[\Sigma(Z)_{bb} - \Sigma(\overline{Z})_{bb} \quad 0\right] = Q'^{\mathbb{T}} \left[H_{bb} \quad H_{bc}\right] Q'' + O(\|H\|^2).$$
(16)

Given the index set a_l for $l \in \{1, \ldots, r\}$ at $\overline{Z} \in \mathbb{V}^{m \times n}$, we define $\mathcal{U}_l : \mathbb{V}^{m \times n} \to \mathbb{V}^{m \times n}$ by

$$\mathcal{U}_l(Z) = \sum_{i \in a_l} u_i v_i^{\mathbb{T}}, \quad Z \in \mathbb{V}^{m \times n} ,$$
(17)

where u_i and v_i are the *i*-th column of U and V, respectively, and $(U, V) \in \mathbb{O}^{m,n}(Z)$. Let $\mathcal{B} \subseteq \mathbb{V}^{m \times n}$ be an open neighborhood of \overline{Z} . By shrinking \mathcal{B} if necessary, we may assume that for any $Z \in \mathcal{B}$, if $i \in a_l$, $1 \leq l \leq r$, then $\sigma_i(Z) > 0$, and if $i \in a_l$, $j \in a_l$ and $1 \leq l \neq l' \leq r$, then $\sigma_i(Z) \neq \sigma_j(Z)$. Therefore, for any $Z \in \mathcal{B}$, we may define matrices $\Gamma_l(Z), \Xi_l(Z) \in \mathbb{R}^{m \times m}$ and $\Upsilon_l(Z) \in \mathbb{R}^{m \times (n-m)}, l = 1, \ldots, r$ by

$$(\Gamma_{l}(Z))_{ij} = \begin{cases} \frac{1}{\sigma_{i}(Z) - \sigma_{j}(Z)} & \text{if } i \in a_{l}, \ j \in a_{l'}, \ l \neq l', \ l' = 1, \dots, r+1, \\ \frac{-1}{\sigma_{i}(Z) - \sigma_{j}(Z)} & \text{if } i \in a_{l'}, \ j \in a_{l}, \ l \neq l', \ l' = 1, \dots, r+1, \\ 0 & \text{otherwise}, \end{cases}$$
(18)

$$(\Xi_{l}(Z))_{ij} = \begin{cases} \frac{1}{\sigma_{i}(Z) + \sigma_{j}(Z)} & \text{if } i \in a_{l}, \ j \in a_{l'}, \ l \neq l', \ l' = 1, \dots, r+1, \\ \frac{1}{\sigma_{i}(Z) + \sigma_{j}(Z)} & \text{if } i \in a_{l'}, \ j \in a_{l}, \ l \neq l', \ l' = 1, \dots, r+1, \\ \frac{2}{\sigma_{i}(Z) + \sigma_{j}(Z)} & \text{if } i, j \in a_{l}, \\ 0 & \text{otherwise}, \end{cases}$$
(19)

$$(\Upsilon_l(Z))_{ij} = \begin{cases} \frac{1}{\sigma_i(Z)} & \text{if } i \in a_l, \ j = 1, \dots, n - m, \\ 0 & \text{otherwise.} \end{cases}$$
(20)

We use " \circ " to denote the usual Hadamard product between two matrices, i.e., for any two matrices A and B in $\mathbb{V}^{m \times n}$ the (i, j)-th entry of $Z := A \circ B \in \mathbb{V}^{m \times n}$ is $Z_{ij} = A_{ij}B_{ij}$. We have the following differential properties of \mathcal{U}_l , $l = 1, \ldots, r$. For the real case, the results have been shown in [14, Proposition 2.11]. By using similar arguments to the real case, one can derive the corresponding results for the complex case.

Proposition 2.6 Let \mathcal{U}_l , l = 1, ..., r be defined by (17). Then, there exists an open neighborhood \mathcal{B} of \overline{Z} such that \mathcal{U}_l is at least twice continuously differentiable in \mathcal{B} , and for any $H \in \mathbb{V}^{m \times n}$, the first order derivative of \mathcal{U}_l at $Z \in \mathcal{B}$ is given by

$$\mathcal{U}_{l}'(Z)H = U[\Gamma_{l}(Z) \circ S(U^{\mathbb{T}}HV_{1}) + \Xi_{l}(Z) \circ T(U^{\mathbb{T}}HV_{1})]V_{1}^{\mathbb{T}} + U(\Upsilon_{l}(Z) \circ U^{\mathbb{T}}HV_{2})V_{2}^{\mathbb{T}},$$
(21)

where $(U, V) \in \mathbb{O}^{m,n}(Z)$ and the linear operators S and T are defined by (11).

3 Continuity, directional and Fréchet differentiability

In this and the next section, we will first focus on the study of spectral operators for the case that $\mathcal{X} \equiv \mathbb{V}^{m \times n}$. The corresponding extensions for the spectral operators defined on the general Cartesian product of several matrix spaces will be presented in Section 5.

Let \mathcal{N} be a given nonempty open set in $\mathbb{V}^{m \times n}$. Suppose that $g : \mathbb{R}^m \to \mathbb{R}^m$ is mixed symmetric, with respect to $\mathcal{P} \equiv \pm \mathbb{P}^m$ (called absolutely symmetric in this case), on an open set $\hat{\sigma}_{\mathcal{N}}$ in \mathbb{R}^m containing $\sigma_{\mathcal{N}} := \{\sigma(X) \mid X \in \mathcal{N}\}$. The spectral operator $G : \mathcal{N} \to \mathbb{V}^{m \times n}$ with respect to g defined in Definition 2.2 then takes the form of

$$G(X) = U[\operatorname{Diag}(g(\sigma(X))) \quad 0] V^{\mathbb{T}}, \quad X \in \mathcal{N},$$

where $(U, V) \in \mathbb{O}^{m,n}(X)$. Let $\overline{X} \in \mathcal{N}$ be given. Consider the SVD (8) for \overline{X} , i.e.,

$$\overline{X} = \overline{U} \begin{bmatrix} \Sigma(\overline{X}) & 0 \end{bmatrix} \overline{V}^{\mathbb{T}}, \qquad (22)$$

where $\overline{V} = \begin{bmatrix} \overline{V}_1 & \overline{V}_2 \end{bmatrix} \in \mathbb{O}^n$ with $\overline{V}_1 \in \mathbb{V}^{n \times m}$ and $\overline{V}_2 \in \mathbb{V}^{n \times (n-m)}$. Let $\overline{\sigma} := \sigma(\overline{X}) \in \mathbb{R}^m$. Let $a, b, c, a_l, l = 1, \ldots, r$ be the index sets defined by (9) and (7) with \overline{Z} being replaced by \overline{X} . Denote $\overline{a} := \{1, \ldots, n\} \setminus a$. For any given vector $y \in \mathbb{R}^m$, let $|y|^{\downarrow}$ be the vector of entries of $|y| = (|y_1|, \ldots, |y_m|)$ being arranged in the non-increasing order $|y|_1^{\downarrow} \ge \ldots \ge |y|_m^{\downarrow}$. The following result follows from the absolutely symmetric property of g on $\hat{\sigma}_{\mathcal{N}}$.

Proposition 3.1 Let $U \in \mathbb{O}^m$ and $V = \begin{bmatrix} V_1 & V_2 \end{bmatrix} \in \mathbb{O}^n$ with $V_1 \in \mathbb{V}^{n \times m}$ and $V_2 \in \mathbb{V}^{n \times (n-m)}$ be given. Let $y \in \hat{\sigma}_N$. Then, for Y := U [Diag(y) 0] $V^{\mathbb{T}}$ it always holds that

$$G(Y) = U [\operatorname{Diag}(g(y)) \quad 0] V^{\mathbb{T}} = U \operatorname{Diag}(g(y)) V_1^{\mathbb{T}}$$

Proof. Let $P \in \pm \mathbb{P}^m$ be a signed permutation matrix such that $Py = |y|^{\downarrow}$. Then, we know that $\sigma(Y) = |y|^{\downarrow}$ and Y has the following SVD

$$Y = U[P^{\mathbb{T}}\text{Diag}(|y|^{\downarrow})W \quad 0]V^{\mathbb{T}} = UP^{\mathbb{T}}\left[\text{Diag}(|y|^{\downarrow}) \quad 0\right]\left[V_{1}W^{\mathbb{T}} \quad V_{2}\right]^{\mathbb{T}},$$

where $W := |P| \in \mathbb{P}^m$ is the *m* by *m* permutation matrix whose (i, j)-th element is the absolute value of the (i, j)-th element of *P*. Then, we know from Definition 2.2 that

$$G(Y) = UP^{\mathbb{T}} \begin{bmatrix} \operatorname{Diag}(g(|y|^{\downarrow})) & 0 \end{bmatrix} \begin{bmatrix} V_1 W^{\mathbb{T}} & V_2 \end{bmatrix}^{\mathbb{T}}.$$

Since g is absolutely symmetric at y, one has

$$\operatorname{Diag}(g(|y|^{\downarrow})) = \operatorname{Diag}(g(Py)) = \operatorname{Diag}(Pg(y)) = P\operatorname{Diag}(g(y))W^{\mathbb{T}}$$

Thus,

$$G(Y) = UP^{\mathbb{T}} \begin{bmatrix} P \operatorname{Diag}(g(y)) W^{\mathbb{T}} & 0 \end{bmatrix} \begin{bmatrix} V_1 W^{\mathbb{T}} & V_2 \end{bmatrix}^{\mathbb{T}} = U \begin{bmatrix} \operatorname{Diag}(g(y)) & 0 \end{bmatrix} V^{\mathbb{T}},$$

which, proves the conclusion.

By using Proposition 2.5, we have the following result on the continuity of the spectral operator G.

Theorem 3.1 Suppose that $\overline{X} \in \mathcal{N}$ has the SVD (22). The spectral operator G is continuous at \overline{X} if and only if g is continuous at $\sigma(\overline{X})$.

Proof. " \Leftarrow " Let $X \in \mathcal{N}$. Denote $H = X - \overline{X}$ and $\sigma = \sigma(X)$. Let $U \in \mathbb{O}^m$ and $V \in \mathbb{O}^n$ be such that $X = \overline{X} + H = U \begin{bmatrix} \Sigma(X) & 0 \end{bmatrix} V^{\mathbb{T}}$. Then, we know from (22) that

$$\begin{bmatrix} \Sigma(\overline{X}) & 0 \end{bmatrix} + \overline{U}^{\mathbb{T}} H \overline{V} = \overline{U}^{\mathbb{T}} U \begin{bmatrix} \Sigma(X) & 0 \end{bmatrix} V^{\mathbb{T}} \overline{V} \,.$$

From (14) in Proposition 2.5, we know that for any X sufficiently close to \overline{X} , there exist $Q \in \mathbb{O}^{|a|}$, $Q' \in \mathbb{O}^{|b|}$ and $Q'' \in \mathbb{O}^{n-|a|}$ such that

$$\overline{U}^{\mathbb{T}}U = \begin{bmatrix} Q & 0\\ 0 & Q' \end{bmatrix} + O(||H||) \quad \text{and} \quad \overline{V}^{\mathbb{T}}V = \begin{bmatrix} Q & 0\\ 0 & Q'' \end{bmatrix} + O(||H||), \tag{23}$$

where $Q = \text{Diag}(Q_1, Q_2, \dots, Q_r), Q_l \in \mathbb{O}^{|a_l|}$. On the other hand, from the definition of the spectral operator G one has

$$U^{\mathbb{T}}(G(X) - G(\overline{X}))V = [\operatorname{Diag}(g(\sigma)) \quad 0] - U^{\mathbb{T}}\overline{U}[\operatorname{Diag}(g(\overline{\sigma})) \quad 0]\overline{V}^{\mathbb{T}}V.$$

Thus, we obtain from (23) and Proposition 2.1 that for any X sufficiently close to \overline{X} ,

$$U^{\mathbb{T}}\left(G(X) - G(\overline{X})\right)V = \left[\operatorname{Diag}(g(\sigma) - g(\overline{\sigma})) \quad 0\right] + O(\|H\|).$$

Thus, since g is assumed to be continuous at $\overline{\sigma}$, we can conclude that the spectral operator G is continuous at \overline{X} .

" \implies " Suppose that G is continuous at \overline{X} . Let $(\overline{U}, \overline{V}) \in \mathbb{O}^{m \times n}(\overline{X})$ be fixed. Choose any $\sigma \in \hat{\sigma}_{\mathcal{N}}$ and denote $X := \overline{U}[\text{Diag}(\sigma) \quad 0]\overline{V}^{\mathbb{T}}$. We know from Proposition 3.1 that $G(X) = \overline{U}\text{Diag}(g(\sigma))\overline{V}_1^{\mathbb{T}}$ and

$$\operatorname{Diag}(g(\sigma) - g(\overline{\sigma})) = \overline{U}^{\mathbb{T}} \left(G(X) - G(\overline{X}) \right) \overline{V}_{1}.$$

Hence, we know from the assumption that g is continuous at $\overline{\sigma}$.

Next, we introduce some notations which are frequently used in later discussions. For any given $X \in \mathcal{N}$, let $\sigma = \sigma(X)$. For the mapping g, we define three matrices $\mathcal{E}_1(\sigma), \mathcal{E}_2(\sigma) \in \mathbb{R}^{m \times m}$ and $\mathcal{F}(\sigma) \in \mathbb{R}^{m \times (n-m)}$ (depending on $X \in \mathcal{N}$) by

$$(\mathcal{E}_1^0(\sigma))_{ij} := \begin{cases} \frac{g_i(\sigma) - g_j(\sigma)}{\sigma_i - \sigma_j} & \text{if } \sigma_i \neq \sigma_j, \\ 0 & \text{otherwise,} \end{cases} \quad i, j \in \{1, \dots, m\},$$

$$(24)$$

$$(\mathcal{E}_2^0(\sigma)_{ij} := \begin{cases} \frac{g_i(\sigma) + g_j(\sigma)}{\sigma_i + \sigma_j} & \text{if } \sigma_i + \sigma_j \neq 0, \\ 0 & \text{otherwise}, \end{cases} \quad i, j \in \{1, \dots, m\},$$
(25)

$$(\mathcal{F}^{0}(\sigma))_{ij} := \begin{cases} \frac{g_{i}(\sigma)}{\sigma_{i}} & \text{if } \sigma_{i} \neq 0, \\ 0 & \text{otherwise,} \end{cases} \quad i \in \{1, \dots, m\}, \quad j \in \{1, \dots, n-m\}.$$

$$(26)$$

Note that when the dependence of $\mathcal{E}_1^0(\sigma)$, $\mathcal{E}_2^0(\sigma)$ and $\mathcal{F}^0(\sigma)$ on σ are clear from the context, we often drop σ from these notations. In particular, let $\overline{\mathcal{E}}_1^0$, $\overline{\mathcal{E}}_2^0 \in \mathbb{V}^{m \times m}$ and $\overline{\mathcal{F}}^0 \in \mathbb{V}^{m \times (n-m)}$ be the matrices defined by (24)-(26) with respect to $\overline{\sigma} = \sigma(\overline{X})$. Since g is absolutely symmetric at $\overline{\sigma}$, we know that for all $i \in a_l$, $1 \leq l \leq r$, the function values $g_i(\overline{\sigma})$ are the same (denoted by \overline{g}_l). Therefore, for any $X \in \mathcal{N}$, define

$$G_S(X) := \sum_{l=1}^r \bar{g}_l \mathcal{U}_l(X) \quad \text{and} \quad G_R(X) := G(X) - G_S(X) \,, \tag{27}$$

where $\mathcal{U}_l(X)$ is given by (17). The following lemma follows from Proposition 2.6 directly.

Lemma 3.1 Let $G_S : \mathcal{N} \to \mathbb{V}^{m \times n}$ be defined by (27). Then, there exists an open neighborhood \mathcal{B} of \overline{X} in \mathcal{N} such that G_S is twice continuously differentiable on \mathcal{B} , and for any $\mathbb{V}^{m \times n} \ni H \to 0$,

$$G_S(\overline{X} + H) - G_S(\overline{X}) = G'_S(\overline{X})H + O(||H||^2)$$

with

$$G'_{S}(\overline{X})H = \overline{U}\left[\overline{\mathcal{E}}_{1}^{0} \circ S(\overline{U}^{\mathbb{T}}H\overline{V}_{1}) + \overline{\mathcal{E}}_{2}^{0} \circ T(\overline{U}^{\mathbb{T}}H\overline{V}_{1}) \quad \overline{\mathcal{F}}^{0} \circ (\overline{U}^{\mathbb{T}}H\overline{V}_{2})\right]\overline{V}^{\mathbb{T}}.$$
(28)

Lemma 3.1 says that in an open neighborhood of \overline{X} , $G(\cdot)$ can be decomposed into a "smooth part" $G_S(\cdot)$ plus a "nonsmooth part" $G_R(\cdot)$. As we will see in the later developments, this decomposition simplifies many of our proofs.

3.1 Directional differentiability

Let \mathcal{Z} and \mathcal{Z}' be two finite dimensional real Euclidean spaces and \mathcal{O} be an open set in \mathcal{Z} . A function $F: \mathcal{O} \subseteq \mathcal{Z} \to \mathcal{Z}'$ is said to be *Hadamard directionally differentiable* at $z \in \mathcal{O}$ if the limit

$$\lim_{t \downarrow 0, h' \to h} \frac{F(z+th') - F(z)}{t} \quad \text{exists for any } h \in \mathcal{Z}.$$
(29)

It is clear that if F is Hadamard directionally differentiable at z, then F is directionally differentiable at z, and the limit in (29) equals the directional derivative F'(z;h) for any $h \in \mathbb{Z}$.

Assume that the g is directionally differentiable at $\overline{\sigma}$. Then, from the definition of directional derivative and the absolutely symmetry of g on the nonempty open set $\hat{\sigma}_{\mathcal{N}}$, it is easy to see that the directional derivative $\phi := g'(\overline{\sigma}; \cdot) : \mathbb{R}^m \to \mathbb{R}^m$ satisfies

$$\phi(Qh) = Q\phi(h) \quad \forall Q \in \pm \mathbb{P}^m_{\overline{\sigma}} \quad \text{and} \quad \forall h \in \mathbb{R}^m,$$
(30)

where $\pm \mathbb{P}_{\overline{\sigma}}^m$ is the subset defined with respect to $\overline{\sigma}$ by

$$\pm \mathbb{P}^m_{\overline{\sigma}} := \{ Q \in \pm \mathbb{P}^m \, | \, \overline{\sigma} = Q \overline{\sigma} \} \,. \tag{31}$$

Note that $Q \in \pm \mathbb{P}_{\overline{\sigma}}^m$ if and only if

$$Q = \operatorname{Diag}(Q_1, \dots, Q_r, Q_{r+1}) \quad \text{with} \quad Q_l \in \mathbb{P}^{|a_l|}, \quad l = 1, \dots, r \quad \text{and} \quad Q_{r+1} \in \pm \mathbb{P}^{|b|}.$$
(32)

For any $h \in \mathbb{R}^m$, we rewrite $\phi(h)$ in the following form $\phi(h) = (\phi_1(h), \ldots, \phi_r(h), \phi_{r+1}(h))$ with $\phi_l(h) \in \mathbb{R}^{|a_l|}$, $l = 1, \ldots, r$ and $\phi_{r+1}(h) \in \mathbb{R}^{|b|}$. Therefore, we know from (30) and (32) that the function $\phi : \mathbb{R}^m \to \mathbb{R}^m$ is a mixed symmetric mapping, with respect to $\mathbb{P}^{|a_1|} \times \ldots \times \mathbb{P}^{|a_r|} \times \pm \mathbb{P}^{|b|}$, over $\mathbb{R}^{|a_1|} \times \ldots \times \mathbb{R}^{|a_r|} \times \mathbb{R}^{|b|}$. Let $\mathcal{W} := \mathbb{S}^{|a_1|} \times \ldots \times \mathbb{S}^{|a_r|} \times \mathbb{V}^{|b| \times (n-|a|)}$. We can define the spectral operator $\Phi : \mathcal{W} \to \mathcal{W}$ with respect to the symmetric mapping ϕ as follows: for any $W = (W_1, \ldots, W_r, W_{r+1}) \in \mathcal{W}$,

$$\Phi(W) := \left(\Phi_1(W), \dots, \Phi_r(W), \Phi_{r+1}(W)\right)$$
(33)

with

$$\Phi_l(W) := \left\{ \begin{array}{ll} \widetilde{P}_l \mathrm{Diag}(\phi_l(\kappa(W))) \widetilde{P}_l^{\mathbb{T}} & \text{if } 1 \leq l \leq r, \\ \widetilde{M} \mathrm{Diag}(\phi_l(\kappa(W))) \widetilde{N}_1^{\mathbb{T}} & \text{if } l = r+1, \end{array} \right.$$

where $\kappa(W) := (\lambda(W_1), \ldots, \lambda(W_r), \sigma(W_{r+1})) \in \mathbb{R}^m$; $\widetilde{P}_l \in \mathbb{O}^{|a_l|}(W_l)$; and $(\widetilde{M}, \widetilde{N}) \in \mathbb{O}^{|b|, n-|a|}(W_{r+1})$, $\widetilde{N} := [\widetilde{N}_1 \quad \widetilde{N}_2]$ with $\widetilde{N}_1 \in \mathbb{V}^{(n-|a|) \times |b|}$, $\widetilde{N}_2 \in \mathbb{V}^{(n-|a|) \times (n-m)}$. From Theorem 2.1, we know that Φ is well defined on \mathcal{W} .

In order to present the directional differentiability results for the spectral operator G, we define the following first divided directional difference $g^{[1]}(\overline{X};H) \in \mathbb{V}^{m \times n}$ of g at \overline{X} along the direction $H \in \mathbb{V}^{m \times n}$ by

$$g^{[1]}(\overline{X};H) := \left[\overline{\mathcal{E}}_1^0 \circ S(\overline{U}^{\mathbb{T}}H\overline{V}_1) + \overline{\mathcal{E}}_2^0 \circ T(\overline{U}^{\mathbb{T}}H\overline{V}_1) \quad \overline{\mathcal{F}}^0 \circ \overline{U}^{\mathbb{T}}H\overline{V}_2\right] + \widehat{\Phi}(D(H)), \tag{34}$$

where $\overline{\mathcal{E}}_1, \overline{\mathcal{E}}_2, \overline{\mathcal{F}}$ are defined as in (24)–(26) at $\overline{\sigma} = \sigma(\overline{X})$,

$$D(H) := \left(S(\overline{U}_{a_1}^{\mathbb{T}} H \overline{V}_{a_1}), \dots, S(\overline{U}_{a_r}^{\mathbb{T}} H \overline{V}_{a_r}), \overline{U}_b^{\mathbb{T}} H[\overline{V}_b \quad \overline{V}_2] \right) \in \mathcal{W}$$
(35)

and for any $W = (W_1, \ldots, W_r, W_{r+1}) \in \mathcal{W}, \ \widehat{\Phi}(W) \in \mathbb{V}^{m \times n}$ is defined by

$$\widehat{\Phi}(W) := \begin{bmatrix} \operatorname{Diag}\left(\Phi_1(W), \dots, \Phi_r(W)\right) & 0\\ 0 & \Phi_{r+1}(W) \end{bmatrix}.$$
(36)

For the directional differentiability of the spectral operator, we have the following result.

Theorem 3.2 Suppose that $\overline{X} \in \mathcal{N}$ has the SVD (22). The spectral operator G is Hadamard directionally differentiable at \overline{X} if and only if g is Hadamard directionally differentiable at $\overline{\sigma} = \sigma(\overline{X})$. In that case, the directional derivative of G at \overline{X} along any direction $H \in \mathbb{V}^{m \times n}$ is given by

$$G'(\overline{X};H) = \overline{U}g^{[1]}(\overline{X};H)\overline{V}^{\mathbb{T}}.$$
(37)

Proof. " \Leftarrow " Let $H \in \mathbb{V}^{m \times n}$ be any given direction. For any $\mathbb{V}^{m \times n} \ni H' \to H$ and $\tau > 0$, denote $X := \overline{X} + \tau H'$. Consider the SVD of X, i.e.,

$$X = U[\Sigma(X) \quad 0]V^{\mathbb{T}}.$$
(38)

Denote $\sigma = \sigma(X)$. For τ and H' sufficiently close to 0 and H, let G_S and G_R be the mappings defined in (27). Then, by Lemma 3.1, we know that

$$\lim_{\tau \downarrow 0, \ H' \to H} \frac{1}{\tau} (G_S(X) - G_S(\overline{X})) = G'_S(\overline{X})H,$$
(39)

where $G'_S(\overline{X})H$ is given by (28). On the other hand, for τ and H' sufficiently close to 0 and H, we have $\mathcal{U}_l(X) = \sum_{i \in a_l} u_i v_i^{\mathbb{T}}, l = 1, \ldots, r$ and

$$G_R(X) = G(X) - G_S(X) = \sum_{l=1}^r \sum_{i \in a_l} [g_i(\sigma) - g_i(\overline{\sigma})] u_i v_i^{\mathbb{T}} + \sum_{i \in b} g_i(\sigma) u_i v_i^{\mathbb{T}}.$$
(40)

For τ and H' sufficiently close to 0 and H, let

$$\Delta_l(\tau, H') = \frac{1}{\tau} \sum_{i \in a_l} [g_i(\sigma) - g_i(\overline{\sigma})] u_i v_i^{\mathbb{T}}, \quad l = 1, \dots, r \quad \text{and} \quad \Delta_{r+1}(\tau, H') = \sum_{i \in b} g_i(\sigma) u_i v_i^{\mathbb{T}}.$$

Firstly, consider the case that $\overline{X} = [\Sigma(\overline{X}) \quad 0]$. Then, from (12) and (13), we know that for any τ and $H' \in \mathbb{V}^{m \times n}$ sufficiently close to 0 and H,

$$\sigma(X) = \sigma(\overline{X}) + \tau \sigma'(\overline{X}; H') + O(\tau^2 ||H'||^2), \qquad (41)$$

where $(\sigma'(\overline{X}; H'))_{al} = \lambda(S(H'_{a_l a_l})), l = 1, ..., r$ and $(\sigma'(\overline{X}; H'))_b = \sigma([H'_{bb} H'_{bc}])$. Denote $h' := \sigma'(\overline{X}; H')$ and $h := \sigma'(\overline{X}; H)$. By using the fact that the singular value functions of a general matrix are globally Lipschitz continuous, we know that

$$\lim_{\tau \downarrow 0, \ H' \to H} \left(h' + O(\tau \| H' \|^2) \right) = h \,. \tag{42}$$

Since g is assumed to be Hadamard directionally differentiable at $\overline{\sigma}$, we have

$$\lim_{\tau \downarrow 0, \ H' \to H} \frac{g(\sigma) - g(\overline{\sigma})}{\tau} = \lim_{\tau \downarrow 0, \ H' \to H} \frac{1}{\tau} [g(\overline{\sigma} + \tau(h' + O(\tau ||H'||^2))) - g(\overline{\sigma})] = g'(\overline{\sigma}; h) = \phi(h),$$

where $\phi \equiv g'(\overline{\sigma}; \cdot) : \mathbb{R}^m \to \mathbb{R}^m$ satisfies the condition (30). Since $u_i v_i^{\mathbb{T}}$, $i = 1, \ldots, m$ are uniformly bounded, we know that for τ and H' sufficiently close to 0 and H,

$$\Delta_l(\tau, H') = U_{a_l} \operatorname{Diag}(\phi_l(h)) V_{a_l}^{\mathbb{T}} + o(1) \quad l = 1, \dots, r ,$$

$$\Delta_{r+1}(\tau, H') = U_b \operatorname{Diag}(\phi_{r+1}(h)) V_b^{\mathbb{T}} + o(1) .$$

By (14) in Proposition 2.5, we know that there exist $Q_l \in \mathbb{O}^{|a_l|}$, $l = 1, \ldots, r, M \in \mathbb{O}^{|b|}$ and $N = \begin{bmatrix} N_1 & N_2 \end{bmatrix} \in \mathbb{O}^{|a_l|}$ with $N_1 \in \mathbb{V}^{(n-|a|) \times |b|}$ and $N_2 \in \mathbb{V}^{(n-|a|) \times (n-m)}$ (depending on τ and H') such that

$$U_{a_{l}} = \begin{bmatrix} O(\tau ||H'||) \\ Q_{l} + O(\tau ||H'||) \\ O(\tau ||H'||) \end{bmatrix}, \quad V_{a_{l}} = \begin{bmatrix} O(\tau ||H'||) \\ Q_{l} + O(\tau ||H'||) \\ O(\tau ||H'||) \end{bmatrix} l = 1, \dots, r,$$
$$U_{b} = \begin{bmatrix} O(\tau ||H'||) \\ M + O(\tau ||H'||) \end{bmatrix}, \quad [V_{b} \quad V_{c}] = \begin{bmatrix} O(\tau ||H'||) \\ N + O(\tau ||H'||) \end{bmatrix}.$$

Thus, we have

$$\Delta_{l}(\tau, H') = \begin{bmatrix} 0 & 0 & 0 \\ 0 & Q_{l} \operatorname{Diag}(\phi_{l}(h)) Q_{l}^{\mathbb{T}} & 0 \\ 0 & 0 & 0 \end{bmatrix} + O(\tau \|H'\|) + o(1), \quad l = 1, \dots, r,$$
(43)

$$\Delta_{r+1}(\tau, H') = \begin{bmatrix} 0 & 0\\ 0 & M \text{Diag}(\phi_{r+1}(h)) N_1^{\mathbb{T}} \end{bmatrix} + O(\tau \| H' \|) + o(1) \,.$$
(44)

We know from (15) and (16) that

$$S(H'_{a_l a_l}) = S(H_{a_l a_l}) + o(1) = \frac{1}{\tau} Q_l[\Sigma(X)_{a_l a_l} - \overline{\nu}_l I_{|a_l|}] Q_l^{\mathbb{T}} + O(\tau ||H'||^2), \quad l = 1, \dots, r,$$
(45)

$$[H'_{bb} \quad H'_{bc}] = [H_{bb} \quad H_{bc}] + o(1) = \frac{1}{\tau} M[\Sigma(X)_{bb} - \overline{\nu}_{r+1} I_{|b|}] N_1^{\mathbb{T}} + O(\tau ||H'||^2).$$
(46)

Since Q_l , l = 1, ..., r, M and N are uniformly bounded, by taking subsequences if necessary, we may assume that when $\tau \downarrow 0$ and $H' \to H$, Q_l , M and N converge to \widetilde{Q}_l , \widetilde{M} and \widetilde{N} , respectively. Therefore, by taking limits in (45) and (46), we obtain from (41) and (42) that

$$S(H_{a_l a_l}) = \widetilde{Q}_l \Lambda(S(H_{a_l a_l})) \widetilde{Q}_l^{\mathbb{T}}, \quad l = 1, \dots, r,$$

$$[H_{bb} \quad H_{bc}] = \widetilde{M} [\Sigma([H_{bb} \quad H_{bc}]) \quad 0] \widetilde{N}^{\mathbb{T}} = \widetilde{M} \Sigma([H_{bb} \quad H_{bc}]) \widetilde{N}_1^{\mathbb{T}}.$$

Hence, by using the notation (33), we know from (40), (43), (44) and (36) that

$$\lim_{\downarrow 0, H' \to H} \frac{1}{\tau} G_R(X) = \lim_{\tau \downarrow 0, H' \to H} \sum_{l=1}^{r+1} \Delta_l(\tau, H') = \widehat{\Phi}(D(H)), \qquad (47)$$

where $D(H) = (S(H_{a_1a_1}), \dots, S(H_{a_ra_r}), H_{b\bar{a}}).$

au

To prove the conclusion for the general case of \overline{X} , rewrite (38) as

$$\begin{bmatrix} \Sigma(\overline{X}) & 0 \end{bmatrix} + \overline{U}^{\mathbb{T}} H' \overline{V} = \overline{U}^{\mathbb{T}} U[\Sigma(X) & 0] V^{\mathbb{T}} \overline{V}$$

Let $\widetilde{U} := \overline{U}^{\mathbb{T}}U$, $\widetilde{V} := \overline{V}^{\mathbb{T}}V$ and $\widetilde{H} = \overline{U}^{\mathbb{T}}H\overline{V}$. Denote $\widetilde{X} := [\Sigma(\overline{X}) \quad 0] + \overline{U}^{\mathbb{T}}H'\overline{V}$. Then, we obtain that $G_R(X) = \overline{U}G_R(\widetilde{X})\overline{V}^{\mathbb{T}}$. Thus, we know from (47) that

$$\lim_{\tau \downarrow 0, \ H' \to H} \frac{1}{\tau} G_R(X) = \overline{U} \widehat{\Phi}(D(\widetilde{H})) \overline{V}^{\mathbb{T}} .$$
(48)

Therefore, by combining (39) and (48) and noting that $G(\overline{X}) = G_S(\overline{X})$, we obtain that for any given $H \in \mathbb{V}^{m \times n}$,

$$\lim_{\downarrow 0, \ H' \to H} \frac{G(X) - G(\overline{X})}{\tau} = \lim_{\tau \downarrow 0, \ H' \to H} \frac{G_S(X) - G_S(\overline{X}) + G_R(X)}{\tau} = \overline{U}g^{[1]}(\overline{X}; \widetilde{H})\overline{V}^{\mathbb{T}}$$

where $g^{[1]}(\overline{X}; \widetilde{H})$ is given by (34). This implies that G is Hadamard directionally differentiable at \overline{X} and (37) holds.

" \Longrightarrow " Suppose that G is Hadamard directionally differentiable at \overline{X} . Let $(\overline{U}, \overline{V}) \in \mathbb{O}^{m \times n}(\overline{X})$ be fixed. For any given direction $h \in \mathbb{R}^m$, suppose that $\mathbb{R}^m \ni h' \to h$. Denote $H' := \overline{U}[\operatorname{Diag}(h') \quad 0]\overline{V}^{\mathbb{T}} \in \mathbb{V}^{m \times n}$ and $H := \overline{U}[\operatorname{Diag}(h) \quad 0]\overline{V}^{\mathbb{T}} \in \mathbb{V}^{m \times n}$. Then, we have $H' \to H$ as $h' \to h$. Since for all $\tau > 0$ and h' sufficiently close to 0 and $h, \sigma := \overline{\sigma} + \tau h' \in \hat{\sigma}_{\mathcal{N}}$, we know from Proposition 3.1 that for all $\tau > 0$ and h' sufficiently close to 0 and $h, G(\overline{X} + \tau H') = \overline{U}\operatorname{Diag}(g(\overline{\sigma} + \tau h'))\overline{V}_1^{\mathbb{T}}$. This implies that

$$\operatorname{Diag}\left(\lim_{\tau \downarrow 0, \ h' \to h} \frac{g(\overline{\sigma} + \tau h') - g(\overline{\sigma})}{\tau}\right) = \overline{U}^{\mathbb{T}}\left(\lim_{\tau \downarrow 0, \ H' \to H} \frac{G(\overline{X} + \tau H') - G(\overline{X})}{\tau}\right) \overline{V}_{1}.$$

Thus, we know from the assumption that $\lim_{\tau \downarrow 0, h' \to h} \frac{g(\overline{\sigma} + \tau h') - g(\overline{\sigma})}{\tau}$ exists and that g is Hadamard directionally differentiable at $\overline{\sigma}$.

Remark 1 Note that for a general spectral operator G, we cannot obtain the directional differentiability at \overline{X} if we only assume that g is directionally differentiable at $\sigma(\overline{X})$. In fact, a counterexample can be found in [27]. However, since $\mathbb{V}^{m \times n}$ is a finite dimensional Euclidean space, it is well-known that for locally Lipschitz continuous functions, the directional differentiability in the sense of Hadamard and Gâteaux are equivalent (see e.g. [37, Theorem 1.13], [13, Lemma 3.2], [19, p.259]). Therefore, if G and g are locally Lipschitz continuous near \overline{X} and $\sigma(\overline{X})$, respectively (e.g., the proximal mapping P_f and its vector counterpart P_{θ}), then G is directionally differentiable at \overline{X} if and only if g is directionally differentiable at $\sigma(\overline{X})$.

3.2 Fréchet differentiability

For a given $X \in \mathcal{N}$, suppose that the given absolutely symmetric mapping g is F(réchet)-differentiable at $\sigma = \sigma(X)$. The following results on the Jacobian matrix $g'(\sigma)$ can be obtained directly from the assumed absolute symmetry of g on $\hat{\sigma}_{\mathcal{N}}$ and the block structure (32) for any $Q \in \pm \mathbb{P}_{\sigma}^{m}$.

Lemma 3.2 For any $X \in \mathcal{N}$, suppose that g is F-differentiable at $\sigma = \sigma(X)$. Then, the Jacobian matrix $q'(\sigma)$ has the following property

$$g'(\sigma) = Q^{\mathbb{T}}g'(\sigma)Q \quad \forall Q \in \pm \mathbb{P}_{\sigma}^{m}.$$

In particular,

$$\begin{cases} (g'(\sigma))_{ii} = (g'(\sigma))_{i'i'} & \text{if } \sigma_i = \sigma_{i'} \text{ and } i, i' \in \{1, \dots, m\}, \\ (g'(\sigma))_{ij} = (g'(\sigma))_{i'j'} & \text{if } \sigma_i = \sigma_{i'}, \ \sigma_j = \sigma_{j'}, \ i \neq j, \ i' \neq j' \text{ and } i, i', j, j' \in \{1, \dots, m\}, \\ (g'(\sigma))_{ij} = (g'(\sigma))_{ji} = 0 & \text{if } \sigma_i = 0, \ i \neq j \text{ and } i, j \in \{1, \dots, m\}. \end{cases}$$

Lemma 3.2 is a simple extension of [28, Lemma 2.1] for symmetric mappings. But one should note that the Jacobian matrix $q'(\sigma)$ of q at the F-differentiable point σ may not be symmetric since here q is not assumed to be the gradient mapping as in [28, Lemma 2.1]. For example, let us consider the absolutely symmetric mapping q defined by (4) in the introduction. Then q is differentiable at x = (2, 1) by taking m=2 and $\tau=\varepsilon=1$ in (5). However, it is easy to see that the Jacobian matrix g'(x) is not symmetric.

Let $\eta(\sigma) \in \mathbb{R}^m$ be the vector defined by

$$(\eta(\sigma))_i := \begin{cases} (g'(\sigma))_{ii} - (g'(\sigma))_{i(i+1)} & \text{if } \exists j \in \{1, \dots, m\} \text{ and } j \neq i \text{ such that } \sigma_i = \sigma_j, \\ (g'(\sigma))_{ii} & \text{otherwise}, \end{cases} \quad i \in \{1, \dots, m\}.$$

$$(49)$$

Define the corresponding divided difference matrix $\mathcal{E}_1(\sigma) \in \mathbb{R}^{m \times m}$, the divided addition matrix $\mathcal{E}_2(\sigma) \in \mathbb{R}^{m \times m}$ $\mathbb{R}^{m \times m}$, the division matrix $\mathcal{F}(\sigma) \in \mathbb{R}^{m \times (n-m)}$, respectively, by

$$(\mathcal{E}_1(\sigma))_{ij} := \begin{cases} \frac{g_i(\sigma) - g_j(\sigma)}{\sigma_i - \sigma_j} & \text{if } \sigma_i \neq \sigma_j, \\ (\eta(\sigma))_i & \text{otherwise}, \end{cases} \quad i, j \in \{1, \dots, m\},$$

$$(50)$$

$$(\mathcal{E}_2(\sigma))_{ij} := \begin{cases} \frac{g_i(\sigma) + g_j(\sigma)}{\sigma_i + \sigma_j} & \text{if } \sigma_i + \sigma_j \neq 0, \\ g'(\sigma))_{ii} & \text{otherwise}, \end{cases} \quad i, j \in \{1, \dots, m\},$$
(51)

$$(\mathcal{F}(\sigma))_{ij} := \begin{cases} \frac{g_i(\sigma)}{\sigma_i} & \text{if } \sigma_i \neq 0, \\ (g'(\sigma))_{ii} & \text{otherwise,} \end{cases} \quad i \in \{1, \dots, m\}, \quad j \in \{1, \dots, n-m\}.$$

$$(52)$$

Define the matrix $\mathcal{C}(\sigma) \in \mathbb{R}^{m \times m}$ to be the difference between $q'(\sigma)$ and $\text{Diag}(\eta(\sigma))$, i.e.,

$$\mathcal{C}(\sigma) := g'(\sigma) - \operatorname{Diag}(\eta(\sigma)).$$
(53)

Note that when the dependence of η , \mathcal{E}_1 , \mathcal{E}_2 , \mathcal{F} and \mathcal{C} on σ is clear from the context, we often drop σ from the corresponding notations.

Let $\overline{X} \in \mathcal{N}$ be given and denote $\overline{\sigma} = \sigma(\overline{X})$. Denote $\overline{\eta} = \eta(\overline{\sigma}) \in \mathbb{R}^m$ to be the vector defined by (49). Let $\overline{\mathcal{E}}_1, \overline{\mathcal{E}}_2, \overline{\mathcal{F}}$ and $\overline{\mathcal{C}}$ be the real matrices defined in (50)–(53) with respect to $\overline{\sigma}$. Now, we are ready to state the result on the F-differentiability of spectral operators.

Theorem 3.3 Suppose that the given matrix $\overline{X} \in \mathcal{N}$ has the SVD (22). Then the spectral operator G is F-differentiable at \overline{X} if and only if q is F-differentiable at $\overline{\sigma}$. In that case, the derivative of G at \overline{X} is given by

$$G'(\overline{X})H = \overline{U}[\overline{\mathcal{E}}_1 \circ S(A) + \operatorname{Diag}\left(\overline{\mathcal{C}}\operatorname{diag}(S(A))\right) + \overline{\mathcal{E}}_2 \circ T(A) \quad \overline{\mathcal{F}} \circ B]\overline{V}^{\mathbb{T}} \quad \forall \ H \in \mathbb{V}^{m \times n},$$
(54)

where $A := \overline{U}^{\mathbb{T}} H \overline{V}_1$ and $B := \overline{U}^{\mathbb{T}} H \overline{V}_2$.

Proof. " \Leftarrow " For any $\mathbb{V}^{m \times n} \ni H = [H_1 \ H_2] \to 0$ with $H_1 \in \mathbb{V}^{m \times m}$ and $H_2 \in \mathbb{V}^{m \times (n-m)}$, denote $X = \overline{X} + H$. Let $U \in \mathbb{O}^m$ and $V \in \mathbb{O}^n$ be such that

$$X = U[\Sigma(X) \quad 0]V^{\mathbb{T}}.$$
(55)

Denote $\sigma = \sigma(X)$. Let $G_S(X)$ and $G_R(X)$ be defined by (27). Then, by Lemma 3.1, we know that for any $H \to 0$,

$$G_{S}(X) - G_{S}(\overline{X}) = G'_{S}(\overline{X})H + O(||H||^{2}) = G'_{S}(\overline{X})H + O(||H||^{2}),$$
(56)

where $G'_S(\overline{X})H$ is given by (28). For $H \in \mathbb{V}^{m \times n}$ sufficiently small, we have $\mathcal{U}_l(X) = \sum_{i \in a_l} u_i v_i^{\mathbb{T}}, l = 1, \ldots, r$. Therefore,

$$G_R(X) = G(X) - G_S(X) = \sum_{l=1}^{r+1} \Delta_l(H),$$
(57)

where $\Delta_l(H) = \sum_{i \in a_l} (g_i(\sigma) - g_i(\overline{\sigma})) u_i v_i^{\mathbb{T}}, \ l = 1, \dots, r \text{ and } \Delta_{r+1}(H) = \sum_{i \in b} g_i(\sigma) u_i v_i^{\mathbb{T}}.$

Firstly, consider the case that $\overline{X} = [\Sigma(\overline{X}) \quad 0]$. Then, from (12) and (13), for any $H \in \mathbb{V}^{m \times n}$ sufficiently small, we have

$$\sigma = \overline{\sigma} + h + O(\|H\|^2), \qquad (58)$$

where $h := \sigma'(\overline{X}; H) \in \mathbb{R}^m$ with

$$(\sigma'(\overline{X};H))_{a_l} = \lambda(S(H_{a_l a_l})), \ l = 1, \dots, r \quad \text{and} \quad (\sigma'(\overline{X};H))_b = \sigma([H_{bb} \quad H_{bc}]).$$
(59)

Since g is F-differentiable at $\overline{\sigma}$ and the singular value functions are globally Lipschitz continuous, we know from (58) that for any $H \in \mathbb{V}^{m \times n}$ sufficiently small,

$$g(\sigma) - g(\overline{\sigma}) = g(\overline{\sigma} + h + O(||H||^2)) - g(\overline{\sigma}) = g'(\overline{\sigma})(h + O(||H||^2)) + o(||h||) = g'(\overline{\sigma})h + o(||H||).$$

Since $u_i v_i^{\mathbb{T}}$, i = 1, ..., m are uniformly bounded, we have for H sufficiently small,

$$\Delta_l(H) = U_{a_l} \operatorname{Diag}((g'(\overline{\sigma})h)_{a_l}) V_{a_l}^{\mathbb{T}} + o(||H||), \ l = 1, \dots, r, \quad \Delta_{r+1}(H) = U_b \operatorname{Diag}((g'(\overline{\sigma})h)_b) V_b^{\mathbb{T}} + o(||H||).$$

By (14) in Proposition 2.5, we know that there exist $Q_l \in \mathbb{O}^{|a_l|}$, $M \in \mathbb{O}^{|b|}$ and $N = [N_1 \quad N_2] \in \mathbb{O}^{n-|a|}$ with $N_1 \in \mathbb{V}^{(n-|a|) \times |b|}$ and $N_2 \in \mathbb{V}^{(n-|a|) \times (n-m)}$ (depending on H) such that

$$U_{a_{l}} = \begin{bmatrix} O(\|H\|) \\ Q_{l} + O(\|H\|) \\ O(\|H\|) \end{bmatrix}, \quad V_{a_{l}} = \begin{bmatrix} O(\|H\|) \\ Q_{l} + O(\|H\|) \\ O(\|H\|) \end{bmatrix}, \quad l = 1, \dots, r,$$
$$U_{b} = \begin{bmatrix} O(\|H\|) \\ M + O(\|H\|) \end{bmatrix}, \quad [V_{b} \quad V_{c}] = \begin{bmatrix} O(\|H\|) \\ N + O(\|H\|) \end{bmatrix}.$$

Therefore, since $||g'(\overline{\sigma})h|| = O(||H||)$, we obtain that

$$\Delta_{l}(H) = \begin{bmatrix} 0 & 0 & 0\\ 0 & Q_{l} \operatorname{Diag}((g'(\overline{\sigma})h)_{a_{l}})Q_{l}^{\mathbb{T}} & 0\\ 0 & 0 & 0 \end{bmatrix} + o(||H||), \quad l = 1, \dots, r,$$
(60)

$$\Delta_{r+1}(H) = \begin{bmatrix} 0 & 0\\ 0 & M \operatorname{Diag}((g'(\overline{\sigma})h)_b)N_1^{\mathbb{T}} \end{bmatrix} + o(||H||).$$
(61)

We know from (49) and Lemma 3.2 that $\overline{\eta}_{a_l} = \overline{\gamma}_l e^{|a_l|}$ for some $\overline{\gamma}_l \in \mathbb{R}$, $l = 1, \ldots, r$ and $\overline{\eta}_{r+1} = \overline{\gamma}_{r+1} e^{|b|}$ for some $\overline{\gamma}_{r+1} \in \mathbb{R}$, where e^p is the vector of all ones in \mathbb{R}^p and $\overline{\mathcal{C}} = g'(\overline{\sigma}) - \text{Diag}(\overline{\eta}) \in \mathbb{R}^{m \times m}$ has the following form

$$\overline{\mathcal{C}} = \begin{bmatrix} c_{11}E^{|a_1||a_1|} & \cdots & c_{1r}E^{|a_1||a_r|} & 0\\ \vdots & \ddots & \vdots & \vdots\\ \overline{c}_{r1}E^{|a_r||a_1|} & \cdots & \overline{c}_{rr}E^{|a_r||a_r|} & 0\\ 0 & \cdots & 0 & 0 \end{bmatrix},$$
(62)

where $E^{pq} \in \mathbb{R}^{p \times q}$ is the p by q matrix of all ones and $\bar{c} \in \mathbb{R}^{r \times r}$. Then we know from (59) that

$$(g'(\overline{\sigma})h)_{a_l} = \begin{cases} (\overline{\mathcal{C}}h)_{a_l} + \overline{\gamma}_l \lambda(S(H_{a_l a_l})) & \text{if } l = 1, \dots, r, \\ \overline{\gamma}_{r+1} \sigma([H_{bb} \quad H_{bc}]) & \text{if } l = r+1, \end{cases}$$

where for $l \in \{1, \ldots, r\}$, $(\overline{\mathcal{C}}h)_{a_l} = \sum_{l'=1}^r \overline{c}_{ll'} \operatorname{tr}(S(H_{a_{l'}a_{l'}})) e^{|a_l|} = (\overline{\mathcal{C}}\operatorname{diag}(S(H_1)))_{a_l}$. On the other hand, we know from (15), (16), (58) and (59) that for H sufficiently close to 0, and $l = 1, \ldots, r$,

$$S(H_{a_{l}a_{l}}) = Q_{l}(\Sigma(X)_{a_{l}a_{l}} - \overline{\nu}_{l}I_{|a_{l}|})Q_{l}^{\mathbb{T}} + O(||H||^{2}) = Q_{l}\Lambda(S(H_{a_{l}a_{l}}))Q_{l}^{\mathbb{T}} + O(||H||^{2}),$$

$$[H_{bb} \quad H_{bc}] = M(\Sigma(X)_{bb} - \overline{\nu}_{r+1}I_{|b|})N_{1}^{\mathbb{T}} + O(||H||^{2}) = M\Sigma([H_{bb} \quad H_{bc}])N_{1} + O(||H||^{2}).$$

Therefore, from (59), (60) and (61), we obtain that

$$\Delta_{l}(H) = \begin{bmatrix} 0 & 0 & 0 \\ 0 & \text{Diag}\left((\overline{\mathcal{C}}\text{diag}(S(H_{1})))_{a_{l}}\right) + \bar{\gamma}_{l}S(H_{a_{l}a_{l}}) & 0 \\ 0 & 0 & 0 \end{bmatrix} + o(||H||), \quad l = 1, \dots, r,$$

$$\Delta_{r+1}(H) = \begin{bmatrix} 0 & 0 & 0 \\ 0 & \overline{\gamma}_{r+1}H_{bb} & \overline{\gamma}_{r+1}H_{bc} \end{bmatrix} + o(||H||).$$

Thus, we know from (57) that for any H sufficiently close to 0,

$$G_{R}(X) = \left[\text{Diag}\left(\overline{\mathcal{C}}\text{diag}(S(H_{1}))\right) \quad 0\right] + \left[\begin{array}{cccc} \overline{\gamma}_{1}S(H_{a_{1}a_{1}}) & 0 & 0 & 0 & 0\\ 0 & \ddots & 0 & 0 & 0\\ 0 & 0 & \overline{\gamma}_{r}S(H_{a_{r}a_{r}}) & 0 & 0\\ 0 & 0 & 0 & \overline{\gamma}_{r+1}H_{bb} & \overline{\gamma}_{r+1}H_{bc} \end{array}\right] + o(||H||) \,.$$
(63)

Next, consider the general $\overline{X} \in \mathbb{V}^{m \times n}$. For any $H \in \mathbb{V}^{m \times n}$ sufficiently close to 0, rewrite (55) as

$$[\Sigma(\overline{X}) \quad 0] + \overline{U}^{\mathbb{T}} H \overline{V} = \overline{U}^{\mathbb{T}} U[\Sigma(X) \quad 0] V^{\mathbb{T}} \overline{V} \,.$$

Denote $\widetilde{U} := \overline{U}^{\mathbb{T}}U$ and $\widetilde{V} := \overline{V}^{\mathbb{T}}V$. Let $\widetilde{X} := [\Sigma(\overline{X}) \quad 0] + \overline{U}^{\mathbb{T}}H\overline{V}$. Then, since \overline{U} and \overline{V} are unitary matrices, we know from (63) that

$$G_R(X) = \overline{U}G_R(\widetilde{X})\overline{V}^{\mathbb{T}} = \overline{U}\left[\text{Diag}\left(\overline{\mathcal{C}}\text{diag}(S(A))\right) \quad 0\right]\overline{V}^{\mathbb{T}} + \overline{U}\left[\begin{array}{cccc} \overline{\gamma}_1 S(A_{a_1a_1}) & 0 & 0 & 0 & 0\\ 0 & \ddots & 0 & 0 & 0\\ 0 & 0 & \overline{\gamma}_r S(A_{a_ra_r}) & 0 & 0\\ 0 & 0 & 0 & \overline{\gamma}_{r+1}A_{bb} & \overline{\gamma}_{r+1}B_{bc} \end{array}\right]\overline{V}^{\mathbb{T}} + o(||H||).$$
(64)

Thus, by combining (56) and (64) with (28) and noting that $G(\overline{X}) = G_S(\overline{X})$, we obtain that for any $H \in \mathbb{V}^{m \times n}$ sufficiently small,

$$G(X) - G(\overline{X}) = \overline{U}[\overline{\mathcal{E}}_1 \circ S(A) + \operatorname{Diag}\left(\overline{\mathcal{C}}\operatorname{diag}(S(A))\right) + \overline{\mathcal{E}}_2 \circ T(A) \quad \overline{\mathcal{F}} \circ B]\overline{V}^{\mathbb{T}} + o(||H||).$$

Therefore, we know that G is F-differentiable at \overline{X} and (54) holds.

" \implies " Suppose that G is F-differentiable at \overline{X} . Let $(\overline{U}, \overline{V}) \in \mathbb{O}^{m \times n}(\overline{X})$ be fixed. For any $h \in \mathbb{R}^m$, let $H = \overline{U}[\operatorname{Diag}(h) \quad 0]\overline{V}^{\mathbb{T}} \in \mathbb{V}^{m \times n}$. We know from Proposition 3.1 that for all h sufficiently close to 0, $G(\overline{X} + H) = \overline{U}\operatorname{Diag}(g(\overline{\sigma} + h))\overline{V}_1^{\mathbb{T}}$. Therefore, we know from the assumption that for all h sufficiently close to 0,

$$\operatorname{Diag}(g(\overline{\sigma}+h)-g(\overline{\sigma})) = \overline{U}^{\mathbb{T}}\left(G(\overline{X}+H)-G(\overline{X})\right)\overline{V}_{1} = \overline{U}^{\mathbb{T}}G'(\overline{X})H\overline{V}_{1} + o(\|h\|).$$

Hence, we know that g is F-differentiable at $\overline{\sigma}$ and $\text{Diag}(g'(\sigma)h) = \overline{U}^{\mathbb{T}}G'(\overline{X})H\overline{V}_1$. The proof is competed.

Finally, we can present the continuous differentiability result of spectral operators in the following theorem.

Theorem 3.4 Suppose that $\overline{X} \in \mathcal{N}$ has the SVD (22). Then, G is continuously differentiable at \overline{X} if and only if g is continuously differentiable at $\overline{\sigma} = \sigma(\overline{X})$.

Proof. " \Leftarrow " By the assumption, we know from Theorem 3.3 that there exists an open neighborhood $\mathcal{B} \subseteq \mathcal{N}$ of \overline{X} such that the spectral operator G is differentiable on \mathcal{B} , and for any $X \in \mathcal{B}$, the derivative G'(X) is given by

$$G'(X)H = U[\mathcal{E}_1 \circ S(A) + \operatorname{Diag}\left(\mathcal{C}\operatorname{diag}(S(A))\right) + \mathcal{E}_2 \circ T(A) \quad \mathcal{F} \circ B]V^{\mathbb{T}} \quad \forall H \in \mathbb{V}^{m \times n},$$
(65)

where $(U, V) \in \mathbb{O}^{m,n}(X)$, $A = U^{\mathbb{T}}HV_1$, $B = U^{\mathbb{T}}HV_2$ and η , \mathcal{E}_1 , \mathcal{E}_2 , \mathcal{F} and \mathcal{C} are defined by (49)–(53) with respect to $\sigma = \sigma(X)$, respectively. Next, we shall prove that

$$\lim_{X \to \overline{X}} G'(X)H \to G'(\overline{X})H \quad \forall H \in \mathbb{V}^{m \times n} .$$
(66)

Firstly, we will show that (66) holds for the special case that $\overline{X} = [\Sigma(\overline{X}) \quad 0]$ and $X = [\Sigma(X) \quad 0] \to \overline{X}$. Let $\{F^{(ij)}\}$ be the standard basis of $\mathbb{V}^{m \times n}$, i.e., for each $i \in \{1, \ldots, m\}$ and $j \in \{1, \ldots, n\}$, $F^{(ij)} \in \mathbb{V}^{m \times n}$ is a matrix whose entries are zeros, except the (i, j)-th entry is 1 or $\sqrt{-1}$. Therefore, we only need to show (66) holds for all $F^{(ij)}$. Note that since $\sigma(\cdot)$ is globally Lipschitz continuous, we know that for X sufficiently close to \overline{X} , $\sigma_i \neq \sigma_j$ if $\overline{\sigma}_i \neq \overline{\sigma}_j$.

For each $i \in \{1, \ldots, m\}$ and $j \in \{1, \ldots, n\}$, write $F^{(ij)}$ in the following form

$$F^{(ij)} = \begin{bmatrix} F_1^{(ij)} & F_2^{(ij)} \end{bmatrix}$$

with $F_1^{(ij)} \in \mathbb{V}^{m \times m}$ and $F_2^{(ij)} \in \mathbb{V}^{m \times (n-m)}$. Next, we consider the following cases.

Case 1: $i, j \in \{1, ..., m\}$ and i = j. In this case, since g' is continuous at $\overline{\sigma}$, we know that if $F^{(ij)}$ is real, then

$$\lim_{X \to \overline{X}} G'(X) F^{(ij)} = \lim_{X \to \overline{X}} [\operatorname{Diag}(g'(\sigma)e_i) \quad 0] = [\operatorname{Diag}(g'(\overline{\sigma})e_i) \quad 0] = G'(\overline{X}) F^{(ij)}$$

where e_i is the vector whose *i*-th entry is one, and zero otherwise; if $F^{(ij)}$ is complex, then

$$\lim_{X \to \overline{X}} G'(X) F^{(ij)} = \lim_{X \to \overline{X}} \left[\frac{g_i(\sigma) + g_j(\sigma)}{\sigma_i + \sigma_j} T(F_1^{(ij)}) \quad 0 \right] = \left[\frac{g_i(\overline{\sigma}) + g_j(\overline{\sigma})}{\overline{\sigma_i} + \overline{\sigma_j}} T(F_1^{(ij)}) \quad 0 \right] = G'(\overline{X}) F^{(ij)} .$$

Case 2: $i, j \in \{1, ..., m\}, i \neq j, \sigma_i = \sigma_j \text{ and } \overline{\sigma}_i = \overline{\sigma}_j > 0$. Therefore, we know that there exists $l \in \{1, ..., r\}$ such that $i, j \in a_l$. Since g' is continuous at $\overline{\sigma}$, we know from (49) that

$$\lim_{X \to \overline{X}} G'(X) F^{(ij)} = \lim_{X \to \overline{X}} \left[((g'(\sigma))_{ii} - (g'(\sigma))_{ij}) S(F_1^{(ij)}) + \frac{g_i(\sigma) + g_j(\sigma)}{\sigma_i(X) + \sigma_j(X)} T(F_1^{(ij)}) \quad 0 \right]$$
$$= \left[((g'(\overline{\sigma}))_{ii} - (g'(\overline{\sigma}))_{ij}) S(F_1^{(ij)}) + \frac{g_i(\overline{\sigma}) + g_j(\overline{\sigma})}{\overline{\sigma}_i + \overline{\sigma}_j} T(F_1^{(ij)}) \quad 0 \right]$$
$$= G'(\overline{X}) F^{(ij)}.$$

Case 3: $i, j \in \{1, \ldots, m\}, i \neq j, \sigma_i \neq \sigma_j \text{ and } \overline{\sigma}_i = \overline{\sigma}_j > 0$. In this case, we know that

$$G'(X)F^{(ij)} = \begin{bmatrix} \underline{g_i(\sigma) - g_j(\sigma)} \\ \overline{\sigma_i - \sigma_j}S(F_1^{(ij)}) + \frac{g_i(\sigma) + g_j(\sigma)}{\sigma_i + \sigma_j}T(F_1^{(ij)}) & 0 \end{bmatrix}.$$

Let $s, t \in \mathbb{R}^m$ be two vectors defined by

$$s_p := \begin{cases} \sigma_p & \text{if } p \neq i, \\ \sigma_j & \text{if } p = i \end{cases} \quad \text{and} \quad t_p := \begin{cases} \sigma_p & \text{if } p \neq i, j, \\ \sigma_j & \text{if } p = i, \\ \sigma_i & \text{if } p = j, \end{cases} \quad p \in \{1, \dots, m\}.$$
(67)

It is clear that both s and t converge to $\overline{\sigma}$ as $X \to \overline{X}$. By noting that g is absolutely symmetric on $\hat{\sigma}_{\mathcal{N}}$, we know from (6) that $g_j(\sigma) = g_i(t)$, since the vector t is obtained from σ by swapping the *i*-th and the *j*-th components. By the mean value theorem (cf. e.g., [38, Page 68-69]), we have

$$\frac{g_i(\sigma) - g_j(\sigma)}{\sigma_i - \sigma_j} = \frac{g_i(\sigma) - g_i(s) + g_i(s) - g_j(\sigma)}{\sigma_i - \sigma_j} = \frac{\frac{\partial g_i(\xi)}{\partial \mu_i}(\sigma_i - \sigma_j) + g_i(s) - g_j(\sigma)}{\sigma_i - \sigma_j} \\
= \frac{\partial g_i(\xi)}{\partial \mu_i} + \frac{g_i(s) - g_i(t) + g_i(t) - g_j(\sigma)}{\sigma_i - \sigma_j} \\
= \frac{\partial g_i(\xi)}{\partial \mu_i} + \frac{\frac{\partial g_i(\hat{\xi})}{\partial \mu_j}(\sigma_j - \sigma_i) + g_i(t) - g_j(\sigma)}{\sigma_i - \sigma_j} = \frac{\partial g_i(\xi)}{\partial \mu_i} - \frac{\partial g_i(\hat{\xi})}{\partial \mu_j},$$
(68)

where $\xi \in \mathbb{R}^m$ lies between σ and s and $\hat{\xi} \in \mathbb{R}^m$ is between s and t. Consequently, we have $\xi \to \overline{\sigma}$ and $\hat{\xi} \to \overline{\sigma}$ as $X \to \overline{X}$. By the continuity of g', we have

$$\lim_{X \to \overline{X}} \frac{g_i(\sigma) - g_j(\sigma)}{\sigma_i - \sigma_j} = (g'(\overline{\sigma}))_{ii} - (g'(\overline{\sigma}))_{ij} \quad \text{and} \quad \lim_{X \to \overline{X}} \frac{g_i(\sigma) + g_j(\sigma)}{\sigma_i + \sigma_j} = \frac{g_i(\overline{\sigma}) + g_j(\overline{\sigma})}{\overline{\sigma}_i + \overline{\sigma}_j}.$$

Therefore, we have

$$\lim_{X \to \overline{X}} G'(X) F^{(ij)} = \left[\left((g'(\overline{\sigma}))_{ii} - (g'(\overline{\sigma}))_{ij} \right) S(F_1^{(ij)}) + \frac{g_i(\overline{\sigma}) + g_j(\overline{\sigma})}{\overline{\sigma}_i + \overline{\sigma}_j} T(F_1^{(ij)}) \quad 0 \right] = G'(\overline{X}) F^{(ij)} .$$

Case 4: $i, j \in \{1, ..., m\}, i \neq j, \sigma_i > 0 \text{ or } \sigma_j > 0 \text{ and } \overline{\sigma}_i \neq \overline{\sigma}_j$. Then, we have $\sigma_i > 0 \text{ or } \sigma_j > 0$ and $\sigma_i \neq \sigma_j$. Since g' is continuous at $\overline{\sigma}$, we know that

$$\lim_{X \to \overline{X}} G'(X) F^{(ij)} = \lim_{X \to \overline{X}} \left[\frac{g_i(\sigma) - g_j(\sigma)}{\sigma_i - \sigma_j} S(F_1^{(ij)}) + \frac{g_i(\sigma) + g_j(\sigma)}{\sigma_i + \sigma_j} T(F_1^{(ij)}) \quad 0 \right]$$
$$= \left[\frac{g_i(\overline{\sigma}) - g_j(\overline{\sigma})}{\overline{\sigma}_i - \overline{\sigma}_j} S(F_1^{(ij)}) + \frac{g_i(\overline{\sigma}) + g_j(\overline{\sigma})}{\overline{\sigma}_i + \overline{\sigma}_j} T(F_1^{(ij)}) \quad 0 \right] = G'(\overline{X}) F^{(ij)}.$$

Case 5: $j \in \{m+1,\ldots,n\}$ and $\overline{\sigma}_i > 0$. Since g' is continuous at $\overline{\sigma}$, we obtain that

$$\lim_{X \to \overline{X}} G'(X) F^{(ij)} = \lim_{X \to \overline{X}} \left[0 \quad \frac{g_i(\sigma)}{\sigma_i} F_2^{(ij)} \right] = \left[0 \quad \frac{g_i(\overline{\sigma})}{\overline{\sigma}_i} F_2^{(ij)} \right] = G'(\overline{X}) F^{(ij)} \,.$$

Case 6: $i, j \in \{1, \ldots, m\}, i \neq j, \overline{\sigma}_i = \overline{\sigma}_j = 0 \text{ and } \sigma_i = \sigma_j > 0.$ Therefore, we know that

$$G'(X)F^{(ij)} = \left[((g'(\sigma))_{ii} - (g'(\sigma))_{ij})S(F_1^{(ij)}) + \frac{g_i(\sigma) + g_j(\sigma)}{\sigma_i + \sigma_j}T(F_1^{(ij)}) \quad 0 \right]$$

We know from (49) and Lemma 3.2 that

$$\lim_{X \to \overline{X}} (g'(\sigma))_{ii} = (g'(\overline{\sigma}))_{ii} = \overline{\eta}_i \quad \text{and} \quad \lim_{X \to \overline{X}} (g'(\sigma))_{ij} = (g'(\overline{\sigma}))_{ij} = 0.$$
(69)

Let $\hat{s}, \hat{t} \in \mathbb{R}^m$ be two vectors defined by

$$\hat{s}_p := \begin{cases} \sigma_p & \text{if } p \neq i, \\ -\sigma_j & \text{if } p = i \end{cases} \quad \text{and} \quad \hat{t}_p := \begin{cases} \sigma_p & \text{if } p \neq i, j, \\ -\sigma_j & \text{if } p = i, \\ -\sigma_i & \text{if } p = j, \end{cases} \quad p \in \{1, \dots, m\}.$$

$$(70)$$

Also, it clear that both \hat{s} and \hat{t} converge to $\overline{\sigma}$ as $X \to \overline{X}$. Again, by noting that g is absolutely symmetric on $\hat{\sigma}_{\mathcal{N}}$, we know from (6) that

$$g_i(\sigma) = -g_j(\hat{t})$$
 and $g_j(\sigma) = -g_i(\hat{t})$.

By using similar arguments for deriving (68), we have

$$\frac{g_i(\sigma) + g_j(\sigma)}{\sigma_i + \sigma_j} = \frac{\partial g_i(\zeta)}{\partial \mu_i} + \frac{\partial g_i(\hat{\zeta})}{\partial \mu_j}, \qquad (71)$$

where $\zeta \in \mathbb{R}^m$ is between σ and \hat{s} and $\hat{\zeta} \in \mathbb{R}^m$ is between \hat{s} and \hat{t} . Consequently, we know that $\zeta, \hat{\zeta} \to \overline{\sigma}$ as $X \to \overline{X}$. By the continuity of g', we know from (49) that

$$\lim_{X \to \overline{X}} \frac{g_i(\sigma) + g_j(\sigma)}{\sigma_i + \sigma_j} = (g'(\overline{\sigma}))_{ii} = \overline{\eta}_i.$$
(72)

Therefore, from (69) and (72), we have

$$\lim_{X \to \overline{X}} G'(X) F^{(ij)} = \begin{bmatrix} \overline{\eta}_i F_1^{(ij)} & 0 \end{bmatrix} = G'(\overline{\sigma}) F^{(ij)} .$$

Case 7: $i, j \in \{1, \ldots, m\}, i \neq j, \overline{\sigma}_i = \overline{\sigma}_j = 0, \sigma_i \neq \sigma_j \text{ and } \sigma_i > 0 \text{ or } \sigma_j > 0.$ Let s, t and \hat{s}, \hat{t} be defined by (67) and (70), respectively. By the continuity of g', we know from (68) and (71) that

$$\lim_{X \to \overline{X}} G'(X) F^{(ij)} = \lim_{X \to \overline{X}} \left[\frac{g_i(\sigma) - g_j(\sigma)}{\sigma_i - \sigma_j} S(F_1^{(ij)}) + \frac{g_i(\sigma) + g_j(\sigma)}{\sigma_i + \sigma_j} T(F_1^{(ij)}) \quad 0 \right]$$
$$= \left[\overline{\eta}_i S(F_1^{(ij)}) + \overline{\eta}_i T(F_1^{(ij)}) \quad 0 \right] = \left[\overline{\eta}_i F_1^{(ij)} \quad 0 \right] = G'(\overline{X}) F^{(ij)} + \overline{\eta}_i T(\overline{X}) F^{(ij)} = 0$$

Case 8: $i \neq j \in \{1, \ldots, m\}, \overline{\sigma}_i = \overline{\sigma}_j = 0 \text{ and } \sigma_i = \sigma_j = 0$. By the continuity of g', we obtain that

$$\lim_{X \to \overline{X}} G'(X) F^{(ij)} = \lim_{X \to \overline{X}} \begin{bmatrix} (g'(\sigma))_{ii} F_1^{(ij)} & 0 \end{bmatrix} = \begin{bmatrix} (g'(\overline{\sigma}))_{ii} F_1^{(ij)} & 0 \end{bmatrix} = \begin{bmatrix} \overline{\eta}_i F_1^{(ij)} & 0 \end{bmatrix} = G'(\overline{X}) F^{(ij)}.$$

Case 9: $j \in \{m+1,\ldots,n\}, \overline{\sigma}_i = 0 \text{ and } \sigma_i > 0$. We know that

$$G'(X)F^{(ij)} = \begin{bmatrix} 0 & \frac{g_i(\sigma)}{\sigma_i}F_2^{(ij)} \end{bmatrix}.$$

Let $\tilde{s} \in \mathbb{R}^m$ be a vector given by

$$\tilde{s}_p := \begin{cases} \sigma_p & \text{if } p \neq i, \\ 0 & \text{if } p = i, \end{cases} \quad p \in \{1, \dots, m\}.$$

Therefore, we have \tilde{s} converges to $\overline{\sigma}$ as $X \to \overline{X}$. Since g is absolutely symmetric on $\hat{\sigma}_{\mathcal{N}}$, we know that $g_i(\tilde{s}) = 0$. Also, by the mean value theorem, we have

$$\frac{g_i(\sigma)}{\sigma_i} = \frac{g_i(\sigma) - g_i(\tilde{s})}{\sigma_i} = \frac{\partial g_i(\rho)}{\partial \mu_i},$$

where $\rho \in \mathbb{R}^m$ is between σ and \tilde{s} . Consequently, we have ρ converges to $\overline{\sigma}$ as $X \to \overline{X}$. By the continuity of g', we know from (49) that

$$\lim_{X \to \overline{X}} \frac{g_i(\sigma)}{\sigma_i} = (g'(\overline{\sigma}))_{ii} = \overline{\eta}_i \,.$$

Thus,

$$\lim_{X \to \overline{X}} G'(X) F^{(ij)} = \lim_{X \to \overline{X}} \left[0 \quad \frac{g_i(\sigma)}{\sigma_i} F_2^{(ij)} \right] = \left[0 \quad \overline{\eta}_i F_2^{(ij)} \right] = G'(\overline{X}) F^{(ij)} \,.$$

Case 10: $j \in \{m+1,\ldots,n\}, \overline{\sigma}_i = 0 \text{ and } \sigma_i = 0$. By the continuity of g', we know that

$$\lim_{X \to \overline{X}} G'(X) F^{(ij)} = \lim_{X \to \overline{X}} \left[0 \quad (g'(\sigma))_{ii} F_2^{(ij)} \right] = \left[0 \quad (g'(\overline{\sigma}))_{ii} F_2^{(ij)} \right] = G'(\overline{X}) F^{(ij)}$$

Finally, we consider the general case that

$$X = U[\Sigma(X) \quad 0] V^{\mathbb{T}}$$
 and $\overline{X} = \overline{U} [\Sigma(\overline{X}) \quad 0] \overline{V}^{\mathbb{T}}$.

By noting from Theorem 3.3 that G is F-differential at X if and only if G is F-differential at $[\Sigma(X) \quad 0]$ and for any $H \in \mathbb{V}^{m \times n}$,

$$G'(X)H = U \begin{pmatrix} G'([\Sigma(X) \quad 0])(U^T H V) \end{pmatrix} V^T,$$

we know from the above analysis that G is continuously differentiable at \overline{X} .

" \Longrightarrow " Suppose that G is continuously differentiable at \overline{X} . Let $(\overline{U}, \overline{V}) \in \mathbb{O}^{m \times n}(\overline{X})$ be fixed. For any $\sigma \in \mathbb{R}^m$, define $X := \overline{U}[\text{Diag}(\sigma) \quad 0]\overline{V}^{\mathbb{T}}$. For any $h \in \mathbb{R}^m$, let $H := \overline{U}[\text{Diag}(h) \quad 0]\overline{V}^{\mathbb{T}}$. From the proof of the second part of Theorem 3.3, we know from the assumption that for all σ sufficiently close to $\overline{\sigma}$,

$$\operatorname{Diag}(g'(\sigma)h) = \overline{U}^{\mathbb{T}}(G'(X)H)\overline{V}_1, \quad h \in \mathbb{R}^m$$

Consequently, g is also continuously differentiable at $\overline{\sigma}$.

Remark 2 In order to compute (54), it appears that one needs to compute and store $\overline{V}_2 \in \mathbb{V}^{n \times (n-m)}$ explicitly, which would incur huge memory cost if $n \gg m$. Fortunately, due to the special form of $\overline{\mathcal{F}}$, the explicit computation of \overline{V}_2 can be avoided as we shall show next. Let $\overline{f} = (\overline{f}_1, \ldots, \overline{f}_m)^T$ be defined by

$$\bar{f}_i = \begin{cases} g_i(\bar{\sigma})/\bar{\sigma}_i & \text{if } \bar{\sigma}_i \neq 0, \\ (g'(\bar{\sigma}))_{ii} & \text{otherwise.} \end{cases}$$

Observe that the term in (54) involving \overline{V}_2 is given by

$$\overline{U}(\overline{\mathcal{F}} \circ (\overline{U}^{\mathbb{T}}H\overline{V}_2))\overline{V}_2^{\mathbb{T}} = \overline{U}\text{Diag}(\overline{f})\overline{U}^{\mathbb{T}}H\overline{V}_2\overline{V}_2^{\mathbb{T}} = \overline{U}\text{Diag}(\overline{f})\overline{U}^{\mathbb{T}}H(I_n - \overline{V}_1\overline{V}_1^{\mathbb{T}}) = \overline{U}\text{Diag}(\overline{f})\overline{U}^{\mathbb{T}}(H - (H\overline{V}_1)\overline{V}_1^{\mathbb{T}}).$$

Thus in numerical implementation, the large matrix \overline{V}_2 is not needed.

4 Lipschitz continuity, Bouligand differentiability, G-semismoothness, and Clarke's generalized Jacobian

4.1 Lipschitz continuity

In this subsection, we analyze the local Lipschitz continuity of the spectral operator G defined on a nonempty set \mathcal{N} . Let $\overline{X} \in \mathcal{N}$ be given. Assume that g is locally Lipschitz continuous near $\overline{\sigma} = \sigma(\overline{X})$ with module L > 0. Therefore, there exists a positive constant $\delta_0 > 0$ such that

$$\|g(\sigma) - g(\sigma')\| \le L \|\sigma - \sigma'\| \quad \forall \, \sigma, \sigma' \in B(\overline{\sigma}, \delta_0) := \{ y \in \hat{\sigma}_{\mathcal{N}} \mid \|y - \overline{\sigma}\| \le \delta_0 \} \; .$$

By using the absolutely symmetric property of g on $\hat{\sigma}_{\mathcal{N}}$, we obtain the following simple proposition.

Proposition 4.1 There exist a positive constant L' > 0 and a positive constant $\delta > 0$ such that for any $\sigma \in B(\overline{\sigma}, \delta)$,

$$g_i(\sigma) - g_j(\sigma)| \leq L' |\sigma_i - \sigma_j| \quad \forall i, j \in \{1, \dots, m\}, \ i \neq j, \ \sigma_i \neq \sigma_j,$$

$$(73)$$

$$|g_i(\sigma) + g_j(\sigma)| \leq L' |\sigma_i + \sigma_j| \quad \forall i, j \in \{1, \dots, m\}, \ \sigma_i + \sigma_j > 0,$$

$$(74)$$

$$|g_i(\sigma)| \leq L'|\sigma_i| \quad \forall i \in \{1, \dots, m\}, \ \sigma_i > 0.$$

$$(75)$$

Proof. It is easy to check that there exists a positive constant $\delta_1 > 0$ such that for any $\sigma \in B(\overline{\sigma}, \delta_1)$,

$$|\sigma_i - \sigma_j| \geq \delta_1 > 0 \quad \forall i, j \in \{1, \dots, m\}, \ i \neq j, \ \overline{\sigma}_i \neq \overline{\sigma}_j ,$$

$$(76)$$

$$|\sigma_i + \sigma_j| \geq \delta_1 > 0 \quad \forall \, i, j \in \{1, \dots, m\}, \ \overline{\sigma}_i + \overline{\sigma}_j > 0,$$

$$(77)$$

$$|\sigma_i| \geq \delta_1 > 0 \quad \forall i \in \{1, \dots, m\}, \ \overline{\sigma}_i > 0.$$

$$(78)$$

Let $\delta := \min\{\delta_0, \delta_1\} > 0$. Denote $\tau := \max_{i,j}\{|g_i(\overline{\sigma}) - g_j(\overline{\sigma})|, |g_i(\overline{\sigma}) + g_j(\overline{\sigma})|, |g_i(\overline{\sigma})|\} \ge 0, L_1 := (2L\delta + \tau)/\delta$ and $L' := \max\{L_1, \sqrt{2}L\}$. Let σ be any fixed vector in $B(\overline{\sigma}, \delta)$.

Firstly, we consider the case that $i, j \in \{1, ..., m\}, i \neq j$ and $\sigma_i \neq \sigma_j$. If $\overline{\sigma}_i \neq \overline{\sigma}_j$, then from (76), we know that

$$|g_{i}(\sigma) - g_{j}(\sigma)| = |g_{i}(\sigma) - g_{i}(\overline{\sigma}) + g_{i}(\overline{\sigma}) - g_{j}(\overline{\sigma}) + g_{j}(\overline{\sigma}) - g_{j}(\sigma)|$$

$$\leq 2||g(\sigma) - g(\overline{\sigma})|| + \tau \leq \frac{2L\delta + \tau}{\delta} |\sigma_{i} - \sigma_{j}| = L_{1}|\sigma_{i} - \sigma_{j}|.$$
(79)

If $\overline{\sigma}_i = \overline{\sigma}_j$, define $t \in \mathbb{R}^m$ by

$$t_p := \begin{cases} \sigma_p & \text{if } p \neq i, j, \\ \sigma_j & \text{if } p = i, \\ \sigma_i & \text{if } p = j, \end{cases} \quad p = 1, \dots, m$$

Then, we have $||t - \overline{\sigma}|| = ||\sigma - \overline{\sigma}|| \leq \delta$. Moreover, since g is absolutely symmetric on $\hat{\sigma}_{\mathcal{N}}$, we have $g_i(t) = g_j(\sigma)$. Therefore

$$|g_i(\sigma) - g_j(\sigma)| = |g_i(\sigma) - g_i(t)| \le ||g(\sigma) - g(t)|| \le L ||\sigma - t|| = \sqrt{2}L|\sigma_i - \sigma_j|.$$
(80)

Thus, the inequality (73) follows from (79) and (80) immediately.

Secondly, consider the case $i, j \in \{1, \ldots, m\}$ and $\sigma_i + \sigma_j > 0$. If $\overline{\sigma}_i + \overline{\sigma}_j > 0$, it follows from (77) that

$$|g_{i}(\sigma) + g_{j}(\sigma)| = |g_{i}(\sigma) - g_{i}(\overline{\sigma}) + g_{i}(\overline{\sigma}) + g_{j}(\overline{\sigma}) - g_{j}(\overline{\sigma}) + g_{j}(\sigma)|$$

$$\leq 2||g(\sigma) - g(\overline{\sigma})|| + \tau \leq \frac{2L\delta + \tau}{\delta}|\sigma_{i} + \sigma_{j}| = L_{1}|\sigma_{i} + \sigma_{j}|.$$
(81)

If $\overline{\sigma}_i + \overline{\sigma}_j = 0$, i.e., $\overline{\sigma}_i = \overline{\sigma}_j = 0$, define the vector $\hat{t} \in \mathbb{R}^m$ by

$$\hat{t}_p := \begin{cases} \sigma_p & \text{if } p \neq i, j, \\ -\sigma_j & \text{if } p = i, \\ -\sigma_i & \text{if } p = j, \end{cases} \quad p = 1, \dots, m.$$

By noting that $\overline{\sigma}_i = \overline{\sigma}_j = 0$, we obtain that $\|\hat{t} - \overline{\sigma}\| = \|\sigma - \overline{\sigma}\| \leq \delta$. Again, since g is absolutely symmetric on $\hat{\sigma}_{\mathcal{N}}$, we have $g_i(t) = g_j(\sigma)$, we have $g_i(\hat{t}) = -g_j(\sigma)$. Therefore,

$$|g_i(\sigma) + g_j(\sigma)| = |g_i(\sigma) - g_i(\hat{t})| \le ||g(\sigma) - g(\hat{t})|| \le L ||\sigma - \hat{t}|| = \sqrt{2}L|\sigma_i + \sigma_j|.$$
(82)

Thus the inequality (74) follows from (81) and (82).

Finally, we consider the case that $i \in \{1, \ldots, m\}$ and $\sigma_i > 0$. If $\overline{\sigma}_i > 0$, then we know from (78) that

$$|g_{i}(\sigma)| = |g_{i}(\sigma) - g_{i}(\overline{\sigma}) + g_{i}(\overline{\sigma})| \le |g_{i}(\sigma) - g_{i}(\overline{\sigma})| + |g_{i}(\overline{\sigma})|$$

$$\le ||g(\sigma) - g(\overline{\sigma})|| + \tau \le \frac{2L\delta + \tau}{\delta} |\sigma_{i}| \le L_{1}|\sigma_{i}|.$$
(83)

If $\overline{\sigma}_i = 0$, define $s \in \mathbb{R}^m$ by

$$s_p := \begin{cases} \sigma_p & \text{if } p \neq i, \\ 0 & \text{if } p = i, \end{cases} \quad p = 1, \dots, m.$$

Then, since $\sigma_i > 0$, we know that $||s - \overline{\sigma}|| < ||\sigma - \overline{\sigma}|| \le \delta$. Moreover, since g is absolutely symmetric on $\hat{\sigma}_N$, we have $g_i(t) = g_j(\sigma)$, we know that $g_i(s) = 0$. Therefore, we have

$$|g_i(\sigma)| = |g_i(\sigma) - g_i(s)| \le ||g(\sigma) - g(s)|| \le L ||\sigma - s|| \le L |\sigma_i|.$$
(84)

Thus, the inequality (73) follows from (83) and (84) immediately. This completes the proof.

For any fixed $0 < \omega \leq \delta_0/\sqrt{m}$ and $y \in B(\overline{\sigma}, \delta_0/(2\sqrt{m})) := \{\|y - \overline{\sigma}\|_{\infty} \leq \delta_0/(2\sqrt{m})\}$, the function g is integrable on $V_{\omega}(y) := \{z \in \mathbb{R}^m \mid \|y - z\|_{\infty} \leq \omega/2\}$ (in the sense of Lebesgue). Therefore, we know that the function

$$g(\omega, y) := \frac{1}{\omega^m} \int_{V_\omega(y)} g(z) dz \tag{85}$$

is well-defined on $(0, \delta_0/\sqrt{m}] \times B(\overline{\sigma}, \delta_0/(2\sqrt{m}))$ and is said to be Steklov averaged function [49] of g. For the sake of convenience, we always define g(0, y) = g(y). Since g is absolutely symmetric on $\hat{\sigma}_N$, we have $g_i(t) = g_j(\sigma)$, it is easy to check that for each fixed $0 < \omega \leq \delta_0/\sqrt{m}$, the function $g(\omega, \cdot)$ is also absolutely symmetric on $B(\overline{\sigma}, \delta_0/(2\sqrt{m}))$. By the definition, we know that $g(\cdot, \cdot)$ is locally Lipschitz continuous on $(0, \delta_0/\sqrt{m}] \times B(\overline{\sigma}, \delta_0/(2\sqrt{m}))$ with the module L. Meanwhile, by elementary calculations, we know that $g(\cdot, \cdot)$ is continuously differentiable on $(0, \delta_0/\sqrt{m}] \times B(\overline{\sigma}, \delta_0/(2\sqrt{m}))$ and for any fixed $\omega \in (0, \delta_0/\sqrt{m}]$ and $y \in B(\overline{\sigma}, \delta_0/(2\sqrt{m}))$,

$$\|g'_{y}(\omega, y)\| \le L.$$

Moreover, we know that $g(\omega, \cdot)$ converges to g uniformly on the compact set $B(\overline{\sigma}, \delta_0/(2\sqrt{m}))$ as $\omega \downarrow 0$. By using the formula (54), the following results can be obtained from Theorem 3.4 and Proposition 4.1 directly.

Proposition 4.2 Suppose that g is locally Lipschitz continuous near $\overline{\sigma}$, Let $g(\cdot, \cdot)$ be the corresponding Steklov averaged function defined in (85). Then, for any given $\omega \in (0, \delta_0/\sqrt{m}]$, the spectral operator $G(\omega, \cdot)$ with respect to $g(\omega, \cdot)$ is continuously differentiable on $B(\overline{X}, \delta_0/(2\sqrt{m})) := \{X \in \mathcal{X} \mid \|\sigma(X) - \overline{\sigma}\|_{\infty} \leq \delta_0/(2\sqrt{m})\}$, and there exist two positive constants $\delta_1 > 0$ and $\overline{L} > 0$ such that

$$\|G'(\omega, X)\| \le \overline{L} \quad \forall \, 0 < \omega \le \min\{\delta_0/\sqrt{m}, \delta_1\} \text{ and } X \in B(\overline{X}, \delta_0/(2\sqrt{m})).$$
(86)

Moreover, $G(\omega, \cdot)$ converges to G uniformly in the compact set $B(\overline{X}, \delta_0/(2\sqrt{m}))$ as $\omega \downarrow 0$.

Proposition 4.2 allows us to derive the following result on the local Lipschitz continuity of spectral operators.

Theorem 4.1 Suppose that \overline{X} has the SVD (22). The spectral operator G is locally Lipschitz continuous near \overline{X} if and only if g is locally Lipschitz continuous near $\overline{\sigma} = \sigma(\overline{X})$.

Proof. " \Leftarrow " Suppose that g is locally Lipschitz continuous near $\overline{\sigma} = \sigma(\overline{X})$ with module L > 0, i.e., there exists a positive constant $\delta_0 > 0$ such that

$$\|g(\sigma) - g(\sigma')\| \le L \|\sigma - \sigma'\| \quad \forall \sigma, \sigma' \in B(\overline{\sigma}, \delta_0).$$

By Proposition 4.2, for any $\omega \in (0, \delta_0/\sqrt{m}]$, the spectral operator $G(\omega, \cdot)$ defined with respect to the Steklov averaged function $g(\omega, \cdot)$ is continuously differentiable. Since $G(\omega, \cdot)$ converges to G uniformly in the compact set $B(\overline{X}, \delta_0/(2\sqrt{m}))$ as $\omega \downarrow 0$, we know that for any $\varepsilon > 0$, there exists a constant $\delta_2 > 0$ such that for any $0 < \omega \leq \delta_2$,

$$|G(\omega, X) - G(X)|| \le \varepsilon \quad \forall X \in B(\overline{X}, \delta_0/(2\sqrt{m})).$$

Fix any $X, X' \in B(\overline{X}, \delta_0/(2\sqrt{m}))$ with $X \neq X'$. By Proposition 4.2, we know that there exists $\delta_1 > 0$ such that (86) holds. Let $\overline{\delta} := \min\{\delta_1, \delta_2, \delta_0/\sqrt{m}\}$. Then, by the mean value theorem, we know that

$$\begin{aligned} \|G(X) - G(X')\| &= \|G(X) - G(\omega, X) + G(\omega, X) - G(\omega, X') + G(\omega, X') - G(X')\| \\ &\leq 2\varepsilon + \|\int_0^1 G'(\omega, X + t(X - X'))dt\| \\ &\leq \overline{L}\|X - X'\| + 2\varepsilon \quad \forall \, 0 < \omega < \overline{\delta} \,. \end{aligned}$$

Since $X, X' \in B(\overline{X}, \delta_0/(2\sqrt{m}))$ and $\varepsilon > 0$ are arbitrary, by letting $\varepsilon \downarrow 0$, we obtain that

$$\|G(X) - G(X')\| \le \overline{L} \|X - X'\| \quad \forall X, X' \in B(\overline{X}, \delta_0/(2\sqrt{m}))$$

Thus G is locally Lipschitz continuous near \overline{X} .

" \implies " Suppose that G is locally Lipschitz continuous near \overline{X} with module L > 0, i.e., there exists an open neighborhood \mathcal{B} of \overline{X} in \mathcal{N} such that for any $X, X' \in \mathcal{B}$,

$$||G(X) - G(X')|| \le L ||X - X'||.$$

Let $(\overline{U}, \overline{V}) \in \mathbb{O}^{m \times n}(\overline{X})$ be fixed. For any $y \in \hat{\sigma}_{\mathcal{N}}$, we define $Y := \overline{U}[\operatorname{Diag}(y) \quad 0] \overline{V}^{\mathbb{T}}$. Then, we know from Proposition 3.1 that $G(Y) = \overline{U}[\operatorname{Diag}(g(y)) \quad 0] \overline{V}^{\mathbb{T}}$. Therefore, we obtain that there exists an open neighborhood $\mathcal{B}_{\overline{\sigma}}$ of $\overline{\sigma}$ in $\hat{\sigma}_{\mathcal{N}}$ such that

$$||g(y) - g(y')|| = ||G(Y) - G(Y')|| \le L||Y - Y'|| = L||y - y'|| \quad \forall y, y' \in \mathcal{B}_{\overline{\sigma}}.$$

This completes the proof.

4.2 Bouligand-differentiability

In this section, we study the ρ -order Bouligand-differentiability of G with $0 < \rho \leq 1$, which is stronger than the directional differentiability. Let \mathcal{Z} be a finite dimensional real Euclidean space equipped with an inner product $\langle \cdot, \cdot \rangle$ and its induced norm $\|\cdot\|$. Let \mathcal{O} be an open set in \mathcal{Z} and \mathcal{Z}' be another finite dimensional real Euclidean space. The function $F : \mathcal{O} \subseteq \mathcal{Z} \to \mathcal{Z}'$ is said to be B(ouligand)-differentiable at $z \in \mathcal{O}$ if for any $h \in \mathcal{Z}$ with $h \to 0$,

$$F(z+h) - F(z) - F'(z;h) = o(||h||).$$

A stronger notion than B-differentiability is ρ -order B-differentiability with $\rho > 0$. The function $F : \mathcal{O} \subseteq \mathcal{Z} \to \mathcal{Z}'$ is said to be ρ -order B-differentiable at $z \in \mathcal{O}$ if for any $h \in \mathcal{Z}$ with $h \to 0$,

$$F(z+h) - F(z) - F'(z;h) = O(||h||^{1+\rho}).$$

Let $\overline{X} \in \mathbb{V}^{m \times n}$ be given. We have the following results on the ρ -order B-differentiability of spectral operators.

Theorem 4.2 Suppose that $\overline{X} \in \mathcal{N}$ has the SVD (22). Let $0 < \rho \leq 1$ be given. Then,

(i) if g is locally Lipschitz continuous near $\sigma(\overline{X})$ and ρ -order B-differentiable at $\sigma(\overline{X})$, then G is ρ -order B-differentiable at \overline{X} ;

(ii) if G is ρ -order B-differentiable at \overline{X} , then g is ρ -order B-differentiable at $\sigma(\overline{X})$.

Proof. Without loss of generality, we only consider the case that $\rho = 1$.

(i) For any $H \in \mathbb{V}^{m \times n}$, denote $X = \overline{X} + H$. Let $U \in \mathbb{O}^m$ and $V \in \mathbb{O}^n$ be such that

$$X = U[\Sigma(X) \quad 0]V^{\mathbb{T}}.$$
(87)

Denote $\sigma = \sigma(X)$. Let $G_S(X)$ and $G_R(X)$ be defined by (27). Therefore, by Lemma 3.1, we know that for any $H \to 0$,

$$G_{S}(X) - G_{S}(\overline{X}) = G'_{S}(\overline{X})H + O(||H||^{2}) = G'_{S}(\overline{X})H + O(||H||^{2}),$$
(88)

where $G'_S(\overline{X})H$ is given by (28). For $H \in \mathbb{V}^{m \times n}$ sufficiently small, we have $\mathcal{U}_l(X) = \sum_{i \in a_l} u_i v_i^{\mathbb{T}}, l = 1, \ldots, r$. Therefore, we know that

$$G_R(X) = G(X) - G_S(X) = \sum_{l=1}^{r+1} \Delta_l(H),$$
(89)

where

$$\Delta_l(H) = \sum_{i \in a_l} (g_i(\sigma) - g_i(\overline{\sigma})) u_i v_i^{\mathbb{T}} \quad l = 1, \dots, r \quad \text{and} \quad \Delta_{r+1}(H) = \sum_{i \in b} g_i(\sigma) u_i v_i^{\mathbb{T}}.$$

We first consider the case that $\overline{X} = [\Sigma(\overline{X}) \quad 0]$. Then, we know from (12) and (13) that for any H sufficiently small,

$$\sigma = \overline{\sigma} + \sigma'(\overline{X}; H) + O(\|H\|^2), \qquad (90)$$

where $\sigma'(\overline{X}; H) = (\lambda(S(H_{a_1a_1})), \ldots, \lambda(S(B_{a_ra_r})), \sigma([H_{bb} \quad H_{bc}])) \in \mathbb{R}^m$. Denote $h := \sigma'(\overline{X}; H)$. Since g is locally Lipschitz continuous near $\overline{\sigma}$ and 1-order B-differentiable at $\overline{\sigma}$, we know that for any H sufficiently small,

$$g(\sigma) - g(\overline{\sigma}) = g(\sigma + h + O(||H||^2)) - g(\overline{\sigma}) = g(\sigma + h) - g(\overline{\sigma}) + O(||H||^2) = g'(\overline{\sigma}; h) + O(||H||^2).$$

Let $\phi = g'(\overline{\sigma}; \cdot)$. Since $u_i v_i^{\mathbb{T}}$, $i = 1, \ldots, m$ are uniformly bounded, we obtain that for H sufficiently small,

$$\Delta_l(H) = U_{a_l} \operatorname{Diag}(\phi_l(h)) V_{a_l}^{\mathbb{T}} + O(||H||^2), \quad l = 1, \dots, r,$$

$$\Delta_{r+1}(H) = U_b \operatorname{Diag}(\phi_{r+1}(h)) V_b^{\mathbb{T}} + O(||H||^2).$$

Again, we know from (14) that there exist $Q_l \in \mathbb{O}^{|a_l|}$, $M \in \mathbb{O}^{|b|}$ and $N = [N_1 \quad N_2] \in \mathbb{O}^{n-|a|}$ with $N_1 \in \mathbb{V}^{(n-|a|) \times |b|}$ and $N_2 \in \mathbb{V}^{(n-|a|) \times (n-m)}$ (depending on H) such that

$$U_{a_{l}} = \begin{bmatrix} O(\|H\|) \\ Q_{l} + O(\|H\|) \\ O(\|H\|) \end{bmatrix}, \quad V_{a_{l}} = \begin{bmatrix} O(\|H\|) \\ Q_{l} + O(\|H\|) \\ O(\|H\|) \end{bmatrix}, \quad l = 1, \dots, r,$$
$$U_{b} = \begin{bmatrix} O(\|H\|) \\ M + O(\|H\|) \end{bmatrix}, \quad [V_{b} \quad V_{c}] = \begin{bmatrix} O(\|H\|) \\ N + O(\|H\|) \end{bmatrix}.$$

Since g is locally Lipschitz continuous near $\overline{\sigma}$ and directionally differentiable at $\overline{\sigma}$, we know from [44, Theorem A.2] or [41, Lemma 2.2] that the directional derivative ϕ is globally Lipschitz continuous on \mathbb{R}^m . Thus, for H sufficiently small, we have $\|\phi(h)\| = O(\|H\|)$. Therefore, we obtain that

$$\Delta_{l}(H) = \begin{bmatrix} 0 & 0 & 0\\ 0 & Q_{l} \operatorname{Diag}(\phi_{l}(h)) Q_{l}^{\mathbb{T}} & 0\\ 0 & 0 & 0 \end{bmatrix} + O(\|H\|^{2}), \quad l = 1, \dots, r,$$
(91)

$$\Delta_{r+1}(H) = \begin{bmatrix} 0 & 0\\ 0 & M \text{Diag}(\phi_{r+1}(h)) N_1^{\mathbb{T}} \end{bmatrix} + O(\|H\|^2).$$
(92)

Again, we know from (15) and (16) that

$$S(H_{a_l a_l}) = Q_l(\Sigma(X)_{a_l a_l} - \overline{\nu}_l I_{|a_l|}) Q_l^{\mathbb{T}} + O(||H||^2), \quad l = 1, \dots, r,$$
(93)

$$[H_{bb} \quad H_{bc}] = M(\Sigma(X)_{bb} - \overline{\nu}_{r+1}I_{|b|})N_1^{\mathbb{T}} + O(||H||^2).$$
(94)

Since q is locally Lipschitz continuous near $\overline{\sigma} = \sigma(\overline{X})$, we know from Theorem 4.1 that the spectral operator G is locally Lipschitz continuous near \overline{X} . Therefore, we know from Theorem 3.2 and Remark 1 that G is directional differentiable at \overline{X} . Thus, from [44, Theorem A.2] or [41, Lemma 2.2], we know that $G'(\overline{X}, \cdot)$ is globally Lipschitz continuous on $\mathbb{V}^{m \times n}$. Thus, the corresponding spectral operator Φ defined by (33) is globally Lipschitz continuous on \mathcal{W} . Hence, we know from (89) that for H sufficiently small,

$$G_R(X) = \widehat{\Phi}(D(H)) + O(||H||^2), \qquad (95)$$

where $D(H) = (S(H_{a_1a_1}), \dots, S(H_{a_ra_r}), H_{b\bar{a}})$ and $\widehat{\Phi}(\cdot)$ is defined by (36). Next, consider the general case that $\overline{X} \in \mathbb{V}^{m \times n}$. For any $H \in \mathbb{V}^{m \times n}$, rewrite (87) as

$$[\Sigma(\overline{X}) \quad 0] + \overline{U}^{\mathbb{T}} H \overline{V} = \overline{U}^{\mathbb{T}} U[\Sigma(Z) \quad 0] V^{\mathbb{T}} \overline{V}.$$

Denote $\widetilde{U} := \overline{U}^{\mathbb{T}}U$ and $\widetilde{V} := \overline{V}^{\mathbb{T}}V$. Let $\widetilde{X} := [\Sigma(\overline{X}) \quad 0] + \overline{U}^{\mathbb{T}}H\overline{V}$. Then, since \overline{U} and \overline{V} are unitary matrices, we know from (95) that

$$G_R(X) = \overline{U}\widehat{\Phi}(D(H))\overline{V}^{\mathbb{T}} + O(||H||^2), \qquad (96)$$

where $D(H) = \left(S(\widetilde{H}_{a_1a_1}), \dots, S(\widetilde{H}_{a_ra_r}), \widetilde{H}_{b\bar{a}}\right)$ and $\widetilde{H} = \overline{U}^{\mathbb{T}}H\overline{V}$. Thus, by combining (88) and (96) and noting that $G(X) = G_S(\overline{X})$, we obtain that for any $H \in \mathbb{V}^{m \times n}$ sufficiently close to 0,

$$G(X) - G(\overline{X}) - G'(\overline{X}; H) = O(||H||^2),$$

where $G'(\overline{X}; H)$ is given by (37). This implies that G is 1-order B-differentiable at \overline{X} .

(ii) Suppose that G is 1-order B-differentiable at \overline{X} . Let $(\overline{U}, \overline{V}) \in \mathbb{O}^{m \times n}(\overline{X})$ be fixed. For any $h \in \mathbb{R}^m$, let $H = \overline{U}[\text{Diag}(h) \quad 0]\overline{V}^{\mathbb{T}} \in \mathbb{V}^{m \times n}$. We know from Proposition 3.1 that for all h sufficiently close to 0, $G(\overline{X} + H) = \overline{U}\text{Diag}(g(\overline{\sigma} + h))\overline{V}_1^{\mathbb{T}}$. Therefore, we know from the assumption that

$$\operatorname{Diag}(g(\overline{\sigma}+h)-g(\overline{\sigma})) = \overline{U}^{\mathbb{T}}\left(G(\overline{X}+H)-G(\overline{X})\right)\overline{V}_1 = \overline{U}^{\mathbb{T}}G'(\overline{X};H)\overline{V}_1 + O(\|H\|^2).$$

This shows that g is 1-order B-differentiable at $\overline{\sigma}$. The proof is competed.

4.3 G-semismoothness

Let \mathcal{Z} and \mathcal{Z}' be two finite dimensional real Euclidean spaces and \mathcal{O} be an open set in \mathcal{Z} . Suppose that $F: \mathcal{O} \subseteq \mathcal{Z} \to \mathcal{Z}'$ is a locally Lipschitz continuous function on \mathcal{O} . Then, according to Rademacher's theorem, F is almost everywhere differentiable (in the sense of Fréchet) in \mathcal{O} . Let \mathcal{D}_F be the set of points in \mathcal{O} where F is differentiable. Let F'(z) be the derivative of F at $z \in \mathcal{D}_F$. Then the B(ouligand)-subdifferential of F at $z \in \mathcal{O}$ is denoted by [40]:

$$\partial_B F(z) := \left\{ \lim_{\mathcal{D}_F \ni z^k \to z} F'(z^k) \right\}$$

and Clarke's generalized Jacobian of F at $z \in \mathcal{O}$ [10] takes the form:

$$\partial F(z) = \operatorname{conv}\{\partial_B F(z)\},\,$$

where "conv" stands for the convex hull in the usual sense of convex analysis [45]. The function F is said to be G-semismooth at a point $z \in \mathcal{O}$ if for any $y \to z$ and $V \in \partial F(y)$,

$$F(y) - F(z) - V(y - z) = o(||y - z||).$$

A stronger notion than G-semismoothness is ρ -order G-semismoothness with $\rho > 0$. The function F is said to be ρ -order G-semismooth at z if for any $y \to z$ and $V \in \partial F(y)$,

$$F(y) - F(z) - V(y - z) = O(||y - z||^{1+\rho}).$$

In particular, the function F is said to be strongly G-semismooth at z if F is 1-order G-semismooth at z. Furthermore, the function F is said to be (ρ -order, strongly) semismooth at $z \in \mathcal{O}$ if (i) the directional derivative of F at z along any direction $d \in \mathbb{Z}$, denoted by F'(z; d), exists; and (ii) F is (ρ -order, strongly) G-semismooth.

The following result taken from [50, Theorem 3.7] provides a convenient tool for proving the G-semismoothness of Lipschitz functions.

Lemma 4.1 Let $F : \mathcal{O} \subseteq \mathcal{Z} \to \mathcal{Z}'$ be a locally Lipschitz continuous function on the open set \mathcal{O} . Let $\rho > 0$ be a constant. F is ρ -order G-semismooth (G-semismooth) at z if and only if for any $\mathcal{D}_F \ni y \to z$,

$$F(y) - F(z) - F'(y)(y - z) = O(||y - z||^{1+\rho}) \quad (= o(||y - z||)).$$
(97)

Let $\overline{X} \in \mathcal{N}$ be given. Assume that g is locally Lipschitz continuous near $\overline{\sigma} = \sigma(\overline{X})$. Thus, from Theorem 4.1 we know that the corresponding spectral operator G is locally Lipschitz continuous near \overline{X} . The following theorem is on the G-semismoothness of the spectral operator G.

Theorem 4.3 Suppose that $\overline{X} \in \mathcal{N}$ has the decomposition (22). Let $0 < \rho \leq 1$ be given. G is ρ -order G-semismooth at \overline{X} if and only if g is ρ -order G-semismooth at $\overline{\sigma}$.

Proof. Without loss of generality, we only consider the case that $\rho = 1$. " \Leftarrow " For any $H \in \mathbb{V}^{m \times n}$, denote $X = \overline{X} + H$. Let $U \in \mathbb{O}^m$ and $V \in \mathbb{O}^n$ be such that

$$X = U[\Sigma(X) \quad 0]V^{\mathbb{T}}.$$
(98)

Denote $\sigma = \sigma(X)$. G_S and G_R are two mappings defined in (27). We know from Lemma 3.1 that there exists an open neighborhood $\mathcal{B} \subseteq \mathcal{N}$ of \overline{X} such that G_S twice continuously differentiable on \mathcal{B} and

$$G_{S}(X) - G_{S}(\overline{X}) = \sum_{l=1}^{r} \bar{g}_{l} \mathcal{U}_{l}'(X)H + O(\|H\|^{2})$$

$$= \sum_{l=1}^{r} \bar{g}_{l} \left\{ U[\Gamma_{l}(X) \circ S(U^{\mathbb{T}}HV_{1}) + \Xi_{l}(X) \circ T(U^{\mathbb{T}}HV_{1})]V_{1}^{\mathbb{T}} + U(\Upsilon_{l}(X) \circ U^{\mathbb{T}}HV_{2})V_{2}^{\mathbb{T}} \right\} + O(\|H\|^{2}),$$
(99)

where for each $l \in \{1, \ldots, r\}$, $\Gamma_l(X)$, $\Xi_l(X)$ and $\Upsilon_l(X)$ are given by (18)-(20) with X, respectively. By taking a smaller \mathcal{B} if necessary, we assume that for any $X \in \mathcal{B}$ and $l, l' \in \{1, \ldots, r\}$,

$$\sigma_i(X) > 0, \quad \sigma_i(X) \neq \sigma_j(X) \quad \forall i \in a_l, \ j \in a_{l'} \text{ and } l \neq l'.$$
 (100)

Since g is locally Lipschitz continuous near $\overline{\sigma}$, we know that for any H sufficiently small,

$$\bar{g}_l = g_i(\sigma) + O(||H||) \quad \forall i \in a_l, \quad l = 1, \dots, r.$$

Let \mathcal{E}_1 , \mathcal{E}_2 and \mathcal{F} (depending on X) be the matrices defined by (24)-(26). By noting that $U \in \mathbb{O}^m$ and $V \in \mathbb{O}^n$ are uniformly bounded, since g is locally Lipschitz continuous near $\overline{\sigma}$, we know that for any $X \in \mathcal{B}$ (shrinking \mathcal{B} if necessary),

$$G_S(X) - G_S(\overline{X}) = U \left[\mathcal{E}_1^0 \circ S(U^{\mathbb{T}} H V_1) + \mathcal{E}_2^0 \circ T(U^{\mathbb{T}} H V_1) \quad \mathcal{F}^0 \circ U^{\mathbb{T}} H V_2 \right] V^{\mathbb{T}} + O(\|H\|^2),$$
(101)

where \mathcal{E}_1^0 , \mathcal{E}_2^0 and \mathcal{F}^0 are the corresponding real matrices defined in (24)-(26), respectively.

Let $X \in \mathcal{D}_G \cap \mathcal{B}$, where \mathcal{D}_G is the set of points in $\mathbb{V}^{m \times n}$ for which G is (F-)differentiable. Define the corresponding index sets in $\{1, \ldots, m\}$ for X by $a' := \{i \mid \sigma_i(X) > 0\}$ and $b' := \{i \mid \sigma_i(X) = 0\}$. By (100), we have

$$a' \supseteq a \quad \text{and} \quad b' \subseteq b \,.$$
 (102)

Let \mathcal{E}_1 , \mathcal{E}_2 , \mathcal{F} and \mathcal{C} be the corresponding real matrices defined in (50)–(53), respectively. We know from Theorem 3.3 that

$$G'(X)H = U[\mathcal{E}_1 \circ S(U^{\mathbb{T}}HV_1) + \mathcal{E}_2 \circ T(U^{\mathbb{T}}HV_1) + \operatorname{Diag}\left(\mathcal{C}\operatorname{diag}(S(U^{\mathbb{T}}HV_1))\right) \quad \mathcal{F} \circ U^{\mathbb{T}}HV_2]V^{\mathbb{T}}, \quad (103)$$

where η , \mathcal{E}_1 , \mathcal{E}_2 , \mathcal{F} and \mathcal{C} are defined by (49)–(53) with respect to σ , respectively. Denote

$$\Delta(H) := G'(X)H - (G_S(X) - G_S(\overline{X})).$$

Moreover, since there exists an integer $j \in \{0, \ldots, |b|\}$ such that |a'| = |a| + j, we can define two index sets $b_1 := \{|a| + 1, \ldots, |a| + j\}$ and $b_2 := \{|a| + j + 1, \ldots, |a| + |b|\}$ such that $a' = a \cup b_1$ and $b' = b_2$. From (101) and (103), we obtain that

$$\Delta(H) = U\widehat{R}(H)V^{\mathbb{T}} + O(||H||^2), \qquad (104)$$

where $\widehat{R}(H) \in \mathbb{V}^{m \times n}$ is defined by

$$\widehat{R}(H) := \begin{bmatrix} \operatorname{Diag}\left(R_1(H), \dots, R_r(H)\right) & 0\\ 0 & R_{r+1}(H) \end{bmatrix},$$

$$R_l(H) = (\mathcal{E}_1)_{a_l a_l} \circ S(U_{a_l}^{\mathbb{T}} H V_{a_l}) + \operatorname{Diag}\left((\mathcal{C}\operatorname{diag}(S(U^{\mathbb{T}} H V_1)))_{a_l a_l} \right), \ l = 1, \dots, r,$$
(105)

$$R_{r+1}(H) = \begin{bmatrix} (\mathcal{E}_1)_{b_1 b_1} \circ S(U_{b_1}^{\mathbb{T}} H V_{b_1}) + \text{Diag} \left((\mathcal{C} \text{diag}(S(U^{\mathbb{T}} H V_1)))_{b_1 b_1} \right) & 0 & 0 \\ 0 & \gamma U_{b_2}^{\mathbb{T}} H V_{b_2} & \gamma U_{b_2}^{\mathbb{T}} H V_2 \end{bmatrix} (106)$$

and $\gamma := (g'(\sigma))_{ii}$ for any $i \in b_2$. By (22), we obtain from (98) that

$$[\Sigma(\overline{X}) \quad 0] + \overline{U}^{\mathbb{T}} H \overline{V} = \overline{U}^{\mathbb{T}} U [\Sigma(X) \quad 0] V^{\mathbb{T}} \overline{V}.$$

Let $\widehat{H} := \overline{U}^{\mathbb{T}} H \overline{V}$, $\widehat{U} := \overline{U}^{\mathbb{T}} U$ and $\widehat{V} := \overline{V}^{\mathbb{T}} V$. Then, $U^{\mathbb{T}} H V = \widehat{U}^{\mathbb{T}} \overline{U}^{\mathbb{T}} H \overline{V} \widehat{V} = \widehat{U}^{\mathbb{T}} \widehat{H} \widehat{V}$. We know from (14) that there exist $Q_l \in \mathbb{O}^{|a_l|}$, $l = 1, \ldots, r$ and $M \in \mathbb{O}^{|b|}$, $N \in \mathbb{O}^{n-|a|}$ such that

$$U_{a_l}^{\mathbb{T}} H V_{a_l} = \widehat{U}_{a_l}^{\mathbb{T}} \widehat{H} \widehat{V}_{a_l} = Q_l^{\mathbb{T}} \widehat{H}_{a_l a_l} Q_l + O(||H||^2), \quad l = 1, \dots, r,$$

$$\begin{bmatrix} U_b^{\mathbb{T}} H V_b & U_b^{\mathbb{T}} H V_2 \end{bmatrix} = \begin{bmatrix} \widehat{U}_b^{\mathbb{T}} \widehat{H} \widehat{V}_b & \widehat{U}_b^{\mathbb{T}} \widehat{H} \widehat{V}_2 \end{bmatrix} = M^{\mathbb{T}} \begin{bmatrix} \widehat{H}_{bb} & \widehat{H}_{bc} \end{bmatrix} N + O(||H||^2).$$

Moreover, from (15) and (16), we obtain that

$$S(U_{a_{l}}^{\mathbb{T}}HV_{a_{l}}) = Q_{l}^{\mathbb{T}}S(\hat{H}_{a_{l}a_{l}})Q_{l} + O(||H||^{2}) = \Sigma(X)_{a_{l}a_{l}} - \Sigma(\overline{X})_{a_{l}a_{l}} + O(||H||^{2}), \quad l = 1, \dots, r,$$
$$[U_{b}^{\mathbb{T}}HV_{b} \quad U_{b}^{\mathbb{T}}HV_{2}] = M^{\mathbb{T}}\left[\hat{H}_{bb} \quad \hat{H}_{bc}\right]N = \left[\Sigma(X)_{bb} - \Sigma(\overline{X})_{bb} \quad 0\right] + O(||H||^{2}).$$

Denote $h = \sigma'(X; H) \in \mathbb{R}^m$. Since the single value functions are strongly semismooth [51], we know that

$$S(U_{a_{l}}^{\mathbb{T}}HV_{a_{l}}) = \text{Diag}(h_{a_{l}}) + O(||H||^{2}), \quad l = 1, \dots, r,$$

$$S(U_{b_{1}}^{\mathbb{T}}HV_{b_{1}}) = \text{Diag}(h_{b_{1}}) + O(||H||^{2}), \quad [U_{b_{2}}^{\mathbb{T}}HV_{b_{2}} \quad U_{b_{2}}^{\mathbb{T}}HV_{2}] = [\text{Diag}(h_{b_{2}}) \quad 0] + O(||H||^{2}).$$

Therefore, since $\mathcal{C} = g'(\sigma) - \text{Diag}(\eta)$, by (105) and (106), we obtain from (104) that

$$\Delta(H) = U \left[\text{Diag}\left(g'(\sigma)h\right) \quad 0 \right] V^{\mathbb{T}} = U \left[\text{Diag}\left(g'(\sigma)h\right) \quad 0 \right] V^{\mathbb{T}} + O(\|H\|^2) \,. \tag{107}$$

On the other hand, for X sufficiently close to \overline{X} , we have $\mathcal{U}_l(X) = \sum_{i \in a_l} u_i v_i^{\mathbb{T}}, l = 1, \ldots, r$. Therefore,

$$G_R(X) = G(X) - G_S(X) = \sum_{l=1}^r \sum_{i \in a_l} [g_i(\sigma) - g_i(\overline{\sigma})] u_i v_i^{\mathbb{T}} + \sum_{i \in b} g_i(\sigma) u_i v_i^{\mathbb{T}}.$$
 (108)

We know from Theorem 3.3 that G is differentiable at X if and only if g is differentiable at σ . Since g is 1-order G-semismooth at $\overline{\sigma}$ and $\sigma(\cdot)$ is strongly semismooth, we obtain that for any $X \in \mathcal{D}_G \cap \mathcal{B}$ (shrinking \mathcal{B} if necessary),

$$g(\sigma) - g(\overline{\sigma}) = g'(\sigma)(\sigma - \overline{\sigma}) + O(||H||^2) = g'(\sigma)(h + O(||H||^2)) + O(||H||^2) = g'(\sigma)h + O(||H||^2).$$

Then, since $U \in \mathbb{O}^m$ and $U \in \mathbb{O}^n$ are uniformly bounded, we obtain from (108) that

$$G_R(X) = U \left[\text{Diag} \left(g'(\sigma)h \right) \quad 0 \right] V^{\mathbb{T}} + O(||H||^2).$$

Thus, from (107), we obtain that $\Delta(H) = G_R(X) + O(||H||^2)$. That is, for any $X \in \mathcal{D}_G$ converging to \overline{X} ,

$$G(X) - G(\overline{X}) - G'(X)H = -\Delta(H) + G_R(X) = O(||H||^2)$$

" \Longrightarrow " Suppose that G is 1-order G-semismooth at \overline{X} . Let $(\overline{U}, \overline{V}) \in \mathbb{O}^{m \times n}(\overline{X})$ be fixed. Assume that $\sigma = \overline{\sigma} + h \in \mathcal{D}_g$ and $h \in \mathbb{R}^m$ is sufficiently small. Let $X = \overline{U}[\operatorname{Diag}(\sigma) \quad 0] \overline{V}^{\mathbb{T}}$ and $H = \overline{U}[\operatorname{Diag}(h) \quad 0] \overline{V}^{\mathbb{T}}$. Then, $X \in \mathcal{D}_G$ and converges to \overline{X} if h goes to zero. We know from Proposition 3.1 that for all h sufficiently close to 0, $G(X) = \overline{U}\operatorname{Diag}(g(\sigma))\overline{V}_1^{\mathbb{T}}$. Therefore, for any h sufficiently close to 0,

$$\operatorname{Diag}(g(\overline{\sigma}+h)-g(\overline{\sigma}))=\overline{U}^{\mathbb{T}}\left(G(X)-G(\overline{X})\right)\overline{V}_{1}=\overline{U}^{\mathbb{T}}G'(X)H\overline{V}_{1}+O(||H||^{2}).$$

Hence, since obviously $\text{Diag}(g'(\sigma)h) = \overline{U}^{\mathbb{T}}G'(X)H\overline{V}_1$, we know that for h sufficiently small, $g(\overline{\sigma} + h) - g(\overline{\sigma}) = g'(\overline{\sigma})h + O(\|h\|^2)$. Then, g is 1-order G-semismooth at $\overline{\sigma}$.

4.4 Characterization of Clarke's generalized Jacobian

Let $\overline{X} \in \mathcal{N}$ be given. In this section, we assume that g is locally Lipschitz continuous near $\overline{\sigma} = \sigma(\overline{X})$ and directionally differentiable at $\overline{\sigma}$. Therefore, from Theorem 4.1, Theorem 3.2 and Remark 1, we know that the corresponding spectral operator G is locally Lipschitz continuous near \overline{X} and directionally differentiable at \overline{X} . Furthermore, we define the function $d : \mathbb{R}^m \to \mathbb{R}^m$ by

$$d(h) := g(\overline{\sigma} + h) - g(\overline{\sigma}) - g'(\overline{\sigma}; h), \quad h \in \mathbb{R}^m.$$
(109)

Thus, since g is absolutely symmetric on the nonempty open set $\hat{\sigma}_{\mathcal{N}}$, we know from (30) and (32) that d is also a mixed symmetric mapping, with respect to $\mathbb{P}^{|a_1|} \times \ldots \times \mathbb{P}^{|a_r|} \times \pm \mathbb{P}^{|b|}$, over $\mathbb{R}^{|a_1|} \times \ldots \times \mathbb{R}^{|a_r|} \times \mathbb{R}^{|b|}$. Moreover, since g is locally Lipschitz continuous near $\overline{\sigma}$ and directional differentiable at $\overline{\sigma}$, we know that g is B-differentiable at $\overline{\sigma}$ (cf. [47]). Thus, d is differentiable at zero with the derivative d'(0) = 0. Furthermore, if we assume that the function d is also strictly differentiable at zero, then we have

$$\lim_{\substack{w,w' \to 0 \\ w \neq w'}} \frac{d(w) - d(w')}{\|w - w'\|} = 0.$$
(110)

Thus, by using the mixed symmetric property of d, one can easily obtain the following results. We omit the details of the proof here.

Lemma 4.2 Let $d : \mathbb{R}^m \to \mathbb{R}^m$ be the function given by (109). Suppose that d is strictly differentiable at zero. Let $\{w^k\}$ be a given sequence in \mathbb{R}^m converging to zero. Then, if there exist $i, j \in a_l$ for some $l \in \{1, \ldots, r\}$ or $i, j \in b$ such that $w_i^k \neq w_j^k$ for all k sufficiently large, then

$$\lim_{k \to \infty} \frac{d_i(w^k) - d_j(w^k)}{w_i^k - w_j^k} = 0;$$
(111)

if there exist $i, j \in b$ such that $w_i^k + w_i^k \neq 0$ for all k sufficiently large, then

$$\lim_{k \to \infty} \frac{d_i(w^k) + d_j(w^k)}{w_i^k + w_j^k} = 0;$$
(112)

and if there exists $i \in b$ such that $w_i^k \neq 0$ for all k sufficiently large, then

$$\lim_{k \to \infty} \frac{d_i(w^k)}{w_i^k} = 0.$$
(113)

Let $\Psi(\cdot) := G'(\overline{X}; \cdot) : \mathbb{V}^{m \times n} \to \mathbb{V}^{m \times n}$ be the directional derivative of G at \overline{X} . We know from (37) that for any $Z \in \mathbb{V}^{m \times n}$,

$$\Psi(Z) = G'(\overline{X}; Z) = \overline{U} \left[\overline{\mathcal{E}}_1^0 \circ S(\overline{U}^{\mathbb{T}} Z \overline{V}_1) + \overline{\mathcal{E}}_2^0 \circ T(\overline{U}^{\mathbb{T}} Z \overline{V}_1) \quad \overline{\mathcal{F}}^0 \circ \overline{U}^{\mathbb{T}} Z \overline{V}_2 \right] \overline{V}^{\mathbb{T}} + \overline{U} \widehat{\Phi}(D(Z)) \overline{V}^{\mathbb{T}}, \quad (114)$$

where $D(Z) = \left(S(\widetilde{Z}_{a_1a_1}), \dots, S(\widetilde{Z}_{a_ra_r}), \widetilde{Z}_{b\overline{a}}\right) \in \mathcal{W}, \widetilde{Z} = \overline{U}^{\mathbb{T}} Z \overline{V} \text{ and } \widehat{\Phi}(\cdot) : \mathcal{W} \to \mathbb{V}^{m \times n} \text{ is given by (36) with } \Phi(\cdot) : \mathcal{W} \to \mathcal{W} \text{ being the spectral operator defined by (33) with respect to the mixed symmetric mapping } \phi(\cdot) := g'(\overline{\sigma}; \cdot).$ Since the spectral operator G is locally Lipschitz continuous near \overline{X} , we know that $\Psi(\cdot) = G'(\overline{X}; \cdot)$ is globally Lipschitz continuous (cf. [44, Theorem A.2] or [41, Lemma 2.2]). Therefore, $\partial_B \Psi(0)$ and $\partial \Psi(0)$ are well-defined. Furthermore, we have the following useful results on the characterization of the B-subdifferential and Clarke's subdifferential of the spectral operator G at \overline{X} .

Theorem 4.4 Suppose that the given $\overline{X} \in \mathcal{N}$ has the decomposition (22). Suppose that there exists an open neighborhood $\mathcal{B} \subseteq \mathbb{R}^m$ of $\overline{\sigma}$ in $\hat{\sigma}_{\mathcal{N}}$ such that $g(\cdot)$ is differentiable at $\sigma \in \mathcal{B}$ if and only if $g'(\overline{\sigma}; \cdot)$ is differentiable at $\sigma - \overline{\sigma}$. Assume further that the function $d : \mathbb{R}^m \to \mathbb{R}^m$ defined by (109) is strictly differentiable at zero. Then, we have

$$\partial_B G(\overline{X}) = \partial_B \Psi(0)$$
 and $\partial G(\overline{X}) = \partial \Psi(0)$.

Proof. We only need to prove the result for the B-subdifferentials. Let \mathcal{V} be any element of $\partial_B G(\overline{X})$. Then, there exists a sequence $\{X^k\}$ in \mathcal{D}_G converging to \overline{X} such that $\mathcal{V} = \lim_{k \to \infty} G'(X^k)$. For each X^k , let $U^k \in \mathbb{O}^m$ and $V^k \in \mathbb{O}^n$ be the matrices such that

$$X^k = U^k [\Sigma(X^k) \quad 0] (V^k)^{\mathbb{T}}.$$

For each X^k , denote $\sigma^k = \sigma(X^k)$. Then, we know from Theorem 3.3 that for each $k, \sigma_k \in \mathcal{D}_g$. For k sufficiently large, we know from Lemma 3.1 that for each k, G_S is twice continuously differentiable at \overline{X} . Thus, $\lim_{k \to \infty} G'_S(X^k) = G'_S(\overline{X})$. Hence, we have for any $H \in \mathbb{V}^{m \times n}$,

$$\lim_{k \to \infty} G'_S(X^k) H = G'_S(\overline{X}) H = \overline{U} \left[\overline{\mathcal{E}}_1^0 \circ S(\overline{U}^{\mathbb{T}} H \overline{V}_1) + \overline{\mathcal{E}}_2^0 \circ T(\overline{U}^{\mathbb{T}} H \overline{V}_1) - \overline{\mathcal{F}}^0 \circ \overline{U}^{\mathbb{T}} H \overline{V}_2 \right] \overline{V}^{\mathbb{T}}.$$
 (115)

Moreover, we know that the mapping $G_R = G - G_S$ is also differentiable at each X^k for k sufficiently large. Therefore, we have

$$\mathcal{V} = \lim_{k \to \infty} G'(X^k) = G'_S(\overline{X}) + \lim_{k \to \infty} G'_R(X^k).$$
(116)

From the continuity of the singular value function $\sigma(\cdot)$, by taking a subsequence if necessary, we assume that for each X^k and $l, l' \in \{1, \ldots, r\}$, $\sigma_i(X^k) > 0$, $\sigma_i(X^k) \neq \sigma_j(X^k)$ for any $i \in a_l$, $j \in a_{l'}$ and $l \neq l'$. Since $\{U^k\}$ and $\{V^k\}$ are uniformly bounded, by taking subsequences if necessary, we may also assume that $\{U^k\}$ and $\{V^k\}$ converge and denote the limits by $U^{\infty} \in \mathbb{O}^m$ and $V^{\infty} \in \mathbb{O}^n$, respectively. It is clear that $(U^{\infty}, V^{\infty}) \in \mathbb{O}^{m,n}(\overline{X})$. Therefore, we know from Proposition 2.3 that there exist $Q_l \in \mathbb{O}^{|a_l|}$, $l = 1, \ldots, r$, $Q' \in \mathbb{O}^{|b|}$ and $Q'' \in \mathbb{O}^{n-|a|}$ such that $U^{\infty} = \overline{U}M$ and $V^{\infty} = \overline{V}N$, where $M = \text{Diag}(Q_1, \ldots, Q_r, Q') \in \mathbb{O}^m$ and $N = \text{Diag}(Q_1, \ldots, Q_r, Q'') \in \mathbb{O}^n$. Let $H \in \mathbb{V}^{m \times n}$ be arbitrarily given. For each k, denote $\widetilde{H}^k := (U^k)^{\mathbb{T}}HV^k$. Since $\{(U^k, V^k)\} \in \mathbb{O}^{m,n}(X^k)$ converges to $(U^{\infty}, V^{\infty}) \in \mathbb{O}^{m,n}(\overline{X})$, we know that $\lim_{k \to \infty} \widetilde{H}^k = (U^{\infty})^{\mathbb{T}}HV^{\infty}$. For the notational simplicity, we denote $\widetilde{H} := \overline{U}^{\mathbb{T}}H\overline{V}$ and $\widehat{H} := (U^{\infty})^{\mathbb{T}}HV^{\infty}$.

For k sufficiently large, we know from (21) and (54) that for any $H \in \mathbb{V}^{m \times n}$, $G'_R(X^k)H = U^k \Delta^k (V^k)^{\mathbb{T}}$ with

$$\Delta^{k} := \begin{bmatrix} \operatorname{Diag} \left(\Delta_{1}^{k}, \dots, \Delta_{r}^{k} \right) & 0 \\ 0 & \Delta_{r+1}^{k} \end{bmatrix} \in \mathbb{V}^{m \times n},$$

where for each k, $\Delta_l^k = (\mathcal{E}_l(\sigma^k))_{a_l a_l} \circ S(\widetilde{H}_{a_l a_l}^k) + \text{Diag}((\mathcal{C}(\sigma)\text{diag}(S(\widetilde{H}^k)))_{a_l}), l = 1, \ldots, r,$

$$\Delta_{r+1}^{k} = \left[(\mathcal{E}_{1}(\sigma^{k}))_{bb} \circ S(\widetilde{H}_{bb}^{k}) + \operatorname{Diag}((\mathcal{C}(\sigma)\operatorname{diag}(S(\widetilde{H}^{k})))_{b}) + (\mathcal{E}_{2}(\sigma^{k}))_{bb} \circ T(\widetilde{H}_{bb}^{k}) \quad (\mathcal{F}_{2}(\sigma^{k}))_{bc} \circ \widetilde{H}_{bc}^{k} \right]$$

and $\mathcal{E}_1(\sigma^k)$, $\mathcal{E}_2(\sigma^k)$, $\mathcal{F}(\sigma^k)$ and $\mathcal{C}(\sigma^k)$ are defined for σ^k by (50)-(52), respectively. Again, since $\{U^k\}$ and $\{V^k\}$ are uniformly bounded, we know that

$$\lim_{k \to \infty} G'_R(X^k) H = U^{\infty} (\lim_{k \to \infty} \Delta^k) (V^{\infty})^{\mathbb{T}} = \overline{U} M (\lim_{k \to \infty} \Delta^k) N^{\mathbb{T}} \overline{V}^{\mathbb{T}} .$$
(117)

Next, we shall show that $\mathcal{V} \in \partial_B \Psi(0)$. For each k, denote $w^k := \sigma^k - \overline{\sigma} \in \mathbb{R}^m$. Moreover, for each k, we can define $W_l^k := Q_l \operatorname{Diag}(w_{a_l}^k) Q_l^{\mathbb{T}} \in \mathbb{S}^{|a_l|}, l = 1, \ldots, r$ and $W_{r+1}^k := Q'[\operatorname{Diag}(w_b^k) \quad 0]Q''^{\mathbb{T}} \in \mathbb{V}^{|b| \times (n-|a|)}$. Therefore, it is clear that for each k, $W^k := (W_1^k, \ldots, W_l^k, W_{r+1}^k) \in \mathcal{W}$ and $\kappa(W^k) = w^k$. Moreover, since $\lim_{k \to \infty} \sigma^k = \overline{\sigma}$, we know that $\lim_{k \to \infty} W^k = 0$ in \mathcal{W} . From the assumption, we know that $\phi(\cdot) = g'(\overline{\sigma}; \cdot)$ and $d(\cdot)$ are differentiable at each w^k and $\phi'(w^k) = g'(\sigma^k) - d'(w^k)$ for all w^k . Since d is strictly differentiable at zero, it can be checked easily that $\lim_{k \to \infty} d'(w^k) = d'(0) = 0$. By taking a subsequence if necessary, we may assume that $\lim_{k \to \infty} g'(\sigma^k)$ exists. Therefore, we have

$$\lim_{k \to \infty} \phi'(w^k) = \lim_{k \to \infty} g'(\sigma^k) \,. \tag{118}$$

Since Φ is the spectral operator with respect to the mixed symmetric mapping ϕ , from Theorem 5.3 in Section 5¹ we know that Φ is differentiable at $W \in W$ if and only if ϕ is differentiable at $\kappa(W)$. Recall that $\widehat{\Phi} : W \to \mathbb{V}^{m \times n}$ is defined by (36). Then, for k sufficiently large, $\widehat{\Phi}$ is differentiable at W^k . Moreover, for each k, we define the matrix $C^k \in \mathbb{V}^{m \times n}$ by

$$C^{k} = \overline{U} \begin{bmatrix} \operatorname{Diag} \left(W_{1}^{k}, \dots, W_{r}^{k} \right) & 0 \\ 0 & W_{r+1}^{k} \end{bmatrix} \overline{V}^{\mathbb{T}}$$

Then, we know that for k sufficiently large, Ψ is differentiable at C^k and $\lim_{k\to\infty} C^k = 0$ in $\mathbb{V}^{m\times n}$. Thus, we know from (114) that for each k,

$$\Psi'(C^k)H = G'_S(\overline{X})H + \overline{U}\left[\widehat{\Phi}'(W^k)D(H)\right]\overline{V}^{\mathbb{T}} \quad \forall \ H \in \mathbb{V}^{m \times n} ,$$

where $D(H) = \left(S(\hat{H}_{a_1a_1}), \ldots, S(\hat{H}_{a_ra_r}), \hat{H}_{b\bar{a}}\right) \in \mathcal{W}$ is defined by (35) and $\hat{\Phi}'(W^k)D(H)$ can be derived from (128). By comparing with (116) and (117), we know that the conclusion then follows if we show that

$$\lim_{k \to \infty} \Delta^k = \lim_{k \to \infty} M^{\mathbb{T}} \widehat{\Phi}'(W^k) D(H) N \,. \tag{119}$$

 $^{^{1}}$ We could present the results in this subsection after introducing Theorem 5.3 in Section 5. We include it here for the sake of readability and notational convenience.

For any $(i, j) \in \{1, \ldots, m\} \times \{1, \ldots, n\}$, consider the following cases.

Case 1: i = j. It is easy to check that for each k,

$$(\Delta^k)_{ii} = (g'(\sigma^k)h^k)_i \text{ and } \left(M^{\mathbb{T}}\widehat{\Phi}'(W^k)D(H)N\right)_{ii} = (\phi'(w^k)\widehat{h})_i,$$

where $h^k = \left(\operatorname{diag}(S(\widetilde{H}_{aa}^k)), \operatorname{diag}(\widetilde{H}_{bb}^k)\right)$ and $\widehat{h} = \left(\operatorname{diag}(S(\widehat{H}_{aa})), \operatorname{diag}(\widehat{H}_{bb})\right)$. Therefore, we know from (118) that

$$\lim_{k \to \infty} (\Delta^k)_{ii} = \lim_{k \to \infty} (g'(\sigma^k)h^k)_i = \lim_{k \to \infty} (\phi'(w^k)\widehat{h})_i = \lim_{k \to \infty} \left(M^{\mathbb{T}}\widehat{\Phi}'(W^k)D(H)N \right)_{ii}$$

Case 2: $i, j \in a_l$ for some $l \in \{1, ..., r\}, i \neq j$ and $\sigma_i^k \neq \sigma_j^k$ for k sufficiently large. We obtain that for k sufficiently large,

$$\begin{split} (\Delta^k)_{ij} &= \frac{g_i(\sigma^k) - g_j(\sigma^k)}{\sigma_i^k - \sigma_j^k} (S(\widetilde{H}_{a_l a_l}^k))_{ij}, \\ \left(M^{\mathbb{T}} \widehat{\Phi}'(W^k) D(H) N \right)_{ij} &= \frac{\phi_i(w^k) - \phi_j(w^k)}{w_i^k - w_j^k} (S(\widehat{H}_{a_l a_l}))_{ij} \,. \end{split}$$

Since $\overline{\sigma}_i = \overline{\sigma}_j$ and $g_i(\overline{\sigma}) = g_j(\overline{\sigma})$, we know that for k sufficiently large,

$$\frac{g_i(\sigma^k) - g_j(\sigma^k)}{\sigma_i^k - \sigma_j^k} = \frac{g_i(\overline{\sigma} + w^k) - g_j(\overline{\sigma} + w^k)}{w_i^k - w_j^k} = \frac{g_i(\overline{\sigma} + w^k) - g_i(\overline{\sigma}) + g_j(\overline{\sigma}) - g_j(\overline{\sigma} + w^k)}{w_i^k - w_j^k} \\
= \frac{d_i(w^k) - d_j(w^k)}{w_i^k - w_j^k} + \frac{\phi_i(w^k) - \phi_j(w^k)}{w_i^k - w_j^k}.$$
(120)

Therefore, we know from (111) that

which implies $\lim_{k \to \infty} (\Delta^k)_{ij} = \lim_{k \to \infty} \left(M^{\mathbb{T}} \widehat{\Phi}'(W^k) D(H) N \right)_{ij}$. **Case 3:** $i, j \in a_l$ for some $l \in \{1, \dots, r\}, i \neq j$ and $\sigma_i^k = \sigma_j^k$ for k sufficiently large. We have for k sufficiently large,

$$(\Delta^k)_{ij} = \left((g'(\sigma^k))_{ii} - (g'(\sigma^k))_{ij} \right) (S(\widetilde{H}^k_{a_l a_l}))_{ij},$$
$$\left(M^{\mathbb{T}} \widehat{\Phi}'(W^k) D(H) N \right)_{ij} = \left((\phi'(w^k))_{ii} - (\phi'(w^k))_{ij} \right) (S(\widehat{H}_{a_l a_l}))_{ij}$$

Therefore, we obtain from (118) that

$$\lim_{k \to \infty} \left((g'(\sigma^k))_{ii} - (g'(\sigma^k))_{ij} \right) (S(\widetilde{H}^k_{a_l a_l}))_{ij} = \lim_{k \to \infty} \left((\phi'(w^k))_{ii} - (\phi'(w^k))_{ij} \right) (S(\widehat{H}_{a_l a_l}))_{ij}.$$

Thus, we have $\lim_{k\to\infty} (\Delta^k)_{ij} = \lim_{k\to\infty} \left(M^{\mathbb{T}} \widehat{\Phi}'(W^k) D(H) N \right)_{ij}$. **Case 4:** $i, j \in b, i \neq j$ and $\sigma_i^k = \sigma_j^k > 0$ for k sufficiently large. We have for k large,

$$(\Delta^{k})_{ij} = \left((g'(\sigma^{k}))_{ii} - (g'(\sigma^{k}))_{ij} \right) (S(\widetilde{H}_{bb}^{k}))_{ij} + \frac{g_{i}(\sigma^{k}) + g_{j}(\sigma^{k})}{\sigma_{i}^{k} + \sigma_{j}^{k}} (T(\widetilde{H}_{bb}^{k}))_{ij}, \\ \left(M^{\mathbb{T}} \widehat{\Phi}'(W^{k}) D(H) N \right)_{ij} = \left((\phi'(w^{k}))_{ii} - (\phi'(w^{k}))_{ij} \right) (S(\widehat{H}_{bb}))_{ij} + \frac{\phi_{i}(w^{k}) + \phi_{j}(w^{k})}{w_{i}^{k} + w_{j}^{k}} (T(\widehat{H}_{bb}))_{ij}.$$

Since $\overline{\sigma}_i = \overline{\sigma}_i = 0$ and $g_i(\overline{\sigma}) = g_i(\overline{\sigma}) = 0$, we get

$$\frac{g_i(\sigma^k) + g_j(\sigma^k)}{\sigma_i^k + \sigma_j^k} = \frac{d_i(w^k) + d_j(w^k)}{w_i^k + w_j^k} + \frac{\phi_i(w^k) + \phi_j(w^k)}{w_i^k + w_j^k}.$$
(121)

Therefore, we know from (112) and (118) that $\lim_{k \to \infty} (\Delta^k)_{ij} = \lim_{k \to \infty} \left(M^{\mathbb{T}} \widehat{\Phi}'(W^k) D(H) N \right)_{ij}$.

Case 5: $i, j \in b, i \neq j$ and $\sigma_i^k \neq \sigma_j^k$ for k sufficiently large. For large k, we have

$$(\Delta^k)_{ij} = \frac{g_i(\sigma^k) - g_j(\sigma^k)}{\sigma_i^k - \sigma_j^k} (S(\widetilde{H}_{bb}^k))_{ij} + \frac{g_i(\sigma^k) + g_j(\sigma^k)}{\sigma_i^k + \sigma_j^k} (T(\widetilde{H}_{bb}^k))_{ij},$$
$$\left(M^{\mathbb{T}}\widehat{\Phi}'(W^k)D(H)N\right)_{ij} = \frac{\phi_i(w^k) - \phi_j(w^k)}{w_i^k - w_j^k} (S(\widehat{H}_{bb}))_{ij} + \frac{\phi_i(w^k) + \phi_j(w^k)}{w_i^k + w_j^k} (T(\widehat{H}_{bb}))_{ij}$$

Thus, by (120) and (121), we know from (111) and (112) that $\lim_{k \to \infty} (\Delta^k)_{ij} = \lim_{k \to \infty} \left(M^{\mathbb{T}} \widehat{\Phi}'(W^k) D(H) N \right)_{ij}.$ **Case 6:** $i, j \in b, i \neq j$ and $\sigma_i^k = \sigma_j^k = 0$ for k sufficiently large. We know for k large,

$$\begin{split} (\Delta^k)_{ij} &= \left((g'(\sigma^k))_{ii} - (g'(\sigma^k))_{ij} \right) (S(\widehat{H}^k_{bb}))_{ij} + (g'(\sigma^k))_{ii} (T(\widehat{H}^k_{bb}))_{ij}, \\ \left(M^{\mathbb{T}} \widehat{\Phi}'(W^k) D(H) N \right)_{ij} &= \left((\phi'(w^k))_{ii} - (\phi'(w^k))_{ij} \right) (S(\widehat{H}_{bb}))_{ij} + (\phi'(w^k))_{ii} (T(\widehat{H}_{bb}))_{ij}. \end{split}$$

Again, we obtain from (118) that $\lim_{k\to\infty} (\Delta^k)_{ij} = \lim_{k\to\infty} \left(M^{\mathbb{T}} \widehat{\Phi}'(W^k) D(H) N \right)_{ij}.$

Case 7: $i \in b, j \in c$ and $\sigma_i^k > 0$ for k sufficiently large. We have for k sufficiently large,

$$(\Delta^k)_{ij} = \frac{g_i(\sigma^k)}{\sigma_i^k} (\widetilde{H}_{bc}^k)_{ij}, \quad \left(M^{\mathbb{T}}\widehat{\Phi}'(W^k)D(H)N\right)_{ij} = \frac{\phi_i(w^k)}{w_i^k} (\widehat{H}_{bc})_{ij}$$

Since $\overline{\sigma}_i = 0$ and $g_i(\overline{\sigma}) = 0$, we get

$$\frac{g_i(\sigma^k)}{\sigma_i^k} = \frac{g_i(\overline{\sigma} + w^k) - g_i(\overline{\sigma})}{w_i^k} = \frac{d_i(w^k)}{w_i^k} + \frac{\phi_i(w^k)}{w_i^k}.$$

Therefore, by (113), we obtain that $\lim_{k \to \infty} (\Delta^k)_{ij} = \lim_{k \to \infty} \left(M^{\mathbb{T}} \widehat{\Phi}'(W^k) D(H) N \right)_{ij}$.

Case 8: $i \in b, j \in c$ and $\sigma_i^k = 0$ for k sufficiently large. We have for k sufficiently large,

$$(\Delta^k)_{ij} = (g'(\sigma^k))_{ii}(\widetilde{H}^k_{bc})_{ij}, \quad \left(M^{\mathbb{T}}\widehat{\Phi}'(W^k)D(H)N\right)_{ij} = (\phi'(w^k))_{ii}(\widehat{H}_{bc})_{ij}.$$

Therefore, by (118), we obtain that $\lim_{k \to \infty} (\Delta^k)_{ij} = \lim_{k \to \infty} \left(M^{\mathbb{T}} \widehat{\Phi}'(W^k) D(H) N \right)_{ij}$.

Thus, we know that (119) holds. Therefore, by (116) and (117), we obtain that $\mathcal{V} \in \partial_B \Psi(0)$.

Conversely, suppose that $\mathcal{V} \in \partial_B \Psi(0)$ is arbitrarily chosen. Then, from the definition of $\partial_B \Psi(0)$, we know that there exists a sequence $\{C^k\} \subseteq \mathbb{V}^{m \times n}$ converging to zero such that Ψ is differentiable at each C^k and $\mathcal{V} = \lim_{k \to \infty} \Psi'(C^k)$. For each k, we know from (114) that Ψ is differentiable at C^k if and only if the spectral operator $\Phi: \mathcal{W} \to \mathcal{W}$ is differentiable at $W^k := D(C^k) = \left(S(\widetilde{C}_{a_1a_1}^k), \dots, S(\widetilde{C}_{a_ra_r}^k), \widetilde{C}_{b\overline{a}}^k\right) \in \mathcal{W}$, where for each $k, \ \widetilde{C}^k = \overline{U}^{\mathbb{T}} C^k \overline{V}$. Moreover, for each k, we have the following decompositions

$$S(\widetilde{C}_{a_la_l}^k) = Q_l^k \Lambda(S(\widetilde{C}_{a_la_l}^k))(Q_l^k)^{\mathbb{T}}, \ l = 1, \dots, r, \quad \widetilde{C}_{b\bar{a}}^k = {Q'}^k \begin{bmatrix} \Sigma(\widetilde{C}_{b\bar{a}}^k) & 0 \end{bmatrix} ({Q''}^k)^{\mathbb{T}},$$

where $Q_l^k \in \mathbb{O}^{|a_l|}, Q'^k \in \mathbb{O}^{|b|}$ and $Q''^k \in \mathbb{O}^{n-|a|}$. For each k, let

$$w^{k} := \left(\lambda(S(\widetilde{C}_{a_{1}a_{1}}^{k})), \dots, \lambda(S(\widetilde{C}_{a_{r}a_{r}}^{k})), \sigma(\widetilde{C}_{b\bar{a}}^{k})\right) \in \mathbb{R}^{m},$$
$$M^{k} := \operatorname{Diag}\left(Q_{1}^{k}, \dots, Q_{r}^{k}, Q'^{k}\right) \in \mathbb{O}^{m}, \quad N^{k} := \operatorname{Diag}\left(Q_{1}^{k}, \dots, Q_{r}^{k}, Q''^{k}\right) \in \mathbb{O}^{n}.$$

Since $\{M^k\}$ and $\{N^k\}$ are uniformly bounded, by taking subsequences if necessary, we know that there exist $Q_l \in \mathbb{O}^{|a_l|}, Q' \in \mathbb{O}^{|b|}$ and $Q'' \in \mathbb{O}^{n-|b|}$ such that

$$\lim_{k \to \infty} M^k = M := \operatorname{Diag}\left(Q_1, \dots, Q_r, Q'\right) \quad \lim_{k \to \infty} N^k = N := \operatorname{Diag}\left(Q_1, \dots, Q_r, Q''\right).$$

For each k, by (128) (in Section 5), we know that for any $H \in \mathbb{V}^{m \times n}$,

$$\Psi'(C^k)H = \overline{U}\left[\overline{\mathcal{E}}_1^0 \circ S(\overline{U}^{\mathbb{T}}H\overline{V}_1) + \overline{\mathcal{E}}_2^0 \circ T(\overline{U}^{\mathbb{T}}H\overline{V}_1) \quad \overline{\mathcal{F}}^0 \circ \overline{U}^{\mathbb{T}}H\overline{V}_2\right]\overline{V}^{\mathbb{T}} + \overline{U}\left[\widehat{\Phi}'(W^k)D(H)\right]\overline{V}^{\mathbb{T}}, \quad (122)$$

where $D(H) \in \mathcal{W}$ is defined by (35). Let $R^k := \Phi'_k(W^k)D(H), k = 1, \dots, r+1$.

For each k, define $\sigma^k := \overline{\sigma} + w^k \in \mathbb{R}^m$. Since $\lim_{k \to \infty} w^k = 0$ and for each k, $w_i^k \ge 0$ for all $i \in b$, we have $\sigma^k \ge 0$ for k sufficiently large. Therefore, for k sufficiently large, we are able to define

$$X^k := \overline{U}M[\operatorname{Diag}(\sigma^k) \quad 0]N^{\mathbb{T}}\overline{V}^{\mathbb{T}} \in \mathbb{V}^{m \times n} \,.$$

For simplicity, denote $U = \overline{U}M \in \mathbb{O}^m$ and $V = \overline{V}N \in \mathbb{O}^n$. It is clear that the sequence $\{X^k\}$ converges to \overline{X} . From the assumption, we know that g is differentiable at each σ^k and d is differentiable at each w^k with $g'(\sigma^k) = \phi'(w^k) + d'(w^k)$ for all σ^k . Therefore, by Theorem 3.3, we know that G is differentiable at each X^k . By taking subsequences if necessary, we may assume that $\lim_{k\to\infty} \phi'(w^k)$ exists. Thus, since d is strictly differentiable at zero, we know that (118) holds. Since the derivative formula (28) is independent of $(U, V) \in \mathbb{O}^{m,n}(\overline{X})$, we know from (54) that for any $H \in \mathbb{V}^{m \times n}$,

$$G'(X^{k})H = \overline{U} \begin{bmatrix} \overline{\mathcal{E}}_{1}^{0} \circ S(\overline{U}^{\mathbb{T}}H\overline{V}_{1}) + \overline{\mathcal{E}}_{2}^{0} \circ T(\overline{U}^{\mathbb{T}}H\overline{V}_{1}) & \overline{\mathcal{F}}^{0} \circ \overline{U}^{\mathbb{T}}H\overline{V}_{2} \end{bmatrix} \overline{V}^{\mathbb{T}} + \overline{U} \begin{bmatrix} \operatorname{Diag}\left(Q_{1}\Omega_{1}^{k}Q_{1}^{\mathbb{T}}, \dots, Q_{r}\Omega_{r}^{k}Q_{r}^{\mathbb{T}}\right) & 0 \\ 0 & Q'\Omega_{r+1}^{k}Q''^{\mathbb{T}} \end{bmatrix} \overline{V}^{\mathbb{T}},$$
(123)

where for each k, $\Omega_l^k = (\mathcal{E}_l(\sigma^k))_{a_l a_l} \circ S(\widehat{H}_{a_l a_l}) + \text{Diag}((\mathcal{C}(\sigma^k)\text{diag}(S(\widehat{H})))_{a_l}), l = 1, \ldots, r \text{ and}$

$$\Omega_{r+1}^{k} = \left[(\mathcal{E}_{1}(\sigma^{k}))_{bb} \circ S(\widehat{H}_{bb}) + \operatorname{Diag}((\mathcal{C}(\sigma^{k})\operatorname{diag}(S(\widehat{H})))_{b}) + (\mathcal{E}_{2}(\sigma^{k}))_{bb} \circ T(\widehat{H}_{bb}) \quad (\mathcal{F}_{2}(\sigma^{k}))_{bc} \circ \widehat{H}_{bc} \right] ,$$

 $\mathcal{E}_1(\sigma^k)$, $\mathcal{E}_2(\sigma^k)$ and $\mathcal{F}(\sigma^k)$ are defined by (50)–(52), respectively and $\widehat{H} := M^{\mathbb{T}}\overline{U}^{\mathbb{T}}H\overline{V}N = M^{\mathbb{T}}\widetilde{H}N$. Therefore, by comparing (122) and (123), we know that the inclusion $\mathcal{V} \in \partial_B G(\overline{X})$ follows if we show that

$$\lim_{k \to \infty} \left(R_1^k, \dots, R_r^k, R_{r+1}^k \right) = \lim_{k \to \infty} \left(Q_1 \Omega_1^k Q_1^{\mathbb{T}}, \dots, Q_r \Omega_r^k Q_r^{\mathbb{T}}, Q' \Omega_{r+1}^k Q''^{\mathbb{T}} \right) \,. \tag{124}$$

Similarly to the proofs to **Cases 1-8** in the first part, by using (118) and (111)–(113) in Lemma 4.2, we can show that (124) holds. For simplicity, we omit the details here. Therefore, we obtain that $\partial_B G(\overline{X}) = \partial_B \Psi(0)$. This completes the proof.

5 Extensions

In this section, we consider the spectral operators defined on the Cartesian product of several real or complex matrices. The corresponding properties, including continuity, directional differentiability, (continuous) differentiability, locally Lipschitzian continuity, ρ -order B-differentiability, ρ -order G-semismoothness and the characterization of Clarke's generalized Jacobian, can be studied in the same fashion as those in Section 3 and Section 4. Instead of presenting the proofs here, we refer the readers to the PhD thesis of Ding [15] to work out details.

Without loss of generality, from now on, we assume that $\mathcal{X} = \mathbb{S}^{m_1} \times \mathbb{V}^{m_2 \times n_2}$ with $m = m_1 + m_2$. For any $X = (X_1, X_2) \in \mathbb{S}^{m_1} \times \mathbb{V}^{m_2 \times n_2}$, denote $\kappa(X) = (\lambda(X_1), \sigma(X_2))$. Let \mathcal{N} be a given nonempty open set in \mathcal{X} . Suppose that $\mathbf{g} : \mathbb{R}^m \to \mathbb{R}^m$ is mixed symmetric, with respect to $\mathcal{P} \equiv \mathbb{P}^{m_1} \times \pm \mathbb{P}^{m_2}$, on an open set $\hat{\kappa}_{\mathcal{N}}$ in \mathbb{R}^m containing $\kappa_{\mathcal{N}} = \{\kappa(X) \mid X \in \mathcal{N}\}$. Let $G : \mathcal{X} \to \mathcal{X}$ be the corresponding spectral operator defined in Definition 2.2.

Let $\overline{X} = (\overline{X}_1, \overline{X}_2) \in \mathcal{N}$ be given. Suppose the given $\overline{X}_1 \in \mathbb{S}^{m_1}$ and $\overline{X}_2 \in \mathbb{V}^{m_2 \times n_2}$ have the following decompositions

$$\overline{X}_1 = \overline{P} \operatorname{Diag}(\lambda(\overline{X}_1)) \overline{P}^{\mathbb{T}} \quad \text{and} \quad \overline{X}_2 = \overline{U} [\operatorname{Diag}(\sigma(\overline{X}_2)) \quad 0] \overline{V}^{\mathbb{T}},$$
(125)

where $\overline{P} \in \mathbb{O}^{m_1}$, $\overline{U} \in \mathbb{O}^{m_2}$ and $\overline{V} = [\overline{V}_1 \quad \overline{V}_2] \in \mathbb{O}^{n_2}$ with $\overline{V}_1 \in \mathbb{V}^{n_2 \times m_2}$ and $\overline{V}_2 \in \mathbb{V}^{n_2 \times (n_2 - m_2)}$. Denote $\overline{\lambda} := \lambda(\overline{X}_1)$, $\overline{\sigma} := \sigma(\overline{X}_2)$ and $\overline{\kappa} := (\overline{\lambda}, \overline{\sigma})$. We use $\overline{\nu}_1 > \ldots > \overline{\nu}_{r_1}$ to denote the distinct eigenvalues of \overline{X}_1 and $\overline{\nu}_{r_1+1} > \ldots > \overline{\nu}_{r_1+r_2} > 0$ to denote the distinct nonzero singular values of \overline{X}_2 . Define the index sets

$$\begin{cases} a_l := \{i \,|\, \overline{\lambda}_i = \overline{\nu}_l, \ 1 \le i \le m_1\} & l = 1, \dots, r_1, \\ a_l := \{i \,|\, \overline{\sigma}_i = \overline{\nu}_l, \ 1 \le i \le m_2\} & l = r_1 + 1, \dots, r_1 + r_2. \end{cases}$$

Define $b := \{i \mid \overline{\sigma}_i = 0, 1 \le i \le m_2\}$. We have the following result on the continuity of spectral operators.

Theorem 5.1 Let $\overline{X} = (\overline{X}_1, \overline{X}_2) \in \mathcal{N}$ be given. Suppose that \overline{X}_1 and \overline{X}_2 have the decompositions (125). The spectral operator G is continuous at \overline{X} if and only if **g** is continuous at $\kappa(\overline{X})$.

In order to present the results on the directional differentiability of spectral operators of matrices, we introduce some notations. For the given mixed symmetric mapping $\mathbf{g} = (\mathbf{g}_1, \mathbf{g}_2) : \mathbb{R}^m \to \mathbb{R}^{m_1} \times \mathbb{R}^{m_2}$, define the matrices $\mathcal{A}^0(\overline{\kappa}) \in \mathbb{S}^{m_1}$, $\mathcal{E}_1^0(\overline{\kappa}) \in \mathbb{S}^{m_2}$, $\mathcal{E}_2^0(\overline{\kappa}) \in \mathbb{V}^{m_2 \times m_2}$ and $\mathcal{F}^0(\overline{\kappa}) \in \mathbb{V}^{m_2 \times (n_2 - m_2)}$ with respect to $\overline{\kappa} = (\overline{\lambda}, \overline{\sigma})$ by

$$\begin{split} (\mathcal{A}^{0}(\overline{\kappa}))_{ij} &:= \begin{cases} \frac{(\mathbf{g}_{1}(\overline{\kappa}))_{i} - (\mathbf{g}_{1}(\overline{\kappa}))_{j}}{\overline{\lambda}_{i} - \overline{\lambda}_{j}} & \text{if } \overline{\lambda}_{i} \neq \overline{\lambda}_{j}, \\ 0 & \text{otherwise,} \end{cases} \quad i, j \in \{1, \dots, m_{1}\}, \\ (\mathcal{E}_{1}^{0}(\overline{\kappa}))_{ij} &:= \begin{cases} \frac{(\mathbf{g}_{2}(\overline{\kappa}))_{i} - (\mathbf{g}_{2}(\overline{\kappa}))_{j}}{\overline{\sigma}_{i} - \overline{\sigma}_{j}} & \text{if } \overline{\sigma}_{i} \neq \overline{\sigma}_{j}, \\ 0 & \text{otherwise,} \end{cases} \quad i, j \in \{1, \dots, m_{2}\}, \\ (\mathcal{E}_{2}^{0}(\overline{\kappa}))_{ij} &:= \begin{cases} \frac{(\mathbf{g}_{2}(\overline{\kappa}))_{i} + (\mathbf{g}_{2}(\overline{\kappa}))_{j}}{\overline{\sigma}_{i} + \overline{\sigma}_{j}} & \text{if } \overline{\sigma}_{i} + \overline{\sigma}_{j} \neq 0, \\ 0 & \text{otherwise,} \end{cases} \quad i, j \in \{1, \dots, m_{2}\} \end{split}$$

and

$$(\mathcal{F}^0(\overline{\kappa}))_{ij} := \begin{cases} \frac{(\mathbf{g}_2(\overline{\kappa}))_i}{\overline{\sigma}_i} & \text{if } \overline{\sigma}_i \neq 0, \\ 0 & \text{otherwise.} \end{cases} \quad i \in \{1, \dots, m_2\}, \quad j \in \{1, \dots, n_2 - m_2\}. \end{cases}$$

Suppose that **g** is directionally differentiable at $\overline{\kappa}$. Then, we know that the directional derivative $\mathbf{g}'(\overline{\kappa}; \cdot) = (\mathbf{g}'_1(\overline{\kappa}; \cdot), \mathbf{g}'_2(\overline{\kappa}; \cdot)) : \mathbb{R}^{m_1+m_2} \to \mathbb{R}^{m_1+m_2}$ satisfies that for any $(Q_1, Q_2) \in \mathcal{P}_{\overline{\kappa}}$ and any $(\mathbf{h}_1, \mathbf{h}_2) \in \mathbb{R}^{m_1} \times \mathbb{R}^{m_2}$,

$$\left(\mathbf{g}_{1}'(\overline{\kappa};(Q_{1}\mathbf{h}_{1},Q_{2}\mathbf{h}_{2})),\mathbf{g}_{2}'(\overline{\kappa};(Q_{1}\mathbf{h}_{1},Q_{2}\mathbf{h}_{2}))\right) = \left(Q_{1}\mathbf{g}_{1}'(\overline{\kappa};(\mathbf{h}_{1},\mathbf{h}_{2})),Q_{2}\mathbf{g}_{2}'(\overline{\kappa};(\mathbf{h}_{1},\mathbf{h}_{2}))\right),\tag{126}$$

where $\mathcal{P}_{\overline{\kappa}}$ is the subset of $\mathcal{P} \equiv \mathbb{P}^{m_1} \times \pm \mathbb{P}^{m_2}$ defined with respect to $\overline{\kappa}$ by

$$\mathcal{P}_{\overline{\kappa}} := \left\{ (Q_1, Q_2) \in \mathbb{P}^{m_1} \times \pm \mathbb{P}^{m_2} \mid (\overline{\lambda}, \overline{\sigma}) = (Q_1 \overline{\lambda}, Q_2 \overline{\sigma}) \right\}$$

It is easy to check that $(Q_1, Q_2) \in \mathcal{P}_{\overline{\kappa}}$ if and only if there exist $Q_1^l \in \mathbb{P}^{|a_l|}$, $l = 1, \ldots, r_1, Q_2^l \in \mathbb{P}^{|a_l|}$, $l = r_1 + 1, \ldots, r_1 + r_2$ and $Q_2^{r_1 + r_2 + 1} \in \pm \mathbb{P}^{|b|}$ such that

$$Q_1 = \text{Diag}\left(Q_1^1, \dots, Q_1^{r_1}\right) \in \mathbb{P}^{m_1} \quad \text{and} \quad Q_2 = \text{Diag}\left(Q_2^{r_1+1}, \dots, Q_2^{r_1+r_2}, Q_2^{r_1+r_2+1}\right) \in \pm \mathbb{P}^{m_2}.$$
 (127)

Denote $\phi(\cdot) := \mathbf{g}'(\overline{\kappa}; \cdot)$. For any $h \in \mathbb{R}^m$, rewrite $\phi(h) \in \mathbb{R}^m$ as $\phi(h) = (\phi_1(h), \dots, \phi_{r_1+r_2}(h), \phi_{r_1+r_2+1}(h))$ with $\phi_l(h) \in \mathbb{R}^{|a_l|}$ for $l = 1, \dots, r_1 + r_2$ and $\phi_{r_1+r_2+1}(h) \in \mathbb{R}^{|b|}$. Therefore, we know from (126) and (127) that the directional derivative ϕ is mixed symmetric mapping, with respect to $\mathbb{P}^{|a_1|} \times \ldots \times \mathbb{P}^{|a_{r_1+r_2}|} \times \pm \mathbb{P}^{|b|}$, over $\mathbb{R}^{|a_1|} \times \ldots \times \mathbb{R}^{|a_{r_1+r_2}|} \times \mathbb{R}^{|b|}$. Denote

$$\mathcal{W} := \mathbb{S}^{|a_1|} \times \ldots \times \mathbb{S}^{|a_{r_1+r_2}|} \times \mathbb{V}^{|b| \times (|b|+n_2-m_2)}.$$

Let $\Phi : \mathcal{W} \to \mathcal{W}$ be the corresponding spectral operator defined in Definition 2.2 with respect to the mixed symmetric mapping ϕ , i.e., for any $W = (W_1, \ldots, W_{r_1+r_2}, W_{r_1+r_2+1}) \in \mathcal{W}$,

$$\Phi(W) = (\Phi_1(W), \dots, \Phi_{r_1+r_2}(W), \Phi_{r_1+r_2+1}(W))$$

with

$$\Phi_l(W) = \begin{cases} \widetilde{R}_l \operatorname{Diag}(\phi_l(\kappa(W))) \widetilde{R}_l^{\mathbb{T}} & \text{if } l = 1, \dots, r_1 + r_2, \\ \widetilde{M} \operatorname{Diag}(\phi_{r_1 + r_2 + 1}(\kappa(W))) \widetilde{N}_1^{\mathbb{T}} & \text{if } l = r_1 + r_2 + 1, \end{cases}$$

where $\kappa(W) = (\lambda(W_1), \dots, \lambda(W_{r_1+r_2}), \sigma(W_{r_1+r_2+1})) \in \mathbb{R}^m$, $\widetilde{R}_l \in \mathbb{O}^{|a_l|}(W_l)$, and $(\widetilde{M}, \widetilde{N}) \in \mathbb{O}^{|b|, |b|+n_2-m_2}(W_{r_1+r_2+1})$, $\widetilde{N} = \begin{bmatrix} \widetilde{N}_1 & \widetilde{N}_2 \end{bmatrix}$ with $\widetilde{N}_1 \in \mathbb{V}^{(|b|+n_2-m_2) \times |b|}$, $\widetilde{N}_2 \in \mathbb{V}^{(|b|+n_2-m_2) \times (n_2-m_2)}$. Then, the first divided directional difference $\mathbf{g}^{[1]}(\overline{X}; H) \in \mathcal{X}$ of \mathbf{g} at \overline{X} along the direction $H = (H_1, H_2) \in \mathcal{X}$ is defined by

$$\mathbf{g}^{[1]}(\overline{X};H) := \left(\mathbf{g}_1^{[1]}(\overline{X};H), \ \mathbf{g}_2^{[1]}(\overline{X};H)\right)$$

with

$$\begin{aligned} \mathbf{g}_{1}^{[1]}(\overline{X};H) &= \mathcal{A}^{0}(\overline{\kappa}) \circ \overline{P}^{\mathbb{T}} H_{1}\overline{P} + \operatorname{Diag}\left(\Phi_{1}(D(H)), \dots, \Phi_{r_{1}}(D(H))\right) \in \mathbb{S}^{m_{1}}, \\ \mathbf{g}_{2}^{[1]}(\overline{X};H) &= \begin{bmatrix} \mathcal{E}_{1}^{0}(\overline{\kappa}) \circ S(\overline{U}^{\mathbb{T}} H_{2}\overline{V}_{1}) + \mathcal{E}_{2}^{0}(\overline{\kappa}) \circ T(\overline{U}^{\mathbb{T}} H_{2}\overline{V}_{1}) & \mathcal{F}^{0}(\overline{\kappa}) \circ \overline{U}^{\mathbb{T}} H_{2}\overline{V}_{2} \end{bmatrix} \\ &+ \begin{bmatrix} \operatorname{Diag}\left(\Phi_{r_{1}+1}(D(H)), \dots, \Phi_{r_{1}+r_{2}}(D(H))\right) & 0 \\ 0 & \Phi_{r_{1}+r_{2}+1}(D(H)) \end{bmatrix} \in \mathbb{V}^{m_{2} \times n_{2}}, \end{aligned}$$

where

$$D(H) = \left(\overline{P}_{a_1}^{\mathbb{T}} H_1 \overline{P}_{a_1}, \dots, \overline{P}_{a_{r_1}}^{\mathbb{T}} H_1 \overline{P}_{a_{r_1}}, S(\overline{U}_{a_{r_1+1}}^{\mathbb{T}} H_2 \overline{V}_{a_{r_1+1}}), \dots, S(\overline{U}_{a_{r_1+r_2}}^{\mathbb{T}} H_2 \overline{V}_{a_{r_1+r_2}}), \overline{U}_b^{\mathbb{T}} H_2 [\overline{V}_b \quad \overline{V}_2]\right) \in \mathcal{W}.$$

Now, we are ready to state the results on the directional differentiability of the spectral operator G.

Theorem 5.2 Let $\overline{X} = (\overline{X}_1, \overline{X}_2) \in \mathcal{N}$ be given. Suppose that \overline{X}_1 and \overline{X}_2 have the decompositions (125). The spectral operator G is Hadamard directionally differentiable at \overline{X} if and only if \mathbf{g} is Hadamard directionally differentiable at \overline{X} and the directional directional directional directional directional directional directional directional directional direction $H \in \mathcal{X}$ is given by

$$G'(\overline{X};H) = \left(\overline{P}\mathbf{g}_1^{[1]}(\overline{X};H)\overline{P}^{\mathbb{T}}, \ \overline{U}\mathbf{g}_2^{[1]}(\overline{X};H)\overline{V}^{\mathbb{T}}\right).$$

In order to present the derivative formulas of spectral operators, we introduce the following notations. For the given $\overline{X} = (\overline{X}_1, \overline{X}_2) \in \mathcal{N}$, suppose that **g** is F-differentiable at $\overline{\kappa} = \kappa(\overline{X})$. Denote by $\mathbf{g}'(\overline{\kappa}) \in \mathbb{R}^{m \times m}$ the Jacobian matrix of **g** at $\overline{\kappa}$. Let $\eta_1(\overline{\kappa}) \in \mathbb{R}^{m_1}$ and $\eta_2(\overline{\kappa}) \in \mathbb{R}^{m_2}$ be the vectors defined by

$$(\eta_1(\overline{\kappa}))_i := \begin{cases} (\mathbf{g}_1'(\overline{\kappa}))_{ii} - (\mathbf{g}_1'(\overline{\kappa}))_{i(i+1)} & \text{if } \exists j \in \{1, \dots, m_1\} \text{ and } j \neq i \text{ such that } \overline{\lambda}_i = \overline{\lambda}_j, \\ (\mathbf{g}_1'(\overline{\kappa}))_{ii} & \text{otherwise}, \end{cases} \quad i \in \{1, \dots, m_1\}$$

and

$$(\eta_2(\overline{\kappa}))_i := \begin{cases} (\mathbf{g}_2'(\overline{\kappa}))_{ii} - (\mathbf{g}_2'(\overline{\kappa}))_{i(i+1)} & \text{if } \exists j \in \{1, \dots, m_2\} \text{ and } j \neq i \text{ such that } \overline{\sigma}_i = \overline{\sigma}_j, \\ (\mathbf{g}_2'(\overline{\kappa}))_{ii} & \text{otherwise}, \end{cases} \quad i \in \{1, \dots, m_2\}$$

Define the corresponding divided difference matrices $\mathcal{A}(\overline{\kappa}) \in \mathbb{R}^{m_1 \times m_1}$ and $\mathcal{E}_1(\overline{\kappa}) \in \mathbb{R}^{m_2 \times m_2}$, the divided addition matrix $\mathcal{E}_2(\overline{\kappa}) \in \mathbb{R}^{m_2 \times m_2}$, the division matrix $\mathcal{F}(\overline{\kappa}) \in \mathbb{R}^{m_2 \times (n_2 - m_2)}$, respectively, by

$$\begin{split} (\mathcal{A}(\overline{\kappa}))_{ij} &:= \begin{cases} \frac{(\mathbf{g}_1(\overline{\kappa}))_i - (\mathbf{g}_1(\overline{\kappa}))_j}{\overline{\lambda}_i - \overline{\lambda}_j} & \text{if } \overline{\lambda}_i \neq \overline{\lambda}_j, \\ (\eta_1(\overline{\kappa}))_i & \text{otherwise,} \end{cases} & i, j \in \{1, \dots, m_1\}, \\ (\mathcal{E}_1(\overline{\kappa}))_{ij} &:= \begin{cases} \frac{(\mathbf{g}_2(\overline{\kappa}))_i - (\mathbf{g}_2(\overline{\kappa}))_j}{\overline{\sigma}_i - \overline{\sigma}_j} & \text{if } \overline{\sigma}_i \neq \overline{\sigma}_j, \\ (\eta_2(\overline{\kappa}))_i & \text{otherwise,} \end{cases} & i, j \in \{1, \dots, m_2\}, \\ (\mathcal{E}_2(\overline{\kappa}))_{ij} &:= \begin{cases} \frac{(\mathbf{g}_2(\overline{\kappa}))_i + (\mathbf{g}_2(\overline{\kappa}))_j}{\overline{\sigma}_i + \overline{\sigma}_j} & \text{if } \overline{\sigma}_i + \overline{\sigma}_j \neq 0, \\ (\mathbf{g}_2'(\overline{\kappa}))_{ii} & \text{otherwise,} \end{cases} & i, j \in \{1, \dots, m_2\}, \\ (\mathcal{F}(\overline{\kappa}))_{ij} &:= \begin{cases} \frac{(\mathbf{g}_2(\overline{\kappa}))_i}{\overline{\sigma}_i} & \text{if } \overline{\sigma}_i \neq 0, \\ (\mathbf{g}_2'(\overline{\kappa}))_{ii} & \text{otherwise,} \end{cases} & i \in \{1, \dots, m_2\}, \quad j \in \{1, \dots, m_2\}. \end{split}$$

Define the matrices $\mathcal{C}_1(\overline{\kappa}) \in \mathbb{R}^{m_1 \times m}$ and $\mathcal{C}_2(\overline{\kappa}) \in \mathbb{R}^{m_2 \times m}$ by

$$C_1(\overline{\kappa}) = \mathbf{g}_1'(\overline{\kappa}) - [\operatorname{Diag}(\eta_1(\overline{\kappa})) \quad 0] \text{ and } C_2(\overline{\kappa}) = \mathbf{g}_2'(\overline{\kappa}) - [0 \operatorname{Diag}(\eta_2(\overline{\kappa}))].$$

Then, we have the following results on the F-differentiability of spectral operators.

Theorem 5.3 Let $\overline{X} = (\overline{X}_1, \overline{X}_2) \in \mathcal{N}$ be given. Suppose that \overline{X}_1 and \overline{X}_2 have the decompositions (125). The spectral operator G is (continuously) differentiable at \overline{X} if and only if \mathbf{g} is (continuously) differentiable at $\overline{K} = \kappa(\overline{X})$. In that case, the derivative of G at \overline{X} is given by for any $H = (H_1, H_2) \in \mathcal{X}$,

$$G'(\overline{X})(H) = \left(\overline{P}[\mathcal{A}(\overline{\kappa}) \circ \overline{P}^{\mathbb{T}} H_1 \overline{P} + \text{Diag}\left(\mathcal{C}_1(\overline{\kappa})h\right)]\overline{P}^{\mathbb{T}}, \\ \overline{U}\left[\mathcal{E}_1(\overline{\kappa}) \circ S(\overline{U}^{\mathbb{T}} H_2 \overline{V}_1) + \text{Diag}\left(\mathcal{C}_2(\overline{\kappa})h\right) + \mathcal{E}_2(\overline{\kappa}) \circ T(\overline{U}^{\mathbb{T}} H_2 \overline{V}_1) \quad \mathcal{F}(\overline{\kappa}) \circ \overline{U}^{\mathbb{T}} H_2 \overline{V}_2\right] \overline{V}^{\mathbb{T}}\right),$$

$$(128)$$

where $h := (\operatorname{diag}(\overline{P}^{\mathbb{T}}H_1\overline{P}), \operatorname{diag}(S(\overline{U}^{\mathbb{T}}H_2\overline{V}_1))) \in \mathbb{R}^m$.

The following theorem is on the locally Lipschitzian continuity of spectral operators of matrices.

Theorem 5.4 Let $\overline{X} = (\overline{X}_1, \overline{X}_2) \in \mathcal{N}$ be given. Suppose that \overline{X}_1 and \overline{X}_2 have the decompositions (125). Then, the spectral operator G is locally Lipschitz continuous near \overline{X} if and only if \mathbf{g} is locally Lipschitz continuous near $\kappa(\overline{X})$.

For the ρ -order B(ouligand)-differentiability of spectral operators, we have the following theorem.

Theorem 5.5 Let $\overline{X} = (\overline{X}_1, \overline{X}_2) \in \mathcal{N}$ be given. Suppose that \overline{X}_1 and \overline{X}_2 have the decompositions (125). Let $0 < \rho \leq 1$ be given. Then,

- (i) if **g** is locally Lipschitz continuous near $\kappa(\overline{X})$ and ρ -order B-differentiable at $\kappa(\overline{X})$, then G is ρ -order B-differentiable at \overline{X} ;
- (ii) if G is ρ -order B-differentiable at \overline{X} , then **g** is ρ -order B-differentiable at $\kappa(\overline{X})$.

Suppose that **g** is locally Lipschitz continuous near $\kappa(\overline{X})$. Thus, we know from Theorem 5.4 that the corresponding spectral operator G is also locally Lipschitz continuous near \overline{X} . We have the following theorem on the G-semismoothness of spectral operators.

Theorem 5.6 Let $\overline{X} = (\overline{X}_1, \overline{X}_2) \in \mathcal{N}$ be given. Suppose that \overline{X}_1 and \overline{X}_2 have the decompositions (125). Let $0 < \rho \leq 1$ be given. Then, G is ρ -order G-semismooth at \overline{X} if and only if \mathbf{g} is ρ -order G-semismooth at $\kappa(\overline{X})$.

Finally, we assume that **g** is locally Lipschitz continuous near $\overline{\kappa} = \kappa(\overline{X})$ and directionally differentiable at $\overline{\kappa}$. From Theorems 5.2 and Theorem 5.4, the spectral operator G is also locally Lipschitz continuous near \overline{X} and directionally differentiable at \overline{X} . Then, we have the following results on the characterization of the B-subdifferential $\partial_B G(\overline{X})$ and Clarke's subdifferential $\partial G(\overline{X})$.

Theorem 5.7 Let $\overline{X} = (\overline{X}_1, \overline{X}_2) \in \mathcal{N}$ be given. Suppose that \overline{X}_1 and \overline{X}_2 have the decompositions (125). Suppose that there exists an open neighborhood $\mathcal{B} \subseteq \mathbb{R}^m$ of $\overline{\kappa}$ in $\hat{\kappa}_{\mathcal{N}}$ such that $\mathbf{g}(\cdot)$ is differentiable at $\kappa \in \mathcal{B}$ if and only if $\phi = \mathbf{g}'(\overline{\kappa}; \cdot)$ is differentiable at $\kappa - \overline{\kappa}$. Assume that the function $d : \mathbb{R}^m \to \mathbb{R}^m$ defined by

$$d(h) = \mathbf{g}(\overline{\kappa} + h) - \mathbf{g}(\overline{\kappa}) - \mathbf{g}'(\overline{\kappa}; h), \quad h \in \mathbb{R}^m$$

is strictly differentiable at zero. Then, we have

$$\partial_B G(\overline{X}) = \partial_B \Psi(0)$$
 and $\partial G(\overline{X}) = \partial \Psi(0)$,

where $\Psi(\cdot) := G'(\overline{X}; \cdot) : \mathcal{X} \to \mathcal{X}$ is the directional derivative of G at \overline{X} .

6 Conclusions

In this paper, we introduced and studied a class of matrix-valued functions, termed spectral operators of matrices, which frequently arise and play a crucial role in various applications including matrix optimization problems, matrix completion, multi-dimensional shape analysis and others. Several fundamental properties of spectral operators, including well-definedness, continuity, directional differentiability, Fréchetdifferentiability, locally Lipschitz continuity, ρ -order B(ouligand)-differentiability ($0 < \rho \leq 1$), ρ -order G-semismooth $(0 < \rho \leq 1)$ and the characterization of Clarke's generalized Jacobian, are studied systematically. These results provide the necessary theoretical foundations for many applications. Consequently, one is able to use these results to design some efficient numerical methods for solving large-scale matrix optimization problems arising from many important applications. For instance, Chen et al. [7] proposed an efficient and robust semismooth Newton-CG dual proximal point algorithm for solving large scale matrix spectral norm approximation problems. In [7], the properties of the spectral operator, such as the semismoothness and the characterization of Clarke's generalized Jacobian, played an important role in the convergence analysis of the proposed algorithm. The work done in this paper on spectral operators of matrices is by no means complete. Due to the rapid advances in the applications of matrix optimization in different fields, spectral operators of matrices will become even more important and many other properties of spectral operators are waiting to be explored.

References

 R.N. BHATTACHARYA AND V. PATRANGENARU. Large sample theory of intrinsic and extrinsic sample means on manifolds. I. *The Annals of Statistics* 31 (2003) 1–29.

- [2] E.J. CANDÈS AND B. RECHT. Exact matrix completion via convex optimization. Foundations of Computational Mathematics 9 (2008) 717–772.
- [3] E.J. CANDÈS AND T. TAO. The power of convex relaxation: near-optimal matrix completion. *IEEE Transactions on Information Theory* 56 (2009) 2053–2080.
- [4] E.J. CANDÈS, X. LI, Y. MA AND J. WRIGHT. Robust principal component analysis? Journal of the ACM 58 (2011) article No. 11.
- [5] E.J. CANDÈS, C.A. SING-LONG AND J.D. TRZASKO. Unbiased risk estimates for singular value thresholding and spectral estimators. Preprint available at http://arxiv.org/abs/1210.4139 (2012).
- [6] V. CHANDRASEKARAN, S. SANGHAVI, P.A. PARRILO AND A. WILLSKY. Rank-sparsity incoherence for matrix decomposition. SIAM Journal of Optimization 21 (2011) 572–596.
- [7] C.H. CHEN, Y.J. LIU, D.F. SUN AND K.C. TOH. A semismooth Newton-CG dual proximal point algorithm for matrix spectral norm approximation problems. Preprint available at http://www.math.nus.edu.sg/~matsundf/SNDPPA-7.pdf (2012).
- [8] X. CHEN, H.D. QI AND P. TSENG. Analysis of nonsmooth symmetric-matrix-valued functions with applications to semidefinite complement problems. *SIAM Journal on Optimization* 13 (2003) 960–985.
- [9] M. CHU, R. FUNDERLIC AND R. PLEMMONS. Structured low rank approximation. *Linear Algebra* and its Applications 366 (2003) 157–172.
- [10] F.H. CLARKE. Optimization and Nonsmooth Analysis. John Wiley & Sons (New York, 1983).
- [11] C. DAVIS. All convex invariant functions of hermitian matrices. Archiv der Mathematik 8 (1957) 276–278.
- [12] C.-A. DELEDALLE, S. VAITER, G. PEYRÉ, J. FADILI AND C. DOSSAL. Risk estimation for matrix recovery with spectral regularization. Preprint available at http://arxiv.org/abs/1205.1482 (2012).
- [13] V.F. DEMYANOV AND A.M. RUBINOV. On quasidifferentiable mappings. Optimization 14 (1983) 3–21.
- [14] C. DING, D.F. SUN AND K.C. TOH. An introduction to a class of matrix cone programming, Mathematical Programming (2012) DOI:10.1007/s10107-012-0619-7.
- [15] C. DING. An Introduction to a Class of Matrix Optimization Problems. PhD thesis. National University of Singapore, http://www.math.nus.edu.sg/~matsundf/DingChao_Thesis_final.pdf (2012).
- [16] C. Ding AND H.D. QI. А computable characterization of the extrinsic mean of reflection shapes and itsasymptotic properties. Preprint available at http://www.maths.soton.ac.uk/staff/Qi/REPORTS/Extrinsic_Mean_Shape.pdf (2013).
- [17] V. DOBRYNIN. On the rank of a matrix associated with a graph. Discrete Mathematics 276 (2004) 169–175.
- [18] I.L. DRYDEN AND K.V. MARDIA. Statistical Analysis of Shape. John Wiley & Sons (Chichester, 1998).
- [19] T.M. FLETT. Differential Analysis. Cambridge University Press (Cambridge, England, 1980).

- [20] Y. Gao D.F. SUN. А AND majorized penalty approach for calibratconstrained problems. rank correlation matrix Preprint available ing at http://www.math.nus.edu.sg/~matsundf/MajorPen_May5.pdf (2010).
- [21] A. GREENBAUM AND L.N. TREFETHEN. GMRES/CR and Arnoldi/Lanczos as matrix approximation problems. SIAM Journal on Scientific Computing 15 (1994) 359–368.
- [22] D. GROSS. Recovering low-rank matrices from few coefficients in any basis. IEEE Transactions on Information Theory 57 (2011) 1548–1566.
- [23] R.H. KESHAVAN, A. MONTANARI AND S. OH. Matrix completion from a few entries. *IEEE Trans*actions on Information Theory 56 (2010) 2980–2998.
- [24] A. KOTLOV, L. LOVÁSZ AND S. VEMPALA. The Colin de Verdière number and sphere representations of a graph. *Combinatorica* 17 (1997) 483–521.
- [25] P. LANCASTER. On eigenvalues of matrices dependent on a parameter. Numerische Mathematik 6 (1964) 377–387.
- [26] A.S. LEWIS. The convex analysis of unitarily invariant matrix functions. Journal of Convex Analysis 2 (1995) 173–183.
- [27] A.S. LEWIS. Derivatives of spectral functions. Mathematics of Operations Research 21 (1996) 576– 588.
- [28] A.S. LEWIS AND H.S. SENDOV. Twice differentiable spectral functions. SIAM Journal on Matrix Analysis and Applications 23 (2001) 368–386.
- [29] A.S. LEWIS AND H.S. SENDOV. Nonsmooth analysis of singular values. Part I: theory. Set-Valued Analysis 13 (2005) 213–241.
- [30] A.S. LEWIS AND H.S. SENDOV. Nonsmooth analysis of singular values. Part II: applications. Set-Valued Analysis 13 (2005) 243–264.
- [31] Y.J. LIU, D.F. SUN, AND K.C. TOH. An implementable proximal point algorithmic framework for nuclear norm minimization. *Mathematical Programming* 133 (2012) 399–436.
- [32] K. LÖWNER. Über monotone matrixfunktionen. Mathematische Zeitschrift 38 (1934) 177–216.
- [33] L. LOVÁSZ. On the Shannon capacity of a graph. IEEE Transactions on Information Theory 25 (1979) 1–7.
- [34] W.M. MIAO, D.F. SUN AND S.H. PAN. A rank-corrected procedure for matrix completion with fixed basis coefficients. Preprint available at http://arxiv.org/abs/1210.3709 (2012).
- [35] W.M. MIAO. Matrix Completion Models with Fixed Basis Coefficients and Rank Regularized Problems with Hard Constraints. PhD Thesis. National University of Singapore, http://www.math.nus.edu.sg/~matsundf/PhDThesis_Miao_Final.pdf (2013).
- [36] J.-J. MOREAU. Proximité et dualité dans un espace hilbertien. Bulletin de la Société Mathématique de France 93 (1965) 1067–1070.
- [37] M.Z. NASHED. Differentiability and related properties of nonlinear operators: some aspects of the role of differentials in nonlinear functional analysis. *Nonlinear Functional Analysis and Applications*, Edited by L. B. Rall. Academic Press (New York, 1971) 103–309.
- [38] J.M. ORTEGA AND W.C. RHEINBOLDT. Iterative Solution of Nonlinear Equations in Several Variables. Society for Industrial Mathematics (1970).

- [39] H.D. QI AND X.Q. YANG. Semismoothness of spectral functions. SIAM Journal on Matrix Analysis and Applications 25 (2003) 766–783.
- [40] L. QI. Convergence analysis of some algorithms for solving nonsmooth equations. Mathematics of Operations Research 18 (1993) 227–244.
- [41] L. QI AND J. SUN. A nonsmooth version of Newton's method. Mathematical Programming 58 (1993) 353–367.
- [42] B. RECHT. A simpler approach to matrix completion. Journal of Machine Learning Research 12 (2011) 3413–3430.
- [43] B. RECHT, M. FAZEL AND P.A. PARRILO. Guaranteed minimum rank solutions to linear matrix equations via nuclear norm minimization. *SIAM Review* 52 (2010) 471–501.
- [44] S.M. ROBINSON. Local structure of feasible sets in nonlinear programming, Part III: stability and sensitivity. *Mathematical Programming Study* 30 (1987) 45–66.
- [45] R.T. ROCKAFELLAR. Convex Analysis. Princeton University Press (Princeton, 1970).
- [46] R.T. ROCKAFELLAR AND R.J-B. WETS. Variational Analysis. Springer (Berlin, 1998).
- [47] A. SHAPIRO. On concepts of directional differentiability. Journal of Optimization Theory and Applications 66 (1990) 477–487.
- [48] C.M. STEIN. Estimation of the mean of a multivariate normal distribution. The Annals of Statistics 9 (1981) 1135–1151.
- [49] V.A. STEKLOV. On the asymptotic representation of certain functions defined by a linear differential equation of the second order, and their application to the problem of expanding an arbitrary function into a series of these functions. *Kharkov* (1957) (In Russian).
- [50] D.F. SUN AND J. SUN. Semismooth matrix-valued functions. Mathematics of Operations Research 27 (2002) 150–169.
- [51] D.F. SUN AND J. SUN. Strong semismoothness of eigenvalues of symmetric matrices and its applications in inverse eigenvalue problems. *SIAM Journal on Numerical Analysis* 40 (2003) 2352–2367.
- [52] D.F. SUN AND J. SUN. Löwner's operator and spectral functions in Euclidean Jordan algebras. Mathematics of Operations Research 33 (2008) 421–445.
- [53] M.J. TODD. Semidefinite optimization. Acta Numerica 10 (2001) 515–560.
- [54] K.C. TOH. GMRES vs. ideal GMRES. SIAM Journal on Matrix Analysis and Applications 18 (1997) 30–36.
- [55] K.C. TOH AND L.N. TREFETHEN. The Chebyshev polynomials of a matrix. SIAM Journal on Matrix Analysis and Applications 20 (1998) 400–419.
- [56] M. TORKI. Second-order directional derivatives of all eigenvalues of a symmetric matrix. Nonlinear Analysis 46 (2001) 1133–1150.
- [57] J. WRIGHT, Y. MA, A. GANESH AND S. RAO. Robust principal component analysis: exact recovery of corrupted low-rank matrices via convex optimization. In Y. Bengio, D. Schuurmans, J. Lafferty and C. Williams, editors, *Advances in Neural Information Processing Systems* 22 (2009).
- [58] B. WU, C. DING, D. F. SUN, AND K. C. TOH. On the Moreau-Yosida regularization of the vector k-norm related functions. *SIAM Journal on Optimization*, to appear.

- [59] Z. YANG. A Study on Nonsymmetric Matrix-valued Functions. Master's thesis. National University of Singapore, http://www.math.nus.edu.sg/~matsundf/Main_YZ.pdf (2009).
- [60] X.Y. ZHAO, D.F. SUN, AND K.C. TOH. A Newton-CG augmented Lagrangian method for semidefinite programming. SIAM Journal on Optimization 20 (2010) 1737–1765.