# Maximal increasing sequences in fillings of almost-moon polyominoes

Svetlana Poznanović<br/>1 and Catherine H.  ${\rm Yan^2}$ 

<sup>1</sup> Department of Mathematical Sciences Clemson University, Clemson, SC 29634

<sup>2</sup>Department of Mathematics Texas A&M University, College Station, TX 77843, USA

#### Abstract

It was proved by Rubey that the number of fillings with zeros and ones of a given moon polyomino that do not contain a northeast chain of size k depends only on the set of columns of the polyomino, but not the shape of the polyomino. Rubey's proof is an adaption of *jeu de taquin* and promotion for arbitrary fillings of moon polyominoes. In this paper we present a bijective proof for this result by considering fillings of almost-moon polyominoes, which are moon polyominoes after removing one of the rows. Explicitly, we construct a bijection which preserves the size of the largest northeast chains of the fillings when two adjacent rows of the polyomino are exchanged. This bijection also preserves the column sum of the fillings. We also present a bijection that preserves the size of the largest northeast chains, the row sum and the column sum if every row of the fillings has at most one 1.

Keywords: maximal chains, moon polyominoes

MSC Classification: 05A19

# 1 Introduction

The systematic study of matchings and set partitions with certain restrictions on their crossings and nestings started in [2], where Chen et al. used Robinson-Schensted-like insertion/deletion processes to show the symmetry between the sizes of the largest crossings and the largest nestings. Lying in the heart of [2] is Greene's Theorem on the relation between increasing and decreasing subsequences in permutations and the shape of the tableaux which are obtained by the Robinson-Schensted correspondence. These results have been put in a larger context of enumeration of fillings of polyominoes where one imposes restrictions on the increasing and decreasing chains of the fillings.

Krettenthaler [13] extended the results of [2] to 01-fillings as well as N-fillings of Ferrers diagrams, and obtained various generalizations using Fomin's growth diagrams [7, 8, 9]. Jonsson, motivated

E-mail addresses: spoznan@clemson.edu (S. Poznanović), cyan@math.tamu.edu (C.H. Yan)

The first author is partially supported by NSF grant DMS-1312817. The second author is partially supported by NSF grant DMS-1161414.

by the problem of counting generalized triangulations with a given size of the maximal crossings, proved that the number of 01-fillings of a stack polyomino that do not contain a northeast chain of size k depends only on the distribution of lengths of the columns of the polyomino. A stack polyomino is a convex polyomino in which the rows are arranged in a descending order from top to bottom. Jonsson [10] proved the result first for maximal 01-fillings using an involved induction, and later with Welker [11] for all 01-fillings with a fixed number of 1's using the machinery of simplicial complexes and commutative algebra. Rubey [15] generalized Jonsson and Welker's result to moon polyominoes, using an adaptation of *jeu de taquin* and promotion for arbitrary fillings of moon polyominoes.

Moon polyominoes are polyominoes that are convex and intersection-free. In moon polyomino the lengths of rows (and columns) are arranged in a unimodal order. For a moon polyomino  $\mathcal{M}$ , let  $\sigma \mathcal{M}$  be another moon polyomino obtained by permuting the rows of  $\mathcal{M}$ . Rubey proved that the number of 01-fillings with the longest northeast chains of size k and exactly  $c_i$  non-zero entries in column i are equal for  $\mathcal{M}$  and  $\sigma \mathcal{M}$ . It is a very interesting property for fillings of moon polyominoes: many combinatorial statistics are invariant under permutations of rows (or columns). In addition to Rubey's results, it is also known for a major index introduced by Chen, Poznanovic, Yan and Yang [3], for the numbers of northeast and southeast chains of length 2 by Kasraoui [12], and for various analogs and generalizations of 2-chains [4, 16].

The purpose of the present paper is to construct bijective proofs of Rubey's result. Inspired by the work on layer polyominoes [14], we seek to extend the moon polyominoes to a general family that would allow us to transform the moon polyomino  $\mathcal{M}$  to  $\sigma \mathcal{M}$  by a sequence of steps that interchange two adjacent rows at each time. For this purpose we introduce the notion of almostmoon polyominoes, which become moon polyominoes after removing one of its rows (see Section 2 for the exact definition). Let  $\mathcal{M}$  and  $\mathcal{N}$  be two almost-moon polyominoes that are related by an interchange of two adjacent rows. We present two bijections. The first is a map  $\phi_{\mathcal{M},\mathcal{N}}$  from 01-fillings of  $\mathcal{M}$  with exactly n 1's to those of  $\mathcal{N}$  such that it preserves the size of the longest northeast chains and the column sum. The second map  $\psi_{\mathcal{M},\mathcal{N}}$  is restricted to fillings where every row has at most one 1, and preserves the size of the longest northeast chains, the row sum, and the column sum.

Rubey's result implies that the number of 01-fillings of a moon polynomial without northeast chains of size k equals those without southeast chains of size k. There are also known combinatorial transformations and bijections between certain families of fillings that avoid northeast chains of size k and southeast chains of size k, for example, by Backelin, West and Xin [1] for 01-fillings of Ferrers diagrams where every row and every column has exactly one 1, and by de Mier [6] for N-fillings of Ferrers diagrams with fixed row sum and column sum. Nevertheless, none of them gives a bijection on the fillings of polyominoes that preserves the size of the longest northeast chains.

The paper is organized as follows. Section 2 contains necessary notations and the statement of main results. In Section 3 we construct the bijection  $\phi_{\mathcal{M},\mathcal{N}}$  in fillings of almost-moon polyominoes which preserves the size of maximal northeast chains and the columns sum. In section 4 we restrict to fillings that have at most one 1 in each row, and describe the bijection  $\psi_{\mathcal{M},\mathcal{N}}$ . We conclude the paper with some comments and counterexamples to a few seeming natural generalizations in Section 5.

# 2 Notation and statements of the main results

A polyomino is a finite subset of  $\mathbb{Z}^2$ , where we represent every element (i, j) of  $\mathbb{Z}^2$  by a square cell. The polyomino is row-convex (column-convex) if its every row (column) is connected. If the polyomino is both row- and column-convex, we say that it is convex. It is intersection-free if every two columns are comparable, i.e., the row-coordinates of one column form a subset of those of the other column. Equivalently, it is intersection-free if every two rows are comparable. A moon polyomino is a convex intersection-free polyomino (e.g. Figure 1a). The length of a row (or a column) is the number of cells in it. Note that in a moon polyomino the lengths of rows from top to bottom form a unimodal sequence. We will say that a row  $\mathcal{R}$  is an exceptional row of a polyomino is a polyomino with comparable convex rows and at most one exceptional row (e.g. Figure 1b). Therefore, every moon polyomino is also an almost-moon polyomino and an almost-moon polyomino is not necessarily column-convex.



Figure 1: A moon polyomino and an almost-moon polyomino with an exceptional row  $\mathcal{R}$  that differ by an interchange of adjacent rows.

In this paper we will consider polyominoes whose cells are filled with zeros and ones. A northeast chain, or shortly ne-chain, of size k in such a filling is a set of k cells  $\{(i_1, j_1), (i_2, j_2), \ldots, (i_k, j_k)\}$  with  $i_1 < \cdots < i_k, j_1 < \cdots < j_k$  filled with 1's such that the  $k \times k$  submatrix

$$\mathcal{G} = \{(i_r, j_s) : 1 \le r \le k, 1 \le s \le k\}$$

is contained in the polyomino (with no restriction on the filling of the other cells). See Figure 2 for an illustration. We will call the ne-chains of size k shortly k-chains. Note that in a moon polyomino  $\mathcal{M}$ , k 1-cells in a north-east direction satisfy the submatrix condition if and only if the corners  $(i_1, j_k)$  and  $(i_k, j_1)$  are contained in  $\mathcal{M}$ , which is equivalent to the whole rectangle determined by these corners being contained in  $\mathcal{M}$ . In an almost-moon polyomino, the submatrix condition is satisfied if and only if the vertices  $(i_1, j_k)$  and  $(i_k, j_1)$  either determine a rectangle which is completely contained in  $\mathcal{M}$  or an almost-rectangle with one exceptional row contained in  $\mathcal{M}$ . In the latter case, the exceptional row does not contain any elements from the ne-chain.

For a 01-filling M of an almost-moon polyomino, we denote by ne(M) the size of its largest ne-chains. Suppose that  $\mathcal{M}$  has k rows and  $\ell$  columns and let  $\mathbf{r} \in \mathbb{N}^k$  and  $\mathbf{c} \in \mathbb{N}^{\ell}$ . We will denote by  $\mathcal{F}(\mathcal{M})$  the set of all 01-fillings of  $\mathcal{M}$ , by  $\mathcal{F}(\mathcal{M}, n)$  those fillings with exactly n 1's, and by  $\mathcal{F}(\mathcal{M}, \mathbf{r}, \mathbf{c})$  the set of all 01-fillings with row sums given by  $\mathbf{r}$  and column sums given by  $\mathbf{c}$ . Our first main result states that if  $\mathcal{M}$  and  $\mathcal{N}$  are two almost-moon polyominoes related by an interchange of adjacent rows (e.g. Figure 1), then the statistic ne is equidistributed over the sets  $\mathcal{F}(\mathcal{M}, *, \mathbf{c})$  and



Figure 2: A 01-filling M of an almost-moon polyomino with ne(M) = 3. The 1's are represented by dots and the 0-cells are drawn empty. The circled dots form the only 3-chain in M.

 $\mathcal{F}(\mathcal{N}, *, \mathbf{c})$  of fillings of  $\mathcal{M}$  and  $\mathcal{N}$ , respectively, with fixed column sums but arbitrary row sums:

$$\sum_{M \in \mathcal{F}(\mathcal{M}, *, \mathbf{c})} q^{\operatorname{ne}(M)} = \sum_{M \in \mathcal{F}(\mathcal{N}, *, \mathbf{c})} q^{\operatorname{ne}(M)}$$

More precisely, we have the following theorem.

**Theorem 1.** Let  $\mathcal{M}$  and  $\mathcal{N}$  be two almost-moon polyominoes such that  $\mathcal{N}$  can be obtained from  $\mathcal{M}$  by an interchange of two adjacent rows. In addition assume that  $\mathcal{M}$  and  $\mathcal{N}$  have no exceptional rows other than the swapped ones. Then there is a bijection

$$\phi_{\mathcal{M},\mathcal{N}}:\mathcal{F}(\mathcal{M})\longrightarrow\mathcal{F}(\mathcal{N})$$

that preserves the column sums of the fillings and such that  $ne(\phi_{\mathcal{M},\mathcal{N}}(M)) = ne(M)$  for  $M \in \mathcal{F}(\mathcal{M})$ . Moreover,  $\phi_{\mathcal{N},\mathcal{M}} \circ \phi_{\mathcal{M},\mathcal{N}} = 1_{\mathcal{F}(\mathcal{M})}$ .

Let  $\mathcal{M}$  be an almost-moon polyomino with k rows,  $\sigma \in S_k$ , and suppose the polyomino  $\sigma \mathcal{M}$  obtained by permuting the rows of  $\mathcal{M}$  according to  $\sigma$  is also an almost-moon polyomino. Note that  $\sigma \mathcal{M}$  can also be obtained by a sequence of steps in which only two adjacent rows are interchanged. Moreover, the order of steps can be chosen so that the intermediate polyominoes are all almost-moon polyominoes with no exceptional rows other than the swapped ones. In other words, the set of all  $\sigma \mathcal{M}$  which are almost-moon polyominoes is connected by transposition of adjacent rows. One way to see this is to note that one can reach the polyomino in which the row lengths are descending from top to bottom by starting from  $\mathcal{M}$  and first moving its exceptional row down until there is no longer rows below it, then moving the shortest row of  $\mathcal{M}$  to the bottom, the second shortest row to the second position from below, etc. Consequently, by composing the maps from Theorem 1 we get the following corollary.

**Corollary 2.** Let  $\mathcal{M}$  be an almost-moon polyomino with k rows and  $\ell$  columns,  $\sigma \in S_k$  be a permutation of the row indices such that  $\sigma \mathcal{M}$  is also an almost-moon polyomino. Let  $\mathbf{c} \in \mathbb{N}^{\ell}$ . Then there is a bijection  $\phi : \mathcal{F}(\mathcal{M}, *, \mathbf{c}) \longrightarrow \mathcal{F}(\sigma \mathcal{M}, *, \mathbf{c})$  such that  $\operatorname{ne}(\phi(M)) = \operatorname{ne}(M)$  for  $M \in \mathcal{F}(\mathcal{M}, *, \mathbf{c})$ . Moreover, the size of  $\{M : M \in \mathcal{F}(\mathcal{M}, n), \operatorname{ne}(M) = k\}$  depends only on the set of lengths of the columns of  $\mathcal{M}$ .

*Proof.* To prove the second part of the statement, suppose  $\mathcal{M}_1$  is the moon polyomino with descending row lengths that can be obtained by reordering the rows of  $\mathcal{M}$  as in the discussion above. Suppose that the same procedure applied to the transpose  $\mathcal{M}_1^t$  of  $\mathcal{M}_1$  yields the polyomino  $\mathcal{M}_2$  with descending row lengths. Then  $\mathcal{M}_2$  is a Ferrers shape and it depends only on the set of column lengths of  $\mathcal{M}$ . Since the transpose of ne-chains are also ne-chains, it follows from the first part that the size of  $\{M : M \in \mathcal{F}(\mathcal{M}, n), \operatorname{ne}(M) = k\}$  also depends only on the set of lengths of the columns of  $\mathcal{M}$ .

The fact that  $|\{M : M \in \mathcal{F}(\mathcal{M}, *, \mathbf{c}), \operatorname{ne}(M) = k\}| = |\{M : M \in \mathcal{F}(\sigma\mathcal{M}, *, \mathbf{c}), \operatorname{ne}(M) = k\}|$  if  $\mathcal{M}$ and  $\sigma\mathcal{M}$  are moon polyominoes as well as the second part of Corollary 2 was proved algebraically by Rubey [15]. Therefore, our results provide a bijective proof of these facts and extend them to a larger set of polyominoes in which these properties hold. In Section 5 we discuss why this extension is in a sense the best possible.

As discussed in [15], one cannot hope to simultaneously preserve both  $\mathbf{r}$  and  $\mathbf{c}$ , i.e., the natural generalization  $|\{M : M \in \mathcal{F}(\mathcal{M}, \mathbf{r}, \mathbf{c}), \operatorname{ne}(M) = k\}| = |\{M : M \in \mathcal{F}(\sigma \mathcal{M}, \sigma \mathbf{r}, \mathbf{c}), \operatorname{ne}(M) = k\}|$  does not hold. However, our second main result implies that ne can be preserved together with both the row and column sums if the fillings are restricted to have at most one 1 in each row.

**Theorem 3.** Let  $\mathcal{M}$  and  $\mathcal{N}$  be two almost-moon polynomial such that  $\mathcal{N}$  can be obtained from  $\mathcal{M}$  by an interchange of two adjacent rows. In addition assume that  $\mathcal{M}$  and  $\mathcal{N}$  have no exceptional rows other than the swapped ones. If  $\mathbf{r} \in \{0,1\}^*$  and  $\mathbf{c} \in \mathbb{N}^*$ , then there is a map

$$\psi_{\mathcal{M},\mathcal{N}}: \mathcal{F}(\mathcal{M},\boldsymbol{r},\boldsymbol{c}) \longrightarrow \mathcal{F}(\mathcal{N},\boldsymbol{r}',\boldsymbol{c})$$

such that  $ne(\psi_{\mathcal{M},\mathcal{N}}(M)) = ne(M)$  for  $M \in \mathcal{F}(\mathcal{M}, \mathbf{r}, \mathbf{c})$ , where  $\mathbf{r}'$  is obtained from  $\mathbf{r}$  by exchanging the entries corresponding to the two swapped rows.

By the same discussion after Theorem 1, we get the following corollary.

**Corollary 4.** Let  $\mathcal{M}$  be an almost-moon polyomino with k rows and  $\ell$  columns,  $\sigma \in S_k$  be a permutation of the row indices such that  $\sigma \mathcal{M}$  is also an almost-moon polyomino. Let  $\mathbf{r} \in \{0,1\}^k$  and  $\mathbf{c} \in \mathbb{N}^l$ . Then there is a bijection  $\psi : \mathcal{F}(\mathcal{M}, \mathbf{r}, \mathbf{c}) \longrightarrow \mathcal{F}(\sigma \mathcal{M}, \sigma \mathbf{r}, \mathbf{c})$  such that  $\operatorname{ne}(\psi(M)) = \operatorname{ne}(M)$  for  $M \in \mathcal{F}(\mathcal{M}, \mathbf{r}, \mathbf{c})$ . Moreover, the size of  $\{M : M \in \mathcal{F}(\mathcal{M}, \mathbf{r}, \mathbf{c}), \operatorname{ne}(M) = k\}$  depends only on the sequence of lengths of the columns of  $\mathcal{M}$ .

*Proof.* To see the second part of the statement, note that exchanging adjacent rows allows us to transform  $\mathcal{M}$  to  $\mathcal{M}_1$ , the polyomino with descending row lengths, whose shape is determined by the sequence of lengths of the columns of  $\mathcal{M}$ .

#### 3 Maximal increasing sequences in 01-fillings with fixed total sum

In this section we will describe the maps  $\phi_{\mathcal{M},\mathcal{N}}$  and prove Theorem 1. To this end, let  $\mathcal{M}$  and  $\mathcal{N}$  be two almost-moon polyominoes related by an interchange of two adjacent rows (e.g. Figure 1). Assume that  $\mathcal{M}$  and  $\mathcal{N}$  have no exceptional rows other than the swapped ones. If the swapped rows are of equal length then  $\mathcal{M} = \mathcal{N}$  and we define  $\phi_{\mathcal{M},\mathcal{N}}$  to be the identity map on  $\mathcal{F}(\mathcal{M})$ .

Otherwise, suppose the lengths of the swapped rows are not equal and let  $\mathcal{R}_s$  and  $\mathcal{R}_l$ , respectively, be the shorter and longer rows. Note that  $\mathcal{M}\backslash\mathcal{R}_s$  and  $\mathcal{N}\backslash\mathcal{R}_s$  have no exceptional rows. Let  $\alpha, \beta, \gamma, \delta$  be the fillings of the regions of  $\mathcal{R}_s$  and  $\mathcal{R}_l$  in  $\mathcal{M}$  as depicted on the left side of Figure 3. Precisely,  $\alpha$  is the filling of the shorter row  $\mathcal{R}_s$ ,  $\beta$  is the filling of the part of the longer row  $\mathcal{R}_l$  which has the same column support as row  $\mathcal{R}_s$ , and  $\gamma$  and  $\delta$  are the fillings of the two ends of row  $\mathcal{R}_l$  so that the whole filling of row  $\mathcal{R}_l$  viewed as a binary string is a concatenation of  $\gamma$ ,  $\beta$ , and  $\delta$ . Note that one of  $\gamma$  and  $\delta$  may be the empty string. First we define a map  $f_{\mathcal{M},\mathcal{N}}: \mathcal{F}(\mathcal{M}) \longrightarrow \mathcal{F}(\mathcal{N})$  as follows.

**Definition 5.** Using the notation above,  $f_{\mathcal{M},\mathcal{N}}(M)$  is the filling of  $\mathcal{N}$  in which

- (1) the filling of the rows in  $\mathcal{N}$  other than  $\mathcal{R}_s$  and  $\mathcal{R}_l$  is the same as in M,
- (2) the filling of the shorter row  $\mathcal{R}_s$  of  $\mathcal{N}$  is  $\beta$ ,
- (3) the filling of the longer row  $\mathcal{R}_l$  of  $\mathcal{N}$  is the concatenation of  $\gamma$ ,  $\alpha$ , and  $\delta$ .

See Figure 3 for an illustration.



Figure 3: The fillings M and  $f_{\mathcal{M},\mathcal{N}}(M)$  differ only in the rows  $\mathcal{R}_s$  and  $\mathcal{R}_l$ .

It follows directly from the definition.

Lemma 6.  $f_{\mathcal{N},\mathcal{M}} \circ f_{\mathcal{M},\mathcal{N}} = 1_{\mathcal{F}(\mathcal{M})}$ .

If it is clear what the polyominoes  $\mathcal{M}$  and  $\mathcal{N}$  are, we will leave out the subscripts and write only f(M).

**Lemma 7.** If the almost-moon polyominoes  $\mathcal{M}$  and  $\mathcal{N}$  are related by an interchange of the rows  $\mathcal{R}_s$  ans  $\mathcal{R}_l$  as above, then for every  $M \in \mathcal{F}(\mathcal{M})$ ,

$$|\mathrm{ne}(M) - \mathrm{ne}(f_{\mathcal{M},\mathcal{N}}(M))| \le 1.$$

*Proof.* Let C be a k-chain in M.

- (i) If C contains no 1-cells from  $\mathcal{R}_s \cup \mathcal{R}_l$  then C is a k-chain in f(M).
- (ii) If C contains a 1-cell from  $\alpha$  then C is a k-chain in f(M).
- (iii) If C contains a 1-cell  $b_0$  from  $\beta$  then  $C \{b_0\}$  is a (k-1)-chain in f(M).
- (iv) If C contains a 1-cell from  $\gamma$  or  $\delta$  then C is a k-chain in f(M).

Therefore,  $ne(f(M)) \ge ne(M) - 1$ . By switching the roles of M and f(M) with a similar analysis, we get  $ne(M) \ge ne(f(M)) - 1$ .

Therefore, every filling  $M \in \mathcal{F}(\mathcal{M})$  satisfies exactly one of the following 3 conditions:

(I) 
$$ne(f(M)) = ne(M)$$
 (II)  $ne(f(M)) = ne(M) + 1$  (III)  $ne(f(M)) = ne(M) - 1$ .

Let  $\mathcal{F}^{I}(\mathcal{M})$ ,  $\mathcal{F}^{II}(\mathcal{M})$ , and  $\mathcal{F}^{III}(\mathcal{M})$ , be the fillings of  $\mathcal{M}$  that satisfy the conditions (I), (II), and (III), respectively. Below we describe how  $\phi_{\mathcal{M},\mathcal{N}}(\mathcal{M})$  is defined on each of these three sets.

**Case I.** For  $M \in \mathcal{F}^{I}(\mathcal{M})$  we define  $\phi_{\mathcal{M},\mathcal{N}}(M) = f_{\mathcal{M},\mathcal{N}}(M)$ . It is clear that this defines a bijection from  $\mathcal{F}^{I}(\mathcal{M})$  onto  $\mathcal{F}^{I}(\mathcal{N})$ .

**Case II.** For  $M \in \mathcal{F}^{II}(\mathcal{M})$  with  $\operatorname{ne}(M) = k$ , we have  $\operatorname{ne}(N') = k + 1$  where N' = f(M). Reasoning as in the proof of Lemma 7, one can see that all (k + 1)-chains in N' contain exactly one cell from  $\mathcal{R}_s \cup \mathcal{R}_l$  and that that cell must be in the part  $\alpha$  of the longer row  $\mathcal{R}_l$  of  $\mathcal{N}$  (see the right filling in Figure 3).

**Definition 8.** Suppose  $M \in \mathcal{F}^{II}(\mathcal{M})$  and  $\operatorname{ne}(f(M)) = \operatorname{ne}(M) + 1 = k + 1$ . The 1-cells in row  $\mathcal{R}_l$  of f(M) which are part of a (k + 1)-chain are called problem cells.

Let  $\alpha_0$  be the set of problem cells in N' = f(M) and let  $\beta_0$  be the set of cells in row  $\mathcal{R}_s$  of N'that share a horizontal edge with the problem cells. We define  $\phi_{\mathcal{M},\mathcal{N}}(M)$  to be the filling N'' of  $\mathcal{N}$ obtained by replacing the problem cells in  $\alpha_0$  by zeros and the cells in  $\beta_0$  by ones. In other words,  $\phi_{\mathcal{M},\mathcal{N}}(M) = N''$  is obtained by a vertical shift of the problem cells in N' from row  $\mathcal{R}_l$  to row  $\mathcal{R}_s$ .

**Case III.** For  $M \in \mathcal{F}^{III}(\mathcal{M})$ , we have  $f_{\mathcal{M},\mathcal{N}}(M) \in \mathcal{F}^{II}(\mathcal{N})$ . In this case we set  $\phi_{\mathcal{M},\mathcal{N}}(M) = f_{\mathcal{M},\mathcal{N}}(\phi_{\mathcal{N},\mathcal{M}}(f_{\mathcal{M},\mathcal{N}}(M)))$ .

In the following we show that  $\phi_{\mathcal{M},\mathcal{N}}$  is a well-defined bijection from  $\mathcal{F}(\mathcal{M}, *, \mathbf{c})$  to  $\mathcal{F}(\mathcal{N}, *, \mathbf{c})$ preserving the statistic ne. Lemmas 9–12 show that  $\phi_{\mathcal{M},\mathcal{N}}$  defined in Case II preserves column sum and ne( $\mathcal{N}''$ ) = k. Lemma 13 shows that  $\phi_{\mathcal{M},\mathcal{N}}$ , when restricted to  $\mathcal{F}^{II}(\mathcal{M})$ , is a bijection onto  $\mathcal{F}^{II}(\mathcal{N})$ . Lemma 14 shows  $\phi_{\mathcal{M},\mathcal{N}}$  restricted to  $\mathcal{F}^{III}(\mathcal{M})$  is a bijection onto  $\mathcal{F}^{III}(\mathcal{N})$ .

We first assume that M is a filling in  $\mathcal{F}^{II}(\mathcal{M})$  with ne(M) = k.

**Lemma 9.** If  $a_0$  in row  $\mathcal{R}_l$  of f(M) is a problem cell then the cell  $b_0$  in row  $R_s$  of f(M) in the same column as  $a_0$  is a 0-cell.

*Proof.* Suppose  $a_0$  is part of the (k + 1)-chain C in f(M). If  $b_0$  is a 1-cell then  $(C - \{a_0\}) \cup \{b_0\}$  is a (k + 1)-chain in M.

Lemma 9 shows that in Case II, the cells in  $\beta_0$  are 0-cells for N'. Hence N'' is a 01-filling of  $\mathcal{N}$  and the column sums of M and N'' are the same. We will prove next that ne(N'') = k.

Since  $\operatorname{ne}(N') = k + 1$ , from the way N'' is constructed it can readily be seen that  $\operatorname{ne}(N'') \ge k$ . Suppose  $\operatorname{ne}(N'') > k$ . Since the ne-chains are preserved under rotation by  $180^\circ$ , in the discussion below we can assume, without loss of generality, that in  $\mathcal{M}$  the shorter row  $\mathcal{R}_s$  is above  $\mathcal{R}_l$ . This will allow us to avoid introducing too much notation, and instead use words such as "below", "above", etc. Let  $\mathcal{S}$  be the intersection of  $\mathcal{M}$  (and therefore  $\mathcal{N}$ ) and the vertical strip determined by the row  $\mathcal{R}_s$ . Precisely, let

 $\mathcal{S} = \{(x, y) : (x, y) \text{ is a cell in } \mathcal{M} \text{ such that there is a cell } (x, y') \in \mathcal{R}_s\}.$ 

**Lemma 10.** Every (k + 1)-chain in N'' contains one 1-cell in each of the rows  $\mathcal{R}_s$  and  $\mathcal{R}_l$ . Thus, every such chain is contained in the strip  $\mathcal{S}$ .

Proof. Since ne(N'') > k, N'' contains a (k+1)-chain C. If none of the elements of C is in  $\mathcal{R}_s \cup \mathcal{R}_l$ , then C is a chain in M, which yields a contradiction with ne(M) = k. If C contains a 1-cell c from  $\mathcal{R}_l$  but not a cell from  $\mathcal{R}_s$  then C is a (k+1)-chain in N' also, which means that c is a problem cell, which contradicts the fact that during the construction of N'' all problem cells in N' are replaced by 0's. If C contains a 1-cell c from  $\mathcal{R}_s$  but not a cell from  $\mathcal{R}_l$  then there are 2 possible cases: (i) c is a 1-cell in N'. In this case C is a (k+1)-chain in M as well. This contradicts ne(M) = k. (ii) c is a 0-cell in N'. In this case the cell d above c in row  $\mathcal{R}_l$  is a problem 1-cell and C is contained in the strip S. Then  $(C - \{c\}) \cup \{d\}$  is a (k+1)-chain in M. This again contradicts ne(M) = k.

Let  $y_1 - c - b - x_1$  be a (k+1)-chain in N'', where c and b are the 1-cells in  $\mathcal{R}_s$  and  $\mathcal{R}_l$  respectively,  $y_1$  is the part of the chain which is southwest of c and  $x_1$  is the part of the chain which is northeast of b. Note that, since b is a 1-cell in N'', it is also a 1-cell in N' but it is not a problem cell.

**Lemma 11.** The cell c is a 0-cell in N' and therefore the cell a in row  $\mathcal{R}_l$  directly above c is a problem cell.

*Proof.* If c is a 1-cell in N' then  $y_1 - c - b - x_1$  is a (k + 1)-chain in N' as well, which implies that b is a problem cell. But then b would be a 0-cell in N" and cannot be a part of a (k + 1)-chain in N".

Therefore, the cell a is a part of a (k + 1)-chain  $y_2 - a - x_2$  in N'. Moreover, this chain is not contained in the strip S. Otherwise,  $y_2 - a - x_2$  is a (k + 1)-chain in M as well. We will show that this implies a contradiction, as stated in Lemma 12. Figure 4 illustrates the positions of the two chains.



Figure 4: The (k + 1)-chain  $y_1 - c - b - x_1$  in N'' is contained in the strip S. The (k + 1)-chain  $y_2 - a - x_2$  in N' is not contained in S. In N' : a = 1, b = 1, c = 0, in N'' : a = 0, b = 1, c = 1.

Lemma 12. Suppose, using the notation above,

$$y_1 - c - b - x_1$$
 is a  $(k+1)$ -chain in  $N''$  contained in  $\mathcal{S}$ , (1)

$$y_2 - a - x_2$$
 is a  $(k+1)$ -chain in N' not contained in S. (2)

Then M has a (k+1)-chain or N' has a (k+1)-chain that contains b.

*Proof.* We will discuss cases according to the relative positions of  $x_1$  and  $x_2$ . For the discussion, it is helpful to visualize a chain as a piece of string directed northeast connecting the centers of

its 1-cells, so that it's piecewise linearly increasing. Then we will say that chains cross if the corresponding strings cross. Let  $r_{\mathcal{M}}(d)$  denote the row in  $\mathcal{M}$  that contains the cell d and define  $r_{\mathcal{M}}(d_1) \leq r_{\mathcal{M}}(d_2)$  if the set of abscissas of the cells of row  $r_{\mathcal{M}}(d_1)$  is contained in the set of abscissas of the cells of  $r_{\mathcal{M}}(d_2)$ . For any almost-moon polyomino, this defines a total order on its rows. Note that condition (2) implies that

$$\mathcal{R}_s \le r_{\mathcal{N}}(d)$$
 for every  $d \in x_2 \cup y_2$ . (3)

Moreover,

C forms a k-chain if and only if every two cells in C form a 2-chain, (4)

- $d_1 \in x_1$  and  $d_2 \in x_2$  form a 2-chain if and only if they are in north-east direction, (5)
- $d_1 \in y_1$  and  $d_2 \in x_2$  such that  $d_2$  is weakly to the left of a cell in  $x_1$  form a 2-chain, (6)

 $d_1 \in y_2$  and  $d_2 \in x_1$  such that  $d_2$  is not higher than the highest point of  $x_2$  form a 2-chain, (7) Suppose first that both  $x_1$  and  $x_2$  are nonempty.

- (1) If the chains b—x₁ and a—x₂ cross, let P be the first intersection point. Let x₃ and x₄, respectively, be the parts of x₁ and x₂, respectively, that are strictly southwest of P. Then
  if |x₄| > |x₃| then y₁—a—x₄—(x₁\x₃) is at least a (k + 1)-chain in M; the submatrix condition is satisfied because of (4), (5), and (6);
  if |x₄| ≤ |x₃| then y₂—b—x₃—(x₂\x₄) is at least a (k + 1)-chain in N'; the submatrix condition is satisfied because of (4), (5), and (7).
- (2) If the chains  $b-x_1$  and  $a-x_2$  do not cross, there are 4 cases to be considered:
  - (a) x<sub>2</sub> ends weakly below and strictly to the left of x<sub>1</sub>. Let x<sub>3</sub> be the part of x<sub>1</sub> that is in the rows weakly below the highest cell of x<sub>2</sub>. Then
     if |x<sub>2</sub>| > |x<sub>3</sub>| then y<sub>1</sub>—a—x<sub>2</sub>—(x<sub>1</sub>\x<sub>3</sub>) is at least a (k + 1)-chain in M; the submatrix condition is satisfied because of (4), (5), and (6);
     if |x<sub>2</sub>| ≤ |x<sub>3</sub>| then y<sub>2</sub>—b—x<sub>3</sub> is at least a (k + 1)-chain in N'; the submatrix condition is satisfied because of (4) and (7).
  - (b) x<sub>2</sub> ends strictly above and weakly to the left of x<sub>1</sub>. Then
     if |x<sub>2</sub>| ≤ |x<sub>1</sub>| then y<sub>2</sub>—b—x<sub>1</sub> is at least a (k + 1)-chain in N'; the submatrix condition is satisfied because of (4) and (7);
     if |x<sub>2</sub>| > |x<sub>1</sub>| then y<sub>1</sub>—a—x<sub>2</sub> is at least a (k + 1)-chain in M; the submatrix condition is satisfied because of (4) and (6).
  - (c) x<sub>2</sub> ends strictly above and to the right of x<sub>1</sub>. Let x<sub>4</sub> be the part of x<sub>2</sub> that is in the columns weakly to the left of the end of x<sub>2</sub>. Then
    if |x<sub>4</sub>| > |x<sub>1</sub>| then y<sub>1</sub>—a—x<sub>4</sub> is at least a (k + 1)-chain in M; the submatrix condition is satisfied because of (4) and (6);
    if |x<sub>4</sub>| ≤ |x<sub>1</sub>| then y<sub>2</sub>—b—x<sub>1</sub>—(x<sub>2</sub>\x<sub>4</sub>) is at least a (k + 1)-chain in N'; the submatrix condition is satisfied because of (4), (5), and (7).
  - (d)  $x_2$  ends weakly below and weakly to the right of  $x_1$ . This case is not possible because then  $x_1$  and  $x_2$  must cross.

Finally, we consider the cases when one of  $x_1$  and  $x_2$  is empty.

- (1) If  $|x_2| = 0$  then  $y_2$ —*b* is a (k + 1)-chain in N'; the submatrix condition is satisfied because of (3).
- (2) If |x<sub>2</sub>| ≠ 0 but |x<sub>1</sub>| = 0, let x<sub>5</sub> be the part of x<sub>2</sub> that is in the columns weakly to the left of b. Then
   if |x<sub>5</sub>| > 0 then y<sub>1</sub>—a—x<sub>5</sub> is at least a (k + 1)-chain in M; the submatrix condition is satisfied because of (4) and (6);

- if  $|x_5| = 0$  then  $y_2 - b - x_2$  is a (k + 1)-chain in N'; the submatrix condition is satisfied because of (3).

To conclude, the construction of N'' and the preceding three lemmas imply the following property: For  $M \in \mathcal{F}^{II}(M)$ , the above constructed filling  $N'' = \phi_{\mathcal{M},\mathcal{N}}(M)$  of  $\mathcal{N}$  has the same column sums as M and  $\operatorname{ne}(N'') = \operatorname{ne}(M)$ .

We next prove some properties of the transformation of the, so far, partially defined map  $\phi_{\mathcal{M},\mathcal{N}}$ . These will be useful in extending the definition of  $\phi_{\mathcal{M},\mathcal{N}}$  to  $\mathcal{F}^{III}(\mathcal{M})$ .

**Lemma 13.** Let  $M \in \mathcal{F}^{II}(\mathcal{M})$  with ne(M) = k. Let  $N'' = \phi_{\mathcal{M},\mathcal{N}}(M)$  be defined as in Case II. Then

$$ne(f_{\mathcal{N},\mathcal{M}}(N'')) = ne(N'') + 1 = k + 1.$$

In other words, if  $M \in \mathcal{F}^{II}(\mathcal{M})$  then  $\phi_{\mathcal{M},\mathcal{N}}(M) \in \mathcal{F}^{II}(\mathcal{N})$ . Moreover,

$$\phi_{\mathcal{N},\mathcal{M}}(\phi_{\mathcal{M},\mathcal{N}}(M)) = M.$$

Consequently,  $\phi_{\mathcal{M},\mathcal{N}}$  restricted on  $\mathcal{F}^{II}(\mathcal{M})$  is a bijection onto  $\mathcal{F}^{II}(\mathcal{N})$ .

*Proof.* Suppose A is the set of 1-cells in row  $\mathcal{R}_l$  of N' = f(M) and  $A_0$  is the subset of A containing the problem cells. Let also B be the set of 1-cells in  $\mathcal{R}_s$  of N' and  $B_0$  the set of cells in  $\mathcal{R}_s$  that share a horizontal edge with a cell in  $A_0$ . By construction, the set of 1-cells in row  $\mathcal{R}_l$  of  $N'' = \phi_{\mathcal{M},\mathcal{N}}(M)$  is  $A_1 = A \setminus A_0$  and the set of 1-cells in row  $\mathcal{R}_s$  is  $B_1 = B \cup B_0$ .

We now consider  $M' = f_{\mathcal{N},\mathcal{M}}(N'') \in \mathcal{F}(\mathcal{M})$ . Let  $a \in A_0$  be a problem cell in N'. Then, by definition, a is a part of a (k+1)-chain y - a - x in N' and the cells in the chain are all contained in the vertical strip determined by row  $\mathcal{R}_l$ . Let  $b \in B_0$  be the neighbor cell of a. Then y - b - x forms a (k+1) chain in M'. Since M' contains a (k+1) chain, Lemma 7 implies that  $\operatorname{ne}(M') = k+1$ . The preceding argument also shows that the 1-cells in  $B_0$  are problem cells in M' and it is not difficult to see that these are the only problem cells. Therefore, to construct  $\phi_{\mathcal{N},\mathcal{M}}(N'') \in \mathcal{F}(\mathcal{M})$ , we replace the 1-cells in  $B_0$  by 0's and the neighboring 0-cells in  $A_0$  by 1's, which yields the initial filling M.

**Lemma 14.** Let  $M \in \mathcal{F}^{III}(\mathcal{M})$ . Then  $\phi_{\mathcal{M},\mathcal{N}}(M)$  is well defined, its column sums are the same as M's, and  $\operatorname{ne}(\phi_{\mathcal{M},\mathcal{N}}(M)) = \operatorname{ne}(M)$ . Moreover,  $\phi_{\mathcal{M},\mathcal{N}}(M) \in \mathcal{F}^{III}(\mathcal{N})$  and  $\phi_{\mathcal{N},\mathcal{M}}(\phi_{\mathcal{M},\mathcal{N}}(M)) = M$ . Consequently,  $\phi_{\mathcal{M},\mathcal{N}}$  restricted on  $\mathcal{F}^{III}(\mathcal{M})$  is a bijection onto  $\mathcal{F}^{III}(\mathcal{N})$ .

Proof. To see that  $\phi_{\mathcal{M},\mathcal{N}}(M)$  is well defined, let  $N' = f_{\mathcal{M},\mathcal{N}}(M)$ . Then  $f_{\mathcal{N},\mathcal{M}}(N') = M$  and  $N' \in \mathcal{F}^{II}(\mathcal{N})$  because  $\operatorname{ne}(f_{\mathcal{N},\mathcal{M}}(N')) = \operatorname{ne}(N') + 1$ . Therefore,  $\phi_{\mathcal{N},\mathcal{M}}(N')$  is well defined and so is  $\phi_{\mathcal{M},\mathcal{N}}(M)$ . Additionally, Lemma 13 implies that  $\phi_{\mathcal{N},\mathcal{M}}(N') \in \mathcal{F}^{II}(\mathcal{M})$ . Therefore,

 $\operatorname{ne}(\phi_{\mathcal{M},\mathcal{N}}(M)) = \operatorname{ne}(\phi_{\mathcal{N},\mathcal{M}}(f_{\mathcal{M},\mathcal{N}}(M))) + 1 = \operatorname{ne}(f_{\mathcal{M},\mathcal{N}}(M)) + 1 = \operatorname{ne}(M).$ 

It is clear that the column sums of M and  $\phi_{\mathcal{M},\mathcal{N}}(M)$  are equal. Finally,  $\phi_{\mathcal{N},\mathcal{M}}(\phi_{\mathcal{M},\mathcal{N}}(M)) = M$  follows from Lemma 6 and Lemma 13.

#### 4 Maximal increasing sequences in fillings with restricted row sum

In this section we restrict to fillings with at most one 1 in each row and prove Theorem 3. Explicitly, let  $\mathcal{M}$  and  $\mathcal{N}$  be two almost-moon polyominoes that can be obtained from each other by an interchange of two adjacent rows. Assume that  $\mathcal{M}$  and  $\mathcal{N}$  have no exceptional rows other than the swapped ones. Let  $\mathbf{r} \in \{0,1\}^*$  and  $\mathbf{c} \in \mathbf{N}^*$ , we shall construct a bijection  $\psi_{\mathcal{M},\mathcal{N}}$  from  $\mathcal{F}(\mathcal{M},\mathbf{r},\mathbf{c})$ to  $\mathcal{F}(\mathcal{M},\mathbf{r}',\mathbf{c})$  that preserves the size of the largest ne-chains, where  $\mathbf{r}'$  is obtained from  $\mathbf{r}$  by exchanging the entries corresponding to the two swapped rows.

If the two swapped rows are of equal length, then  $\mathcal{M} = \mathcal{N}$  and we can simply take  $\psi_{\mathcal{M},\mathcal{N}}$  to be the identity map. In the following, we assume that the two swapped rows are  $\mathcal{R}_s$  and  $\mathcal{R}_l$ , where the length of  $\mathcal{R}_s$  is smaller than that of  $\mathcal{R}_l$ . We keep the notations as in the previous section. For any fillings M let  $\alpha, \beta, \gamma, \delta$  be as defined in Figure 3. Note that for  $M \in \mathcal{F}(\mathcal{M}, \mathbf{r}, \mathbf{c})$ , there is at most one 1 in  $\alpha$ , as well as in the union of  $\beta, \gamma$  and  $\delta$ . Define the *coupling filling* of M to be the filling M' of  $\mathcal{M}$  which is obtained from M by exchanging the fillings  $\alpha$  and  $\beta$ . Let N (resp. N') be obtained from M (resp. M') by swapping the rows  $\mathcal{R}_s$  and  $\mathcal{R}_l$  together with their fillings. In other words,  $N = f_{\mathcal{M},\mathcal{N}}(M')$  and  $N' = f_{\mathcal{M},\mathcal{N}}(M)$ . Clearly N and N' are coupling fillings of each other.

We need the following lemma, which is the crucial observation for the construction of  $\psi_{\mathcal{M},\mathcal{N}}$ .

**Lemma 15.** Let fillings (M, M') be a pair of coupling fillings of in  $\mathcal{F}(\mathcal{M}, \mathbf{r}, \mathbf{c})$ , and  $(N, N') = (f_{\mathcal{M},\mathcal{N}}(M'), f_{\mathcal{M},\mathcal{N}}(M))$  be fillings of  $\mathcal{N}$ . Then  $\operatorname{ne}(M) = \operatorname{ne}(N)$  or  $\operatorname{ne}(M) = \operatorname{ne}(N')$ .

We postpone the proof of Lemma 15 and explain how it helps us construct the bijection  $\psi_{\mathcal{M},\mathcal{N}}$ . Lemma 15 implies that

$$\{ne(M), ne(M')\} = \{ne(N), ne(N')\}$$
(8)

as multisets. To see this, note that if ne(N) = ne(N') = k, then applying Lemma 15 to both Mand M' yields ne(M) = ne(M') = k. Otherwise, if  $ne(N) \neq ne(N')$ , then applying Lemma 15 to N, N' yields that one of ne(M), ne(M') equals ne(N), and the other equals ne(N'). Equation (8) allows us to construct an ne-preserving bijection between the coupling fillings  $\{M, M'\}$  and  $\{N, N'\}$ . Combining the bijections of all the coupling fillings we get a desired bijection. Explicitly, we can describe the map  $\psi_{\mathcal{M},\mathcal{N}}$  as follows:

Let M be a filling in  $\mathcal{F}(\mathcal{M}, \mathbf{r}, \mathbf{c})$ .

- 1. If either  $\alpha$  or  $\beta$  has no 1, then let  $\psi_{\mathcal{M},\mathcal{N}}(M) = N$ , the filling of  $\mathcal{N}$  obtained by swapping the two rows together with their fillings.
- 2. If both  $\alpha$  and  $\beta$  contain a 1 in the same column, then again let  $\psi_{\mathcal{M},\mathcal{N}}(M) = N$ .

3. If each of  $\alpha$  and  $\beta$  contain a unique 1 in a distinct column,

$$\psi_{\mathcal{M},\mathcal{N}}(M) = \begin{cases} N & \text{if } \operatorname{ne}(M) = \operatorname{ne}(N) \\ N' & \text{if } \operatorname{ne}(M) \neq \operatorname{ne}(N). \end{cases}$$

It is clear that the map  $\psi_{\mathcal{M},\mathcal{N}}$  is well-defined and preserving the statistic ne. In addition,  $\psi_{\mathcal{M},\mathcal{N}}$  and  $\psi_{\mathcal{N},\mathcal{M}}$  are inverse to each other.

Proof of Lemma 15. If either  $\alpha$  or  $\beta$  contains no 1, or they both contain a 1 in the same column, then swapping the two rows with their fillings will not change any ne-chains, and hence ne(M) = ne(N). In the following we assume that each of  $\alpha$  and  $\beta$  contains a unique 1-cell in a distinct column, and there is no 1 in  $\gamma$  or  $\delta$ . Furthermore we assume that in  $\mathcal{M}$ , the row  $\mathcal{R}_s$  is above the row  $\mathcal{R}_l$ . The opposite case can be treated similarly.

We consider the relative positions of the two 1-cells in  $R_s$  and  $\mathcal{R}_l$ . See Figure 5 for an illustration.



Figure 5: Positions of the 1-cells in the coupling fillings.

**Case I.** In the filling M the two 1-cells in rows  $\mathcal{R}_s$  and  $\mathcal{R}_l$  form an ne-chain of size 2. That is, the 1-cells of  $\mathcal{M}$  in rows  $\mathcal{R}_s$  and  $\mathcal{R}_l$  are b and c as in the left figure of Figure 5. Then N is the filling of  $\mathcal{N}$  in which the only 1-cells in rows  $\mathcal{R}_s$  and  $\mathcal{R}_l$  are a and d, as in the right figure of Figure 5. Clearly  $ne(N) \leq ne(M)$  as every ne-chain of N is still an ne-chain of M.

Assume ne(M) = k. If  $ne(N) \neq ne(M)$  then ne(N) = k - 1, and every ne-chain of length k in M contains both b and c. In particular, N', the coupling filling of N which contains b and c in  $\mathcal{N}$ , has ne-chains of length k. It follows that  $ne(N') \geq k$ .

Comparing fillings N and N'. Let C be an l-chain in N'.

- (i) If C contains no 1-cells from  $\mathcal{R}_s \cup \mathcal{R}_l$  then C is an *l*-chain in N.
- (ii) If C contains the 1-cell c, then  $C \cup \{a\} \{c\}$  is an *l*-chain in N.
- (iii) If C contains the 1-cell b, then  $C \{b\}$  is an (l-1)-chain in N.
- (iv) If C contains both b and c, then  $C \cup \{a\} \{b, c\}$  is an (l-1)-chain in N.

Therefore  $ne(N) \ge ne(N') - 1$  and hence  $ne(N') \le ne(N) + 1 \le k$ . Thus we must have ne(N') = k = ne(M).

**Case II.** In the filling M the two 1-cells in rows  $\mathcal{R}_s$  and  $\mathcal{R}_l$  do not form an ne-chain of size 2. Then M has 1-cells a, d in  $\mathcal{M}, N$  has 1-cells b, c and N' has 1-cells a, d in  $\mathcal{N}$ . Similarly as in Case 1 we have

$$ne(M) \le ne(N) \le ne(M) + 1$$
$$ne(N) - 1 \le ne(N').$$

Using Lemma 7 we also have  $|\operatorname{ne}(M) - \operatorname{ne}(N')| \leq 1$ . Assume that  $\operatorname{ne}(M) = k$  is not equal to  $\operatorname{ne}(N)$  or  $\operatorname{ne}(N')$ . Combining these three inequalities, we derive that  $\operatorname{ne}(N) = \operatorname{ne}(N') = k+1$ . In addition,

- (1) Every (k+1)-chain in N contains both b and c, and hence lies in the strip S, where S is the union of columns that intersect the row  $\mathcal{R}_s$ , as defined in previous section.
- (2) Every (k+1)-chain in N' contains a, and is not lying inside S.

This is exactly the situation described in Lemma 12. Comparing with Figure 4 and applying Lemma 12, also noting that the filling N' does not have b as a 1-cell, we conclude that M has an ne-chain of length k + 1. This is a contradiction.

Although Lemma 12 plays an important role in the proofs of both Theorem 1 and 3, we remark that the map  $\psi_{\mathcal{M},\mathcal{N}}$  is different than the map  $\phi_{\mathcal{M},\mathcal{N}}$  restricted to the set  $\mathcal{F}(\mathcal{M},\mathbf{r},\mathbf{c})$ . For one thing,  $\phi_{\mathcal{M},\mathcal{N}}$  does not always reserve the row sum, hence not necessarily maps fillings of  $\mathcal{F}(\mathcal{M},\mathbf{r},\mathbf{c})$  to  $\mathcal{F}(\mathcal{N},\mathbf{r}',\mathbf{c})$  when  $\mathbf{r} \in \{0,1\}^*$ .

### 5 Concluding remarks

We conclude this paper with some comments and counterexamples to a few seemingly natural generalizations of Theorem 1 and 3.

• Symmetry of (ne, se) for fillings with  $\mathbf{r} \in \{0, 1\}^*$ .

A southeast chain, or shortly se-chain, of size k in a 01-filling M is a set of k cells

$$\{(i_1, j_k), (i_2, j_{k-1}), \dots, (i_k, j_1)\}$$

with  $i_1 < \cdots < i_k$ ,  $j_1 < \cdots < j_k$  filled with 1's such that the  $k \times k$  submatrix

$$\mathcal{G} = \{(i_r, j_s) : 1 \le r \le k, 1 \le s \le k\}$$

is contained in the polyomino. We denote by se(M) the size of the largest se-chains of M. By the symmetry of ne and se we have that Lemma 15 also holds for se.

It is known that in 01-fillings of a Ferrers shape with fixed row sum and column sum in **N**, the pair (ne, se) may not distribute symmetrically (e.g. see [15, 5]). On the other hand, when both  $\mathbf{r}, \mathbf{c} \in \{0, 1\}$ , (ne, se) does have a symmetric joint distribution. This was proved for Ferrers shapes by Krattenthaler [13], and implied for moon polyominoes by Rubey [15, Section 4.2].

The above results raised the question whether for almost-moon polyominoes the pair of statistics (ne, se) has a symmetric joint distribution when one or both of  $\mathbf{r}, \mathbf{c}$  are in  $\{0, 1\}$ , and whether the distribution of (ne, se) is unchanged when one swaps two adjacent rows.

The answers are all negative. In the following we give two set of counterexamples. The first one is for the case that  $\mathbf{r} \in \{0, 1\}$  but  $\mathbf{c} \in \mathbf{N}$ . The involved polyominoes are of small sizes, and we can list all the fillings explicitly. The second is for the case when both  $\mathbf{r}$  and  $\mathbf{c}$  are in  $\{0, 1\}$ . We found the counterexample by running a computer program, and we will just describe the results without listing all the details.

**Example 16.** Figures 6 and 7 list all the 01-fillings of three polyominoes, where the polyominoes in Figure 7 are obtained from the moon polyomino in Figure 6 by moving down the first row. In all the fillings we require that  $\mathbf{r} = (1, 1, 1, 1)$  and  $\mathbf{c} = (2, 1, 1)$ . The data (ne, se) is given under each filling.

Figure 6 shows that even for moon polyomino with  $\mathbf{r} \in \{0, 1\}$  but  $\mathbf{c} \in \mathbb{N}$ , the distribution of (ne, se) is not necessarily symmetric. Figure 7 gives an example that the distribution of the pair (ne, se) is not preserved when two adjacent rows are swapped in an almost-moon polyomino. Note that the first two fillings (M, M') in Figure 7 are coupling fillings, whose corresponding coupling fillings are the first two fillings (N, N') in the second row of Figure 7. For these two pairs Lemma 15 does not hold for (ne, se), i.e.,

$$\{(ne(M), se(M)), (ne(M'), se(M')\} \neq \{(ne(N), se(N)), (ne(N'), se(N'))\}.$$



Figure 6: Fillings of a moon polyomino with  $\mathbf{r} = (1, 1, 1, 1)$  and  $\mathbf{c} = (2, 1, 1)$ .



Figure 7: Fillings of almost-moon polyominoes. The first row is for an almost-moon polyomino  $\mathcal{M}$ , and the second row is for  $\mathcal{N}$  which differ from  $\mathcal{M}$  by swapping the second and the third rows.

**Example 17.** Our second example is restricted to 01-fillings where each row as well as each column has exactly one 1. We say that such fillings are restricted. Let  $\mathcal{M}_1$ ,  $\mathcal{M}_2$  and  $\mathcal{M}_3$  be the polyominoes

whose cells are given by

$$\mathcal{M}_1 = \{(i,j): 1 \le i \le 6, 1 \le j \le 6\} - \{(1,6)\}.$$
  
$$\mathcal{M}_2 = \{(i,j): 1 \le i \le 6, 1 \le j \le 6\} - \{(1,5)\}.$$
  
$$\mathcal{M}_3 = \{(i,j): 1 \le i \le 6, 1 \le j \le 6\} - \{(1,4)\}.$$

Each polyomino has 600 restricted fillings. Let  $G_1(x, y) = \sum_M x^{\operatorname{ne}(M)} y^{\operatorname{se}(M)}$  be the joint distribution of (ne, se) over restricted fillings of  $\mathcal{M}_1$ . Similarly define  $G_2(x, y)$  and  $G_3(x, y)$  for restricted fillings in  $\mathcal{M}_2$  and  $\mathcal{M}_3$ . With the help of a computer program we obtained

$$G_1(x,y) = G_2(x,y) = xy^5 + x^5y + 72(x^2y^4 + x^4y^2) + 48(x^3y^4 + x^4y^3) + 50(x^2y^3 + x^3y^2) + 8(x^2y^5 + x^5y^2) + 242x^3y^3$$
(9)

and

$$G_{3}(x,y) = xy^{5} + x^{5}y + 72x^{2}y^{4} + 73x^{4}y^{2} + 48x^{3}y^{4} + 47x^{4}y^{3} + 50x^{2}y^{3} + 49x^{3}y^{2} + 8x^{2}y^{5} + 8x^{5}y^{2} + 243x^{3}y^{3}.$$
 (10)

Equation (9) is symmetric with respect to x, y, which is expected for  $\mathcal{M}_1$  since it is a moon polyomino. Equation (10) shows that the joint distribution of (ne, se) over almost-moon polyominoes is not necessarily symmetric, even if we require that every row and every column has exactly one 1. The difference between the two equations implies that the distribution of (ne, se) may not be preserved when two adjacent rows are swapped.

• Coupling fillings with  $\mathbf{r} \in \mathbb{N}^*$ .

Another question is whether we can extend the idea of *coupling* in Theorem 3 to construct a bijection for Theorem 1. The following example shows that the direct application does not work.

**Example 18.** Consider fillings in  $\mathcal{F}(\mathcal{M}, *, \mathbf{c})$  where a row may have multiple 1s. For a filling M of  $\mathcal{M}$ , we couple with it the filling M' obtained from M by swapping the fillings  $\alpha$  and  $\beta$ , just as what we did in Section 4. Again let  $N = f_{\mathcal{M},\mathcal{N}}(M')$  and  $N' = f_{\mathcal{M},\mathcal{N}}(M)$ . In the fillings shown in Figure 8,  $\operatorname{ne}(M) = 3$ ,  $\operatorname{ne}(M') = 4$  while  $\operatorname{ne}(N) = \operatorname{ne}(N') = 4$ . Lemma 15 does not hold for this case.



Figure 8: Coupling fillings in  $\mathcal{F}(\mathcal{M}, \mathbf{r}, \mathbf{c})$  with  $\mathbf{r} \in \mathbb{N}^*$  and  $\mathbf{r} \in \{0, 1\}^*$ . The circled dots form 4-chains in the polynominoes.

It is still open whether Theorem 3 holds for the family of fillings that  $\mathbf{r} \in \mathbb{N}^*$  but  $\mathbf{c} \in \{0, 1\}^*$  We point out that Lemma 15 does not hold in this case either. Given a filling M with multiple 1-cells



Figure 9: Coupling fillings in  $\mathcal{F}(\mathcal{M}, \mathbf{r}, \mathbf{c})$  with  $\mathbf{r} \in \mathbb{N}^*$  and  $\mathbf{c} \in \{0, 1\}^*$ .

in a row, the natural way to define the coupling with the same row sum is to let M' be obtained from M by keeping the empty columns of  $\alpha \cup \beta$  and reversing the fillings in the remaining columns of  $\alpha \cup \beta$ . Then (N, N') are obtained from (M, M') by exchanging the two rows  $\mathcal{R}_s$  and  $R_l$  with their fillings.

**Example 19.** The fillings in Figure 9 gives an example where ne(M) = 2, ne(M') = 3 while ne(N) = ne(N') = 3.

#### • Statistic ne in fillings of general polyominoes

It is natural to ask whether Theorem 1 or Theorem 3 can be extended to a more general family of polyominoes. The following example shows that if there are more than one exceptional rows in the polyomino, then the distribution of ne(M) may not be the same after a swap of two adjacent rows.

**Example 20.** This example was first given in [14] for layer polyominoes, which are polyominoes that are row-convex and row-intersection-free. The right one is almost-moon but the left one is not. Let  $G(x, \mathcal{F}) = \sum_{M \in \mathcal{F}} x^{\operatorname{ne}(M)}$  be the generating function for the statistic ne over the fillings in a set  $\mathcal{F}$ . First consider 01-fillings where every row and every column has exactly one 1. We obtained the following generating functions. For the left polyomino  $\mathcal{M}_1$ ,

$$G(x, \mathcal{F}(\mathcal{M}_1, \mathbf{r}, \mathbf{c})) = x + 37x^2 + 31x^3 + 3x^4$$

while for the right polyomino  $\mathcal{M}_2$ ,

$$G(x, \mathcal{F}(\mathcal{M}_2, \mathbf{r}, \mathbf{c})) = p + 36x^2 + 32x^3 + 3x^4,$$

where  $\mathbf{r} = \mathbf{c} = (1, 1, 1, 1, 1)$ . This provides a counterexample of Theorem 3 for general polyominoes.

The above example also implies that Theorem 1 can not hold for general polyominoes. Namely, take the same two polyominoes and consider all the fillings in which each column has exactly one 1, but there is no constraint on the row. Note that



Figure 10: Two general polyominoes related by an interchange of two adjacent rows.

- Empty rows do not affect the statistic ne and can be ignored.
- Any sub-polyomino of  $\mathcal{M}_1$  or  $\mathcal{M}_2$  containing three rows is an almost-moon polyomino. By Theorem 1 rearranging rows for such three-row polyominoes does not change the distribution of ne( $\mathcal{M}$ ) over  $\mathcal{F}(\mathcal{M}, *, \mathbf{c})$ .

Now let  $\mathbf{c} = (1, 1, 1, 1, 1)$  we have

- 1. Either  $G(\text{ne}, \mathcal{F}(\mathcal{M}_1, *, \mathbf{c})) \neq G(\text{ne}, \mathcal{F}(\mathcal{M}_2, *, \mathbf{c}))$  and we have the desired counterexample, or
- 2.  $G(\operatorname{ne}, \mathcal{F}(\mathcal{M}_1, *, \mathbf{c})) = G(\operatorname{ne}, \mathcal{F}(\mathcal{M}_2, *, \mathbf{c}))$ . But then the example from the previous paragraph implies that the distribution of ne over 01-fillings in  $\mathcal{F}(\mathcal{M}_1, *, \mathbf{c})$  and  $\mathcal{F}(\mathcal{M}_2, *, \mathbf{c})$  with empty rows are different. Note that the set of fillings of a polyomino  $\mathcal{M}$  with empty rows can be obtained as the union of set  $\mathcal{F}_i(\mathcal{M}, \mathbf{r}, \mathbf{c})$ , which consists of fillings in which the *i*-th row is empty. An application of the inclusion-exclusion principle implies that there is a subpolyomino  $\mathcal{N}_1$  of  $\mathcal{M}_1$  consisting of 4 rows such that  $G(\operatorname{ne}, \mathcal{F}(\mathcal{N}_1, *, \mathbf{c})) \neq G(\operatorname{ne}, \mathcal{F}(\mathcal{N}_2, *, \mathbf{c}))$ where  $\mathcal{N}_2$  is the sub-polyomino of  $\mathcal{M}_2$  consisting of the same four rows as in  $\mathcal{N}_1$ .

# Acknowledgment

We thank Hua Peng for writing a computer program to help us get the counterexamples in Example 17.

## References

- J. Backelin, J. West, and G. Xin. Wilf-equivalence for singleton classes. Adv. in Appl. Math., 38(2):133–148, 2007.
- [2] W. Y. C. Chen, E. Y. P. Deng, R. R. X. Du, R. P. Stanley and C. H. Yan, Crossings and nestings of matchings and partitions, *Trans. Amer. Math. Soc.*, 359 (2007), 1555–1575.
- [3] W. Y. C. Chen, S. Poznanović, C. H. Yan and A. L. B. Yang, Major index for 01-fillings of moon polyominoes, J. Combin. Theory Ser. A, 117 (2010), 1058–1081.
- [4] W. Y. C. Chen, A. Y. Z. Wang, C. H. Yan and A. F. Y. Zhao, Mixed statistics on 01-fillings of moon polyominoes, SIAM J. Discrete Math., 24 (2010), 1272–1290.

- [5] A. de Mier, On the symmetry of the distribution of crossings and nestings in graphs, Electron. J. Combin., 13 (2006), N21.
- [6] A. de Mier, k-noncrossing and k-nonnesting graphs and fillings of Ferrers diagrams, Combinatorica, 27 (2007), 699–720.
- [7] S. V. Fomin. The generalized Robinson-Schensted-Knuth correspondence. Zap. Nauchn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI), 155(Differentsialnaya Geometriya, Gruppy Li i Mekh. VIII):156–175, 195, 1986.
- [8] S. V. Fomin. Duality of graded graphs. J. Algebraic Combin., 3(4):357–404, 1994.
- [9] S. V. Fomin. Schensted algorithms for dual graded graphs. J. Algebraic Combin., 4(1):5–45, 1995.
- [10] J. Jonsson, Generalized triangulations and diagonal-free subsets of stack polyominoes, J. Combin. Theory Ser. A, 112 (2005), 117–142.
- [11] J. Jonsson and V. Welker, A spherical initial ideal for Pfaffians, *Illinois J. Math.*, 51 (2007), 1397–1407.
- [12] A. Kasraoui, Ascents and descents in 01-fillings of moon polyominoes, European J. Combin., 31 (2010), 87–105.
- [13] C. Krattenthaler, Growth diagrams, and increasing and decreasing chains in fillings of Ferrers shapes, Adv. in Appl. Math., 37 (2006), 404–431.
- [14] M. Phillipson, C. H. Yan and J. Yeh, 2-Chains in 01-fillings of layer polyominoes, *Electron. J. Combin.*, 20(2013), R51.
- [15] M. Rubey, Increasing and decreasing sequences in fillings of moon polyominoes, Adv. in Appl. Math., 47 (2011), 57–87.
- [16] A. Wang and C. H. Yan, Positive and negative chains in charged moon polyominoes, Adv. in Appl. Math., 51(2013), 467–482.