## Rational Solutions of First-Order Algebraic Ordinary Difference Equations

Thieu N. Vo $,^*$ Yi Zhang $^\dagger$ 

February 5, 2019

We propose an algebraic geometric approach for studying rational solutions of first-order algebraic ordinary difference equations (AO $\Delta$ Es). For an autonomous first-order AO $\Delta$ E, we give an upper bound for the degrees of its rational solutions, and thus derive a complete algorithm for computing corresponding rational solutions.

*Keywords*— algebraic ordinary difference equations; strong rational general solutions; parametrization; separable difference equation; resultant theory; algorithms

### 1 Introduction

An algebraic ordinary difference equation (AO $\Delta E$ ) is a difference equation of the form

$$F(x, y(x), y(x+1), \cdots, y(x+m)) = 0,$$

where F is a nonzero polynomial in  $y(x), y(x+1), \dots, y(x+m)$  with coefficients in the field  $\mathbb{K}(x)$ of rational functions over an algebraically closed field  $\mathbb{K}$  of characteristic zero, and  $m \in \mathbb{N}$ . We say that an AO $\Delta E$  is *autonomous* if the independent variable x does not appear in it explicitly. For computational purpose, we may choose  $\mathbb{K} = \overline{\mathbb{Q}}$ , the field of algebraic numbers. AO $\Delta E$ s naturally appear from various problems, such as symbolic summation [30, 24], factorization of linear difference operators [10], analysis of time or space complexity of computer programs with recursive calls [35]. Thus, to determine (closed form) solutions of a given AO $\Delta E$  is a fundamental problem in difference algebra and is of general interest.

<sup>\*</sup>Fractional Calculus, Optimization and Algebra Research Group, Faculty of Mathematics and Statistics, Ton Duc Thang University, Ho Chi Minh City, Vietnam. Email: vongocthieu@tdtu.edu.vn

<sup>&</sup>lt;sup>†</sup>Department of Mathematical Sciences, The University of Texas at Dallas (UTD), USA. Supported by the UT Dallas Program: P-1-03246. Email: zhangy@amss.ac.cn

Constructive approaches for finding symbolic solutions of linear difference equations and their applications have been extensively investigated. There are well-known algorithms for computing polynomial [4], rational [1, 2, 3, 36] and hypergeometric [5, 8, 17, 23, 28, 27, 26, 29, 37] solutions for linear difference equations. Besides, Karr [20, 21], Kauers and Schneider [22] developed algorithms for determining closed form formulas for finite sums and indefinite nested summation. The corresponding implementations are available in computer algebra systems Maple and Mathematica. For more general types of solutions of linear difference equations, we refer to [19, 42]. Last but not least, Bronstein [9] and Schneider [32] gave algorithms for calculating solutions of parameterized linear difference equations within  $\Pi\Sigma$ -fields.

Nevertheless, there are only a few steps in attacking the problem of computing symbolic solutions of (nonlinear) AO $\Delta$ E. Cohn [11] provided a general algebraic framework for investigating structures of AO $\Delta$ Es and their solutions. Elaydi [13] summarized some useful techniques for transforming certain nonlinear AO $\Delta$ E (such as difference equations of general Riccati type) into linear ones. In [16], the authors gave a polynomial time algorithm for finding polynomial solutions of first-order autonomous AO $\Delta$ Es by utilizing parametrization theory of plane algebraic curves. Using symmetric polynomial theory, Shkaravska and Eekelen [34, 35] gave a degree bound for polynomial solutions of high-order non-autonomous AO $\Delta$ Es under a sufficient condition.

We are mainly interested in rational solutions of first-order AO $\Delta$ Es. In [16], Feng, Gao and Huang proposed an algorithm for computing a rational solution for a first-order autonomous AO $\Delta$ E provided that a bound for the degree of the rational solution is given. They also pointed out that they could not bound the degrees of rational solutions through the parametrization technique because the difference version of [15, Theorem 3.7] is not always true (see [16, Example 4.1]). We overcome this missing part and present an algorithm for computing such a degree bound. It is seen that if  $y(x) \in \mathbb{K}(x)$  is a nonzero rational solution of an autonomous first-order AO $\Delta$ E, then  $y(x+c) \in \mathbb{K}(x,c) \setminus \mathbb{K}(x)$ , where c is a constant in a difference extension of  $\mathbb{K}$ , is again a solution of the given difference equation. In this paper, we consider a more general problem:

#### Problem 1.1.

Let  $F \in \mathbb{K}[x, y, z]$  be an irreducible polynomial. Determine a solution  $s \in \mathbb{K}(x, c) \setminus \mathbb{K}(x)$ , where c is a transcendental constant, for the following difference equation

$$F(x, y(x), y(x+1)) = 0.$$
 (1)

A solution in  $\mathbb{K}(x,c)\setminus\mathbb{K}(x)$  is called a *strong rational general solution* (compare with [16, Definition 2.2]). We prove (Theorem 2.2) that if the difference equation (1) admits a strong rational general solution then its corresponding algebraic curve in  $\mathbb{A}^2(\overline{\mathbb{K}(x)})$  defined by F(x, y, z) = 0 is of genus zero. Thus, we can take use of parametric representations of rational curves to transform the original difference equation into an associated difference equation which is of simpler form (see Theorem 2.2 and Proposition 2.4). The latter difference equation has a special form and it is called a *separable difference equation* (Remark 2.8). We prove (Theorem 2.5) that there is a one-to-one corresponding between the strong rational general solutions of the given difference equation and those of the associated separable difference equation. Therefore, the problem of determining a strong rational general solution for a first-order AO $\Delta E$  is reduced to that of computing a strong rational general solution of the corresponding separable one.

For an autonomous first-order AO $\Delta E$ , we give a bound for the degrees of rational solutions of its associated separable difference equation. Thus, we derive a complete algorithm

for computing rational solutions of autonomous first-order AO $\Delta$ Es (see Algorithm 4.2). To derive a degree bound, we first transform the problem of determining a rational solution of an autonomous separable AO $\Delta$ E to that of computing a pair of polynomial solutions of an autonomous first-order system (Subsection 3.2). Secondly, we use tools from resultant theory to eliminate one dependent variable from the system (Subsection 3.3), and then obtain a nontrivial autonomous homogeneous second-order AO $\Delta$ E for the other dependent variable. By using the difference analog of the combinatorial approach in [39, 41], we finally present a degree bound for polynomial solutions of an autonomous homogeneous second-order AO $\Delta$ E (Subsection 3.4), which is the last key for deriving the complete algorithm.

The rest of the paper is organized as follows. Section 2 is devoted to present an algebraic geometric approach to first-order AO $\Delta$ Es. We propose a constructive approach in Section 3 and a complete algorithm in Section 4 for computing rational solutions of autonomous first-order AO $\Delta$ Es.

### 2 An algebraic geometric approach to first-order AO $\Delta$ Es

In this section, we study first-order AO $\Delta$ Es from an algebraic geometric point of view. The idea is inherited from [14, 15, 6, 40]. Assume that  $F \in \mathbb{K}[x, y, z]$  is a nonzero trivariate polynomial. We associate the difference equation F(x, y(x), y(x+1)) = 0 with the corresponding algebraic curve in the two dimensional affine plane over the field  $\mathbb{K}(x)$  of algebraic functions defined by F(x, y, z) = 0. We prove that if the given difference equation admits a strong rational general solution then the corresponding algebraic curve is of genus zero. Therefore we may apply algebraic tools from the theory of rational curves. In particular, we use rational parametric representations of the algebraic curve to transform the original difference equation to a simpler one (see Remark 2.8). A one-to-one correspondence between strong rational general solutions of the original difference equation and those of the new one is established.

We start this section with a formal definition of strong rational general solution.

#### Definition 2.1. (See [40, Definition 3.3])

Let  $F \in \mathbb{K}[x, y, z]$  be a nonzero trivariate polynomial. A solution s of the algebraic difference equation F(x, y(x), y(x + 1)) = 0 is called a strong rational general solution if  $s = s(x, c) \in \mathbb{K}(x, c) \setminus \mathbb{K}(x)$  for some constant c which is transcendental over  $\mathbb{K}(x)$ .

The following theorem gives a necessary condition for a first-order algebraic difference equation having a strong rational general solution.

#### Theorem 2.2.

Let F be an irreducible polynomial in  $\mathbb{K}[x, y, z] \setminus \mathbb{K}[x, y]$  and consider the difference equation F(x, y(x), y(x+1)) = 0. If the difference equation admits a strong rational general solution, then

- (i) F is irreducible as a polynomial in  $\overline{\mathbb{K}(x)}[y, z]$ , and
- (ii) the algebraic curve in  $\mathbb{A}^2\left(\overline{\mathbb{K}(x)}\right)$  defined by F(x, y, z) = 0 is of genus zero.

*Proof.* (i) Let s(x,c) be a strong rational solution of the difference equation F = 0. Consider the ring homomorphism:

$$\begin{array}{cccc} \phi: & \overline{\mathbb{K}(x)}[y,z] & \longrightarrow & \overline{\mathbb{K}(x)}(c) \\ & & G(x,y,z) & \longmapsto & G(x,s(x,c),s(x+1,c)) \end{array}$$

Let I be the kernel of  $\phi$ . By assumption, we know that  $F \in I$ . Thus, I is nonzero. To prove that F is irreducible over  $\overline{\mathbb{K}(x)}$ , we show that I is a principal prime ideal, and that F generates I.

Since the zero set of I contains a parametric class of points (s(x, c), s(x+1, c)), the ideal I is neither a maximal ideal nor the whole ring. On the other hand, the homomorphism  $\phi$  induces an isomorphism between  $\overline{\mathbb{K}(x)}[y, z]/I$  and a subring of  $\overline{\mathbb{K}(x)}(c)$ , which is an integral domain. Therefore, the ideal I is a prime ideal. Since the Krull dimension of  $\overline{\mathbb{K}(x)}[y, z]$  is 2, it follows that I is of height 1. By [18, prop.1.12A, p.7], we conclude that I is principal.

Next, using the technique of Gröbner basis, we construct a generator of I with coefficients in  $\mathbb{K}(x)$ . In order to do that, we rewrite I as the following form:

$$I = \left\{ H \in \overline{\mathbb{K}(x)}[y, z] \,|\, H(x, s(x, c), s(x+1, c)) = 0 \right\}.$$

Let  $s(x,c) = \frac{P(x,c)}{Q(x,c)}$  and  $s(x+1,c) = \frac{R(x,c)}{S(x,c)}$  where  $P,Q,R,S \in \mathbb{K}[x,c]$  such that gcd(P,Q) = gcd(R,S) = 1. By the technique of implicitization [12, Thm.2, p.138], we know that

$$I = \langle yQ - P, zS - R, 1 - QSt \rangle \cap \overline{\mathbb{K}(x)}[y, z],$$

where the first component J in the right hand side is an ideal in  $\mathbb{K}(x)[c, t, y, z]$  generated by polynomials yQ - P, zS - R and 1 - QSt. Let  $\succ$  be the lexicographic order for monomials of  $\overline{\mathbb{K}(x)}[c, t, y, z]$  such that  $c \succ t \succ y \succ z$ . Using Buchberger's algorithm, one can determine a reduced Gröbner basis  $\mathbf{G}$  for J with respect to  $\succ$ . Then  $\mathbf{G}$  only contains polynomials in c, t, y, z with coefficients in  $\mathbb{K}(x)$ . After discarding all polynomials involving c, t from  $\mathbf{G}$ , we obtain a reduced Gröbner basis  $\tilde{\mathbf{G}}$  for I which contains only polynomials in  $\mathbb{K}(x)[y, z]$ . Since I is principal and prime, the basis  $\tilde{\mathbf{G}}$  contains only one element, say  $G \in \mathbb{K}(x)[y, z]$ , and G is irreducible over  $\overline{\mathbb{K}(x)}$ .

Recall that F is irreducible over  $\mathbb{K}(x)$  and  $F \in I = \langle G \rangle$ . This implies that F differs from G by a multiplication of a nonzero element in  $\mathbb{K}(x)$ . Hence, F is also irreducible over  $\overline{\mathbb{K}(x)}$ .

(ii) As a consequence, the algebraic equation F(x, y, z) = 0 defines an irreducible algebraic curve in the affine plane  $\mathbb{A}^2(\overline{\mathbb{K}(x)})$ . Moreover, this curve can be parametrized by the pair of rational functions (s(x, c), s(x+1, c)). Hence, by [33, Theorem 4.7, p.93], the curve is rational. We conclude from [33, Theorem 4.11, p.95] that its genus is zero.

The above theorem motivates the following concept.

#### Definition 2.3.

Let F be a nonzero polynomial in  $\mathbb{K}[x, y, z]$ . The algebraic curve  $\mathcal{C}_F \subset \mathbb{A}^2(\overline{\mathbb{K}(x)})$  defined by F(x, y, z) = 0 is called the corresponding algebraic curve of the first-order algebraic difference equation F(x, y(x), y(x+1)) = 0.

Due to Theorem 2.2, if a first-order AO $\Delta E$  admits a strong rational general solution, then its corresponding algebraic curve must be of genus zero. Therefore, we may apply algebraic tools from parametrization theory of rational curves.

Given a field  $\mathbb{L}$  of characteristics zero and a non-constant polynomial  $G \in \mathbb{L}[y, z]$ , the algebraic equation G(y, z) = 0 implicitly defines an algebraic curve, say  $\mathcal{C}_G$ , in the affine plane  $\mathbb{A}^2(\overline{\mathbb{L}})$  over the algebraic closure  $\overline{\mathbb{L}}$  of  $\mathbb{L}$ . It is well-known that  $\mathcal{C}_G$  is of genus zero if and only if there exists a birational transformation

$$\begin{array}{rcl} \mathcal{P}: & \mathbb{A}^1(\overline{\mathbb{L}}) & \to & \mathcal{C}_G \subset \mathbb{A}^2(\overline{\mathbb{L}}) \\ & t & \mapsto & (p_1(t), p_2(t)) \end{array}$$

for some rational functions  $p_1, p_2 \in \overline{\mathbb{L}}(t)$ . To construct such a birational transformation is one of the most important problems in parametrization theory of algebraic curves. For details about parametrization theory of algebraic curves, we refer to [33]. In particular, there is an algorithm for determining a birational transformation from the affine line to a genus zero algebraic curve over the field  $\mathbb{L} = \mathbb{K}(x)$  (see [38, 40]).

The following proposition is a direct consequence of [40, Proposition 4.3].

#### Proposition 2.4.

Let  $F \in \mathbb{K}(x)[y,z]$  be a non-constant polynomial such that the corresponding algebraic curve  $\mathcal{C}_F \subset \mathbb{A}^2(\overline{\mathbb{K}(x)})$  defined by F(x,y,z) = 0 is of genus zero. Then there exists a birational transformation  $\mathcal{P} : \mathbb{A}^1(\overline{\mathbb{K}(x)}) \to \mathcal{C}_F$  defined by  $\mathcal{P}(x,t) = (p_1(x,t), p_2(x,t))$  for some rational functions  $p_1(x,t), p_2(x,t) \in \mathbb{K}(x,t)$ .

*Proof.* Choose the birational transformation  $\mathcal{P}$  to be an optimal parametrization of  $\mathcal{C}_F$  (see [40, Section 4]).

Note that there exists an algorithm [40, Algorithm 1] for determining a birational transformation (or more precisely, an optimal parametrization) for  $C_F$  in the above proposition. Assume that  $r \in \mathbb{K}[x]$ . The degree of r is defined to be the maximum of the degree of its numerator and that of its denominator with respect to x, and we denote it by  $\deg_x(r)$  or simply  $\deg(r)$ .

#### Theorem 2.5.

Let F(x, y(x), y(x + 1)) = 0 be an  $AO\Delta E$  such that its corresponding curve  $C_F$  is of genus zero. Assume that  $\mathcal{P}(x, t) = (p_1(x, t), p_2(x, t)) \in \mathbb{K}(x, t)^2$  is a birational transformation from the affine line  $\mathbb{A}^1(\overline{\mathbb{K}(x)})$  to  $C_F$ . Consider the following difference equation

$$p_1(x+1,\omega(x+1)) = p_2(x,\omega(x)).$$
(2)

- 1. If  $s(x,c) \in \mathbb{K}(x,c) \setminus \mathbb{K}(x)$  is a strong rational general solution of the given difference equation F(x, y(x), y(x+1)) = 0, then there exists a strong rational general solution  $\omega(x,c) \in \mathbb{K}(x,c) \setminus \mathbb{K}(x)$  of (2) such that  $s(x,c) = p_1(x, \omega(x,c))$ .
- 2. Conversely, if  $\omega(x,c) \in \mathbb{K}(x,c) \setminus \mathbb{K}(x)$  is a strong rational general solution of (2), then

$$s(x,c) = p_1(x,\omega(x,c)) \in \mathbb{K}(x,c) \setminus \mathbb{K}(x)$$

is a strong rational general solution of F(x, y(x), y(x+1)) = 0.

*Proof.* 1. Assume that  $s(x,c) \in \mathbb{K}(x,c) \setminus \mathbb{K}(x)$  is a strong rational general solution of the given difference equation, i.e.,

$$F(x, s(x, c), s(x + 1, c)) = 0.$$

Then  $\mathcal{Q}(x,t) = (s(x,t), s(x+1,t))$  is a parametric representation for  $\mathcal{C}_F$ . By [33, Lemma 4.17, p. 97], if follows that there exists a function  $\omega(x,t) \in \overline{\mathbb{K}(x)}(t)$  such that  $\mathcal{Q}(x,t) = \mathcal{P}(x, \omega(x,t))$ . In particular, we have

$$s(x,t) = p_1(x,\omega(x,t)), \text{ and } s(x+1,t) = p_2(x,\omega(x,t)).$$

Therefore, we have  $p_1(x+1, \omega(x+1,t)) = p_2(x, \omega(x,t))$ . Thus, it follows that  $\omega(x,c)$  is a solution of the difference equation (2), and  $s(x,c) = p_1(x, \omega(x,c))$ . Due to the proof of [33, Lemma 4.17, p.19], we can choose  $\omega(x,t) = \mathcal{P}^{-1} \circ \mathcal{Q}(x,t)$ . Hence, we conclude that  $\omega(x,c) \in \mathbb{K}(x,c)$ . Since  $\deg_c(s) > 0$  and  $s(x,c) = p_1(x, \omega(x,c))$ , it follows that  $\omega(x,c) \in \mathbb{K}(x,c) \setminus \mathbb{K}(x)$ .

2. Assume that  $\omega(x,c) \in \mathbb{K}(x,c) \setminus \mathbb{K}(x)$  is a solution of the difference equation (2), i.e.,

$$p_1(x+1,\omega(x+1,c)) = p_2(x,\omega(x,c)).$$

Since  $\mathcal{P}$  is a birational transformation of  $\mathcal{C}_F$ , we have  $F(x, p_1(x, t), p_2(x, t)) = 0$ . Substituting t by  $\omega(x, c)$ , we obtain

$$0 = F(x, p_1(x, \omega(x, c)), p_2(x, \omega(x, c)))$$
  
=  $F(x, p_1(x, \omega(x, c)), p_1(x + 1, \omega(x + 1, c))).$ 

Set  $s(x,c) = p_1(x,\omega(x,c))$ . It follows from the above equalities that s(x,c) is a rational solution of F(x,y(x),y(x+1)) = 0. Since  $\deg_t(p_1) > 0$  and  $\deg_c(\omega) > 0$ , it follows from [7, Propostion 1.2, item 11] that  $\deg_c(s) > 0$ . In other word, we have  $s(x,c) \in \mathbb{K}(x,c) \setminus \mathbb{K}(x)$ .

The above theorem motivates the following definition.

#### Definition 2.6.

Using notations in Theorem 2.5, we call equation (2) the associated difference equation of F(x, y(x), y(x + 1)) = 0.

From experiments, we find that the associated difference equation is usually simpler (see Example 2.9) than the original one. In next sections, we will present a degree bound for rational solutions of the associated difference equation of an autonomous first-order AO $\Delta E$ , and thus derive a complete algorithm for determining rational solutions of the original one. The existence of an upper bound for rational solutions of non-autonomous first-order AO $\Delta E$  is still open.

The following proposition is a generalization of [16, Lemma 4.2], which refines the shapes of both the AO $\Delta E$  with strong rational general solutions and the associated one.

#### Proposition 2.7.

Let  $F \in \mathbb{K}[x, y, z] \setminus \mathbb{K}[x, y]$  be an irreducible polynomial. If the algebraic difference equation F(x, y(x), y(x+1)) = 0 admits a strong rational general solution, then we have that  $\deg_y F = \deg_z F$ . Furthermore, in this case, the associated difference equation exists and it must be of the form

$$p_1(x, \omega(x+1)) = p_2(x, \omega(x)),$$

for some rational functions  $p_1, p_2 \in \mathbb{K}(x, y)$  such that

$$\deg_u p_1 = \deg_u p_2 = \deg_z F = \deg_u F.$$

*Proof.* By item (ii) of Theorem 2.2, the corresponding algebraic curve  $C_F$  is of genus zero. By Proposition 2.4, there exists a birational transformation  $\mathcal{P}$  from  $\mathbb{A}^1(\overline{\mathbb{K}(x)})$  to  $\mathcal{C}_F$  defined by  $\mathcal{P}(x,t) = (p_1(x,t), p_2(x,t))$  for some  $p_1(x,t), p_2(x,t) \in \mathbb{K}(x,t)^2$ . Due to [33, Theorem 4.21, p.98], we have

 $\deg_t p_1 = \deg_z F, \quad \text{and} \quad \deg_t p_2 = \deg_u F. \tag{3}$ 

By Theorem 2.5, we know that the associated difference equation also admits a strong rational general solution, say  $\omega(x,c) \in \mathbb{K}(x,c) \setminus \mathbb{K}(x)$ , *i.e.*,

$$p_1(x+1,\omega(x+1,c)) = p_2(x,\omega(x,c)).$$

By [7, Proposition 1.2, item 11], it follows that the degrees of the rational functions in both sides of the above equation with respect to c is equal to

$$\deg_t p_1 \cdot \deg_c \omega(x+1,c) = \deg_t p_2 \cdot \deg_c \omega(x,c).$$

Since  $\deg_c \omega(x+1,c) = \deg_c \omega(x,c) \ge 1$ , we have  $\deg_t p_1 = \deg_t p_2$ . Therefore, we conclude from (3) that  $\deg_t p_1 = \deg_t p_2 = \deg_z F = \deg_y F$ .

#### Remark 2.8.

As a consequence of Theorem 2.5 and Proposition 2.7, in order to solve Problem 1.1, we only need to consider the class of difference equations of the form

$$P(x, y(x+1)) = Q(x, y(x)).$$
(4)

for some rational functions  $P, Q \in \mathbb{K}(x, z)$  such that  $\deg_z P = \deg_z Q$ , and determine their strong rational general solutions. We call (4) a (rational) separable difference equation. If furthermore P and Q are in  $\mathbb{K}(z)$ , then we call (4) the autonomous separable difference equation.

#### Example 2.9.

Consider the following non-autonomous first-order  $AO\Delta E$ :

$$\begin{aligned} F(x,y(x),y(x+1)) &= & (y(x)+x^2+2x+1)y(x+1)^2 - \\ & & (y(x)+2x^2+2x)y(x)y(x+1)+x^2y(x)^2 = 0. \end{aligned}$$

The corresponding algebraic curve in the affine plane  $\mathbb{A}^2(\overline{\mathbb{K}(x)})$  is defined by

$$F(x, y, z) = (y + x^{2} + 2x + 1)z^{2} - (y + 2x^{2} + 2x)yz + x^{2}y = 0.$$

This curve is of genus zero and it admits the following optimal parametrization:

$$\mathcal{P}(x,t) = (p_1(x,t), p_2(x,t)) = \left(\frac{(xt-1)^2}{t}, \frac{(xt-1)^2}{t(t+1)}\right)$$

The associated separable difference equation is

$$\frac{((x+1)\omega(x+1)-1)^2}{\omega(x+1)} = \frac{(x\omega(x)-1)^2}{\omega(x)(\omega(x)+1)}.$$

By a detailed greatest common divisors argument, we can show that  $\omega = \frac{1}{x+c}$  is a strong rational general solution of the above equation, where c is an arbitrary constant. Therefore, it follows from Theorem 2.5 that  $y(x) = p_1(x, \omega(x)) = \frac{c^2}{x+c}$  is a strong rational general solution of F(x, y(x), y(x+1)) = 0.

### 3 Rational solutions of autonomous first-order AO $\Delta$ Es

In this section, we restrict our consideration to the class of autonomous first-order AO $\Delta$ Es and provide a constructive approach for solving the following problem.

#### Problem 3.1.

Let  $F \in \mathbb{K}[y, z] \setminus \mathbb{K}[y]$  be an irreducible polynomial. Find all rational solutions of the difference equation F(y(x), y(x+1)) = 0 if there is any.

We first apply the results from the previous section to reduce an autonomous first-order AO $\Delta E$  to an autonomous separable difference equation. Next, we prove (Proposition 3.5 and Theorem 3.7) that the problem of finding rational solutions of an autonomous separable difference equation can be transformed to that of determining polynomial solutions of a difference system of order one in two dependent variables. Using resultant theory, one can always eliminate one dependent variable from that difference system, and thus obtain a nontrivial (Theorem 3.15 and Corollary 3.17) autonomous homogeneous second-order AO $\Delta E$  with respect to the other dependent variable. Finally, we give a degree bound (Proposition 3.20) for polynomial solutions of that second-order difference equation.

#### 3.1 Reduce to an autonomous separable difference equation

Let us consider the following autonomous first-order AO $\Delta E$ :

$$F(y(x), y(x+1)) = 0,$$
(5)

where  $F \in \mathbb{K}[y, z] \setminus \mathbb{K}[y]$  is an irreducible polynomial. Assume that  $y(x) \in \mathbb{K}(x)$  is a rational solution. First, we observe that every constant function solutions of (5) are solutions of the algebraic equation F(x, x) = 0. Therefore, to avoid triviality, we can assume that the degree of y(x) is at least one. In this case, the function y(x + c) is a strong rational general solution, where c is a transcendental element over  $\mathbb{K}(x)$ . Thus, we can conclude that the problem of finding non-constant rational solutions of an autonomous first-order AO $\Delta E$  is equivalent to that of finding its strong rational general solutions.

Now let us assume that y(x) is a non-constant rational solution of the given difference equation. Then the difference equation admits a strong rational general solution. By Theorem 2.2, the corresponding algebraic curve  $C_F \subset \mathbb{A}^2(\overline{\mathbb{K}(x)})$  defined by F(y,z) = 0 is of genus zero. Since the coefficients of F do not involve x, there exists a birational transformation defined by  $\mathcal{P}(t) = (P(t), Q(t)) \in \mathbb{K}(t) \times \mathbb{K}(t)$  from the affine line  $\mathbb{A}^1(\overline{\mathbb{K}})$  to  $\mathcal{C}_F$ . This birational transformation can be chosen to be an optimal parametrization of the algebraic curve in  $\mathbb{A}^2(\mathbb{K})$  defined by F(y, z) = 0 (see [33]). Furthermore, by Proposition 2.7, we have deg  $P = \deg Q = \deg_y F$ . By Theorem 2.5, we conclude that finding rational solutions of the associated autonomous separable difference equation P(y(x+1)) = Q(y(x)) of (5) is enough for solving Problem 3.1.

## 3.2 Reduce to the problem of finding polynomial solutions of an autonomous first-order difference system

Based on arguments of the previous subsection, we now restrict our consideration further to the class of autonomous separable difference equations.

#### Problem 3.2.

Let  $P_1, P_2, Q_1, Q_2$  be polynomials in  $\mathbb{K}[z] \setminus \{0\}$  such that  $gcd(P_1, Q_1) = gcd(P_2, Q_2) = 1$  and  $deg \frac{P_1}{Q_1} = deg \frac{P_2}{Q_2} = n \ge 1$ . Find all rational solutions of the difference equation

$$\frac{P_1(y(x+1))}{Q_1(y(x+1))} = \frac{P_2(y(x))}{Q_2(y(x))}.$$
(6)

If n = 1, then the difference equation (6) is of general Riccati type (see [13, Section 2.6]). A difference equation of general Riccati type can be solved by transforming it into a second-order linear difference equation with polynomial coefficients and then solving the latter one. Details about the transformation can be found in [13, Section 2.6]. Unfortunately, this method can not be generalized immediately to the arbitrary degree case.

We give a new method that works for the general case. The first step of this approach is to reduce Problem 3.2 to that of determining polynomial solutions of a system of autonomous first-order AO $\Delta$ Es (see Problem 3.10). For doing that, we first need some technical lemmas.

#### Lemma 3.3.

Let  $P, Q, R, S \in \mathbb{K}[x]$  be nonzero polynomials such that gcd(P,Q) = gcd(R,S) = 1 and  $\frac{P}{Q} = \frac{R}{S}$ . Then there exists a constant c in  $\mathbb{K}$  such that P = cR and Q = cS.

Proof. Straightforward.

#### Lemma 3.4.

Let  $R, S \in \mathbb{K}[z, w]$  be homogeneous polynomials such that gcd(R, S) = 1, and A, B be polynomials in  $\mathbb{K}[x]$  such that gcd(A, B) = 1. Then

$$gcd\left(R(A(x), B(x)), S(A(x), B(x))\right) = 1.$$

*Proof.* To avoid triviality, we may assume that R and S are of degrees at least 1. We first prove the lemma for the case deg  $R = \deg S = 1$ . In this case, without loss of generality, we can assume that R = z + aw and S = bz + cw for some  $a, b, c \in \mathbb{K}$  such that  $(b, c) \neq (0, 0)$ . Since  $\gcd(R, S) = 1$ , we have  $c - ab \neq 0$ . Then we have

$$gcd(R(A, B), S(A, B)) = gcd(R(A, B), S(A, B) - bR(A, B))$$
$$= gcd(A + aB, (c - ab)B)$$
$$= gcd(A + aB, B) = gcd(A, B) = 1.$$

Next, we assume that R and S have positive degrees. Let deg R = m, deg S = n. Since  $\mathbb{K}$  is algebraically closed, the homogeneous polynomials R and S can be factored into the products of linear homogeneous polynomials, say

$$R = R_1^{r_1} \cdot \ldots \cdot R_m^{r_m}$$
, and  $S = S_1^{s_1} \cdot \ldots \cdot S_n^{s_n}$ ,

where  $R_i, S_j$  are linear homogeneous polynomials in  $\mathbb{K}[z, w]$  and  $r_i, s_j$  are positive integers. Since R and S are coprime,  $R_i$  and  $S_j$  are also coprime for every i, j. Using the above argument, we have that  $gcd(R_i(A, B), S_j(A, B)) = 1$  for every i, j. Hence, we conclude that gcd(R(A, B), S(A, B)) = 1.

#### Proposition 3.5.

Using notations in Problem 3.2, we set

$$\tilde{P}_i(z,w) = w^n P_i\left(\frac{z}{w}\right), \quad and \quad \tilde{Q}_i(z,w) = w^n Q_i\left(\frac{z}{w}\right),$$

which are homogeneous polynomials of degree n in  $\mathbb{K}[z, w]$ , i = 1, 2. Assume that the function  $\frac{A(x)}{B(x)}$  is a solution of equation (6), where  $A, B \in \mathbb{K}[x]$  and gcd(A, B) = 1. Then there exists a constant  $c \in \mathbb{K}$  such that

$$\begin{cases} \tilde{P}_1(A(x+1), B(x+1)) = c \cdot \tilde{P}_2(A(x), B(x)), \\ \tilde{Q}_1(A(x+1), B(x+1)) = c \cdot \tilde{Q}_2(A(x), B(x)). \end{cases}$$
(7)

*Proof.* Substituting  $\frac{A(x)}{B(x)}$  into (6) and clearing numerators and denominators of the both sides, we obtain

$$\frac{P_1(A(x+1), B(x+1))}{\tilde{Q}_1(A(x+1), B(x+1))} = \frac{P_2(A(x), B(x))}{\tilde{Q}_2(A(x), B(x))}.$$
(8)

Next, we prove that the numerators and the denominators of the right hand side for the above equation are coprime. By Lemma 3.4, it suffices to prove that  $\tilde{P}_2$  and  $\tilde{Q}_2$  are coprime polynomials in  $\mathbb{K}[z,w]$ . Since deg  $\frac{P_2}{Q_2} = n$ , without loss of generality, we can assume that deg  $P_2(z) = m$ , deg  $Q_2(z) = n$  and  $m \leq n$ . In this case, we have

$$\tilde{P}_2(z,w) = w^{n-m} \cdot \left[ w^m P_2\left(\frac{z}{w}\right) \right], \text{ and } \tilde{Q}_2(z,w) = w^n Q_2\left(\frac{z}{w}\right).$$
(9)

Note that  $w^m P_2\left(\frac{z}{w}\right)$  and  $w^n Q_2\left(\frac{z}{w}\right)$  are homogenizations of  $P_2(z)$  and  $Q_2(z)$ , respectively. Since  $P_2(z)$  and  $Q_2(z)$  are coprime, we have  $w^m P_2\left(\frac{z}{w}\right)$  and  $\tilde{Q}_2(z,w)$  are coprime. Furthermore, on account of deg  $Q_2(z) = n$ , we see that  $\tilde{Q}_2(z,w) = az^n + wQ(z,w)$  for some  $a \in \mathbb{K} \setminus \{0\}$  and  $Q(z,w) \in \mathbb{K}[z,w]$ . This implies that  $w^{n-m}$  and  $\tilde{Q}_2(z,w)$  are coprime, too. Hence, we conclude from (9) that  $\tilde{P}_2(z,w)$  and  $\tilde{Q}_2(z,w)$  are coprime.

Similarly, we can also show that the numerator and the denominator of the left hand side of (8) are coprime. Therefore, it follows from Lemma 3.3 that the claim of this proposition holds.

It is straightforward to see that any pair of solutions (A(x), B(x)) of (7) gives rise to a rational solution of (6). Therefore, it suffices to consider polynomial solutions of (7). However, there is a new unknown constant appeared in (7). In order to determine the exact values for the constant c in the above proposition, we introduce the following definition.

#### Definition 3.6.

Using notations in Problem 3.2, we rewrite  $P_1, Q_1, P_2, Q_2 \in \mathbb{K}[z]$  as follows:

$$P_{1}(z) = r_{1,high} \cdot z^{k_{1,high}} + \dots + r_{1,low} \cdot z^{k_{1,low}},$$
  

$$Q_{1}(z) = s_{1,high} \cdot z^{\ell_{1,high}} + \dots + s_{1,low} \cdot z^{\ell_{1,low}},$$
  

$$P_{2}(z) = r_{2,high} \cdot z^{k_{2,high}} + \dots + r_{2,low} \cdot z^{k_{2,low}},$$
  

$$Q_{2}(z) = s_{2,high} \cdot z^{\ell_{2,high}} + \dots + s_{2,low} \cdot z^{\ell_{2,low}},$$

where  $k_{i,high} > k_{i,low}$ , and  $\ell_{i,high} > \ell_{i,low}$ , i = 1, 2. Define a set  $\mathcal{C} \subseteq \mathbb{K}$  associated to (6) recursively as follows:

- (1) Set  $C = \{c \in \mathbb{K} \mid \exists \alpha \in \mathbb{K} : P_1(\alpha) = cP_2(\alpha) \text{ and } Q_1(\alpha) = cQ_2(\alpha)\}.$
- (2) If  $k_{1,high} = k_{2,high}$  then add  $\frac{r_{1,high}}{r_{2,high}}$  and 0 to C,
- (3) If  $l_{1,high} = l_{2,high}$  then add  $\frac{s_{1,high}}{s_{2,high}}$  and 0 to C,
- (4) If  $k_{1,low} = k_{2,low}$  then add  $\frac{r_{1,low}}{r_{2,low}}$  and 0 to C,
- (5) If  $l_{1,low} = l_{2,low}$  then add  $\frac{s_{1,low}}{s_{2,low}}$  and 0 to  $\mathcal{C}$ .

We call C the set of constant candidates of (6).

#### Theorem 3.7.

The constant c in Proposition 3.5 is exactly one element in the set of constant candidates of (6).

*Proof.* We keep the notations of Proposition 3.5 and Definition 3.6. Let  $\frac{A(x)}{B(x)}$  be a solution of (6), where A(x) and B(x) are the same as that in Proposition 3.5. Without loss of generality, we may further assume that the leading coefficient of A(x) is  $\alpha \in \mathbb{K} \setminus \{0\}$  and that of B(x) is 1. Let  $P_1, P_2, Q_1, Q_2$  be polynomials in Definition 3.6. And C is the set of constant candidates of (6). Consider the following three cases.

**Case 1.** deg  $A = \deg B$ . By comparing the leading coefficients of polynomials in system (7), we obtain

$$\begin{cases} \tilde{P}_1(\alpha, 1) = c \cdot \tilde{P}_2(\alpha, 1), \\ \tilde{Q}_1(\alpha, 1) = c \cdot \tilde{Q}_2(\alpha, 1). \end{cases}$$

In other words, we have

$$\begin{cases} P_1(\alpha) = c \cdot P_2(\alpha), \\ Q_1(\alpha) = c \cdot Q_2(\alpha). \end{cases}$$

Hence, it follows from Definition 3.6 that  $c \in \mathcal{C}$  .

**Case 2.** deg A >deg B. The following table shows degrees and leading coefficients of polynomials appearing in system (7).

polynomial	degree	leading coefficient
$\tilde{P}_1(A(x+1), B(x+1))$	$k_{1,\mathbf{high}} \deg A + (n - k_{1,\mathbf{high}}) \deg B$	$r_{1,\mathbf{high}} lpha^{k_{1,\mathbf{high}}}$
$\tilde{Q}_1(A(x+1), B(x+1))$	$\ell_{1,\mathbf{high}} \deg A + (n - \ell_{1,\mathbf{high}}) \deg B$	$s_{1,\mathbf{high}} lpha^{\ell_{1,\mathbf{high}}}$
$ ilde{P}_2(A(x),B(x))$	$k_{2,\mathbf{high}} \deg A + (n - k_{2,\mathbf{high}}) \deg B$	$r_{2,{f high}}lpha^{k_{2,{f high}}}$
$ ilde{Q_2}(A(x),B(x))$	$\ell_{2,\mathbf{high}} \deg A + (n - \ell_{2,\mathbf{high}}) \deg B$	$s_{2,{f high}}lpha^{\ell_{2,{f high}}}$

Consider the following three subcases.

**Subcase 2.1.** deg  $\tilde{P}_1(A(x+1), B(x+1)) \le \deg \tilde{P}_2(A(x), B(x))$ .

This implies that

$$k_{1,\mathbf{high}} \deg A + (n - k_{1,\mathbf{high}}) \deg B \le k_{2,\mathbf{high}} \deg A + (n - k_{2,\mathbf{high}}) \deg B.$$

Or equivalently, we have  $k_{1,\text{high}} \leq k_{2,\text{high}}$ . By comparing the coefficients of terms with degree deg  $\tilde{P}_2(A(x), B(x))$  in the first equation of system (7), we obtain

$$c \cdot r_{2,\mathbf{high}} \alpha^{k_{2,\mathbf{high}}} = \begin{cases} r_{1,\mathbf{high}} \alpha^{k_{1,\mathbf{high}}} & \text{if } k_{1,\mathbf{high}} = k_{2,\mathbf{high}}, \\ 0 & \text{if } k_{1,\mathbf{high}} < k_{2,\mathbf{high}}. \end{cases}$$

Hence, we have

$$c = \begin{cases} \frac{r_{1,\text{high}}}{r_{2,\text{high}}} & \text{if } k_{1,\text{high}} = k_{2,\text{high}}, \\ 0 & \text{if } k_{1,\text{high}} < k_{2,\text{high}}, \end{cases}$$

which belongs to  $\mathcal{C}$ .

Subcase 2.2 deg  $\tilde{Q}_1(A(x+1), B(x+1)) \leq \deg \tilde{Q}_2(A(x), B(x))$ . This inequality is equivalent to  $l_{1,high} \leq l_{2,high}$ . Using the similar argument as that in the above subcase, we obtain

$$c = \begin{cases} \frac{s_{1,\text{high}}}{s_{2,\text{high}}} & \text{if } l_{1,\text{high}} = l_{2,\text{high}}, \\ 0 & \text{if } l_{1,\text{high}} < l_{2,\text{high}}, \end{cases}$$

which belongs to  $\mathcal{C}$ .

**Subcase 2.3.** None of the above two subcases happen. In this case, we have  $k_{1,\text{high}} > k_{2,\text{high}}$  and  $s_{1,\text{high}} > s_{2,\text{high}}$ . Thus, we have

$$n = \deg \frac{P_1}{Q_1} = \max\{k_{1,\text{high}}, s_{1,\text{high}}\}$$
$$> \max\{k_{2,\text{high}}, s_{2,\text{high}}\} = \deg \frac{P_2}{Q_2} = n.$$

This is impossible. Hence, this subcase can not happen.

**Case 3.** deg  $A < \deg B$ . The following table shows degrees and leading coefficients of polynomials appearing in system (7).

polynomial	degree	leading coefficient
$\tilde{P}_1(A(x+1), B(x+1))$	$k_{1,\mathbf{low}} \deg A + (n - k_{1,\mathbf{low}}) \deg B$	$r_{1,\mathbf{low}} \alpha^{k_{1,\mathbf{low}}}$
$\tilde{Q}_1(A(x+1), B(x+1))$	$\ell_{1,\mathbf{low}} \deg A + (n - \ell_{1,\mathbf{low}}) \deg B$	$s_{1,\mathbf{low}} \alpha^{\ell_{1,\mathbf{low}}}$
$ ilde{P}_2(A(x),B(x))$	$k_{2,\mathbf{low}} \deg A + (n - k_{2,\mathbf{low}}) \deg B$	$r_{2,\mathbf{low}} \alpha^{k_{2,\mathbf{low}}}$
$ ilde{Q_2}(A(x),B(x))$	$\ell_{2,\mathbf{low}} \deg A + (n - \ell_{2,\mathbf{low}}) \deg B$	$s_{2,\mathbf{low}} \alpha^{\ell_{2,\mathbf{low}}}$

Consider the following three subcases.

**Subcase 3.1.** deg  $\tilde{P}_1(A(x+1), B(x+1)) \le \deg \tilde{P}_2(A(x), B(x))$ .

This means that

 $k_{1,\mathbf{low}} \deg A + (n - k_{1,\mathbf{low}}) \deg B \le k_{2,\mathbf{low}} \deg A + (n - k_{2,\mathbf{low}}) \deg B.$ 

Or equivalently,  $k_{1,low} \ge k_{2,low}$ . Using a similar argument as that in **Subcase 2.1**, we obtain

$$c = \begin{cases} \frac{r_{1,\mathbf{low}}}{r_{2,\mathbf{low}}} & \text{if } k_{1,\mathbf{low}} = k_{2,\mathbf{low}}, \\ 0 & \text{if } k_{1,\mathbf{low}} > k_{2,\mathbf{low}}, \end{cases}$$

which belongs to  $\mathcal{C}$ .

**Subcase 3.2.** deg  $\tilde{Q}_1(A(x+1), B(x+1)) \le \deg \tilde{Q}_2(A(x), B(x))$ .

This inequality is equivalent to  $l_{1,low} \ge l_{2,low}$ . Using a similar argument as that in **Subcase 2.1**, we have

$$c = \begin{cases} \frac{s_{1,\text{low}}}{s_{2,\text{low}}} & \text{if } s_{1,\text{low}} = s_{2,\text{low}}, \\ 0 & \text{if } s_{1,\text{low}} > s_{2,\text{low}}, \end{cases}$$

which also belongs to  $\mathcal{C}$ .

Subcase 3.3. None of the subcases 3.1 and 3.2 occur.

In this case, we have  $k_{1,low} < k_{2,low}$  and  $l_{1,low} < l_{2,low}$ . Thus, we have  $k_{2,low} \ge 1$ and  $l_{2,low} \ge 1$ . In other words, both polynomials  $P_2(z)$  and  $Q_2(z)$  are divisible by z. This contradicts to the assumption that  $gcd(P_2, Q_2) = 1$ . Hence, this subcase cannot happen.

Above all, we conclude that in each case we always have  $c \in \mathcal{C}$ .

It might happen that the set of constant candidates C of (6) is an infinite set. In this case, there are infinitely many choices for the constant c appearing in the difference system (7). We will see in Proposition 3.8 and 3.9 below that the given difference equation (6) in this case is rather simple and it only has constant solutions.

#### Proposition 3.8.

Let  $P_i(z), Q_i(z) \in \mathbb{K}[z]$  be polynomials in (6), i = 1, 2. Then the set of constant candidates of (6) is an infinite set if and only if  $\frac{P_1(z)}{Q_1(z)} = \frac{P_2(z)}{Q_2(z)}$ .

*Proof.* Assume that C is the set of constant candidates of (6). From Definition 3.6, it is clear that C is an infinite set if and only if the set

$$\bar{\mathcal{C}} = \{ c \in \mathbb{K} \, | \, \exists \alpha \in \mathbb{K} \, : \, P_1(\alpha) = cP_2(\alpha) \text{ and } Q_1(\alpha) = cQ_2(\alpha) \}$$

is an infinite set.

Note that if  $P_1(\alpha) = cP_2(\alpha)$  and  $Q_1(\alpha) = cQ_2(\alpha)$  for some  $\alpha \in \mathbb{K}$ , then we always have

$$\det \begin{pmatrix} P_1(\alpha) & P_2(\alpha) \\ Q_1(\alpha) & Q_2(\alpha) \end{pmatrix} = 0.$$
(10)

For each  $\alpha \in \mathbb{K}$  satisfying (10), since  $gcd(P_2, Q_2) = 1$ , there exists a unique  $c \in \mathbb{K}$  such that  $P_1(\alpha) = cP_2(\alpha)$  and  $Q_1(\alpha) = cQ_2(\alpha)$ . Therefore,  $\overline{C}$  is an infinite set if and only if the algebraic equation

$$\det \begin{pmatrix} P_1(z) & P_2(z) \\ Q_1(z) & Q_2(z) \end{pmatrix} = 0$$

has infinitely many solutions. The latter happens if and only if the left hand side is actually the zero polynomial. In other words, we have  $\frac{P_1(z)}{Q_1(z)} = \frac{P_2(z)}{Q_2(z)}$ .

In (6), if  $\frac{P_1(z)}{Q_1(z)} \neq \frac{P_2(z)}{Q_2(z)}$ , then we can construct  $\overline{C}$  of the above proof in an algorithmic way. There, we can compute the set of constant candidates of (6) in this case.

The following proposition provides an answer for Problem 3.2 if  $\frac{P_1(z)}{Q_1(z)} = \frac{P_2(z)}{Q_2(z)}$ .

#### **Proposition 3.9.**

Let  $f(z) \in \mathbb{K}(z)$  be a non-constant rational function. Then rational solutions of the difference equation f(y(x+1)) = f(y(x)) are only constants in  $\mathbb{K}$ .

Proof. Assume that s(x) is a rational solution of the given difference equation. Then we have f(s(x + 1)) = f(s(x)) for every  $x \in \mathbb{K}$  except finitely many points in  $\mathbb{K}$ . Let us fix a constant  $x_0 \in \mathbb{K}$  such that  $f(s(x_0))$  is defined, and define the new rational function g by  $g(x) = f(s(x)) - f(s(x_0))$ . It is clear that  $g(n + x_0) = 0$  for every  $n \in \mathbb{Z}$  except finitely many points in  $\mathbb{Z}$ . This only happens when g is the zero function. Therefore, there exists a constant  $c \in \mathbb{K}$  such that f(s(x)) - c is the zero function. This means that the function s(x) is a zero of the nonzero rational function  $f(z) - c \in \mathbb{K}(z)$ . Thus, the rational function s(x) is algebraic over  $\mathbb{K}$ . Since  $\mathbb{K}$  is algebraically closed, we conclude that  $s(x) \in \mathbb{K}$ .

Using the above proposition, we obtain a simple answer for Problem 3.2 if  $\frac{P_1}{Q_1} = \frac{P_2}{Q_2}$ . In fact, in this case, every rational solution of the given autonomous separable difference equation is a constant function.

Now we can assume that  $\frac{P_1}{Q_1} \neq \frac{P_2}{Q_2}$ . By Theorem 3.7, the set of constant candidates C of (6) is always a finite set. As we have seen in Proposition 3.5, in order to find a rational solution  $\frac{A(x)}{B(x)}$  of the given separable difference equation (6), we need to determine a pair of polynomials (A(x), B(x)) satisfying the difference system (7) for each c in the finite set C. Therefore, we reduce Problem 3.2 to that of finding polynomial solutions of finitely many difference systems of the form (7). The latter one is addressed in next subsection.

## 3.3 Reduce to the problem of finding polynomial solutions of an autonomous second-order AO $\Delta$ E

In this subsection, we consider the problem of finding polynomial solutions of a difference system of the form (7). Let  $\tilde{P}_2, \tilde{Q}_2 \in \mathbb{K}[z]$  be polynomials and c be the constant in (7). By Theorem 3.7, we can compute all possible values of c by calculating the set of constant candidates of (6). Assume that c is given in (7). Note that if c = 0, then it follows from the fact that  $\mathbb{K}$  is algebraically closed that system (7) only has constant solutions in this case. Therefore, we may further assume that  $c \neq 0$ . Then we can replace  $c\tilde{P}_2$  by  $\bar{P}_2$  and  $c\tilde{Q}_2$  by  $\bar{Q}_2$ . Thus, without loss of generality, we can also assume that c = 1. To be more specific, we focus on the following question.

#### Problem 3.10.

Let  $P_1, P_2, Q_1, Q_2 \in \mathbb{K}[z] \setminus \{0\}$  be polynomials such that  $gcd(P_1, Q_1) = gcd(P_2, Q_2) = 1$  and  $deg \frac{P_1}{Q_1} = deg \frac{P_2}{Q_2} = n \ge 1$ . Set

$$\tilde{P}_i(z,w) = w^n P_i\left(\frac{z}{w}\right), \quad and \quad \tilde{Q}_i(z,w) = w^n Q_i\left(\frac{z}{w}\right),$$

which are homogeneous polynomials of degree n in  $\mathbb{K}[z, w]$ , i = 1, 2. Determine all polynomials  $A(x), B(x) \in \mathbb{K}[x]$  with gcd(A, B) = 1 such that

$$\begin{cases} \tilde{P}_1(A(x+1), B(x+1)) = \tilde{P}_2(A(x), B(x)), \\ \tilde{Q}_1(A(x+1), B(x+1)) = \tilde{Q}_2(A(x), B(x)). \end{cases}$$
(11)

Next, we will present an algorithm for deriving nonzero second-order AO $\Delta$ Es for A(x) and B(x) from system (11), respectively. The technique we use is similar to the prolongation-relaxation approach used in differential algebra (see [31, Section 87-88]).

#### Algorithm 3.11.

Given difference system (11), compute nonzero second-order  $AO\Delta Es$  for A(x) and B(x), respectively.

(1) Let  $I \subseteq \mathbb{K}[w_0, w_1, w_2, z_0, z_1, z_2]$  be the ideal generated by the following polynomials

$$\begin{split} \tilde{P}_1(z_1,w_1) &- \tilde{P}_2(z_0,w_0), \quad \tilde{Q}_1(z_1,w_1) - \tilde{Q}_2(z_0,w_0), \\ \tilde{P}_1(z_2,w_2) &- \tilde{P}_2(z_1,w_1), \quad \tilde{Q}_1(z_2,w_2) - \tilde{Q}_2(z_1,w_1). \end{split}$$

Using Gröbner bases or resultants, compute nonzero elements  $F_A \in I \cap \mathbb{K}[z_0, z_1, z_2]$  and  $F_B \in I \cap \mathbb{K}[w_0, w_1, w_2]$ .

(2) Return  $F_A(A(x), A(x+1), A(x+1)) = 0$  and  $F_B(B(x), B(x+1), B(x+2)) = 0$ .

The termination of the above algorithm is clear. The aim of this subsection is to prove the correctness of the above algorithm. To be more specific, we will prove that one can always derive nonzero second-order AO $\Delta$ Es for A(x) and B(x) from system (11), respectively.

In our arguments, we use polynomial resultants to derive those nonzero second-order AO $\Delta$ Es for A(x) and B(x), respectively. For polynomials P(z) and Q(z) in  $\mathbb{K}[z]$ , we denote by  $\operatorname{Res}_{z}(P(z),Q(z))$  the resultant of P and Q with respect to z. The resultant is a polynomial over  $\mathbb{Z}$  in the coefficients of P and Q. For details about resultant theory, we refer to [12, Chapter 3, §6].

#### Proposition 3.12.

Let f(z, w) and g(z, w) be nonzero polynomials in  $\mathbb{K}[z, w]$ . Let  $R(z) = \operatorname{Res}_w(f, g) \in \mathbb{K}[z]$  be the resultant of f and g with respect to w. Then for any  $c \in \mathbb{K}$  we have

 $R(c) = \operatorname{Res}_w(f(c, w), g(c, w)).$ 

In other words, the resultant computation always commutes with the substitution of polynomials.

*Proof.* It is clear from the definition of resultants (see [12, Definition 2, p.162]).

#### Lemma 3.13.

Let  $P_1, P_2, Q_1, Q_2$  be nonzero polynomials in  $\mathbb{K}[z]$  with  $gcd(P_1, Q_1) = gcd(P_2, Q_2) = 1$ . Set

$$R = \operatorname{Res}_{z_2} \left( P_1(z_1) - w P_2(z_2), Q_1(z_1) - w Q_2(z_2) \right)$$

Then R is a nonzero polynomial in  $\mathbb{K}[w, z_1]$ .

*Proof.* Suppose that R = 0. Due to [12, Proposition 3, p. 163], there exists  $z_{2,0} \in \overline{\mathbb{K}(w, z_1)}$  such that

$$P_1(z_1) = w P_2(z_{2,0}), \text{ and } Q_1(z_1) = w Q_2(z_{2,0}).$$
 (12)

Since  $gcd(P_2, Q_2) = 1$ , either  $P_2(z_{2,0})$  or  $Q_2(z_{2,0})$  is nonzero. Without loss of generality, we can assume that  $Q_2(z_{2,0}) \neq 0$ . From (12), we obtain  $\frac{P_2(z_{2,0})}{Q_2(z_{2,0})} = \frac{P_1(z_1)}{Q_1(z_1)}$ . This implies that  $z_{2,0}$  is a zero of the nonzero rational function  $\frac{P_2(z_2)}{Q_2(z_2)} - \frac{P_1(z_1)}{Q_1(z_1)}$  in  $\mathbb{K}(z_1, z_2)$ . Therefore, we have  $z_{2,0} \in \overline{\mathbb{K}(z_1)}$ . It follows that  $Q_2(z_{2,0}) \in \overline{\mathbb{K}(z_1)}$ . However, since w is transcendental over  $\overline{\mathbb{K}(z_1)}$ , the only possibility for  $Q_1(z_1) = wQ_2(z_{2,0})$  to be true is that  $Q_2(z_{2,0}) = 0$ , a contradiction.  $\Box$ 

#### Lemma 3.14.

Let  $P_i, Q_i$  be nonzero polynomials in  $\mathbb{K}[z]$  with  $gcd(P_i, Q_i) = 1$ , i = 1, 2. Set J to be the ideal of  $\mathbb{K}[w_0, w_1, w_2, z_0, z_1, z_2]$  generated by the following polynomials:

$$w_1 P_1(z_1) - w_0 P_2(z_0), \quad w_1 Q_1(z_1) - w_0 Q_2(z_0), w_2 P_1(z_2) - w_1 P_2(z_1), \quad w_2 Q_1(z_2) - w_1 Q_2(z_1).$$

Then  $J \cap \mathbb{K}[w_0, w_1, w_2] \neq \{0\}.$ 

*Proof.* Set  $p_i = w_i P_1(z_i) - w_{i-1} P_2(z_{i-1})$ ,  $q_i = w_i Q_1(z_i) - w_{i-1} Q_2(z_{i-1})$ , i = 1, 2. By assumption, the ideal J is generated by  $p_1, q_1, p_2, q_2$ . Next, we construct a nonzero polynomial R in  $J \cap \mathbb{K}[w_0, w_1, w_2]$  as follows:

- (1) Set  $R_0 = \operatorname{Res}_{z_0}\left(\frac{p_1}{w_1}, \frac{q_1}{w_1}\right)$  and  $R_2 = \operatorname{Res}_{z_2}\left(\frac{p_2}{w_1}, \frac{q_2}{w_1}\right)$ ;
- (2) Set  $R_1 = \operatorname{Res}_{z_1}(R_0, R_2)$ .

By the above construction, we have  $R_0 \in \mathbb{K}[\frac{w_0}{w_1}, z_1], R_2 \in \mathbb{K}[\frac{w_2}{w_1}, z_1]$ , and  $R_1 \in \mathbb{K}[\frac{w_0}{w_1}, \frac{w_2}{w_1}]$ . By [12, Proposition 5, p. 164], we see that  $R_1$  is a  $\mathbb{K}[\frac{w_0}{w_1}, \frac{w_2}{w_1}, z_0, z_1]$ -linear combination of polynomials  $\frac{p_1}{w_1}, \frac{q_1}{w_1}, \frac{p_2}{w_1}$  and  $\frac{q_2}{w_1}$ , *i.e.*,

$$R_1 = h_1 \cdot \frac{p_1}{w_1} + h_2 \cdot \frac{q_1}{w_1} + h_3 \cdot \frac{p_2}{w_1} + h_4 \cdot \frac{q_2}{w_1}$$

for some  $h_i \in \mathbb{K}[\frac{w_0}{w_1}, \frac{w_2}{w_1}, z_0, z_1]$ , i = 1, 2, 3, 4. Then there exists a large enough natural number N such that  $H_i = w_1^{N-1} \cdot h_i \in \mathbb{K}[w_0, w_1, w_2, z_0, z_1, z_2]$  for i = 1, 2, 3, 4. Set  $R = w_1^N \cdot R_1$ . Then we have

$$R = H_1 \cdot p_1 + H_2 \cdot q_1 + H_3 \cdot p_2 + H_4 \cdot q_2$$

for some  $H_i \in \mathbb{K}[w_0, w_1, w_2, z_0, z_1, z_2]$ , i = 1, 2, 3, 4. Therefore, we have  $R \in J$ . On the other hand, since  $R_1 \in \mathbb{K}[\frac{w_0}{w_1}, \frac{w_2}{w_1}]$  and  $R = w_1^N \cdot R_1$ , we conclude that  $R \in \mathbb{K}[w_0, w_1, w_2]$ . Next, we prove that  $R \neq 0$ .

Suppose that R = 0. Then we have  $R_1 = 0$ . By [12, Proposition 3, p.163], there exists  $y_{1,0} \in \overline{\mathbb{K}(w_0, w_1, w_2)}$  such that  $R_0(w_0, w_1, y_{1,0}) = R_2(w_0, w_2, y_{1,0}) = 0$ . It follows from Lemma 3.13 that  $R_0 \in \mathbb{K}[\frac{w_0}{w_1}, y_1] \setminus \{0\}$  and  $R_2 \in \mathbb{K}[\frac{w_2}{w_1}, y_1] \setminus \{0\}$ . Therefore, we see that  $y_{1,0} \in \overline{\mathbb{K}(\frac{w_0}{w_1})} \cap \overline{\mathbb{K}(\frac{w_2}{w_1})}$ . Moreover, since  $\frac{w_0}{w_1}$  and  $\frac{w_2}{w_0}$  are algebraically independent over  $\mathbb{K}$ , we have  $\overline{\mathbb{K}(\frac{w_0}{w_1})} \cap \overline{\mathbb{K}(\frac{w_2}{w_1})} = \mathbb{K}$ .

By Proposition 3.12, we have

$$\operatorname{Res}_{y_0}\left(\frac{p_1(w_0, w_1, y_0, y_{1,0})}{w_1}, \frac{q_1(w_0, w_1, y_0, y_{1,0})}{w_1}\right) = R_0(w_0, w_1, y_{1,0}) = 0.$$

This shows that there exists  $y_{0,0} \in \overline{\mathbb{K}(w_0, w_1)}$  such that

$$\frac{p_1(w_0, w_1, y_{0,0}, y_{1,0})}{w_1} = \frac{q_1(w_0, w_1, y_{0,0}, y_{1,0})}{w_1} = 0,$$

or equivalently,

$$P_1(y_{1,0}) = \frac{w_0}{w_1} P_2(y_{0,0}), \text{ and } Q_1(y_{1,0}) = \frac{w_0}{w_1} Q_2(y_{0,0}).$$
 (13)

Since  $gcd(P_2, Q_2) = 1$ , the polynomials  $P_2(y_{0,0})$  and  $Q_2(y_{0,0})$  cannot be both zero functions. Without loss of generality, we can assume that  $Q_2(y_{0,0}) \neq 0$ . From (13), we see that  $\frac{P_1(y_{1,0})}{Q_1(y_{1,0})} = \frac{P_2(y_{0,0})}{Q_2(y_{0,0})}$ . Therefore, the function  $y_{0,0}$  is a zero of the rational function  $\frac{P_1(y_{1,0})}{Q_1(y_{1,0})} - \frac{P_2(z_0)}{Q_2(z_0)} \in \mathbb{K}(z_0)$ . Since  $\mathbb{K}$  is algebraically closed, we conclude that  $y_{0,0} \in \mathbb{K}$ . However, since  $\frac{w_0}{w_1}$  is transcendental over  $\mathbb{K}$ , the only possibility for  $Q_1(y_{1,0}) = \frac{w_0}{w_1}Q_2(y_{0,0})$  to be true is that  $Q_2(y_{0,0}) = 0$ , a contradiction.

#### Theorem 3.15.

Let  $P_i, Q_i$  be polynomials in  $\mathbb{K}[z] \setminus \{0\}$  such that  $gcd(P_i, Q_i) = 1$  and  $deg \frac{P_i}{Q_i} = n \ge 1$ , where i = 1, 2. Set I to be the ideal of  $\mathbb{K}[w_0, w_1, w_2, z_0, z_1, z_2]$  generated by the following polynomials:

$$w_{1}^{n}P_{1}\left(\frac{z_{1}}{w_{1}}\right) - w_{0}^{n}P_{2}\left(\frac{z_{0}}{w_{0}}\right), \quad w_{1}^{n}Q_{1}\left(\frac{z_{1}}{w_{1}}\right) - w_{0}^{n}Q_{2}\left(\frac{z_{0}}{w_{0}}\right),$$
$$w_{2}^{n}P_{1}\left(\frac{z_{2}}{w_{2}}\right) - w_{1}^{n}P_{2}\left(\frac{z_{1}}{w_{1}}\right), \quad w_{2}^{n}Q_{1}\left(\frac{z_{2}}{w_{2}}\right) - w_{1}^{n}Q_{2}\left(\frac{z_{1}}{w_{1}}\right).$$

Then  $I \cap \mathbb{K}[w_0, w_1, w_2] \neq \{0\}.$ 

Proof. Set

$$f_{i} = w_{i}^{n} P_{1}\left(\frac{z_{i}}{w_{i}}\right) - w_{i-1}^{n} P_{2}\left(\frac{z_{i-1}}{w_{i-1}}\right), \quad g_{i} = w_{i}^{n} Q_{1}\left(\frac{z_{i}}{w_{i}}\right) - w_{i-1}^{n} Q_{2}\left(\frac{z_{i-1}}{w_{i-1}}\right),$$

where i = 1, 2. Then I is generated by  $f_1, g_1, f_2, g_2$ . Furthermore, set

$$p_i = w_i P_1(z_i) - w_{i-1} P_2(z_{i-1}), \quad q_i = w_i Q_1(z_i) - w_{i-1} Q_2(z_{i-1}),$$

where i = 1, 2.

Set  $\mathbb{E} = \overline{\mathbb{K}(w_0, w_1, w_2)}$ . Consider the following two algebraic systems over the ground field  $\mathbb{E}$ ,

$$f_1 = g_1 = f_2 = g_2 = 0, (14)$$

and

$$p_1 = q_1 = p_2 = q_2 = 0, (15)$$

where  $z_0, z_1, z_2$  are indeterminates.

First, we prove that the above two algebraic systems have the same consistent property, *i.e.*, (14) has solutions in  $\mathbb{E}^3$  if and only if (15) has solutions in  $\mathbb{E}^3$ . Assume that  $(y_0, y_1, y_2) \in \mathbb{E}^3$  is a solution of system (14), then it is clear to see that

$$\left(\frac{y_0(\sqrt[n]{w_0},\sqrt[n]{w_1},\sqrt[n]{w_2})}{\sqrt[n]{w_0}},\frac{y_1(\sqrt[n]{w_0},\sqrt[n]{w_1},\sqrt[n]{w_2})}{\sqrt[n]{w_1}},\frac{y_2(\sqrt[n]{w_0},\sqrt[n]{w_1},\sqrt[n]{w_2})}{\sqrt[n]{w_2}}\right) \in \mathbb{E}^3$$

is a solution of system (15). Conversely, if  $(y_0, y_1, y_2) \in \mathbb{E}^3$  is a solution of system (15), then

$$(w_0 \cdot y_0(w_0^n, w_1^n, w_2^n), w_1 \cdot y_1(w_0^n, w_1^n, w_2^n), w_2 \cdot y_2(w_0^n, w_1^n, w_2^n)) \in \mathbb{E}^3$$

is a solution of system (14).

Due to Lemma 3.14, we can derive a consequence for system (15) of the form

$$p(w_0, w_1, w_2) = 0$$

for some nonzero polynomial  $p \in \mathbb{K}[w_0, w_1, w_2]$ . Since p is a nonzero element in the ground field  $\mathbb{E}$ , the above equation is equivalent to 1 = 0. Therefore, system (15) has no solution in  $\mathbb{E}^3$ . By the consistent property, so is system (14). Due to the weak version of Hilbert Nullstellensatz [25, Theorem 5.4, p. 33], the ideal of  $\mathbb{K}(w_0, w_1, w_2)[z_0, z_1, z_2]$  generated by I contains 1. In other words, we have

$$1 = h_1 f_1 + h_2 g_1 + h_3 f_2 + h_4 g_2 \tag{16}$$

for some polynomials  $h_1, h_2, h_3, h_4$  in  $\mathbb{K}(w_0, w_1, w_2)[z_0, z_1, z_2]$ . Let  $h \in \mathbb{K}[w_0, w_1, w_2]$  be the common denominator of the coefficients (in  $\mathbb{K}(w_0, w_1, w_2)$ ) of  $h_1, h_2, h_3, h_4$ . Multiplying both sides of (16) by h, we see that  $h \in I$ . Hence, we conclude that h is a nonzero polynomial in  $I \cap \mathbb{K}[w_0, w_1, w_2]$ .

Note that if  $P_i, Q_i$  are polynomials in  $\mathbb{K}[z] \setminus \{0\}$  of degrees at most  $n \geq 1$  such that  $gcd(P_i, Q_i) = 1$  for i = 1, 2, then the claim in the above theorem still holds. However, the condition that  $\deg \frac{P_i}{Q_i} = n \geq 1$  is necessary for the next corollary, where i = 1, 2.

#### Corollary 3.16.

Let I be the ideal defined in Theorem 3.15. Then  $I \cap \mathbb{K}[z_0, z_1, z_2] \neq \{0\}$ .

*Proof.* For  $i \in \{1, 2\}$ , we denote new polynomials  $\hat{P}, \hat{Q}$  in  $\mathbb{K}[z] \setminus \{0\}$  as follows:

$$\hat{P}_i(z) = z^n P_i\left(\frac{1}{z}\right)$$
 and  $\hat{Q}_i(z) = z^n Q_i\left(\frac{1}{z}\right)$ .

Since  $\deg(\frac{P_i}{Q_i}) = n$  and  $\gcd(P_i, Q_i) = 1$ , we have  $\deg(\frac{\hat{P}_i}{\hat{Q}_i}) = n$  and  $\gcd(\hat{P}_i, \hat{Q}_i) = 1$ . The four generators of I can be rewritten in terms of  $\hat{P}_i, \hat{Q}_i$  as

$$z_{1}^{n}\hat{P}_{1}\left(\frac{w_{1}}{z_{1}}\right) - z_{0}^{n}\hat{P}_{2}\left(\frac{w_{0}}{z_{0}}\right), \quad z_{1}^{n}\hat{Q}_{1}\left(\frac{w_{1}}{z_{1}}\right) - z_{0}^{n}\hat{Q}_{2}\left(\frac{w_{0}}{z_{0}}\right),$$
$$z_{2}^{n}\hat{P}_{1}\left(\frac{w_{2}}{z_{2}}\right) - z_{1}^{n}\hat{P}_{2}\left(\frac{w_{1}}{z_{1}}\right), \quad z_{2}^{n}\hat{Q}_{1}\left(\frac{w_{2}}{z_{2}}\right) - z_{1}^{n}\hat{Q}_{2}\left(\frac{w_{1}}{z_{1}}\right).$$

We conclude from Theorem 3.15 that  $I \cap \mathbb{K}[z_0, z_1, z_2] \neq \{0\}$ .

#### Corollary 3.17.

Algorithm 3.11 is correct.

*Proof.* Assume that  $(A(x), B(x)) \in \mathbb{K}[x]^2$  is a solution of (11) with gcd(A, B) = 1. Set

It is clear to see that  $\phi$  is a ring homomorphism. Applying the usual shift operator to system (11), we obtain

$$\begin{cases} \tilde{P}_1(A(x+1), B(x+1)) = \tilde{P}_2(A(x), B(x)), \\ \tilde{Q}_1(A(x+1), B(x+1)) = \tilde{Q}_2(A(x), B(x)), \\ \tilde{P}_1(A(x+2), B(x+2)) = \tilde{P}_2(A(x+1), B(x+1)), \\ \tilde{Q}_1(A(x+2), B(x+2)) = \tilde{Q}_2(A(x+1), B(x+1)). \end{cases}$$

The above system implies that  $I \subseteq \text{ker}(\phi)$ . By Theorem 3.15 and Corollary 3.16, there exist nonzero elements  $F_A \in I \cap \mathbb{K}[z_0, z_1, z_2]$  and  $F_B \in I \cap \mathbb{K}[w_0, w_1, w_2]$ . Therefore, we conclude that  $\phi(F_A) = 0$  and  $\phi(F_B) = 0$ .

From experiments, we observe that resultant computation is much more efficient than that of Gröbner bases in Step 1 of Algorithm 3.11. Since generators of the ideal I of Algorithm 3.11 are homogenous, the output difference equations are also homogenous. We will present an algorithm for computing polynomial solutions of second-order autonomous homogenous AO $\Delta$ Es in the next subsection.

## 3.4 Polynomial solutions of autonomous second-order homogenous AO $\Delta$ Es

In this subsection, we consider the following problem:

#### Problem 3.18.

Let  $F \in \mathbb{K}[y, z, w]$  be a trivariate homogeneous polynomial. Find all polynomial solutions of the difference equation F(y(x), y(x+1), y(x+2)) = 0.

An answer for the above problem is the last key for deriving a complete algorithm for finding rational solutions of an autonomous first-order AO $\Delta E$ . In literature, there exists an algorithm for computing polynomial solutions of absolutely irreducible autonomous first-order AO $\Delta E$ s [16, Section 3] and also a degree bound [34] for polynomial solutions of general AO $\Delta E$ s under a certain sufficient condition. However, none of them can give a direct answer for Problem 3.18. We will give a degree bound for polynomial solutions of autonomous secondorder homogeneous AO $\Delta E$ s, and thus it can be used to derive an algorithm for computing the corresponding polynomial solutions. The approach is a difference analog of that in [39, 41].

Assume that  $F \in \mathbb{K}[y, z, w]$  is a homogenous polynomial of total degree D. Consider the following second-order AO $\Delta E$ :

$$F(y(x), y(x+1), y(x+2)) = 0.$$
(17)

Let  $\Delta y(x) = y(x+1) - y(x)$ . Then we have

$$y(x+1) = \Delta y(x) + y(x)$$
  
$$y(x+2) = \Delta^2 y(x) + 2\Delta y(x) + y(x).$$

Substituting the above two formulae into (17), we obtain the algebraic equation:

$$\tilde{F}(y(x), \Delta y(x), \Delta^2 y(x)) = 0, \tag{18}$$

where

$$F(y, z, w) = F(y, y + z, y + 2z + w)$$

is also a homogenous polynomial of total degree D in  $\mathbb{K}[y, z, w]$ .

For each  $I = (i_1, i_2, i_3) \in \mathbb{N}^3$ , we define  $||I|| = i_1 + i_2 + i_3$ . Now we may write

$$\tilde{F} = \sum_{||I||=D} c_I y^{i_1} z^{i_2} w^{i_3}, \tag{19}$$

where  $c_I \in \mathbb{K}$ . Set

$$\mathcal{E}(\tilde{F}) = \{I \in \mathbb{N}^3 | c_I \neq 0\},\$$
  

$$m(\tilde{F}) = \min\{i_2 + 2i_3 | I \in \mathcal{E}(\tilde{F})\},\$$
  

$$\mathcal{M}(\tilde{F}) = \{I \in \mathcal{E}(\tilde{P}) | i_2 + 2i_3 = m(\tilde{F})\},\$$
  

$$\mathcal{P}_{\tilde{F}}(t) = \sum_{I \in \mathcal{M}(\tilde{F})} c_I t^{i_2} [t(t-1)]^{i_3}.\$$

We call  $\mathcal{P}_{\tilde{F}}(t)$  the *indicial polynomial* of  $\tilde{F}$  (at infinity).

#### Proposition 3.19.

Let  $\mathcal{P}_{\tilde{F}}(t)$  be the indicial polynomial of  $\tilde{F}$  at infinity. Then  $\mathcal{P}_{\tilde{F}}(t) \neq 0$ . Proof. Set  $f(t) = \frac{\mathcal{P}_{\tilde{F}}(t)}{t^{m(\tilde{F})}}$ . Then

$$f(t) = \sum_{I \in \mathcal{M}(\tilde{F})} c_I \left(\frac{t-1}{t}\right)^{i_3}$$

Let  $T = \frac{t-1}{t}$ . Then f(t) is the evaluation of the nonzero univariate polynomial

$$\sum_{I \in \mathcal{M}(\tilde{F})} c_I x^{i_3} \text{ at } T.$$

Since T is transcendental over K, we conclude that f(t) is nonzero. Thus,  $\mathcal{P}_{\tilde{F}}(t) \neq 0$ .

Assume that  $p(x) = \sum_{i=0}^{d} a_i x^i \in \mathbb{K}(c)[x]$  is a nonzero polynomial solution of (18), where c is transcendental over  $\mathbb{K}(x)$ , d and  $a_i$ 's are unkown. Then

$$\Delta p(x) = a_d \cdot d \cdot x^{d-1} + \text{ lower terms in } x,$$
  
$$\Delta^2 p(x) = a_d \cdot d \cdot (d-1) \cdot x^{d-2} + \text{ lower terms in } x.$$

Thus, for each  $I = (i_1, i_2, i_3) \in \mathbb{N}^3$  with ||I|| = D, we have

$$p^{i_1}(\Delta p)^{i_2}(\Delta^2 p)^{i_3} = (a_d x^d)^{i_1} (a_d dx^{d-1})^{i_2} (a_d d(d-1)x^{d-2})^{i_3} + \text{ lower terms in } x$$
$$= a_d^{i_1+i_2+i_3} d^{i_2} [d(d-1)]^{i_3} x^{d(i_1+i_2+i_3)-(i_2+2i_3)} + \text{ lower terms in } x$$
$$= a_d^D d^{i_2} [d(d-1)]^{i_3} x^{dD-(i_2+2i_3)} + \text{ lower terms in } x.$$
(20)

Based on the above argument, we have the following proposition.

#### Proposition 3.20.

Let p(x) be a nonzero polynomial solution of (18) with degree d. Then  $\mathcal{P}_{\tilde{F}}(d) = 0$ . Proof. Since p(x) is a nonzero polynomial solution of (18) with degree d, we have that

$$[x^{dD-m(\tilde{F})}]\left(\tilde{F}(p(x),\Delta p(x),\Delta^2 p(x))\right) = 0$$

By (20), the above equation is equivalent to  $a_d^D \cdot \mathcal{P}_{\tilde{F}}(d) = 0$ . On account of  $a_d \neq 0$ , we conclude that  $\mathcal{P}_{\tilde{F}}(d) = 0$ .

By Proposition 3.19 and the above one, we can compute a degree bound of polynomial solutions of  $\tilde{F}$  by computing non-negative integers solutions of the indicial polynomial at infinity. Afterwards, we may compute the polynomial solutions by making an ansatz and then solving the corresponding algebraic equations by using Gröbner bases.

# 4 Algorithms for computing rational solutions of autonomous first-order AO $\Delta$ Es

Based on the results of the previous section, we summarize an algorithm for determining rational solutions of an autonomous first-order AO $\Delta E$ .

#### Algorithm 4.1.

Given a separable difference equation  $\frac{P_1(y(x+1))}{Q_1(y(x+1))} = \frac{P_2(y(x))}{Q_2(y(x))}$  with  $gcd(P_i, Q_i) = 1$  and  $deg \frac{P_1}{Q_1} = deg \frac{P_2}{Q_2} \ge 1$ , i = 1, 2, compute a bound N for the degrees of its rational solutions.

- 1. Set N = 0. If  $\frac{P_1(z)}{Q_2(z)} = \frac{P_1(z)}{Q_2(z)}$ , then output N. Otherwise, go to the next step.
- 2. Compute the set of constant candidates C of the given separable difference equation by Definition 3.6.
- 3. Let  $c_1, c_2, \ldots, c_m$  be nonzero elements in C. Let

$$\tilde{P}_j(z,w) = w^n P_j\left(\frac{z}{w}\right), \quad and \quad \tilde{Q}_j(z,w) = w^n Q_j\left(\frac{z}{w}\right), \qquad j = 1,2$$

For i from 1 to m do

a) Consider the difference system

$$\begin{cases} \tilde{P}_1(A(x+1), B(x+1)) = c_i \cdot \tilde{P}_2(A(x), B(x)), \\ \tilde{Q}_1(A(x+1), B(x+1)) = c_i \cdot \tilde{Q}_2(A(x), B(x)). \end{cases}$$
(21)

where A, B are unknown functions. Derive the following two nonzero autonomous second-order  $AO\Delta Es$  for A(x) and B(x) from the above equations by using Algorithm 3.11:

$$F_{i,A}(A(x), A(x+1), A(x+2)) = 0,$$
 and  $F_{i,B}(B(x), B(x+1), B(x+2)) = 0,$ 

where  $F_{i,A}$  and  $F_{i,B}$  are homogeneous polynomials in  $\mathbb{K}[y, z, w] \setminus \{0\}$ .

b) Determine the indicial polynomials  $\mathcal{P}_{F_{i,A}}$  and  $\mathcal{P}_{F_{i,B}}$  of  $F_{i,A}$  and  $F_{i,B}$ , respectively. Let

> $D_{i,A} = \{non-negative \text{ integer solutions of } \mathcal{P}_{F_{i,A}}(t)\},\$  $D_{i,B} = \{non-negative \text{ integer solutions of } \mathcal{P}_{F_{i,B}}(t)\}.$

- c) Set  $N = \max\{\{N\} \cup D_{i,A} \cup D_{i,B}\}.$
- 4. Return N.

The termination of the above algorithm is evident. The correctness is a consequence of Proposition 3.9, Theorem 3.7, the correctness of Algorithm 3.11 and Proposition 3.20.

To avoid triviality, we only consider non-constant rational solutions below.

#### Algorithm 4.2.

Given an irreducible autonomous first-order  $AO\Delta E F(y(x), y(x+1)) = 0$ , compute a nonconstant rational solution or return NULL.

- 1. If  $\deg_u(F) \neq \deg_z(F)$ , then output NULL. Otherwise, go to the next step.
- 2. Compute the genus g of the corresponding curve  $C_F$  defined by F(y,z) = 0. If  $g \neq 0$ , then output NULL. Otherwise, go to the next step.
- 3. By using [40, Algorithm 1], determine an optimal parametrization for  $C_F$ , say  $\mathcal{P}(t) = (p_1(t), p_2(t))$ .
- 4. Apply Algorithm 4.1 to compute a bound N for the degrees of rational solutions of the associated separable difference equation  $p_1(y(x+1)) = p_2(y(x))$ .
- 5. Set  $M = N \cdot \deg_t p_1$ . Use [16, Algorithm 4.16] to determine a non-constant rational solution of the given  $AO\Delta E$  whose degree at most M. Return the rational solution if there is any. Otherwise return NULL.

The termination of the above algorithm is clear.

#### Theorem 4.3.

Algorithm 4.2 is correct.

Proof. Assume that the difference equation F(y(x), y(x + 1)) = 0 admits a rational solution. Then it has a strong rational general solution. By Proposition 2.7, we have  $\deg_y F = \deg_z F$ . By Theorem 2.2, the genus of the corresponding curve  $C_F$  is of genus zero. In this case,  $C_F$ admits an optimal parametrization  $\mathcal{P}(t) = (p_1(t), p_2(t)) \in \mathbb{K}(t) \times \mathbb{K}(t)$  and the given difference equation has an associated separable difference equation. By Theorem 2.5, we see that if N is a bound for the degrees of rational solutions of the associated separable difference equation, then  $N \cdot \deg_t p_1$  is a bound for the degrees of rational solutions of the given difference equation. The correctness of step 5 follows from that of [16, Algorithm 4.16].

#### Example 4.4.

Consider the following irreducible autonomous first-order  $AO\Delta E$ :

$$F = (12y(x) + 49)y(x+1)^2 - (12y^2 + 62y + 56)y(x+1) + y(x)^2 + 8y(x) + 16 = 0.$$
 (22)

It is clear to see that  $\deg_y(F) = \deg_z(F) = 2$ . The corresponding algebraic curve is of genus zero and it has an optimal parametrization

$$\mathcal{P}(t) = (p_1(t), p_2(t)) = \left(\frac{9t^2 - 12t + 4}{12t}, \frac{9t^2 + 36t + 4}{12(t+4)}\right)$$

Using the above parametrization, we can derive the following associated separable difference equation of (22):

$$\frac{9y(x+1)^2 - 12y(x+1) + 4}{y(x+1)} = \frac{9y(x)^2 + 36y(x) + 4}{y(x) + 4},$$

where  $P_1(z) = 9z^2 - 12z + 4$ ,  $Q_1(z) = z$ ,  $P_2(z) = 9z^2 + 36z + 4$  and  $Q_2(z) = z + 4$ . It is clear that  $\frac{P_1(z)}{Q_2(z)} \neq \frac{P_1(z)}{Q_2(z)}$ . So we can skip step 1 of Algorithm 4.1. By computation,

It is clear that  $\frac{1}{Q_2(z)} \neq \frac{1}{Q_2(z)}$ . So we can skip step 1 of Algorithm 4.1. By computation, we find that the set of candidate constants  $C = \{7 + 4\sqrt{3}, 7 - 4\sqrt{3}, 0, 1\}$ . Using other steps of Algorithm 4.1, we see that the degree bound for rational solutions of the associated separable difference equation is 2. Thus, the degrees of rational solutions of the given difference equation are bounded by 4. By applying [16, Algorithm 4.1], we can determine a rational solution, say

$$y(x) = \frac{(1 - 4x + 2x^2)^2}{2x(1 - 3x + 2x^2)}$$

## Acknowledgement

We thank Georg Grasegger and Matteo Gallet for valuable suggestions and comments on revising our paper.

### References

- S. A. Abramov. Rational solutions of linear differential and difference equations with polynomial coefficients. USSR Computational Mathematics and Mathematical Physics, 29(6):7–12, 1989.
- [2] S. A. Abramov. Rational solutions of linear difference and q-difference equations with polynomial coefficients. In Proceedings of the 1995 International Symposium on Symbolic and Algebraic Computation, ISSAC '95, pages 285–289, New York, NY, USA, 1995. ACM.
- [3] S. A. Abramov. Rational solutions of first order difference systems. In Proceedings of the 1998 International Symposium on Symbolic and Algebraic Computation, ISSAC '98, pages 124–131, New York, NY, USA, 1998. ACM.
- [4] S. A. Abramov, M. Bronstein, and M. Petkovšek. On polynomial solutions of linear operators equations. In *Proceedings of the 1995 International Symposium on Symbolic and Algebraic Computation*, ISSAC '95, pages 290–295, New York, NY, USA, 1995. ACM.
- [5] S. A. Abramov, P. Paule, and M. Petkovšek. q-hypergeometric solutions of q-difference equations. Discrete Mathematics, 100:3–22, 1998.
- [6] J. M. Aroca, J. Cano, R. Feng, and X. S. Gao. Algebraic general solutions of algebraic ordinary differential equations. In *Proceedings of the 2005 International Symposium on Symbolic and Algebraic Computation*, ISSAC '05, pages 29–36, New York, NY, USA, 2005. ACM.
- [7] F. Binder. Fast computations in the lattice of polynomial rational function fields. In Proceedings of the 1996 International Symposium on Symbolic and Algebraic Computation, ISSAC '96, pages 43–48, New York, NY, USA, 1996. ACM.
- [8] H. Böing and W. Koepf. Algorithms for q-hypergeometric summation in computer algebra. Journal of Symbolic Computation, 28(6):777–799, 1999.
- [9] M. Bronstein. On solutions of linear ordinary difference equations in their coefficient field. Journal of Symbolic Computation, 29(6):841–877, 2000.
- [10] M. Bronstein and M. Petkovšek. An introduction to pseudo-linear algebra. Theoretical Computer Science, 157:3–33, 1996.
- [11] R. M. Cohn. Difference Algebra, volume 17 of Tracts in Mathematics. Interscience, New York, 1965.
- [12] D. Cox, J. Little, and D. O'Shea. Ideals, Varieties, and Algorithms. Springer International Publishing, 4th edition, 2015.

- [13] S. Elaydi. An Introduction to Difference Equations. Undergraduate Texts in Mathematics. Springer New York, 3th edition, 2004.
- [14] R. Feng and X. S. Gao. Rational general solutions of algebraic ordinary differential equations. In Proceedings of the 2004 International Symposium on Symbolic and Algebraic Computation, ISSAC '04, pages 155–162, New York, NY, USA, 2004. ACM.
- [15] R. Feng and X. S. Gao. A polynomial time algorithm for finding rational general solutions of first order autonomous odes. *Journal of Symbolic Computation*, 41(7):739–762, 2006.
- [16] R. Feng, X. S. Gao, and Z. Huang. Rational solutions of ordinary difference equations. Journal of Symbolic Computation, 43(10):746 – 763, 2008.
- [17] R. W. Gosper. Decision procedure for indefinite hypergeometric summation. Proceedings of the National Academy of Sciences of the United States of America, 75(1):40–42, 1978.
- [18] R. Hartshorne. Algebraic Geometry, volume 52 of Graduate Texts in Mathematics. Springer-Verlag, New York Berlin Heidelberg, 1977.
- [19] P. A. Hendriks and M. F. Singer. Solving difference equations in finite terms. Journal of Symbolic Computation, 27(3):239–259, 1999.
- [20] M. Karr. Summation in finite terms. Journal of the Association for Computing Machinery, 28(2):305–350, 1981.
- [21] M. Karr. Theory of summation in finite terms. Journal of Symbolic Computation, 1(3):303–315, 1985.
- [22] M. Kauers and C. Schneider. Indefinite summation with unspecified summands. Discrete Mathematics, 306(17):2073–2083, 2006.
- [23] W. Koepf. Algorithms for m-fold hypergeometric summation. Journal of Symbolic Computation, 20(3):399–417, 1995.
- [24] C. Koutschan. Advanced applications of the holonomic systems approach. PhD thesis, Johannes Kepler University Linz, 2009.
- [25] H. Matsumura. Commutative Algebra, volume 120. WA Benjamin New York, 1970.
- [26] P. Paule and A. Riese. A Mathematica q-analogue of Zeilberger's algorithm based on an algebraically motivated approach to q-hypergeometric telescoping. Special Functions, q-series, and Related Topics, 14:179–210, 1997.
- [27] P. Paule. Greatest factorial factorization and symbolic summation. Journal of Symbolic Computation, 20(3):235–268, 1995.
- [28] P. Paule and M. Schorn. A Mathematica version of Zeilberger's algorithm for proving binomial coefficient identities. *Journal of Symbolic Computation*, 20(5-6):673–698, 1995. Symbolic computation in combinatorics  $\Delta_1$  (Ithaca, NY, 1993).
- [29] M. Petkovšek. Hypergeometric solutions of linear recurrences with polynomial coefficients. Journal of Symbolic Computation, 14(2-3):243–264, 1992.

- [30] M. Petkovšek, H. S. Wilf, and D. Zeilberger. A = B. A K Peters Ltd., Wellesley, MA, 1996. With a foreword by Donald E. Knuth.
- [31] J. F. Ritt. Differential equations from the algebraic standpoint, volume 14. American Mathematical Soc., 1932.
- [32] C. Schneider. Solving parameterized linear difference equations in terms of indefinite nested sums and products. *Journal of Difference Equations and Applications*, 11(9):799– 821, 2005.
- [33] J. R. Sendra, F. Winkler, and S. Pérez-Díaz. Rational Algebraic Curves, A Computer Algebra Approach, volume 22 of Algorithms and Computation in Mathematics. Springer-Verlag, Berlin Heidelberg, 2008.
- [34] O. Shkaravska and M. van Eekelen. Univariate polynomial solutions of algebraic difference equations. *Journal of Symbolic Computation*, 60:15–28, 2014.
- [35] O. Shkaravska and M. van Eekelen. Polynomial solutions of algebraic difference equations and homogeneous symmetric polynomials. 2018.
- [36] M. van Hoeij. Rational solutions of linear difference equations. In Proceedings of the 1998 International Symposium on Symbolic and Algebraic Computation, ISSAC '98, pages 120– 123, New York, NY, USA, 1998. ACM.
- [37] M. van Hoeij. Finite singularities and hypergeometric solutions of linear difference equations. Journal of Pure and Applied Algebra, 139:109–131, 1999.
- [38] M. van Hoeij and J. Cremona. Solving conics over function fields. Journal de Théorie des Nombres de Bordeaux, 18:595–606, 2006.
- [39] T. N. Vo and Y. Zhang. Rational solutions of high-order algebraic ordinary differential equations. arXiv: 1709.04174, pages 1–15, 2018.
- [40] T. N. Vo, G. Grasegger, and F. Winkler. Deciding the existence of rational general solutions for first-order algebraic odes. *Journal of Symbolic Computation*, pages –, 2017. To appear.
- [41] T. N. Vo, G. Grasegger, and F. Winkler. Computation of all rational solutions of firstorder algebraic odes. Advances in Applied Mathematics, 98:1–24, 2018.
- [42] D. A. Wolfram. A formula for the general solutions of a constant-coefficient difference equations. *Journal of Symbolic Computation*, 29(1):79–82, 2000.