

An Efficient Algorithm of Logarithmic Transformation to Hirota Bilinear Form of KdV-type Bilinear Equation

Yichao Ye, Lihong Wang, Zhaowei Chang, Jingsong He*

Department of Mathematics, Ningbo University, Ningbo Zhejiang 315211, P.R. China

Abstract

In this paper, an efficient algorithm of logarithmic transformation to Hirota bilinear form of the KdV-type bilinear equation is established. In the algorithm, some properties of Hirota operator and logarithmic transformation are successfully applied, which helps to prove that the linear terms of the nonlinear partial differential equation play a crucial role in finding the Hirota bilinear form. Experimented with various integro-differential equations, our algorithm is proven to be more efficient than the algorithm referred by Zhou, Fu, and Li in getting the Hirota bilinear form, especially in achieving the coefficient of the logarithmic transformation.

Keywords: Hirota bilinear form, KdV-type bilinear equation, D-operator, Logarithmic transformation, Symbolic computation.

1. Introduction

In 1971, Hirota developed a direct method, Hirota bilinear method, to construct the exact multi-soliton solution of integrable nonlinear partial differential equation (NPDE) [1-3]. Once the Hirota bilinear form(HBF) of a NPDE is given, there are several ways to solve it. The solutions of NPDE can be constructed from HBF by the typical perturbation expansion method [1] and others [4,5]. Therefore, the key step of the Hirota direct method is to transform the NPDE into the Hirota bilinear form (HBF). Hietarinta designed a program on searching for integrable bilinear equations such as the KdV-type [6], mKdV-type [7], SG-type [8], NLS-type [9] equations. In 1992, Hereman and Zhuang [10] gave a summary on types of bilinear equations. In recent years, some algorithms for generating bilinear form of NPDE are described and Maple packages Bilinearization and HBTrans are established by Zhou, Fu, Li [11, 12] and Yang, Ruan [13], respectively. The package Bilinearization can construct the HBF of many NPDEs by solving a system of over-determined algebraic equations with respect to the combinatorial coefficients. However, the general ansatz of the bilinear form of NPDE in Ref. [11, 12], which relies on the **WTC** method [14] and the **HBM** method [15], is so complicated that the efficiency of performance is relatively low. Without depending on these two methods, our method can obtain the HBF of the KdV-type equations within shorter time. In this paper, some properties of Hirota-operator are taken advantage of, which brings forward to a more efficient algorithm for finding

*Corresponding author

Email address: hejingsong@nbu.edu.cn (Jingsong He)

Preprint submitted to Applied Mathematics and Computation

March 21, 2018

the HBF of KdV-type equations in NPDE. We take a series of classic KdV-type equations for instance, to demonstrate the validity of our algorithm. Furthermore, the implementation of the algorithm in Maple is applied to automate the tedious computation for the construction of the HBF of KdV-type equations in NPDE.

2. Hirota Bilinear Method

Now, let us briefly review the Hirota derivatives. In 1971, Hirota developed the Hirota derivative, which is also called D-operator [1]. For $1 + 1$ dimensions, the D-operator is defined by

$$D_x^n D_t^m f \cdot g = (\partial_x - \partial_{x'})^n (\partial_t - \partial_{t'})^m f(x, t) g(x', t') \Big|_{x'=x, t'=t}, \quad m, n = 0, 1, 2, 3, \dots \quad (2.1)$$

where $f(x, t)$ and $g(x, t)$ are differentiable functions of x and t , respectively.

From the definition, there are some properties [1] of the D-operator,

$$D_x^n D_t^m f \cdot g = D_t^m D_x^n f \cdot g = D_x^{n-1} D_t^m D_x f \cdot g \quad (2.2)$$

$$D_x^n D_t^m f \cdot 1 = \partial_x^n \partial_t^m f, \quad \text{where } \partial_x^n \equiv \partial^n / \partial x^n. \quad (2.3)$$

$$D_x^n D_t^m f \cdot g = (-1)^{m+n} D_x^n D_t^m g \cdot f \quad (2.4)$$

$$D_x^n D_t^m f \cdot f = 0, \quad \text{if } m + n \text{ is odd}; \quad (2.5)$$

furthermore, for arbitrary independent variable x , a recursive definition of D-operator is,

$$D_x^n f \cdot g = \begin{cases} fg, & \text{for } n = 0 \\ D_x^{n-1} f_x \cdot g - D_x^{n-1} f \cdot g_x, & \text{for } n > 0. \end{cases} \quad (2.6)$$

As an example, the Korteweg de Vries (KdV) equation

$$u_{xxx} + 6uu_x + u_t = 0 \quad (2.7)$$

where $u = u(x, t)$, can be transformed through the dependent variable transformation,

$$u = 2(\ln f)_{xx} \quad (2.8)$$

into

$$\frac{\partial}{\partial x} \left(\frac{f_{xt}f - f_x f_t + f_{xxxx}f - 4f_{xxx}f_x + 3f_{xx}^2}{f^2} \right) = 0, \quad (2.9)$$

from which the bilinear equation below is obtained

$$f_{xt}f - f_x f_t + f_{xxxx}f - 4f_{xxx}f_x + 3f_{xx}^2 = cf^2, \quad (2.10)$$

where c is a constant of integration. Equation (2.10), with $c = 0$, may also be written concisely in terms of D-operators as

$$(D_x^4 + D_x D_t)f \cdot f = 0. \quad (2.11)$$

Thus, the Hirota bilinear form of the KdV equation is obtained.

3. Principles Of Our Algorithm

3.1. The Properties of Hirota Derivatives

Introducing vector notation

$$\vec{D} = (D_t, D_x, D_y, \dots), \quad (3.1)$$

consider the KdV-type bilinear equation [1]

$$P(D_t, D_x, D_y, \dots) f \cdot f = 0, \quad (3.2)$$

where P is a general polynomial in D_t, D_x, D_y, \dots .

According to (2.4) and (2.5), it follows that

$$P(-\vec{D}) = P(\vec{D}). \quad (3.3)$$

Now assume that the degree of every term in P is even.

When n is even, it's true that

$$\begin{aligned} D_x^n f \cdot f &= \left(\sum_{i=0}^n C_n^i \partial_x^{n-i} (-\partial_{x'}) \right) f(x) f(x') \Big|_{x'=x} \\ &= \sum_{i=0}^n (-1)^i C_n^i (\partial_x^{n-i} f(x)) (\partial_{x'}^i f(x')) \\ &= 2 \sum_{i=0}^{n/2-1} (-1)^i C_n^i (\partial_x^{n-i} f(x)) (\partial_{x'}^i f(x)) + (-1)^{n/2} C_n^{n/2} (\partial_x^{n/2} f(x))^2 \\ &= 2 (\partial_x^n f) f + H, \end{aligned} \quad (3.4)$$

where $H = H(f_x, f_t, \dots)$ is a polynomial in f_x, f_t, \dots , WITHOUT f , and $H = D_x^n f \cdot f - 2 (\partial_x^n f) f$. This simple observation implies the following crucial formula

$$P(D_t, D_x, D_y, \dots) f \cdot f = 2 \left(P(\partial_t, \partial_x, \partial_y, \dots) f \right) f + \tilde{H} = 2 \left(P(\vec{\partial}) f \right) f + \tilde{H}, \quad (3.5)$$

where \tilde{H} is a polynomial in f_x, f_t, \dots , WITHOUT f , and $P(\vec{\partial}) = P(\partial_t, \partial_x, \partial_y, \dots)$. Some simple examples are listed as follows to illustrate equation (3.5)

$$D_x^2 f \cdot f = 2(f_{xx}f - f_x^2), \quad \tilde{H} = -2f_x^2, \quad (3.6)$$

$$D_x^4 f \cdot f = 2(f_{4x}f - 4f_{3x}f_x + 3f_{xx}^2), \quad \tilde{H} = -8f_{3x}f_x + 6f_{xx}^2, \quad (3.7)$$

$$D_x D_t f \cdot f = 2(f_{tx}f - f_t f_x), \quad \tilde{H} = -2f_t f_x, \quad (3.8)$$

where $f_{kx} \equiv \partial_x^k f \equiv \frac{\partial^k f}{\partial x^k}$, $k \in \mathbb{N}$. Furthermore, using equation (3.5), it is easy to find $P(\vec{D})f \cdot f$ is a quadratic polynomial in f, f_x, f_t, \dots , and then to obtain

$$\frac{P(\vec{D})f \cdot f}{f^2} = \frac{P_1}{f} + \frac{P_2}{f^2}. \quad (3.9)$$

Here P_1 and P_2 are polynomials in f_x, f_t, \dots , WITHOUT f . $P_1 = 2P(\vec{\partial})f$, is a linear polynomial; $P_2 = \tilde{H}$ is a quadratic polynomial.

3.2. Logarithmic Transformation

Consider the derivation formula,

$$\left(\frac{f}{g}\right)_x = \frac{f_x g - f g_x}{g^2} = \frac{f_x}{g} - \frac{f g_x}{g^2}. \quad (3.10)$$

Furthermore, for arbitrary independent variable t , it's easy to show that

$$\left(\frac{f_x}{f}\right)_{kt} = \frac{\bar{P}_1}{f} + \frac{\bar{P}_2}{f^2} + \cdots + \frac{\bar{P}_{k+1}}{f^{k+1}}, \quad (3.11)$$

where $k \in \mathbb{N}$ and $\bar{P}_i (i = 1, 2, \dots, k+1)$ is a homogeneous polynomial of degree i in f_x, f_t, \dots , WITHOUT f , and

$$\bar{P}_1 = \partial_t^k f_x, \quad \bar{P}_{k+1} = (-1)^k k! f_x f_t^k. \quad (3.12)$$

are obtained by using equation (3.10) repeatedly.

Furthermore, when $m+n \geq 1$, it shows that

$$\partial_x^n \partial_t^m (\ln f) = \frac{\partial_x^n \partial_t^m f}{f} + \cdots + (-1)^{m+n-1} (m+n-1)! \frac{f_x^n f_t^m}{f^{m+n}} \quad (3.13)$$

Still, by using P as the denotation of a general polynomial and letting the lowest degree of the terms in P be equal to or greater than 1, we have

$$P(\vec{\partial})(\ln f) = \frac{\tilde{P}_1}{f} + \frac{\tilde{P}_2}{f^2} + \cdots + \frac{\tilde{P}_r}{f^r}, \quad (3.14)$$

where $r = \text{degree}(P)$ and $\tilde{P}_1 = P(\vec{\partial})f$. Here, $\tilde{P}_i (i = 1, 2, \dots, r)$ also denotes a homogeneous polynomial of degree i in f_x, f_t, \dots , WITHOUT f .

Using the logarithmic transformation $u = 2 \ln f$, the following expressions are obtained:

$$u_{xx} = 2 \left(\frac{f_{xx}}{f} - \frac{f_x^2}{f^2} \right) = \frac{D_x^2 f \cdot f}{f^2} \quad (3.15)$$

$$u_{xt} = 2 \left(\frac{f_{xt}}{f} - \frac{f_x f_t}{f^2} \right) = \frac{D_x D_t f \cdot f}{f^2} \quad (3.16)$$

$$u_{tt} = 2 \left(\frac{f_{tt}}{f} - \frac{f_t^2}{f^2} \right) = \frac{D_t^2 f \cdot f}{f^2} \quad (3.17)$$

$$u_{4x} = 2 \left(\frac{f_{xxxx}}{f} - \frac{4f_{xxx}f_x + 3f_{x,x}^2}{f^2} + \frac{12f_{xx}f_x^2}{f^3} - \frac{6f_x^4}{f^4} \right) = \frac{D_x^4 f \cdot f}{f^2} - 3 \left(\frac{D_x^2 f \cdot f}{f^2} \right)^2 \quad (3.18)$$

$$u_{6x} = 2 \left(\frac{f_{xxxxxx}}{f} + \cdots - \frac{120f_x^6}{f^6} \right) = \frac{D_x^6 f \cdot f}{f^2} - 15 \frac{D_x^4 f \cdot f}{f^2} \frac{D_x^2 f \cdot f}{f^2} + 30 \left(\frac{D_x^2 f \cdot f}{f^2} \right)^3. \quad (3.19)$$

Clearly, the expressions above can lead us to finding the KdV-type bilinear form associated with logarithmic transformation in NPDE.

3.3. The Relationship Between Logarithmic Transformation and KdV-type Bilinear Equation

In this paper, we consider the logarithmic transformation

$$u = 2\alpha (\ln f)_{nx}, \quad n = 0, 1, 2 \quad (3.20)$$

where $f = f(x, t, \dots)$ and α is a nonzero constant.

Substituting (3.20) to a NPDE

$$W(u, u_x, u_t, \dots) = 0, \quad (3.21)$$

we get

$$\tilde{W}(f, f_x, f_t, \dots; \alpha) \equiv W(u, u_x, u_t, \dots)|_{u=2\alpha(\ln f)_{nx}}. \quad (3.22)$$

If NPDE $\tilde{W}(f, f_x, f_t, \dots; \alpha)$ has the KdV-type bilinear form

$$\left(\frac{P(\vec{D})f \cdot f}{f^2} \right)_{mx} = 0, \quad (3.23)$$

where m is a nonnegative integer, using the properties of Hirota operator and logarithmic transformation, our algorithm will find the undetermined parameters $n, m, P(\vec{D})$, and α .

Because of (3.14), the terms in NPDE with degree k generate the homogeneous expression with logarithmic transformation (3.20)

$$\frac{\tilde{P}_k}{f^k} + \frac{\tilde{P}_{k+1}}{f^{k+1}} + \dots. \quad (3.24)$$

That is, with logarithmic transformation (3.20), the terms in NPDE with degree k do NOT generate the terms $\frac{\tilde{P}_i}{f^i}$ ($i = 1, 2, 3, \dots, k-1$).

Consider KdV-type bilinear form

$$\begin{aligned} \left(\frac{P(\vec{D})f \cdot f}{f^2} \right)_{mx} &= \left(\frac{P_1 f + P_2}{f^2} \right)_{mx} = \left(\frac{P_1}{f} + \frac{P_2}{f^2} \right)_{mx} = \left(\frac{P_1}{f} \right)_{mx} + \left(\frac{P_2}{f^2} \right)_{mx} \\ &= \left(\frac{P_{1mx}}{f} + \dots + \frac{(-1)^m m! P_1 f_x^m}{f^{m+1}} \right) + \left(\frac{P_{2mx}}{f^2} + \dots + \frac{(-1)^m (m+1)! P_2 f_x^m}{f^{m+2}} \right) \\ &= \frac{P_{1mx}}{f} + \dots + \frac{(-1)^m (m+1)! P_2 f_x^m}{f^{m+2}} = \frac{2\partial_x^m (P(\vec{D})f)}{f} + \dots + \frac{(-1)^m (m+1)! P_2 f_x^m}{f^{m+2}}. \end{aligned} \quad (3.25)$$

Therefore, if a equation has the KdV-type bilinear form, with logarithmic transformation (3.20), the term $\frac{P_{1mx}}{f}$ must be generated from the linear terms in NPDE (3.21).

Denoting the linear part in NPDE (3.21) by $P_l(\vec{\partial})u$, with logarithmic transformation (3.20), according to (3.14), the linear part is transformed into

$$P_l(\vec{\partial})u \xrightarrow{u=2\alpha(\ln f)_{nx}} \frac{2\alpha \partial_x^n P_l(\vec{\partial})f}{f} + \dots.$$

Thus, it follows that

$$P(\vec{\partial}) = \alpha \partial_x^{n-m} P_l(\vec{\partial}). \quad (3.26)$$

From the above equation, we find it important that the linear terms of the NPDE play a crucial role in finding the HBF. Hence, from the linear part in NPDE, P , the function of HBF, can be obtained.

Let

$$\left(\frac{P(\mathbf{D})f \cdot f}{f^2} \right)_{mx} = \tilde{W}(f, f_x, f_t, \dots; \alpha), \quad (3.27)$$

then the difference should be zero, namely,

$$Res_{\tilde{W}} \triangleq \tilde{W}(f, f_x, f_t, \dots; \alpha) - \left(\frac{P(\vec{D})f \cdot f}{f^2} \right)_{mx} = 0. \quad (3.28)$$

Clearly, n is a finite enumerated integer and m is also a finite enumerated integer (See **Step 5** of algorithm in **Section 5**) and $P(\vec{\partial}) = \alpha \partial_x^{n-m} P_l(\vec{\partial})$.

With P satisfying (3.26), the coefficient of $\frac{1}{f}$ in $Res_{\tilde{W}}$ is equal to zero. Then the homogeneous expression in f and its derivatives follows as:

$$Res_{\tilde{W}} = \frac{\hat{P}_2}{f^2} + \dots + \frac{\hat{P}_r}{f^r}. \quad (3.29)$$

Here, r is a finite positive integer.

Solving the numerators of (3.29), if there exists the nonzero numerical solution α , the undetermined parameters $n, m, P(\vec{D})$ are also achieved. It can be proven that α has no more than one nonzero numerical solution which satisfies (3.28) with certain n and m . The proof will be given in our next paper.

In summary, if a NPDE has the KdV-type bilinear form associated with the logarithmic transformation (3.20), our algorithm will surely find it.

4. An Example Of our Algorithm

Now, let us take the KdV equation as an example.

$$u_{xxx} + 6uu_x + u_t = 0. \quad (4.1)$$

Here, $P_l(\vec{\partial}) = \partial_t + \partial_x^3$.

First substitute $u = 2\alpha \ln f$ to (4.1) and simplify the equation, the homogeneous equation in f and its derivatives follows as:

$$\frac{2\alpha(f_{xxx} + 12\alpha(\ln f)f_x + f_t)}{f} - \frac{6\alpha f_{xx}f_x}{f^2} + \frac{4\alpha f_x^3}{f^3} = 0.$$

Since the above equation has $\ln(f)$ term, we goto the next logarithmic transformation $u = 2\alpha(\ln f)_x$.

Likewise, substitute $u = 2\alpha(\ln f)_x$ to (4.1) and simplify the equation, the homogeneous equation in f and its derivatives follows as:

$$\frac{2\alpha(f_{xxxx} + f_{tx})}{f} - \frac{2\alpha(-12\alpha f_{xx}f_x + 4f_{xxx}f_x + 3f_{xx}^2 + f_x f_t)}{f^2} + \frac{24\alpha f_x^2(-\alpha f_x + f_{xx})}{f^3} - \frac{12\alpha f_x^4}{f^4} = 0.$$

From the above equation, the coefficient of $\frac{1}{f}$ is acquired: $P_1 = 2\alpha(f_{xxxx} + f_{tx}) = 2\alpha(\partial_x^4 + \partial_x^2\partial_t)f = 2\alpha\partial_x P_l(\vec{\partial})f$. Because the order of derivative is even, the Hirota bilinear form can be obtained: $P(\vec{D})f \cdot f = \alpha(D_x^4 + D_x D_t)f \cdot f$, with which divided by f^2 and subtracts the above equation the difference is as follows:

$$\frac{12f_{xx}\alpha(-2\alpha f_x + f_{xx})}{f^2} - \frac{24\alpha f_x^2(-\alpha f_x + f_{xx})}{f^3} + \frac{12\alpha f_x^4}{f^4} = 0.$$

Solving the numerators of the difference equation, it only has $\alpha = 0$. So we goto the next logarithmic transformation $u = 2\alpha(\ln f)_{xx}$.

Similarly, substitute $u = 2\alpha(\ln f)_{xx}$ to (4.1) and simplify the equation, the homogeneous equation in f and its derivatives follows as:

$$\begin{aligned} & \frac{2\alpha(f_{txx} + f_{xxxx})}{f} - \frac{2\alpha(-12\alpha f_{xxx}f_{xx} + f_{xx}f_t + 5f_{xxxx}f_x + 10f_{xxx}f_{xx}) + 2f_x f_{tx}}{f^2} \\ & - \frac{4f_x\alpha(18\alpha f_{xx}^2 + 6\alpha f_{xxx}f_x - 15f_{xx}^2 - 10f_{xxx}f_x - f_x f_t)}{f^3} + \frac{120\alpha f_{xx}f_x^3(\alpha - 1)}{f^4} - \frac{48\alpha f_x^5(\alpha - 1)}{f^5} = 0. \end{aligned}$$

From the above equation, the coefficient of $\frac{1}{f}$ is obtained: $P_1 = 2\alpha(f_{xxxx} + f_{txx}) = 2\alpha(\partial_x^5 + \partial_x^2\partial_t)f$. Because the order of derivative is odd, we integrate P_1 and get the Hirota bilinear form $P(\vec{D})f \cdot f = \alpha(D_x^4 + D_x D_t)f \cdot f$, with which divided by f^2 and derived in terms of x one time then subtracts the above equation, the difference is obtained:

$$\frac{-24\alpha f_{xxx}f_{xx}(\alpha - 1)}{f^2} + \frac{24f_x\alpha(\alpha - 1)(3f_{xx}^2 + f_{xxx}f_x)}{f^3} - \frac{120\alpha f_{xx}f_x^3(\alpha - 1)}{f^4} + \frac{48\alpha f_x^5(\alpha - 1)}{f^5} = 0.$$

Solving the numerators of the difference equation, a nonzero solution, $\alpha = 1$, is achieved.

In conclusion, by using the logarithmic transformation $u = 2(\ln f)_{xx}$, it shows that the equation (4.1) is equivalent to

$$\left(\frac{(D_x^4 + D_x D_t)f \cdot f}{f^2} \right)_x = 0.$$

This example illustrates our method intuitively.

5. An Algorithm of Bilinear

Consider the general NPDE

$$W(u, u_x, u_t, \dots) = 0,$$

where $u = u(x, t, \dots)$, and W is a polynomial in u and its derivatives.

The algorithm for obtaining HBF goes in the following steps:

Step 1 Let $n = 0$, $f = f(x, t, \dots)$ and α is an undetermined nonzero constant.

Step 2 Using logarithmic transformation $u = 2\alpha(\ln f)_{nx}$, it follows that

$$\widetilde{W} = \widetilde{W}(f, f_x, f_t, \dots) = W(u, u_x, u_t, \dots)|_{u=2\alpha(\ln f)_{nx}}$$

Step 3 If \tilde{W} has $\ln(f)$ term, then goto **Step 6**.

Step 4 Simplify the equation \tilde{W} to get the homogeneous expressions in f and its derivatives

$$\tilde{W} = \frac{P_1}{f} + \frac{P_2}{f^2} + \cdots + \frac{P_r}{f^r},$$

where $P_i = P_i(f_x, f_t, \dots)$ ($i = 1, 2, \dots, r$) is a homogeneous polynomial of degree i in f_x, f_t, \dots , WITHOUT f .

Step 5 Get the coefficient of $\frac{1}{f}$ in \tilde{W} : $P_1 = P_1(f_x, f_t, \dots)$. Let the lowest order of the derivative of f with respect to x in P_1 is k , and the corresponding Hirota bilinear form is $P(D_x, D_t, \dots) f \cdot f$. Then the following equation is obtained:

$$P_1(f_x, f_t, \dots) = 2\partial_x^m P(f_x, f_t, \dots),$$

where $0 \leq m \leq k$ and the degrees of the terms in $P(f_x, f_t, \dots)$ are even. That is,

- If the lowest order derivative of f is even, then $m = 0, 2, 4, \dots$, and $m \leq k$.
- If the lowest order derivative of f is odd, then $m = 1, 3, 5, \dots$, and $m \leq k$.

Step 5.1 Let $m = 0$, if the lowest order of the derivative of f is even, or else let $m = 1$.

Step 5.2 Integrating P_1 by m times in terms of x , we get $P(f_x, f_t, \dots) = \frac{1}{2}\partial_x^{-m} P_1(f_x, f_t, \dots)$. Hence the HBF is $P(D_x, D_t, \dots) f \cdot f$.

Step 5.3 Calculate the difference, $Res_{\tilde{W}} = \tilde{W} - \left(\frac{P(D_x, D_t, \dots) f \cdot f}{f^2} \right)_{mx}$, and simplify the difference $Res_{\tilde{W}}$ to get the homogeneous expressions in f and its derivatives

$$Res_{\tilde{W}} = \frac{\widehat{P}_2}{f^2} + \cdots + \frac{\widehat{P}_r}{f^r}.$$

Step 5.4 Get the equations $\widehat{P}_i = 0$ ($i = 2, 3, \dots, r$), and solve the equations. If a nonzero α exists, then using $u = 2\alpha(\ln f)_{nx}$, the NPDE is equal to $\left(\frac{P(D_x, D_t, \dots) f \cdot f}{f^2} \right)_{mx} = 0$. Therefore, the HBF is $P(D_x, D_t, \dots) f \cdot f$, and exit of program.

Step 5.5 Let $m = m + 2$. If $m \leq k$, goto **Step 5.2**, or else goto **Step 6**.

Step 6 Let $n = n + 1$. If $n \leq 2$ then goto **Step 2**, or else the NPDE has no HBF with logarithmic transformation, end program and exit.

6. Applications

Example 1. Boussinesq equation [16, 17]

$$u_{tt} - u_{xx} - 3u_{xx}^2 - u_{4x} = 0 \quad (6.1)$$

Solution. From the equation, *Bilinear* within 0.s outputs $\alpha = 1$, $u = 2 \ln(f)$ and

$$\frac{(D_t^2 - D_x^2 - D_x^4)f \cdot f}{f^2} = 0.$$

Example 2. Sawada-Kotera equation [18, 19]

$$u_t + 45u_x u^2 - 15u_{xx} u_x - 15u_{xxx} u + u_{5x} = 0 \quad (6.2)$$

Solution. Within 0.032s our program outputs $\alpha = -1$, $u = -2(\ln f)_{xx}$ and

$$\left(\frac{(-D_x^6 - D_x D_t) f \cdot f}{f^2} \right)_x = 0.$$

Example 3. Kadomtsev-Petviashvili equation [20]

$$(u_{3x} + 6uu_x + u_t)_x + 3\delta^2 u_{yy} = 0 \quad (6.3)$$

Solution. *Bilinear* within 0.016s outputs $\alpha = 1$, $u = 2(\ln f)_{xx}$ and

$$\left(\frac{(D_x^4 + D_x D_t + 3\delta^2 D_y^2) f \cdot f}{f^2} \right)_{xx} = 0.$$

Example 4. The shallow water waves equation [21]

$$u_t - u_{xxt} - 3uu_t + 3u_x \int_x^\infty u_t dx + u_x = 0 \quad (6.4)$$

Solution. By substituting $u = w_x$ and using the boundary condition $u_t|_{x \rightarrow \infty} = 0$ (6.4) can be converted into the differential form

$$w_{xt} - w_{3xt} - 3w_x w_{xt} - 3w_{xx} w_t + w_{xx} = 0, \quad (6.5)$$

Bilinear within 0.016s outputs $\alpha = 1$, $w = 2(\ln f)_x$ and

$$\left(\frac{(-D_x^3 D_t + D_x^2 + D_x D_t) f \cdot f}{f^2} \right)_x = 0.$$

Example 5. Ito equation [22]

$$u_{tt} + u_{3xt} + 6u_x u_t + 3uu_{xt} + 3u_{xx} \int_{-\infty}^x u_t dx = 0 \quad (6.6)$$

Solution. By substituting $u = w_x$ and using the boundary condition $u_t|_{x \rightarrow -\infty} = 0$ (6.6) is transformed into the differential form

$$w_{xtt} + w_{4xt} + 6w_{xx} w_{xt} + 3w_x w_{xxt} + 3w_{3x} w_t = 0, \quad (6.7)$$

Bilinear within 0.031s outputs $\alpha = 1$, $w = 2(\ln f)_x$ and

$$\left(\frac{(D_x^3 D_t + D_t^2) f \cdot f}{f^2} \right)_{xx} = 0.$$

Example 6. (2+1)-dimensional breaking soliton equation [23]

$$u_t + \beta u_{xxy} + 4\beta uu_y + 4\beta u_x \int_{-\infty}^x u_y dx = 0 \quad (6.8)$$

Solution. The equation can be written as (6.9) by substituting $u = w_x$ and using the boundary condition $u_y|_{x \rightarrow -\infty} = 0$

$$w_{xt} + \beta w_{3xy} + 4\beta w_x w_{xy} + 4\beta w_{xx} w_y = 0, \quad (6.9)$$

Bilinear within 0.016s outputs $\alpha = \frac{3}{4}$, $w = \frac{3}{2}(\ln f)_x$ and

$$\left(\frac{3(D_x D_t + \beta D_x^3 D_y) f \cdot f}{f^2} \right)_x = 0.$$

Example 7. The Bidirectional SK equation [24]

$$5 \int_{-\infty}^x u_{tt} dx + 5u_{xxt} - 15uu_t - 15u_x \int_{-\infty}^x u_t dx - 45u_x u^2 + 15u_{xx} u_x + 15u_{3x} u - u_{5x} = 0 \quad (6.10)$$

Solution. Similarly, we convert (6.10) into the differential form by substituting $u = w_x$ and using the boundary condition $u_t|_{x \rightarrow -\infty} = 0$, $u_{tt}|_{x \rightarrow -\infty} = 0$

$$5w_{tt} + 5w_{3xt} - 15w_x w_{xt} - 15w_{xx} w_t - 45w_{xx} w_x^2 + 15w_{3x} w_{xx} + 15w_{4x} w_x - w_{6x} = 0, \quad (6.11)$$

Bilinear within 0.015s outputs $\alpha = -1$, $w = -2(\ln f)_x$ and

$$\left(\frac{(-5D_t^2 + D_x^6 - 5D_x^3 D_t) f \cdot f}{f^2} \right)_x = 0.$$

Example 8. (2+1)-dimensional SK equation [25]

$$9u_t + u_{5x} + 15u_{xx} u_x + 15u_{3x} u + 45u_x u^2 - 5 \int_{-\infty}^x u_{yy} dx - 15uu_y - 15u_x \int_{-\infty}^x u_y dx - 5u_{xxy} = 0 \quad (6.12)$$

Solution. With the substitution $u = w_x$ and the boundary condition $u_y|_{x \rightarrow -\infty} = 0$, $u_{yy}|_{x \rightarrow -\infty} = 0$ (6.12) acquires a differential form

$$9w_{xt} + w_{6x} + 15w_{3x} w_{xx} + 15w_{4x} w_x + 45w_{xx} w_x^2 - 5w_{yy} - 15w_x w_{xy} - 15w_{xx} w_y - 5w_{3xy} = 0, \quad (6.13)$$

Bilinear within 0.032s outputs $\alpha = 1$, $w = 2(\ln f)_x$ and

$$\left(\frac{(D_x^6 - 5D_x^3 D_y - 5D_y^2 + 9D_x D_t) f \cdot f}{f^2} \right)_x = 0.$$

Example 9. (3+1)-dimensional KdV equation [13]

$$u_t + 6u_x u_y + u_{xxy} + u_{4xz} + 60u_x^2 u_z + 10u_{3x} u_z + 20u_x u_{xz} = 0 \quad (6.14)$$

Solution. Bilinear within 0.016s outputs $\alpha = \frac{1}{2}$, $u = (\ln f)_x$ and

$$\frac{1}{2} \frac{(D_x D_t + D_x^5 D_z + D_x^3 D_y) f \cdot f}{f^2} = 0.$$

7. The Program Code of Bilinear

```

with(PDEtools): with(DEtools):

## Hirota Bilinear Method
## Bilinear Derivative / Hirota Operator

BD:=proc(FF,DD) local f,g,x,m,opt;
if nargs=1 then return `*(FF[]); fi; f,g:=FF[]; x,m:=DD[];
opt:=args[3..-1]; if m=0 then return procname(FF,opt); fi;
procname([diff(f,x),g],[x,m-1],opt)-procname([f,diff(g,x)], [x,m-1],opt);
end:

`print/BD`:=proc(FF,DD) local f,g,x,m,i; f,g:=FF[];
f:=cat(f, ' ', g); g:=product(D[args[i][1]]^args[i][2], i=2..nargs);
if g<>1 then f:='`*(g)*`*(f); fi; f; end:

## collect(expr,f); first!
getFnumer:=proc(df,f,pow::posint:=1) local i,g,fdenom;
if type(df,'+') then
g:=[op(df)];
fdenom:=map(denom,g);
for i to nops(fdenom) while fdenom[i]<>f^pow do od;
if i>nops(fdenom) then lprint(fdenom);
error "no term(s) or numer=0 when denom=%1",op(0,f)^pow fi;
g:=numer(g[i]);
if not type(expand(g),'+') then lprint(g);
error "Expected more than 1 term about Hirota D-operator" fi;
return g; fi; lprint(df);
error "expected 1st argument be type '+.'"; end:

getvarpow:=proc(df::function) local i,f,var,dif,pow; if
op(0,df)<>diff then lprint(df); error "expected diff function" fi;
f:=convert(df,D); var:=[op(f)]; dif:=[op(op([0,0],f))];
pow:=[0$nops(var)]; f:=op(op(0,f))(var[]); for i to nops(var) do
dif:=selectremove(member,dif,{i});
pow[i]:=nops(dif[1]);
dif:=dif[2];
od; pow:=zip((x,y)->[x,y],var,pow); pow:=remove(has,pow,{0});
[[f,f],pow[]]; end:

#convert to Hirota Bilinear Form
HBF:=proc(df) local i,c,f; if type(df,'+') then
f:=[op(df)]; return map(procname,f); fi;
if type(df,'*') then f:=[op(df)];
f:=selectremove(hasfun,f,diff); c:=f[2]; f:=f[1];
if nops(f)<>1 then lprint(df); error "need only one diff function factor." fi;

```

```

f:=f[]; c:='*(c[]); f:=getvarpow(f); f:=[c,f]; return f; fi;
if op(0,df)=diff then f:=getvarpow(df); f:=[1,f]; return f; fi;
lprint(df); error "unexpected type."; end:

printHBF:=proc(PL::list) local j,DD,f,C,tmp,gcdC; C:=map2(op,1,PL);
gcdC:=1; if nops(C)>1 then tmp:=[seq(cat(_Z,i),i=1..nops(C))];
gcdC:=tmp *~ C; gcdC:='+'(gcdC[]); gcdC:=factor(gcdC);
tmp:=selectremove(has,gcdC,tmp); gcdC:=tmp[2];
if gcdC=0 then gcdC:=1 fi; gcdC:=gcdC*content(tmp[1]); fi;
if gcdC<>1 then C:=C /~ gcdC; fi; DD:=map2(op,2,PL);
f:=op(0,DD[1][1][1]);
DD:=map(z->product(D[z[i][1]]^z[i][2],i=2..nops(z)),DD);
DD:=zip('*',C,DD); DD:='+'(DD[]); gcdC * ``(DD) * cat(f,' ',f);
end:

## print Hirota Bilinear Transform
printHBT:=proc(uf,u,f,i,j,PL,alpha:=1) local DD,g,C,tmp,pl;
pl:=printHBF(PL); if j>0 then print(u=2*alpha*'diff'(ln(f),x$j));
else print(u=2*alpha*ln(f)); fi;
if i>0 then print('diff'(pl/f^2,x$i)=0); else
print(pl/f^2=0); fi; NULL; end;

guessdifforder:=proc(PL::list,x::name)
local L,minorder,maxorder,tmp; L:=map2(op,2,PL);
L:=map(z->z[2..-1],L); tmp:=map(z->map2(op,2,z),L);
tmp:=map(z->'+'(z[]),tmp); tmp:=selectremove(type,tmp,even);
minorder:=0; if nops(tmp[1])<nops(tmp[2]) then minorder:=1 fi;
tmp:=map(z->select(has,z,{x}),L); tmp:=map(z->map2(op,2,z),tmp); if
has(tmp,{[]}) then maxorder:=0; else tmp:=map(op,tmp);
maxorder:=min(tmp[]); fi;
if type(maxorder-minorder,odd) then maxorder:=maxorder-1 fi;
[minorder,maxorder]; end;

guessalpha:=proc(Res,uf,u,f,i,j,PL,alpha) local tmp,res,pl,flag,k;
flag:=1; tmp:=[op(Res)]; tmp:=map(numer,tmp);
tmp:=gcd(tmp[1],tmp[-1]); if type(tmp,'*') then
tmp:=remove(has,tmp,f); fi; if tmp<>0 and has(tmp,{alpha}) then
tmp:=solve(tmp/alpha^difforder(uf),{alpha});
if tmp<>NULL and has(tmp,{alpha}) then lprint(tmp);
for k to nops([tmp]) while flag=1 do
res:=collect(expand(subs(tmp[k],Res)),f,factor);
if res=0 then pl:=subs(tmp[k],PL);
printHBT(uf,u,f,i,j,pl,rhs(tmp[k]));
flag:=0; fi; od; fi; fi; PL; end;

Bilinear:=proc(uf,u,f,x,alpha) local su,h,i,j,g1,CB,PL,gdo,DD,Res;
if hasfun(uf,int) then error "Do not support integral function yet.

```

```

Please substitute int function." fi; for j from 0 to 2 do
  Res:=1; su:=u=2*alpha*diff(ln(f),[x$j]);
  h:=collect(expand(dsubs(su,uf)),f,factor);
  if hasfun(h,ln) then next; fi;
  g1:=getFnumer(h,f)/2; g1:=expand(g1); CB:=HBF(g1);
  gdo:=guessdifforder(CB,x);
  for i from gdo[1] by 2 to gdo[2] do
    if i=0 then PL:=CB; else PL:=HBF(int(g1,x$i)); fi;
    DD:=add(PL[i][1]*BD(PL[i][2][]),i=1..nops(PL));
    Res:=collect(expand(diff(DD/f^2,[x$i])-h),f,factor);
    if Res=0 then printHBT(uf,u,f,i,j,PL,alpha); break;
    elif type(alpha,name) and has(DD,alpha) then
      Res:=guessalpha(Res,uf,u,f,i,j,PL,alpha);
    fi; od; if Res=0 then break; fi; od; PL; end:

```

8. Conclusions

To sum up, an algorithm for generating the Hirota bilinear form of NPDE with logarithm transformation has been proposed in this paper, and the bilinear forms of a class of NPDEs are obtained by using package Bilinear. Then we illuminate the availability of the algorithm by illustrating some examples.

Acknowledgement

The work has been partially supported by the Natural Science Foundation of China (No. 10971109), K.C.Wong Magna Fund in Ningbo University and Student Research of Ningbo University. Jingsong He is also supported by Program for NCET under Grant No.NCET-08-0515.

References

- [1] R. Hirota, The direct method in soliton theory (Cambridge University Press, Cambridge, 2004).
- [2] A. Scott, Encyclopedia of Nonlinear Science, Taylor and Francis, Routledge, New York, 2005.
- [3] Wenxiu Ma, Ruguang Zhou, Liang Gao, Exact one-periodic and two-periodic wave solutions to Hirota bilinear equations in (2+1) dimensions, Mod. Phys. Lett. A 21 (2009) 1677-1688.
- [4] Wenxiu Ma, Engui Fan, Linear superposition principle applying to Hirota bilinear equations, Comput. Math. Appl. 61 (2011) 950-959.
- [5] J. Hietarinta, Introduction to the Hirota bilinear method. Integrability of nonlinear systems (Pondicherry, 1996) 95-103, Lecture Notes in Physics, 495 (Springer, Berlin, 1997).
- [6] J. Hietarinta, A search for bilinear equations passing Hirota's three-soliton condition. I. KdV-type bilinear equations. J. Math. Phys. 28 (1987) 1732-1742.
- [7] J. Hietarinta, A search for bilinear equations passing Hirota's three-soliton condition. II. mKdV-type bilinear equations. J. Math. Phys. 28 (1987) 2094-2101.
- [8] J. Hietarinta, A search for bilinear equations passing Hirota's three-soliton condition. III. Sine-Gordon-type bilinear equations. J. Math. Phys. 28 (1987) 2586-2592.
- [9] J. Hietarinta, A search for bilinear equations passing Hirota's three-soliton condition. IV. Complex bilinear equations. J. Math. Phys. 29 (1988) 628-635.
- [10] W. Hereman and W. Zhuang, Symbolic computation of solitons with Macsyma. Computational and applied mathematics, II (Dublin, 1991), 287-296 (North-Holland, Amsterdam, 1992).
- [11] Zhenjiang Zhou, Jingzhi Fu and Zhibin Li, An implementation for the algorithm of Hirota bilinear form of PDE in the Maple system. Appl. Math. Comput. 183 (2006) 872-877.

- [12] Zhenjiang Zhou, Jingzhi Fu and Zhibin Li, Maple packages for computing Hirota's bilinear equation and multisiton solutions of nonlinear evolution equations. *Appl. Math. Comput.* (2010) 92-104.
- [13] Xudong Yang and Hangyu Ruan, A Maple package on symbolic computation of Hirota bilinear form for nonlinear equations. *Commun. Theor. Phys. (Beijing, China)* 52 (2009) 801-807.
- [14] J. Weiss, M. Tabor and G. Carnevale, The Painleve property for partial differential equations. *J. Math. Phys.* 24 (1983) 522-526.
- [15] Mingliang Wang, Yubin Zhou and Zhibin Li, Application of a homogeneous balance method to exact solutions of nonlinear equations in mathematical physics. *Phys. Lett. A* 216 (1996) 67-75.
- [16] G. B. Whitham, *Linear and Nonlinear Waves* (New York, Wiley, 1974) 9.
- [17] D. Zwillinger, *Handbook of Differential Equations*, 3rd ed. (Boston, MA: Academic Press, 1997) 129-130.
- [18] K. Sawada and T. Kotera, A method for finding N -soliton solutions of the K.d.V. equation and K.d.V.-like equation. *Progr. Theoret. Phys.* 51 (1974) 1355-1367.
- [19] P. J. Caudrey, R. K. Dodd and J. D. Gibbon, A new hierarchy of Korteweg-de Vries equations. *Proc. Roy. Soc. London Ser. A* 351 (1976) 407-422.
- [20] B. B. Kadomtsev and V. I. Petviashvili, On the stability of solitary waves in weakly dispersive media. *Sov. Phys. Dokl.* 15 (1970) 539-541.
- [21] R. Hirota and J. Satsuma, N -soliton solutions of model equations for shallow water waves. *J. Phys. Soc. Japan.* 40 (1976) 611-612.
- [22] M. Ito, An extension of nonlinear evolution equations of the K-dV (mK-dV) type to higher orders. *J. Phys. Soc. Japan.* 49 (1980) 771-778.
- [23] WenHua Huang, YuLu Liu and JieFang Zhang, Doubly periodic propagating wave for (2+1)-dimensional breaking soliton equation. *Commun. Theor. Phys. (Beijing)* 49 (2008) 268-274.
- [24] J. M. Dye and A. Parker, A bidirectional Kaup-Kupershmidt equation and directionally dependent solitons. *J. Math. Phys.* 43 (2002) 4921-4949.
- [25] Jingsong He and Xiaodong Li, Solutions of the (2 + 1)-dimensional KP, SK and KK equations generated by gauge transformations from nonzero seeds. *J. Nonlinear Math. Phys.* 16 (2009) 179-194.