Defective incidence coloring of graphs **

Huimin Bi Xin Zhang[‡]

School of Mathematics and Statistics, Xidian University, Xi'an, 710071, China

February 9, 2022

Abstract

We define the *d*-defective incidence chromatic number of a graph, generalizing the notion of incidence chromatic number, and determine it for some classes of graphs including trees, complete bipartite graphs, complete graphs, and outerplanar graphs. Fast algorithms for constructing the optimal *d*-defective incidence colorings of those graphs are presented.

Keywords: incidence coloring; defective coloring; Latin square; outerplanar graph; polynomial-time algorithm.

I Introduction

The incidence coloring of graphs, introduced by Brualdi and Quinn Massey [7] in 1993, has been attracting attention of many researchers (see an online survey of Sopena [15] for the recent progresses of the study of the incidence coloring). This coloring has many applications in theoretical computer science and information science as it can model the multi-frequency assignment problem where each transceiver can be simultaneously in sending and receiving modes [6].

Formally, let *G* be a graph with vertex set V(G) and edge set E(G). An *incidence* of *G* is a vertex-edge pair (v, e) such that the vertex *v* is incident with the edge *e*. For a vertex $u \in V(G)$ and its neighbor *v* in *G* (say $v \in N_G(u)$), the incidence (u, uv) is a *strong incidence* of *u*, and the incidence (v, uv) is a *weak incidence* of *u*. We use I_u and A_u to denote the set of strong incidences and weak incidences of *u*, respectively.

Brualdi and Quinn Massey [7] defined two *adjacent incidences* as (u, e) and (w, f) such that u = w or $uw \in \{e, f\}$ (it may happen that e = f). They also defined the following.

Definition 1. A proper incidence k-coloring of G is a mapping φ from the set I(G) of all incidences of G to the set $[k] := \{1, 2, ..., k\}$ of integers in such a way that two adjacent incidences receive different colors.

^{*}Supported by the National Natural Science Foundation of China (No. 11871055).

[†]Mathematics Subject Classification (2010): 05C15, 68R10

[‡]Corresponding author. Email: xzhang@xidian.edu.cn.

We think about the proper incidence coloring in another view of point, by giving an equivalent definition as follows.

Definition 2. A proper incidence k-coloring of G is a mapping $\varphi : I(G) \longrightarrow [k]$ such that the following conditions hold for every $u \in V(G)$:

(a2) $\varphi(u, uv) \neq \varphi(u, uw)$ for any two distinct vertices $v, w \in N_G(u)$;

(b2) every color in $\varphi(I_u) := \{\varphi(u, uv) \mid v \in N_G(u)\}$ does not appear among A_u .

We similarly define $\varphi(A_u) := \{\varphi(v, uv) \mid v \in N_G(u)\}$ and $\varphi(I_u \cup A_u) := \varphi(I_u) \cup \varphi(A_u)$, which will be frequently used throughout this paper.

The *incidence chromatic number* of *G*, denoted by $\chi_i(G)$, is the minimum integer *k* such that *G* has a proper incidence *k*-coloring. From either Definition 1 or Definition 2, one can easily see that $\chi_i(G) \ge \Delta(G) + 1$ for every graph *G*. The best known upper bound for $\chi_i(G)$ is $\Delta(G) + 20 \log \Delta(G) + 84$, due to Guiduli [10], who disproved a conjecture of Brualdi and Quinn Massey [7] that $\chi_i(G) \le \Delta(G) + 2$ for every graph *G*. This upper bound is also asymptotically sharp [1, 10].

There are many interesting variations of incidence coloring of graphs, including incidence list coloring [5], incidence game coloring [2, 3], interval incidence coloring [11], fractional incidence coloring [17], and oriented incidence coloring [9].

Motivated by Definition 2, we introduce the defective incidence coloring of graphs in this paper.

Definition 3. A *d*-defective incidence *k*-coloring of *G* is a mapping $\varphi : I(G) \longrightarrow [k]$ such that the following conditions hold for every $u \in V(G)$:

- (a3) $\varphi(u, uv) \neq \varphi(u, uw)$ for any two distinct vertices $v, w \in N_G(u)$;
- (b3) $\varphi(u, uv) \neq \varphi(v, uv)$ for every $v \in N_G(u)$;
- (c3) every color in $\varphi(I_u)$ appears at most d times among A_u .

The minimum number of colors used among all *d*-defective incidence colorings of a graph *G*, denoted by $\chi_i^d(G)$, is the *d*-defective incidence chromatic number.

Comparing Definition 2 with Definition 3, one can easily see that the 0-defective incidence coloring is coincide with the proper incidence coloring, and thus $\chi_i^0(G) = \chi_i(G)$. Moreover,

$$\chi_i^0(G) \ge \chi_i^1(G) \ge \chi_i^2(G) \ge \dots \ge \Delta(G).$$
(1.1)

This motivates us to define the *incidence defectivity* $def_i(G)$ of a graph G. Formally, we let

$$def_i(G) = \min\{k \mid \chi_i^k(G) = \Delta(G)\}.$$

This paper is organized as follows.

In Section 2, we show that the *d*-defective incidence chromatic number of a tree or a complete bipartite graph is its maximum degree whenever $d \ge 1$. Moreover, we construct linear-time algorithm to compute a *d*-defective incidence Δ -coloring for every such graph with maximum degree Δ . In Section 3, we establish

an interesting relationship between the 1-defective incidence colorings of the complete graph K_n and the $n \times n$ Latin squares, and then prove that K_n has a *d*-defective incidence (n - 1)-coloring for every integer $d \ge 1$ whenever $n \ne 2, 4$. Thereafter, based on our new results on Latin squares, a quadratic-time algorithm is designed for constructing a *d*-defective incidence (n - 1)-coloring of K_n with $n \ne 2, 4$. In Section 4, we move our attention to the class of outerplanar graphs, a special graph class with bounded treewidth. We show that every outerplanar graph with maximum degree $\Delta \ge 4$ admits a 1-defective incidence Δ -coloring, and this bound for Δ is sharp. Furthermore, we prove that every outerplanar graph with maximum degree Δ admits a *d*-defective incidence Δ -coloring whenever $d \ge 2$, unless the graph is isomorphic to disjoint copies of K_1 and K_2 . All colorings mentioned above can be constructed in polynomial time according to the proofs in Section 4.

2 Trees and complete bipartite graphs

It is easy to observe that every path and every cycle has 1-defective incidence chromatic number exactly 2, as we can color its incidences alternately by two colors following a fixed direction. Hence we have the following.

Theorem 2.1. If $n \ge 3$ is an integer, then $\chi_i^d(P_n) = 2$ and $\chi_i^d(C_n) = 2$ for every integer $d \ge 1$.

A *rooted tree* is a connected acyclic graph with a special vertex that is called the *root* of the tree and every edge directly or indirectly originates from the root. In a rooted tree, every vertex v, except the root, has exactly one *parent vertex u*, which is the first vertex traversed on the path from v to the root. The vertex v is called a *child* of u. We use Par(v) to denote the parent vertex of v in a rooted tree. An *ordered tree* is a rooted tree in which an ordering is specified for the children of each vertex [4]. The *depth* of a vertex in an ordered tree is the length of the (unique) path from the root to the vertex. Note that the root has depth 0.

Theorem 2.2. If T is a tree, then $\chi_i^d(T) = \Delta(T)$ for every integer $d \ge 1$.

Proof. It is sufficient to show that $\chi_i^1(T) = \Delta(T)$. Assume that *T* has already been embdded as an ordered tree whose root *r* has the maximum degree Δ . We construct an incidence Δ -coloring φ of *T* as follows.

First of all, for each child u_i of r, let $\varphi(r, ru_i) = i - 1 \pmod{\Delta}$ and $\varphi(u_i, ru_i) = i \pmod{\Delta}$, where u_i denotes the *i*-th child of r, And then for each vertex u at depth $\ell \ge 1$, let $\varphi(u, uw_i) = \varphi(u, uv) + i \pmod{\Delta}$ and $\varphi(w_i, uw_i) = \varphi(v, uv) + i \pmod{\Delta}$, where w_i is the *i*-th child of u and v is the parent of u.

We show that this incidence coloring of *T* is 1-defective. Clearly, (*a*3) and (*b*3) by the construction of the coloring, and one can see that every two incidences of A_u for any $u \in V(T)$ are colored distinctly. Hence (*c*3) with d = 1 holds naturally.

Below we release a linear-time algorithm to construct a 1-defective incidence $\Delta(T)$ -coloring of *T* based on the proof of Theorem 2.2. Figure 1 shows an instance of Algorithm 1 on how to construct a 1-defective incidence 3-coloring of a given tree with maximum degree three.



Fig. 1: Constructing a 1-defective incidence 3-coloring using breadth-first search

Algorithm 1: 1-defective incidence coloring of T

Input: A tree *T* with maximum degree Δ .

Output: A 1-defective incidence coloring of T using Δ colors.

- 1 Find a vertex r of maximum degree in T
- 2 Root *T* at *r* using breadth-first search so that *r* has a labelling f(r) = (0) and each vertex *u* at depth $\ell \ge 1$ has a labelling with an $(\ell + 1)$ -array $(x_0, \ldots, x_{\ell-1}, x_\ell)$ such that $(x_0, \ldots, x_{\ell-1})$ is the labelling of the parent of *u* and *u* is the $(x_\ell + 1)$ -th child of its parent

/* p(u) denotes below the last entry of the labelling of u. */ 3 for $\ell = 0$ to Dep(T) do

```
if \ell = 0 then
 4
              for each child u of r do
 5
                    \varphi(r, ru) \leftarrow p(u) \pmod{\Delta}
 6
                    \varphi(u, ru) \leftarrow p(u) + 1 \pmod{\Delta}
 7
         else
8
              for each vertex u at depth \ell do
 9
                    v \leftarrow \operatorname{Par}(u)
10
                    for each child w of u do
11
                          \varphi(u, uw) \leftarrow \varphi(u, uv) + p(u) + 1 \pmod{\Delta}
12
                          \varphi(w, uw) \leftarrow \varphi(v, uv) + p(u) + 1 \pmod{\Delta}
13
```

Theorem 2.3. $\chi^d(K_{m,n}) = \max\{m, n\} = \Delta(K_{m,n})$ for every integer $d \ge 1$.

Proof. Let $\{u_1, u_2, \ldots, u_m\}$ and $\{v_1, v_2, \ldots, v_n\}$ be the bipartition of $K_{m,n}$. Assume, without loss of generality, that $m \ge n$. We construct an incidence *m*-coloring *m* of $K_{m,n}$ by coloring $(v_i, u_j v_i)$ with $i + j - 1 \pmod{m}$, and $(u_j, u_j v_i)$ with $i + j \pmod{m}$ for each $i \in [n]$ and $j \in [m]$. Now we prove that this incidence coloring is 1-defective.

Clearly, (a3) and (b3) hold by the construction of φ .

For each vertex v_i with $i \in [n]$ and for each color $\ell \in \varphi(I_{v_i})$, if $\ell \in \varphi(A_{v_i})$, then by the construction of φ , there is a vertex u_k with $k \in [m]$ such that $\varphi(u_k, u_k v_i) = \ell = i + k \pmod{m}$. Since $k \in [m]$, $k = \ell - i \pmod{m}$. This implies that the vertex u_k is uniquely determined. Hence the color ℓ appears exactly once among A_{v_i} .

On the other hand, for each vertex u_j with $j \in [m]$ and each color $\ell \in \varphi(I_{u_j})$, there is a vertex v_i with $i \in [n]$ such that $\varphi(u_j, u_j v_i) = \ell = i + j \pmod{m}$ by the construction of φ . If $\ell \in \varphi(A_{u_j})$, then by (b3), there is a vertex v_k with $k \in [n] \setminus \{i\}$ such that $\varphi(v_k, u_j v_k) = \ell = k + j - 1 \pmod{m}$. This implies $k - 1 = i \pmod{m}$. Since $1 \le k \le n \le m$, $k = i + 1 \pmod{m}$. Hence the vertex u_k is uniquely determined if it exists (note that it may happen that i = m - 1 and n < m, and thus there is no solution for k). Hence the color ℓ appears at most once among A_{u_i} .

Therefore, (c3) holds in each case, and thus $m \ge \chi^d(K_{m,n}) \ge \Delta(K_{m,n}) = m$ for every integer $d \ge 1$. \Box

Note that the proof of Theorem 2.3 naturally yields a linear-time algorithm to construct a 1-defective incidence coloring of $K_{m,n}$ using $\Delta(K_{m,n})$ colors.

We conclude the following from Theorems 2.1, 2.2, and 2.3.

Corollary 2.4. def_{*i*}(G) = 1 *if* G *is a path, a cycle, a tree, or a bipartite graph.*

Algorithm 2:	1-defective	incidence	coloring	of $K_{m,n}$
--------------	-------------	-----------	----------	--------------

Input: Integers *m* and *n* with $m \ge n$; Output: A 1-defective incidence coloring of $K_{m,n}$ using colors *m*. /* { $u_1, u_2, \ldots u_m$ } and { $v_1, v_2, \ldots v_n$ } are the bipartition of $K_{m,n}$. 1 for i = 1 to *n* do 2 for j = 1 to *m* do 3 $\varphi(v_i, u_j v_i) \leftarrow i + j - 1 \pmod{m}$ 4 $\varphi(u_j, u_j v_i) \leftarrow i + j \pmod{m}$

3 Complete graphs and Latin squares

Lemma 3.1. φ is a 1-defective incidence (n - 1)-coloring of K_n if and only if every two strong incidences and every two weak incidences of any vertex are colored differently under φ , and (u, uv) and (v, uv) receive distinct colors for each pair of vertices u and v. **Proof.** The sufficiency is obvious since (*a*3), (*b*3), and (*c*3) with d = 1 hold trivially. For the necessary, we know that every two strong incidences of any vertex are colored differently under φ by (*a*3), and (*u*, *uv*) and (*v*, *uv*) receive distinct colors for each pair of vertices *u* and *v* by (*b*3). For each vertex *v*, since $\varphi(I_v) = [\Delta]$, there is no color appearing twice on A_v by (*c*3) with d = 1. Since $|A_v| = n - 1$ and φ has exactly n - 1 colors, every two weak incidences of *v* are colored differently under φ .

A *Latin square* is an $n \times n$ square matrix whose entries consist of n symbols such that each symbol appears exactly once in each row and each column. An *intercalate* in a Latin square L is a 2 × 2 Latin subsquare, that is, two rows and two columns whose intersection includes only two symbols. A *principal intercalate* in a Latin square L is an intercalate obtained by striking out from L the same rows as columns.

Lemma 3.2. K_n has a 1-defective incidence (n - 1)-coloring if and only if there is an $n \times n$ Latin square without principal intercalates whose entries at the main diagonal are the same.

Proof. Let *L* be an $n \times n$ Latin square without principal intercalates whose entries on the main diagonal are the same. Permute (if necessary) the symbols 0, 1, ..., n - 1 of *L* so that the entries at the main diagonal are 0. Clearly, the resulting Latin square \tilde{L} still has no principal intercalates.

Let v_1, v_2, \ldots, v_n be vertices of K_n . We color the incidence $(v_i, v_i v_j)$ with the entry $\widetilde{L}(i, j)$ of the Latin square that appears at row *i* and column *j*, and denote this coloring by φ . Now for each vertex $v_i, \varphi(I_{v_i}) = {\widetilde{L}(i, j) | j \in [n] \setminus \{i\}\}} = [n - 1]$ and $\varphi(A_{v_i}) = {\widetilde{L}(j, i) | j \in [n] \setminus \{i\}\}} = [n - 1]$, since \widetilde{L} is a Latin square with $\widetilde{L}(i, i) = 0$ for each $i \in [n]$. This implies that every two strong incidences and every two weak incidences of v_i are colored differently under φ . For each pair of vertices v_i and $v_j, \varphi(v_i, v_i v_j) = \widetilde{L}(i, j)$ and $\varphi(v_j, v_i v_j) = \widetilde{L}(j, i)$. Since $\widetilde{L}(i, i) = \widetilde{L}(j, j) = 0$ and \widetilde{L} has no principal intercalates, $\widetilde{L}(i, j) \neq \widetilde{L}(j, i)$, and thus $\varphi(v_i, v_i v_j) \neq \varphi(v_j, v_i v_j)$. By Lemma 3.1, φ is a 1-defective incidence (n - 1)-coloring of K_n .

In the other direction, if φ is a 1-defective incidence *n*-coloring of K_n , then we construct an $n \times n$ Latin square *L* whose entry L(i, j) at row *i* and column *j* with $i \neq j$ is $\varphi(v_i, v_i v_j)$, and entries at the main diagonal are all *n*. Since for each pair of *i* and *j*, $L(i, j) = \varphi(v_i, v_i v_j) \neq \varphi(v_j, v_i v_j) = L(j, i)$ and L(i, i) = L(j, j) = 0, *L* has no principal intercalates.

Lemma 3.3. [12–14] There exist $n \times n$ Latin squares with no intercalates if and only if $n \neq 2, 4$,

Lemma 3.4. A 4 × 4 Latin square with same entries at the main diagonal has no intercalate if and only if *it has no principal intercalate.*

Proof. The necessary is trivial so we prove the sufficiency. Suppose for a contradiction *L* has an intercalate appearing at the intersection of rows *i*, *j* (*i* < *j*) and columns *m*, *n* (*m* < *n*). Assume that L(i, m) = L(j, n) = a and L(i, n) = L(j, m) = b. Since this intercalate is not principal, $L(1, 1) \notin \{a, b\}$. It follows that $i \notin \{m, n\}$, $j \notin \{m, n\}$ and thus $\{i, j, m, n\} = [4]$. Therefore, L(i, j) = L(j, i) and thus there is a principal intercalate appearing at the intersection of rows *i*, *j* and columns *i*, *j*, a contradiction.

Theorem 3.5.

$$\chi^{1}(K_{n}) = \begin{cases} n-1 & \text{if } n \neq 2, 4, \\ n & \text{otherwise.} \end{cases}$$

Proof. First, it is trivial $\chi^1(K_2) = 2$. If $n \neq 2, 4$, then there exists an $n \times n$ Latin square *L* with no intercalates by Lemma 3.3. Permute (if necessary) the rows of *L* so that the entries at the main diagonal are the same. One can easily observe that the resulting Latin square \tilde{L} does not has any intercalate either, and thus has no principal intercalates. Hence K_n has a 1-defective incidence (n - 1)-coloring by Lemma 3.2. It follows that $\Delta(K_n) \leq \chi^1(K_n) \leq n - 1 = \Delta(K_n)$.

If K_4 has a 1-defective incidence 3-coloring, then by Lemma 3.2, there is a 4×4 Latin square *L* without principal intercalates whose entries at the main diagonal are the same, contradicting Lemma 3.4. Hence $\chi^1(K_4) \ge 4$. On the other hand, one can easily construct a 1-defective incidence 4-coloring of K_4 , implying $\chi^1(K_4) \le 4$.

To construct a 1-defective incidence coloring of K_n (n = 3 or $n \ge 5$) with n - 1 colors, we need by Lemma 3.2 to generate an $n \times n$ Latin square without principal intercalates whose entries at the main diagonal are the same. Lemma 3.3 guarantees the existence of such an Latin square.

In the following, we discuss the recursive construction of such Latin squares and design a quadratictime algorithm to generate an $n \times n$ Latin square satisfying those properties for every $n \neq 2, 4$.

An $n \times n$ matrix $M = (m_{ij})$ is a *circulant* if it has the form $m_{ij} = a_{j-i}$ for some $a_0, a_1, \ldots, a_{n-1}$, where the subscript j - i is taken modulo n. In this paper we denote such a matrix by $(a_0, a_1, \ldots, a_{n-1})_{\text{circ}}$.

Lemma 3.6. Every $n \times n$ circulant matrix with n being odd is a Latin square without principal intercalates.

Proof. Let $M = (a_0, a_1, \dots, a_{n-1})_{\text{circ}}$ where *n* is odd. Clearly, $m_{ii} = a_0$ for each $1 \le i \le n$. It is sufficient to check that $a_{ij} \ne a_{ji}$ if i > j. Since *n* is odd, $j - i \ne n + i - j$. This implies $m_{ij} = a_{j-i} \ne a_{n+i-j} = m_{ji}$.

Lemma 3.7. For a positive even integer n that is not a power of 2, there exists an odd $m \ge 3$ such that the *quotient of n divided by m is a positive power of 2*

Proof. Let *t* be the largest integer such that $2^t | n$ and let $n = 2^t \cdot m$. If *m* is even then $2^{t+1} | n$, contradicting the choice of *t*. If m = 1, then $n = 2^t$, contradicting the choice of *n*. Hence $m \ge 3$ is an odd.

Let A be an $n \times n$ matrix and let

$$A\nabla A = \begin{bmatrix} A & A + nJ \\ B & A \end{bmatrix},$$

where

$$B = (A + nJ)^T \begin{bmatrix} 0 & E_{n-1} \\ 1 & 0 \end{bmatrix}$$

and J is an $n \times n$ matrix in which each element is 1.

We use $A^{(0\nabla)}$ and $A^{(1\nabla)}$ to denote A and $A\nabla A$, respectively. By $A^{(t\nabla)}$ with integer $t \ge 2$, we denote $A^{((t-1)\nabla)}\nabla A^{((t-1)\nabla)}$. Clearly, $A^{(t\nabla)}$ is a $2^t n \times 2^t n$ matrix.

Lemma 3.8. If A is a Latin square without principal intercalates, then so does $A^{(t\nabla)}$ for each $t \ge 1$.

Proof. It is sufficient to show that $A\nabla A$ is a Latin square without principal intercalates.

Let $A = (a_{ij})_{n \times n}$ and $A \nabla A = (m_{ij})_{2n \times 2n}$. According to the construction of $A \nabla A$,

$$m_{ij} = \begin{cases} a_{ij} & \text{if } 1 \le i \le n \text{ and } 1 \le j \le n; \\ a_{i,j-n} + n & \text{if } 1 \le i \le n \text{ and } n+1 \le j \le 2n; \\ a_{n,i-n} + n & \text{if } n+1 \le i \le 2n \text{ and } j=1; \\ a_{j-1,i-n} + n & \text{if } n+1 \le i \le 2n \text{ and } 2 \le j \le n; \\ a_{i-n,j-n} & \text{if } n+1 \le i \le 2n \text{ and } n+1 \le j \le 2n. \end{cases}$$

To complete the proof, we show that $m_{ij} \neq m_{ji}$ for each pair of *i* and *j* with $i \neq j$.

If $1 \le i \le n$ and $1 \le j \le n$, then $m_{ij} = a_{ij}$ and $m_{ji} = a_{ji}$. Since *A* is a Latin square without principal intercalates, $a_{ij} \ne a_{ji}$. It follows that $m_{ij} \ne m_{ji}$.

If $1 \le i \le n$ and $n + 1 \le j \le 2n$, then $m_{ij} = a_{i,j-n} + n$, $m_{ji} = a_{i-1,j-n} + n$ if $i \ne 1$, and $m_{ji} = a_{n,j-n} + n$ if i = 1. Since $a_{i,j-n}$ and $a_{i-1,j-n}$ (resp. $a_{1,j-n}$ and $a_{n,j-n}$) are in the same column and not in the same row of A, they are not equal and thus $m_{ij} \ne m_{ji}$.

If $n + 1 \le i \le 2n$ and $n + 1 \le j \le 2n$, then $m_{ij} = a_{i-n,j-n} \ne a_{j-n,i-n} = m_{ji}$, since A is a Latin square without principal intercalates.

Hence in each case we have $m_{ij} \neq m_{ji}$, as desired.

The following Algorithm 3 outputs in $O(n^2)$ time an $n \times n$ Latin square *L* without principal intercalates whose entries at the main diagonal are 0 whenever we input an integer $n \neq 2, 4$. It works by Lemmas 3.6, 3.7, and 3.8. Afterwards, by Lemma 3.2, we can construct a 1-defective incidence coloring of K_n using colors from $\{1, 2, ..., n\}$ according to Algorithm 4, which also runs in $O(n^2)$ time.

To close this section, we present the following.

Theorem 3.9. $\chi^d(K_n) = n - 1$ for every integer $d \ge 2$ and $n \ne 2$.

Proof. Theorem 3.5 and (1.1) imply the result for $n \neq 1, 4$. The case of n = 1 is trivial and a 2-defective 3-coloring of K_4 can be easily constructed. Hence $\chi^d(K_4) = 3$ for every integer $d \ge 2$.

Combining Theorems 3.5 with 3.9, we conclude the following.

Corollary 3.10.

$$def_{i}(K_{n}) = \begin{cases} 1 & if \ n \neq 2, 4, \\ 2 & if \ n = 4, \\ \infty & if \ n = 2. \end{cases}$$

Note that $\chi^d(K_2) = 2$ for every integer $d \ge 1$.

Algorithm 3: LATIN-SQARE(n)

Input: An integer $n \neq 2, 4$; **Output:** An $n \times n$ Latin square *L* without principal intercalates whose entries at the main diagonal are 0.

```
1 if n is odd then
       L \leftarrow (0, 1, \ldots, n-1)_{\text{circ}}
2
3 else
       if n = 2^t for some integer t \ge 3 then
4
                  [0 1 2 3 4 5 6 7]
                   2 0 7 4 5 3
                                          1 6
                  A \leftarrow
5
                   6 7 3 5 1 4
                                          2
                                              0
           L \leftarrow A^{((t-3)\nabla)}
6
       else
7
           Find the largest integer t such that 2^t | n and let m = n/2^t
8
           A \leftarrow (0, 1, \ldots, m-1)_{\text{circ}}
9
            L \leftarrow A^{(t\nabla)}
10
```

Algorithm 4: COLOR-COMPLETE-GRAPH(n)

Input: An integer $n \neq 2, 4$; Output: A 1-defective incidence coloring of K_n using colors from $\{1, 2, ..., n\}$. /* The set of vertices of K_n is $\{v_1, v_2, ..., v_n\}$. 1 $L \leftarrow \text{LATIN-SQARE}(n)$ 2 for i = 1 : n do 3 for j = 1 : n and $j \neq i$ do 4 Color the incidence $(v_i, v_i v_j)$ with L(i, j)

*/

4 Outerplanar graphs

Definition 4. A conditional incidence Δ -coloring of G is an incidence Δ -coloring such that

- (a4) (u, uv) and (v, uv) receive distinct colors for each edge uv;
- (b4) each color appears at most once among A_u for each vertex u;
- (c4) each color appears at most once among I_u for each vertex u;
- (d4) each color appears at least once among $A_u \cup I_u$ for each vertex u with $\deg_G(u) \ge \Delta 1$.

Observation 1. Any conditional incidence Δ -coloring of G is a 1-defective incidence coloring.

An *outplanar graph* is a graph that can be embedded in the plane in such a way that all vertices lie on the outer face. In this section we show the following theorem.

Theorem 4.1. If G is an outerplanar graph with $\Delta(G) \leq \Delta$ and $\Delta \geq 4$, then G has a conditional incidence Δ -coloring.

Combining Observation 1 and Theorem 4.1, we immediately obtain the following.

Theorem 4.2. $\chi^1(G) = \Delta(G)$ if G is an outerplanar graph with $\Delta(G) \ge 4$.

Remark 1. The bound 4 in Theorems 4.1 and 4.2 are both sharp. To see this, we look at the outerplanar graph H derived from a cycle uxvy of length four via adding an edge uv and a pendant edge xz. It has maximum degree 3. Suppose that H has a 1-defective incidence 3-coloring φ . Assume by symmetry that $\varphi(u, ux) = 1$, $\varphi(u, uv) = 2$ and $\varphi(u, uy) = 3$. It follows that $\varphi(v, uv) \neq 2$. If $\varphi(v, uv) = 1$, then $\varphi(x, xv) = 1$, $\varphi(y, uy) = \varphi(v, vy) = 2$, and $\varphi(x, ux) = \varphi(v, vx) = \varphi(y, vy) = 3$, which forces $\varphi(x, xz) = \varphi(z, xz) = 2$. If $\varphi(v, uv) = 3$, then $\varphi(y, uy) = \varphi(v, vy) = \varphi(x, vx) = 1$, $\varphi(v, xv) = \varphi(x, ux) = 2$, and $\varphi(y, vy) = 3$, forcing $\varphi(x, xz) = \varphi(z, xz) = 3$. Each case contradicts (b3) of Definition 3. Hence there exist outerplanar graphs with maximum degree 3 that are not 1-defectively incidence 3-colorable, and thus not conditionally incidence 3-colorable.

A graph *H* is a *minor* of a graph *G* if a copy of *H* can be obtained from *G* via repeated edge deletion and/or edge contraction. Chartrand and Harary [8] first pointed out that a graph is outerplanar if and only if it is $\{K_4, K_{2,3}\}$ -minor-free. The following classic structural theorem on outerplanar graphs were applied in quite a lot of papers.

Lemma 4.3. [16] Every outerplanar graph G contains one of the following configurations

- (C1) a vertex of degree at most 1;
- (C2) an edge uv with $\deg_G(u) = \deg_G(v) = 2$;
- (C3) a triangle uvw with $\deg_G(u) = 2$ and $\deg_G(v) = 3$;
- (C4) two intersecting triangle uvx and uwy such that $\deg_G(u) = 4$ and $\deg_G(v) = \deg(w) = 2$.

In the following, we show a series of self-contained lemmas that not only support the proof of Theorem 4.1 but also can be applied in other places.

A family \mathcal{G} of graphs is *hereditary* (resp. *minor-closed*) if every subgraph (resp. minor) H of each graph $G \in \mathcal{G}$ belongs to \mathcal{G} . A graph G is ρ - Δ -*critical* (resp ρ - Δ -*critical*) if $\Delta(G) \leq \Delta$ and G has no conditional incidence Δ -coloring but every subgraph (resp. minor) H of G does. Since every subgraph is a minor, we have the following.

Lemma 4.4. A ρ - Δ -critical graph is definitely ρ - Δ -critical.

Lemma 4.5. If G is a ρ - Δ -critical graph, then G is connected.

Proof. If *G* has two components G_1 and G_2 , then G_1 and G_2 are conditionally incidence Δ -colorable. Combining the conditional incidence Δ -colorings of them, we obtain a conditional incidence Δ -coloring of *G*, contradicting the fact that *G* is ρ - Δ -critical.

Lemma 4.6. If G is a ρ - Δ -critical graph, then $\delta(G) \ge 2$.

Proof. Suppose for a contradiction that *G* has a vertex *v* of degree at most 1. By Lemma 4.5, we may assume $N_G(v) = \{u\}$. Since *G* is ρ - Δ -critical, G' = G - v admits a conditional incidence Δ -coloring φ' .

We extend φ' to a conditional incidence Δ -coloring φ of G as follows.

If $\varphi'(I_u \cup A_u) = [\Delta]$, then $\varphi'(I_u) \setminus \varphi'(A_u) \neq \emptyset$ and $\varphi'(A_u) \setminus \varphi'(I_u) \neq \emptyset$, since $\deg_{G'}(u) \leq \Delta(G) - 1 \leq \Delta - 1$. Hence we can color (u, uv) with $a \in \varphi'(A_u) \setminus \varphi'(I_u)$ and (v, uv) with $b \in \varphi'(I_u) \setminus \varphi'(A_u)$ to complete φ .

If $|\varphi'(I_u \cup A_u)| < \Delta$, then $\deg_{G'}(u) \le \Delta - 2$. If $\deg_{G'}(u) < \Delta - 2$, then color (u, uv) with $a \in [\Delta] \setminus \varphi'(I_u)$. and (v, uv) with $b \in [\Delta] \setminus (\varphi'(A_u) \cup \{a\})$. If $\deg_{G'}(u) = \Delta - 2$, then $[\Delta] \setminus \varphi'(I_u) = \{a, b\}$ has one element, say b, that does not appear in $\varphi'(A_u)$. Hence we can color (u, uv) with a and (v, uv) with b to complete φ . Note that this extension guarantees $\varphi(I_u \cup A_u) = [\Delta]$.

Remark 2. If G is an outerplanar graph with $\Delta(G) \leq \Delta$ such that G has no 1-defective incidence Δ coloring but every subgraph H of G does, then the idea of proving Lemma 4.6 cannot be applied to prove $\delta(G) \geq 2$. Actually, if G has an edge uv with $\deg_G(v) = 1$ and $\deg_G(u) = \Delta$, then G - v has a 1-defective incidence Δ -coloring φ' by the minimality of G. However, it may happen that $\varphi'(I_u) = \varphi'(A_u) = [\Delta - 1]$ and thus we have to color both (u, uv) and (v, uv) with Δ while extending φ' to G and then return a failure. This is indeed the reason why we introduce the notion of conditional incidence coloring and then prove Theorem 4.1 instead of proving Theorem 4.2 directly.

Lemma 4.7. If G is a ρ - Δ -critical graph, then G does not contain an edge uv with $\deg_G(u) = 2$ and $\deg_G(v) \le \Delta - 2$.

Proof. Suppose for a contradiction that *G* has such an edge *uv*. By Lemma 4.6, we have $2 \le \deg_G(v) \le \Delta - 2$ and thus $\Delta \ge 4$. Let *w* be the other neighbor of *u* besides *v*. Since *G* is a ρ - Δ -critical, G' = G - uv admits a conditional incidence Δ -coloring φ' .

Assume $\varphi(w, uw) = a$ and $\varphi(u, uw) = b$. Since $\deg_{G'}(v) \le (\Delta - 2) - 1 \le \Delta - 3$, we are able to color (v, uv) with a color $a' \in [\Delta] \setminus (\varphi'(I_v) \cup \{a\})$ and (u, uv) with a color from $[\Delta] \setminus (\varphi'(A_v) \cup \{a', b\})$. This extends φ' to a conditional incidence Δ -coloring of G.

Lemma 4.8. If G is a ρ - Δ -critical graph and $\Delta \ge 4$, then G does not contain a vertex with $\deg_G(u) = 2$ such that $N_G(u) = \{v, w\}$ and $\deg_G(v) = \deg_G(w) = \Delta - 1$.

Proof. Suppose for a contradiction that *G* contains such a vertex. Since *G* is ρ - Δ -critical, G' = G - uw has a conditional incidence Δ -coloring φ' . Assume $\varphi'(v, uv) = a$, $\varphi'(u, uv) = b$, $[\Delta] \setminus \varphi'(I_w) = \{a', c'\}$, and $[\Delta] \setminus \varphi'(A_w) = \{b', d'\}$. We extend φ' to an incidence Δ -coloring φ as follows.

Case 1. $\{a', c'\} \cap \{b', d'\} = \emptyset$

Since $\deg_{G'}(w) = \Delta -2$, $\varphi'(I_w \cup A_w) = [\Delta]$ under this case. So we color (w, uw) with a color in $\{a', c'\} \setminus \{a\}$ and (u, uw) with a color in $\{b', d'\} \setminus \{b\}$ to complete φ . It is easy to see that φ is a conditional incidence Δ -coloring of *G*.

Case 2. $a \in \{a', c'\} \cap \{b', d'\}$.

Assume, without loss of generality, that a' = b' = a. Coloring (u, uw) with a and (w, uw) with c', we obtain a conditional incidence Δ -coloring φ of G as $\varphi(I_w \cup A_w) \supseteq \varphi'(I_w) \cup \{a, c'\} = \varphi'(I_w) \cup \{a', c'\} = [\Delta]$, $\varphi(u, uw) = a \neq b = \varphi(u, uv)$, and $\varphi(w, uw) = c' \neq a' = a = \varphi(v, uv)$.

Case 3. $a \notin \{a', c'\} \cap \{b', d'\} \neq \emptyset$.

Assume, without loss of generality, that $a' = b' \neq a$.

Subcase 3.1. $b \neq d'$.

We color (w, uw) with a' and (u, uw) with d'. This extended coloring φ of G satisfies $\varphi(I_w \cup A_w) \supseteq \varphi'(A_w) \cup \{a', d'\} = \varphi'(A_w) \cup \{b', d'\} = [\Delta], \varphi(w, uw) = a' \neq a = \varphi(v, uv), \text{ and } \varphi(u, uw) = d' \neq b = \varphi(u, uv),$ and thus is a conditional incidence Δ -coloring of G.

Subcase 3.2. $a \neq c'$.

We color (w, uw) with c' and (u, uw) with b'. This extended coloring φ of G satisfies $\varphi(I_w \cup A_w) \supseteq \varphi'(A_w) \cup \{b', c'\} = \varphi'(A_w) \cup \{a', c'\} = [\Delta], \varphi(w, uw) = c' \neq a = \varphi(v, uv), \text{ and } \varphi(u, uw) = b' \neq b = \varphi(u, uv).$ It follows that φ is a conditional incidence Δ -coloring of G.

Subcase 3.3. b = d' and a = c'.

This is equivalent to say $[\Delta] \setminus \varphi'(I_w) = \{a, a'\}$ and $[\Delta] \setminus \varphi'(A_w) = \{b, a'\}$.

Erase the colors of (u, uv) and (v, uv). Now we have two ways to transfer φ' to a conditional incidence Δ -coloring of G'' = G - uv. The first way is to color (w, uw) with a' and (u, uw) with b, while the second way is to color (w, uw) with a and (u, uw) with a'. We denote by φ_1 and φ_2 be those two colorings respectively.

Assume $[\Delta] \setminus \varphi_1(I_v) = \{a, c''\}$ and $[\Delta] \setminus \varphi_1(A_v) = \{b, d''\}$. Using the same proof strategies in Cases 1 and 2 and Subcases 3.1 and 3.2, one can find that the only obstacle while extending φ_1 to a conditional incidence Δ -coloring of *G* is the case that $[\Delta] \setminus \varphi_1(I_v) = \{a', e\}$ and $[\Delta] \setminus \varphi_1(A_v) = \{b, e\}$ for some $e \in [\Delta]$. In particular, since $a \neq a'$, we deduce e = d'' = a and a' = c''.

Therefore, $[\Delta] \setminus \varphi_2(I_v) = [\Delta] \setminus \varphi_1(I_v) = \{a', a\}$ and $[\Delta] \setminus \varphi_2(A_v) = [\Delta] \setminus \varphi_1(A_v) = \{a, b\}$ if φ_1 is not extendable. However, we earn chance to extend φ_2 now. This can be done by coloring (u, uv) with a and (v, uv) with a'. Since the resulting coloring φ satisfies $\varphi(I_v \cup A_v) \supseteq \varphi_2(I_v) \cup \{a, a'\} = [\Delta], \varphi(u, uv) = a \neq a' = \varphi_2(u, uw)$, and $\varphi(v, uv) = a' \neq a = \varphi_2(w, uw), \varphi$ is a conditional incidence Δ -coloring of G. \Box



Fig. 2: The configuration T_1 , where xy is a non-edge and we may have $\{x_1, x_2\} \cap \{y_1, y_2\} \neq \emptyset$, and the operation $T_1 - \{v, w\} + xy$.

Lemma 4.9. If *G* is a ρ - Δ -critical graph and $\Delta \ge 4$, then *G* does not contain a triangle uvw with $\deg_G(u) = 2$, $\deg_G(v) = 3$ and $\deg_G(w) = 4$.

Proof. Suppose for a contradiction that such a triangle exists in *G*. By Lemma 4.7, we have $\Delta = 4$. Since *G* is ρ -4-critical, G - uv has a conditional incidence 4-coloring φ' . Let $N_G(v) = \{u, w, x\}$ and $N_G(w) = \{u, v, y, z\}$. Assume the colors on (v, vx), (x, vx), (v, vw), (w, vw), (w, uw) and (u, uw) are 1, *a*, 2, *b*, *c*, *d*, $(a \neq b, a \neq 1, b \neq 2, b \neq c, c \neq d$ and $d \neq 2$) respectively.

If $|\{1, 2\} \cup \{a, b\}| = 2$, then a = 2 and b = 1. If $d \notin \{1, 2\}$, say, d = 3, then $c \neq 3$ and thus we can color (u, uv) with 4 and (v, uv) with 3. If $d \in \{1, 2\}$, then d = 1 and color (v, uv) with $a' \in \{3, 4\} \setminus \{c\}$ and (u, uv) with $b' \in \{3, 4\} \setminus \{a'\}$.

If $|\{1, 2\} \cup \{a, b\}| = 3$, then assume, without loss of generality, that $4 \notin \{1, 2\} \cup \{a, b\}$.

If a = 3, then b = 1. If c = 4, then $d \neq 4$, and we color (u, uv) with 4 and (v, uv) with color 3. If $c \neq 4$, then we are able to color (u, uv) with 2 as $d \neq 2$ and (v, uv) with color 4.

If b = 3, then a = 2. If c = 4, then $d \neq 4$, and we color (u, uv) with 4 and (v, uv) with color 3. If $c \neq 4$ and $d \neq 1$, then we color (u, uv) with 1 and (v, uv) with color 4. If $c \neq 4$ and d = 1, then c = 2 and we can color (u, uv) with 4 and (v, uv) with 3.

If $|\{1, 2\} \cup \{a, b\}| = 4$, then assume by symmetry that a = 3 and b = 4. Since $d \neq 2$, we are able to color (u, uv) with 2 and (v, uv) with a color in $\{3, 4\} \setminus \{c\}$.

In each of the above cases, the resulting extended coloring is a conditional incidence 4-coloring φ of *G* as the property that $\varphi(I_v \cup A_v) = [4]$ is satisfied throughout the extension.

In the following, we use T_1 to stand for the configuration as shown by the left picture of Figure 2.

Lemma 4.10. If G is a ρ - Δ -critical graph and $\Delta \geq 4$, then G does not contain the configuration T_1 .

Proof. Suppose to the contrary that *G* contains a copy of T_1 . By Lemmas 4.7 and 4.8, we have $\Delta = 4$. Since *G* is ρ -4-critical and $G' = G - \{v, w\} + xy$ is a minor of *G*, *G'* has a conditional incidence 4-coloring φ' . The right picture of Figure 2 shows a partial coloring of φ' .

The idea of the proof is to restrict φ' to a partial coloring of $G - \{u, v, w\}$ and then extend it to G by coloring the remaining 12 incidences using colors t_1, t_2, \ldots, t_{12} as shown by the first picture of Figure 2. In order to make the extended coloring being a conditional incidence 4-coloring of G, we need choose each t_i carefully from [4] such that

$$\Omega = \prod_{i=1}^{12} \prod_{\lambda \in \Lambda_i} (t_i - \lambda) \neq 0,$$

where $\Lambda_1 = \{t_2, t_5, t_{10}, c, d\}, \Lambda_2 = \{t_3, t_6, t_9, t_{12}\}, \Lambda_3 = \{t_4, t_7, t_9, t_{12}\}, \Lambda_4 = \{t_8, t_{11}, g, h\}, \Lambda_5 = \{t_6, t_9, a, b\}, \Lambda_6 = \{t_7, t_{10}, t_{11}\}, \Lambda_7 = \{t_8, t_{10}, t_{11}\}, \Lambda_8 = \{t_{12}, e, f\}, \Lambda_9 = \{t_{10}, t_{12}, a, b\}, \Lambda_{10} = \{t_{11}, c, d\}, \Lambda_{11} = \{t_{12}, g, h\},$ and $\lambda_{12} = \{e, f\}$. If such t_i 's exist, then we say that (a, b, c, d, e, f, g, h) is extendable.

We traverse all cases of (a, b, c, d, e, f, g, h) and check whether all of them are extendable by computer assistance. Algorithm 5 defined a function CHECK(a,b,c,d,e,f,g,h). This function returns a non-zero vector if and only if we input integers $a, b, c, d, e, f, g, h \in [4]$ such that (a, b, c, d, e, f, g, h) is extendable.

Assume a = 1 and b = 2. We run Algorithm 6 on a usual personal computer using MATLAB. It returns a zero matrix M in less than one minute. Since (a, b, c, d, e, f, g, h) comes from the conditional incidence 4-coloring φ' , it naturally holds that $c \notin \{1, d\}, d \neq 2, g \notin \{e, h\}$, and $f \notin \{e, h\}$. So lines 1–10 of Algorithm 6 returns a matrix such that if (a, b, c, d, e, f, g, h) is not extendable then it is coincide with some row of this matrix. Since Algorithm 6 finally returns a zero matrix, we conclude that if (a, b, c, d, e, f, g, h) is not extendable then either

- $\{1,2\} = \{e,f\}, |\{c,d\} \cap \{g,h\}| = 1 \text{ and } (\{c,d\} \cap \{g,h\}) \cap \{1,2\} = \emptyset, \text{ or }$
- $\{1,2\} = \{e,f\}, |\{c,d\} \cap \{g,h\}| = 1, (\{c,d\} \cap \{g,h\}) \cap \{1,2\} \neq \emptyset \text{ and } (\{c,d\} \oplus \{g,h\}) \cap \{1,2\} = \emptyset, \text{ or } \{g,h\} \cap \{g,h$
- $\{c,d\} = \{g,h\}, |\{1,2\} \cap \{e,f\}| = 1 \text{ and } (\{1,2\} \cap \{e,f\}) \cap \{c,d\} = \emptyset, \text{ or }$
- $\{c,d\} = \{g,h\}, |\{1,2\} \cap \{e,f\}| = 1, (\{1,2\} \cap \{e,f\}) \cap \{c,d\} \neq \emptyset \text{ and } (\{1,2\} \oplus \{e,f\}) \cap \{c,d\} = \emptyset.$

Here \oplus is the operation of symmetric difference.

We look back at the conditional incidence 4-coloring φ' .

If the first case occurs, then assume, without loss of generality, that e = 1, f = 2, and c = g = 3. It follows $3 \notin \{s_5, s_6\}$ and thus $s_5 = s_6 = 4$, a contradiction.

If the second case occurs, then assume, without loss of generality, that e = d = h = 1, f = 2, c = 3, and g = 4. It follows $s_1 = s_2 = 2$, a contradiction.

Similarly, we still have contradiction if we meet the third or the fourth case.

Hence every valid (a, b, c, d, e, f, g, h) is extendable.

We denote by T_2 the configuration described by the picture of Figure 3.

Algorithm 5: CHECK(*a*,*b*,*c*,*d*,*e*,*f*,*g*,*h*)





Fig. 3: The configuration T_2 , where we may have $x_1 = y_1$.

Algorithm 6: REDUCIBILITY-INSPECTION()

Input: Null Output: An inspection matrix M $1 i \leftarrow 1$ **2** for c = 1 to 4 do for d = 1 to 4 do 3 for e = 1 to 4 do 4 **for** f = 1 to 4 **do** 5 for g = 1 to 4 do 6 for h = 1 to 4 do 7 **if** $c \notin \{1, d\}, d \neq 2, g \notin \{e, h\}, f \notin \{e, h\}, and$ 8 CHECK(1, 2, c, d, e, f, g, h) = [0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0] then M(i,:) = [1, 2, c, d, e, f, g, h]9 i = i + 110 11 $s \leftarrow$ the number of rows of M **12** for j = 1 to s do $c \leftarrow M(j,3)$ 13 $d \leftarrow M(j, 4)$ 14 $e \leftarrow M(j, 5)$ 15 $f \leftarrow M(j, 6)$ 16 $q \leftarrow M(j,7)$ 17 $h \leftarrow M(j, 8)$ 18 if $\{1, 2\} = \{e, f\}$ and $|\{c, d\} \cap \{g, h\}| = 1$ then 19 **if** $(\{c, d\} \cap \{g, h\}) \cap \{1, 2\} = \emptyset$ or $(\{c, d\} \oplus \{g, h\}) \cap \{1, 2\} = \emptyset$ **then** 20 $M(j,:) \leftarrow [0,0,0,0,0,0,0,0]$ 21 if $\{c, d\} = \{g, h\}$ and $|\{1, 2\} \cap \{e, f\}| = 1$ then 22 **if** $(\{1,2\} \cap \{e,f\}) \cap \{c,d\} = \emptyset$ or $(\{1,2\} \oplus \{e,f\}) \cap \{c,d\} = \emptyset$ **then** 23 $M(j,:) \leftarrow [0,0,0,0,0,0,0,0]$ 24



Fig. 4: The configuration T_3 , where max{deg_G(x), deg_G(y)} ≤ 5 .

Lemma 4.11. If G is a ρ - Δ -critical graph and $\Delta \geq 4$, then G does not contain the configuration T_2 .

Proof. Suppose for a contradiction that *G* contains a copy of T_2 . By Lemmas 4.7 and 4.8, we have $\Delta = 4$. Since *G* is ρ -4-critical, $G - \{u, v, w\}$ has a conditional incidence 4-coloring φ' . Assume $\varphi'(x, xx_1) = 1$, $\varphi'(x_1, xx_1) = 2$, $\varphi'(y, yy_1) = c$, and $\varphi(y_1, yy_1) = d$. We are to extend φ' to *G* by coloring the remaining 14 incidences as marked in the picture with colors t_1, t_2, \ldots, t_{14} . Below we distinguish six non-isomorphic cases and show what those t_i 's are. Let $T = [t_1, t_2, \ldots, t_{14}]$.

If c = 1 and d = 2, then choose T = [3, 1, 4, 1, 4, 2, 3, 2, 2, 1, 4, 3, 3, 4]. If c = 1 and d = 3, then choose T = [3, 1, 4, 1, 4, 3, 2, 3, 3, 1, 4, 2, 2, 4]. If c = 2 and d = 1, then choose T = [3, 4, 3, 4, 4, 3, 4, 3, 2, 1, 2, 1, 3, 4]. If c = 2 and d = 3, then choose T = [1, 2, 1, 4, 4, 3, 2, 1, 3, 4, 1, 4, 2, 3]. If c = 3 and d = 2, then choose T = [1, 3, 1, 4, 2, 4, 2, 1, 4, 3, 1, 2, 3, 4]. If c = 3 and d = 4, then choose T = [3, 4, 2, 1, 4, 2, 4, 2, 1, 4, 3, 1, 2, 3, 4]. One can check that the resulting coloring in each case is a conditional incidence Δ -coloring of G.

One can check that the resulting coloring in each case is a conditional incluence Δ -coloring of O.

We denote by T_3 the configuration described by the picture of Figure 4.

Lemma 4.12. If G is a ρ - Δ -critical graph and $\Delta \geq 5$, then G does not contain the configuration T_3 .

Proof. Suppose for a contradiction that *G* contains a copy of T_3 . By Lemma 4.7, we have $\Delta = 5$. Since *G* is a ρ -5-critical graph, G - uv has a conditional incidence 5-coloring φ' . Figure 4 shows a partial coloring of φ' .

If $\{c, d, e\} = \{1, 2, 3\}$, then we can color (u, uv) with $a' \in \{4, 5\} \setminus \{a\}$ and (v, uv) with $b' \in \{4, 5\} \setminus \{a'\}$ to complete a conditional incidence 5-coloring of *G* provided $b \notin \{4, 5\}$. Hence $b \in \{4, 5\}$. Assume by symmetry that b = 4. It follows that $a \neq 4$ and thus we are able to color (u, uv) with 4 and (v, uv) with 5 to finish a conditional incidence 5-coloring of *G*.

If $|\{1, 2, 3\} \cup \{c, d, e\}| = 4$, then assume by symmetry that $4 \in \{c, d, e\}$ and $5 \notin \{c, d, e\}$. If a = 5, then $b \neq 5$ and we can color (u, uv) with 4 and (v, uv) with 5 and obtain a conditional incidence 5-coloring of *G*. Hence $a \neq 5$. If we are able to color (u, uv) with $a' \in \{4, 5\} \setminus \{a\}$ and (v, uv) with $b' \in [5] \setminus \{b, c, d, e, a'\}$ such that $5 \in \{a', b'\}$, then we obtain a conditional incidence 5-coloring φ of *G* as $\varphi(I_u \cup A_u) = [5]$. If this

is impossible, then a = 4 and $\{b, c, d, e\} = [4]$. It follows that $b, c \notin \{4, 5\}$. Recolor (u, ux) with 5 and color (u, uv) with 1 and (v, uv) with 5. This completes a conditional incidence 5-coloring of *G*.

If $|\{1, 2, 3\} \cup \{c, d, e\}| = 5$, then color (u, uv) with $a' \in \{4, 5\} \setminus \{a\}$ and (v, uv) with $b' \in [5] \setminus \{b, c, d, e, a'\}$. This is possible as if $\{b, c, d, e, a'\} = [5]$ then $a' \notin \{c, d, e\}$, which contradicts the assumption that $\{4, 5\} \subseteq \{c, d, e\}$.

The proof of Theorem 4.1. Suppose for a contradiction that *G* is a minimal outerplanar graph in terms of |V(G)| + |E(G)| with $\Delta(G) \le \Delta$ such that *G* is not conditional incidence Δ -colorable. It follows that *G* is a ρ - Δ -critical graph. As an outerplanar graph, *G* contains one of the four configurations among (C1), (C2), (C3), and (C4) by Lemma 4.3. However, *G* does not contain (C1) by Lemmas 4.4 and 4.6, (C2) by Lemmas 4.4 and 4.7, (C3) by Lemmas 4.4, 4.7, 4.8 and 4.9, or (C4) by Lemmas 4.4, 4.6, 4.9, 4.10, 4.11 and 4.12. This contradiction completes the proof.

To close this section, we prove the following.

Theorem 4.13. $\chi^d(G) = \Delta(G)$ for every integer $d \ge 2$ if G is an outerplanar graph with $\Delta(G) \ge 2$.

Proof. It is sufficient to prove $\chi^2(G) \le 3$ for every subcubic outerplanar graph by (1.1) and by Theorems 2.1 and 4.2. Suppose for a contradiction that *G* is minimal counterexample in terms of |V(G)| + |E(G)| to this result. Clearly, *G* is connected, so by Lemma 4.3, *G* contains a vertex *u* of degree at most 2.

If $\deg_G(u) = 1$, then assume $N_G(u) = \{v\}$. By the minimality of G, G - u has a 2-defective incidence 3coloring φ' . We extend φ' to a 2-defective incidence 3-coloring of G by coloring (v, uv) with $a \in [3] \setminus \varphi'(I_v)$ and (u, uv) with $b \in [3] \setminus (S \cup \{a\})$, where S is the set of colors used at most twice among A_v under φ' . This is possible as $|\varphi'(I_v)| = \deg_G(v) - 1 \le 2$ and $|S| \le 1$.

If $\deg_G(u) = 2$, then assume $N_G(u) = \{v, w\}$. By the minimality of G, G - uv has a 2-defective incidence 3-coloring φ' . Since $|\varphi'(I_v)| = \deg_G(v) - 1 \le 2$, we are able to color (v, uv) with $a \in [3] \setminus \varphi'(I_v)$. If $a \in \varphi'(A_v)$, then color (u, uv) with $b \in [3] \setminus \{\varphi'(u, uw), a\}$ to complete a 2-defective incidence 3-coloring of G. If $a \notin \varphi'(A_v)$, then $|\varphi'(A_v)| = 2$ and therefore we are still able to color (u, uv) with $b \in [3] \setminus \{\varphi'(u, uw), a\}$ to complete the desired coloring.

Remark 1 implies that the condition of $d \ge 2$ in Theorem 4.13 is necessary. Combining Theorems 2.1, 4.2, and 4.13 together, we deduce the following.

Corollary 4.14.

$$def_i(G) = \begin{cases} 1 & if \Delta \neq 1, 3, \\ 2 & if \Delta = 3, \\ \infty & if \Delta = 1. \end{cases}$$

if G is an outerplanar graph with maximum degree Δ *.*

Remark 3. Since the proofs in this section are all constructive, they yield a polynomial-time algorithm for outputting a d-defective incidence Δ -coloring whenever we input an outplanar graph with maximum degree Δ and an integer d such that $\Delta \ge 4$ and $d \ge 1$, or $\Delta \in \{2, 3\}$ and $d \ge 2$.

We leave an open problem to close this paper.

Problem 4.15. Does every bridgeless subcubic outerplanar graph has a 1-defective incidence 3-coloring?

It would be interesting to point out that not every bridgeless subcubic graph is 1-defectively incidence 3-colorable. In the literature, a *snark* is a simple, connected, bridgeless cubic graph with chromatic index equal to 4. There are many well-known snarks including the Petersen graph, which is the smallest snark. We claim that $\chi^1(S) \ge 4$ for every snark *S*.

Suppose for a contradiction that *S* admits a 1-defective incidence 3-coloring φ . For every edge *uv*, let $c(uv) := \varphi(u, uv) + \varphi(v, uv) \pmod{3}$. For every two edges *ux* and *uy* incident with *u*, if c(ux) = c(uy), then $\{\varphi(u, ux), \varphi(x, ux)\} = \{\varphi(u, uy), \varphi(y, uy)\}$, which implies $\varphi(u, ux) = \varphi(y, uy), \varphi(u, uy) = \varphi(x, ux)$, and thus $\varphi(u, uz) = \varphi(z, uz)$ for the third neighbor *z* of *u*, a contradiction. Hence the images under *c* of every two adjacent edges of *S* are distinct and thus the mapping $c : E(S) \longrightarrow \{0, 1, 2\}$ is a proper edge 3-coloring of *S*, contradicting the fact that the chromatic index of *S* is 4.

Note that this argument cannot return a negative answer to Problem 4.15, as every subcubic outerplanar graph has chromatic index 3.

References

- [1] I. Algor and N. Alon. The star arboricity of graphs. *Discrete Mathematics*, 75(1):11–22, 1989.
- [2] S. D. Andres. The incidence game chromatic number. *Discrete Appl. Math.*, 157(9):1980–1987, 2009.
- [3] S. D. Andres. Erratum to: The incidence game chromatic number [Discrete Appl. Math. 157 (9) (2009) 1980–1987] [mr2522464]. *Discrete Appl. Math.*, 158(6):728, 2010.
- [4] E. A. Bender. *Lists, Decisions and Graphs With an Introduction to Probability.* S. Gill Williamson, 2010.
- [5] B. Benmedjdoub, I. Bouchemakh, and E. Sopena. Incidence choosability of graphs. *Discrete Appl. Math.*, 265:40–55, 2019.
- [6] H. Bi and X. Zhang. Incidence coloring of Mycielskians with fast algorithm. *Theoret. Comput. Sci.*, 878(879):1–10, 2021.
- [7] R. A. Brualdi and J. J. Q. Massey. Incidence and strong edge colorings of graphs. *Discrete Math.*, 122(1-3):51–58, 1993.
- [8] G. Chartrand and F. Harary. Planar Permutation Graphs. *Annales de l'I.H.P. Probabilités et statis-tiques*, 3(4):433–438, 1967.

- [9] C. Duffy, G. MacGillivray, P. Ochem, and A. Raspaud. Oriented incidence colourings of digraphs. *Discuss. Math. Graph Theory*, 39(1):191–210, 2019.
- [10] B. Guiduli. On incidence coloring and star arboricity of graphs. *Discrete Math.*, 163(1-3):275–278, 1997.
- [11] R. Janczewski, A. Mał afiejska, and M. Mał afiejski. Interval incidence coloring of bipartite graphs. *Discrete Appl. Math.*, 166:131–140, 2014.
- [12] A. Kotzig, C. Lindner, and A. Rosa. Latin squares with no subsquares of order two and disjoint steiner triple systems. *Utilitas Mathematica*, 7:287–294, 1975.
- [13] A. Kotzig and J. Turgeon. On certain constructions for latin squares with no latin subsquares of order two. *Discrete Mathematics*, 16(3):263–270, 1976.
- [14] M. McLeish. On the existence of latin squares with no subsquares of order two. *Utilitas Mathematica*, 8:41–53, 1975.
- [15] E. Sopena. The incidence coloring page. https://www.labri.fr/perso/sopena. Page last modified on January 28, 2021.
- [16] K. Z. W. Wang. Δ-matchings and edge-face chromatic numbers. Acta Math. Appl. Sin., 22:236–242, 1999.
- [17] D. Yang. Fractional incidence coloring and star arboricity of graphs. *Ars Combin.*, 105:213–224, 2012.