A note on one-dimensional time fractional ODEs

Yuanyuan Feng*¹, Lei Li^{†2}, Jian-Guo Liu^{‡3}, and Xiaoqian Xu ^{§4}

Abstract

In this note, we prove or re-prove several important results regarding one dimensional time fractional ODEs following our previous work [4]. Here we use the definition of Caputo derivative proposed in [8, 10] based on a convolution group. In particular, we establish generalized comparison principles consistent with the new definition of Caputo derivatives. In addition, we establish the full asymptotic behaviors of the solutions for $D_c^{\gamma}u = Au^p$. Lastly, we provide a simplified proof for the strict monotonicity and stability in initial values for the time fractional differential equations with weak assumptions.

1 Introduction

The fractional calculus in time has been used widely in physics and engineering for memory effect, viscoelasticity, porous media etc [5, 7, 2, 1, 9]. There is a huge amount of literature discussing time fractional differential equations. For instance, one can find some results in [3, 2] using the classic Caputo derivatives. In this paper, we study the following time fractional ODE:

$$D_c^{\gamma} u = f(t, u), \ u(0) = u_0,$$
 (1.1)

for $\gamma \in (0,1)$ and f measurable. Here $D_c^{\gamma}u$ is the generalized Caputo derivative introduced in [8, 10]. As we will see later, this generalized definition is theoretically more convenient, since it allows us to take advantage of the underlying group structure.

As in [8], we use the following distributions $\{g_{\beta}\}$ as convolution kernels for $\beta \in (-1,0)$:

$$g_{\beta}(t) = \frac{1}{\Gamma(1+\beta)} D\left(\theta(t)t^{\beta}\right).$$

Here $\theta(t)$ is the standard Heaviside step function, $\Gamma(\cdot)$ is the gamma function, and D means the distributional derivative on \mathbb{R} . Indeed, g_{β} can be defined for $\beta \in \mathbb{R}$ (see [8]) so that $\{g_{\beta} : \beta \in \mathbb{R}\}$ forms a convolution group. In particular, we have

$$g_{\beta_1} * g_{\beta_2} = g_{\beta_1 + \beta_2}. \tag{1.2}$$

Here since the support of g_{β_i} (i=1,2) is bounded from left, the convolution is well-defined. Now we are able to give the generalized definition of fractional derivatives:

^{1,4}Department of Mathematics, Carnegie Mellon University, Pittsburgh, PA 15213, USA

²Department of Mathematics, Duke University, Durham, NC 27708, USA.

³ Departments of Mathematics and Physics, Duke University, Durham, NC 27708, USA.

^{*}yuanyuaf@andrew.cmu.edu

[†]leili@math.duke.edu

[‡]jliu@phy.duke.edu

[§]xxu@math.cmu.edu

Definition 1.1 ([8, 10]). Let $0 < \gamma < 1$. Consider $u \in L^1_{loc}[0,T)$. Given $u_0 \in \mathbb{R}$, we define the γ -th order generalized Caputo derivative of u, associated with initial value u_0 , to be a distribution in $\mathscr{D}'(-\infty,T)$ with support in [0,T), given by

$$D_c^{\gamma} u = g_{-\gamma} * ((u - u_0)\theta(t)).$$

If $\lim_{t\to 0+} \frac{1}{t} \int_0^t |u(s) - u_0| ds = 0$, we call $D_c^{\gamma}u$ the Caputo derivative of u.

As in [8], if the function u is absolutely continuous, the generalized definition reduces to the classical definition. However, the generalized definition is theoretically useful because it reveals the underlying group structure (see Proposition 1.1).

Definition 1.2. Let T>0. A function $u\in L^1_{loc}[0,T)$ is a weak solution to (1.1) on [0,T) with initial value u_0 , if $f(t,u(t))\in \mathscr{D}'(-\infty,T)$ and the equality holds in the distributional sense. We call a weak solution u a strong solution if (i). $\lim_{t\to 0+} \frac{1}{t} \int_0^t |u(s)-u_0| ds=0$; (ii). both $D_c^{\gamma}u$ and f(t,u(t)) are locally integrable on [0,T).

By the group property (1.2), we have

Proposition 1.1 ([8]). Suppose $f \in L^{\infty}_{loc}([0,\infty) \times \mathbb{R}; \mathbb{R})$. Fix T > 0. Then, $u(t) \in L^{1}_{loc}[0,T)$ with initial value u_0 is a strong solution of (1.1) on (0,T) if and only if $\lim_{t\to 0_+} \frac{1}{t} \int_0^t |u(s) - u_0| ds = 0$ and it solves the following integral equation

$$u(t) = u_0 + \frac{1}{\Gamma(\gamma)} \int_0^t (t - s)^{\gamma - 1} f(s, u(s)) ds, \ \forall t \in (0, T).$$
 (1.3)

Using this integral formulation, the following has been shown in [8]

Proposition 1.2. Suppose $f:[0,\infty)\times(\alpha,\beta)\to\mathbb{R}$ is continuous and locally Lipschitz continuous in u. For any given initial value $u_0\in(\alpha,\beta)$, there is a unique strong solution, which either exists globally on $[0,\infty)$ or approaches the boundary of (α,β) in finite time. Moreover, this solution is continuous on the interval of existence.

Below in Section 2, we will establish some generalized comparison principles consistent with the new definition of Caputo derivatives. In Section 3, we establish the full asymptotic behaviors of the solutions for $D_c^{\gamma}u = Au^p$. In Section 4, we provide a new proof for the strict monotonicity and stability in initial values with weak assumptions.

2 Generalized comparison principles

The comparison principles are important in the analysis of time fractional PDEs (See [11]). There are many versions of comparison principles proved in literature using various definitions of Caputo derivatives. In [8], the authors assumed $f(t,\cdot)$ to be non-decreasing. In [15, Lemma 2.6], $f(t,\cdot)$ was assumed to be non-increasing. In [14, Theorem 2.3], there is no assumption on the monotonicity of $f(t,\cdot)$, but the function v is assumed to be C^1 so that the pointwise value of $D_c^{\gamma}v$ can be defined. Combining these ideas and establishing a crucial lemma (Lemma 2.1), we prove some generalized comparison principles in this section. Similar to [8], we define the inequality in the distributional sense:

Definition 2.1. Let U be an open interval. We say $f \in \mathscr{D}'(U)$ is a nonpositive (nonnegative) distribution if for any $\varphi \in C_c^{\infty}(U)$ with $\varphi \geq 0$, we have $\langle f, \varphi \rangle \leq 0$ ($\langle f, \varphi \rangle \geq 0$). We say $f_1 \leq f_2$ in the distributional sense for $f_1, f_2 \in \mathscr{D}'(U)$, if $f_1 - f_2$ is nonpositive. We say $f_1 \geq f_2$ in the distributional sense if $f_1 - f_2$ is nonnegative.

In order to prove the comparison principle, we first prove the following auxiliary lemma:

Lemma 2.1. Suppose $u \in L^1_{loc}[0,T)$ and $\lim_{t\to 0^+} \frac{1}{t} \int_0^t |u(s) - u_0| ds = 0$. If there exists a function $f \in L^1_{loc}(0,T)$ such that on interval (0,T) we have in the distributional sense that $D_c^{\gamma} u \leq f$, then for any given $A \in \mathbb{R}$, we have in the distributional sense

$$D_c^{\gamma}(u-A)^+ \leq \chi(u \geq A)f$$
, on $(0,T)$.

Proof. First, recall the following result in [8, Proposition 3.11]: if $u \in C[0,T) \cap C^1(0,T)$ and $u \mapsto E(u)$ is C^1 and convex, we have

$$D_c^{\gamma} E(u) \leq E'(u) D_c^{\gamma} u.$$

Now let us consider $\eta \in C_c^{\infty}(-1,0)$ with $\eta \geq 0$ and $\int \eta \, dt = 1$. Define $\eta^{\epsilon}(t) = \frac{1}{\epsilon} \eta(\frac{t}{\epsilon})$ and $u^{\epsilon} = \eta^{\epsilon} * u$. As showed in [8, Proposition 3.11], $u^{\epsilon}(0) \to u_0$ and $u^{\epsilon}(t) \to u(t)$ in $L^1_{\text{loc}}[0,T)$. Denote $E(u) = (u-A)^+$ and define $E^{\delta}(u) = (E*\eta^{\delta})(u)$. Clearly, $(E^{\delta})'(u) = \eta^{\delta} * \chi(u \geq A)$

Denote $E(u) = (u-A)^+$ and define $E^{\delta}(u) = (E*\eta^{\delta})(u)$. Clearly, $(E^{\delta})'(u) = \eta^{\delta}*\chi(u \ge A)$ is nonnegative and increasing, which implies that E^{δ} is a convex increasing function. Then, we have

$$D_c^{\gamma} E^{\delta}(u^{\epsilon}) \le (E^{\delta})'|_{u^{\epsilon}} D_c^{\gamma} u^{\epsilon}. \tag{2.1}$$

It is not hard to see $\limsup_{\epsilon \to 0} (E^{\delta})'|_{u^{\epsilon}} D_c^{\gamma} u^{\epsilon} \leq (E^{\delta})'|_{u} f(t)$. Since $E^{\delta}(u^{\epsilon})$ converges to $E^{\delta}(u)$ in L^{1}_{loc} and $E^{\delta}(u^{\epsilon}(0))$ converges to $E^{\delta}(u_{0})$, according to Definition 1.1, $D_{c}^{\gamma} E^{\delta}(u^{\epsilon}) \to D_{c}^{\gamma} E^{\delta}(u)$ as distributions. Moreover, notice that the inequality is preserved in the distributional sense (Definition 2.1). We have $D_{c}^{\gamma} E^{\delta}(u) \leq (E^{\delta})'|_{u} f(t)$. Taking $\delta \to 0$, similarly we have $D_{c}^{\gamma} E^{\delta}(u)$ converges as distributions to $D_{c}^{\gamma}(u-A)^{+}$. Then the right hand side of (2.1) converges to $\chi(u \geq A) f(t)$, and the inequality is preserved in the distributional sense. \square

As is well-known, if $u \in H^1(0,T)$, $D(u-A)^+ = \chi(u-A)Du$. Since Caputo derivative is nonlocal, the equality is no longer true in general. However, we have similar inequalities and Lemma 2.1 provides an answer.

Corollary 2.1. Suppose u(t) is a locally integrable function with $\lim_{t\to 0^+} \frac{1}{t} \int_0^t |u(s)-u_0| ds = 0$. Let $A \in \mathbb{R}$ and $t_1 \in (0,T)$ is a Lebesgue point. If $u \leq A$ for a.e. $t \leq t_1$, and on the interval (t_1,T) we have $D_c^{\gamma}u \leq 0$ in the distributional sense, then we have $u \leq A$, a.e. (0,T).

Let u^{ϵ} be the mollification in the proof of Lemma 2.1. Consider $v^{\epsilon} = u^{\epsilon} - \frac{C(\epsilon)\theta(t)}{\Gamma(1+\gamma)}t^{\gamma}$ such that $v^{\epsilon} \leq A$ for $t \in [0, t_1 + \epsilon]$. $C(\epsilon) \to 0$ since t_1 is a Lebesgue point. Applying Lemma 2.1, $D_c^{\gamma}(v^{\epsilon} - A)^+ \leq \chi(t \geq t_1 + \epsilon)(D_c^{\gamma}u^{\epsilon} - C(\epsilon)) \leq \chi(t \geq t_1 + \epsilon)(D_c^{\gamma}u^{\epsilon} - \eta_{\epsilon} * D_c^{\gamma}u)$. Taking $\epsilon \to 0$ yields $D_c^{\gamma}(u - A)^+ \leq 0$. The details are left to readers. Now several versions of comparison principles can be stated as follows:

Theorem 2.1. (i) Suppose $u_i \in L^1_{loc}[0,T)$ with $\lim_{t\to 0^+} \frac{1}{t} \int_0^t |u_i(s) - u_{i,0}| \, ds = 0$ (i = 1,2). Suppose $u_1(t) \leq u_2(t)$ on $[0,t_1]$ for a Lebesgue point t_1 , and the γ -th Caputo derivatives of u_1 , u_2 on $[0,t_1]$ are locally integrable. Define

$$h_i(t) = u_{i,0} + \frac{1}{\Gamma(\gamma)} \int_0^{t \wedge t_1} (t-s)^{\gamma-1} D_c^{\gamma} u_i(s) \, ds, \ i = 1, 2.$$

Then, $h_1(t) \leq h_2(t)$ for all $t \in [0,T]$. Moreover, assume there exists a measurable function f(t,u) such that (i) $f(\cdot,u_i(\cdot))$ (i = 1,2) is locally integrable on $[t_1,T)$; (ii) $f(t,\cdot)$ is non-decreasing on $[t_1,T)$; (iii) $D_c^{\gamma}u_1 \leq f(t,u_1)$ and $D_c^{\gamma}u_2 \geq f(t,u_2)$ in the distributional sense on (t_1,T) , then $u_1 \leq u_2$ a.e. on [0,T).

- (ii) Suppose $u_i \in L^1_{loc}[0,T)$ with $\lim_{t\to 0^+} \frac{1}{t} \int_0^t |u_i(s) u_{i,0}| \, ds = 0$ (i=1,2). If $u_1 \leq u_2$ on $[0,t_1]$ for a Lebesgue point t_1 and $D_c^{\gamma}(u_1-u_2) \leq f(t,u_1) f(t,u_2)$ as distributions on (t_1,T) , with $f(t,\cdot)$ being non-increasing on (t_1,T) and $f(\cdot,u_i(\cdot))$ (i=1,2) being locally integrable on $[t_1,T)$, then $u_1 \leq u_2$ a.e on [0,T).
- (iii) Suppose u(t) is a continuous function on [0,T]. If $u(t_1) = \sup_{0 \le s \le t_1} u(s)$ for some $t_1 \in (0,T]$ and $f(t) = D_c^{\gamma} u(t)$ is a continuous function, then $f(t_1) \ge 0$.

Proof. (i). Clearly, $D_c^{\gamma} h_i = D_c^{\gamma} u_i$ for $t \leq t_1$ and $D_c^{\gamma} h_i = 0$ for $t > t_1$. Let $u = h_1 - h_2$, A = 0 in Corollary 2.1, we find $h_1 \leq h_2$. On $[t_1, T)$, we have

$$u_1(t) \le h_1(t) + \frac{1}{\Gamma(\gamma)} \int_{t_1}^t (t-s)^{\gamma-1} f(s, u_1) \, ds, \ u_2(t) \ge h_2(t) + \frac{1}{\Gamma(\gamma)} \int_{t_1}^t (t-s)^{\gamma-1} f(s, u_2) \, ds.$$

As $h_1(t) \le h_2(t)$ and $f(t, \cdot)$ is non-decreasing, one has $u_1(t) \le u_2(t)$ (see [8, Theorem 4.10]).

(ii). Apply Lemma 2.1 for $u_1 - u_2$ and A = 0. (The proof is similar as in Corollary 2.1.)

(iii). Consider $u^{\epsilon}(t) = u(t) + \frac{\epsilon \theta(t)}{\Gamma(1+\gamma)} t^{\gamma}$, where $\epsilon > 0$. Then, t_1 is the unique maximizer of u^{ϵ} on $[0, t_1]$. Let $f^{\epsilon} = D_c^{\gamma} u^{\epsilon} = f + \epsilon$. It suffices to show

$$f^{\epsilon}(t_1) \ge 0, \ \forall \epsilon > 0.$$
 (2.2)

Otherwise, there is an $\epsilon_0 > 0$ such that $f^{\epsilon_0}(t_1) < 0$. Since f^{ϵ_0} is continuous, we can find $\delta > 0$ such that on $[t_1 - \delta, t_1]$ f^{ϵ_0} is negative and $u^{\epsilon_0}(t) \le u^{\epsilon_0}(t_1 - \delta)$ for $t \le t_1 - \delta$. Applying Corollary 2.1, we have $u^{\epsilon_0}(t) \le u^{\epsilon_0}(t_1 - \delta)$ for $t \in [t_1 - \delta, t_1]$, which is a contradiction. Taking $\epsilon \to 0$ then gives the result.

Remark 2.1. Though the conditions here are weaker under the new definition of Caputo derivative, (ii) is essentially [15, Lemma 2.6] and (iii) is well-known for C^1 functions (see, for example [12, 14]).

Now, we establish a generalized Grönwall inequality (or another version of comparison principle), consistent with the new definition of Caputo derivative. The main construction is inspired by [14].

Theorem 2.2. Suppose f(t,u) is continuous and locally Lipschitz in u. Let v(t) be a continuous function. If $D_c^{\gamma}v \leq f(t,v)$ in the distributional sense, and $D_c^{\gamma}u = f(t,u)$, with $v_0 \leq u_0$. Then, $v \leq u$ on the common interval. Similarly, if we have $D_c^{\gamma}v \geq f(t,v)$ as distributions and $v_0 \geq u_0$, then $v \geq u$ on the common interval.

Proof. We only prove the first claim (the proof for the other is similar). By Proposition 1.2, $D_c^{\gamma}u = f(t, u)$ with initial value $u(0) = u_0$ has a unique solution on the interval $[0, T_b)$, where T_b is the largest time of existence. Moreover, u is continuous on $[0, T_b)$.

Fix $T \in (0, T_b)$. Pick M large enough so that u(t) and v(t) fall into $[0, T] \times [-M, M]$. Let L be the Lipschitz constant of $f(t, \cdot)$ for the region $[0, T] \times [-2M, 2M]$. Consider

$$v^{\epsilon} = v - \epsilon w.$$

Here $w=E_{\gamma}(2Lt^{\gamma})$ is the solution to $D_c^{\gamma}w=2Lw$ with initial value 1, where $E_{\gamma}(z)=\sum_{n=0}^{\infty}\frac{z^n}{\Gamma(n\gamma+1)}$ is the Mittag-Leffler function [6, 13]. Clearly, if ϵ is sufficiently small, v^{ϵ} falls into $[0,T]\times[-2M,2M]$. Then, we find that in the distributional sense

$$D_c^{\gamma} v^{\epsilon} = D_c^{\gamma} v - \epsilon 2Lw \le f(t, v) - \epsilon 2Lw \le f(t, v^{\epsilon}) - \epsilon Lw.$$

We claim that for all such small ϵ ,

$$v^{\epsilon}(t) < u(t), \forall t \in [0, T]. \tag{2.3}$$

If not, define

$$t_1 = \sup\{t \in (0,T] : v^{\epsilon}(s) \le u(s), \ \forall s \in [0,t]\}.$$

Since $v^{\epsilon}(0) = v_0 - \epsilon < u_0$, by continuity we have $t_1 > 0$. By assumption, (2.3) is not true, and we have $t_1 < T$. Consequently, there exists $\delta_1 > 0$, such that $v^{\epsilon}(t_1) = u(t_1)$ and $v^{\epsilon}(t) > u(t)$ for $t \in (t_1, t_1 + \delta_1)$. Moreover,

$$D_c^{\gamma}(v^{\epsilon} - u) \le f(t, v^{\epsilon}) - \epsilon Lw - f(t, u).$$

By continuity, for some $\delta_2 \in (0, \delta_1)$, $D_c^{\gamma}(v^{\epsilon} - u)$ is a nonpositive distribution on the interval $(t_1, t_1 + \delta_2)$. By Corollary 2.1, we have $v^{\epsilon}(t) \leq u(t)$ for $t \in (t_1, t_1 + \delta_2)$, which is a contradiction. Hence, (2.3) is true. Taking $\epsilon \to 0$ in (2.3) yields the result on [0, T]. Since T is arbitrary, the result is true.

3 Asymptotic behaviors for a class of fractional ODEs

In this section, we study the solution curves to the following autonomous fractional ODEs:

$$D_c^{\gamma} u = A u^p, \ u(0) = u_0 > 0.$$
 (3.1)

The monotonicity of the solutions to (3.1) and some partial results for the asymptotic behaviors have been established in our previous work [4]. The asymptotic behaviors of the solutions for the A < 0, p > 0 case have also been discussed in [15, Theorem 7.1]. However, the discussion on all the range of A and p is not complete. Here, we will give a complete description on asymptotic behaviors of the solution curves.

By Proposition 1.2, the strong solution u to (3.1) exists on $[0, T_b)$ for $T_b \in (0, \infty]$. If $T_b < \infty$, either $\lim_{t \to T_b^-} u(t) = 0$ or $\lim_{t \to T_b^-} u(t) = \infty$. We give a complete description regarding the solutions curves to (3.1):

Theorem 3.1. Consider (3.1). If A = 0, then $u(t) = u_0$. If A > 0, then all the solutions are strictly increasing on $(0, T_b)$. If A < 0, then all solutions are strictly decreasing before they touch 0.

- (i) Suppose A > 0. If p > 1, then $T_b < \infty$ and $u(t) \sim \left[\frac{\Gamma(\frac{p\gamma}{p-1})}{A\Gamma(\frac{\gamma}{p-1})}\right]^{\frac{1}{p-1}} (T_b t)^{-\frac{\gamma}{p-1}}$, as $t \to T_b^-$. If p = 1, then $u(t) = u_0 E_{\gamma}(At^{\gamma})$. If p < 1, then there exist $c_1 > 0$ and $c_2 > 0$ such that $c_1 t^{\frac{\gamma}{1-p}} \le u(t) \le c_2 t^{\frac{\gamma}{1-p}}$, $t \ge 1$.
- (ii) Suppose A < 0. If p < 0, the solution curve touches u = 0 in finite time where the right hand side blows up. If p = 0, then $u = u_0 + Ag_{1+\gamma}$. If p > 0, then $T_b = \infty$, and there exist $c_1 > 0$, $c_2 > 0$ such that $c_1 t^{-\frac{\gamma}{p}} \le u(t) \le c_2 t^{-\frac{\gamma}{p}}$, $t \ge 1$.

Proof. The A=0 or p=0 cases are trivial. The monotonicity has been proved in [4]. The A>0, p>1 case has also been discussed there. Indeed, there is also an accurate estimate of T_b in [4]. The p=1 case is trivial. The A<0, p>0 case has been discussed in [15, Theorem 7.1]. In fact, they established a version of comparison principle and used a subsolution and a supersolution to get $c_1 t^{-\frac{\gamma}{p}} \leq u(t) \leq c_2 t^{-\frac{\gamma}{p}}, \quad t \geq 1$. For the case A<0, p<0, since the solution is decreasing, we have $D_c^{\gamma} u \leq A u_0^p < 0$ before u touches zero. Hence, the claim follows.

Now, we establish the results for A>0, p<1 case. First, let us construct the subsolution as follows:

$$\omega(t) = \begin{cases} u_0, & t \in [0, t_0], \\ at^{\frac{\gamma}{1-p}}, & t \ge t_0. \end{cases}$$

Here a > 0 is to be determined and t_0 is determined by $at_0^{\frac{\gamma}{1-p}} = u_0$. Clearly, ω is absolutely continuous on any finite interval. For $t < t_0$, $D_c^{\gamma} \omega = 0 \le A \omega^p$. For $t \ge t_0$, we have

$$D_c^{\gamma}\omega = \frac{a\gamma}{(1-p)\Gamma(1-\gamma)} \int_{t_0}^{t} \frac{\tau^{\frac{\gamma}{1-p}-1}}{(t-\tau)^{\gamma}} d\tau < \frac{a\gamma B(\frac{\gamma}{1-p},1-\gamma)}{(1-p)\Gamma(1-\gamma)} t^{\frac{\gamma p}{1-p}} = \frac{a\Gamma(\gamma/(1-p)+1)}{\Gamma(\gamma p/(1-p)+1)} t^{\frac{p\gamma}{1-p}},$$

where $B(\cdot, \cdot)$ is the Beta function. Clearly, if we choose a > 0 such that $\frac{a\Gamma(\gamma/(1-p)+1)}{\Gamma(\gamma p/(1-p)+1)} \le Aa^p$, then $D_c^{\gamma}\omega \le A\omega^p$. Such a exists because p < 1.

For the super-solution, let us consider

$$v(t) = \begin{cases} u_0 + B_1 \frac{t^{\gamma}}{\Gamma(1+\gamma)}, & t \in [0,1], \\ B_2 t^{\frac{\gamma}{1-p}}, & t \ge 1. \end{cases}$$

 B_2 is determined by $B_2 = u_0 + \frac{B_1}{\Gamma(1+\gamma)}$. This choice of B_2 makes v absolutely continuous on any finite interval. We now determine B_1 . On [0,1], one has $D_c^{\gamma}v = B_1$. For t > 1, we have

$$D_c^{\gamma} v = \frac{B_1 \gamma}{B(1+\gamma, 1-\gamma)} \int_0^1 \frac{\tau^{\gamma-1}}{(t-\tau)^{\gamma}} d\tau + \frac{B_2}{\Gamma(1-\gamma)} \frac{\gamma}{1-p} \int_1^t \frac{\tau^{\frac{\gamma}{1-p}-1}}{(t-\tau)^{\gamma}} d\tau.$$

On [1,2], one has $D_c^{\gamma} v > \frac{B_1 \gamma}{B(1+\gamma,1-\gamma)} \int_0^1 \frac{\tau^{\gamma-1}}{(2-\tau)^{\gamma}} d\tau = B_1 C_1(\gamma)$. For t > 2, we have

$$D_c^{\gamma} v > B_2 \frac{1}{\Gamma(1-\gamma)} \frac{\gamma}{1-p} t^{\frac{\gamma_p}{1-p}} \int_{\frac{1}{t}}^1 \frac{\tau^{\frac{\gamma}{1-p}-1}}{(1-\tau)^{\gamma}} d\tau \ge B_2 t^{\frac{\gamma_p}{1-p}} C_2(p,\gamma).$$

It is clear that there exists $M_1(A, p, \gamma)$ such that as long as $B_2 \ge M_1$, $D_c^{\gamma} v \ge A v^p$ for $t \ge 2$ since p < 1. For v to be a super-solution, one needs

$$u_0 + B_1 \frac{1}{\Gamma(1+\gamma)} \ge M_1, \ B_1 \min(1, C_1(\gamma)) \ge A \max\left(u_0^p, \left(u_0 + \frac{B_1}{\Gamma(1+\gamma)}\right)^p 2^{\frac{p\gamma}{1-p}}\right).$$

Such B_1 exists since p < 1. Hence, applying comparison principle Theorem 2.2 yields the result.

4 Strict monotonicity and stability in initial values

It is well-known that solution curves for well-behaved ODEs do not touch each other. However, for fractional ODEs, similar results are not trivial since the dynamics is non-Markovian. By the comparison principles (or generalized Grönwall inequality), if f(t,u) in (1.1) is continuous and locally Lipschitz in u, u(0) < v(0) implies $u(t) \le v(t)$ for $t \ge 0$. However we do not have strict inequality. In [2, Theorem 6.12], the strict inequality has been established following a series of contraction techniques. Using our new definition of Caputo derivative, we provide a new proof of that solutions are strict monotone in initial values, by assuming $f \in L^{\infty}_{loc}$.

The following lemma (a variant of [4, Lemma 3.4] or [16, Theorem 1]), is important:

Lemma 4.1. Let $r_{\lambda}(t) = -\frac{d}{dt}E_{\gamma}(-\lambda\Gamma(\gamma)t^{\gamma})$ be the resolvent for kernel $\lambda t^{\gamma-1}$ (in other words, $r_{\lambda}(t) + \lambda \int_{0}^{t} (t-s)^{\gamma-1} r_{\lambda}(s) ds = \lambda t^{\gamma-1}$). Let T > 0. Assume $h \in L^{1}[0,T]$, h > 0 a.e., satisfying

$$h(t) - \int_0^t r_{\lambda}(t-s)h(s)ds > 0, \ a.e., \quad \forall \lambda > 0.$$

Suppose $v \in L^{\infty}[0,T]$, then the integral equation

$$y(t) + \int_0^t (t-s)^{\gamma-1} v(s)y(s)ds = h(t)$$
 (4.1)

 $has \ a \ unique \ solution \ y(t) \in L^1[0,T]. \ Moreover, \ y(t) > 0, a.e..$

The proof is exactly the same as [4, Lemma 3.4], though we only assume $v \in L^{\infty}[0,T]$ here. Next, we provide a new proof for the strict monotonicity in initial value. We also prove the stability of solutions with respect to initial values.

Theorem 4.1. Assume that $f(\cdot, \cdot) \in L^{\infty}_{loc}([0, \infty) \times \mathbb{R})$. Moreover, assume for every compact set K, there is $L_K > 0$ such that $|f(t, u) - f(t, v)| \leq L_K |u - v|$ for a.e. $(t, u), (t, v) \in K$. Then, for a given initial value u_0 , the solution in $L^{\infty}_{loc}[0, T_b)$ is unique. Further, we have

- Any two solutions $u_i \in L^{\infty}_{loc}[0, T_b^i)$ (i = 1, 2) with initial values $u_{1,0} < u_{2,0}$ satisfy $u_1(t) < u_2(t)$ on $[0, \min(T_b^1, T_b^2))$.
- For any T > 0, M > 0, there exists C(M,T) > 0 such that any two solutions with $||u_i||_{L^{\infty}[0,T]} \le M$ (i = 1,2) and initial values $u_{1,0}, u_{2,0}$ satisfy

$$||u_1 - u_2||_{L^{\infty}[0,T]} \le C(M,T)|u_{1,0} - u_{2,0}|.$$

Proof. Fix $T \in (0, \min(T_b^1, T_b^2))$. There exists K compact such that for a.e $t \in [0, T]$, $(t, u_i(t)) \in K$. By Proposition 1.1, one has

$$u_i(t) = u_{i,0} + \frac{1}{\Gamma(\gamma)} \int_0^t (t-s)^{\gamma-1} f(s, u_i(s)) ds.$$

The boundedness of $f(s, u_i(s))$ implies that $u_i(t) \in C[0, T]$. If $u_{1,0} = u_{2,0}$, by taking the difference, $|u_1(t) - u_2(t)| \le C \int_0^t (t-s)^{\gamma-1} |u_1(s) - u_2(s)| ds$ and the uniqueness therefore follows.

Now, assume $u_{1,0} \neq u_{2,0}$. Define $y(t) = (u_2(t) - u_1(t))/(u_{2,0} - u_{1,0})$, we have

$$y(t) + \int_0^t (t-s)^{\gamma-1} v(s) y(s) ds = 1$$
, where $v(s) = -\frac{1}{\Gamma(\gamma)} \frac{f(s, u_2(s)) - f(s, u_1(s))}{u_2(s) - u_1(s)}$.

If $u_1(s) = u_2(s)$, we define v(s) = 0. Note that $|v| \le L_K/\Gamma(\gamma)$ a.e. for $t \in (0, T)$. By setting h = 1 in Lemma 4.1, one has

$$1 - \int_0^t r_{\lambda}(t - s) \, ds = E_{\gamma}(-\lambda \Gamma(\gamma)t^{\gamma}) > 0.$$

By Lemma 4.1, y(t) > 0. Since y is continuous, satisfying

$$y(t) \le 1 + \int_0^t (t-s)^{\gamma-1} ||v||_{L^{\infty}[0,T]} y(s) ds,$$

we have $y(t) \leq C(\|v\|_{L^{\infty}}, T)$ by [4, Proposition 5]. This verifies the last claim.

Acknowledgements

The work of J.-G Liu was partially supported by KI-Net NSF RNMS11-07444 and NSF DMS-1514826. Y. Feng was supported by NSF DMS-1252912.

References

- [1] M. Allen, L. Caffarelli, and A. Vasseur. A parabolic problem with a fractional time derivative. *Arch. Ration. Mech. An.*, 221(2):603–630, 2016.
- [2] K. Diethelm. The analysis of fractional differential equations: An application-oriented exposition using differential operators of Caputo type. Springer, 2010.
- [3] K. Diethelm and N. J. Ford. Analysis of fractional differential equations. *J. Math. Anal. Appl.*, 265(2):229–248, 2002.
- [4] Y. Feng, L. Li, J.-G. Liu, and X. Xu. Continuous and discrete one dimensional autonomous fractional ODEs. *Discrete and Continuous Dynamical Systems -B*, Published online. doi: 10.3934/dcdsb.2017210.
- [5] R. Gorenflo and F. Mainardi. Fractional Calculus. Springer, 1997.
- [6] H. J. Haubold, A. M. Mathai, and R. K. Saxena. Mittag-Leffler functions and their applications. *J. Appl. Math.*, 2011, 2011.
- [7] A. A. Kilbas, H. M. Srivastava, and J. J. Trujillo. Theory and applications of fractional differential equations. *North-Holland Mathematics Studies*, 204:vii x, 2006.
- [8] L. Li and J.-G. Liu. A generalized definition of Caputo derivatives and its application to fractional ODEs. arXiv preprint arXiv:1612.05103, 2017.
- [9] L. Li and J.-G. Liu. A note on deconvolution with completely monotone sequences and discrete fractional calculus. Q. Appl. Math., 78:189–198, 2017.
- [10] L. Li and J.-G. Liu. Some compactness criteria for weak solutions of time fractional PDEs. arXiv preprint arXiv:1708.08384, 2017.
- [11] L. Li, J.-G. Liu, and L. Wang. Cauchy problems for Keller-Segel type time-space fractional diffusion equation. *J. Differ. Equations. Accepted*, 2018.

- [12] Y. Luchko. Maximum principle for the generalized time-fractional diffusion equation. J. Math. Anal. Appl., 351(1):218–223, 2009.
- [13] F. Mainardi and R. Gorenflo. On Mittag-Leffler-type functions in fractional evolution processes. *J. Comput. Appl. Math.*, 118(1):283–299, 2000.
- [14] J.D. Ramirez and A.S. Vatsala. Generalized monotone iterative technique for Caputo fractional differential equation with periodic boundary condition via initial value problem. *Int. J. Differ. Equations*, 2012, 2012.
- [15] V. Vergara and R. Zacher. Optimal decay estimates for time-fractional and other nonlocal subdiffusion equations via energy methods. SIAM J. Math. Anal., 47(1):210–239, 2015.
- [16] D. G. Weis. Asymptotic behavior of some nonlinear Volterra integral equations. J. Math. Anal. Appl., 49(1):59–87, 1975.