Triggered Gradient Tracking for Asynchronous Distributed Optimization

Guido Carnevale, Ivano Notarnicola, Lorenzo Marconi, Giuseppe Notarstefano

Abstract—This paper proposes ASYNCHRONOUS TRIGGERED GRADIENT TRACKING, i.e., a distributed optimization algorithm to solve consensus optimization over networks with asynchronous communication. As a building block, we devise the continuous-time counterpart of the recently proposed (discretetime) distributed gradient tracking called CONTINUOUS GRA-DIENT TRACKING. By using a Lyapunov approach, we prove exponential stability of the equilibrium corresponding to agents' estimates being consensual to the optimal solution, with arbitrary initialization of the local estimates. Then, we propose two triggered versions of the algorithm. In the first one, the agents continuously integrate their local dynamics and exchange with neighbors their current local variables in a synchronous way. In ASYNCHRONOUS TRIGGERED GRADIENT TRACKING, we propose a totally asynchronous scheme in which each agent sends to neighbors its current local variables based on a triggering condition that depends on a locally verifiable condition. The triggering protocol preserves the linear convergence of the algorithm and avoids the Zeno behavior, i.e., an infinite number of triggering events over a finite interval of time is excluded. By using the stability analysis of CONTINUOUS GRADIENT TRACKING as a preparatory result, we show exponential stability of the equilibrium point holds for both triggered algorithms and any estimate initialization. Finally, the simulations validate the effectiveness of the proposed methods on a data analytics problem, showing also improved performance in terms of interagent communication.

I. INTRODUCTION

As the devices with computation and communication capabilities are growing all around us, several contexts ranging from machine learning to autonomous vehicles and cooperative robotics, need to control networks of communication and computing agents. Tasks must be performed in an independent and cooperative way, without resorting to a centralized unit. Often, these tasks can be mathematically posed as distributed optimization problems. For this reason, distributed optimization has become an active research area, see [1]–[3] for recent overviews. Some applications tasks can be performed relying on distributed continuous-time optimization schemes. In [4], a distributed continuous-time optimization algorithm is proposed to solve a cost-coupled convex problem over a weighted digraph. A constrained convex problem is solved in [5] for a network of agents having local, second-order dynamics. In [6], a nonsmooth convex optimization problem with local constraints is solved by using a novel distributed continuoustime projected algorithm. In [7], a distributed optimization problem with general differentiable convex objective functions is studied for continuous-time multi-agent systems with singleintegrator dynamics. Objective functions subjected to bounds, equality, and inequality constraints have been taken into account in [8] and addressed with a proportional-integral protocol. A distributed optimization problem on directed graphs with nonconvex local objective functions is tackled in [9] by using an adaptive continuous-time algorithm. In [10], a passivity-based approach is used to prove the convergence of distributed continuous-time schemes for both unconstrained and constrained scenarios, also in presence of communication delays. In [11], a continuous-time distributed optimization algorithm is shown to possess exponential convergence properties by decomposing it into a set of input feedforward passive systems that interact with each other. Paper [12] proposes a continuous-time optimization algorithm designed by taking inspiration from the existing discrete-time algorithm known as Newton-Raphson method. In fact, a part of works is recently trying to study the convergence properties of dynamic systems representing the continuous counterpart of existing iterative optimization algorithms. This line of research starts with the work [13], where a second order differential equation is presented as the continuous counterpart of Nesterov's accelerated gradient method. The authors of [14] propose a systematic way to go from continuous-time curves generated by a Lagrangian functional to a family of discrete-time accelerated algorithms. A connection between a Lyapunov approach and the socalled estimating sequence analysis (typically adopted for momentum-based algorithms) is proposed in [15] to analyze accelerated optimization methods. In [16], high resolution ordinary differential equations are introduced to better understand the continuous-time counterpart of Nesterov algorithm and heavy ball algorithm. Paper [17] studies the first order mirror descent algorithm by deriving ordinary differential equations from duality gaps.

1

Distributed continuous-time schemes rely on continuoustime communication among the network agents. Being such a mechanism not implementable in practice, a research effort has been also devoted to study continuous-time distributed algorithms with discrete-time communication. In this regard, the authors of [18] propose a continuous-time optimization algorithm for solving a cost-coupled problem in a distributed manner. Moreover, they also propose variants of the plain algorithm in which the communication between agents occurs both in a periodic and event-triggered fashion. In [19], the same scenario is addressed with a continuous-time algorithm

The authors are with the Department of Electrical, Electronic and Information Engineering, University of Bologna, Bologna, Italy, {name.lastname}@unibo.it. This result is part of a project that has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No 638992 - OPT4SMART).

in which the agents can only provide quantized information to neighbors with an event-triggered protocol. In [20], a continuous-time algorithm with event-triggered communication inspired from [18] is extended to reject external disturbances by relying on internal model concepts. Paper [21] employs a subgradient method is employed with an eventtriggered communication policy. In [22] a continuous-time distributed optimization algorithm with second order dynamics both with continuous communication and event-triggered communication between agents is proposed. The work [23] considers a quadratic optimization problem considered and a continuous-time algorithm with event-triggered communication is proposed to solve it. Finally, nonconvex optimization problems are addressed in [24] with an event-triggered implementation of the distributed gradient descent.

The main contribution of this paper is the development of three distributed algorithms for strongly convex consensus optimization problems over networks along with the theoretical proof of their linear convergence rate¹. Specifically, the development of two of them rely on alternative communication paradigms that allow for a practical implementation of continuous-time-based schemes on real devices. The first algorithm is called CONTINUOUS GRADIENT TRACKING and has been designed as the continuous-time counterpart of the recently proposed (discrete-time) distributed gradient tracking. In the second algorithm, agents send their own variables to neighbors ruled by a synchronous triggering condition. Whereas, the third algorithm implements an asynchronous communication protocol among the agents that trigger according to locally verifiable triggering conditions. By relying on a Lyapunov approach, we first develop a system theoretical stability analysis to show that the equilibrium of a dynamical system equivalent to CONTINUOUS GRADIENT TRACKING is globally exponentially stable. The equilibrium point corresponds to agents' solution estimates being consensual and equal to the optimal solution of the original optimization problem. Then, this result is extended to handle the stability analysis of the two triggered algorithms. Indeed, the stability of their equilibrium is addressed by recasting the triggered algorithms as perturbed versions of CONTINUOUS GRADI-ENT TRACKING. Under suitable conditions on the triggering protocol, exponential stability of CONTINUOUS GRADIENT TRACKING can be proved with arbitrary initial estimates. The asynchronous version of the algorithm uses an additional auxiliary variable and a specific triggering condition. These features exclude the Zeno behavior (i.e., an infinite number of events in a finite period of time) while preserving the linear convergence. As a side result, we provide a robust stability guarantee of our algorithms against inexactness computations and/or communications.

As mentioned above, the discrete-time method called gradient tracking is instrumental for our novel triggered algorithms. It has been obtained as an extension of [25], [26], in which the gradient method is combined with consensus, by introducing a "tracking action". Such tracking action is based on the dynamic average consensus (see [27], [28]) in order to let each agent estimate the gradient of a total cost function, which is only partially locally known. There exist several variants of the gradient tracking, see [29]–[38]. A controlbased analysis of this algorithm has been proposed in [39], where a suitable change of coordinates is considered and turns out to be fundamental also for the analysis performed in this paper. Finally, the approach in [39] has been exploited to design gradient tracking algorithms with sparse (non-necessarily diagonal) gains in [40].

The paper is organized as follows. The problem setup is given in Section II. In Section III the CONTINUOUS GRADIENT TRACKING is derived along with its convergence properties. In Section IV the triggered algorithms are proposed and analyzed. Section V discusses robustness aspects, while numerical simulations are provided in Section VI. The proofs are all deferred to the Appendix.

Notation: The matrix $M \in \mathbb{R}^{n \times n}$ is said to be Hurwitz if its spectrum, denoted as $\sigma(M)$, lies in the open lefthalf complex plane. The identity matrix in $\mathbb{R}^{m \times m}$ is I_m , while 0_m is the all-zero matrix in $\mathbb{R}^{m \times m}$. The vector of Nones is denoted by 1_N , while **1** denotes $1_N \otimes I_d$ with \otimes being the Kronecker product. The vertical concatenation of the column vectors v_1 and v_2 is $\operatorname{col}(v_1, v_2)$. We denote as $\operatorname{diag}(m_1, \ldots, m_N) \in \mathbb{R}^{N \times N}$ the diagonal matrix with diagonal elements m_1, \ldots, m_N . Similarly, $\operatorname{blkdiag}(M_1, \ldots, M_N)$ is the block diagonal matrix whose *i*-th block is $M_i \in \mathbb{R}^{d_i \times d_i}$.

II. PROBLEM SET-UP AND PRELIMINARIES

A. Optimization and Network Setup

In this paper, we consider a network of N agents that aim at solving optimization problems in the form

$$\min_{x \in \mathbb{R}^d} \sum_{i=1}^N f_i(x), \tag{1}$$

with the cost function $f_i : \mathbb{R}^d \to \mathbb{R}$ known to agent *i* only, for all $i \in \{1, \ldots, N\}$. Let the following hold.

Assumption 2.1: For all *i*, the function $f_i : \mathbb{R}^d \to \mathbb{R}$ is strongly convex with coefficient $\alpha > 0$.

Assumption 2.2: For all *i*, the function $f_i : \mathbb{R}^d \to \mathbb{R}$ has Lipschitz continuous gradient with constant L > 0.

Notice that Assumption 2.1 ensures that problem (1) has a unique optimal solution, denoted by $x^* \in \mathbb{R}^d$.

The N agents cooperate by exchanging information among neighbors to solve problem (1). Their interaction is modeled as a graph $\mathcal{G} = (\{1, \ldots, N\}, \mathcal{E})$, with $\mathcal{E} \subset \{1, \ldots, N\}^2$ being the edge set. If an edge (i, j) belongs to \mathcal{E} , then agents i and j can exchange information, otherwise not. The set of neighbors of an agent i is $\mathcal{N}_i := \{j \in \{1, \ldots, N\} \mid (i, j) \in \mathcal{E}\}$ and includes i itself, i.e., \mathcal{G} contains self-loops. We associate to the graph \mathcal{G} a symmetric weighted adjacency matrix $\mathcal{W} \in \mathbb{R}^{N \times N}$ whose entries match the graph, i.e., $[\mathcal{W}]_{ij} > 0$ whenever $(i, j) \in \mathcal{E}$ and $[\mathcal{W}]_{ij} = 0$ otherwise. The weighted degree of an agent is defined as $d_i = \sum_{j \in \mathcal{N}_i} [\mathcal{W}]_{ij}$. Finally, we associate to \mathcal{G} the so-called Laplacian matrix defined as $\mathcal{L} := \mathcal{D} - \mathcal{W}$, where $\mathcal{D} := \text{diag}(d_1, \ldots, d_N) \in \mathbb{R}^{N \times N}$. Let the following assumption of the network holds.

¹"Linear convergence to an optimum" in optimization and "Exponential stability of an equilibrium" in control are two expressions that refer to closely related concepts. In our paper they are equivalent.

Assumption 2.3: G is undirected and connected.

B. Discrete Gradient Tracking

Let us recall the discrete-time gradient tracking, recently proposed in the literature [29]–[38]. At iteration $k \in \mathbb{N}$, each agent *i* maintains a local estimate $\mathbf{x}_{i,k} \in \mathbb{R}^d$ of the optimal solution of problem (1) and an auxiliary state $\mathbf{s}_{i,k} \in \mathbb{R}^d$. These states are iteratively updated by agent *i* based on (*i*) the current value of the its local cost function gradient $\nabla f_i(\mathbf{x}_{i,k})$ and (*ii*) the information received from its neighbors according to

$$\mathbf{x}_{i,k+1} = \sum_{j \in \mathcal{N}_i} [\mathcal{W}]_{ij} \mathbf{x}_{j,k} - \gamma \,\mathbf{s}_{i,k} \tag{2a}$$

$$\mathbf{s}_{i,k+1} = \sum_{j \in \mathcal{N}_i} [\mathcal{W}]_{ij} \mathbf{s}_{j,k} + \nabla f_i(\mathbf{x}_{i,k+1}) - \nabla f_i(\mathbf{x}_{i,k}), \qquad (2b)$$

where $\gamma > 0$ is a stepsize, while each $[\mathcal{W}]_{ij}$ is the (i, j)-th element of a symmetric doubly stochastic matrix \mathcal{W} , i.e., such that $[\mathcal{W}]_{ij} \geq 0$ and $\sum_{i=1}^{N} [\mathcal{W}]_{ij} = \sum_{i=1}^{N} [\mathcal{W}]_{ji} = 1$ for all $j = \{1, \ldots, N\}$. The distributed algorithm (2) can be written into an aggregate form as

$$\mathbf{x}_{k+1} = W\mathbf{x}_k - \gamma \,\mathbf{s}_k \tag{3a}$$

$$\mathbf{s}_{k+1} = W\mathbf{s}_k + \nabla \mathbf{f}(\mathbf{x}_{k+1}) - \nabla \mathbf{f}(\mathbf{x}_k).$$
(3b)

where $\mathbf{x}_k := \operatorname{col}(\mathbf{x}_{1,k}, \dots, \mathbf{x}_{N,k})$, $\mathbf{s}_k := \operatorname{col}(\mathbf{s}_{1,k}, \dots, \mathbf{s}_{N,k})$, and $\nabla \mathbf{f}(\mathbf{x}_k) := \operatorname{col}(\nabla f_1(\mathbf{x}_{1,k}), \dots, \nabla f_N(\mathbf{x}_{N,k}))$ collect all the local variables of the agents while we set $W := \mathcal{W} \otimes I_d$. It can be proved that under the initialization $\mathbf{s}_0 = \nabla \mathbf{f}(\mathbf{x}_0)$, the algorithm in (3) converges to a point corresponding to a consensual optimal solution of problem (1), see [29]–[38].

III. CONTINUOUS GRADIENT TRACKING

In this section, we introduce CONTINUOUS GRADIENT TRACKING which is a novel continuous-time, distributed algorithm to solve problem (1). We first show how to derive it as the continuous-time counterpart of the discrete-time algorithm recalled in (3). Then, the convergence result to the optimal solution of problem (1) is provided.

A. From Discrete to Continuous

Let us first introduce an alternative, causal formulation of the (discrete) gradient tracking. Consider the nonlinear change of coordinates proposed in [39] that moves the gradient term at k + 1 to the left-hand side in (3). Therefore, let $z_{i,k} :=$ $s_{i,k} - \nabla f_i(x_{i,k})$ and rewrite (3) as

$$\begin{aligned} \mathbf{x}_{i,k+1} &= \sum_{j \in \mathcal{N}_i} [\mathcal{W}]_{ij} \mathbf{x}_{j,k} - \gamma \, \mathbf{z}_{i,k} - \gamma \, \nabla f_i(\mathbf{x}_{i,k}) \\ \mathbf{z}_{i,k+1} &= \sum_{j \in \mathcal{N}_i} [\mathcal{W}]_{ij} \mathbf{z}_{j,k} + \sum_{j \in \mathcal{N}_i} [\mathcal{W}]_{ij} \nabla f_j(\mathbf{x}_{j,k}) - \nabla f_i(\mathbf{x}_{i,k}). \end{aligned}$$

The latter can be also written into an aggregate form as

$$\mathbf{x}_{k+1} = W\mathbf{x}_k - \gamma \,\mathbf{z}_k - \gamma \,\nabla \mathbf{f}(\mathbf{x}_k) \tag{5a}$$

$$\mathbf{z}_{k+1} = W\mathbf{z}_k - (I_{Nd} - W)\nabla \mathbf{f}(\mathbf{x}_k), \tag{5b}$$

where $z_k := col(z_{1,k}, ..., z_{N,k})$ while x_k , $\nabla f(x_k)$, W and γ are the same as in Section II-B.

Following the arguments proposed in [13] for a centralized optimization algorithm, we interpret the sequences x_k and z_k of (5) as sampled versions of two continuous-time signals x(t) and z(t). These signals are assumed to be smooth. According to this interpretation, the stepsize $\gamma > 0$ can be then seen as the sampling time characterizing a discretization procedure. Figure 1 shows a graphical representation of the discretization of the continuous signal x(t) resulting in the discrete sequence x_k .




Fig. 1. x_k as sampled version of x(t).

 \square

Informally, we start from the intuition

$$\mathbf{x}_k \approx x(t) \Big|_{t=k\gamma}, \qquad \mathbf{z}_k \approx z(t) \Big|_{t=k\gamma}$$

in which the discrete time k is obtained by setting $k := t/\gamma$ with t being the continuous time. For any fixed t, by choosing an arbitrarily small stepsize γ , we can consider the following approximations

$$x(t) \approx \mathbf{x}_{t/\gamma} = \mathbf{x}_k, \quad x(t+\gamma) \approx \mathbf{x}_{(t+\gamma)/\gamma} = \mathbf{x}_{k+1}.$$

The same clearly holds also for the sequence z_k . With these approximations in mind, we can write the following Taylor expansions

$$\mathbf{x}_{k+1} = x(t) \Big|_{t=(k+1)\gamma} = x(t) \Big|_{t=k\gamma} + \gamma \dot{x}(t) \Big|_{t=k\gamma} + o(\gamma) \quad (6a)$$

$$z_{k+1} = z(t) \Big|_{t=(k+1)\gamma} = z(t) \Big|_{t=k\gamma} + \gamma \dot{z}(t) \Big|_{t=k\gamma} + o(\gamma), \quad (6b)$$

where $o(\gamma)$ collects the higher order terms of the expansions. As γ goes to zero, the higher order terms in (6) can be neglected, leading to

$$\dot{x}(t) = \frac{1}{\gamma} (\mathbf{x}_{k+1} - \mathbf{x}_k) = \frac{W - I_{Nd}}{\gamma} x(t) - z(t) - \nabla \mathbf{f}(x(t))$$
$$\dot{z}(t) = \frac{1}{\gamma} (\mathbf{z}_{k+1} - \mathbf{z}_k) = \frac{W - I_{Nd}}{\gamma} z(t) - \frac{W - I_{Nd}}{\gamma} \nabla \mathbf{f}(x(t)),$$

which can be rewritten as

$$\dot{x}(t) = -L_{\gamma}x(t) - z(t) - \nabla \mathbf{f}(x(t))$$
(7a)

$$\dot{z}(t) = -L_{\gamma}z(t) - L_{\gamma}\nabla\mathbf{f}(x(t)), \tag{7b}$$

where $L_{\gamma} := (I_{Nd} - W)/\gamma$.

B. Algorithm Definition and Convergence

The Ordinary Differential Equation (ODE) in (7) involves matrices whose structure depends on the preceding derivation. However, one may consider any weighted Laplacian matrix \mathcal{L} associated to \mathcal{G} . Therefore, we define the CONTINUOUS GRADIENT TRACKING as

$$\begin{bmatrix} \dot{x}(t) \\ \dot{z}(t) \end{bmatrix} = \begin{bmatrix} -L & -I_{Nd} \\ 0 & -L \end{bmatrix} \begin{bmatrix} x(t) \\ z(t) \end{bmatrix} - \begin{bmatrix} I_{Nd} \\ L \end{bmatrix} \nabla \mathbf{f}(x(t)), \quad (8)$$

Fig. 2. Block diagram representation of system (8).

where $L \in \mathbb{R}^{Nd \times Nd}$ is given by $L := \mathcal{L} \otimes I_d$.

It is useful to also provide a local view of algorithm (8), i.e., from the perspective of a generic agent *i*. The *i*-th blockcomponents of x(t) and z(t) corresponds, respectively, to the local states $x_i(t)$ and $z_i(t)$ of agent *i*. The state $x_i(t)$ represents the local estimate at time *t* of the optimal solution of problem (1) while $z_i(t) \in \mathbb{R}^d$ is an auxiliary state. Set $\mathcal{W} := \mathcal{D} - \mathcal{L}$ (with \mathcal{D} being the degree matrix of \mathcal{G}) and let w_{ij} be its (i, j)-th entry. Exploiting the sparsity in \mathcal{W} , the *i*-th block-components of (8) can be then written as

$$\dot{x}_{i}(t) = -\sum_{j \in \mathcal{N}_{i}} w_{ij} \left(x_{i}(t) - x_{j}(t) \right) - z_{i}(t) - \nabla f_{i}(x_{i}(t))$$
(9a)

$$\dot{z}_{i}(t) = -\sum_{j \in \mathcal{N}_{i}} w_{ij} \left(z_{i}(t) - z_{j}(t) \right) -\sum_{j \in \mathcal{N}_{i}} w_{ij} \left(\nabla f_{i}(x_{i}(t)) - \nabla f_{j}(x_{j}(t)) \right).$$
(9b)

The following theorem establishes the convergence properties of CONTINUOUS GRADIENT TRACKING.

Theorem 3.1: Consider the CONTINUOUS GRADIENT TRACKING distributed algorithm described by (8). Let Assumptions 2.1, 2.2, 2.3 hold and pick any col(x(0), z(0)) such that $\mathbf{1}^{\top} z(0) = 0$. Then, there exist $a_1 > 0$ and $a_2 > 0$ such that

$$||x_i(t) - x^*|| \le a_1 \exp(-a_2 t), \quad \forall i \in \{1, \dots, N\}.$$

See Appendix A for the proof.

We underline that both a_1 and a_2 in Theorem 3.1 depend on (*i*) the distance between the initial conditions and the system equilibrium and (*ii*) the problem properties as, e.g., the network connectivity, the strong convexity parameter of the cost and the Lipschitz constants of the cost gradients. The same observation consistently applies to the subsequent results.

We point out that the initialization $\mathbf{1}^{\top} z(0) = 0$ can be obtained in a fully distributed way by simply setting each $z_i(0) = 0$, for all $i \in \{1, ..., N\}$. The proof of Theorem 3.1 is based on a Lyapunov analysis relying on the feedback structure of (8) as represented in Figure 2.

Specifically, noticing that $col(1x^*, -\nabla f(1x^*))$ is the unique equilibrium point for system (8) and by exploiting the initialization, we perform a sequence of coordinate changes to obtain an equivalent, reduced system formulation. The equivalent system is characterized by a (marginally stable) linear part, associated to the consensus mechanism, which is perturbed by a nonlinear (feedback) term depending on the gradient ∇f . We then prove the exponential stability of the equilibrium by

designing a quadratic Lyapunov function based on the linear part only and bounding the nonlinear gradient term using the strong convexity and the Lipschitz continuity.

Remark 3.2: The expression of the discrete-time dynamics as in (5) turns out to be crucial in the derivation of its continuous-time version. In fact, one can check that

$$\begin{aligned} \nabla \mathbf{f}(\mathbf{x}_{k+1}) &= \nabla \mathbf{f}\left(x(t)\Big|_{t=(k+1)\gamma}\right) \\ &= \nabla \mathbf{f}\left(x(t)\Big|_{t=k\gamma} + \gamma \dot{x}(t)\Big|_{t=k\gamma} + o(\gamma)\right) \\ &= \nabla \mathbf{f}\left(x(t)\Big|_{t=k\gamma}\right) + \gamma \nabla^2 \mathbf{f}\left(x(t)\Big|_{t=k\gamma}\right) \dot{x}(t)\Big|_{t=k\gamma} + o(\gamma). \end{aligned}$$

Thus, the arguments presented in Section III-A applied to the algorithm in its original coordinates (3) results in

$$\begin{bmatrix} \dot{x}(t) \\ \dot{s}(t) \end{bmatrix} = \begin{bmatrix} -L_{\gamma} & -I_{Nd} \\ -\nabla^2 \mathbf{f}(x(t))L_{\gamma} & -(L_{\gamma} + \nabla^2 \mathbf{f}(x(t))) \end{bmatrix} \begin{bmatrix} x(t) \\ s(t) \end{bmatrix},$$

where s(t) is the continuous counterpart of s_k while $\nabla^2 \mathbf{f}(x(t)) := \text{blkdiag}(\nabla^2 f_1(x_1(t)), \dots, \nabla^2 f_N(x_N(t))) \in \mathbb{R}^{Nd \times Nd}$. Notice that these coordinates involve the second-order matrix $\nabla^2 \mathbf{f}(\cdot)$ which usually requires a non-negligible computational complexity and, in certain applications, is not even known.

IV. TRIGGERED GRADIENT TRACKING

The CONTINUOUS GRADIENT TRACKING would require communication among agents at all $t \ge 0$. Clearly, this prevents its practical implementation on real devices that require time-slotted communication. This issue is addressed next by proposing two alternative schemes in which interagent communication is triggered synchronously and asynchronously, respectively.

Let $\{t_i^{k_i}\}_{k_i \in \mathbb{N}}$, be the sequence of time instants at which agent *i* sends its states (x_i, z_i) and ∇f_i to its neighbors $j \in \mathcal{N}_i$. Consistently, at time $t_j^{k_j}$, agent *i* receives the updated variables from its neighbor $j \in \mathcal{N}_i$. Let $\{\tilde{t}^k\}_{k\geq 0}$ be the ordered sequence of all the triggering times that occurred in the network. Then, given any $t \in [\tilde{t}^k, \tilde{t}^{k+1})$, let us introduce, for all $i \in \{1, \ldots, N\}$, the shorthands

$$\begin{aligned} \hat{x}_{i}^{k} &:= x_{i}(t) \Big|_{t=\inf_{k_{i} \in \mathbb{N}} \left\{ t_{i}^{k_{i}} \ge \tilde{t}^{k} \right\}} \\ \hat{z}_{i}^{k} &:= z_{i}(t) \Big|_{t=\inf_{k_{i} \in \mathbb{N}} \left\{ t_{i}^{k_{i}} \ge \tilde{t}^{k} \right\}} \\ \nabla f_{i}^{k} &:= \nabla f_{i}(x_{i}(t)) \Big|_{t=\inf_{k_{i} \in \mathbb{N}} \left\{ t_{i}^{k_{i}} \ge \tilde{t}^{k} \right\}}. \end{aligned}$$
(10)

Quantities in (10) represent the most updated values in the network within the considered time interval. Under the described communication paradigm, we propose to modify the local dynamics in (9) as follows

$$\dot{x}_{i}(t) = -\sum_{j \in \mathcal{N}_{i}} w_{ij} \left(\hat{x}_{i}^{k} - \hat{x}_{j}^{k} \right) - z_{i}(t) - \nabla f_{i}(x_{i}(t))$$
(11a)

$$\dot{z}_i(t) = -\sum_{j \in \mathcal{N}_i} w_{ij} \left(\hat{z}_i^k - \hat{z}_j^k \right) - \sum_{j \in \mathcal{N}_i} w_{ij} \left(\nabla f_i^k - \nabla f_j^k \right), \quad (11b)$$

for all $t \in [\tilde{t}^k, \tilde{t}^{k+1})$. Within the k-th period, the variable z_i behaves as an integrator. As for the variable x_i , it is a local gradient flow compensated with an integral action z_i and a constant consensus-error-like term. Agent *i* does not use its own variables $x_i(t)$, $z_i(t)$, and $\nabla f_i(x_i(t))$ in the consensus mixing terms, but it rather uses their sampled version. This fact allows one to preserve the theoretical consensus properties of the original scheme (9).

As one can expect, the specific rule to choose the triggering time $t_i^{k_i}$ will play a crucial role in the convergence properties of the resulting algorithms.

A. SYNCHRONOUS TRIGGERED GRADIENT TRACKING

We start by presenting SYNCHRONOUS TRIGGERED GRA-DIENT TRACKING, obtained by imposing a synchronous communication among agents. Specifically, in this protocol each agent $i \in \{1, ..., N\}$ sends its local variables to its neighbors at common instants of time chosen according to

$$t_i^{k_i+1} := t_i^{k_i} + \Delta, \tag{12}$$

for some common $\Delta > 0$ and with $t^{0_i} = t_0$ for all $i \in \{1, \ldots, N\}$. Intuitively, the greater Δ , the more inter-agent communication reduces. On the other hand, the greater Δ , the more the triggered algorithmic evolution moves away from the behavior of CONTINUOUS GRADIENT TRACKING. The next theorem gives theoretical guarantees about the maximum admissible value for Δ .

Theorem 4.1: Consider the algorithm in (11) with the synchronous communication protocol given by (12). Let Assumptions 2.1, 2.2, 2.3 hold and pick any col(x(0), z(0)) such that $\mathbf{1}^{\top} z(0) = 0$. Then, there exist $\Delta^* > 0$, $a_3 > 0$, and $a_4 > 0$ such that for any $\Delta \in (0, \Delta^*)$, it holds

$$||x_i(t) - x^*|| \le a_3 \exp(-a_4 t), \quad \forall i \in \{1, \dots, N\}.$$

See Appendix A for the proof.

For all $t \in [\tilde{t}^k, \tilde{t}^{k+1})$, the aggregate form of (11) reads as

$$\begin{bmatrix} \dot{x}(t) \\ \dot{z}(t) \end{bmatrix} = H \begin{bmatrix} x(t) \\ z(t) \end{bmatrix} + B_1 \nabla \mathbf{f}(x(t)) + B_2 \begin{bmatrix} \hat{x}^k \\ \hat{z}^k \\ \nabla \mathbf{f}^k \end{bmatrix}, \quad (13)$$

where $\hat{x}^k := col(\hat{x}_1^k, ..., \hat{x}_N^k), \, \hat{z}^k := col(\hat{z}_1^k, ..., \hat{z}_N^k), \, \nabla \mathbf{f}^k := col(\nabla f_1^k, ..., \nabla f_N^k)$, and

$$H := \begin{bmatrix} 0 & -I \\ 0 & 0 \end{bmatrix}, \quad B_1 := \begin{bmatrix} -I \\ 0 \end{bmatrix}, \quad B_2 := \begin{bmatrix} -L & 0 & 0 \\ 0 & -L & -L \end{bmatrix}$$

The proof of Theorem 4.1 relies on a proper reformulation of the dynamics (13) as a perturbed instance of the CONTINUOUS GRADIENT TRACKING system (8). In particular, we show that the periodic triggering law (12) gives rise to a perturbation term that vanishes at the equilibrium point (see, e.g., [41, Chapter 9] for the notion of vanishing perturbation) and that can be arbitrarily bounded through the parameter Δ . Thus, we can consider the same Lyapunov function V used in the proof of Theorem 3.1 and pick a sufficiently small Δ to show that the perturbation does not alter the sign of the derivative of V to conclude the proof.

B. ASYNCHRONOUS TRIGGERED GRADIENT TRACKING

We now investigate the case in which the agents choose their triggering time $t_i^{k_i}$ in a fully asynchronous way giving rise to an algorithm termed ASYNCHRONOUS TRIGGERED GRADI-ENT TRACKING. This scheme is motivated by the fact that the synchronous communication executed according to (12) is rather conservative with a consequent non-efficient usage of the available resources. An asynchronous communication protocol allows agents to exchange information only when really needed. This requires a modification of the synchronous scheme. In particular, each agent has to check a local triggering condition and to maintain an additional auxiliary variable. The latter is important to take into account the so-called Zeno behavior. Specifically, an infinite number of triggerings over a finite interval of time must be avoided. Indeed, for agent *i*, a triggering law suffers from the Zeno effect if

$$\lim_{k_i \to \infty} t_i^{k_i} = \sum_{k_i=0}^{\infty} (t_i^{k_i+1} - t_i^{k_i}) = t_i^{\infty},$$

for some (finite) $t_i^{\infty} > 0$ termed the Zeno time.

The local dynamics is again described by (11). But, in order to perform communication only when needed, each agent *i* chooses the next triggering time instant $t_i^{k_i+1}$ according to a locally verifiable condition. A possible choice for such a condition may be

$$k_{i}^{k_{i}+1} := \inf_{t > t_{i}^{k_{i}}} \left\{ \left\| e_{i}(t) \right\| > \lambda \left\| h_{i}(t) \right\| \right\},$$
(14)

with $e_i(t) := col(x_i(t) - \hat{x}_i^k, z_i(t) - \hat{z}_i^k, \nabla f_i(x_i(t)) - \nabla f_i^k),$ $h_i(t) := z_i(t) + \nabla f_i(x_i(t))$, and $\lambda > 0$ a constant to be properly specified later. The rationale for the triggering mechanism is to (i) keep the triggered scheme close to the original dynamics (8), and (ii) avoid the Zeno behavior. To this end, the righthand side of the inequality within (14) must be asymptotically vanishing when the algorithm approaches a steady state. This, in turn, gives rise to a vanishing quantity on the left term of the inequality. Indeed, looking also to the discrete-time version (2), the (local) quantity $z_i(t) + \nabla f_i(x_i(t))$ can be seen as a proxy for $\sum_{i=1}^{N} \nabla f_i(x_i(t))$, i.e., a quantity that vanishes at a consensual optimal solution. However, $||h_i(t)||$ vanishes not only when the algorithm approaches the equilibrium, but also if $(x_i(t), z_i(t)) \in \mathcal{S}_i := \{(x_i, z_i) \in \mathbb{R}^{2d} \mid z_i = -\nabla f_i(x_i)\},\$ possibly giving rise to the Zeno behavior. Thus, in order to exclude this situation, the triggering condition (14) is further modified as

$$t_i^{k_i+1} := \inf_{t > t_i^{k_i}} \left\{ \|e_i(t)\| > \lambda \|h_i(t)\| + |\xi_i(t)| \right\},$$
(15)

where $\xi_i \in \mathbb{R}$ is a local, auxiliary variable maintained by each agent *i* evolving as

$$\xi_i(t) = -\nu \xi_i(t), \tag{16}$$

where $\nu > 0$ is a parameter ruling the decay of $\xi_i(t)$.

As formally shown next, if the ξ_i are initialized to nonzero values, then algorithm (11) with triggering law (15) does not incur in the Zeno behavior.

Theorem 4.2: Consider the algorithm described by (11) with the asynchronous communication protocol given

by (15). Let Assumptions 2.1, 2.2, 2.3 hold and pick any $\operatorname{col}(x(0), z(0), \xi(0))$ such that $\mathbf{1}^{\top} z(0) = 0$ and with $\xi(0) = \operatorname{col}(\xi_1(0), \ldots, \xi_N(0)) \neq 0$. Then, there exist $\lambda^* > 0$, $\nu^* > 0$, $a_5 > 0$, and $a_6 > 0$ such that for any $\lambda \in (0, \lambda^*)$ in (15) and any $\nu > \nu^*$, it holds

$$||x_i(t) - x^*|| \le a_5 \exp(-a_6 t), \quad \forall i \in \{1, \dots, N\}.$$

Moreover, system (11) does not exhibit the Zeno behavior. \Box See Appendix A for the proof.

As for Theorem 4.1, also this proof is based on a proper reformulation of the aggregate form of ASYNCHRONOUS TRIG-GERED GRADIENT TRACKING (which is still given by (13)) as a perturbed instance of the CONTINUOUS GRADIENT TRACKING dynamics (8) with a vanishing perturbation. For this asynchronous triggering law, (15), an upper bound on the perturbation magnitude is provided. It is proportional to (i)the term $\lambda \|z(t) + \nabla \mathbf{f}(x(t))\|$, which, as already stated, represents a surrogate for the distance from the equilibrium point $col(\mathbf{1}x^{\star}, -\nabla \mathbf{f}(\mathbf{1}x^{\star}))$, and *(ii)* to the exponentially decaying term $\|\xi\|$. Thus, considering a Lyapunov function derived from the one used in Theorem 3.1, it is possible to show that, by picking suitable λ and ν , the perturbation does not affect the sign of the Lyapunov derivative. A specific choice for λ^* and ν^{\star} can be obtained by exploiting the bounds derived in the proof.

V. ROBUSTNESS AGAINST INEXACT COMPUTATION

Let us consider a more general scenario in which agents can access only inexact evaluations of their local state (x_i, z_i) and/or of the local gradients ∇f_i . Let $v_{i,\nabla}(t) \in \mathbb{R}^d$ represents the mismatch between the exact value of $\nabla f_i(x_i(t))$ and the one available to agent *i* for the local updates. The presence of this mismatch may be due to several reasons as, e.g., quantization errors of the computing unit, measurement errors in the sensor providing $\nabla f_i(x_i(t))$, or model uncertainties affecting the available gradient. Similarly, also mismatches affecting the states x_i and z_i can be considered. Thus, we consistently introduce $v_{i,x}(t) \in \mathbb{R}^d$ and $v_{i,z}(t) \in \mathbb{R}^d$ to model such uncertainties. This framework can be formalized by writing

$$\begin{bmatrix} \dot{x}(t) \\ \dot{z}(t) \end{bmatrix} = \begin{bmatrix} -L & -I \\ 0 & -L \end{bmatrix} \begin{bmatrix} x(t) \\ z(t) \end{bmatrix} - \begin{bmatrix} I \\ L \end{bmatrix} \nabla \mathbf{f}(x(t)) + \delta_1 B_2 e(t) + B_3 v(t), \quad (17)$$

where $e(t) := \operatorname{col}(\hat{x}^k - x(t), \hat{z}^k - z(t), \nabla \mathbf{f}^k - \nabla \mathbf{f}(x(t)))$ collects (possible) mismatches due to discrete-time communication and $v(t) := \operatorname{col}(v_{\nabla}(t), v_x(t), v_z(t))$ collects the mentioned local mismatches between the gradients, the solution estimates and the auxiliary variables, the matrix B_2 has the same meaning as in (13), and B_3 is defined as

$$B_3 := \begin{bmatrix} -L & -I & -I \\ 0 & -L & -L \end{bmatrix}.$$
 (18)

Finally, δ_1 is equal to 0 for CONTINUOUS GRADIENT TRACK-ING and equal to 1 for both the SYNCHRONOUS and ASYN-CHRONOUS TRIGGERED GRADIENT TRACKING. Similarly, we denote as δ_2 is equal to 0 for both CONTINUOUS GRADI-ENT TRACKING and SYNCHRONOUS TRIGGERED GRADIENT TRACKING and equal to 1 for ASYNCHRONOUS TRIGGERED GRADIENT TRACKING.

Next, the robustness of the algorithm in terms of input-tostate stability is studied.

Proposition 5.1: Consider the algorithm described by (17). Let Assumptions 2.1, 2.2, 2.3 hold and pick any $\operatorname{col}(x(0), z(0))$ such that $\mathbf{1}^{\top} z(0) = 0$. Then, there exist a \mathcal{KL} function $g_1(\cdot)$ and a \mathcal{K}_{∞} function $g_2(\cdot)$ such that for any $x(0) \in \mathbb{R}^{Nd}$ it holds $\|x(t) - \mathbf{1}x^{\star}\| \leq g_1(\|\chi(0)\|, t) + g_2(\|v(\cdot)\|_{\infty})$, with $\chi(0) := \operatorname{col}(x(0) - \mathbf{1}x^{\star}, z(0) + \nabla \mathbf{f}(\mathbf{1}x^{\star}), \delta_2\xi(0))$ and for any $v(\cdot) \in \mathcal{L}_{\infty}^{3Nd}$.

See Appendix A for the proof.

Proposition 5.1 guarantees that within the framework modeled by (17), the proposed algorithms behave as input-tostate stable systems. Therefore, in presence of mismatches on variables and gradients, the distance between the solution of problem (1) and the computed estimates stay bounded according to the error magnitude.

VI. NUMERICAL SIMULATIONS

We next present numerical simulations to confirm and support the theoretical findings. The simulations are done using Matlab with its numerical solver "ode45" to integrate the CONTINUOUS GRADIENT TRACKING.

We consider a network of agents that want to cooperatively solve a data analytics problem in which a linear classifier must be trained. Each agent *i* is equipped with $m_i \in \mathbb{N}$ points $p_{i,1}, \ldots, p_{i,m_i} \in \mathbb{R}^d$ with binary labels $l_{i,h} \in \{-1,1\}$ for all $h \in \{1, \ldots, m_i\}$. We consider a logistic regression problem given by

$$\min_{w,b} \sum_{i=1}^{N} \sum_{h=1}^{m_i} \log \left(1 + \exp(-l_{i,h}(w^\top p_{i,h} + b)) \right) + \frac{C(\|w\|^2 + b^2)}{2},$$

where the optimization variables $w \in \mathbb{R}^{d-1}$ and $b \in \mathbb{R}$ define the separating hyperplane, while C > 0 is the socalled regularization parameter. Notice that the presence of the regularization makes the cost function strongly convex. In our simulations, we pick d = 3, $m_i = 10$ for all $i \in \{1, \ldots, N\}$, and C = 0.1.

A. CONTINUOUS GRADIENT TRACKING

In this subsection, the effectiveness of CONTINUOUS GRA-DIENT TRACKING is shown on a network of N = 50 agents communicating according to an undirected and connected Erdős-Rényi graph with parameter 0.4. In Figure 3 the convergence performances of CONTINUOUS GRADIENT TRACKING algorithm are shown. Specifically, the distance of the local estimates $x(t) := \operatorname{col}(x_1(t), \ldots, x_N(t))$ from the optimum $||x(t) - \mathbf{1}x^*||$, converges to zero exponentially fast as expected from Theorem 3.1.

²See [41, Chapter 4] for the function classes' definitions.




Fig. 3. Evolution of the distance from the optimum of local estimates generated by CONTINUOUS GRADIENT TRACKING.

B. SYNCHRONOUS AND ASYNCHRONOUS TRIGGERED GRADIENT TRACKING

In this subsection, the effectiveness of the triggered algorithms is shown for a network of N = 10 agents communicating according to an undirected and connected Erdős-Rényi graph with parameter 0.4. We tested SYNCHRONOUS TRIGGERED GRADIENT TRACKING and ASYNCHRONOUS TRIGGERED GRADIENT TRACKING for different values of their key parameters Δ and λ , respectively. Moreover, we experimentally tuned the stepsize for the discrete graident tracking as $\gamma = 0.1$ in order to optimize its convergence rate. Finally, we set $\nu = 5$ for the dynamics of ξ_i in (16). For the simulation of ASYNCHRONOUS TRIGGERED GRADIENT TRACKING, the triggering condition (cf. (15)) is checked every 0.001 seconds. Figure 4 compares the evolution of the optimality error obtained with different Δ and λ , for SYNCHRONOUS TRIGGERED GRADIENT TRACKING, ASYN-CHRONOUS TRIGGERED GRADIENT TRACKING, and the discrete gradient tracking. Specifically, the comparison is done in terms of communication rounds. The plot considers the performances of the most efficient agent, say i_{\star} , that performs the smallest number of neighboring communications in Asynchronous Triggered Gradient Tracking. As for the discrete gradient tracking, we denote, with a slight of abuse of notation, $x_{i_{\star}}(t_{i_{\star}}^{k_{i_{\star}}}) = x_{i_{\star}}^{k}$, with the sequence $\{x_{i}^{k}\}_{k\geq 0}$ generated by (5). As Figure 4 clearly highlights, the communication rounds decrease as λ increases. The same applies to Δ . In particular, we underline that ASYNCHRONOUS TRIGGERED GRADIENT TRACKING results more efficient in finding the optimal solution with respect to both SYN-CHRONOUS TRIGGERED GRADIENT TRACKING and discrete gradient tracking.

Finally, in Figure 5 each cross represents when the triggering condition occurred for each agent while running the ASYNCHRONOUS TRIGGERED GRADIENT TRACKING with $\lambda = 0.1$. The plots demonstrate how event-triggered communication effectively reduces inter-agents communication.

VII. CONCLUSIONS

In this paper, we addressed consensus optimization by proposing three novel distributed optimization algorithms. First, CONTINUOUS GRADIENT TRACKING is derived as the continuous-time counterpart of the existing discrete-time




Fig. 4. Comparison among ASYNCHRONOUS TRIGGERED GRADIENT TRACKING (ATGT), SYNCHRONOUS TRIGGERED GRADIENT TRACKING (STGT) and the discrete gradient tracking (DGT) in terms of evolution of the optimality error.




Fig. 5. Occurrence of the triggering conditions in the ASYNCHRONOUS TRIGGERED GRADIENT TRACKING.

gradient tracking algorithm. Then, by specifying proper interagent communication protocols, two triggered algorithms are derived and analyzed, a synchronous and an asynchronous one, with an algorithmic structure inspired by the continuous version. The convergence analysis of all algorithms exploited a system-theoretical approach based on a suitably defined quadratic Lyapunov function. The theoretical findings have been supported through numerical simulations.

REFERENCES

- A. Nedić and J. Liu, "Distributed optimization for control," Annual Review of Control, Robotics, and Autonomous Systems, vol. 1, pp. 77– 103, 2018.
- [2] G. Notarstefano, I. Notarnicola, and A. Camisa, "Distributed optimization for smart cyber physical net-works," *Foundations and Trends in Systems and Control*, vol. 7, no. 3, pp. 253–383, 2019.
- [3] T. Yang, X. Yi, J. Wu, Y. Yuan, D. Wu, Z. Meng, Y. Hong, H. Wang, Z. Lin, and K. H. Johansson, "A survey of distributed optimization," *Annual Reviews in Control*, vol. 47, pp. 278–305, 2019.
- [4] B. Gharesifard and J. Cortés, "Distributed continuous-time convex optimization on weight-balanced digraphs," *IEEE Transactions on Automatic Control*, vol. 59, no. 3, pp. 781–786, 2013.
- [5] Q. Liu and J. Wang, "A second-order multi-agent network for boundconstrained distributed optimization," *IEEE Transactions on Automatic Control*, vol. 60, no. 12, pp. 3310–3315, 2015.
- [6] X. Zeng, P. Yi, and Y. Hong, "Distributed continuous-time algorithm for constrained convex optimizations via nonsmooth analysis approach," *IEEE Transactions on Automatic Control*, vol. 62, no. 10, pp. 5227– 5233, 2017.
- [7] P. Lin, W. Ren, and J. A. Farrell, "Distributed continuous-time optimization: nonuniform gradient gains, finite-time convergence, and convex constraint set," *IEEE Transactions on Automatic Control*, vol. 62, no. 5, pp. 2239–2253, 2017.
- [8] S. Yang, Q. Liu, and J. Wang, "A multi-agent system with a proportionalintegral protocol for distributed constrained optimization," *IEEE Transactions on Automatic Control*, vol. 62, no. 7, pp. 3461–3467, 2017.

- [9] Z. Li, Z. Ding, J. Sun, and Z. Li, "Distributed adaptive convex optimization on directed graphs via continuous-time algorithms," *IEEE Transactions on Automatic Control*, vol. 63, no. 5, pp. 1434–1441, 2018.
- [10] T. Hatanaka, N. Chopra, T. Ishizaki, and N. Li, "Passivity-based distributed optimization with communication delays using pi consensus algorithm," *IEEE Transactions on Automatic Control*, vol. 63, no. 12, pp. 4421–4428, 2018.
- [11] M. Li, G. Chesi, and Y. Hong, "Input-feedforward-passivity-based distributed optimization over jointly connected balanced digraphs," *IEEE Transactions on Automatic Control*, vol. 66, no. 9, pp. 4117–4131, 2021.
- [12] H. Moradian and S. S. Kia, "A distributed continuous-time modified newton-raphson algorithm," *Automatica*, vol. 136, p. 109886, 2022.
- [13] W. Su, S. Boyd, and E. Candes, "A differential equation for modeling nesterov's accelerated gradient method: Theory and insights," in *Advances in Neural Information Processing Systems*, pp. 2510–2518, 2014.
- [14] A. Wibisono, A. C. Wilson, and M. I. Jordan, "A variational perspective on accelerated methods in optimization," *Proceedings of the National Academy of Sciences*, vol. 113, no. 47, pp. E7351–E7358, 2016.
- [15] A. C. Wilson, B. Recht, and M. I. Jordan, "A lyapunov analysis of accelerated methods in optimization," *Journal of Machine Learning Research*, vol. 22, no. 113, pp. 1–34, 2021.
- [16] B. Shi, S. S. Du, M. I. Jordan, and W. J. Su, "Understanding the acceleration phenomenon via high-resolution differential equations," *Mathematical Programming*, pp. 1–70, 2021.
- [17] J. Diakonikolas and L. Orecchia, "The approximate duality gap technique: A unified theory of first-order methods," *SIAM Journal on Optimization*, vol. 29, no. 1, pp. 660–689, 2019.
- [18] S. S. Kia, J. Cortés, and S. Martínez, "Distributed convex optimization via continuous-time coordination algorithms with discrete-time communication," *Automatica*, vol. 55, pp. 254–264, 2015.
- [19] S. Liu, L. Xie, and D. E. Quevedo, "Event-triggered quantized communication-based distributed convex optimization," *IEEE Transactions on Control of Network Systems*, vol. 5, no. 1, pp. 167–178, 2016.
- [20] Z. Deng, X. Wang, and Y. Hong, "Distributed optimisation design with triggers for disturbed continuous-time multi-agent systems," *IET Control Theory & Applications*, vol. 11, no. 2, pp. 282–290, 2017.
- [21] Y. Kajiyama, N. Hayashi, and S. Takai, "Distributed subgradient method with edge-based event-triggered communication," *IEEE Transactions on Automatic Control*, vol. 63, no. 7, pp. 2248–2255, 2018.
- [22] X. Yi, L. Yao, T. Yang, J. George, and K. H. Johansson, "Distributed optimization for second-order multi-agent systems with dynamic eventtriggered communication," in *IEEE Conference on Decision and Control* (CDC), pp. 3397–3402, 2018.
- [23] Z. Zhao, G. Chen, and M. Dai, "Distributed event-triggered scheme for a convex optimization problem in multi-agent systems," *Neurocomputing*, vol. 284, pp. 90–98, 2018.
- [24] T. Adachi, N. Hayashi, and S. Takai, "Distributed gradient descent method with edge-based event-driven communication for non-convex optimization," *IET Control Theory & Applications*, 2021.
- [25] A. Nedić and A. Ozdaglar, "Distributed subgradient methods for multiagent optimization," *IEEE Transactions on Automatic Control*, vol. 54, no. 1, p. 48, 2009.
- [26] A. Nedić, A. Ozdaglar, and P. A. Parrilo, "Constrained consensus and optimization in multi-agent networks," *IEEE Transactions on Automatic Control*, vol. 55, no. 4, pp. 922–938, 2010.
 [27] M. Zhu and S. Martínez, "Discrete-time dynamic average consensus,"
- [27] M. Zhu and S. Martínez, "Discrete-time dynamic average consensus," *Automatica*, vol. 46, no. 2, pp. 322–329, 2010.
- [28] S. S. Kia, B. Van Scoy, J. Cortés, R. A. Freeman, K. M. Lynch, and S. Martínez, "Tutorial on dynamic average consensus: The problem, its applications, and the algorithms," *IEEE Control Systems Magazine*, vol. 39, no. 3, pp. 40–72, 2019.
- [29] D. Varagnolo, F. Zanella, A. Cenedese, G. Pillonetto, and L. Schenato, "Newton-Raphson consensus for distributed convex optimization," *IEEE Transactions on Automatic Control*, vol. 61, no. 4, pp. 994–1009, 2016.
- [30] P. Di Lorenzo and G. Scutari, "Next: In-network nonconvex optimization," *IEEE Transactions on Signal and Information Processing over Networks*, vol. 2, no. 2, pp. 120–136, 2016.
- [31] A. Nedić, A. Olshevsky, and W. Shi, "Achieving geometric convergence for distributed optimization over time-varying graphs," *SIAM Journal on Optimization*, vol. 27, no. 4, pp. 2597–2633, 2017.
- [32] G. Qu and N. Li, "Harnessing smoothness to accelerate distributed optimization," *IEEE Transactions on Control of Network Systems*, vol. 5, no. 3, pp. 1245–1260, 2018.
- [33] J. Xu, S. Zhu, Y. C. Soh, and L. Xie, "Convergence of asynchronous distributed gradient methods over stochastic networks," *IEEE Transactions* on Automatic Control, vol. 63, no. 2, pp. 434–448, 2017.

- [34] C. Xi, R. Xin, and U. A. Khan, "ADD-OPT: Accelerated distributed directed optimization," *IEEE Transactions on Automatic Control*, vol. 63, no. 5, pp. 1329–1339, 2017.
- [35] R. Xin and U. A. Khan, "A linear algorithm for optimization over directed graphs with geometric convergence," *IEEE Control Systems Letters*, vol. 2, no. 3, pp. 315–320, 2018.
- [36] G. Scutari and Y. Sun, "Distributed nonconvex constrained optimization over time-varying digraphs," *Mathematical Programming*, vol. 176, no. 1-2, pp. 497–544, 2019.
- [37] S. Pu, W. Shi, J. Xu, and A. Nedić, "Push-pull gradient methods for distributed optimization in networks," *IEEE Transactions on Automatic Control*, vol. 66, no. 1, pp. 1–16, 2020.
- [38] G. Carnevale, F. Farina, I. Notarnicola, and G. Notarstefano, "Distributed online optimization via gradient tracking with adaptive momentum," *preprint arXiv*:2009.01745, 2020.
- [39] M. Bin, I. Notarnicola, L. Marconi, and G. Notarstefano, "A system theoretical perspective to gradient-tracking algorithms for distributed quadratic optimization," in *IEEE Conference on Decision and Control* (CDC), pp. 2994–2999, 2019.
- [40] G. Carnevale, M. Bin, I. Notarnicola, L. Marconi, and G. Notarstefano, "Enhanced gradient tracking algorithms for distributed quadratic optimization via sparse gain design," *IFAC-PapersOnLine*, vol. 53, no. 2, pp. 2696–2701, 2020.
- [41] H. K. Khalil, "Nonlinear systems," Upper Saddle River, 2002.
- [42] E. D. Sontag, "Input to state stability: Basic concepts and results," in Nonlinear and optimal control theory, pp. 163–220, Springer, 2008.

APPENDIX

We first observe that, in light of Assumption 2.2, the ODE in (8) is well posed for any $t \ge 0$ and admits a unique solution, see [41, Theorem 3.2]. By inspecting system (8), we can assert that it has a unique equilibrium point at

$$\begin{bmatrix} x_{\mathsf{eq}} \\ z_{\mathsf{eq}} \end{bmatrix} := \begin{bmatrix} \mathbf{1} x^{\star} \\ -\nabla \mathbf{f}(\mathbf{1} x^{\star}) \end{bmatrix},$$

which represents the situation in which the N agents have a consensual solution estimate equal to the optimal solution x^* of the optimization problem (1). In order to use a Lyapunov approach, we put the system in error coordinates. Let

$$\begin{bmatrix} x \\ z \end{bmatrix} \longmapsto \begin{bmatrix} \tilde{x} \\ \tilde{z} \end{bmatrix} := \begin{bmatrix} x \\ z \end{bmatrix} - \begin{bmatrix} x_{\text{eq}} \\ z_{\text{eq}} \end{bmatrix}.$$
(A.19)

Then, system (8) can be rewritten as

$$\begin{bmatrix} \tilde{x} \\ \dot{\tilde{z}} \end{bmatrix} = \begin{bmatrix} -L & -I \\ 0 & -L \end{bmatrix} \begin{bmatrix} \tilde{x} \\ \tilde{z} \end{bmatrix} + \begin{bmatrix} I \\ L \end{bmatrix} u(\tilde{x}), \tag{A.20}$$

where the role played by the "input" term $u(\tilde{x}) := \nabla \mathbf{f}(\mathbf{1}x^*) - \nabla \mathbf{f}(\tilde{x}+\mathbf{1}x^*)$ has been highlighted. Indeed, it can be interpreted as a nonlinear feedback of the output $\tilde{y} = \tilde{x}$ and suggests to introduce a further change of coordinates given by

$$\begin{bmatrix} \tilde{x} \\ \tilde{z} \end{bmatrix} \longmapsto \begin{bmatrix} \tilde{y} \\ \tilde{\eta} \end{bmatrix} := \underbrace{\begin{bmatrix} I & 0 \\ L & -I \end{bmatrix}}_{T_1} \begin{bmatrix} \tilde{x} \\ \tilde{z} \end{bmatrix}.$$
(A.21)

Since T_1 is an involutory matrix (i.e., it coincides with its inverse), the change of coordinates (A.21) transforms (A.20) in

$$\begin{bmatrix} \dot{\tilde{y}} \\ \ddot{\tilde{\eta}} \end{bmatrix} = \begin{bmatrix} -2L & I \\ -L^2 & 0 \end{bmatrix} \begin{bmatrix} \tilde{y} \\ \tilde{\eta} \end{bmatrix} + \begin{bmatrix} I \\ 0 \end{bmatrix} u(\tilde{y}).$$
(A.22)

Before studying the stability of the origin for (A.22), the effect of the initialization $\mathbf{1}^{\top} z(0) = 0$ in the new coordinates $(\tilde{y}, \tilde{\eta})$ is investigated. We observe that the subspace

$$\mathcal{S} := \{ (\tilde{y}, \tilde{\eta}) \mid \mathbf{1}^{\top} \tilde{\eta} = 0 \}$$

is invariant for (A.22). In light of (A.21), it holds

$$0 = \mathbf{1}^{\top} \tilde{\eta} = \mathbf{1}^{\top} (L \tilde{x} - \tilde{z}) = \mathbf{1}^{\top} z,$$

where the last equality holds in light of (A.19) and since $\mathbf{1}^{\top} \nabla \mathbf{f}(\mathbf{1} x^{\star}) = 0$ and $\mathbf{1}^{\top} L = 0$ (cf. Assumption 2.3). Therefore the initialization of z(0) guarantees that $\tilde{\eta}(0) \in S$. Hence, we can perform a final change of coordinates to isolate the invariant state to further restrict the dynamics. Let

$$\begin{bmatrix} \tilde{y} \\ \tilde{\eta} \end{bmatrix} \longmapsto \begin{bmatrix} \tilde{y} \\ \tilde{\psi} \\ \tilde{\eta}_{\text{avg}} \end{bmatrix} := T_2 \begin{bmatrix} \tilde{y} \\ \tilde{\eta} \end{bmatrix}, \qquad (A.23)$$

in which

$$T_2 := \begin{bmatrix} T_{\tilde{y}} \\ T_{\tilde{\eta}} \end{bmatrix}, \ T_{\tilde{y}} := \begin{bmatrix} I & 0 \\ 0 & R^{\top} \end{bmatrix}, \ T_{\tilde{\eta}} := \begin{bmatrix} 0 & \frac{1}{\sqrt{N}} \mathbf{1}^{\top} \end{bmatrix},$$
(A.24)

with $R \in \mathbb{R}^{Nd \times (N-1)d}$ such that $R^{\top}R = I$, $R^{\top}\mathbf{1} = 0$ and ||R|| = 1. The following useful relations holds true

$$RR^{\top} = I - \frac{1}{N} \mathbf{1} \mathbf{1}^{\top}.$$
 (A.25)

It is easy to check that $T_2^{-1} = T_2^{\top}$, thus (A.22) can be rewritten as

$$\begin{bmatrix} \dot{\tilde{y}} \\ \dot{\tilde{\psi}} \\ \dot{\tilde{\eta}}_{\text{avg}} \end{bmatrix} = \begin{bmatrix} -2L & R & \frac{1}{\sqrt{N}} \\ -R^{\top}L^2 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} \tilde{y} \\ \tilde{\psi} \\ \tilde{\eta}_{\text{avg}} \end{bmatrix} + \begin{bmatrix} I \\ 0 \\ 0 \end{bmatrix} u(\tilde{y}). \quad (A.26)$$

In light of the invariance of S, it holds $\tilde{\eta}_{avg}(t) \equiv 0$. Then we can consider only $\zeta := \operatorname{col}(\tilde{y}, \tilde{\psi}) \in \mathbb{R}^n$, with n := (2N-1)d. The dynamics (A.26) can be written as

$$\dot{\zeta} = A\zeta + Bu(\tilde{y}), \tag{A.27a}$$

with

$$A := \begin{bmatrix} -2L & R\\ -R^{\top}L^2 & 0 \end{bmatrix}, \quad B := \begin{bmatrix} I\\ 0 \end{bmatrix}.$$
(A.28)

Next, consider a quadratic, candidate Lyapunov function V: $\mathbb{R}^n \to \mathbb{R}$ given by

$$V(\zeta) := \zeta^{\top} P \zeta, \qquad (A.29)$$

with $P \in \mathbb{R}^{n \times n}$ such that $P = P^{\top} > 0$ and arranged in blocks as

$$P := \begin{bmatrix} P_1 & P_2 \\ P_2^\top & P_3 \end{bmatrix}, \tag{A.30}$$

where $P_1 \in \mathbb{R}^{Nd \times Nd}$, $P_2 \in \mathbb{R}^{Nd \times d}$, and $P_3 \in \mathbb{R}^{(Nd-d) \times (Nd-d)}$. Next, it is shown how to choose P in order to prove global exponential stability of the origin of (A.27). Let m > 0 and set

$$P_1 = mI, \quad P_2 = -R, \quad P_3 = mR^{\top} (L^2)^{\dagger} R,$$
 (A.31)

where $(\cdot)^{\dagger}$ denotes the Moore-Penrose pseudoinverse. By the Schur complement lemma, P > 0 imposes that m must satisfy

$$\begin{cases} mI > 0\\ mR^{\top}(L^2)^{\dagger}R - \frac{1}{m}I > 0 \end{cases} \implies m > \frac{1}{\sqrt{\min\{\sigma(R^{\top}(L^2)^{\dagger}R)\}}}.$$
(A.32)

The time-derivative of V along trajectories of (A.27) is

$$\dot{V}(\zeta) = \zeta^{\top} \underbrace{(A^{\top}P + PA)}_{-Q} \zeta + 2\zeta^{\top}PBu.$$
(A.33)

The choices (A.31) yield to

$$Q = \begin{bmatrix} 4mL - 2L^2 & 2LR \\ 2R^{\top}L & 2I \end{bmatrix}, \quad PB = \begin{bmatrix} mI \\ -R^{\top} \end{bmatrix}. \quad (A.34)$$

We separately study the quadratic term $-\zeta^{\top}Q\zeta$ and the cross term $2\zeta^{\top}PBu$ as a function of m to show that $\dot{V}(\zeta)$ can be made negative definite for a sufficiently large m.

As for the first term in (A.33), we observe that Q is a solution to a Lyapunov equation associated to a marginally stable matrix. Therefore, it can only be positive semidefinite. Indeed, the upper-left block within the expression (A.34) has the kernel spanned by 1 for any choice of m. By the Schur complement lemma, imposing $Q \ge 0$ is equivalent to

$$4mL - 2L^2 - 2LRR^{\top}L \ge 0.$$
 (A.35)

In light of (A.25) and since $L\mathbf{1} = 0$, condition (A.35) reduces to $4mL - 4L^2 \ge 0$. Since L and L^2 have the same kernel, the latter condition is fulfilled by any m such that

$$m \ge \frac{\max\{\sigma(L^2)\}}{\min\{\sigma(L) \setminus \{0\}\}}.$$
(A.36)

Moreover, L positive semidefinite, condition (A.36) can be satisfied with m > 0.

Next, the second term in (A.33) is considered to show $\dot{V} < 0$. In light of (A.34), it holds

$$2\zeta^{\top} PBu = 2m\tilde{y}^{\top}u - 2\tilde{\psi}^{\top}R^{\top}u$$

$$\stackrel{(a)}{\leq} -2m\alpha \|\tilde{y}\|^{2} - 2\tilde{\psi}^{\top}R^{\top}u, \qquad (A.37)$$

where in (a) we use the strong convexity of the cost functions (cf. Assumption 2.1). Using the Cauchy-Schwarz inequality, condition (A.37) can be manipulated as

$$2\zeta^{\top} PBu \leq -2m\alpha \|\tilde{y}\|^{2} + 2 \|R\| \|\tilde{\psi}\| \|u\|$$

$$\stackrel{(a)}{\leq} -2m\alpha \|\tilde{y}\|^{2} + 2\beta \|\tilde{\psi}\| \|\tilde{y}\|$$

$$\stackrel{(b)}{\leq} -2m\alpha \|\tilde{y}\|^{2} + \frac{\beta}{\epsilon} \|\tilde{y}\|^{2} + \beta\epsilon \|\tilde{\psi}\|^{2}$$

$$\stackrel{(c)}{\equiv} \zeta^{\top} \underbrace{\left[\left(-2m\alpha + \frac{\beta}{\epsilon} \right)I \quad 0 \\ 0 \quad \beta\epsilon I \right]}_{Q_{0}} \zeta, \quad (A.38)$$

where in (a) we use the Lipschitz continuity of the gradient of the cost functions (cf. Assumption 2.2) and the fact that ||R|| = 1, while in (b) we use the Young's inequality with $\epsilon > 0$, and in (c) we introduce the matrix Q_0 . Indeed, we want to show that the zero eigenvalues of Q can be moved inside the open left-half plane through Q_0 . Thus, by plugging (A.38) in (A.33), it holds

$$\dot{V}(\zeta) \le -\zeta^{\top} \tilde{Q} \zeta,$$
 (A.39)

where $\tilde{Q} := Q - Q_0$, i.e., it holds

$$\tilde{Q} := \begin{bmatrix} 4mL - 2L^2 + \left(2m\alpha - \frac{\beta}{\epsilon}\right)I & 2LR\\ 2R^{\top}L & (2 - \beta\epsilon)I \end{bmatrix}. \quad (A.40)$$

By the Schur complement lemma, $\tilde{Q} > 0$ is equivalent to

$$\begin{cases} 2-\beta\epsilon > 0\\ 4mL - 2\left(\frac{2}{2-\beta\epsilon} + 1\right)L^2 + \frac{2m\alpha\epsilon-\beta}{\epsilon}I > 0, \end{cases}$$

which is verified for every $\epsilon < \frac{2}{\beta}$ and

$$m > \max\left\{\frac{\left(1 + \frac{1}{2 - \beta\epsilon}\right) \max\{\sigma(L^2)\}}{2\min\{\sigma(L) \setminus \{0\}\}}, \frac{\beta}{2\alpha\epsilon}\right\}.$$
 (A.41)

Therefore, we can conclude that $\dot{V}(\zeta) < -\min\{\sigma(\tilde{Q})\} \|\zeta\|^2$ which implies that the origin is globally exponentially stable for system (A.27) (cf. [41, Theorem 4.10]). Specifically, there exist $a_7, a_2 > 0$ such that

$$\|\zeta(t)\| \le a_7 \,\|\zeta(0)\| \exp(-a_2 t),\tag{A.42}$$

for any $\zeta(0) \in \mathbb{R}^n$. By noticing that $||x_i(t) - x^*|| \leq ||x(t) - \mathbf{1}x^*|| = ||y(t)|| \leq ||\zeta(t)||$, the proof follows by (A.42) by setting $a_1 = a_7 ||\zeta(0)||$.

The dynamics (11) of SYNCHRONOUS TRIGGERED GRA-DIENT TRACKING can be reformulated as a perturbed instance of the nominal dynamics of CONTINUOUS GRADI-ENT TRACKING described by (8). Clearly, the perturbation expresses the impact of the triggering mechanism on the algorithmic evolution. Thus, by adding and subtracting the term $B_2 \operatorname{col}(x(t), z(t), \nabla \mathbf{f}(x(t)))$ in the dynamics (13), we get

$$\begin{bmatrix} \dot{x} \\ \dot{z} \end{bmatrix} = \begin{bmatrix} -L & -I \\ 0 & -L \end{bmatrix} \begin{bmatrix} x \\ z \end{bmatrix} + \begin{bmatrix} -I \\ -L \end{bmatrix} \nabla \mathbf{f}(x) + B_2 e, \quad (A.43)$$

where e has the same meaning as in (17). By performing the same changes of coordinates defined in (A.19), (A.21), and (A.23), the dynamics (A.43) can be equivalently reformulated as the following (restricted) dynamics

$$\zeta = A\zeta + Bu + Ee_{\zeta,\nabla},\tag{A.44}$$

where the vectors $\zeta \in \mathbb{R}^n$, $u \in \mathbb{R}^{Nd}$ and the matrices $A \in \mathbb{R}^{n \times n}$ and $B \in \mathbb{R}^{n \times Nd}$ are as in (A.27), while the quantities associated to the perturbation are

$$\begin{split} E &:= T_{\tilde{y}}^{\top} T_1 B_2 T_1 T_{\tilde{y}} = \begin{bmatrix} -L & 0 & 0\\ 0 & -R^{\top} L R & R^{\top} L \end{bmatrix}, \\ (A.45a) \\ e_{\zeta,\nabla} &:= \operatorname{col}(\hat{\tilde{y}} - \tilde{y}, \hat{\tilde{\psi}} - \tilde{\psi}, e_{\nabla}) \end{split}$$

$$:= \operatorname{col}(\hat{\tilde{y}} - \tilde{y}, \hat{\tilde{\psi}} - \tilde{\psi}, \nabla \mathbf{f}^{k} - \nabla \mathbf{f}(\tilde{y} + x^{\star})) \quad (A.45b)$$

with T_1 and $T_{\tilde{y}}$ defined in (A.21) and (A.24), respectively. Let us consider a quadratic, candidate Lyapunov function $V(\zeta) = \zeta^{\top} P \zeta$ as in (A.29) with the blocks of P set as in (A.31). The time-derivative of V along the trajectories of (A.43) satisfies

$$\dot{V}(\zeta) = \zeta^{\top} (A^{\top}P + PA)\zeta + 2\zeta^{\top} PBu + 2\zeta^{\top} PEe_{\zeta,\nabla}$$

$$\leq -\zeta^{\top} \tilde{Q}\zeta + 2\zeta^{\top} PEe_{\zeta,\nabla}, \qquad (A.46)$$

where \hat{Q} is as in (A.40) so that the inequality holds in light the previous proof of Theorem 3.1 (cf. (A.39)). By using the Young's inequality with $\epsilon > 0$, we can further upper bound (A.46) as

$$\dot{V}(\zeta) \leq -\zeta^{\top} \tilde{Q}\zeta + \epsilon \zeta^{\top} PP\zeta + \frac{1}{\epsilon} e_{\zeta,\nabla}^{\top} E^{\top} Ee_{\zeta,\nabla}$$
$$\stackrel{(a)}{=} -\zeta^{\top} \left(\tilde{Q} - \epsilon P^2 \right) \zeta + \frac{1}{\epsilon} e_{\zeta,\nabla}^{\top} E^{\top} Ee_{\zeta,\nabla}, \qquad (A.47)$$

where in (a) the terms in ζ have been grouped. In light of the sufficient condition in (A.41) to get a positive definite \tilde{Q} , we can always take ϵ such that

$$0 < \epsilon < \frac{\min\{\sigma(Q)\}}{\max\{\sigma(P^2)\}}$$

in order to impose also $\tilde{Q} - \epsilon P^2$ positive definite. Thus, by denoting as q > 0 the smallest eigenvalue of the matrix $\tilde{Q} - \epsilon P^2$ and by applying the Cauchy-Schwarz inequality to the quadratic term in $e_{\zeta,\nabla}$ of (A.47), we bound (A.47) as

$$\dot{V}(\zeta) \leq -q \|\zeta\|^{2} + \frac{1}{\epsilon} \|E^{\top}E\| \|e_{\zeta,\nabla}\|^{2}
\stackrel{(a)}{=} -q \|\zeta\|^{2} + \frac{1}{\epsilon} \|E^{\top}E\| (\|e_{\zeta}\|^{2} + \|e_{\nabla}\|^{2})
\stackrel{(b)}{\leq} -q \|\zeta\|^{2} + \frac{1}{\epsilon} \|E^{\top}E\| (\|e_{\zeta}\|^{2} + \beta^{2} \|\hat{\tilde{y}} - \tilde{y}\|^{2})
\stackrel{(c)}{\leq} -q \|\zeta\|^{2} + \underbrace{\frac{1}{\epsilon} \|E^{\top}E\| (1 + \beta^{2})}_{c_{1}} \|e_{\zeta}\|^{2}, \quad (A.48)$$

where in (a) we introduce $e_{\zeta} := \operatorname{col}(\hat{\hat{y}} - \tilde{y}, \tilde{\psi} - \tilde{\psi})$ to write $\|e_{\zeta,\nabla}\|^2 = \|e_{\zeta}\|^2 + \|e_{\nabla}\|^2$, in (b) we use the Lipschitz continuity of the gradients of the cost functions (cf. Assumption 2.2) to bound $\|e_{\nabla}\|^2 \leq \beta^2 \|\hat{\hat{y}} - \tilde{y}\|^2$, and in (c) we rely on the fact that $\hat{\hat{y}} - \tilde{y}$ is a component of e_{ζ} .

The proof continues by deriving an upper bound for $||e_{\zeta}||^2$ in (A.48). We start by defining

$$r := \frac{\|e_{\zeta}\|}{\|\zeta\|}.\tag{A.49}$$

Moreover, recall that in each interval $[\tilde{t}^k, \tilde{t}^{k+1})$, the error e_{ζ} is set to zero at \tilde{t}^k and grows until \tilde{t}^{k+1} when it is reset again to zero. Hence, the goal is to establish a lower bound on the needed time for r(t) to reach $\sqrt{q/c_1}$. By computing the time derivative of (A.49), it follows

$$\dot{r} = \frac{e_{\zeta}^{\top} \dot{e}_{\zeta}}{\|e_{\zeta}\| \|\zeta\|} - \frac{\|e_{\zeta}\| \zeta^{\top} \dot{\zeta}}{\|\zeta\|^3}.$$
(A.50)

Using the Cauchy-Schwarz inequality, we bound \dot{r} as

$$\dot{r} \leq \frac{\|e_{\zeta}\| \|\dot{e}_{\zeta}\|}{\|e_{\zeta}\| \|\zeta\|} + \frac{\|e_{\zeta}\| \|\zeta\|}{\|\zeta\|^{3}} \leq \frac{\|\dot{\zeta}\|}{\|\zeta\|} + \frac{\|e_{\zeta}\| \|\dot{\zeta}\|}{\|\zeta\|^{2}} \stackrel{(b)}{=} (1+r) \frac{\|\dot{\zeta}\|}{\|\zeta\|}$$
(A.51)

where in (a) we use the identity $\dot{e}_{\zeta} = -\dot{\zeta}$ while in (b) we exploit the definition of r in (A.49). Then, in light of the dynamics of ζ in (A.44), it holds

$$\dot{r} \leq (1+r) \frac{\|A\zeta + Bu + Ee_{\zeta,\nabla}\|}{\|\zeta\|} \\ \stackrel{(a)}{\leq} (1+r) \frac{\|A\| \|\zeta\| + \|u\| + \|E\| \|e_{\zeta}\| + \|E\| \|e_{\nabla}\|}{\|\zeta\|},$$
(A.52)

where in (a) we use the triangle and the Cauchy-Schwarz inequalities combined with ||Bu|| = ||u||. Next, by using the Lipschitz continuity of the gradients of the cost functions (cf. Assumption 2.2), we have $||u|| \le \beta ||\zeta||$ and $||e_{\nabla}|| \le \beta ||e_{\zeta}||$. Thus, (A.52) becomes

$$\begin{split} \dot{r} &\leq (1+r) \frac{(\|A\|+\beta) \|\zeta\| + (1+\beta) \|E\| \|e_{\zeta}\|}{\|\zeta\|} \\ &\stackrel{(a)}{=} (1+r) \frac{\beta \|\zeta\|}{\|\zeta\|} + (1+r) \frac{\|A\| \|\zeta\| + (1+\beta) \|E\| \|e_{\zeta}\|}{\|\zeta\|} \\ &\stackrel{(b)}{=} \beta (1+r) + (1+r) \frac{c_2 \|\zeta\| + c_2 \|e_{\zeta}\|}{\|\zeta\|} \\ &\stackrel{(c)}{=} \beta (1+r) + c_2 (1+r)^2, \end{split}$$
(A.53)

where in (a) we simply rearrange the terms, in (b) we introduce $c_2 := \max\{||A||, (1 + \beta) ||E||\}$, and in (c) we use the definition of r in (A.49).

Using the Comparison Lemma (see [41, Lemma 3.4]) the bound (A.53) translates in the following inequality

$$r(t, r(\tilde{t}^k)) \le \bar{r}(t, \bar{r}(\tilde{t}^k)), \tag{A.54}$$

where $r(t, r(\tilde{t}^k))$ denotes the solution of (A.50) with initial condition at $t = t_k$ given by $r(t_k)$ while $\bar{r}(t, \bar{r}(\tilde{t}^k))$ denotes the solution of

$$\dot{\bar{r}}(t) = \beta (1 + \bar{r}(t)) + c_2 (1 + \bar{r}(t))^2,$$
 (A.55)

for some initial condition initial condition at $t = \tilde{t}^k$ given by $\bar{r}(\tilde{t}^k)$ such that $r(\tilde{t}^k) \leq \bar{r}(\tilde{t}^k)$. Recalling that the protocol (12) imposes $r(\tilde{t}^k) = 0$ at the beginning of each time interval $[\tilde{t}^k, \tilde{t}^{k+1})$, then we select $\bar{r}(\tilde{t}^k) = 0$. The solution of (A.55) can be shown to be (cf. [18])

$$\bar{r}(t,0) = \frac{(\beta + c_2)(\exp(\beta(t - \tilde{t}^k)) - 1)}{-c_2 \exp(\beta(t - \tilde{t}^k)) + \beta + c_2}.$$
 (A.56)

Notice that $\bar{r}(t,0)$ starts from 0 at $t = \tilde{t}^k$ and monotonically increases within the interval $\left[0, t_k + \ln\left(\frac{\beta+c_2}{c_2}\right)/\beta\right)$. Thus, we can always find a triggering value $t = \Delta^* > 0$ such that $\bar{r}(\Delta^*, 0) = \sqrt{q/c_1}$. Hence, by choosing any $\Delta \in (0, \Delta^*)$ in (12), the inequality (A.54) ensures

$$|r(t)| = r(t) < \sqrt{\frac{q}{c_1}},$$
 (A.57)

for all $t \in [\tilde{t}^k, \tilde{t}^{k+1})$, where the equality holds because r is always positive, see its definition in (A.49). With this result in mind, the inequality (A.48) can be rewritten as

$$\dot{V}(\zeta) \leq -\left(q - \frac{|r|^2}{c_1}\right) \left\|\zeta\right\|^2,$$

which allows us to use (A.57) to conclude that the origin is globally exponentially stable for system (A.44) (cf. [41, Th. 4.10]). Specifically, there exist $a_4, a_8 > 0$ such that

$$\|\zeta(t)\| \le a_8 \|\zeta(0)\| \exp(-a_4 t),$$
 (A.58)

for any $\zeta(0) \in \mathbb{R}^n$. By noticing that $||x_i(t) - x^*|| \leq ||x(t) - \mathbf{1}x^*|| = ||y(t)|| \leq ||\zeta(t)||$, the proof follows by (A.58) by setting $a_3 = a_8 ||\zeta(0)||$.

The proof of Theorem 4.2 traces the same initial steps in Section A. Specifically, we reformulate the ASYNCHRONOUS TRIGGERED GRADIENT TRACKING as a perturbed, extended version of CONTINUOUS GRADIENT TRACKING in which the perturbation is due to the event-triggered communication. By exploiting the steps leading to (A.44), the aggregate form of (11) and (16) reads

$$\dot{\zeta} = A\zeta + Bu + De$$
 (A.59a)

$$\dot{\xi} = -\nu\xi,$$
 (A.59b)

where the vectors $\zeta \in \mathbb{R}^n$, $u \in \mathbb{R}^{Nd}$ and the matrices $A \in \mathbb{R}^{n \times n}$ and $B \in \mathbb{R}^{n \times Nd}$ are as in (A.27), $e \in \mathbb{R}^{3Nd}$ has the same meaning as in (17), $\xi := \operatorname{col}(\xi_1, \ldots, \xi_N) \in \mathbb{R}^N$, while the matrix D is given by

$$D := T_{\tilde{y}}^{\top} T_1 B_2 = \begin{bmatrix} -L & 0 & 0\\ -R^{\top} L^2 & R^{\top} L & R^{\top} L \end{bmatrix}, \quad (A.60)$$

where the matrices T_1 , $T_{\tilde{y}}$ and B_2 are as in (A.21), (A.24), and (13), respectively. We underline that the dynamics of ζ and ξ are decoupled while both quantities affect the triggering law (15).

Next, we show how to properly choose the value for ν in (16) and for λ in the triggering law (15) to guarantee that the perturbation term De and the auxiliary variable ξ do not alter the stability property associated the nominal system $\dot{\zeta} = A\zeta + Bu$ (cf. Theorem 3.1). To this end, an upper bound for ||De||, proportional to $||\zeta||$ and $||\xi||$, is derived. We start by using the Cauchy-Schwarz inequality to write $||De|| \leq ||D|| ||e|| \leq c_3 \sum_{i=1}^{N} ||e_i||$, with $c_3 := ||D||$. In light of the triggering law (15), the latter inequality can be upper bounded as

$$\|De\| \leq \lambda c_{3} \sum_{i=1}^{N} \|z_{i} + \nabla f_{i}(x_{i})\| + c_{3} \sum_{i=1}^{N} |\xi_{i}|$$

$$\stackrel{(a)}{\leq} \lambda c_{3} \sqrt{N} \|z + \nabla \mathbf{f}(x)\| + c_{3} \sqrt{N} \|\xi\|$$

$$\stackrel{(b)}{=} \lambda c_{4} \|\tilde{z} + \nabla \mathbf{f}(\tilde{x} + \mathbf{1}x^{*}) - \nabla \mathbf{f}(\mathbf{1}x^{*})\| + c_{4} \|\xi\|$$

$$\stackrel{(c)}{\leq} \lambda c_{4} \|\tilde{z}\| + \lambda c_{4}\beta \|\tilde{x}\| + c_{4} \|\xi\|, \quad (A.61)$$

where in (a) we apply the basic algebraic relation $\sum_{i=1}^{N} \|\theta_i\| \leq \sqrt{N} \|\theta\|$ for a vector $\theta = \operatorname{col}(\theta_1, \ldots, \theta_N)$, in (b) we perform the change of coordinates given in (A.19) and introduce the constant $c_4 := c_3\sqrt{N}$, and in (c) we use the triangle inequality and the Lipschitz continuity of the gradients of the cost functions (cf. Assumption 2.2). According to (A.21) and (A.23), it holds

$$\begin{bmatrix} \tilde{x} \\ \tilde{z} \end{bmatrix} = T_1 T_2^\top \begin{bmatrix} \zeta \\ \tilde{\eta}_{\text{avg}} \end{bmatrix} = T_1 T_2^\top \begin{bmatrix} \zeta \\ 0 \end{bmatrix}, \qquad (A.62)$$

where we use the fact that the initialization z(0) leads to $\tilde{\eta}_{avg}(t) \equiv 0$. We rearrange the inequality (A.61) to reconstruct the term $\|col(\tilde{x}, \tilde{z})\|$ as

$$\|De\| \leq \lambda c_4 \max\{1, \beta\} \sqrt{2} \|\operatorname{col}(\tilde{x}, \tilde{z})\| + c_4 \|\xi\|$$

$$\stackrel{(a)}{\leq} \lambda c_5 \|\zeta\| + c_4 \|\xi\|, \qquad (A.63)$$

where in (a) we combine (A.62) with the Cauchy-Schwarz inequality and set $c_5 := c_4 \max\{1, \beta\}\sqrt{2} ||T_1T_2^\top||$. Given the linear bound in (A.63), we can pursue a Lyapunov approach to conclude the global exponential stability of the origin. Let us consider a quadratic, candidate Lyapunov function $\tilde{V}(\zeta, \xi) = \zeta^\top P \zeta + \frac{1}{2} \xi^\top \xi$. derived from the one considered in (A.29) with the blocks of P set as in (A.31). Using similar arguments leading to (A.48), the time-derivative of \tilde{V} along trajectories of (A.59) can be upper bounded as

$$\dot{\tilde{V}}(\zeta,\xi) \le -\tilde{q} \|\zeta\|^2 + 2\zeta^{\top} P D e - \nu \|\xi\|^2$$
. (A.64)

By using the Cauchy-Schwarz inequality, we can plug (A.63) in (A.64) to obtain

$$\tilde{V}(\zeta,\xi) \leq -\tilde{q} \|\zeta\|^{2} + 2 \|\zeta\| \|P\| \|De\|
\leq -\tilde{q}_{\lambda} \|\zeta\|^{2} + 2c_{4} \|P\| \|\zeta\| \|\xi\| - \nu \|\xi\|^{2}, \quad (A.65)$$

where we introduce $\tilde{q}_{\lambda} := (\tilde{q} - 2\lambda c_5 ||P||)$. Then, for any $\lambda < \frac{\tilde{q}}{2c_5} ||P|| =: \lambda^*$, it holds $\tilde{q}_{\lambda} > 0$. Setting $c_6 := c_4 ||P||$, the inequality (A.65) can be arranged in a matrix form as

$$\dot{\tilde{V}}(\zeta,\xi) \leq -\begin{bmatrix} \|\zeta\| \\ \|\xi\| \end{bmatrix}^{\perp} \underbrace{\begin{bmatrix} \tilde{q}_{\lambda} & -c_6 \\ -c_6 & \nu \end{bmatrix}}_{U} \begin{bmatrix} \|\zeta\| \\ \|\xi\| \end{bmatrix}.$$
(A.66)

Being $U \in \mathbb{R}^{2 \times 2}$ symmetric, by the Sylvester criterion U > 0if and only if $\tilde{q}_{\lambda}\nu > c_6^2$. Therefore, by taking any $\nu > \nu^* := \frac{c_6^2}{q_{\lambda}}$, the matrix U is positive definite. Thus the inequality (A.66) guarantees that the origin is globally exponentially stable for system (A.59) (cf. [41, Lemma 4.10]). Specifically, there exist $a_6, a_9 > 0$ such that

$$\|\operatorname{col}(\zeta(t),\xi(t))\| \le \underbrace{a_9 \|\operatorname{col}(\zeta(0),\xi(0))\|}_{a_5} \exp(-a_6 t), \quad (A.67)$$

for any $col(\zeta(0), \xi(0)) \in \mathbb{R}^{N+n}$. By noticing that

$$||x_i(t) - x^*|| \le ||x(t) - \mathbf{1}x^*|| = ||y(t)|| \le ||\operatorname{col}(\zeta(t), \xi(t))||,$$

the proof of the first part of the theorem follows by (A.67).

Next, we prove by contradction that (11) does not exhibit the Zeno behavior. Suppose, without loss of generality, that an agent *i* exhibits the Zeno behavior, namely

$$\lim_{k_i \to \infty} t_i^{k_i} = t_i^{\infty}.$$
 (A.68)

For any $t \ge 0$, we have

$$\frac{d}{dt} \|e_i(t)\| = \frac{e_i^\top \dot{e}_i}{\|e_i(t)\|} \stackrel{(a)}{\leq} \|\dot{e}_i(t)\| \\
\stackrel{(b)}{=} \|\operatorname{col}(\dot{x}_i(t), \dot{z}_i(t), \nabla^2 f_i(x_i(t)) \dot{x}_i(t))\| \\
\stackrel{(c)}{=} \|\operatorname{col}(\dot{\tilde{x}}_i(t), \dot{\tilde{z}}_i(t), \nabla^2 f_i(\tilde{x}_i(t) + x^*) \dot{\tilde{x}}_i(t))\|, \quad (A.69)$$

where in (a) we use the Cauchy-Schwarz inequality, in (b) we use the definition of $e_i(t)$, and in (c) we locally perform the change of variables given in (A.19). Combining the latter change of variables with (11), it holds

$$\dot{\tilde{x}}_{i}(t) = -\sum_{j \in \mathcal{N}_{i}} w_{ij}(\hat{\tilde{x}}_{i}^{k} - \hat{\tilde{x}}_{j}^{k}) - \tilde{z}_{i}(t) + u_{i}(\tilde{x}_{i}(t))$$
(A.70a)

$$\dot{\tilde{z}}_i(t) = -\sum_{j \in \mathcal{N}_i} w_{ij} (\hat{\tilde{z}}_i^k - \hat{\tilde{z}}_j^k) - \sum_{j \in \mathcal{N}_i} w_{ij} (\nabla f_i^k - \nabla f_j^k), \quad (A.70b)$$

where we use $u_i(\tilde{x}_i(t)) := (\nabla f_i(\tilde{x}_i(t) + x^*) - \nabla f_i(x^*))$ and the local components of the shorthands given in (10). By (A.67), the variables $\tilde{x}_i(t)$ and $\tilde{z}_i(t)$ are bounded for all $i \in \{1, \ldots, N\}$ and $t \ge 0$. Then, by defining $c_7 := \max_{i,t} ||\tilde{x}_i(t)||$ and $c_8 := \max_{i,t} ||\tilde{z}_i(t)||$, (A.70a) and the triangle inequality can be combined to get

$$\|\dot{\tilde{x}}_{i}(t)\| \leq \sum_{j \in \mathcal{N}_{i}} w_{ij} 2c_{7} + c_{8} + \|u_{i}(\tilde{x}_{i}(t))\| \stackrel{(a)}{\leq} (2c_{9} + \beta)c_{7} + c_{8},$$

where in (a) we introduce $c_9 := \sum_{j \in \mathcal{N}_i} w_{ij}$ and we use the Lipschitz continuity of the gradients of the cost functions (cf. Assumption 2.2). Using again the boundedness of the quantities, and by adding and subtracting $\nabla f_i(x^*)$ within the second sum of (A.70b), it holds

$$\|\tilde{z}_i(t)\| \leq 2c_9(c_8 + \beta c_7).$$

Moreover, the Lipschitz continuity of the gradients of the cost functions (cf. Assumption 2.2) also ensures that $\|\nabla^2 f_i(v)\| \leq \beta$, for all $v \in \mathbb{R}^d$ and all $i \in \{1, \ldots, N\}$. By combining the latter with the two previous equations, the inequality (A.69) can be upper bounded as

$$\frac{d}{dt} \|e_i(t)\| \le c_{10},$$
 (A.71)

with $c_{10} := (1+\beta)(2c_9+\beta)c_7+c_8+2c_9(c_8+\beta c_7).$

Since the protocol (15) imposes $e_i(t) = 0$ at the beginning of each time interval $[t_i^{k_i}, t_i^{k_i+1})$, then by also using (A.71), we can write

$$e_i(t) = e_i(t_i^{k_i}) + \int_{t_i^{k_i}}^t \frac{d \|e_i(\tau)\|}{d\tau} d\tau \le c_{10}(t - t_i^{k_i}).$$
 (A.72)

By (16), it holds $\xi_i(t) = \xi_i(0) \exp(-\nu t)$ for all $t \ge 0$. Thus, being $\lambda ||h_i(t)|| \ge 0$ for any $t \ge 0$, the bound in (A.72) imposes, as a necessary condition to satisfy the triggering in (15), that

$$c_{10}(t_i^{k_i+1} - t_i^{k_i}) \ge |\xi_i(0)| \exp(-\nu t_i^{k_i+1})$$
(A.73)

From (A.68), for all $\epsilon > 0$ there exists $k_{i,\epsilon} \in \mathbb{N}$ such that

$$t_i^{k_i} \in [t_i^\infty - \epsilon, t_i^\infty], \quad \forall k_i \ge k_{i,\epsilon}.$$
 (A.74)

Set

$$\epsilon := \frac{|\xi_i(0)|}{2c_{10}} \exp(-\nu t_i^{\infty}), \tag{A.75}$$

and suppose that the $k_{i,\epsilon}$ -th triggering time of agent *i*, namely $t_i^{k_{i,\epsilon}}$, has occurred. Let $t_i^{k_{i,\epsilon}+1}$ be the next triggering time determined by (15). Then, using the necessary condition (A.73) we can write

$$t_{i}^{k_{i,\epsilon}+1} - t_{i}^{k_{i,\epsilon}} \geq \frac{|\xi_{i}(0)|}{c_{10}} \exp(-\nu t_{i}^{k_{i,\epsilon}+1})$$

$$\stackrel{(a)}{\geq} \frac{|\xi_{i}(0)|}{c_{10}} \exp(-\nu t_{i}^{\infty}) \stackrel{(b)}{=} 2\epsilon, \quad (A.76)$$

where in (a) we use $t_i^{\infty} \ge t_i^{k_{i,\epsilon}+1}$, while in (b) we use (A.75). However (A.76) implies

$$t_i^{k_{i,\epsilon}} \le t_i^{k_{i,\epsilon}+1} - 2\epsilon \le t_i^\infty - 2\epsilon,$$

which contradicts (A.74) and concludes the proof.

The proof of Proposition 5.1 traces the same initial steps in Section A and Section A. Using the change of coordinates in (A.19), (A.21), (A.23), system (17) can be recast as

$$\dot{\zeta} = A\zeta + Bu + \delta_1 E e_{\zeta,\nabla} + T_{\tilde{y}}^\top T_1 B_3 v_{xz\nabla}, \qquad (A.77)$$

with $\zeta \in \mathbb{R}^n$, $u \in \mathbb{R}^{Nd}$, where $A \in \mathbb{R}^{n \times n}$ and $B \in \mathbb{R}^{n \times Nd}$ are as in (A.27), E and $e_{\zeta,\nabla}$ are as in (A.45a) and (A.45b), B_3 , $T_{\tilde{y}}$, and T_1 are as in (18), (A.21), and (A.24), while $v_{xz\nabla} := \operatorname{col}(v_x, v_z, v_{\nabla})$. We remark that $e_{\zeta,\nabla}$ changes according to the implemented communication protocol. Moreover, when ASYNCHRONOUS TRIGGERED GRADIENT TRACKING is considered, also dynamics (16) has to be taken into account. However, when $v_{\nabla} \equiv v_{xz} \equiv 0$, then $v_{xz\nabla} \equiv 0$ and system (A.77) reduces to

$$\dot{\zeta} = A\zeta + Bu + \delta_1 Ee_{\zeta,\nabla}.\tag{A.78}$$

Theorems 3.1, 4.1, and 4.2 ensure that the origin is globally exponentially stable for (A.78) for both $\delta_1, \delta_2 \in \{0, 1\}$ and for both communication protocols (12) and (15). In light of [41, Lemma 4.6], this condition is sufficient to assert that system (A.77) is input-to-state stable and the proof follows (cf. [42, Section 2.9]).