Program in C for studying characteristic properties of two-body interactions in the framework of spectral distribution theory

K. D. Launey, S. Sarbadhicary, T. Dytrych, and J. P. Draayer Department of Physics and Astronomy, Louisiana State University, Baton Rouge, Louisiana 70803, USA

Abstract

We present a program in C that employs spectral distribution theory for studies of characteristic properties of a many-particle quantum-mechanical system and the underlying few-body interaction. In particular, the program focuses on twobody nuclear interactions given in a JT-coupled harmonic oscillator basis and calculates correlation coefficients, a measure of similarity of any two interactions, as well as Hilbert-Schmidt norms specifying interaction strengths. An important feature of the program is its ability to identify the monopole part (centroid) of a 2-body interaction, as well as its 'density-dependent' one-body and twobody part, thereby providing key information on the evolution of shell gaps and binding energies for larger nuclear systems. As additional features, we provide statistical measures for 'density-dependent' interactions, as well as a mechanism to express an interaction in terms of two other interactions. This, in turn, allows one to identify, e.g., established features of the nuclear interaction (such as pairing correlations) within a general Hamiltonian. The program handles the radial degeneracy for 'density-dependent' one-body interactions and together with an efficient linked list data structure, facilitates studies of nuclear interactions in large model spaces that go beyond valence-shell applications.

Keywords: Spectral distribution theory, Similarity of interactions, Properties of realistic and schematic interactions, Monopole interaction

PROGRAM SUMMARY

Program title: sdt

Catalogue identifier: AEQG_v1_0

Program summary URL: http://cpc.cs.qub.ac.uk/summaries/AEQG_v1_0.html

Program obtainable from: CPC Program Library, Queens University,

Belfast, N. Ireland

Licensing provisions: Standard CPC licence,

http://cpc.cs.qub.ac.uk/licence/licence.html

No. of lines in distributed program, including test data, etc.: 10 888 No. of bytes in distributed program, including test data, etc.: 88 778

Distribution format: tar.gz Programming language: C.

Computer: Laptop, Workstation.

Operating system: Linux [tested on Linux (Kernel 2.6.9) with a gcc, version 3.4.6]

RAM: Less than 10 MB Classification: 17.15.

Nature of problem: The program calculates second-order energy moments, such as variances and correlation coefficients, widely used as measures of the overall strength of an interaction and its similarity to other interactions. It allows for studies of the physical properties of various interactions and their effect on many-particle systems.

Solution method: Calculations are based on spectral distribution theory and invoke statistical measures provided by the theory.

Running time: Less than 20 min (typically, several seconds) using a $1.80~\mathrm{GHz}$ processor.

References:

J. B. French and K. F. Ratcliff, *Phys. Rev.* C 3, 94 (1971);

F. S. Chang, J. B. French, and T. H. Thio, Ann. Phys. (N.Y.) 66, 137 (1971);

K. T. Hecht and J. P. Draayer, Nucl. Phys. A223, 285 (1974);

K. D. Sviratcheva, J. P. Draayer, and J. P. Vary, Nucl. Phys. A 786, 31 (2007).

1. Introduction

Spectral distribution theory (SDT) [1-4] originated as a complement to conventional configuration-interaction spectroscopy (for a review, see [5]). The efficacy of the theory stems from the finding that low-order energy moments of a microscopic interaction typically capture the dominant characteristic features of a many-particle system. While not the primary focus of SDT, convergence of the theory to exact eigensolution results improves, in principle, as higher-order energy moments are taken into account, and/or in the limit of the many particles occupying a much larger available single-particle (s.p.) space. The theory also provides a relatively simple means to calculate important average properties, such as level densities, the degree of symmetry violation, and various other important features (e.g., [6-16]). In particular, in nuclear physics, the SDT approach has been successfully applied to studies of energy spectra and reactions for p-, sd-, and fp-shell nuclei [17–22], as well as for understanding dominant features and differences among sd-shell realistic effective interactions [23, 24]. This has been Preprint submitted to Elsevier March 6, 2022 achieved without the need for carrying out large-scale shell-model calculations, which are not always feasible. Recent applications include explorations on quantum chaos, nuclear structure, and parity/time-reversal violation (for example, see [25–32, 5]).

We focus on second-order energy moments, such as variances and correlation coefficients, widely used as measures of the overall strength of an interaction and its similarity to other interactions. Furthermore, these moments can be propagated straightforwardly beyond the defining two-nucleon system to derivative systems with larger numbers of nucleons [3] and higher values of isospin [4]. As a result, SDT allows one to gain further insight into the physical properties of various interactions and above all, into their effect on many-particle nuclear systems. For example, correlation coefficients can be used to extract important information on how well pairing/rotational features develop in a nucleus given a specific interaction [33].

In addition, SDT gives an exact and simple prescription for identifying 'density-dependent' monopole (centroid), one-body (induced single-particle energies), and its residual, irreducible two-body parts. The theory is readily extensible to 3-body interactions and beyond and can be of special interest when such interactions are invoked, e.g., [34–36]. Hence, SDT framework provides important information on the evolution of the shell structure, shell gaps, binding energies and contribution of 3-body forces with increasing number of particles [37–39].

In this paper, we present a detailed explanation of a computer code in C that utilizes SDT. The program is applicable to microscopic two-body nuclear interactions, regardless of whether they are realistic (like N³LO [40], Argonne 18 [41], and CD-Bonn [42]) or schematic (like pairing interaction), given in the JTcoupled basis of the harmonic oscillator (HO) potential or an arbitrary spherically symmetric potential. The program, as implemented, can be straightforwardly modified to accommodate other coupling schemes (e.g., uncoupled, m-scheme, basis [39]). The computer program is also extensible to 3-body interactions and beyond (see, e.g., Ref. [39] for derivations in the m scheme), and, in principle, can be expanded to include higher-order energy moments. The program introduces a new feature, namely, it handles the radial degeneracy of the 'density-dependent' one-body interactions and together with an efficient linked-list data structure, facilitates studies in large model spaces that go beyond valence-shell or two-shell applications of earlier Fortran programs [9, 43, 44]. We also provide additional features that allow for studies of 'density-dependent' interactions, as well as of interactions that can be expressed in terms of two other interactions that are mutually orthogonal. This, in turn, can provide, for example, a key indication regarding how well a realistic nucleon-nucleon interaction may or may not reproduce – without actually employing shell-model calculations – prominent features of nuclei, such as pairing gaps in nuclear energy spectra or enhanced electric quadrupole transitions in collective rotational bands. The code is an essential computational tool that we have utilized in recent explorations to provide important nuclear structure information [32, 33, 39] and can enable further studies of interest to nuclear theory (e.g., as suggested in [45]).

2. Spectral Distribution Theory – notation and definitions

Spectral distribution theory (or statistical spectroscopy) is well documented in the literature [2, 3, 6, 4, 9] and is accompanied by early computational codes [9, 43] for evaluating various measures. In this section, we specify the notation we use as well as present the formulae used in the program to calculate the measures by following the appendix of Ref. [32]. In addition, we include the radial degeneracy concept [3] to accommodate many HO major shells, together with a discussion on orthogonalizing an interaction with respect to a reference Hamiltonian and on expanding a Hamiltonian in terms of two other Hamiltonians.

In a standard second quantized form, a one- plus two-body isospin-conserving Hamiltonian is given in terms of fermion creation $a^{\dagger}_{\eta j m(1/2)\sigma}$ and annihilation $\tilde{a}_{\eta j - m(1/2) - \sigma} = (-1)^{j - m + 1/2 - \sigma} a_{\eta j m(1/2)\sigma}$ tensors, where the operator a^{\dagger} (a) creates (annihilates) a particle of type $\sigma = \pm 1/2$ (proton/neutron) in a state of oscillator quantum number η ($\eta = 0, 1, 2, \ldots$ for s, p, sd, \ldots oscillator shells, respectively), total angular momentum j (half integer) with projection m in a finite space $2\Omega = \Sigma_j(2j+1)$,

$$H = \sum_{r} \sqrt{[r]} \varepsilon_r \{a_r^{\dagger} \otimes \tilde{a}_r\}^{(00)}$$
$$-\frac{1}{4} \sum_{retu\Gamma} \sqrt{(1+\delta_{rs})(1+\delta_{tu})[\Gamma]} W_{rstu}^{\Gamma} \{\{a_r^{\dagger} \otimes a_s^{\dagger}\}^{\Gamma} \otimes \{\tilde{a}_t \otimes \tilde{a}_u\}^{\Gamma}\}^{(00)}, (1)$$

where the labels are $r = \{\eta_r, j_r, \tau_r = \frac{1}{2}\}$, $[r] = 2(2j_r + 1)$, and $[\Gamma] = (2J + 1)(2T + 1)$. In Eq. (1), ε_r is an (external) single-particle energy and $W^{JT}_{rstu} = \langle rsJTMT_0|H|tuJTMT_0\rangle$ is a two-body antisymmetric matrix element for JT-coupled normalized basis states, $|tuJTMT_0\rangle = \frac{1}{\sqrt{1+\delta_{tu}}}\{a_t^{\dagger} \otimes a_u^{\dagger}\}^{JTMT_0}|0\rangle$, with

$$W_{rstu}^{\Gamma} = -(-)^{r+s-\Gamma}W_{srtu}^{\Gamma} = -(-)^{t+u-\Gamma}W_{rsut}^{\Gamma} = (-)^{r+s-t-u}W_{srut}^{\Gamma} = W_{turs}^{\Gamma}. \quad (2)$$

The underlying principle of SDT is the mapping of Hamiltonians onto a multidimensional linear vector space with an inner product. That is, a Hamiltonian Hcan be realized as a vector with coordinates specified by the independent matrix elements of H. For the one- plus two-body H, this space is spanned by a complete

¹Equivalently, the isospin-conserving part of a charge-dependent interaction (see Sec. 3).

set of unit tensors (basis), $\{a_r^{\dagger} \otimes \tilde{a}_r\}^{(00)}$ and $\{\{a_r^{\dagger} \otimes a_s^{\dagger}\}^{\Gamma} \otimes \{\tilde{a}_t \otimes \tilde{a}_u\}^{\Gamma}\}^{(00)}$, and all such Hamiltonians expand along this basis according to Eq. (1). In this multidimensional space, an inner product, (H, H'), for a pair of Hamiltonians H and H', is introduced (defined below), which, in turn, defines important measures; namely, a norm of H, $\sigma_H = \sqrt{(H, H)}$ ("size" of a Hamiltonian), an angle between two Hamiltonians, $\cos \theta_{H,H'} = (H, H')/\sigma_H \sigma_{H'}$ ("orientation" of one Hamiltonian relative to the other), and a metric ("distance"), $d(H, H') = \sigma_{(H-H')}$. For example, for the 3-dimensional coordinate space, an inner product is introduced by the scalar product of two vectors, $(\mathbf{x}, \mathbf{x}') = \sum_{i=1}^3 x_i x_i' = \text{Tr}(\mathbf{x}\mathbf{x}')$, which defines $\sigma_{\mathbf{x}} = \sqrt{x_1^2 + x_2^2 + x_3^2}$, the length of a vector, and a metric $d(\mathbf{x}, \mathbf{x}') = \sqrt{\sum_{i=1}^3 (x_i - x_i')^2}$, the distance between two points in space. Note that these measures are independent of the basis representation, but are properties of the vectors.

The inner product in SDT is defined in terms of traceless operators, $H - \langle H \rangle^{(\alpha)}$, where

$$\langle H \rangle^{(\alpha)} = \frac{1}{\mathcal{N}_d} \text{Tr}^{(\alpha)} H$$
 (3)

is the expectation value of H averaged over the subset of basis states α (distributions) with dimensionality \mathcal{N}_d and thus, is related to the trace of the operator. E.g., for the matrix representation of a Hamiltonian with matrix elements $H_{fi} = \langle f|H|i\rangle$ for a final $|f\rangle$ and initial $|i\rangle$ basis state, $\langle H\rangle^{(\alpha)} = \sum_{i\subset\alpha} H_{ii}/\sum_{i\subset\alpha} 1$, and clearly, $H - \langle H\rangle^{(\alpha)}$ is traceless, such that $\mathrm{Tr}^{(\alpha)}(H - \langle H\rangle^{(\alpha)}) = \sum_{i\subset\alpha} H_{ii} - \langle H\rangle^{(\alpha)}\sum_{i\subset\alpha} 1 = 0$. We consider two distributions α , namely, a scalar distribution (denoted by "n" in the formulae) for a set of all states with a fixed n particle number, as well as isospin-scalar distribution (denoted by "n, T") for a set of all states with fixed n particle number and T isospin value². For a spectral distribution α (α is n or n, T), the inner product is defined as,

$$(H, H')^{(\alpha)} = \langle (H^{\dagger} - \langle H^{\dagger} \rangle^{\alpha})(H' - \langle H' \rangle^{\alpha}) \rangle^{\alpha} = \langle H^{\dagger} H' \rangle^{\alpha} - \langle H^{\dagger} \rangle^{\alpha} \langle H' \rangle^{\alpha}. \tag{4}$$

Hence, the correlation coefficient between two Hamiltonian operators, H and H', is given as

$$\zeta_{H,H'}^{(\alpha)} = \frac{(H,H')^{(\alpha)}}{\sigma_H^{(\alpha)}\sigma_{H'}^{(\alpha)}} = \frac{\langle H^{\dagger}H'\rangle^{\alpha} - \langle H^{\dagger}\rangle^{\alpha}\langle H'\rangle^{\alpha}}{\sigma_H^{(\alpha)}\sigma_{H'}^{(\alpha)}},\tag{5}$$

²A technical detail is that the model space in spectral distribution theory is partitioned according to particular group symmetries and each subsequent subgroup partitioning yields finer and more detailed spectral estimates. Specifically, for n particles distributed over 4Ω single-particle states, the scalar distribution averaged over all n-particle states is associated with the $U(4\Omega)$ group structure and the isospin-scalar distribution averaged over the ensemble of all n-particle states of isospin T is associated with $U(2\Omega) \otimes U(2)_T$.

where the "width" of H (related to the Hilbert-Schmidt norm) is the positive square root of the variance,

$$(\sigma_H^{(\alpha)})^2 = (H, H)^{(\alpha)} = \langle H^2 \rangle^{\alpha} - (\langle H \rangle^{\alpha})^2. \tag{6}$$

The steps for computing these quantities are outlined in Sections 2.1 and 2.2. The significance of a positive correlation coefficient is given by Cohen [46] and later revised to the following, $\zeta = 0.00 - 0.09$ represents a trivial correlation, $\zeta = 0.10 - 0.29$, 0.30 - 0.49, 0.50 - 0.69, and 0.70 - 0.89 represent small, medium, large and very large correlations, respectively, while $\zeta = 0.90 - 0.99$ and 1.00 represent nearly perfect and perfect correlations, respectively.

From a geometrical perspective, in spectral distribution theory every interaction is associated with a vector (Fig. 1) of length σ (Eq. 6) and the correlation coefficient ζ (Eq. 5) defines the angle (via a normalized scalar product) between two vectors. Hence, $\zeta_{H,H'}$ gives the normalized projection of H' onto the H interaction (or H onto H'). In addition, $(\zeta_{H,H'})^2$ gives the percentage of H' that reflects the characteristic properties of the H interaction. As for σ_H , it is a natural measure of the H operator size and realizes the spread of the H eigenvalue distribution. As is well-known, the smaller the σ_H (the weaker the interaction), the more compressed the energy spectrum for H [2].




Figure 1: Geometrical interpretation of two Hamiltonians in the framework of spectral distribution theory. Each Hamiltonian (H and H') is represented by a vector of length given by its respective norm σ ; the angle between H and H' is specified by the corresponding correlation coefficient, $\theta_{H,H'} = \arccos \zeta_{H,H'}$. H'_H is the projection of H' onto H and reflects the spectral characteristics of H.

As mentioned above, H and H' of Fig. 1 lie in a multi-dimensional space, spanned by a complete set of unit tensors. This allows one to identify, within this multi-dimensional space, a set of Hamiltonians $\{H\}$ (multi-dimensional vector H in Fig. 1) and then project any H' onto the space spanned by $\{H\}$, thereby providing an expansion of H' in terms of other Hamiltonians, $H' = \oplus H + H_{\text{res}}$. Here, H_{res} is the part of H' that lies outside the $\{H\}$ space (multi-dimensional dashed line in Fig. 1). In this study, we focus on the case of a two-dimensional $\{H\}$ space (Section 2.3).

The similarity between two interactions can be further tracked in manynucleon systems [4, 47] through propagation formulae. The latter determine how the averages extracted from the two-nucleon matrix elements get carried forward into many-nucleon systems. This propagation of information is modelindependent. In order to calculate energy moments and their propagation in many-particle systems, that is, for higher n (and T) values, each Hamiltonian H, which consists of one- (k=1) and two-body (k=2) interactions, is expressed as a linear combination of terms of definite particle rank (irreducible tensors $\mathcal{H}_k(\nu)$ of rank $\nu = 0, 1, 2$, that is as a collection of pure zero-, one- and two-body interactions. For example, in the case of scalar distribution and for n particles [6], the Hamiltonian can be rendered³,

$$H = n\mathcal{H}_1(0) + \binom{n}{2}\mathcal{H}_2(0) + \mathcal{H}_1(1) + (n-1)\mathcal{H}_2(1) + \mathcal{H}_2(2). \tag{7}$$

For then the inner product (Eq. 4) that defines the correlation coefficient (Eq. 5) or the variance (Eq. 6) is easily computed for different particle numbers n using the irreducible tensors and their expectation values in a many-particle basis,

$$\zeta_{H,H'}^{(\alpha)} \sigma_H^{(\alpha)} \sigma_{H'}^{(\alpha)} = \langle [\mathcal{H}^{\dagger}_{1}(1) + (n-1)\mathcal{H}^{\dagger}_{2}(1)][\mathcal{H}'_{1}(1) + (n-1)\mathcal{H}'_{2}(1)] \rangle^{\alpha}
+ \langle \mathcal{H}^{\dagger}_{2}(2)\mathcal{H}'_{2}(2) \rangle^{\alpha},$$
(8)
$$(\sigma_H^{(\alpha)})^2 = \langle [\mathcal{H}_{1}(1) + (n-1)\mathcal{H}_{2}(1)]^2 \rangle^{\alpha} + \langle [\mathcal{H}_{2}(2)]^2 \rangle^{\alpha}.$$
(9)

2.1. One-body interaction and single-particle energies SPE's

For a set s of s.p. states (typically, referred to as orbits) of degeneracy \mathcal{N}_s , the average single-particle energy ε and the traceless single-particle energy, $\tilde{\varepsilon}_r$, of the r^{th} orbit are given as,

$$\varepsilon = \frac{1}{\mathcal{N}} \sum_{s} \varepsilon_s \mathcal{N}_s \tag{10a}$$

$$\tilde{\varepsilon}_r = \varepsilon_r - \varepsilon = \varepsilon_r - \frac{1}{\mathcal{N}} \sum_s \varepsilon_s \mathcal{N}_s,$$
 (10b)

³ As explained by French [6], for a scalar distribution ($\alpha = n$), a 2-body Hamiltonian can be constructed in terms of simpler 0- and 1-body terms with the use of the one-body operator n. E.g., with (n-1) being (0+1)-body, if one finds a pure 1-body $\mathcal{H}_2(1)$ [that vanishes in the zero-particle space, $\mathcal{H}_2(1)|0\rangle = 0$, $(n-1)\mathcal{H}_2(1)$ can only be, in general, (1+2)-body - and as it vanishes in the one-particle space, for n = 1, it is in fact 2-body; similarly for $\binom{n}{2}\mathcal{H}_2(0)$ (it vanishes for n=0 and n=1). Hence, the generalization for constructing a k-body interaction in terms of pure particle-rank interactions is given as, $H = \sum_{\nu=0}^{k} {n-\nu \choose k-\nu} \mathcal{H}_k(\nu) =$ $\sum_{\nu=0}^{k} \frac{(n-\nu)(n-\nu-1)\dots(n-k+1)}{(k-\nu)!} \mathcal{H}_k(\nu) - \text{clearly, it vanishes for } n = \nu, \nu+1, \dots, k-1 \text{ due to}$ the *n*-dependent polynomial and for $n < \nu$ due to the ν -body $\mathcal{H}_k(\nu)$. Note that for an isospin-scalar distribution $(\alpha = n, T)$, two operators are invoked, n and \mathbf{T}^2 , such that, e.g., they vanish for n=0, T=0 and n=1, T=1/2).

where $\mathcal{N} = \sum_{s} \mathcal{N}_{s}$ and for a jj-coupled HO s.p. basis, $\mathcal{N}_{s} = 2\sum_{s}(2j_{s} + 1)$. A quick check for $\tilde{\varepsilon}$, $\operatorname{Tr}\tilde{\varepsilon} = \sum_{r} \mathcal{N}_{r}\tilde{\varepsilon}_{r} = \sum_{r} \mathcal{N}_{r}(\varepsilon_{r} - \varepsilon) = \mathcal{N}\varepsilon - \varepsilon \sum_{r} \mathcal{N}_{r} = 0$, confirms that it is traceless.

These quantities construct pure 0-body and 1-body terms (irreducible interaction tensors of particle ranks 0 and 1) for a one-body interaction, namely,

$$\mathcal{H}_1(0) = \varepsilon \mathcal{H}_1(1) = \sum_r \sqrt{[r]} \tilde{\varepsilon}_r \{ a_r^{\dagger} \otimes \tilde{a}_r \}^{(00)}.$$

With these definitions and using that $n = \sum_r \sqrt{[r]} \{a_r^{\dagger} \otimes \tilde{a}_r\}^{(00)}$, one can verify that a one-body $H = n\mathcal{H}_1(0) + \mathcal{H}_1(1)$, as shown in (7).

2.2. Two-body interaction and two-body matrix elements TBME's Scalar distribution. – In this case, degeneracy dimensionalities are given as,

$$\mathcal{N} = \sum_{r} \mathcal{N}_r$$
, where $\mathcal{N}_r = 2(2j_r + 1)$. (11)

For a two-particle system, the monopole moment (centroid), which is the average expectation value of the two-body interaction, is defined as

$$V_c = \frac{1}{\binom{N}{2}} \sum_{r \le s} \Gamma [\Gamma] W_{rsrs}^{\Gamma}, \tag{12}$$

where the Γ -sum goes over all possible (J,T) for given r and s, while $\binom{\mathcal{N}}{2} = \sum_{r \leq s,\Gamma} [\Gamma]$. The traceless induced single-particle energy is constructed as (that is, by contraction of the two-body interaction into an effective one-body operator under the particular group structure),

$$\lambda_{rt} = \frac{1}{\mathcal{N}_r} \sum_{s,\Gamma} [\Gamma] W_{rsts}^{\Gamma} \sqrt{(1+\delta_{rs})(1+\delta_{ts})} \hat{\delta}_{rt} - \frac{\delta_{rt}}{\mathcal{N}} \sum_{tu,\Gamma} [\Gamma] W_{tutu}^{\Gamma} (1+\delta_{tu}), \quad (13)$$

where $\hat{\delta}_{rt}$ is 1 if $j_r = j_t$ and $(-1)^{\eta_r} = (-1)^{\eta_t}$. We thus take into account a radial degeneracy [3, 2], that is, not every s.p. state is distinguishable by the j angular momentum and parity, thereby leading to an effective 1-body interaction that mixes different oscillator quantum numbers (equivalently, radial quantum numbers). This, for example, introduces a nonzero λ_{rt} with $r = \{\eta_r = 0, j_r = \frac{1}{2}, \tau_r = \frac{1}{2}\}$ and $t = \{\eta_t = 2, j_t = \frac{1}{2}, \tau_t = \frac{1}{2}\}$.

In turn, the traceless pure two-body interaction is defined through the antisymmetrized matrix elements,

$$W_{rstu}^{\Gamma}(2) = W_{rstu}^{\Gamma} - V_c \delta_{rt} \delta_{su} - \frac{\lambda_{rt} \delta_{su} + \lambda_{su} \delta_{rt} - (-1)^{r+s-\Gamma} (\lambda_{st} \delta_{ru} + \lambda_{ru} \delta_{st})}{(\mathcal{N} - 2)\sqrt{(1 + \delta_{rs})(1 + \delta_{ts})}}.$$
(14)

The quantities defined above, V_c , λ , and W(2), specify the corresponding tensors of particle ranks 0, 1, and 2 for a two-body interaction, namely,

$$\mathcal{H}_{2}(0) = V_{c},$$

$$\mathcal{H}_{2}(1) = \sum_{rs} \sqrt{[r]} \frac{\lambda_{rs}}{\mathcal{N} - 2} \{a_{r}^{\dagger} \otimes \tilde{a}_{s}\}^{(00)},$$

$$\mathcal{H}_{2}(2) = \sum^{*} \frac{\sqrt{[\Gamma]}}{\sqrt{(1 + \delta_{rs})(1 + \delta_{tu})}} W_{rstu}^{\Gamma}(2) \{\{a_{r}^{\dagger} \otimes a_{s}^{\dagger}\}^{\Gamma} \otimes \{\tilde{a}_{t} \otimes \tilde{a}_{u}\}^{\Gamma}\}^{(00)},$$

$$(15)$$

where the sum \sum^* goes over $r \leq s, t \leq u$ and $\Gamma = (J, T)$.

Hence, the quantity that defines the correlation coefficient (Eq. 5) and the norm for H = H' (Eq. 6) is easily computed for a larger number of particles, n,

$$\zeta_{H,H'}^{(n)}\sigma_H^{(n)}\sigma_{H'}^{(n)} = \langle H^{\dagger}H'\rangle^n - \langle H^{\dagger}\rangle^n \langle H'\rangle^n = \frac{n(\mathcal{N}-n)}{(\mathcal{N}-1)} \left(\frac{1}{\mathcal{N}} \sum_r \mathcal{N}_r \tilde{\varepsilon}_r \tilde{\varepsilon}_r'\right) +$$
(16a)

$$\frac{n(\mathcal{N}-n)(n-1)}{(\mathcal{N}-1)(\mathcal{N}-2)} \left(\frac{1}{\mathcal{N}} \sum_{r} \mathcal{N}_r(\tilde{\varepsilon}_r \lambda_r' + \lambda_r \tilde{\varepsilon}_r') \right) + \tag{16b}$$

$$\frac{n(\mathcal{N}-n)(n-1)^2}{(\mathcal{N}-1)(\mathcal{N}-2)^2} \left(\frac{1}{\mathcal{N}} \sum_{rs} \mathcal{N}_r \lambda_{rs} \lambda'_{rs} \right) + \tag{16c}$$

$$\frac{n(n-1)(\mathcal{N}-n)(\mathcal{N}-n-1)}{2(\mathcal{N}-2)(\mathcal{N}-3)} \left(\frac{1}{\binom{\mathcal{N}}{2}} \sum^* [\Gamma] W_{rstu}^{\Gamma}(2) W_{rstu}^{\prime \Gamma}(2) \right). (16d)$$

Isospin-scalar distribution. – In this case, degeneracy dimensionalities are given as.

$$\mathcal{N} = \sum_{r} \mathcal{N}_r$$
, where $\mathcal{N}_r = 2j_r + 1$, (17)

and the centroid is,

$$V_c^T = \frac{2}{\mathcal{N}(\mathcal{N} + (-1)^T)} \sum_{r \le s, J} [J] W_{rsrs}^{JT}.$$
 (18)

The λ_{rt}^T traceless induced single-particle energies, which specify $\mathcal{H}_2(1)$, and the $W_{rstu}^{JT}(2)$ two-body matrix elements [9] that specify the traceless pure 2-body interaction $\mathcal{H}_2(2)$ are defined as,

$$\lambda_{rt}^{T} = \frac{1}{\mathcal{N}_r} \sum_{s,I} [J] W_{rsts}^{JT} \sqrt{(1+\delta_{rs})(1+\delta_{ts})} \hat{\delta}_{rt} - \frac{\delta_{rt}}{\mathcal{N}} \sum_{tu,I} [J] W_{tutu}^{JT} (1+\delta_{tu}), \quad (19)$$

$$W_{rstu}^{JT}(2) = W_{rstu}^{JT} - V_c^T \delta_{rt} \delta_{su} - \frac{\lambda_{rt}^T \delta_{su} + \lambda_{su}^T \delta_{rt} - (-1)^{r+s-J-T} (\lambda_{st}^T \delta_{ru} + \lambda_{ru}^T \delta_{st})}{(\mathcal{N} + 2(-1)^T) \sqrt{(1 + \delta_{rs})(1 + \delta_{ts})}}.$$
 (20)

The quantities defined above are used to calculate the correlation coefficient $\zeta^{(n,T)}$ (Eq. 5) and the variance $(\sigma^{(n,T)})^2$ for H = H' (Eq. 6) for higher values of n and T, by employing

$$\zeta_{H,H'}^{(n,T)}\sigma_H^{(n,T)}\sigma_{H'}^{(n,T)} = \langle H^{\dagger}H'\rangle^{n,T} - \langle H^{\dagger}\rangle^{n,T}\langle H'\rangle^{n,T} =$$

$$\mathcal{P}_1(n,T)\left(\frac{1}{\mathcal{N}}\sum_r \mathcal{N}_r\tilde{\varepsilon}_r\tilde{\varepsilon}_r'\right) +$$
(21a)

$$\sum_{\tau} \mathcal{P}_1(n, T, \tau) \left(\frac{1}{\mathcal{N}} \sum_{r} \mathcal{N}_r(\tilde{\varepsilon}_r \lambda_r' + \lambda_r \tilde{\varepsilon}_r') \right) + \tag{21b}$$

$$\sum_{\{\tau_1,\tau_2\}} \mathcal{P}_1(n,T,\tau_1,\tau_2) \left(\frac{1}{\mathcal{N}} \sum_{rs} \mathcal{N}_r \frac{1}{2} \left[\lambda_{rs}^{\tau_1} \lambda_{rs}^{\prime \tau_2} + \lambda_{rs}^{\tau_2} \lambda_{rs}^{\prime \tau_1} \right] \right) + \tag{21}c)$$

$$\sum_{\tau} \mathcal{P}_2(n, T, \tau) \left(\frac{2}{\mathcal{N}(\mathcal{N} + (-1)^{\tau})} \sum^* [J] W_{rstu}^{J\tau}(2) W_{rstu}^{J\tau}(2) \right), \quad (21d)$$

where τ is 0 or 1, and the set $\{\tau_1, \tau_2\}$ is $\{0, 0\}, \{0, 1\}$ or $\{1, 1\}$. The sum \sum^* goes over $r \leq s, t \leq u$ and J. The propagator functions are derived in [48, 4] and shown below for completeness:

$$\mathcal{P}_{1}(n,T) = \frac{n(\mathcal{N}+2)(\mathcal{N}-\frac{n}{2})-2\mathcal{N}T(T+1)}{(\mathcal{N}-1)(\mathcal{N}+1)}, \tag{22}$$

$$\mathcal{P}_{1}(n,T,\tau) = \frac{4\mathcal{N}T(T+1)(1-n)(1-(-1)^{\tau})+(\mathcal{N}+2)(\mathcal{N}-\frac{n}{2})[(2\tau+1)n(n+2(-1)^{\tau})-4T(T+1)(-1)^{\tau}]}{4(\mathcal{N}-1)(\mathcal{N}+1)(\mathcal{N}+2(-1)^{\tau})},$$

$$\mathcal{P}_{1}(n,T,\tau_{1}=\text{odd},\tau_{2}) = \frac{8\mathcal{N}T(T+1)(n-1)(\mathcal{N}-2n+4)+[(2\tau_{1}+1)n(n-2)+4T(T+1)][\frac{(2\tau_{2}+1)(n+2(-1)^{\tau_{2}})}{2}(\mathcal{N}-\frac{n}{2})+T(T+1)(-1)^{\tau_{2}}][\mathcal{N}-2(-1)^{\tau_{2}}]}{8(\mathcal{N}-1)(\mathcal{N}+1)(\mathcal{N}-2)^{2}}.$$

$$\mathcal{P}_1(n, T, \tau_1 = \text{even}, \tau_2) =$$

$$\frac{[(2\tau_1+1)n(n+2)-4T(T+1)][\frac{(2\tau_2+1)(n+2(-1)^{\tau_2})}{2}(\mathcal{N}-\frac{n}{2})+T(T+1)(-1)^{\tau_2}][\mathcal{N}-2(-1)^{\tau_2}]}{8(\mathcal{N}-1)(\mathcal{N}+1)(\mathcal{N}-2)(\mathcal{N}+2)}$$

$$\mathcal{P}_2(n,T,\tau=0) = \frac{[n(n+2)-4T(T+1)][(\mathcal{N}-\frac{n}{2})(\mathcal{N}-\frac{n}{2}+1)-T(T+1)]}{8\mathcal{N}(\mathcal{N}-1)},$$

$$\mathcal{P}_2(n, T, \tau = 0) = \frac{[n(n+2) - 4T(T+1)][(\mathcal{N} - \frac{n}{2})(\mathcal{N} - \frac{n}{2} + 1) - T(T+1)]}{8\mathcal{N}(\mathcal{N} - 1)},$$

$$\mathcal{P}_2(n,T,\tau=1) = \frac{T^2(T+1)^2 \frac{3\mathcal{N}^2 - 7\mathcal{N} + 6}{2} + \frac{3n(n-2)}{8}(\mathcal{N} - \frac{n}{2})(\mathcal{N} - \frac{n}{2} - 1)(\mathcal{N} + 1)(\mathcal{N} + 2)}{\mathcal{N}(\mathcal{N} + 1)(\mathcal{N} - 2)(\mathcal{N} - 3)} + \frac{\frac{T(T+1)}{2}[(5\mathcal{N} - 3)(\mathcal{N} + 2)n(\frac{n}{2} - \mathcal{N}) + \mathcal{N}(\mathcal{N} - 1)(\mathcal{N} + 1)(\mathcal{N} + 6)]}{\mathcal{N}(\mathcal{N} + 1)(\mathcal{N} - 2)(\mathcal{N} - 3)}.$$

2.3. Orthogonal Hamiltonians

Orthogonal Hamiltonians are important in SDT, as they provide an orthogonal "basis", along which a Hamiltonian in consideration, K, can be projected. Large overlaps of K with a "basis" Hamiltonian implies that features of the latter are also well realized in K, while the orthogonality ensures that other "basis" Hamiltonians point to complementary features. For example, if K projects onto a $Q \cdot Q$ interaction and an L^2 interaction on an equal footing, this does not necessarily imply that both $Q \cdot Q$ and L^2 are required to describe related features of K, as $Q \cdot Q$ and L^2 can indeed highly overlap.

In the SDT framework, as shown in Ref. [11], one can straightforwardly extract the traceless 2-body H_{\perp} orthogonal part of a Hamiltonian H with respect to a reference one, $H_{\rm Ref}$, such that the condition $\langle H_{\perp}^{\dagger}H_{\rm Ref}\rangle=0$ is satisfied. This is done in a way analogous to a Gram-Schmidt orthogonalization of two vectors \vec{h} (reference) and \vec{g} , so that a new vector orthogonal to \vec{h} is given as, $\vec{g}_{\perp}=\vec{g}-\frac{\vec{g}\cdot\vec{h}}{\vec{h}\cdot\vec{h}}\vec{h}=\vec{g}-\frac{\zeta_{g,h}\sigma_g}{\sigma_h}\vec{h}$. Similarly, the traceless 2-body H_{\perp} is constructed using,

$$W_{rstu}^{JT}(2)_H - \frac{\zeta_{H,H_{\text{Ref}}}\sigma_H}{\sigma_{H_{\text{Ref}}}}W_{rstu}^{JT}(2)_{H_{\text{Ref}}}.$$
 (23)

Then, as illustrated in Fig. 2, any interaction K can be projected (K_H) [49] onto a plane defined by the two 'orthogonal vectors', $K_H = c_{H_{Ref}} H_{Ref} + c_{H_{\perp}} H_{\perp}$ $(K = K_H + K_{res})$ with a residual interaction K_{res} perpendicular to this plane), where the projection coefficients are given by,

$$c_{H_{\perp}} = \zeta_{K,H_{\perp}} \sigma_K / \sigma_{H_{\perp}} \tag{24a}$$

$$c_{H_{\text{Ref}}} = \zeta_{K,H_{\text{Ref}}} \sigma_K / \sigma_{H_{\text{Ref}}},$$
 (24b)

and the cosine of the angle between K and the plane (or equally, and K_H) is,

$$\zeta_{K,K_H} = \sqrt{\zeta_{K,H_{\text{Ref}}}^2 + \zeta_{K,H_{\perp}}^2}.$$
 (24c)

The norm of K_H can be then calculated as $\sigma_{K_H} = \sigma_K \zeta_{K,K_H}$. A more interesting case is a normalized projection of K (which preserves the norm of H_{Ref} , see Fig. 2), $K_{H(\text{Ref})} = \frac{K_H}{c_{H_{\text{Ref}}}} = H_{\text{Ref}} + \frac{c_{H_{\perp}}}{c_{H_{\text{Ref}}}} H_{\perp}$, with a norm,

$$\sigma_{K_{H(\text{Ref})}} = \sigma_{H_{\text{Ref}}} \sqrt{\left(1 + \zeta_{K,H_{\perp}}^2 / \zeta_{K,H_{\text{Ref}}}^2\right)}.$$
 (24*d*)

This allows for studies that utilize H_{Ref} that has been already deduced from selected nuclear properties, such as a standard pairing Hamiltonian with a pairing strength adjusted to reproduce pairing gaps in nuclei (also see [33]).




Figure 2: Geometrical interpretation of expanding a Hamiltonian K in terms of two Hamiltonians H_{Ref} and H_{\perp} , that is, projecting a vector K onto a plane defined by the two 'orthogonal vectors'. K_H is the projection of K onto the plane, while $K_{H(Ref)}$ is the portion that does not involve H_{Ref} renormalization.

6

10

3. Program Performance

According to the current default (see <calculateMomentsDr.h>), the program can handle up to 45 single-j orbits (that is, 9 major HO shells, $\eta =$ $0, 1, \dots, 8$), as well as up to 20 Hamiltonians that are simultaneously considered in a given calculation. These limits can be increased if an application requires more orbits and/or a larger set of Hamiltonians. In addition, a subset of orbits can be selected.




Figure 3: Run-time as a function of (a) the size of the s.p. basis using 3 Hamiltonians and (b) the number of Hamiltonians that are simultaneously considered in a given calculation employing 15 single-j orbits.

The computer program has been applied to various interactions including the realistic interactions N³LO [40], JISP16 [50], CD-Bonn [42] and AV18 [41], as well as their effective counterparts. Only the isospin-conserving part,

$$W_{rstu}^{JT(IC)} = \begin{cases} W_{rstu}^{JT(pn)} &, T = 0\\ \frac{1}{3}(W_{rstu}^{JT(pp)} + W_{rstu}^{JT(nn)} + W_{rstu}^{JT(pn)}) &, T = 1 \end{cases}$$
(25)

has been considered for charge-dependent interactions, that is, with different pp $(T_0 = 1)$, nn $(T_0 = -1)$, and pn $(T_0 = 0)$ T = 1 matrix elements, $W_{rstu}^{JT(T_0)} = \langle rsJTMT_0|H|tuJTMT_0\rangle$. The much smaller isospin-mixing part of nuclear interactions cannot be treated in the current program, but could be included, as the associated formulae have been already derived in Ref. [4]. In addition, several other interactions have been utilized, e.g., GXPF1 [37], as well as conventional pn- plus like-particle pairing and $Q \cdot Q$ interactions, and the Sp(4) pairing Hamiltonian [51]. Interactions may be specified for a large set of orbits and only a subset of orbits can be selected in the program.

The program run-time is studied as a function of the number of s.p. states, as well as the number of Hamiltonians considered (Fig. 3). While the runtime increases exponentially, as expected, for larger model spaces (more orbits considered), it remains reasonably small even when major shells up to the sdgik shell are considered (45 single-j orbits). For a broad range of problems of interest that explore up through the sdg shell (15 orbits), the program is indeed very fast (Fig. 3a). Furthermore, the run-time is only linearly dependent on the number of Hamiltonians considered, as shown in Fig. 3b for 15 orbits. This makes the program easily applicable to a large set of Hamiltonians, which in turn benefits calculations of projecting a given interaction K simultaneously along many distinct interactions $\{H\}$, especially when symmetry-preserving terms are involved. Such a project is currently being implemented as an extension of the present computational code.

Finally, the program benefits from an efficient way for storing TBME's, namely, matrix elements are stored in linked lists that are grouped, while being read, according to the role each W_{rstu}^{JT} plays in various sums that are needed to construct irreducible tensors. For example, one group consists of all W_{rsrs}^{JT} and another group consists of all W_{rsts}^{JT} ($r \neq t$), thereby facilitating fast computations of sums that enter in V_c and λ 's, respectively.

4. Program Description

In addition to the <main.c> file, the package consists of eight additional files that are described below (omitting functions that are self-explanatory).

4.1. File <calculateMomentsDr.c>

This file includes drivers that read Hamiltonians, calculate centroids, λ 's, W(2)'s, norms, and correlation coefficients, and write calculated quantities to output files.

InitModelOrb – Performs initialization of variables for handling the orbits, Hamiltonians, and statistical measures.

Hamiltonian – Reads all Hamiltonians [their respective SPE's (1-body) and TBME's (2-body)].

CalculateStatistics – Calculates statistical moments for all Hamiltonians (including orthogonalized interactions): first, the irreducible interaction tensors (pure 0-, 1-, and 2-body) are determined; next, second-order energy moments are calculated and stored for the base case [n-independent averages]. Calculates correlation coefficients and norms for $n \leq 4\Omega - 1$ [and $T \leq \min(\frac{n}{2}, 2\Omega - \frac{n}{2})$ in the isospin-scalar case], according to user's choice.

NewModelTBMEOrthog – Introduces a new Hamiltonian that contains the orthogonal part of a given Hamiltonian with respect to a reference one and for the new Hamiltonian, calculates quantities needed for statistical moments.

AllStatsOfModel – Stores centroids and λ 's for all Hamiltonians.

StatsOfModel – Stores centroids and λ 's for a given Hamiltonian.

NullZeroLambdas - Handles computational errors and zeroes $\lambda \leq 10^{-5}$.

AllAssignNorm - Stores the norm for all 2-body interactions (for n=2 particles).

AssignNorm - Stores the norm for a given 2-body interaction (for n=2 particles).

ClearAll - Clears memory.

SetClock - Resets the clock.

RunTime - Calculates program run-time.

4.2. File <calculateMoments.c>

This file includes functions that calculate centroids, λ 's, W(2)'s, norms, and correlation coefficients.

TracelessSPE – Calculates the average [Eq. (10a)] and traceless energies [Eq. (10b)] for external SPE's.

Int0b_1b - Calculates the centroids [Eqs. (12) & (18)] and λ 's [Eqs. (13) & (19)] for a given 2-body interaction.

SumV - Calculates sums linear in W_{rstu}^{JT} 's [needed for computing V_c and λ].

AssnV2 — Calculates W(2) [Eqs. (14) & (20)] for the irreducible 2-body part of a given 2-body interaction.

Avg2b - Calculates Eqs. (16d) & (21d) for $\langle \mathcal{H}^{\dagger}_{2}(2)\mathcal{H}'_{2}(2)\rangle$.

SumAvg2b - Calculates the *n*-independent sum in parenthesis of Eqs. (16*d*) & (21*d*) for $\langle \mathcal{H}^{\dagger}_{2}(2)\mathcal{H}'_{2}(2)\rangle$.

SumV2 – Calculates a sum quadratic in W_{rstu}^{JT} 's [needed for computing second-order energy moments].

Avg1b_lmlm - Calculates Eqs. (16c) & (21c) for $\langle \mathcal{H}^{\dagger}_{2}(1)\mathcal{H}'_{2}(1)\rangle$.

SumAvg1b_lmlm - Calculates the *n*-independent sum in parenthesis of Eqs. (16*c*) & (21*c*) for $\langle \mathcal{H}^{\dagger}_{2}(1)\mathcal{H}'_{2}(1)\rangle$.

Avg1b_spelm - Calculates Eqs. (16b) & (21b) for $\langle \mathcal{H}^{\dagger}_{1}(1)\mathcal{H}'_{2}(1)\rangle$.

SumAvg1b_spelm - Calculates the *n*-independent sum in parenthesis of Eqs. (16b) & (21b) for $\langle \mathcal{H}^{\dagger}_{1}(1)\mathcal{H}'_{2}(1)\rangle$.

 ${\tt Avg1b_lmspe} \ - Calculates \ Eqs. \ (16b) \ \& \ (21b) \ for \ \langle \mathcal{H^\dagger}_2(1)\mathcal{H'}_1(1)\rangle.$

SumAvg1b_lmspe - Calculates the *n*-independent sum in parenthesis of Eqs. (16b) & (21b) for $\langle \mathcal{H}^{\dagger}_{2}(1)\mathcal{H}'_{1}(1)\rangle$.

- Avg1b_spe Calculates Eqs. (16a) & (21a) for $\langle \mathcal{H}^{\dagger}_{1}(1)\mathcal{H}'_{1}(1)\rangle$.
- SumAvg1b_spe Calculates the *n*-independent sum in parenthesis of Eqs. (16a) & (21a) for $\langle \mathcal{H}^{\dagger}_{1}(1)\mathcal{H}'_{1}(1)\rangle$.
- AvgTot Calculates total averages [Eqs. (16) & (21)] used to compute second-order energy moments for any n and T (calculates the variance [norm squared] in the case when H = H').
- AssignAvgBaseCase Allocates memory and stores all n-independent sums in parenthesis of Eqs. (16) & (21) for all Hamiltonians ('base case').
- SetSumAvg Stores all n-independent sum in parenthesis of Eqs. (16) & (21) for two given Hamiltonians ('base case').
- CorrelCoeff Calculates correlation coefficients for given two Hamiltonians, space partitioning (scalar or isospin-scalar distribution) and number of particles.
- CreateModelTBMEOrthogonal Creates a new traceless 2-body Hamiltonian (with zero SPE's, V_c and λ 's, by construction) that contains only the orthogonal part of a given Hamiltonian with respect to a reference Hamiltonian according to Eq.(23).
- DestroyNewModel Clears the last Hamiltonian that has been added during run-time to the user's list of Hamiltonians.
- StatsRefPlusOrthog Calculates correlation coefficients (24c), norms (24d), and projection coefficients (24a) & (24b) for interactions projected onto a reference and orthogonalized interactions.
- NameNewMode Provides an index for an orthogonalized Hamiltonian based on the indices of the Hamiltonian to be orthogonalized (id) and the reference one (id_{Ref}). E.g. the orthogonal of id = 23 with respect to id_{Ref} = 22 is indexed as 2223.
- STmodel Introduces three new labels for a given Hamiltonian that correspond to a scalar distribution (all T), an isospin-scalar distribution for T=0, and an isospin-scalar distribution for T=1.

4.3. File <tools.c>

This file contains a number of auxiliary functions, which are designed to handle s.p. basis states and TBME's. The s.p. basis states are indexed by positive integer numbers. All SPE's and TBME's are given in terms of these indices as an input. For nuclear Hamiltonians in a HO jj-coupled basis, indices k ($k=1,2,3,\ldots$) are related to the η (referred to as n in the program) and j quantum numbers of a s.p. state by $k=\frac{\eta(\eta+1)}{2}+j+\frac{1}{2}$. For example, k=1 ($0s_{1/2}$), k=2 ($1p_{1/2}$), k=3 ($1p_{3/2}$), and so on. Two-body matrix elements W_{ijkl}^{JT} are supplied only for indices.

$$i \leq j, \, i \leq k, \, \mathrm{and} \left\{ \begin{array}{l} j \leq l, \, \mathrm{if} \, \, i = k \\ k \leq l, \, \mathrm{if} \, \, i < k \end{array} \right. \, ,$$

as all the TBME's can be then obtained using relations (2). The TBME's are sorted, while being read, in ascending order in the i-j-k-l-J quantum numbers and grouped according to T.

- GroupME Groups TBME's according to their indices (e.g., ijij and ijkj) to reduce computational time for calculating various sums.
- allConfigurationsList Creates a model (called "modelAllConfig") with all possible configurations, $\{ijkl, JT\}$, that enter V_c and λ 's, as well as provides checks for dimensionalities

- FactorME Provides the ability to rescale TBME's by an n-dependent factor for a given Hamiltonian in calculating second-order energy moments. Current default: returns 1. Example given in comments is for GXPF1 [37] with TBME's decreasing as $[42/(40+n)]^{0.3}$.
- ScalingME Provides the ability to rescale TBME's by a factor. Current default: a scaling example, $W_{rstu}^{JT} \to W_{rstu}^{JT} / \sqrt{(\eta_r + \eta_s)(\eta_t + \eta_u)}$, for nonzero $\eta_r, \eta_s, \eta_t, \eta_u$ and for Hamiltonians with indices between 80-89 (the rescaling is done while reading the TBME's from a file).

4.4. File c>

This file contains propagation functions, n-dependent functions in Eqs. (16a-16d) for the scalar case and Eq. (22) for the isospin-scalar case, used to calculate second-order energy moments for a many-particle system (for any number of particles n and isospin T).

4.5. I/O Files <readFile.c> and <writeFile.c>

The functionalities in <readFile.c> include reading the user input menu as well as reading and storing (in arrays or linked lists) SPE's and TBME's of a set of Hamiltonians specified by the user

The <writeFile.c> consists of functions for writing the output. In the output files, the number of particles is denoted as m and the isospin values are given as twice the value, 2T.

PrintListModel - Prints TBME's to a file for all T for a given Hamiltonian.

PrintListModelT - Prints TBME's to a file for a specific T for a given Hamiltonian.

PrintLambda – Prints λ 's for a given Hamiltonian and space partitioning (and isospin).

PrintLambdaH – Prints λ 's for all Hamiltonians and a given space partitioning.

PrintStatsOfModel - Prints centroids, λ 's, and norms of a Hamiltonian.

PrintStatsOfAllModel – Prints centroids, λ 's, and norms for all Hamiltonians.

PrintNorm – Prints norms for all Hamiltonians given a space partitioning for all n (and T).

PrintNormPairsModel – Prints norms for all Hamiltonians for a given n (and T).

PrintCorrelCoef – Prints correlation coefficients for all Hamiltonians given a space partitioning for all n (and T).

PrintcCPairsModel – Prints correlation coefficients for all pairs of Hamiltonians for a given n (and T), e.g., (H_1, H_2) (H_1, H_3) (H_2, H_3) .

PrintCCPairsOneModel – Prints correlation coefficients for a given Hamiltonian for a given n (and T), e.g., for H_1 , (H_1, H_2) (H_1, H_3) .

PrintRefOrthogQnty – Prints correlation coefficients (24c), norms (24d), and projection coefficients (24a) & (24b) for interactions projected onto a reference and orthogonalized interactions.

PrintCorrelCoefEachUnitaryRk – Prints correlation coefficients for irreducible 1-body and 2-body interaction tensors.

PrintList2mdls – (additional) Prints TBME's of two models for the same set of quantum numbers [zero TBME's are included if missing in one model, but existing in the other; also, every TBME of J and T is printed (2J+1)(2T+1) times].

TestOrthogMdls - Provides a test if an orthogonalized Hamiltonian is orthogonal to its reference Hamiltonian.

PrintInitNorm, PrintInitCCheader, PrintInitCC(OneModel), PrintInitRefOrthogQnty - Write various headers that provide labels for each type of calculations in the output file.

4.6. Files <arrayFns.c> and <linkList.c>

These files contain conventional array and linked-list manipulations.

4.7. Data-type definitions and variables

- caseST Data type for distributions: kScalar (or 0) if scalar distribution; kTScalar (or 1)
 if isospin-scalar distribution.
- orbType Data type for indices: jqn0rb refers to the 2j angular momentum of a s.p. state; index0rb refers to the index k = 1, 2, 3, ... of a s.p. state.
- intPart Data type for irreducible tensors: hAll specifies the entire 1-body+2-body Hamiltonian (all irreducible tensors), hLm2b refers to $\mathcal{H}_2(1) + \mathcal{H}_2(2)$, h2b refers to $\mathcal{H}_2(2)$, and hLm refers to $\mathcal{H}_2(1)$.
- meType Data type for TBME's W^{JT}_{ijkl} : kiOrb, kjOrb, kkOrb, klOrb [s.p. indices $i,\ j,\ k,\ l$]; kJ and kT [2J and 2T]; dimJ and dimJT [2J+1 and (2J+1)(2T+1)]; vInt and vInt2 $[W^{JT}_{ijkl}$ and $W^{JT}_{ijkl}(2)$]; numericME and numericME2 [true if W^{JT}_{ijkl} and $W^{JT}_{ijkl}(2)$ assigned].
- statType Data type for key characteristics of a Hamiltonian, namely, V_c , λ 's and norms.
- avgBCType Data type for base-case averages.
- arrayModels, arrayOrbits, arrayDimOrb, arrayRadDeg, numRadDeg, arraySPE Arrays to store Hamiltonian id's, orbits, dimensionalities, radial-degenerate orbits, number of radial-degenerate orbits for a given orbit, and SPE's, respectively.
- arrayStat, arrayAvgBaseCase Arrays to store V_c , λ 's, and norms of Hamiltonians (arrayStat), as well as base-case averages (arrayAvgBaseCase).
- head Array of linked-list heads for TBME's of all Hamiltonians.
- kModels, kOrbits, kShells, kOrthogMdls Provides the total number of Hamiltonians, orbits, major HO shells, and orthogonalized Hamiltonians, respectively.
- numModels, numOrbits, orthogInModels Provides the maximum allowed number of Hamiltonians [including orthogonalized Hamiltonians and modelAllConfig that is automatically created], orbits, and orthogonalized Hamiltonians, respectively.

Acknowledgments

This work was supported by the U.S. National Science Foundation (OCI-0904874), the U.S. Department of Energy (DE-SC0005248), and the Southeastern Universities Research Association.

Appendix

Sample input files for Hamiltonians

```
1 1 1 4 2 0 -7.684559
1 1 2 2 2 0 -1.161099
1 1 2 7 2 0 -1.377213
1 2 1 2 2 0 5.997909
2 9 7 8 4 2 -5.486247
For further details, see Sec. 4.3.
Program input file
Test input (file) for the output shown below:
                     /* NO. of orbits
3
4 5 6
                     /* no. of orbits
/* orbits
/* NO. of interactions
                     /* NU. of interactions
/* interactions (TBME)
/* interactions (TBME)
/* 0 sets all external s.p.energies=0 for all interactions; 1 reads from files
/* output: (1) CC; (2) CC+norms; (3) 2 + additional CC/norms, (4) 1+test
/* NO. of interactions to be orthogonalized
/* Href & Interaction to be orthogonalized to Href
21 22 23
22 23
23 21
Sample input (file) for TBME rescaling:
sp4_10int81_22_23
                                /* NO. of orbits
4 5 6 7 8 9 10
                                /* orbits /* NO. of interactions MESCALED
81 22 23
                                /* interactions (TBME)
                                /* Interactions (IBME)
/* O sets all external s.p.energies=O for all interactions; 1 reads from files
/* output: (1) CC; (2) CC+norms; (3) 2 + additional CC/norms, (4) 1+test
/* NO. of interactions to be orthogonalized
/* Href & Interaction to be orthogonalized to Href
1
22 23
User input menu:
Enter the name to label output files
USER INPUT MENU
MAXIMUM ALLOWED ORBITS: 45
Enter -1 to terminate program
Sample Input :-
NO. OF ORBITS => 3
ORBITS => 1 2 3
 Enter number of orbits:
Enter orbits in ASCENDING order:
ORBITS READ: 4 5 6
**************** INTERACTIONS ****************
Interactions are represented by numerical indices corresponding to their respective input files of two-body matrix elements, TBMEs (e.g. '8' --> TBME8.dat, '13' --> TBME13.dat)
Interactions with indices between 80-89 can be rescaled
(refer to ScalingME in tools.c).
For rescaling: add a comment containing MESCALED after entering number of interactions.
MAX. ALLOWED INTERACTIONS: 20
Enter -1 to terminate program
Sample Input :-
NO. OF INTERACTIONS => 2
INTERACTIONS => 21 23
Enter number of interactions (followed by MESCALED for rescaling):
```

1 1 1 1 2 0 0.803376

Enter interactions:

```
INTERACTIONS READ: 21 22 23
```

```
****** SINGLE PARTICLE ENERGIES (SPE's) ********
Single particle energies will be read from files corresponding to the interaction numerical indices % \left( 1\right) =\left( 1\right) +\left( 1\right)
 (e.g. '21'---> SPE21.dat, '22' ---> SPE22.dat)
0. No external SPE's (for all interactions)
   1. Reads SPE's from files (for all interactions)
 Enter (1/0/-1):
SPE's INCLUDED

    Correlation Coefficients (CC's)
    Correlation Coefficients (CC's) and Norms

 3. 2 + CC's for Irreducible (1-body and 2-body) Parts of Interactions
4. 1 + Tests for orthogonality
 Enter (1/2/3/4):
CHOICE FOR OUTPUT: 2
    ******* INTERACTIONS TO BE ORTHOGONALIZED *******
Interactions must be chosen from Interactions specified above.
MAX. ALLOWED PAIRS: 3
 Enter -1 to terminate program
 NO. OF INTERACTIONS TO BE ORTHOGONALIZED => 2
REF. AND ORTHOG. MODELS #1 => 22 23
REF. AND ORTHOG. MODELS #2 => 23 21
   *****************
 Enter number of interactions to be orthogonalized:
 Enter pairs: Reference Interaction Href and Interaction to be orthogonalized to Href
 Enter pair #1:
Enter pair #2:
 INTERACTIONS TO BE ORTHOGONALIZED: 23 (to 22) 21 (to 23)
 ::InitModelOrb:: Finish reading input ...
  ::Hamiltonian:: Reading SPE/TBME...
::CalculateStatistics:: Calculating moments...
 ::CalculateStatistics:: Scalar distribution...Finish calculating...
::CalculateStatistics:: Isospin-scalar distribution...Finish calculating...
 ::ClearAll:: End.
 Test output
 File <out_sp4_6int21_22_23.dat>, shown only for Scalar distribution (format: semicolon-delimited):
 ***********************************
 *Scalar distribution*
 Hamiltonian; 21; 22; 23; Vc; -2.81556; 0.11436; -1.66531;
                                                                                                                                                                              2223; 2321;
-0.00000; -0.00000;
                                                   9.82673;
                                                                                          9.43303; 10.35063; 10.18719;
                                                                                                                                                                                                                       9.81476;
 lambda:
                                             14.51213; -0.91548; 12.97585; 5.99441; -24.58659; -4.52222; -8.83365; 16.69622; -1.31047;
   4 4;
5 5;
                                                                                                                                                                                    0.00000;
                                                                                                                                                                                                                              0.00000;
                                                                                                                                                                                    0.00000:
                                                                                                                                                                                                                             0.00000:
                                                                                                                                                                                   0.00000;
                                                                                                                                                                                                                            0.00000;
Correlation Coefficients (H1,H2), Projection Norm | Hproj=Href+c*Hperp| and Projection Coefficients cH_H1 of H along H1:

m; (21,22); (21,223); (21,2223); (21,22+2223); |21proj|; c21_22; c21_2223; (21,23); (21,231); (21,2321); (21,23+2321); |21proj|; c21_23; c21_2321;

2; 0.07529; -0.0278; 0.08026; 1.006e+01; 0.07843; -0.0268; -0.0160; 0.99878; 0.99891; 6.464e+02; -0.0152; 1.00000;

3; 0.06932; -0.0278; 0.07469; 1.693e+01; 0.07154; -0.0268; -0.0149; 0.99977; 0.99988; 1.147e+03; -0.0141; 1.00000;

4; 0.06269; -0.0278; 0.06858; 2.332e+01; 0.06419; -0.0268; -0.0137; 0.99882; 0.99892; 1.680e+03; -0.0103; 1.00000;

5; 0.05546; -0.0278; 0.06202; 2.949e+01; 0.06532; -0.0268; -0.0123; 0.99725; 0.99733; 2.282e+03; -0.0117; 1.00000;

6; 0.04755; -0.0277; 0.05504; 3.573e+01; 0.04787; -0.0268; -0.0108; 0.99528; 0.99533; 3.010e+03; -0.0103; 1.00000;
```

```
-0.0276;
                                     0.04772; 4.275e+01; 0.03879; -0.0268;
                                                                                                0.99295;
         0.03890;
                                                                                   -0.0091;
                                                                                                               0.99299; 3.974e+03; -0.0087; 1.00000;
         0.02938;
                        -0.0276;
                                     0.04028; 5
                                                 .252e+01; 0.02900;
                                                                                   -0.0072;
                                                                                                 0.99026;
                                                                                                               0.99029;
                                                                                                                        5.437e+03;
                                                                                                                                    -0.0069;
  9:
         0.01886:
                        -0.0275
                                     0.03333:
                                                 .284e+01: 0.01840:
                                                                     -0.0268:
                                                                                   -0.0051:
                                                                                                 0.98717:
                                                                                                               0.98718:
                                                                                                                        8.203e+03:
                                                                                                                                    -0.0048:
                                                                                                                                              1.00000:
         0.00717
                       -0.0274;
                                     0.02830;
                                                                                   -0.0026;
                                                                                                 0.98360;
                                                                                                               0.98361;
                                                                                                                         1.656e+04;
                                                                                                                                    -0.0025;
                                                 .722e+02;
                                                          0.00691
                                                                     -0.0268
                                                                                                                                               .00000
 11:
         -0.0059
                       -0.0273:
                                     0.02790:
                                                 .146e+02: -0.0056:
                                                                     -0.0268
                                                                                   0.00023:
                                                                                                0.97946:
                                                                                                               0.97946:
                                                                                                                        1.903e+05:
                                                                                                                                    0.00022:
                                                                                                                                              1.00000:
                                                                                                               0.97463;
                                                                                                                                    0.00341:
         -0.0206
                        -0.0271
                                     0.03408
                                                           -0.0193:
                                                                                   0.00355
                                                                                                 0.97462
                                                 724e+01:
                                                                                                                         1.251e+04:
                                                                                   0.00746;
 13;
         -0.0374;
                       -0.0270;
                                     0.04607; 5.861e+01; -0.0343;
                                                                     -0.0268;
                                                                                                 0.96889;
                                                                                                               0.96892;
                                                                                                                        5.883e+03; 0.00719;
                                                                                                                                              1.00000;
 14;
15;
                                                                                                                        3.514e+03;
                                     0.06255;
                                                                                                 0.96200;
          -0.0565
                                                                                   0.01216
                                                                                                               0.96208;
         -0.0788
                       -0.0265:
                                                                                                               0.95375:
                                     0.08312: 4.999e+01: -0.0692:
                                                                     -0.0268
                                                                                   0.01789:
                                                                                                 0.95359:
                                                                                                                        2.284e+03;
                                                                                                                                    0.01739:
                                                                                                                                              1.00000:
                        -0.0263
                                     0.10815;
                                               4.787e+01;
                                                           -0.0897
                                                                                   0.02506
                                                                                                 0.94307;
                                                                                                               0.94341;
                                                                                                                        1.529e+03;
                                                                                                                                    0.02449;
 17;
         -0.1362;
                        -0.0259;
                                     0.13863; 4.571e+01; -0.1126;
                                                                     -0.0268
                                                                                   0.03426;
                                                                                                 0.92957;
                                                                                                               0.93020; 1.023e+03;
                                                                                                                                    0.03374;
                                                                                                                                              1.00000;
 18;
         -0.1744
                       -0.0254;
                                     0.17624; 4.319e+01; -0.1385
                                                                     -0.0268
                                                                                   0.04651:
                                                                                                 0.91160;
                                                                                                               0.91279:
                                                                                                                        6.693e+02:
                                                                                                                                    0.04628:
                                                                                                                                              1.00000
 19;
         -0.2224;
                       -0.0247;
                                     0.22377;
                                               4.014e+01; -0.1679;
                                                                     -0.0268
                                                                                   0.06365;
                                                                                                 0.88649;
                                                                                                               0.88877;
                                                                                                                        4.168e+02;
                                                                                                                                    0.06423;
                                                                                                                                              1.00000;
                                                                                                               0.85357;
 20;
21;
         -0.2851
                        -0.0236
                                     0.28605
                                               3.642e+01; -0.2017
                                                                                   0.08933
                                                                                                 0.84888
                                                                                                                        2.381e+02;
                                                                                                                                    0.09209;
         -0.3716;
                       -0.0219;
                                     0.37225; 3.185e+01; -0.2410;
                                                                                   0.13215;
                                                                                                 0.78609;
                                                                     -0.0268;
                                                                                                               0.79712; 1.164e+02; 0.14115; 1.00000;
 22.
         -0.5025
                        -0.0183:
                                     0.50287: 2.613e+01: -0.2870:
                                                                     -0.0268
                                                                                   0.21849
                                                                                                 0.65796
                                                                                                               0.69328: 4.129e+01: 0.25049:
                                                                                                                                             1 00000
 23;
                       0.00000;
                                                                                   0.49577;
                                                                                                0.00000;
          -0.7390;
                                     0.73901; 1.847e+01; -0.3419; 0.00000;
                                                                                                               0.49577; 5.978e+00; 0.70846; 0.00000;
Norms
      |H|:
          211:
 m;
                       221;
                                   231;
                                            [2223];
 2; 9.827e+00; 9.433e+00; 1.035e+01; 1.019e+01; 9.815e+00
3; 1.621e+01; 1.571e+01; 1.711e+01; 1.682e+01; 1.621e+01
                                        1.682e+01; 1.621e+01;
  4: 2.183e+01:
                 2.132e+01: 2.303e+01:
                                        2.263e+01:
                                                    2 180e+01:
  5; 2.678e+01; 2.637e+01; 2.823e+01;
                                        2.772e+01;
                                                    2.670e+01;
  6; 3.108e+01; 3.088e+01; 3.274e+01;
                                        3.211e+01:
                                                    3.094e+01:
  7; 3.476e+01; 3.485e+01; 3.656e+01;
                                        3.582e+01:
                                                    3.451e+01;
  8; 3.780e+01; 3.830e+01; 3.970e+01;
                                        3.885e+01:
                                                    3.743e+01:
  9; 4.022e+01; 4.122e+01; 4.217e+01;
                                        4.121e+01;
 10: 4.201e+01: 4.360e+01: 4.397e+01:
                                        4 289e+01 ·
                                                      132e+01:
                 4.545e+01;
 12; 4.373e+01; 4.676e+01; 4.554e+01;
                                        4.424e+01:
                                                    4.262e+01:
                 4.752e+01; 4.531e+01;
 14: 4.296e+01: 4.773e+01: 4.441e+01:
                                        4.289e+01:
                                                    4.132e+01:
 15; 4.164e+01;
                 4.738e+01;
                            4.285e+01;
                                        4.121e+01;
 16; 3.969e+01;
                 4.644e+01; 4.060e+01;
                                        3.885e+01;
                                                    3.743e+01;
 17; 3.713e+01; 4.491e+01; 3.769e+01;
                                        3.582e+01;
 18; 3.394e+01; 4.274e+01; 3.411e+01;
                                        3.211e+01: 3.094e+01:
 19; 3.012e+01; 3.990e+01; 2.985e+01;
                                        2.772e+01;
 20: 2.568e+01: 3.629e+01: 2.491e+01: 2.263e+01: 2.180e+01:
     2.062e+01; 3.180e+01; 1.931e+01;
                                        1.682e+01:
 22; 1.492e+01; 2.612e+01; 1.301e+01; 1.019e+01; 9.815e+00;
     8.543e+00; 1.847e+01; 5.978e+00; 0.000e+00; 0.000e+00;
```

This output file provides the centroid (listed as "Vc"), the norm, as well as the density-dependent 1-body part (listed as "lambda") for each Hamiltonian, together with various second-order moments listed for increasing number of particles (shown as "m") for the scalar distribution, and also for twice the isospin value (shown as "2T") for the isospin-scalar distribution. In a complete space (e.g., 24 particles for the output shown above, $m = 4\Omega$), moments are zero by definition, Eqs. (16) and (22), and the corresponding output is omitted.

Additional information, including dimensionality checks, set of input parameters, run-time, and messages on rescaling and orthogonalization, are written into a corresponding log-file <0.sp4.6int21.22.23.log>.

References

- [1] J. B. French, Phys. Lett. 23 (1966) 248; J. B. French, Phys. Lett. 26B (1967) 75
- [2] J. B. French and K. F. Ratcliff, Phys. Rev. C 3 (1971) 94
- [3] F. S. Chang, J. B. French, and T. H. Thio, Ann. Phys. (N.Y.) 66 (1971) 137
- [4] K. T. Hecht and J. P. Draayer, Nucl. Phys. A223 (1974) 285
- [5] V. K. B. Kota and R. U. Haq, "Spectral Distributions in Nuclei and Statistical Spectroscopy," (World Scientific Publishing Co., 2010)
- [6] J. B. French, in *Dynamic Structure of Nuclear States*, ed. D. J. Rowe et al. (Univ. of Toronto Press, Toronto, 1972) p.154
- [7] J. P. Draayer, Nucl. Phys. A216 (1973) 457
- [8] J. C. Parikh, Group Symmetries in Nuclear Structure (Plenum, New York, 1978)

- [9] V. K. B. Kota, Phys. Rev. C 20 (1979) 347; Fortran Programs for Statistical Spectroscopy Calculations
- [10] C. R. Countee, J. P. Draayer, T. R. Halemane, and K. Kar, Nucl. Phys. A356 (1981) 1
- [11] V. Potbhare, Nucl. Phys. A289 (1977) 373
- [12] J. B. French, Nucl. Phys. A396 (1983) 87c
- [13] J. P. Draayer and G. Rosensteel Phys. Lett. 124B (1983) 281; J. P. Draayer and G. Rosensteel, Phys. Lett. 125B (1983) 237; G. Rosensteel and J. P. Draayer, Nucl. Phys. A436 (1985) 445
- [14] S. Sarkar, K. Kar, and V. K. B. Kota, Phys. Rev. C 36 (1987) 2700
- [15] J. B. French, V. K. B. Kota, A. Pandey, and S. Tomsovic, Ann. Phys. (N.Y.) 181 (1988) 235
- [16] V. K. B. Kota and D. Majumdar, Z. Phys. A 351 (1995) 365; Z. Phys. A 351 (1995) 377
- [17] K. F. Ratcliff, Phys. Rev. C 3 (1971) 117
- [18] J. N. Ginnochio, Phys. Rev. Lett. **31** (1973) 1260
- [19] J. P. Draayer, J. B. French, M. Prasad, V. Potbhare, and S. S. M. Wong, Phys. Lett. 57B (1975) 130
- [20] J. B. French and V. K. B. Kota, Phys. Rev. Lett. 51 (1983) 2183
- [21] B. Strohmaier, S. M. Grimes and H. Satyanarayana, Phys. Rev. C 36 (1987) 1604
- [22] A. Abzouzi, E. Caurier, and A. P. Zuker, Phys. Rev. Lett. 66 (1991) 1134
- [23] T. R. Halemane, K. Kar, and J. P. Draayer, Nucl. Phys. A311 (1978) 1134
- [24] B. J. Dalton, W. J. Baldridge, and J. P. Vary, Phys. Rev. C 20 (1979) 1908
- [25] L. Benet, T. Rupp, and H. A. Weidenmiller, Phys. Rev. Lett. 87 (2001) 010601
- [26] S. Tomsovic, M. B. Johnson, A. Hayes, and J. D. Bowman, Phys. Rev. C 62 (2000) 054607
- [27] J. M. G. Gomez, K. Kar, V. K. B. Kota, R. A. Molina, and J. Retamosa, Phys. Lett. B 567 (2003) 251
- [28] M. Horoi, M. Ghita, and V. Zelevinsky, Phys. Rev. C 69 (2004) 041307(R); M. Horoi, J. Kaiser, and V. Zelevinsky, Phys. Rev. C 67 (2003) 054309
- [29] V. K. B. Kota, Phys. Rev. C 71 (2005) 041304(R)
- [30] Y. M. Zhao, A. Arima, N. Yoshida, K. Ogawa, N. Yoshinaga, and V. K. B. Kota, Phys. Rev. C 72 (2005) 064314
- [31] E. Teran and C. W. Johnson, Phys. Rev. C 74 (2006) 067302
- [32] K. D. Sviratcheva, J. P. Draayer, and J. P. Vary, Nucl. Phys. A 786 (2007) 31
- [33] K. D. Sviratcheva, J. P. Draayer, and J. P. Vary, Phys. Rev. C 73 (2006) 034324
- [34] A. F. Lisetskiy, B. R. Barrett, M. K. G. Kruse, P. Navratil, I. Stetcu, and J. P. Vary, Phys. Rev. C 78 (2008) 044302
- [35] G. Hagen et al., Phys. Rev. C 76 (2007) 034302
- [36] K. Tsukiyama, S. K. Bogner, and A. Schwenk, Phys. Rev. Lett. 106 (2011) 222502
- [37] M. Honma, T. Otsuka, B. A. Brown, and T. Mizusaki, Phys. Rev. C 69 (2004) 034335
- [38] T. Otsuka et al., Phys. Rev. Lett. 87 (2001) 082502
- [39] K. D. Launey, T. Dytrych, and J. P. Draayer, Phys. Rev. C 85 (2012) 044003
- $[40]\,$ D. R. Entem and R. Machleidt, Phys. Rev. C $\mathbf{68}$ (2003) 041001
- [41] R.B. Wiringa, V.G.J. Stoks, and R. Schiavilla, Phys. Rev. C 51 (1995) 38
- [42] R. Machleidt, Phys. Rev. C 63 (2001) 024001
- [43] B. D. Chang, J. P. Draayer, and S. S. M. Wong, Comput. Phys. Commun. 28 (1982) 41
- [44] S. M. Grimes, S. D. Bloom, H. K. Vonach, R. F. Hausman, Jr., Phys. Rev. C 27 (1983) 2893
- [45] M. A. Caprio, F. Q. Luo, K. Cai, V. Hellemans, C. Constantinou, Phys. Rev. C 85 (2012) 034324
- [46] J. Cohen, Statistical Power Analysis for the Behavioral Sciences (Lawrence Erlbaum Associates, Hillsdale, NJ) (1988); J. Cohen, P. Cohen, S. G. West, and L. S. Aiken, Applied multiple regression/correlation analysis for the behavioral sciences, 2nd ed. (Lawrence Erl-

- baum Associates, Hillsdale, NJ) (2003)
- [47] V. K. B. Kota, S. P. Pandya, and V. Potbhare, Nucl. Phys. A349 (1980) 397
- [48] J. B. French, in *Isospin in Nuclear Physics*, ed. D. H. Wilkinson (North Holland, Amsterdam, 1969), p.259
- [49] J. P. Draayer and G. Rosensteel, Nucl. Phys. A386 (1982) 189
- [50] A. M. Shirokov et al., Phys. Letts. B **644** (2007) 33
- [51] K. D. Sviratcheva, A. I. Georgieva, and J. P. Draayer, Phys. Rev. C 70 (2004) 064302